

phases prodrugs which improve the pharmacokinetic properties of anticancer agents and, in particular, prodrugs which are activated selectively in tumour cells to the active drug.

## 2. Prodrugs with altered chemical stability, solubility or better tissue penetration

The chemical instability of some anticancer agents when dissolved in physiological saline is well recognised and is overcome, for example, in the case of reactive alkylating agents by ensuring rapid dissolution and injection, and, in the case of formulations of dimethyltriazenoimidazole carboxamide that are light unstable, by the use of dark vessels and the avoidance of sunlight. Other agents may decompose less slowly, but nevertheless this may be a problem when they are used as infusions over long periods of time. This problem can usually be overcome by a study of the mechanism of decomposition, followed by appropriate prodrug design to increase stability. Azacytidine, for example, is used in the treatment of acute myeloid leukaemia. When given by an intravenous bolus it causes severe and dose-limiting gastrointestinal toxicity. However, if the drug is slowly infused over a 5-day period this toxicity is eliminated, but such infusions are not really practicable because of the instability of azacytidine in aqueous solutions. Azacytidine is hydrolysed reversibly to the ring-opened formyl derivative, which is then converted slowly and irreversibly to the guanylurea (Fig. 1). While both the formyl and the urea derivatives have lower toxicity, they do not have antitumour activity so that over the long infusion period a gradually reducing concentration of azacytidine is being infused,




Fig. 1

which i  
mechan  
bond, f  
across t  
reverts  
azacyti  
ing mo  
more, l  
oral ad  
Similar  
whose  
arabino  
employ  
mg/ml,  
litres o  
it is co  
deamin





Fig. 2

Som  
ed clin  
12/17  
has on  
likely  
clinical  
drugs s  
terest  
several  
conditi  
Wate  
associa  
sensiti

hydrophilic nitroimidazoles concentrate less in neural tissues and are excreted more rapidly, both properties reducing neurotoxicity. These nitroimidazoles may also be suitable for oral administration, but since decreased lipophilicity will decrease oral absorption a prodrug may be required to obtain optimal activity [9]. A nitroimidazole has been synthesised which has a very low octanol/water partition coefficient and might be expected to be less neurotoxic than earlier radiation sensitisers such as misonidazole, which has dose-limiting neurotoxicity. In order to increase absorption from the gut the prodrug acetyl ester was synthesised and shown to be converted completely to the drug in the first pass through the liver (Fig. 3).

In some cases it is necessary to reduce the water solubility of a drug by the synthesis of a more fat-soluble prodrug. Anticancer agents such as 6-mercaptopurine are also useful in the treatment of psoriasis, but, because they do not penetrate the skin, must be given systemically, with resulting toxicity. A soft alkylated derivative of 6-mercaptopurine may be used as a topical treatment of psoriasis [10] since it is transported effectively through epidermal barriers and is then converted, firstly by esterases and then spontaneously to the active thiopurine (Fig. 4).

### 3. Prodrugs which overcome acquired resistance

Often if the mechanism by which a cell acquires resistance to a drug is understood it is possible to design a prodrug to overcome this resistance. Thus, 6-mercaptopurine, like many antipurines and antypyrimidines, requires intracellular conversion to a nucleotide before it exerts its inhibitory action. 6-Mercaptopurine is converted by hypoxanthine, guanine phosphoribosyl transferase (HGPRT) to the nucleoside monophosphate (Fig. 5), which interferes with cell growth by a number of




Fig. 3




Fig. 4




Fig. 5

mechanism... become... HGPR... 6-merc... degrad... 3',5'-p... 6-merc... cells an... acylase




Fig. 6

In th... tion. S... transp... might... phenyl... [13], a... melph

#### 4. Prodrugs with improved pharmacokinetic properties

Most anticancer drugs are cell cycle-specific, meaning that they are particularly toxic to cells in cycle. Some drugs are even more specific, being cytotoxic to cells only in a particular phase of the cell cycle. For these agents the correct pharmacokinetic profile is essential if selective inhibition of tumour cells is to be obtained. Cytosine arabinoside (Ara-C), for example, acts after its intracellular conversion to the triphosphate, in which form it is an inhibitor of DNA polymerase. Therefore, it is S phase-specific and to achieve its optimal anti-tumour effect it must be present in the tumour environment at a toxic concentration for the period of time that it takes all the cells of the asynchronously growing tumour to enter and pass through the S phase. However, Ara-C as well as being activated to its triphosphate is also a substrate for cytidine deaminase and is converted by this enzyme to an inactive product (Fig. 7). This would result in a rapid clearance of Ara-C from the body and its profile after intraperitoneal or intravenous injection would be as indicated diagrammatically in Figure 8. After either route of administration the phar-




Fig. 7




Fig. 8

maco  
inject  
Any c  
fects  
sion o  
patient  
drugs  
pharm  
An  
9), wh  
tive c  
substi  
cytidin  
lipoph  
increa  
has b  
Howe  
prodr

R O CH  
K

Fig. 9

Ad  
deam  
C, m  
and i  
ticula  
poter  
Ara-  
arabi  
of Ar  
A s






Fig. 10

from steroids such as cortisone, prednisolone and a range of anti-inflammatory steroids [24]. Some but not all were superior to Ara-C alone in tumour-bearing animals, and it is of interest that prodrug activation appeared to take place inside the cells rather than in the blood. Clearly, this could form the basis for the design of prodrugs with activity against cells, with acquired resistance to Ara-C as a result of loss of the activating enzyme. Some prodrugs of Ara-C, such as Ara-CTP-L-dipalmitine, are not substrates for cytidine deaminase and are very effective against tumour cells in culture [25]. They might be readily taken up by cells and then degraded to Ara-CTP. In such a situation cells resistant to Ara-C because of loss of the kinase would remain sensitive to this type of prodrug. In fact, diacylglycerol derivatives of cytosine arabinoside do have an effect against some mouse tumours which have acquired resistance to Ara-C [26]. Other analogues of Ara-C which have been studied include phospholipids and polyglutamates and an  $N^4$ -palmitoyl derivative which was effective when given orally to tumour-bearing mice [25–30].

Following similar lines esters, glutamyl conjugates, amides, hydrazides and hydroxamic derivatives have been made of methotrexate [31–33]. Although in many cases the rationale has been to produce slow-release prodrugs forms of methotrexate, others have been made which may be taken up readily by cells by passive diffusion, and thus be active in cells resistant to methotrexate in which the active transport mechanism has been lost. In the course of these studies some interesting findings have been made. Most importantly, diesters of methotrexate may be cytotoxic to leukaemia cells in their own right, an effect which is only partially reversed by folinic acid, indicating that they may be acting by more than one pathway. Furthermore, in mouse serum there is a regioselective hydrolysis of the diester, the  $\alpha$ -carboxyl ester being hydrolysed much more readily than the  $\gamma$  [31]. Studies on esterases as prodrugs have emphasised that the rodent may not be a satisfactory model for humans since serum esterases in the former are much more active than in humans.

$N,N$ -Dibenzyl-daunorubicin has no activity against tumour cells in vitro but is active in vivo, suggesting that in whole animals it acts as a prodrug, with possible dif-

ferenc  
The  
it sele  
site o  
dinitr

5. S

Many  
which  
first e  
which  
[36] (

NaOPO<sub>2</sub>

Fig. 11

The  
alter  
(5-FU  
tumo  
to its  
B lym  
do no  
its n  
phosp  
as the  
The  
best i  
titum  
cytoto  
metab




Fig. 12

and cytotoxicity. Highly reactive nitrogen mustards are likely to be very toxic (both to tumour cells and to normal tissues) and poorly reactive ones very much less toxic. Figure 13 shows that as the alkylating activity varies over a 25-fold range, the  $LD_{50}$  varies from 40 to 3000  $\mu\text{mol}/\text{kg}$ . Very small changes in structure, e.g., *p*-hydroxylation, can greatly alter reactivity, and hence toxicity [40]. On this basis prodrugs can be designed which are poor alkylating agents and non-toxic, but which act as substrates for enzymes which transform them to highly reactive and cytotoxic agents. Clearly, for there to be selective tumour cell kill, the prodrug must be a good substrate for the enzyme under physiological conditions and the enzyme ideally should be present uniquely in the tumour cells, or at least at much higher levels in the tumour than in normal tissues, especially those which are also sensitive to this type of agent. A further prerequisite is that once formed from the prodrug, the drug

Compound	Alkylating activity ( $K_{10} \times 10^3$ )	Toxicity ( $LD_{50}$ , $\mu\text{mole}/\text{kg}$ )
	4.9	3000
	4.6	915
	13.0	367
	48.6	74
	123.0	39

Fig. 13

should  
alkylat  
that th  
would  
that it  
produ  
Follo  
reduct  
designe  
which  
agent s  
react in




Fig. 14

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.