

Journal of Pharmaceutical Sciences

JANUARY 1977

VOLUME 66 NUMBER 1



REVIEW ARTICLE

Pharmaceutical Salts

STEPHEN M. BERGE *†, LYLE D. BIGHLEY *, and
DONALD C. MONKHOUSE *

Keyphrases □ Pharmaceutical salts—general pharmacy, physicochemical properties, bioavailability, pharmaceutical properties, toxicology, review □ Salts, pharmaceutical—general pharmacy, physicochemical properties, bioavailability, pharmaceutical properties, toxicology, review □ Physicochemical properties—dissolution, solubility, stability, and organoleptic properties of pharmaceutical salts, review □ Bioavailability—formulation effects, absorption alteration and pharmacokinetics of pharmaceutical salts, review □ Toxicology—pharmaceutical salts, review

CONTENTS

Potentially Useful Salts	2
Physicochemical Studies	4
Dissolution Rate	5
Solubility	7
Organoleptic Properties	8
Stability	9
Miscellaneous Properties	10
Bioavailability	10
Formulation Effects	11
Absorption Alteration	11
Pharmacokinetics	13
General Pharmacy	14
Pharmacological Effect	14
Dialysis	14
Miscellaneous	14
Toxicological Considerations	15
Toxicity of Salt Ion	15
Toxicity of Salt Form	15
Conclusions	16
References	16

The chemical, biological, physical, and economic characteristics of medicinal agents can be manipulated and, hence, often optimized by conversion to a salt form. Choosing the appropriate salt, however, can be a very difficult task, since each salt imparts unique properties to the parent compound.

Salt-forming agents are often chosen empirically. Of the many salts synthesized, the preferred form is selected by pharmaceutical chemists primarily on a practical basis: cost of raw materials, ease of crystallization, and percent yield. Other basic considerations include stability, hygroscopicity, and flowability of the resulting bulk drug. Unfortunately, there is no reliable way of predicting the influence of a particular salt species on the behavior of the parent compound. Furthermore, even after many salts of the same basic agent have been prepared, no efficient screening techniques exist to facilitate selection of the salt most likely to exhibit the desired pharmacokinetic, solubility, and formulation profiles.

Some decision-making models have, however, been developed to help predict salt performance. For example, Walkling and Appino (1) described two techniques, "decision analysis" and "potential problem analysis," and applied them to the selection of the most suitable derivative of an organic acid for development as a tablet. The derivatives considered were the free acid and the potassium, sodium, and calcium salts. Both techniques are based on the chemical, physical, and biological properties of these specific derivatives and offer a promising avenue for developing optimal salt forms.

Information on salts is widely dispersed throughout the pharmaceutical literature, much of which addresses the use of salt formation to prolong the release of the active component, thereby eliminating various undesirable drug properties (2–6). This review surveys literature of the last 25 years, emphasizing comparisons between the properties of different salt forms of the same compound. Included also is a discussion of potentially useful salt forms. Our purpose is twofold: to present an overview of the many different salts from which new drug candidates can be chosen and

Table I—FDA-Approved Commercially Marketed Salts

Anion	Percent ^a	Anion	Percent ^a
Acetate	1.26	Iodide	2.02
Benzenesulfonate	0.25	Isethionate ⁱ	0.88
Benzoate	0.51	Lactate	0.76
Bicarbonate	0.13	Lactobionate	0.13
Bitartrate	0.63	Malate	0.13
Bromide	4.68	Maleate	3.03
Calcium edetate	0.25	Mandelate	0.38
Camsylate ^h	0.25	Mesylate	2.02
Carbonate	0.38	Methylbromide	0.76
Chloride	4.17	Methylnitrate	0.38
Citrate	3.03	Methylsulfate	0.88
Dihydrochloride	0.51	Mucate	0.13
Eddate	0.25	Napsylate	0.25
Edisylate ^j	0.38	Nitrate	0.64
Estolate ^d	0.13	Pamoate (Embonate)	1.01
Esylate ^v	0.13	Pantothenate	0.25
Fumarate	0.25	Phosphate/diphosphate	3.16
Gluceptate ^f	0.18	Polygalacturonate	0.13
Gluconate	0.51	Salicylate	0.88
Glutamate	0.25	Stearate	0.25
Glycolylarsanilate ^g	0.13	Subacetate	0.38
Hexylresorcinate	0.13	Succinate	0.38
Hydrabamine ^h	0.25	Sulfate	7.46
Hydrobromide	1.90	Tannate	0.88
Hydrochloride	42.98	Tartrate	3.54
Hydroxynaphthoate	0.25	Teoclate ^j	0.13
		Triethiodide	0.13
Cation	Percent ^a	Cation	Percent ^a
Organic:		Metallic:	
Benzathine ^k	0.66	Aluminum	0.66
Chlorprocaine	0.33	Calcium	10.49
Choline	0.33	Lithium	1.64
Diethanolamine	0.98	Magnesium	1.31
Ethylenediamine	0.66	Potassium	10.82
Meglumine ^l	2.29	Sodium	61.97
Procaine	0.66	Zinc	2.95

^a Percent is based on total number of anionic or cationic salts in use through 1974. ^b Camphorsulfonate. ^c 1,2-Ethanedisulfonate. ^d Lauryl sulfate. ^e Ethanesulfonate. ^f Glucoheptonate. ^g p-Glycollamidophenylarsonate. ^h N,N'-Di(dehydroabietyl)ethylenediamine. ⁱ 2-Hydroxyethanesulfonate. ^j 8-Chlorotheophyllinate. ^k N,N'-Dibenzylethylenediamine. ^l N-Methylglucamine.

to assemble data that will provide, for the student and practitioner alike, a rational basis for selecting a suitable salt form.

POTENTIALLY USEFUL SALTS

Salt formation is an acid-base reaction involving either a proton-transfer or neutralization reaction and is therefore controlled by factors influencing such reactions. Theoretically, every compound that exhibits acid or base characteristics can participate in salt formation. Particularly important is the relative strength of the acid or base—the acidity and basicity constants of the chemical species involved. These factors determine whether or not formation occurs and are a measure of the stability of the resulting salt.

The number of salt forms available to a chemist is large; surveys of patent literature show numerous new salts being synthesized annually. Various salts of the same compound often behave quite differently because of the physical, chemical, and thermodynamic properties they impart to the parent compound. For example, a salt's hydrophobicity and high crystal lattice energy can affect dissolution rate and, hence, bioavailability. Ideally, it would be desirable if one could predict how a pharmaceutical agent's properties would be affected by salt formation.

Tables I and II list all salts that were commercially marketed through 1974. The list was compiled from all agents listed in "Martindale The Extra Pharmacopoeia."

**Table II—
Salts**

Adipate
Alginate
Aminosalicylate
Anhydromercaptopropionate
Arecoline
Aspartate
Bisulfate
Butylbromide
Camphorate
Digluconate
Dihydrobromide
Disuccinate
Glycerophosphate
Hemisulfate
Hydrofluoride
Hydroiodide
Methylenebisacrylamide
Napadisylate
Oxalate
Pectinate
Persulfate
Phenylethylsulfate
Picrate
Propionate
Thiocyanate
Tosylate
Undecanoate

^a Percent is based on total number of anionic or cationic salts in use through 1974.

^d 1-p-Chlorodioxymethyl

using pamoate (Embonate) action preventatives are dispersants.

Alginic acid is a pharmaceutical polymer and shows some promise in the treatment of certain diseases. A study has shown that of pilocarpine hydrochloride dried to a powder, the preparation is more effective than the liquid. While liquid pilocarpine hydrochloride salts are more effective than solid ones, the solid form is more effective. More significantly, the solid form is more soluble in water. Solid dose forms are more effective than liquid forms because it is easier to disperse the drug which holds the drug in a more immediate release form.

Málek et al. (10) have shown that the absorption of ibuprofen from a sustained-release preparation is increased by merely adding a small amount of the drug molecules to the aqueous solution.

26th ed. (7). Table I categorizes all salt forms approved by the Food and Drug Administration (FDA), while Table II lists those not approved by the FDA but in use in other countries. (Only salts of organic compounds are considered because most drugs are organic substances.) The relative frequency with which each salt type has been used is calculated as a percentage, based on the total number of anionic or cationic salts in use through 1974. Because of simple availability and physiological reasons, the monoprotic hydrochlorides have been by far the most frequent choice of the available anionic salt-forming radicals, outnumbering the sulfates nearly six to one. For similar reasons, sodium has been the most predominant cation.

Knowledge that one salt form imparts greater water solubility, is less toxic, or slows dissolution rate would greatly benefit chemists and formulators. In some cases, such generalizations can be made. Miller and Heller (8) discussed some properties associated with specific classes of salt forms. They stated that, in general, salt combinations with monocarboxylic acids are insoluble in water and lend themselves to repository preparations, while those of dicarboxylic acids confer water solubility if one carboxylic group is left free. Pamoic acid, an aromatic dicarboxylic acid, is an exception since it is used as a means of obtaining prolonged action by forming slightly soluble salts with certain basic drugs. Saisas et al. (9) reviewed the use of this salt form in preparing sustained-release preparations. More recently, latentiation of dihydrostreptomycin (10)

Table I—FDA-Approved Commercially Marketed Salts

Anion	Percent ^a	Anion	Percent ^a
Acetate	1.26	Iodide	2.02
Benzenesulfonate	0.25	Isethionate ^b	0.88
Benzoate	0.51	Lactate	0.76
Bicarbonate	0.13	Lactobionate	0.13
Bitartrate	0.63	Malate	0.13
Bromide	4.68	Maleate	3.03
Calcium edetate	0.25	Mandelate	0.38
Camsylate ^b	0.25	Mesylate	2.02
Carbonate	0.38	Methylbromide	0.76
Chloride	4.17	Methylnitrate	0.38
Citrate	3.03	Methylsulfate	0.88
Dihydrochloride	0.51	Mucate	0.13
Edetate	0.25	Napsylate	0.25
Edisylate ^c	0.38	Nitrate	0.64
Estolate ^d	0.13	Pamoate (Embonate)	1.01
Esylate ^e	0.13	Pantothenate	0.25
Fumarate	0.25	Phosphate/diphosphate	3.16
Gluceptate ^f	0.18	Polygalacturonate	0.13
Gluconate	0.51	Salicylate	0.88
Glutamate	0.25	Stearate	0.25
Glycolylarsanilate ^g	0.13	Subacetate	0.38
Hexylresorinate	0.13	Succinate	0.38
Hydrabamine ^h	0.25	Sulfate	7.46
Hydrobromide	1.90	Tannate	0.88
Hydrochloride	42.98	Tartrate	3.54
Hydroxynaphthoate	0.25	Teoclate ^j	0.13
		Triethiodide	0.13
Cation	Percent ^a	Cation	Percent ^a
Organic:		Metallic:	
Benzathine ^k	0.66	Aluminum	0.66
Chlorprocaine	0.33	Calcium	10.49
Choline	0.33	Lithium	1.64
Diethanolamine	0.98	Magnesium	1.31
Ethylenediamine	0.66	Potassium	10.82
Meglumine ^l	2.29	Sodium	61.97
Procaine	0.66	Zinc	2.95

^a Percent is based on total number of anionic or cationic salts in use through 1974. ^b Camphorsulfonate. ^c 1,2-Ethanedisulfonate. ^d Lauryl sulfate.^e Ethanesulfonate. ^f Glucoheptonate. ^g p-Glycollamidophenylarsonate. ^h N,N'-Di(dehydroabietyl)ethylenediamine. ⁱ 2-Hydroxyethanesulfonate.^j 8-Chlorotheophyllinate. ^k N,N'-Dibenzylethylenediamine. ^l N-Methylglucamine.

to assemble data that will provide, for the student and practitioner alike, a rational basis for selecting a suitable salt form.

POTENTIALLY USEFUL SALTS

Salt formation is an acid-base reaction involving either a proton-transfer or neutralization reaction and is therefore controlled by factors influencing such reactions. Theoretically, every compound that exhibits acid or base characteristics can participate in salt formation. Particularly important is the relative strength of the acid or base—the acidity and basicity constants of the chemical species involved. These factors determine whether or not formation occurs and are a measure of the stability of the resulting salt.

The number of salt forms available to a chemist is large; surveys of patent literature show numerous new salts being synthesized annually. Various salts of the same compound often behave quite differently because of the physical, chemical, and thermodynamic properties they impart to the parent compound. For example, a salt's hydrophobicity and high crystal lattice energy can affect dissolution rate and, hence, bioavailability. Ideally, it would be desirable if one could predict how a pharmaceutical agent's properties would be affected by salt formation.

Tables I and II list all salts that were commercially marketed through 1974. The list was compiled from all

26th ed. (7). Table I categorizes all salt forms approved by the Food and Drug Administration (FDA), while Table II lists those not approved by the FDA but in use in other countries. (Only salts of organic compounds are considered because most drugs are organic substances.) The relative frequency with which each salt type has been used is calculated as a percent, based on the total number of anionic or cationic salts in use through 1974. Because of simple availability and physiological reasons, the monoprotic hydrochlorides have been by far the most frequent choice of the available anionic salt-forming radicals, outnumbering the sulfates nearly six to one. For similar reasons, sodium has been the most predominant cation.

Knowledge that one salt form imparts greater water solubility, is less toxic, or slows dissolution rate would greatly benefit chemists and formulators. In some cases, such generalizations can be made. Miller and Heller (8) discussed some properties associated with specific classes of salt forms. They stated that, in general, salt combinations with monocarboxylic acids are insoluble in water and lend themselves to repository preparations, while those of dicarboxylic acids confer water solubility if one carboxylic group is left free. Pamoic acid, an aromatic dicarboxylic acid, is an exception since it is used as a means of obtaining prolonged action by forming slightly soluble salts with certain basic drugs. Sajas *et al.* (9) reviewed the use of this salt form in preparing sustained-release preparations.

Table II—
Salts

Adipate
Alginate
Aminosalicylate
Anhydrometh-
Arecoline
Aspartate
Bisulfate
Butylbromide
Camphorate
Digluconate
Dihydrobrom-
Disuccinate
Glycerophosp-
Hemisulfate
Hydrofluorid-
Hydroiodide
Methylenebis-
Napadisylate
Oxalate
Pectinate
Persulfate
Phenylethylb-
Picrate
Propionate
Thiocyanate
Tosylate
Undecanoate

Cation

Organic:
Benethamini-
Clemizole ^d
Diethylam-
Piperazine
Tromethar-

^a Percent is through 1974.
^d 1-p-Chloro-
droxymethyl-

using pamoic acid as an example, the action preparation is dispersible.

Alginic acid is a polymer found in many pharmaceuticals and shown to be dispersible in aqueous solutions. A strong gel is formed that of pilocarpine hydrochloride is dried to a solid form and prepared for injection. While liquid alginic acid salts are dispersible, solid pamoic acid is more effective than the aqueous form. Solid dose forms are dispersible because it is a polymer which holds the drug in immediate release.

Málek *et al.* (10) studied the action of pamoic acid in the distribution of the drug in the body by merely dispersing the molecules.

Table II—Non-FDA-Approved Commercially Marketed Salts

Percent ^a	Anion	Percent ^a
2.02		
0.88	Adipate	0.13
0.76	Alginate	0.13
0.13	Aminosalicylate	0.25
0.13	Anhydromethylenecitrate	0.13
3.03	Arecoline	0.13
0.38	Aspartate	0.25
2.02	Bisulfate	0.25
0.76	Butylbromide	0.13
0.38	Camphorate	0.13
0.88	Digluconate	0.13
0.13	Dihydrobromide	0.13
0.25	Disuccinate	0.13
0.64	Glycerophosphate	0.88
1.01	Hemisulfate	0.13
0.25	Hydrofluoride	0.13
3.16	Hydroiodide	0.25
0.13	Methylenebis(salicylate)	0.13
0.88	Napadisylate ^b	0.13
0.25	Oxalate	0.25
0.38	Pectinate	0.13
7.46	Persulfate	0.13
0.88	Phenylethylbarbiturate	0.13
3.54	Picrate	0.13
0.13	Propionate	0.13
0.13	Thiocyanate	0.13
	Tosylate	0.13
	Undecanoate	0.13
Percent ^a	Cation	Percent ^a
0.66		
10.49	Organic:	
1.64	Benethamine ^c	0.33
1.31	Clemizole ^d	0.33
10.82	Diethylamine	0.33
61.97	Piperazine	0.98
2.95	Tromethamine ^e	0.33
	Metallic:	
	Barium	0.33
	Bismuth	0.98

^a Percent is based on total number of anionic and cationic salts in use through 1974. ^b 1,5-Naphthalenedisulfonate. ^c N-Benzylphenethylamine. ^d 1-p-Chlorobenzyl-2-pyrrolidin-1'-ylmethylbenzimidazole. ^e Tris(hydroxymethyl)aminomethane.

using pamoic acid resulted in the formation of a delayed-action preparation. Numerous studies using pamoate salts are dispersed throughout the literature (11–15).

Alginic acid also has been used to prepare long-acting pharmaceuticals. Streptomycin alginate was prepared (16) and shown to be effective in sustained-release preparations. A striking example of a long-acting alginate salt is that of pilocarpine. When dispersed in sterile water and dried to a solid gel, this compound was found useful in the preparation of long-acting ophthalmic dosage forms (17). While liquid preparations of the alginate and hydrochloride salts possess similar miotic activity, studies showed that solid pilocarpine alginate flakes constricted pupil size more effectively and increased the duration of miosis significantly when compared with the liquid preparations. Solid dose pilocarpine may be more uniformly available, because it diffuses more slowly through the gel matrix which holds the drug in reserve. In contrast, drops of the commonly employed solution dosage form release the dose immediately to the conjunctival fluid.

Málek *et al.* (18) devised a unique way of prolonging action through salt formation; they showed that the distribution of several antibiotics could be markedly altered by merely preparing macromolecular salts. Since macromolecules and colloidal particles have an affinity for the lymphatic system, streptomycin, neomycin, viomycin, and

streptothrycin were combined with high molecular weight compounds such as polyacrylic acids, sulfonic or phosphorylated polysaccharides, and polyuronic derivatives. Parenteral administration of these compounds produced low blood levels of the antibiotic for long periods, while lymph levels were high. (In comparison, streptomycin sulfate gave high blood levels but low lymph levels.) This alteration in distribution caused the streptomycin to prolong its passage through the body, since lymphatic circulation is quite slow.

The appropriate choice of a salt form has been found to reduce toxicity. It can be rationalized that any compound associated with the normal metabolism of food and drink must be essentially nontoxic. The approach of choosing organic radicals that are readily excreted or metabolized opened up a new class of substances from which to select a salt form. For example, certain salts of the strong base choline have proven to be considerably less toxic than their parent compound. The preparation and properties of choline salts of a series of theophylline derivatives were reported (19), and it was shown that choline theophyllinate possessed a greater LD₅₀ than theophylline or its other salts (20). It was postulated that this agent would be less irritating to the GI tract than aminophylline, because "its basic constituent, choline, is an almost completely non-toxic substance of actual importance to the physiologic economy." This evidence led to the preparation of choline salicylate (21) as an attempt to reduce the GI disturbances associated with salicylate administration. Clinical studies indicated that choline salicylate elicited a lower incidence of GI distress, was tolerated in higher doses, and was of greater benefit to the patient than was acetylsalicylic acid (aspirin).

Amino acids and acid vitamins also have been used as salt-forming agents. Based on the evidence that coadministration of amino acids with aminoglycoside antibiotics reduced their toxicity, a series of amino acid salts of dihydrostreptomycin was prepared (22). In all but one case, the acute toxicities of these salts were lower than the toxicity of the sulfate. The ascorbate and pantothenate also were synthesized and shown to be less toxic than the sulfate. Of the salts prepared, the ascorbate had the highest LD₅₀.

The vitamins most commonly used for forming salts exhibiting reduced toxicity are ascorbic and pantothenic acids. Keller *et al.* (23) were the first to use pantothenic acid as a means of "detoxifying" the basic streptomycetes antibiotics. Parenteral administration of the pantothenates of streptomycin and dihydrostreptomycin had a significantly reduced incidence of acute neurotoxicity in cats as compared with the sulfates. Subsequent studies (24–28) supported this finding and showed that the pantothenates of neomycin and viomycin also are less toxic. The ascorbate of oleandomycin was synthesized and its pharmacological properties were reported (29). Upon intramuscular injection in rats, it produced less irritation than the phosphate.

p-Acetamidobenzoic acid, an innocuous metabolite of folic acid present in normal blood and urine, has been used in preparing salts. In particular, it yields stable salts with amines that otherwise tend to form hygroscopic products with conventional acid components (30).

Often the salt form is chosen by determining a salt

Table III—

Salt

Acetylaminoo
N-Acetyl-L-a
N-Acetylcyst
Adamantoic a
Adipic acid
N-Alkylsulfur

Anthraquinon
Arabogalacta
Arginine

Aspartate
Betaine
Bis(2-carboxy
Carnitine
4-Chloro-m-t
Decanoate
Diacyl sulfat
Dibenzylethyl
Diethylamine
Diguaicayl ph
Diocetyl sulfos
Embonic (pan

Fructose 1,6-d

Glucose 1-ph
6-phosphor
L-Glutamine
Hydroxynaph
2-(4-Imidazol
Isobutanolam
Lauryl sulfate
Lysine

Methanesulfo
N-Methylgluc

N-Methylpipe
Morpholine
2-Naphthalen
Octanoate
Probenecid
Tannic acid
Theobromine
3,4,5-Trimetho

Tromethamine

tion, the rel
ministration

In a review
“different sa
logically; the
properties.”
panded upon
nature of the
salts of the s
ciably, the in

The salt f
cochemical p
dissolution r
These pro
mulation ch
pharmaceuti
extensive pr
properties of
suitable form
concerning s
tion studies
salt form on

component that will pharmacologically antagonize an unfavorable property or properties exhibited by the basic agent. Salts of *N*-cyclohexylsulfamic acid are an example of the practical application of this approach. *N*-Cyclohexylsulfamic acid salts, better known as cyclamates, have a characteristic sweet, pleasing taste. Although presently under investigation by the FDA for potentially carcinogenic properties, salts incorporating this compound can render unpleasant or bitter-tasting drugs acceptable. For example, the cyclamates of dextromethorphan and chlorpheniramine exhibit greatly improved bitterness thresholds compared to commonly occurring salts (31). Furthermore, their stability in aqueous solution was described as good when maintained at a pH not greater than 4.

N-Cyclohexylsulfamic acid salts of thiamine hydrochloride and lincomycin also have been synthesized. Thiamine *N*-cyclohexylsulfamate hydrochloride was reported to have a more pleasant taste than other thiamine salts while having an equal or greater stability (32). Lincomycin cyclamate, shown to possess an enhanced thermal stability over its hydrochloride, was prepared (33) to test the hypothesis that reduced lincomycin absorption in the presence of small quantities of cyclamates was due to a simple metathetic reaction. However, this assumption was found not to be true. An extensive study of the preparation and characterization of cyclamic acid salts of several widely used classes of drugs including antihistamines, antibiotics, antitussives, myospasmolytics, and local anesthetics was reported (34, 35).

Various salts of penicillin and basic amine compounds have been formulated in an effort to produce a long-acting, nonallergenic form of penicillin. Since antihistamines appear to mitigate the symptomatology of penicillin reactions in some patients, coadministration of the two has been advocated. The preparation of the benzhydralamine salt of penicillin was an attempt to produce a repository form of penicillin with antiallergic properties (36). Blood levels achieved with this salt were comparable to those of penicillin G potassium; however, its antiallergic properties were not evaluated. In fact, the investigators noted that antihistamines can actually cause sensitization at times and stated that “despite their occasionally favorable influence on the symptoms of penicillin sensitivity, they contribute directly to the potential of drug sensitivity when co-administered with penicillin.”

Silver salts of sulfanilamide, penicillin, and other antibiotics have been prepared and represent cases where the species (ions) are complementary. When aqueous solutions of the salts were applied topically to burned tissue, they yielded the combined benefits of the oligodynamic action of silver and the advantages of the antibacterial agents (37).

The use of 8-substituted xanthines, particularly the 8-substituted theophyllines, as salt-forming agents was first reported in the preparation of a series of antihistamine salts (38–41). Synthesis of these xanthine salts was an attempt to find a drug to counteract the drowsiness caused by the antihistamines with the stimulant properties of the xanthines. When an electronegative group is introduced into the xanthine molecule at the 8-position, the electron-drawing capacity of the substituent results in the

moderately strong acidic compounds can undergo salt formation with various organic bases.

The 8-halotheophyllines were the first group of xanthines studied as potential salt-forming agents. Since the report on the preparation of the 8-chlorotheophylline salt of diphenhydramine (42), synthesis of the 8-halotheophyllines of a number of organic bases has been attempted. The 8-chlorotheophylline salts of quinine, ephedrine, and strychnine were prepared and characterized (43). These salts were less water soluble than the corresponding free alkaloidal bases. In a similar report, the 8-chlorotheophyllines of three synthetic narcotics, meperidine, levorphanol, and metopon, were prepared (44).

Pharmacological and clinical studies involving the 8-bromotheophylline pyrilamine salt revealed the unusual diuretic properties associated with the 8-halotheophylline portion of the compound (45, 46). This finding initiated an investigation into the preparation of a soluble 8-bromotheophylline salt of high diuretic activity. With readily available amines, over 30 salts were synthesized and screened for diuretic activity (47). When tested against theophylline salts of the same amines, the 8-bromotheophyllines showed greater activity in every case.

With the successful formation of 8-halotheophyllines of organic bases, Morozowich and Bope (48) proposed that, if the halogen moiety was replaced with a more electronegative substituent such as a nitro group, a more acidic compound would be formed. Presumably, more stable salts would result and precipitation of the free xanthine derivative in the stomach would be less likely to occur. On this premise, they successfully prepared pharmacologically effective 8-nitrotheophyllines of several pharmaceutically useful bases.

Duesel *et al.* (19), in their study of choline theophyllinate, prepared the 8-chloro-, 8-bromo-, and 8-nitrotheophylline salts of choline. Oral toxicity studies in mice showed that the LD₅₀ of the 8-nitrotheophyllinate was much greater than that of either 8-halotheophylline. In fact, it remained nonlethal at doses as high as 5 g.

Polygalacturonic acid, a derivative of pectin, has been used to prepare quinidine salts exhibiting reduced toxicity (49, 50). The compound possesses special demulcent properties and inhibits mucosal irritation. The rationale for use of this agent is to reduce the ionic shock to the GI mucosa resulting from the flood of irritating ions liberated by rapid dissociation of the conventional inorganic quinidine salts. Studies have shown that it is four times less toxic orally than the sulfate. This difference was attributed to the slower release of quinidine from the polygalacturonate.

Other compounds reported to be potentially useful as pharmaceutical salt forms are listed in Table III.

PHYSICOCHEMICAL STUDIES

Biological activity of a drug molecule is influenced by two factors: its chemical structure and effect at a specific site and its ability to reach—and then be removed from—the site of action. Thus, a knowledge of the physicochemical properties of a compound that influence its absorption, distribution, metabolism, and excretion is essential for a

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.