

JIA is the most common cause of chronic arthritis in children and affects children worldwide, with no racial predilection.

Approximately 1/1,000 children, or more than 300,000 children in the U.S. have JIA. Although children of all ages may get JIA, it is rare before six months of age.

Arthritis may affect any joint, even your jaw and spine.

JIA may be categorized into several subtypes, some of which may be associated with other diseases, which may affect the eyes or gastrointestinal tract, for example.

Although there are several sub-types of JIA, all patients are affected by chronic arthritis.

The severity of arthritis is variable, and patients may have an unpredictable flare (worsening) of their arthritis. Sometimes this flare may relate to the weather, viral infections or inactivity.

Learn More

- How is JIA diagnosed?
- What causes JIA?
- Types of JIA
- · How is JIA treated?

Ethics • For Faculty • For Staff • Disclaimer • Privacy Policy • Contact Webmaster

Call us for more information:

http://pediatrics.med.nyu.edu/rheumatology/patient-care/juvenile-idiopathic-arthritis-jia[11/18/14 1:00:28 PM]

DOCKE

RM

Medac Exhibit 2078

Find authenticated court documents without watermarks at docketalarm.com.



© 2013 Division of Pediatric Rheumatology | 550 First Avenue, New York, NY 10016

DOCKET A L A R M Find authenticated court documents without watermarks at <u>docketalarm.com</u>.