



Published in final edited form as:

Prostate. 2008 October 1; 68(14): 1546–1554. doi:10.1002/pros.20814.

Nonhuman primates as models for studies of prostate specific antigen and prostatic diseases

James N. Mubiru¹, Gene B. Hubbard¹, Edward J. Dick Jr.¹, Jaime Furman³, Dean A. Troyer³, and Jeffrey Rogers^{1,2}

¹Southwest National Primate Research Center, Southwest Foundation for Biomedical Research, San Antonio, Texas

²Department of Genetics, Southwest Foundation for Biomedical Research, San Antonio, Texas

³Department of Pathology, University of Texas Health Science Center at San Antonio, Texas

Abstract

Background—Because prostate specific antigen (PSA) is released at increased levels into the blood early in the development of prostate cancer, benign prostatic hyperplasia (BPH) and prostatitis, it is widely used as a marker for these diseases. However, PSA has clinical limitations as a screen for prostatic diseases due to its low sensitivity and specificity. There is a strong need to better understand the biology of PSA and factors affecting its serum levels.

Methods—We evaluated cynomolgus macaques, rhesus macaques, baboons, and marmosets for their suitability as models for the study of PSA biology and prostatic diseases.

Results—Prostates of several nonhuman primates are anatomically similar to the human counterpart. Anti-human PSA antibody detected PSA antigens in all the Old World monkeys (cynomolgus macaques, rhesus macaques and baboons) but not in marmosets. Of the Old World monkeys, cynomolgus macaques have the highest serum PSA levels; baboons have the lowest. Serum PSA levels from macaques includes a number of outlier samples with unusually high values. We also report two cases of abnormal pathologies in macaques accompanied by high serum PSA levels. One case consisted of prostatic hyperplasia involving both glandular and basal cells in a cynomolgus macaque and another of glandular hyperplasia and atrophy in a rhesus macaque. The finding that pathological changes in the prostate of macaques may lead to increases in serum PSA is worthy of further exploration.

Conclusion—Cynomolgus macaques and rhesus macaques are promising animal models for PSA biology studies.

Keywords

PSA; nonhuman primates; prostate; macaque; baboon

Introduction

A major hindrance to the study of the pathogenesis of prostate diseases such as cancer, benign prostatic hyperplasia (BPH), and prostatitis is the lack of appropriate animal models.

Corresponding author: James N. Mubiru, Southwest National Primate Research Center, Southwest Foundation for Biomedical Research, P.O. Box 760549, San Antonio, TX 78245-0549, Telephone: (210) 258-9744, Fax: (210) 670-3323, E-mail: jmubiru@sfbf.org

In the case of prostate cancer, animal models for this condition are very limited due to the very few species (lions and dogs) that are known to develop prostate cancer spontaneously. However, the usefulness of these animal models is questionable due to either availability or the differences in anatomy and histology of the prostates of these animals compared to humans.

Although nonhuman primates are phylogenetically closer to humans than any other laboratory animals and therefore should be considered suitable models for human prostatic diseases, they have not been studied systematically as models of prostatic diseases mainly because of the widely held view that nonhuman primates do not develop prostatic diseases [1].

Presently, a number of proteins are used in the diagnosis, monitoring and management of prostatic diseases. In healthy males, prostate specific antigen (PSA) is restricted to the prostate gland and very little leaks into the bloodstream. Increased amounts are detected in the blood in cases of prostatic diseases and therefore PSA is used for screening, diagnosing and monitoring of prostate cancer. Alpha-methyl-acyl-CoA racemase (AMACR) is a mitochondrial and peroxisome enzyme that is overexpressed in prostate cancer and is presently used as an auxiliary immunohistochemical test for prostate cancer [2].

We evaluated four nonhuman primate species commonly used in research for their suitability as models of PSA biology and prostatic diseases. The study also examined AMACR, a protein that is clinically important in the diagnosis of prostate cancer. Two cases of prostatic lesions which were accompanied by elevated serum PSA are also reported.

Materials and Methods

Animals and Tissue Collection

Cynomolgus macaques (*Macaca fascicularis*), rhesus macaques (*Macaca mulatta*), baboons (*Papio hamadryas anubis*), and common marmosets (*Callithrix jacchus*) maintained at the Southwest National Primate Research Center, Southwest Foundation for Biomedical Research (SFBR) in San Antonio, Texas, were used for this study. A total of 114 baboons, 66 cynomolgus macaques, 12 rhesus macaques and 5 marmosets were examined.

We opportunistically sampled animals that presented for necropsy and an attempt was made to use animals across the whole lifespan. Male baboons reach sexual maturity between 6 and 7 years of age, male cynomolgus macaques and rhesus macaques between 4–5 years of age, and male marmosets reach sexual maturity at between 9–13 months.

Body weights were obtained either from SFBR animal records or when animals underwent necropsy. All animals were humanely euthanized. Only male animals were used in these studies. Complete necropsies were carried out for each animal. Male reproductive organs were removed and weighed. Tissue samples were either frozen at -80°C or fixed in neutral buffered formalin and processed for routine histological examination. All histopathological slides were examined by two board-certified veterinary pathologists (GBH and EJD). Two slides with microscopic lesions and elevated serum PSA were also sent to the Armed Forces Institute of Pathology (AFIP, Washington, DC) for evaluation.

Serum PSA Assay and Cloning

Serum was obtained from animals prior to euthanasia or from the archived serum bank maintained at SFBR. A radioimmunoassay with the lowest standard curve point of 0.01 ng/ml was used to assay PSA in serum (AniLytics, Inc., Gaithersburg, MD).

For cloning of the baboon PSA cDNA, total RNA was extracted using Trizol reagent (Invitrogen, Carlsbad, CA) and reverse-transcribed at 42°C with Moloney murine leukemia virus (MMLV) reverse transcriptase and random decamers following the suppliers protocol (RETROscript kit, Ambion, Austin, TX). The resulting cDNA was then used as a template for PCR reaction using nested primers and touchdown conditions [3]. The primer sequences were based on the published rhesus macaque sequence. The primer sequences used were, PSA outer forward: 5'CTG GAC AGC TGT GTC ACC AT3'; PSA inner forward: 5'CTG TGT CAC CAT GTG GGT TC3'; PSA outer reverse primer: 5'TGA GAT GTC TCC AGG CAT GA3'; PSA inner reverse primer: 5'AGG ACA CGG AGA GGA CAA AA3'. The PCR products were then cloned into the pCR2.1-TOPO vector (Invitrogen, Carlsbad, CA). Positive clones were identified by restriction enzyme analysis and nucleotide sequencing was carried out by the Advanced Nucleic Acid Core Facility at University of Texas Health Science Center at San Antonio. The PSA cDNA of both the rhesus macaque and the cynomolgus macaque were not cloned as they were already reported in the literature [4,5].

PSA Immunoblotting

Prostate tissue samples were homogenized in sample buffer, separated by 12% SDS-PAGE, transferred to nitrocellulose membranes, and probed using standard conditions. Membranes were probed at room temperature for 1 hour with either a monoclonal mouse anti-human PSA, clone ER-PR8 (Dakocytomation), or a rabbit monoclonal alpha-methyl-acyl-CoA racemase (AMACR, P504S Zeta Corporation, Sierra Madre, CA). Goat anti-mouse secondary antibody linked to horseradish peroxidase (Pierce) were added to the blots at a dilution of 1:5000 and incubated for 1 hour at room temperature. Bound antibodies were detected by chemiluminescence (Supersignal West Pico, Pierce Biotechnology, Rockford, IL).

Results

Prostate Anatomy and Body Weight

The prostate of the adult cynomolgus macaque, rhesus macaque, baboon, and marmoset is located at the distal end of the urinary bladder. On gross observation the prostate of cynomolgus macaques, rhesus macaques, and baboons is divided into two parts which are labeled cranial and caudal according to their proximity to the bladder and seminal vesicles. For the marmoset, although the prostate can be seen on gross examination, it cannot be divided into two lobes with a naked eye. However, histologically differences in acinar size and hematoxylin-eosin staining characteristics can be seen. On the basis of these differences the marmoset prostate can also be divided into two lobes which correspond with the other three species. In all these monkey species, the glandular prostate does not completely surround the urethra (Figure 1) and in this regard differs from what has been reported in humans [6,7].

There is a major difference in the size of the two lobes of the prostate between the baboon and the two macaque species (rhesus macaque and cynomolgus macaque). In the macaque species the two lobes of the prostate are of almost equal sizes. The cranial lobe constitutes 59% of weight of the prostate in the cynomolgus while it constitutes 47% in the rhesus macaque. In the case of the baboon the two lobes are quite different in size; the cranial lobe is much smaller, contributing only 36% of the total weight of the prostate (Figure 1). Since the marmoset prostate is quite small and the two lobes are not visible on gross examination, their relative proportions were not quantified.

The average adult body weight of the animals studied were 5.82, 10.06, 27.14, and 0.29 kg for cynomolgus macaques, rhesus macaques, baboons, and marmosets, respectively (Table 1).

When the weight of the prostate is compared to body weight (g/kg), the macaque species have a higher ratio than the baboon. The prostate-to-body-weight ratios are 0.51, 0.68, and 0.31 for cynomolgus, rhesus, and baboon, respectively (Table 1).

Testis and Epididymis

The rhesus and cynomolgus macaques had similar testis-to-body-weight ratios, which were in turn higher than that of the baboon. The testis-to-body-weight ratios were 2.17 and 2.36 (g/kg), respectively, in the cynomolgus and rhesus macaque. Baboons, on the other hand, had a testis-to-body-weight ratio of only 1.2. Similar trends were observed for the epididymis for which the epididymis weight-to-body-weight ratios were 0.33, 0.42, and 0.22 (g/kg) for cynomolgus, rhesus, and baboons, respectively (Table 1).

Growth and Development of the Prostate and Testis Across the Lifespan

Changes in body, prostate, and testes weights of the Old World monkeys across the lifespan are shown in Figure 2. The weights increased with age up to around 6 years in cynomolgus macaque, rhesus macaques, and baboons and thereafter there was minimal change in the weights. The changes in growth of the prostate and testis were initially slow or undetectable, but at around 5 years in baboons and rhesus macaques, and at around 4 years of age for cynomolgus macaques, rapid changes occurred (Figure 2).

Serum Prostate Specific Antigen Assay

Serum PSA was detected in all the three Catarrhine (Old World) monkeys (cynomolgus macaques, rhesus macaques, and baboons) studied but not in marmosets, which are members of the Platyrrhine (New World) monkeys. Serum PSA concentration in cynomolgus macaques ranged from 0.04 to 6.2 ng/ml, with a median value of 1.2 ng/ml. Serum PSA values of the rhesus macaques ranged from 0.07 to 2.35 ng/ml with a median value of 0.31, while those of baboons ranged from 0.05 to 0.78 ng/ml with a median value of 0.32 ng/ml. Serum PSA could not be detected in marmosets.

Serum levels of PSA correlated positively with age. The coefficient of correlation (R^2) was 0.30 for cynomolgus macaques, 0.183 for rhesus macaques, and 0.059 for baboons. This correlation was statistically significant for both cynomolgus macaques and rhesus macaques but not for baboons (Figure 3). For both the cynomolgus macaques and rhesus macaques, the serum PSA data include a subset of animals whose values are higher than the rest and are therefore outliers (See supplementary Table 2). When the serum PSA values for the three Old World monkey species are plotted together on the same graph, the cynomolgus macaques and rhesus macaques values are higher and follow a different trend from those of baboons (Figure 3).

Histopathology Lesions

The most common pathological change in the animals studied was inflammation. The inflammatory response was either lymphocytic, suppurative, or a combination of both. Most of the inflammatory changes occurred in the caudal prostate.

A case of prostate hyperplasia in a 17-year-old cynomolgus macaque was studied further and it was found that the hyperplasia involved both the glandular and basal cells (Figure 4). The prostate in this animal contained a single nodular, minimally expansive area composed of closely packed solid nests and glands of cells supported by a thin fibrovascular stroma.

Prostate. Author manuscript; available in PMC 2009 October 1.

The cells had indistinct cell borders, clear to slightly basophilic cytoplasm, round to oval nuclei with lightly stippled chromatin and prominent nucleoli. The cells lining the glandular structures frequently formed papillary structures that extended into the lumen. There was occasional eosinophilic fluid within the lumen of these glands. Mitoses were not observed (Figure 4). An independent examination of this case by the Armed Forces Institute of Pathology (AFIP, Washington, DC) confirmed our earlier diagnosis of hyperplasia involving both the glandular and basal cells. Immunohistochemical staining done by the AFIP showed basal cells that were immunopositive for p63 but not K903. Immunostaining for prostatic acid phosphatase was negative. The serum PSA values at the time of sacrifice for this animal that had prostate hyperplasia was 2.07 ng/ml. Serum PSA in this animal rose 1.12 units in a period of 10 months. A case of focal glandular atrophy, acute prostatitis with subsequent reactive change of mild prostatic hyperplasia in a 20-year-old rhesus macaque was also further studied. This case had a serum PSA level of 2.35 ng/ml. Most prostatic glands in this animal were distended by clear space and lined by attenuated epithelium. Others contained luminal cellular debris and were lined by plump cuboidal cells with increased amounts of basophilic cytoplasm and prominent nuclei with lightly stippled chromatin (Figure 4).

Baboon PSA Cloned and Compared with Other Primate PSA Sequences

Using cloning primers which were based on the published rhesus macaque sequence, the baboon PSA cDNA was cloned. The cloned baboon cDNA sequence has been deposited in the GenBank under accession number EF676031. This sequence was compared to the already published cynomolgus macaque, rhesus macaque, and human PSA sequences. The PSA protein sequences of the rhesus, cynomolgus macaque, and baboon were 89.65%, 89.65%, and 90.42% similar to human PSA, respectively. The rhesus and cynomolgus protein sequences were 99.23% similar to each other and 98.45% similar to their baboon counterpart (Figure 5).

PSA and AMACR Expression in the Prostate of the Three Species

Western blots done on prostate tissue showed that PSA and AMACR were both expressed in cynomolgus macaques, rhesus macaques, and baboons. The macaque species seemed to express more PSA and AMACR per tissue than baboons (Figure 6).

Discussion

The aims of this project were to study the male reproductive systems of nonhuman primates that are commonly used in research in order to evaluate their suitability as models for PSA biology studies in particular and for prostatic diseases in general. Preliminary data reported in this study suggest the macaque species are more suitable for these studies.

New World monkeys like the marmoset are not useful models for studies of PSA biology and prostatic diseases because they do not have the PSA gene. A previous study had reported the presence of the PSA gene in a number of nonhuman primates including orangutans, chimpanzees, gorillas, and macaques but not in dogs, mice, rats, or cows [8]. However, the nonhuman primate species examined in these studies did not include New World monkeys. More recent studies have reported the absence of the PSA gene in both the marmoset and cotton-top tamarin [9–11]. Our failure to detect PSA in marmoset serum in this study confirms the absence of the PSA gene. However, the possibility that the human antibodies used in our studies cannot detect marmoset PSA cannot be ruled out. Probably the PSA gene arose in the Catarrhine primates (Old World monkeys, anthropoid apes and humans) after their phylogenetic lineage separated from the Platyrrhines (New World primates) approximately 35 to 45 million years ago [12,13].

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.