

1995

USP 23

NF 18

Contents

THE UNITED STATES PHARMACOPEIA

THE NATIONAL FORMULARY

By authority of the United States Pharmacopeial Convention, Inc., meeting at Washington, D.C., March 8-10, 1990. Prepared by the Committee of Revision and published by the Board of Trustees

Official from January 1, 1995



UNITED STATES PHARMACOPEIAL CONVENTION, INC.
12601 Twinbrook Parkway, Rockville, MD 20852

NOTICE AND WARNING

Concerning U.S. Patent or Trademark Rights

The inclusion in the Pharmacopeia or in the National Formulary of a monograph on any drug in respect to which patent or trademark rights may exist shall not be deemed, and is not intended as, a grant of, or authority to exercise, any right or privilege protected by such patent or trademark. All such rights and privileges are vested in the patent or trademark owner, and no other person may exercise the same without express permission, authority, or license secured from such patent or trademark owner.

Concerning Use of USP or NF Text

Attention is called to the fact that USP and NF text is fully copyrighted. Authors and others wishing to use portions of the text should request permission to do so from the Secretary of the USPC Board of Trustees.

© 1994 The United States Pharmacopeial Convention, Inc.
12601 Twinbrook Parkway, Rockville, MD 20852.

All rights reserved

ISSN 0195-7996

ISBN 0-913595-76-4 (cloth)

0-913595-81-0 (leather)

Printed by Rand McNally, 1133 County Street, Taunton, MA 02780-3795

SUSPENSIONS

Suspensions are liquid preparations that consist of solid particles dispersed throughout a liquid phase in which the particles are not soluble. Dosage forms officially categorized as Suspensions are designated as such if they are not included in other more specific categories of suspensions, such as Oral Suspensions, Topical Suspensions, etc. (see these other categories). Some suspensions are prepared and ready for use, while others are prepared as solid mixtures intended for constitution just before use with an appropriate vehicle. Such products are designated "for Oral Suspension," etc. The term, Milk, is sometimes used for suspensions in aqueous vehicles intended for oral administration (e.g., *Milk of Magnesia*). The term, Magma, is often used to describe suspensions of inorganic solids such as clays in water, where there is a tendency for strong hydration and aggregation of the solid, giving rise to gel-like consistency and thixotropic rheological behavior (e.g., *Bentonite Magma*). The term, Lotion, has been used to categorize many topical suspensions and emulsions intended for application to the skin (e.g., *Calamine Lotion*). Some suspensions are prepared in sterile form and are used as Injectables, as well as for ophthalmic and otic administration. These may be of two types, ready to use or intended for constitution with a prescribed amount of Water for Injection or other suitable diluent before use by the designated route. Suspensions should not be injected intravenously or intrathecally.

Suspensions intended for any route of administration should contain suitable antimicrobial agents to protect against bacteria, yeast, and mold contamination (see *Emulsions* for some consideration of antimicrobial preservative properties that apply also to Suspensions). By its very nature, the particular matter in a suspension may settle or sediment to the bottom of the container upon standing. Such sedimentation may also lead to caking and solidification of the sediment with a resulting difficulty in redispersing the suspension upon agitation. To prevent such problems, suitable ingredients that increase viscosity and the gel state of the suspension, such as clays, surfactants, polyols, polymers, or sugars, should be added. It is important that suspensions always be shaken well before use to ensure uniform distribution of the solid in the vehicle, thereby ensuring uniform and proper dosage. Suspensions require storage in tight containers.

Oral Suspensions

Oral Suspensions are liquid preparations containing solid particles dispersed in a liquid vehicle, with suitable flavoring agents, intended for oral administration. Some suspensions labeled as Milks or Magmas fall into this category.

Topical Suspensions

Topical Suspensions are liquid preparations containing solid particles dispersed in a liquid vehicle, intended for application to the skin. Some suspensions labeled as Lotions fall into this category.

Otic Suspensions

Otic Suspensions are liquid preparations containing micronized particles intended for instillation in the outer ear.

Ophthalmic Suspensions

(See *Ophthalmic Preparations*).

SYRUPS

See *Solutions*.

SYSTEMS

In recent years, a number of dosage forms have been developed using modern technology that allows for the uniform release or targeting of drugs to the body. These products are commonly called delivery systems. The most widely used of these are Transdermal Systems.

Transdermal Systems

Transdermal drug delivery systems are self-contained, discrete dosage forms that, when applied to intact skin, are designed to deliver the drug(s) through the skin to the systemic circulation. Systems typically comprise an outer covering (barrier), a drug reservoir, which may have a rate controlling membrane, a contact adhesive applied to some or all parts of the system and the system/skin interface, and a protective liner that is removed before applying the system. The activity of these systems is defined in terms of the release rate of the drug(s) from the system. The total duration of drug release from the system and the system surface area may also be stated.

Transdermal drug delivery systems work by diffusion: the drug diffuses from the drug reservoir, directly or through the rate controlling membrane and/or contact adhesive if present, and then through the skin into the general circulation. Typically, modified-release systems are designed to provide drug delivery at a constant rate, such that a true steady state blood concentration is achieved and maintained until the system is removed. At that time, blood concentration declines at a rate consistent with the pharmacokinetics of the drug.

Transdermal drug delivery systems are applied to body areas consistent with the labeling for the product(s). As long as drug concentration at the system/skin interface remains constant, the amount of drug in the dosage form does not influence plasma concentrations. The functional lifetime of the system is defined by the initial amount of drug in the reservoir and the release rate from the reservoir.

NOTE—Drugs for local rather than systemic effect are commonly applied to the skin embedded in glue on a cloth or plastic backing. These products are defined traditionally as plasters or tapes.

Ocular System

Another type of system is the ocular system, which is intended for placement in the lower conjunctival fornix from which the drug diffuses through a membrane at a constant rate over a seven-day period (e.g., *Pilocarpine Ocular System*).

Intrauterine System

An intrauterine system, based on a similar principle but intended for release of drug over a much longer period of time, i.e., one year, is also available (e.g., *Progesterone Intrauterine Contraceptive System*).

TABLETS

Tablets are solid dosage forms containing medicinal substances with or without suitable diluents. They may be classed, according to the method of manufacture, as compressed tablets or molded tablets.

The vast majority of all tablets manufactured are made by compression, and compressed tablets are the most widely used dosage form in this country. Compressed tablets are prepared by the application of high pressures, utilizing steel punches and dies, to powders or granulations. Tablets can be produced in a wide variety of sizes, shapes, and surface markings, depending upon the design of the punches and dies. Capsule-shaped tablets are commonly referred to as caplets. Boluses are large tablets intended for veterinary use, usually for large animals.

Molded tablets are prepared by forcing dampened powders under low pressure into die cavities. Solidification depends upon crystal bridges built up during the subsequent drying process, and not upon the compaction force.

Tablet triturates are small, usually cylindrical, molded or compressed tablets. Tablet triturates were traditionally used as dispensing tablets in order to provide a convenient, measured quantity of a potent drug for compounding purposes. Such tablets are rarely used today. Hypodermic tablets are molded tablets made from completely and readily water-soluble ingredients and formerly were intended for use in making preparations for hypodermic injection. They are employed orally, or where rapid drug availability is required such as in the case of *Nitroglycerin Tablets*, sublingually.

Buccal tablets are intended to be inserted in the buccal pouch, and sublingual tablets are intended to be inserted beneath the