

*Brit. J. Ophthalm.* (1955), 39, 90.

## KERATO-CONJUNCTIVITIS SICCA TREATED WITH CORTISONE AND ACTH\*

BY

STELLA EADIE AND MALCOLM THOMPSON

*Edinburgh*

THE effects of cortisone and ACTH in cases of kerato-conjunctivitis sicca (KCS) have been recorded by several authors (Table I). Reports of their value have been conflicting, probably because most observers only treated a single case, and because in several instances only limited supplies of the hormones were available, so that treatment may have been inadequate.

TABLE I  
RESULTS OF TREATMENT BY CORTISONE AND ACTH ALREADY RECORDED

Authors	Date	No. of Cases	Treatment	Result
Offret and Forest	1950, 1951	1	Cortisone	Unchanged
Phillips	1952	1		Unchanged
Cristini	1952	1		Improved
Beiglböck and Hoff	1952	1		Improved
Fernández y Fernandes and Marañón	1952	1		Improved
Forestier and others	1951	1		Improved
Cadman and Robertson	1952	2	ACTH	Unchanged
Cristini	1952	2		Unchanged
Sjögren and Eriksen	1952	4		Unchanged
Fitzgerald and others	1951	1		Unchanged
Stephens	1950	1		Improved
Forestier and others	1951	1		Improved
Frenkel and others	1951	1		Improved
Bourne	1952	1		Cortisone Eye Drops
Phillips	1952	1	Improved	

The purpose of this paper is to report on twelve cases of kerato-conjunctivitis sicca treated with cortisone drops, systemic cortisone, or ACTH. The cortisone drops used were a 0.625 per cent. solution of cortisone acetate (Merck) in a buffered base, except where stated otherwise.

\*Received for publication August 20, 1954.

**CORTISONE AND ACTH IN KERATO-CONJUNCTIVITIS SICCA 91****Case Notes**

**Case 1, a female aged 41 years**, had had rheumatic fever in childhood with resultant mitral stenosis and aortic regurgitation. She had had rheumatoid arthritis for 28 years, latterly associated with KCS, xerostoma, salivary gland enlargement, and a persistent dry cough with small areas of pulmonary atelectasis and recurrent laryngitis. The ocular symptoms were relieved during a course of oral cortisone in December, 1951.

An exacerbation of ocular symptoms occurred in October, 1952, when Schirmer's test gave 5 mm. in both eyes, and vision was 6/60 in the right eye, and 6/18 in the left. She was treated with cortisone drops three times daily and obtained relief within 48 hours. In November, 1952, vision was 6/18 in the right eye and 6/5 in the left. Staining was still present but less marked.

In March, 1953, a course of Acthar gel relieved her arthritic and ocular symptoms. The parotid and submandibular swellings subsided. Schirmer's test was not significantly improved (8 mm. maximum). A relapse occurred on cessation of ACTH therapy (April, 1953). The subsequent course is shown in Table II.

**TABLE II**  
**PROGRESS OF CASE 1**

Date of Examination	Right Eye				Left Eye			
	Symptoms	Staining	Vision	Schirmer's Test (mm.)	Symptoms	Staining	Vision	Schirmer's Test (mm.)
13.11.53	+++	+++	6/36	2	+++	+++	6/36	2
	No treatment				No treatment			
20.11.53	+++	+++	6/36	0	+++	+++	6/24pt.	3
	Cortisone drops 3-hrly				Cortisone drops 3-hrly			
27.11.53	++	+++	6/12	2	+	++	6/18+	4
	Cortisone drops three times daily				Cortisone drops three times daily			
11.12.53	++	+++	6/24	0	++	++	6/18+	0
	Cortisone drops three times daily				Cortisone drops three times daily			
23.12.53	++	+++	6/18+	4	+	++	6/9	3
	Hydrocortisone drops three times daily				Cortisone drops three times daily			
8.1.54	0	++	6/12	0	0	+++	6/12pt.	3
	Hydrocortisone drops three times daily				Cortisone drops three times daily			
22.1.54	0	++	6/9pt.	0	0	+++	6/18+	3
	Hydrocortisone drops three times daily				Cortisone drops three times daily			
14.2.54	0	++	6/9+	—	0	++	6/9	—
	Hydrocortisone drops three times daily				Cortisone drops three times daily			
26.2.54	+	+++	6/9pt.	2	+	+++	6/9pt.,	3
	Lacrimal puncta sealed				Nil			
31.5.54	0	++	6/6+	5	++	+++	6/24	5

**Case 2, a female aged 64 years,** had had severe rheumatoid arthritis for 25 years, and intermittent soreness of the eyes for 8 years. Schirmer's test gave 3 mm. in both eyes. In March, 1953, vision was 6/18 in the right eye and 6/60 in the left. She was treated with cortisone drops three times daily, and in April, 1953, vision was 6/6 in the right eye and 6/36 in the left. When treatment was discontinued in July, 1953, she relapsed, but she improved again when treatment was resumed. In November, 1953, while she was still receiving drops, vision was 6/5 in the right eye and 6/9 in the left. Schirmer's test showed 4 mm. in both eyes. The cortisone drops were then withdrawn. The subsequent course is shown in Table III.

TABLE III  
PROGRESS OF CASE 2

Date of Examination	Right Eye				Left Eye			
	Symptoms	Staining	Vision	Schirmer's Test (mm.)	Symptoms	Staining	Vision	Schirmer's Test (mm.)
10.12.53	+++	—	6/5	5	+++	—	6/12	4
	Cortisone drops three times daily				Cortisone drops three times daily			
15.12.53	+	+++	6/6pt.	—	+	+++	6/12+	—
	Nil				Nil			
18.1.54	+	+++	6/6pt.	10	+	+++	6/12+	10
	Nil				Nil			
9.2.54	+	+++	6/6pt.	6	+++	+++	6/60	3
	Nil				Cortisone drops four times daily			
19.2.54	+	+++	6/5pt.	4	+	+++	6/18	8
	Nil				Cortisone drops four times daily			
25.3.54	+	++	6/6	—	+	+++	6/18+	—
	Nil				Hydrocortisone drops four times daily			
6.5.54	+	+++	6/6pt.	0	+	+++	6/36	1
	Nil				Lacrimal puncta sealed			
21.5.54	++	+++	6/6pt.	1	0	++	6/9	4
	Cortisone drops three times daily				Puncta still sealed			
17.6.54	0	—	6/5pt.	—	0	—	6/6pt.	—

**Case 3, a female aged 47 years,** had had xerostoma and inflamed eyes for 5 years, with no rheumatoid arthritis, but anaemia and achlorhydria. She obtained no relief from large doses of vitamins and oral iron, nor from antibiotic eye drops. In June, 1953, she had severe KCS; vision was 6/18 in the right eye and 6/9+ in the left. Treatment with cortisone drops three times a day gave symptomatic relief and the corneal staining decreased. Cortisone drops were discontinued in November, 1953. The subsequent course is shown in Table IV (opposite).

**CORTISONE AND ACTH IN KERATO-CONJUNCTIVITIS SICCA** 93TABLE IV  
PROGRESS OF CASE 3

Date of Examination	Right Eye				Left Eye			
	Symptoms	Staining	Vision	Schirmer's Test (mm.)	Symptoms	Staining	Vision	Schirmer's Test (mm.)
5.11.53	+++	+++	6/18	2	+++	+++	6/9	6
	Nil				Nil			
10.11.53	+++	+++	6/18pt.	0	+++	+++	6/9pt.	0
	Intensive vitamin therapy				Intensive vitamin therapy			
5.1.54	+++	+++	6/24pt.	2	+++	+++	6/9pt.	0
	Cortisone drops 3-hrly for 2 days then four times daily				Cortisone drops 3-hrly for 2 days then four times daily			
19.1.54	++	+++	6/18pt.	0	+	++	6/9pt.	0
	Cortisone drops four times daily				Cortisone drops four times daily			
2.2.54	++	+++	6/24pt.	0	+	+++	6/12	0
	Lacrimal puncta sealed				Cortisone drops four times daily			
16.2.54	+	+++	6/18+	6	+	++	6/12+	0
	Lacrimal puncta sealed				Cortisone drops four times daily			
26.2.54	+	++	6/12pt.	2	+	+++	6/12+	0
	Lacrimal puncta re-sealed				Lacrimal puncta sealed			
13.3.54	0	—	—	4	0	—	—	6
	Lacrimal puncta still sealed				Lacrimal puncta still sealed			
4.5.54	+	+	6/12pt.	6	0	++	6/9pt.	6

**Case 4, female aged 54 years**, had had severe rheumatoid arthritis for 25 years, sore eyes intermittently for 18 months, xerostoma, and bilateral parotid swelling. In December, 1953, the right eye was enucleated after perforation of a corneal ulcer. The left eye showed KCS, Schirmer's test gave 4 mm., and vision in the left eye was 6/12. Cortisone drops three times a day gave only slight subjective improvement. The staining and Schirmer's test did not improve and vision in the left eye deteriorated to 6/18. Cortisone was instilled 5 times daily without further improvement. In February, 1954, the left lacrimal puncta and canaliculi were sealed, giving marked symptomatic improvement. In May, 1954, the improvement was maintained, vision was 6/6, and Schirmer's test gave 3 mm., but the corneal staining was still present.

**Case 5, female aged 44 years**, had had rheumatoid arthritis for 22 years, recurrent laryngitis, and inflamed eyes for one year. Various eye drops had not given any relief. In February, 1954, vision was 6/18 in both eyes, KCS was present, and Schirmer's test gave 3 mm. in the right eye and 1 mm. in the left. Cortisone drops 1.5 per cent. solution were given four times a day to the right eye and hydrocortisone drops 1.5 per cent. solution four times a day to the left eye. After 2 weeks the symptoms in the right eye were relieved, but no improvement was seen in the left eye. Vision was 6/9 in the right eye

and 6/24 in the left eye. Later cortisone drops 1·5 per cent. solution were given four times daily to both eyes, but there was no change in the condition of the left eye.

Treatment was discontinued in April, 1954, and a full relapse occurred. In May, 1954, cortisone drops four times daily were combined with cortisone ointment applied to the eyes at night. After 2 weeks, the right eye had again responded but there was no improvement in the left eye. In June, 1954, the left lacrimal puncta were sealed and cortisone to the left eye was stopped. The eye improved with complete symptomatic relief and vision improved to 6/9. This improvement has been maintained.

**Case 6, a female aged 38 years**, had had rheumatoid arthritis for 17 years, with Raynaud's disease, xerostoma, intermittent bilateral parotid swellings, recurrent laryngitis, and attacks of "conjunctivitis". In November, 1954, she was thought to have KCS; Schirmer's test gave 0 mm. in the right eye and 1 mm. in the left eye. Vision was 6/9 in both eyes. Cortisone drops were given four times daily with marked subjective relief in both eyes and reduction of staining in the right. The cortisone was reduced to three times daily and the improvement was maintained. In January, 1954, 1 per cent. hydrocortisone drops were given four times daily to the left eye and cortisone 1 per cent. four times daily to right eye. There was no further improvement in either eye.

**Case 7, a female aged 46 years**, had had alopecia totalis since 1940, dry skin and trophic changes in the nails since 1946, xerostoma and dry cough since 1948, recurrently sore eyes since April, 1951, and severe sinusitis and rheumatoid arthritis since September, 1951. In February, 1952, she was given oral cortisone 100 mg. daily, with relief of ophthalmic symptoms and improvement in general condition. Schirmer's test gave 8 mm. in both eyes. The visual acuity was 6/6 in each eye. After one year of therapy, Schirmer's test gave 2 mm. in the right eye and 3 mm. in the left eye. Staining with rose bengal solution was present throughout.

After 2 years of treatment, oral cortisone therapy was withdrawn during January, 1954, and the ocular symptoms recurred within a few days of withdrawal, with bilateral conjunctival injection. Cortisone eye-drops were given, and it was necessary to instil the drops 2-hourly to relieve the patient's symptoms completely. Schirmer's test continued to give 2 mm. in both eyes.

Oral cortisone therapy, 75 mg. daily, was resumed in February, and this partially controlled the ocular and arthritic symptoms; when the dosage was increased to 100 mg. daily, the symptoms were completely relieved.

**Case 8, a female aged 66 years**, had had severe rheumatoid arthritis for 4 years, and xerostoma and KCS for 1 year. In July, 1952, she was treated with cortisone eye-drops three times daily and chloramphenicol eye ointment at night. Relief was obtained but the patient relapsed when the cortisone was stopped. In October, 1952, cortisone eye-drops were given to the left eye three times daily with marked symptomatic improvement.

**Case 9, female aged 59 years**, had had rheumatoid arthritis for 13 years, recurrent attacks of conjunctival injection for 3 years, and xerostoma. In January, 1954, examination revealed Sjögren's syndrome with signs of Felty's syndrome.

When mild exacerbation of ocular symptoms occurred in February, 1954, Schirmer's test gave 3 mm. in both eyes, and 1 per cent. cortisone eye-drops four times daily gave relief within 3 days; after 9 days of treatment the drops were withdrawn and relapse occurred. Further therapy with 1 per cent. cortisone eye-drops four times daily was then given for 3 weeks with prompt relief; since the end of course the patient has had no further ocular symptoms.

**Case 10, a female aged 60 years**, had had severe rheumatoid arthritis for 20 years, and sore eyes and photophobia for 1 year, with xerostoma; KCS was diagnosed and Schirmer's test gave 3 mm. in the right eye and 5 mm. in the left; vision was 6/24 in the right eye and 6/9 in the left.

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.