

A Wise Owl Once Asked,  
"Why Are You Still Running  
Your Own Westerns?"



Auto-Western Service

Learn More

 RayBiotech  
Empowering your proteomics

 The Journal of  
Immunology

## Desiccating Stress Induces T Cell-Mediated Sjögren's Syndrome-Like Lacrimal Keratoconjunctivitis

This information is current as  
of June 26, 2017.

Jerry Y. Niederkorn, Michael E. Stern, Stephen C.  
Pflugfelder, Cintia S. De Paiva, Rosa M. Corrales, Jianping  
Gao and Karyn Siemasko

*J Immunol* 2006; 176:3950-3957; ;  
doi: 10.4049/jimmunol.176.7.3950  
<http://www.jimmunol.org/content/176/7/3950>

- 
- References** This article **cites 20 articles**, 9 of which you can access for free at:  
<http://www.jimmunol.org/content/176/7/3950.full#ref-list-1>
- Subscription** Information about subscribing to *The Journal of Immunology* is online at:  
<http://jimmunol.org/subscription>
- Permissions** Submit copyright permission requests at:  
<http://www.aai.org/About/Publications/JI/copyright.html>
- Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at:  
<http://jimmunol.org/alerts>

Downloaded from <http://www.jimmunol.org/> by guest on June 26, 2017

---

*The Journal of Immunology* is published twice each month by  
The American Association of Immunologists, Inc.,  
1451 Rockville Pike, Suite 650, Rockville, MD 20852  
Copyright © 2006 by The American Association of



**DOCKET  
ALARM**

Find authenticated court documents without watermarks at [docketalarm.com](http://docketalarm.com).

# Desiccating Stress Induces T Cell-Mediated Sjögren's Syndrome-Like Lacrimal Keratoconjunctivitis<sup>1</sup>

Jerry Y. Niederkorn,<sup>2,3\*</sup> Michael E. Stern,<sup>2†</sup> Stephen C. Pflugfelder,<sup>2‡</sup> Cintia S. De Paiva,<sup>‡</sup> Rosa M. Corrales,<sup>‡</sup> Jianping Gao,<sup>†</sup> and Karyn Siemasko<sup>†</sup>

Chronic dry eye syndrome affects over 10 million people in the United States; it is associated with inflammation of the lacrimal gland (LG) and in some cases involves T cell infiltration of the conjunctiva. We demonstrate that environmental desiccating stress (DS) elicits T cell-mediated inflammation of the cornea, conjunctiva, and LG, but not other organs in mice. The lacrimal keratoconjunctivitis (LKC) was mediated by CD4<sup>+</sup> T cells, which, when adoptively transferred to T cell-deficient nude mice, produced inflammation in the LG, cornea, and conjunctiva, but not in any other organ. Adoptively transferred CD4<sup>+</sup> T cells produced LKC even though recipients were not exposed to DS. LKC was exacerbated in euthymic mice depleted of CD4<sup>+</sup>CD25<sup>+</sup> forkhead/winged helix transcription factor<sup>+</sup> regulatory T cells. The results suggest that DS exposes shared epitopes in the cornea, conjunctiva, and LG that induce pathogenic CD4<sup>+</sup> T cells that produce LKC, which under normal circumstances is restrained by CD4<sup>+</sup>CD25<sup>+</sup> forkhead/winged helix transcription factor<sup>+</sup> regulatory T cells. *The Journal of Immunology*, 2006, 176: 3950–3957.

**D**ry eye is a common ocular disease and is one of the leading causes of patients seeking ophthalmic care. Dry eye or lacrimal keratoconjunctivitis (LKC)<sup>4</sup> has a tremendous impact on quality of life and can cause debilitating eye pain and loss of vision. In animal models and humans, chronic dry eye disease is often associated with T cell infiltration of the conjunctiva (1–3). T cell infiltration has also been observed in dry eye patients with or without a systemic autoimmune disease, such as Sjögren's syndrome (SS) (3). It has been proposed that in some forms of dry eye disease, autoreactive T cells alter the function of the ocular surface epithelium and also cause reduced tear secretion (4). The presence of CD4<sup>+</sup> T cells in the epithelium and stroma of the conjunctiva and the salutary effect of the T cell-immunosuppressive drug, cyclosporine, in some patients support the proposition that dry eye is an immune-mediated disease (2, 5). However, others have argued that inflammation of the cornea and conjunctiva is the result, not the cause, of dry eye disease.

It is now recognized that the ocular surface and lacrimal glands function as an integrated unit, termed the lacrimal functional unit (LFU), which is linked by their sensory and autonomic nerves (4). Inflammation in any component of the LFU has the capability to compromise the function of other components through soluble me-

diators, generation of autoreactive T cells to shared Ags, and inhibition of neural transmission (4). SS causes profound dysfunction of the LFU, which includes decreased tear secretion, loss of conjunctival goblet cells (GC) transformation of the ocular surface epithelium to a keratinized surface. Immunopathological changes, including immune-mediated inflammation and increased production of inflammatory cytokines and chemokines, have been detected in the dysfunctional lacrimal glands and ocular surface tissues in SS (6–8).

It has been proposed that a wide range of factors that alter the integrity of tissues create danger signals that activate APCs and promote the generation of immune responses to self Ags (9). The danger hypothesis has been implicated in the pathogenesis of SS (10) and is consistent with the observation that desiccating stress (DS) to the ocular surface induces the expression of proinflammatory cytokines (11). With this in mind, we tested the hypothesis that environmental DS to the ocular surface would break self-tolerance and induce the development of T cell-mediated inflammation directed against epitopes expressed on the ocular surface and the lacrimal gland (LG).

## Materials and Methods

### Mice

Female BALB/c, T cell-deficient nude BALB/c (BALB/cByJ-Hfh11<v>), and C57BL/6 mice, 6–8 wk old, were purchased from The Jackson Laboratory or Charles River Laboratories. All animal experiments were approved by the institutional animal care and use committees at Allergan and Baylor College of Medicine. All studies adhered to the Association for Research in Vision and Ophthalmology statement for the Use of Animals in Ophthalmic and Vision Research.

### Induction of DS in mice

DS was induced by s.c. injection of scopolamine hydrobromide (0.5 mg/0.2 ml; Sigma-Aldrich) four times a day (0800, 1200, 1400, and 1700 h), alternating between the left and right flanks of 4- to 6-wk-old mice, as previously described (12). Up to five mice were placed in a cage with a perforated plastic screen on one side to allow airflow from a fan (Cafrano) placed 6 in. in front of it for 16 h/day. Room humidity was maintained at 30–35% and temperature at 80°F. DS was induced for either 5 or 12 consecutive days. This model of dry eye disease has been used previously with no discernible ill effects from the scopolamine treatment or low humidity (11, 12). Changes in corneal permeability were assessed by measuring corneal staining by Oregon Green dextran (OGD; 70,000 m.w.; Molecular Probes) as previously described (13). Aqueous tear production was assessed

\*Department of Ophthalmology, University of Texas Southwestern Medical Center, Dallas, TX 75390; †Department of Biological Sciences, Allergan, Irvine, CA 92623; and ‡Department of Ophthalmology, Baylor College of Medicine, Houston, TX 77030  
Received for publication December 8, 2005. Accepted for publication January 19, 2006.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported by Grant EY11915 and unrestricted grants from Research to Prevent Blindness (New York, NY) and Allergan.

<sup>2</sup> J.Y.N., M.E.S., and S.C.P. contributed equally to this study.

<sup>3</sup> Address correspondence and reprint requests to Dr. Jerry Y. Niederkorn, Department of Ophthalmology, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75390. E-mail address: jerry.niederkorn@utsouthwestern.edu

<sup>4</sup> Abbreviations used in this paper: LKC, lacrimal keratoconjunctivitis; CLN, cervical lymph node; DS, desiccating stress; FoxP3, forkhead/winged helix transcription factor; GC, goblet cell; LFU, lacrimal functional unit; LG, lacrimal gland; NS, non-stressed; OGD, Oregon Green dextran; SS, Sjögren's syndrome.

using cotton thread (Quick Thread; FCI Ophthalmics) as previously described (14). Control mice were maintained in a nonstressed (NS) environment containing 50–75% relative humidity without exposure to forced air. Tear production was measured with phenol red-impregnated cotton threads (Zone-Quick; Oasis) placed into the tear meniscus of the lateral canthus for 30 s.

#### Measurement of GC density

Surgically excised eyes were fixed in 10% formalin and embedded in paraffin. Sections (6  $\mu\text{m}$ ) were stained with periodic acid-Schiff reagent. Sections from three eyes in each group (in two different sets of experiments) were examined and photographed with a Nikon Eclipse E400 microscope equipped with a Nikon DXM 1200 digital camera. GC density was measured in the superior and inferior bulbar and tarsal conjunctiva and expressed as number per 100  $\mu\text{m}$  using Metavue 6.24r software (Molecular Devices).

#### In vivo analysis of adoptively transferred CFSE-labeled lymphocytes

Spleen cells ( $1 \times 10^7/\text{ml}$ ) were collected from BALB/c mice subjected to DS and were labeled with 10  $\mu\text{M}$  CFSE (Molecular Probes) for 15 min at room temperature. Cells were washed three times in RPMI 1640 and were injected i.p. into BALB/c mice ( $1 \times 10^7$  spleen cells/0.1 ml/mouse).

#### Immunohistochemistry

OCT-embedded globes with attached lids were sectioned at 10- $\mu\text{m}$  thickness and stained for the expression of mouse CD4 (rat anti-mouse CD4; rat IgG2a, k; clone H129.19; BD Pharmingen) using Vectastain Elite ABC reagents (Vector Laboratories).

#### Adoptive cell transfer and anti-Thy 1.2 Ab treatment

Spleens and cervical lymph nodes (CLN) were collected from mice subjected to DS and NS, and one donor-equivalent of either spleen or CLN cells was transferred i.p. to syngeneic nude mice. One donor-equivalent is defined as the number of cells remaining after the respective in vitro manipulation (e.g., anti-Thy 1.2 Ab treatment or CD4<sup>+</sup> T cell enrichment) of a single spleen or CLN from a single donor. The remaining cells represent the total lymphocyte population for that spleen cell category for a single donor. One splenic equivalent of T cells was equal to  $\sim 5 \times 10^7$  cells. Spleen cells were depleted of T cells by in vitro treatment with anti-Thy 1.2 Ab (BD Pharmingen) in the presence of complement. Control aliquots of spleen cells were treated with complement alone. Cells were washed with RPMI 1640 medium before being adoptively transferred in 0.1 ml of RPMI 1640 medium.

#### CD4<sup>+</sup> and CD25<sup>+</sup> T cell enrichment

BALB/c CLN cell suspensions were enriched for CD4<sup>+</sup> T cells by positive selection using rat anti-mouse CD4-conjugated magnetic microbeads (MACS system; Miltenyi Biotec) as described previously (15). The CD4-enriched cell suspensions contained  $>87\%$  CD4<sup>+</sup> T cells as determined by flow cytometry. CD4<sup>+</sup>CD25<sup>+</sup> T cells were enriched using a mouse CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cell magnetic isolation kit (Miltenyi Biotec) according to the manufacturer's instructions.

Permeabilized CD4<sup>+</sup>CD25<sup>+</sup> cells were tested for the expression of cytoplasmic forkhead/winged helix transcription factor (Foxp3) protein. Briefly, CD4<sup>+</sup>CD25<sup>+</sup> cells were washed three times in HBSS, resuspended in Cytotfix/Cytoperm solution (BD Pharmingen), and incubated for 30 min at room temperature. Cells were washed three times in Perm/Wash buffer and resuspended with an FITC-labeled rat anti-mouse Foxp3 Ab (eBiosciences) at a concentration of 1  $\mu\text{g}/\text{ml}$  for 30 min on ice. Cells were washed in Perm/Wash buffer three times, incubated with PE-labeled secondary Ab for 20–30 min at 4°C, washed three additional times in Perm/Wash buffer, fixed in 1% paraformaldehyde, and assessed for fluorescence in a FACScan flow cytometer (BD Biosciences). The results were analyzed using CellQuest version 3.1f software (BD Biosciences). CD4<sup>+</sup>CD25<sup>+</sup> cells were found to be 89% Foxp3<sup>+</sup> (data not shown).

#### In vivo depletion of CD25<sup>+</sup> cells


In vivo depletion of CD25<sup>+</sup> regulatory T cells was achieved as described previously (16). Briefly, anti-CD25 Ab (hybridoma PC61; American Type Culture Collection) was administered i.p. to normal BALB/c mice 7 days before and on the day DS was initiated.

#### Statistics

## Results

### DS induces immune-mediated inflammation of the ocular surface and lacrimal gland

C57BL/6 mice exposed to DS displayed altered corneal epithelial barrier function, which was reflected in an 85% increase in the uptake of the OGD label that was used to measure the integrity of the corneal epithelial barrier (Fig. 1). Histopathological analysis revealed reduced numbers of conjunctival GC, decreased tear production, and CD4<sup>+</sup> T cell infiltration of the conjunctival basal




**FIGURE 1.** Increased corneal permeability induced by DS. C57BL/6 mice were subjected to 5 days of DS or NS conditions. Mouse corneas were viewed and photographed under cobalt blue light 10 min after application of 1  $\mu\text{l}$  of the fluorescent molecule OGD to the ocular surface. *A*, Ocular surface of mouse subjected to 5 days of DS. *B*, Ocular surface of untreated, NS mouse. *C*, Quantification of OGD uptake by mouse of NS control

epithelium (Fig. 2, A–E). The greater severity of keratoconjunctivitis in C57BL/6 mice than BALB/c mice is consistent with the observation that BALB/c mice have a larger repertoire of CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells compared with C57BL/6 mice (17). The role of CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells was tested by treating BALB/c mice with anti-CD25 Ab to deplete natural regulatory T cells before subjecting the mice to DS. Anti-CD25-treated BALB/c mice developed significant keratoconjunctivitis and displayed a steep reduction in the density of conjunctival GC compared with the isotype control mice and mice subjected to DS without Ab treatment (Fig. 2F).

*DS-induced lacrimal keratoconjunctivitis can be adoptively transferred with CD4<sup>+</sup> T cells*


Adoptive transfer experiments were performed to determine the role of T cells in the ocular inflammation elicited by DS. Draining

CLN cells were collected from BALB/c mice after a 5-day exposure to DS and were adoptively transferred to athymic (nude) BALB/c mice that were maintained under NS conditions (50–75% relative humidity). Even though the adoptive cell transfer recipients were NS, they had reduced tear production and developed LKC after receiving CLN cells from donors treated with DS (Fig. 3). Inflammation was immune mediated, because the nude mouse recipients of CLN cells from NS donors maintained normal tear production and did not develop inflammation of the conjunctiva (Fig. 3, B and C). Importantly, inflammation in recipients of CLN cells from DS donors was restricted to the lacrimal functional unit (cornea, conjunctiva, and LG), and no significant inflammatory cell infiltrates were found in the salivary gland, oral mucosa, thyroid, heart, lung, colon, spleen, adrenal gland, or vagina (data not shown). Histopathological analysis revealed significant LG inflammation and loss of conjunctival GC, which coincided with a steep



**FIGURE 2.** DS induces LKC. C57BL/6 mice (A–E) and BALB/c mice (A, B, and F) were maintained in either NS conditions (50–75% relative humidity) or DS (30–35% relative humidity) for either 5 days (A–E) or 12 days (A and C). Five days of DS consistently produced LKC in C57BL/6 mice. A, Reduced numbers of conjunctival GC; B, diminished tear production; C, CD4<sup>+</sup> cell density in conjunctivae of NS mice. D, Mice treated with DS; CD4<sup>+</sup> T cells (brown staining) in conjunctivae of DS mice. E, CD4<sup>+</sup> T cell density in conjunctivae of DS and NS mice. F, Reduced conjunctival GC density in BALB/c mice treated with DS, DS plus anti-CD25 Ab, or DS plus isotype Ab.

Downloaded from <http://www.jimmunol.org/> by guest on June 26, 2017




**FIGURE 3.** Adoptive T cell transfer of LKC. Euthymic BALB/c mice (normal) were subjected to either DS or NS conditions for 5 days, and one donor-equivalent of CLN cells was adoptively transferred to each athymic (nude) BALB/c nude mouse recipient. Nude mice were maintained under NS conditions after adoptive cell transfer and were assessed for tear production

reduction in tear production (Fig. 3A) and the appearance of ocular surface inflammation (Fig. 3, B and C). Immunohistochemical staining demonstrated a significant infiltrate of CD4<sup>+</sup> T cells in the LG in the adoptive cell transfer recipients (Fig. 4). Inflammation was T cell dependent, because in vitro treatment of CLN cell suspensions with anti-Thy 1.2 Ab plus complement before adoptive transfer prevented the development of LKC (Fig. 5A). Moreover, histopathological examination of the eyes and LGs of recipients of CLN cell suspensions depleted of T cells demonstrated significantly reduced mononuclear cellular infiltrates in the conjunctivae and the sparing of 80% of the normal tear production (Fig. 5, B and C). To determine whether the CD4<sup>+</sup> T cells in the conjunctivae and LGs contributed to LKC, adoptive transfer experiments were performed in which either CD4<sup>+</sup> T cell-enriched or CD4<sup>+</sup> T cell-depleted CLN cell suspensions were collected from mice subjected to DS and transferred to syngeneic BALB/c nude mice. Recipients of CD4<sup>+</sup> T cell-enriched CLN cells developed intense LG inflammation (Fig. 5D), with significant increases in the number of both conjunctival mononuclear cells and neutrophils (data not shown). Moreover, CD4<sup>+</sup> T cells were detected in the LG of the adoptive transfer recipients (Fig. 5G). Additional evidence of CD4<sup>+</sup> T cell involvement was demonstrated in experiments in which CD4<sup>+</sup> T cells were collected from mice subjected to DS, labeled with CFSE, and adoptively transferred to nude mice that were maintained in an NS environment. Accumulations of CFSE-labeled CD4<sup>+</sup> T cells were found in the conjunctival epithelium of the nude mice that received adoptive cell transfers (Fig. 5H), but were absent in nonocular organs (data not shown). By contrast, adoptive transfer of CFSE-labeled CD4<sup>+</sup> T cells from normal donors did not preferentially localize in the corneas or conjunctivae of nude mouse recipients (Fig. 6), suggesting that mice subjected to DS develop CD4<sup>+</sup> T cells that preferentially localize in the eye and LGs, where they produce inflammatory disease, GC loss, and LG dysfunction.

*Role of CD4<sup>+</sup>CD25<sup>+</sup> T cells in mitigating LKC*

The inflammation produced in the adoptive cell transfer recipients was consistently more severe in nude mice than euthymic mice, suggesting the possible mitigating effects of one or more regulatory T cell populations in the euthymic host. To examine the role of natural regulatory T cells in LKC, euthymic BALB/c mice were treated with anti-CD25 mAb to deplete putative natural regulatory T cells before receiving adoptively transferred T cells from BALB/c donors treated with DS. Euthymic BALB/c



**FIGURE 4.** CD4<sup>+</sup> T cell infiltrate in lacrimal glands in recipients of adoptively transferred CD4<sup>+</sup> T cells from mice subjected to DS. A, CD4<sup>+</sup> T cells (brown) in LGs in recipients of CLN cells from DS donors. B,

Downloaded from <http://www.jimmunol.org/> by guest on June 26, 2017

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.