

Acute megaloblastic anemia: homocysteine levels are useful for diagnosis and follow-up.

By: Carrasco, M; Remacha, A; Sureda, A; Sarda, P; Martino, R; Sierra, J
View ResearcherID and ORCID (provided by Thomson Reuters)

Haematologica
Volume: 84 Issue: 8 Pages: 767-8
Published: 1999-Aug

Author Information

Categories / Classification

Research Areas: Pathology; Hematology; Biochemistry & Molecular Biology (provided by Thomson Reuters)
MeSH Terms:

Heading	Qualifier
Acute Disease	
Anemia, Megaloblastic	*blood diagnosis physiopathology
Biological Markers	
Homocysteine	*blood
Humans	
Male	
Middle Aged	
Prognosis	

Chemical:

Registry Number	Substance
0	Biological Markers
0LVT1QZ0BA	Homocysteine

Document Information

Document Type: Case Reports; Letter
Language: English
PubMed ID: 10457427
NLM Unique ID: 0417435
Date Created: 12 Oct 1999 Date Completed: 12 Oct 1999 Date Revised: 21 Nov 2013
Country: ITALY
ISSN: 0390-6078

Journal Information

Table of Contents: Current Contents Connect®
Impact Factor: Journal Citation Reports®

Other Information

Citation Subset: Index Medicus
Status: MEDLINE

Citation Network

2 Times Cited
10 Cited References
View Related Records
View Citation Map
Create Citation Alert
(data from Web of Science™ Core Collection)

View PubMed Related Articles

All Times Cited Counts
2 in All Databases
2 in Web of Science Core Collection
2 in BIOSIS Citation Index
0 in Chinese Science Citation Database
0 in Data Citation Index
0 in Russian Science Citation Index
0 in SciELO Citation Index

Usage Count
Last 180 Days: 0
Since 2013: 0
Learn more

Most Recent Citation
Ventura, P. A role for homocysteine increase in haemolysis of megaloblastic anaemias due to vitamin B-12 and folate deficiency: results from an in vitro experience. BIOCHIMICA ET BIOPHYSICA ACTA-MOLECULAR BASIS OF DISEASE, DEC 24 2004.
View All

This record is from: MEDLINE®

Suggest a correction
If you would like to improve the quality of the data in this record, please suggest a correction.