

CONCERNING PERIODIC POINTS IN MAPPINGS OF CONTINUA

W. T. INGRAM

(Communicated by Dennis Burke)

ABSTRACT. In this paper we present some conditions which are sufficient for a mapping to have periodic points.

THEOREM. *If f is a mapping of the space X into X and there exist subcontinua H and K of X such that (1) every subcontinuum of K has the fixed point property, (2) $f[K]$ and every subcontinuum of $f[H]$ are in class W , (3) $f[K]$ contains H , (4) $f[H]$ contains $H \cup K$, and (5) if n is a positive integer such that $(f|H)^{-n}(K)$ intersects K , then $n = 2$, then K contains periodic points of f of every period greater than 1.*

Also included is a fixed point lemma:

LEMMA. *Suppose f is a mapping of the space X into X and K is a subcontinuum of X such that $f[K]$ contains K . If (1) every subcontinuum of K has the fixed point property, and (2) every subcontinuum of $f[K]$ is in class W , then there is a point x of K such that $f(x) = x$.*

Further we show that: If f is a mapping of $[0, 1]$ into $[0, 1]$ and f has a periodic point which is not a power of 2, then $\lim\{[0, 1], f\}$ contains an indecomposable continuum. Moreover, for each positive integer i , there is a mapping of $[0, 1]$ into $[0, 1]$ with a periodic point of period 2^i and having a hereditarily decomposable inverse limit.

1. Introduction. In his book, *An Introduction to Chaotic Dynamical Systems* [3, Theorem 10.2, p. 62], Robert L. Devaney includes a proof of Sarkovskii's Theorem. Consider the following order on the natural numbers: $3 \triangleright 5 \triangleright 7 \triangleright \cdots \triangleright 2 \cdot 3 \triangleright 2 \cdot 5 \triangleright \cdots \triangleright 2^2 \cdot 3 \triangleright 2^2 \cdot 5 \triangleright \cdots \triangleright 2^3 \cdot 3 \triangleright 2^3 \cdot 5 \triangleright \cdots \triangleright 2^3 \triangleright 2^2 \triangleright 2 \triangleright 1$. Suppose $f: R \rightarrow R$ is continuous. If $k \triangleright m$ and f has a periodic point of prime period k , then f has a periodic point of period m . In working through a proof of this theorem for $k = 3$, the author discovered the main result of this paper—Theorem 2. For an alternate proof of Sarkovskii's Theorem for $k = 3$, see also [7]. For a further look at this theorem for ordered spaces see [13].

By a *continuum* we mean a compact connected metric space and by a *mapping* we mean a continuous function. By a *periodic point* of period n for a mapping f of a continuum M into M is meant a point x such that $f^n(x) = x$. The statement that x has prime period n means that n is the least integer k such that $f^k(x) = x$. A continuum M is said to have the *fixed point property* provided if f is a mapping of M into M there is a point x such that $f(x) = x$. A mapping f of a continuum X onto a continuum M is said to be *weakly confluent* provided for each subcontinuum K of

Received by the editors May 22, 1987 and, in revised form, September 14, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54F20, 54H20; Secondary 54F62, 54H25, 54F55.

Key words and phrases. Periodic point, fixed point property, class W , indecomposable continuum, inverse limit.

M some component of $f^{-1}(K)$ is thrown by f onto K . A continuum is said to be in Class W provided every mapping of a continuum onto it is weakly confluent. The continuum T is a *triod* provided there is a subcontinuum K of T such that $T - K$ has at least three components. A continuum is *atriodic* provided it does not contain a triod. A continuum M is *unicoherent* provided if M is the union of two subcontinua H and K , then the common part of H and K is connected. A continuum is *hereditarily unicoherent* provided each of its subcontinua is unicoherent. If f is a mapping of a space X into X , the *inverse limit* of the inverse limit sequence $\{X_i, f_i\}$ where, for each i , X_i is X and f_i is f will be denoted $\lim\{X, f\}$. For the inverse sequence $\{X_i, f_i\}$, the inverse limit is the subset of the product of the sequence of spaces X_1, X_2, \dots to which the point (x_1, x_2, \dots) belongs if and only if $f_i(x_{i+1}) = x_i$.

There has been considerable interest in periodic homeomorphisms of continua where a homeomorphism h is called periodic provided there is an integer n such that h^n is the identity. Wayne Lewis has shown [8] that for each n there is a chainable continuum with a periodic homeomorphism of period n . A theorem of Michel Smith and Sam Young [14] should be compared with Theorem 3 of this paper. Smith and Young show that if a chainable continuum M has a periodic homeomorphism of period greater than 2, then M contains an indecomposable continuum. In this paper we consider the question of the existence of periodic points in mappings of continua.

2. A fixed point theorem. The problem of finding a periodic point of period n for a mapping f is, of course, the same as the problem of finding a fixed point for f^n . Not surprisingly, we need a fixed point theorem as a lemma to the main theorem of this paper. The following theorem, which the author finds interesting in its own right, should be compared with an example of Sam Nadler [11] of a mapping with no fixed point of a disk to a containing disk. A corollary to Theorem 1 is the well-known corresponding result for mappings of intervals.

THEOREM 1. *Suppose X is a space, f is a mapping of X into X , and K is a subcontinuum of X such that $f[K]$ contains K . If (1) every subcontinuum of K has the fixed point property, and (2) every subcontinuum of $f[K]$ is in Class W , then there is a point x of K such that $f(x) = x$.*

PROOF. Since $f[K]$ is in Class W and K is a subset of $f[K]$, there is a subcontinuum K_1 of K such that $f[K_1] = K$. Then $f|_{K_1:K_1} \rightarrow K$ is weakly confluent since every subcontinuum of $f[K]$ is in Class W ; thus there is a subcontinuum K_2 of K_1 such that $f[K_2] = K_1$. Since K_1 is in Class W , $f|_{K_2:K_2} \rightarrow K_1$ is weakly confluent; therefore there is a subcontinuum K_3 of K_2 such that $f[K_3] = K_2$. Continuing this process there exists a monotonic decreasing sequence K_1, K_2, K_3, \dots of subcontinua of K such that $f[K_{i+1}] = K_i$ for $i = 1, 2, 3, \dots$. Let H denote the common part of all the terms of this sequence and note that $f[H] = H$, since $f[H] = f[\bigcap_{i>0} K_i] = \bigcap_{i>0} f[K_i] = \bigcap_{i>0} K_i = H$. Since $f|_H$ throws H onto H and H has the fixed point property, there exists a point x of H (and therefore of K) such that $f(x) = x$.

REMARK. Note that (1) and (2) of the hypothesis of Theorem 1 are met if $f[K]$ is chainable ([12, Theorem 4, p. 236 and 4], respectively), while (2) is met if $f[K]$ is

atriodic and acyclic [1] and (1) is met by planar, tree-like continua such that each two points of a subcontinuum L lie in a weakly chainable subcontinuum of L [10].

3. Periodic points. In this section we prove the main result of the paper.

THEOREM 2. *If f is a mapping of the space X into X and there exist subcontinua H and K of X such that (1) every subcontinuum of K has the fixed point property, (2) $f[K]$ and every subcontinuum of $f[H]$ are in class W , (3) $f[K]$ contains H , (4) $f[H]$ contains $H \cup K$, and (5) if n is a positive integer such that $(f|H)^{-n}(K)$ intersects K , then $n = 2$, then K contains periodic points of f of every period greater than 1.*

PROOF. Suppose $n \geq 2$. There is a sequence H_1, H_2, \dots, H_{n-1} of subcontinua of H such that $f[H_1] = K$ (note that $f|H$ is weakly confluent) and $f[H_{i+1}] = H_i$ for $i = 1, 2, \dots, n - 2$ (in case $n > 2$). There is a subcontinuum K_n of K so that $f[K_n] = H_{n-1}$. Thus, $f^n[K_n] = K$ and so $f^n[K_n]$ contains K_n , so, by Theorem 1, there is a point x of K_n such that $f^n(x) = x$. We must show that if $j < n$ then $f^j(x)$ is not x . If $j < n$ and $f^j(x) = x$, then $j = n - 2$ and x is in H_2 . Since $f^n(x) = x$ and $f^{n-2}(x) = x$, $f^2(x) = x$. Since x is in $(f|H)^{-2}(K)$, x is in $(f|H)^{-4}(K)$ and in K contrary to (5) of the hypothesis. Therefore, x is periodic of prime period n .

REMARK. If f is a mapping of the continuum M into itself and f has a periodic point of period k , then the mapping of $\text{lim}\{M, f\}$ induced by f has periodic points of period k , e.g. $(x, f^{k-1}(x), \dots, f(x), x, \dots)$. Thus, although Theorem 2 does not directly apply to homeomorphisms, it may be used to conclude the existence of homeomorphisms with periodic points.

COROLLARY. *If M is a chainable continuum, f is a mapping of M into M , and there are subcontinua H and K of M such that $f[K] = H$, $f[H]$ contains $H \cup K$, and if $(f|H)^{-n}(K)$ intersects K then $n = 2$ then f has periodic points of every period.*




FIGURE 1

EXAMPLE. Let f be the mapping of the simple triod T to itself given in [5]. The mapping f is represented in Figure 1 above. Letting $H = [0, A/2]$ and $K = [B/3, B/2]$ it follows from Theorem 2 that f has periodic points of every period.

EXAMPLE. Let f be the mapping of the simple triod T to itself given in [2]. The mapping f is represented in Figure 2 below. Letting $H = [0, 3B/8]$ and $K = [C/32, C/8]$, it follows from Theorem 2 that f has periodic points of every period.




FIGURE 2

EXAMPLE. Let f be the mapping of the unit circle S^1 to itself given by $f(z) = z^2$. Letting $H = \{e^{i\theta} | 0 \leq \theta \leq 3\pi/4\}$ and $K = \{e^{i\theta} | \pi \leq \theta \leq 3\pi/2\}$, it follows from Theorem 2 that f has periodic points of every period. Similarly, if f is a mapping of S^1 onto itself which is homotopic to z^n for some $n > 1$, then f has periodic points of every period.




FIGURE 3

COROLLARY. If f is a mapping of an interval to itself with a periodic point of period 3, then f has periodic points of every period.

PROOF. To see this it is a matter of noting that the hypothesis of Theorem 2 is met. We indicate the proof for one of two cases and leave the second similar case to the reader. Suppose a , b and c are points of the interval with $a < b < c$ and $f(a) = b$, $f(b) = c$ and $f(c) = a$ [the other case is $f(a) = c$, $f(b) = a$ and $f(c) = b$].

If $f^{-1}(c)$ is nondegenerate, then there exist mutually exclusive intervals H and K lying in $[b, c]$ and $[a, b]$, respectively, so that $f[H]$ is $[a, c]$ and $f[K]$ is $[b, c]$ and Theorem 2 applies.

Suppose $f^{-1}(c) = \{b\}$. Choose K lying in $[a, b]$ and H lying in $[b, c]$ so that $f[K] = [b, c]$ and $f[H] = [a, c]$. For each i , denote by H_i the set $(f[H])^{-1}(K)$. Note that a is not in H_i for $i = 1, 2, 3, \dots$ so c is not in H_i for $i = 2, 3, 4, \dots$ and thus b is not in H_i for $i = 3, 4, \dots$. Further, b is not in H_1 since c is not in K . Thus, if H_i intersects K , then $i = 2$. Consequently, the hypothesis of Theorem 2 is met.

REMARK. Condition (5) of Theorem 2 seems a bit artificial. A more natural condition the author experimented with in its place is a requirement that H and K be mutually exclusive. In fact, in each of the examples, the H and K given are mutually exclusive. However, replacing condition (5) with this proved to be undesirable in that the Sarkovskii Theorem for $k = 3$ is not a corollary to Theorem 2 if the alternate condition is used. That condition (5) may not be replaced by the assumption that H and K are mutually exclusive can be seen by the following. For the function $f: [0, 1] \rightarrow [0, 1]$, which is piecewise linear and contains the points $(0, \frac{1}{2})$, $(\frac{1}{2}, 1)$ and $(1, 0)$, there do not exist mutually exclusive intervals H and K such that $f[H]$ contains $H \cup K$ and $f[K]$ contains H . To see this suppose H and K are such mutually exclusive intervals. By Theorem 2, K contains a periodic point of f of period 3. Note that f^3 has only four fixed points: 0 , $\frac{1}{2}$, $\frac{2}{3}$, and 1 . Since $\frac{2}{3}$ is a fixed point for f , K must contain one of 0 , $\frac{1}{2}$, and 1 . We complete the proof by showing that each of these possibilities leads to a contradiction.

(1) Suppose 0 is in K . Then 1 is in H since $f^{-1}(0) = \{1\}$ and $f[H]$ contains K . But since $f^{-1}(1) = \{\frac{1}{2}\}$, $\frac{1}{2}$ is in both H and K .

(2) Suppose 1 is in K . Since $f^{-1}(1) = \{\frac{1}{2}\}$, $\frac{1}{2}$ is in H . Since $f^{-1}(\frac{1}{2}) = \{0, \frac{3}{4}\}$ and H and K do not intersect 0 is in H and $\frac{3}{4}$ is in K . But, $f^{-1}(0) = \{1\}$ so 1 is in H .

(3) Suppose $\frac{1}{2}$ is in K . As before, one of 0 and $\frac{3}{4}$ is in H . Since $f^{-1}(0) = \{1\}$, if 0 is in H then 1 is in both H and K . Thus $\frac{3}{4}$ is in H . Then $f^{-1}(\frac{3}{4})$ contains two points, $\frac{5}{8}$ and one less than $\frac{1}{2}$, so $P_1 = \frac{5}{8}$ is in H . Since $f^{-1}(P_1)$ contains two points, $\frac{1}{8}$ and one between $\frac{5}{8}$ and $\frac{3}{4}$, $\frac{1}{8}$ is in K . Thus, $f^{-1}(\frac{1}{8}) = \frac{15}{16}$ is in H . Since $f^{-1}(\frac{15}{16})$ contains two points, $\frac{17}{32}$ and one less than $\frac{1}{2}$, $P_2 = \frac{17}{32}$ is in H . Continuing this process, we get a sequence P_1, P_2, \dots of points of H which converges to $\frac{1}{2}$. Thus $\frac{1}{2}$ is in H .

4. Periodic points and indecomposability. In this section we show that under certain conditions the existence of a periodic point of period three in a mapping of a continuum M to itself implies that $\lim\{M, f\}$ contains an indecomposable continuum. Of course the result is not true in general since a rotation of S^1 by 120 degrees yields a homeomorphism of S^1 and a copy of S^1 for the inverse limit.

THEOREM 3. *Suppose f is a mapping of the continuum M into itself and x is a point of M which is a periodic point of f of period three. If M is atriodic and hereditarily unicoherent, then $\lim\{M, f\}$ contains an indecomposable continuum.*

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.