

**U.S. Food and Drug Administration**  
Protecting and Promoting *Your* Health

# Development of New Stereoisomeric Drugs

Publication Date: 5/1/1992

## I. INTRODUCTION AND BACKGROUND

Stereoisomers are molecules that are identical in atomic constitution and bonding, but differ in the three-dimensional arrangement of the atoms. For the purpose of this document, the stereoisomeric pairs of greatest interest are those with one or more asymmetric (chiral) centers whose enantiomers (individual stereoisomers) are mirror images. They have essentially identical physical (except for optical rotatory) and chemical (except in a chiral environment) properties.

This document focuses on issues relating to the study and pharmaceutical development of individual enantiomers and racemates. Such stereoisomers usually require specialized chiral techniques for their correct identification, characterization, separation and measurement. They are often readily distinguished by biological systems, however, and may have different pharmacokinetic properties (absorption; distribution, biotransformation, and excretion) and quantitatively or qualitatively different pharmacologic or toxicologic effects.

When stereoisomers are biologically distinguishable, they might seem to be different drugs, yet it has been past practice to develop racemates (i.e., compounds with 50:50 proportion of enantiomers). The properties of the individual enantiomers have not generally been well studied or characterized. Whether separated enantiomers should be developed was largely an academic question because commercial separation of racemates was difficult. Now that technological advances (large scale chiral separation procedures or asymmetric syntheses) permit production of many single enantiomers on a commercial scale, it is appropriate to consider what FDA's policy with respect to stereoisomeric mixtures should be. Development of racemates raises issues of acceptable manufacturing control of synthesis and impurities, adequate pharmacologic and toxicologic assessment, proper characterization of metabolism and distribution, and appropriate clinical evaluation.

It should be noted that the term "stereoisomers" is a general one for all isomers that differ only in the orientation of the atoms in space. Stereoisomers include not only the mirror image enantiomers, but also geometric (cis/trans) isomers and diastereoisomers (isomers of drugs with more than one chiral center that are not mirror images of one another). Diastereoisomers and geometric isomers are both chemically distinct and pharmacologically different (unless they are interconverted *in vivo*) and are generally readily separated without chiral techniques. Geometric isomers and diastereoisomers therefore should, with the rare exception of cases where *in vivo* interconversion occurs, be treated as separate drugs and developed accordingly. There is no reason to consider developing mixtures of geometric isomers or diastereoisomers unless they fortuitously represent a reasonable fixed dose combination (see 21 CFR 300.50). Even in that case, whether the optimal ratio of the two isomers is the ratio produced by an undirected or unmodified synthesis should be critically examined. In general, geometric isomers have been developed as single isomers. Practice with respect to diastereoisomers has been variable. These categories of stereoisomers will not be considered further in this document. Examination of cases in which the properties of enantiomers have been evaluated reveals instances in which both members had similar desirable activities:

- both enantiomers of dobutamine are positive inotropes;
- both ibuprofen enantiomers are anti-inflammatory agents;
- both enantiomers of warfarin and phenprocoumon are anticoagulants;
- the enantiomers of bupivacaine both produce local anesthesia;
- the enantiomers of the quinolones and the b-lactam antibiotics are all antibacterial instances in which one member of a pair was pharmacologically active and the other inactive (l-propranolol is a b-blocker; d-propranolol is not),
- the enantiomers had completely different activities (d-sotalol is a type 3 antiarrhythmic while l-sotalol is a b-blocker) or
- had different concentration-response relationships for a given property.

While inactivity of one member of a pair might be considered trivial, there are instances in which toxicity has been linked to one member of a pair of stereoisomers, not necessarily the active isomer (granulocytopenia is related to the d-isomer of levodopa; vomiting is caused by the d-isomer of levamisole; and myasthenia gravis symptoms were no longer observed when the d-isomer was removed from d,l-carnitine), and there are examples of an effect on the disposition of

one member of a pair by the other. In addition, there are many cases in which enantiomers have been shown to have different pharmacokinetic behavior. Differences in pharmacokinetic behavior may not pose a major therapeutic problem although it can make non-chiral blood level assays difficult to interpret with respect to activity and confuse interpretation of non-clinical data if the pharmacokinetic properties of the isomers in animals differ from those in humans.

While some enantiomeric pairs have had interesting and useful therapeutic properties (e.g., dl-sotalol, dl-dobutamine), there is no reason to expect the optimum ratio of the components to be the 1:1 ratio of a racemate (i.e., the dose response curves would not usually be expected to be congruent).

Despite the problems identified with some racemates, the common practice of developing racemates has resulted in few recognized adverse consequences. Although it is now technologically feasible to prepare purified enantiomers, development of racemates may continue to be appropriate. However, currently available information suggests that the following should be considered in product development:

1. Appropriate manufacturing and control procedures should be used to assure stereoisomeric composition of a product, with respect to identity, strength, quality and purity. Manufacturers should notify compendia of these specifications and tests.
2. Pharmacokinetic evaluations that do not use a chiral assay will be misleading if the disposition of the enantiomers is different. Therefore, techniques to quantify individual stereoisomers in pharmacokinetic samples should be available early. If the pharmacokinetics of the enantiomers are demonstrated to be the same or to exist as a fixed-ratio in the target population, an achiral assay or an assay that monitors one of the enantiomers may be used, subsequently.

## II. POLICY IN GENERAL

The stereoisomeric composition of a drug with a chiral center should be known and the quantitative isomeric composition of the material used in pharmacologic, toxicologic, and clinical studies known. Specifications for the final product should assure identity; strength, quality, and purity from a stereochemical viewpoint.

To evaluate the pharmacokinetics of a single enantiomer or mixture of enantiomers, manufacturers should develop quantitative assays for individual enantiomers in *in vivo* samples early in drug development. This will allow assessment of the potential for interconversion and the absorption, distribution, biotransformation, and excretion (ADBE) profile of the individual isomers. When the drug product is a racemate and the pharmacokinetic profiles of the isomers are different,

manufacturers should monitor the enantiomers individually to determine such properties as dose linearity and the effects of altered metabolic or excretory function and drug-drug interactions. If the pharmacokinetic profile is the same for both isomers or a fixed ratio between the plasma levels of enantiomers is demonstrated in the target population, an achiral assay or an assay that monitors one of the stereoisomers should suffice for later evaluation. In vivo measurement of individual enantiomers should be available to help assess toxicologic findings, but if this cannot be achieved, it would be sufficient in some cases to establish the kinetics of the isomers in humans.

Unless it proves particularly difficult, the main pharmacologic activities of the isomers should be compared in *in vitro* systems, in animals and/or in humans. A relatively benign toxicologic profile using the racemate would ordinarily support further development without separate toxicologic evaluation of the individual enantiomers. If, however, there are toxic findings other than those that are natural extensions of the pharmacologic effects of the drug, and especially if they are unusual or occur near the effective dose in animals or near the planned human exposure, toxicologic evaluation of the individual isomers in the study where the toxicity was detected should be undertaken.

FDA invites discussion with sponsors concerning whether to pursue development of the racemate or the individual enantiomer. All information developed by the sponsor or available from the literature that is relevant to the chemistry, pharmacology, toxicology, or clinical actions of the stereoisomers should be included in the IND and NDA submissions.

### III. CHEMISTRY, MANUFACTURING AND CONTROLS

The chemistry section of the application should contain the requisite information to assure the identity, quality, purity and strength of the drug substance and drug product. In addition, the following considerations should be taken into account when dealing with chiral drug substances and drug products.

#### Methods and Specifications

##### *Drug Substance*

Applications for enantiomeric and racemic drug substances should include a stereochemically specific identity test and/or a stereochemically selective assay method. The choice of the controls should be based upon the substance's method of manufacture and stability characteristics.

##### *Drug Product*

Applications for drug products that contain an enantiomer or racemic drug substance should include a stereochemically specific identity test and/or a stereochemically selective assay method. The choice of the controls should be based upon the product's composition, method of manufacture and stability characteristics.

## Stability

The stability protocol for enantiomeric drug substances and drug products should include a method or methods capable of assessing the stereochemical integrity of the drug substance and drug product. However, once it has been demonstrated that stereochemical conversion does not occur, stereoselective tests might not be needed.

## Labeling

The labeling should include a unique established name and a chemical name with the appropriate stereochemical descriptors.

## Pharmacology/Toxicology

### *Pharmacology*

The pharmacologic activity of the individual enantiomers should be characterized for the principal pharmacologic effect and any other important pharmacological effect, with respect to potency, specificity, maximum effect, etc.

### *Pharmacokinetic Profile*

To monitor in vivo interconversion and disposition, the pharmacokinetic profile of each isomer should be characterized in animals and later compared to the clinical pharmacokinetic profile obtained in phase 1.

### *Toxicology*

It is ordinarily sufficient to carry out toxicity studies on the racemate. If toxicity other than that predicted from the pharmacologic properties of the drug occurs at relatively low multiples of the exposure planned for clinical trials, the toxicity study where the unexpected toxicity occurred should be repeated with the individual isomers to ascertain whether only one enantiomer was responsible for the toxicity. If toxicity of significant concern can be eliminated by development of single isomer with the desired pharmacologic effect, it would in general be desirable to do so. The agency would be pleased to discuss any cases where questions exist regarding the definition of "significant toxicity."

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.