

Excipients and Their Use in Injectable Products

SANDEEP NEMA*, R. J. WASHKUH, and R. J. BRENDL

Mallinckrodt Medical, Incorporated, Saint Louis, Missouri

ABSTRACT: Formulation of a new drug product with excipients, that have been previously added to an approved injectable product, may save pharmaceutical companies developmental time and cost. The Physicians' Desk Reference (PDR) and Handbook on Injectable Drugs were reviewed, extracting all information on excipients. The information was consolidated into eight tables, categorizing excipients as 1) Solvents and Co-solvents, 2) Solubilizing, Wetting, Suspending, Emulsifying or Thickening agents, 3) Chelating Agents, 4) Antioxidants and Reducing Agents, 5) Antimicrobial Preservatives, 6) Buffers and pH Adjusting Agents, 7) Bulking Agents, Protectants, and Tonicity Adjustors, and 8) Special Additives. Where applicable, tables list frequency of use, concentration, and an example of a commercial product containing the excipient. Excipients which are included in the 1996 FDA 'Inactive Ingredient Guide,' but do not appear in the PDR or Handbook on Injectable Drugs, were included as a separate list.

Introduction

Injectable products require a unique formulation strategy. The formulated product has to be sterile, pyrogen free and, in the case of solutions, free of particulate matter. Preferably, the formulation will be isotonic, and depending on the route of administration (for instance, for intra-spinal or intracisternal routes), antioxidants and preservatives may not be allowed. For a given drug, the risk of adverse events is higher if it is administered as an injection versus a non-parenteral route. The requirement for sterility demands that the excipients be able to withstand autoclaving or other sterilization processes. These factors limit the choice of excipients available to the formulators.

Generally, a knowledge of which excipients have been deemed safe by the FDA or are already present in a marketed product provides increased assurance to the formulator that these excipients will probably be safe for their new drug product. However, there is no guarantee that the new drug product will be safe as excipients are combined with other additives and/or with a new drug, creating unforeseen potentiation or synergistic toxic effects. Regulatory bodies may view an excipient previously approved in an injectable dosage form favorably, and will frequently require less safety data. A new additive in a formulated product will always require additional studies adding to the cost and timeline of product development.

The purpose of this paper is to present the various excipients that have been included in the formulation of injectable products marketed in the USA. This information is not readily available. A literature search indicates that the last paper dealing with this was published in 1980 (1). Products approved outside the US are not covered in this

review. Also, sterile dosage forms not administered parenterally, such as solutions for irrigation, ophthalmic or otic drops, and ointments were excluded.

Methodology

Physicians' Desk Reference published in 1994 & 1996 (2, 3), and Handbook on Injectable Drugs (4) were used as the primary source of information. Entries on all injectable drugs were summarized in an Excel worksheet. Each product was classified by Manufacturer, Trade name, Drug name, Route of Administration, SVP/LVP, pH of Product, Solvent Used, Solubilizing/Suspending Agent, Preservative, Antioxidant, Chelator and Other Formulation Additives.

The resulting Excel sheet had information on more than 700 products. This information was condensed into easy-to-read tables. Each table has been categorized based on the primary function of excipient in the formulation. For example, citrates are classified as buffers and not as chelating agents, and ascorbates are categorized as antioxidants, although they can serve as buffers. This classification system was based on our experience in formulation development and on the published literature. Such simplification avoids duplication of entries and provides the audience with easy-to-read tables.

Some duplication was unavoidable. Tables VII and VIII contain some excipients which may have also been listed in the first six tables. Whenever the reference specifically designated a specific function to an ingredient it was re-listed in Tables VII and VIII. For example, glycine can be used as a buffer or as a stabilizing (protecting) agent. Therefore, glycine is listed in Tables VI and VII. Methyl paraben is a preservative (Table V) but also has a special function in Adriamycin RDF[®] formulation (Table VIII).

The concentration of excipients is listed as percentages weight by volume (w/v) or volume by volume (v/v). If the product was listed as lyophilized or powder, these percent-

Received October 1, 1996. Accepted for publication May 16, 1997.

*Author to whom correspondence should be addressed: P.O. Box 5840, St. Louis, MO, 63134

TABLE I
Solvents and Co-solvents

| Excipient | Frequency | Range | Example |
|-----------------------|-----------|-------------|---|
| Benzyl Benzoate | 2 | 20% v/v | Depo-Testosterone® (Upjohn) 20% v/v |
| Cottonseed Oil | 1 | 73.6% w/v | Depo-Testosterone® (Upjohn) 73.6% w/v |
| N,N Dimethylacetamide | 1 | 6% w/v | Vumon® (Bristol Myers) 6% w/v |
| Ethanol | 24 | 0.6-80% | Prograf® (Fujisawa) 80% v/v |
| Glycerin (Glycerol) | 9 | 1.6-70% w/v | Multitest CMI® (Connaught) 70% w/v |
| Peanut oil | 1 | * | Bal in Oil® (Becton Dickinson) |
| Polylethylene glycol | | | |
| PEG | 4 | 0.15-50% | Secobarbital sodium (Wyeth-Ayerst) 50% |
| PEG 300 | 2 | 50-65% | VePesid® (Bristol Myers) 65% w/v |
| PEG 400 | 2 | * | Ativan® (Wyeth-Ayerst) |
| PEG 3350 | 5 | 0.3-3% | Depo-Medrol® (Upjohn) 2.95% w/v |
| Poppyseed oil | 1 | 1% | Ethiodol® (Savage) 1% |
| Propylene Glycol | 25 | 0.2-75.2% | Terramycin Solution (Roerig) 75.2% |
| Safflower oil | 2 | 5-10% | Liposyn II® (Abbott) 10% |
| Sesame oil | 6 | * | Solganal Inj.® (Schering) |
| Soybean oil | 4 | 5-20% w/v | Intralipid® (Clintec) 20% |
| Vegetable oil | 2 | * | Virtion IM Inj.® (Star Pharmaceuticals) |

* No data available.

ages were derived based on the reconstitution volume commonly used. The tables list the range of concentration used, typical or most common concentration employed, and examples of products containing the excipient, specifically those which use extremely low or high concentrations.

Discussions

Table I list solvents and co-solvents used in parenteral products. Water for injection is the most common solvent but may be combined or substituted with a co-solvent to improve the solubility or stability of drugs. Oils like safflower and soybean are used in total parenteral nutrition products where they serve as a fat source and as carriers for fat-soluble vitamins. Ethanol and propylene glycol are used, either alone or in combination with other solvents, in more than 50% of parenteral co-solvent systems. It is surprising to see propylene glycol used more often than polyethylene

glycols (PEGs) in spite of its higher myotoxicity and hemolyzing effects (5, 6). Probably, the presence or generation of peroxides in PEGs is a major limitation.

Table II includes a broad category of excipients whose function in formulation could be—(1) Viscosity imparting or suspending agents like carboxy methyl cellulose, sodium carboxy methyl cellulose, sorbitol, acacia, Povidone, hydrolyzed gelatin; (2) Solubilizing, wetting or emulsifying agents like Cremophore EL, sodium desoxycholate, Polysorbate 20 or 80, PEG 40 castor oil, PEG 60 castor oil, sodium dodecyl sulfate, lecithin or egg yolk phospholipid; (3) Aluminum monostearate which is added to fixed oil to form viscous or gel-like suspending medium. Polysorbate 80 is the most common and versatile solubilizing, wetting and emulsifying agent.

Only a limited number of chelating agents are used in parenteral products (Table III). They serve to complex heavy

TABLE II
Solubilizing, Wetting, Suspending, Emulsifying or Thickening Agents

| Excipient | Frequency | Range | Example |
|--|-----------|--------------|--------------------------------------|
| Acacia | 2 | 7% | Tuberculin Old Test® (Lederle) 7% |
| Aluminum monostearate | 1 | 2% | Solganal Inj.® (Schering) 2% |
| Carboxy methyl cellulose | 4 | 1% | Bicillin® (Wyeth-Ayerst) 0.55% |
| Carboxy methyl cellulose, sodium | 9 | 0.1-0.75% | Lupron Depot® (TAP) 0.75% w/v |
| Cremophore EL* | 3 | 50-65% w/v | Sandimmune® (Sandoz) 65% w/v |
| Desoxycholate sodium | 1 | 0.4% w/v | Fungizone® (Bristol Myers) 0.41% w/v |
| Egg yolk phospholipid | 3 | 1.2% | Intralipid® (Clintec) 1.2% |
| Gelatin, Hydrolyzed | 1 | 16% w/v | Contone® (Merck) 16% w/v |
| Lecithin | 7 | 0.4-1.2% w/v | Diprivan® (Zeneca) 1.2% w/v |
| Polyoxyethylated fatty acid | 1 | 7% w/v | AquaMephyton® (Merck) 7% w/v |
| Polysorbate 80 (Tween 80) | 31 | 0.01-12% | Cardarone X Lv.® (Wyeth-Ayerst) 10% |
| Polysorbate 20 (Tween 20) | 5 | 0.01-0.4% | Calcijex® (Abbott) 0.4% w/v |
| PEG 40 castor oil** | 1 | 11.5% v/v | Monistat® (Janssen) 11.5% v/v |
| PEG 60 castor oil*** | 1 | 20% w/v | Prograf® (Fujisawa) 20% w/v |
| Povidone (Polyvinyl pyrrolidone) | 6 | 0.5-0.6% w/v | Bicillin® (Wyeth-Ayerst) 0.6% w/v |
| Sodium dodecyl sulfate (Na lauryl sulfate) | 1 | 0.018% w/v | Proleukin® (Cetus) 0.018% w/v |
| Sorbitol | 3 | 25-50% | Aristrospan® (Fujisawa) 50% w/v |

* Cremophor EL: Etoas 35, polyoxyethylated castor oil, polyoxyethylene 35 castor oil.

** PEG 40 castor oil; polyoxyl 40 castor oil, castor oil POE-40, Crodarec 40, polyoxyethylene 40 castor oil, Protachem CA-40.

*** PEG 60 hydrogenated castor oil; Cremophor RH 60, hydrogenated castor oil POE-60, Protachem CAH-60.

TABLE III
Chelating Agents

| Excipient | Frequency | Range | Example |
|------------------------|-----------|-----------|---------------------------------|
| Calcium disodium EDTA* | 9 | 0.01-0.1% | Wydase® (Wyeth-Ayerst) 0.1% w/v |
| Disodium EDTA | 34 | 0.01-0.1% | Calcijex® (Abbott) 0.11% w/v |
| Sodium EDTA | 1 | 0.20% | Folvite® (Lederle) 0.2% |
| DTPA** | 1 | 0.01% | Magnevist® (Berlex) 0.04% |

* EDTA = Ethylenediaminetetraacetic acid.

** DTPA = Diethylenetriaminepentaacetic acid; Pentetic acid.

metals and therefore can improve the efficacy of antioxidants or preservatives. In our opinion, calcium EDTA has an advantage over tetrasodium salt by not contributing sodium and not chelating calcium from the blood.

An antioxidant as a class is defined as those compounds that can act as reducing agents or may serve as free radical scavengers. Table IV summarizes the antioxidants, their frequency of use, concentration range and examples of products containing them. Sulfite, bisulfite, and metabisulfite constitute the majority of antioxidants used in parenteral products despite several reports of incompatibilities and

toxicity (7, 8). Butylated hydroxy anisole, butylated hydroxy toluene and propyl gallate are primarily used in semi/non-aqueous vehicles because of their low aqueous solubility. Ascorbic acid/sodium ascorbate may serve as an antioxidant, buffer, and chelating agent in the same formulation.

Benzyl alcohol was the most common antimicrobial preservative present in parenteral formulations (Table V). This is consistent with other surveys (9). Parabens are the next most common preservatives. Thirty-nine products had a combination of methyl and propyl parabens; eleven had only methyl, and one had only propyl paraben. Thimerosal was surprisingly common, especially in vaccines, even though some individuals have sensitivity to mercurics. Chlorocresol is purported to be a good preservative for parenterals, but our survey did not find any examples of commercial products containing chlorocresol.

Table VI lists buffers and chemicals used to adjust the pH of formulations. Phosphate, citrate, and acetate are the most common buffers used in parenteral products. Mono and diethanolamine are added to adjust pH and form corresponding salts. Hydrogen bromide, sulfuric acid, benzene sulfonic acid and methane sulfonic acids are added to drugs which are bromide (Scopolamine HBr, Hyoscine HBr, UDL), sulfate (Nebcin, Tobramycin sulfate, Lilly), besylate

TABLE IV
Antioxidants and Reducing Agents

| Excipient | Frequency | Range | Example |
|---|-----------|-------------------|---------------------------------------|
| Acetone sodium bisulfite | 4 | 0.2-0.4% w/v | Novocaine® (Sanofi-Winthrop) 0.4% w/v |
| Ascorbate (sodium/acid) | 7 | 0.1-4.8% w/v | Vibramycin® (Roerig) 4.8% w/v |
| Bisulfite sodium | 28 | 0.02-0.66% w/v | Amikin® (Bristol Myers) 0.66% w/v |
| Butylated hydroxy anisole (BHA) | 3 | 0.00028-0.03% w/v | Aquasol® (Astra) 0.03% |
| Butylated hydroxy toluene (BHT) | 3 | 0.00116-0.03% w/v | Aquasol® (Astra) 0.03% |
| Cystein/Cysteinate HCl | 2 | 0.07-0.10% w/v | Aethar Gel® (Rhône-Poulanc) 0.1% w/v |
| Dithionite sodium (Na hydrosulfite, Na sulfite oxylate) | 1 | 0.10% | Numorphan® (DuPont) 0.10% |
| Gentisic acid | 1 | 0.02% w/v | OctreoScan® (Mallinckrodt) |
| Gentisic acid ethanolamine | 1 | 2% | M.V.L. 12® (Astra) 2% |
| Glutamate monosodium | 2 | 0.1% w/v | Varivas® (Merck) 0.1% w/v |
| Formaldehyde sulfoxylate sodium | 9 | 0.075-0.5% w/v | Terramycin Solution (Roerig) 0.5% w/v |
| Metabisulfite potassium | 1 | 0.10% | Vasoxyl® (Glaxo-Wellcome) 0.10% |
| Metabisulfite sodium | 29 | 0.02-1% w/v | Intropin® (DuPont) 1% w/v |
| Monothioglycerol (Thioglycerol) | 6 | 0.1-1% | Terramycin Solution (Roerig) 1% |
| Propyl gallate | 2 | 0.02% | Navane® (Roerig) |
| Sulfite sodium | 7 | 0.05-0.2% w/v | Enion® (Ohmeda) 0.2% w/v |
| Thioglycolate sodium | 1 | 0.66% w/v | Sus-Pirine® (Forest) 0.66% w/v |

TABLE V
Antimicrobial Preservatives

| Excipient | Frequency | Range | Example |
|------------------------------------|-----------|-------------------|--|
| Benzalkonium chloride | 1 | 0.02% w/v | Celestone Soluspan® (Schering) 0.02% w/v |
| Benzethonium chloride | 4 | 0.01% | Benadryl® (Parke-Davis) 0.01% w/v |
| Benzyl alcohol | 74 | 0.75-5% | Dimenhydrinate® (Steris) 5% |
| Chlorobutanol | 17 | 0.25-0.5% | Codine phosphate (Wyeth-Ayerst) 0.5% |
| m-Cresol | 3 | 0.1-0.3% | Humatrope® (Lilly) 0.30% |
| Myristyl gamma-picolinium chloride | 2 | 0.0195-0.169% w/v | Depo-Provera® (Upjohn) 0.169% w/v |
| Paraben methyl | 50 | 0.05-0.18% | Inapsine® (Janssen) 0.18% w/v |
| Paraben propyl | 40 | 0.01-0.1% | Xylocaine w/Epinephrine (Astra) 0.1% w/v |
| Phenol | 48 | 0.2-0.5% | Calcimar® (Rhône Poulanc) 0.5% w/v |
| 2-Phenoxyethanol | 3 | 0.50% | Havrix® (SmithKline Beecham) 0.50% w/v |
| Phenyl mercuric nitrate | 3 | 0.001% | Antivenin® (Wyeth-Ayerst) 0.001% |
| Thimerosal | 46 | 0.003-0.01% | Atgam® (Upjohn) 0.01% |

TABLE VI
Buffers and pH Adjusting Agents

| Excipient | Example |
|------------------------|--|
| Acetate | |
| Sodium | Miacalcin Injection® (Sandoz) |
| Acetic acid | Miacalcin Injection® (Sandoz) |
| Glacial acetic acid | Brevibloc Injection® (Ohmeda) |
| Ammonium | Bumex Injection® (Roche) |
| Ammonium hydroxide | Triostat Injection® (SmithKline Beecham) |
| Benzene sulfonic acid | Tracrium Injection® (Glaxo-Wellcome) |
| Benzoate Sodium/sodium | Valium Injection® (Roche) |
| Bicarbonate Sodium | Cefotan Injection® (Zeneca) |
| Carbonate Sodium | HypoRho-D® (Bayer) |
| Citrate | |
| Acid | DTIC-Dome® (Bayer) |
| Sodium | Ceredase® (Genzyme) |
| Disodium | Cerezyme® (Genzyme) |
| Trisodium | Cerezyme® (Genzyme) |
| Diethanolamine | Bacrim IV® (Roche) |
| Glucosyl delta lactone | Quinidine® (Lilly) |
| Glycine | Hep-B Gammax® (Merck) |
| Hydrochloric acid | Amicar® (Immunex) |
| Hydrogen bromide | Scopolamine (UDL) |
| Lactate acid/Sodium | Fentanyl citrate & Droperidol (Astra) |
| Lysine | Eminase Injection® (Roberts) |
| Maleic acid | Librium Injection® (Roche) |
| Methanesulfonic acid | DHE-45 Injection® (Sandoz) |
| Monoethanolamine | Terramycin Solution (Roerig) |
| Phosphate | |
| Acid (phosphoric) | Humegon® (Organon) |
| Monobasic potassium | Zantac Injection® (Glaxo-Wellcome) |
| Monobasic sodium* | Pregnyl® (Organon) |
| Dibasic sodium** | Prolastin® (Bayer) |
| Tribasic sodium | Synthroid® (Knoll) |
| Sodium hydroxide | Optiray® (Mallinckrodt) |
| Sulfuric acid | Nebcin® (Lilly) |
| Tartrate acid/sodium | Methergine Injection® (Sandoz) |
| Tromethamine | Optiray® (Mallinckrodt) |

* Sodium biphosphate, Sodium dihydrogen phosphate or Na dihydrogen orthophosphate.

** Sodium phosphate, Disodium hydrogen phosphate.

(Tracrium Inj., Atropium besylate) or mesylate (DHE 45 Injection, Dihydroergotamine mesylate) salts. Glucosyl delta lactone is used to adjust the pH of Quinidine gluconate (Lilly). Benzoate buffer, at a concentration of 5%, is used in Valium Injection. Citrates are common buffers that can have a dual role as chelating agents. Lysine and glycine are amino acids which function as buffers and stabilize protein and peptide formulations. These amino acids are also used as lyo-additives and may prevent cold denaturation. Lactate and tartrate are occasionally used as buffer systems.

Table VII lists additives which are used to modify osmolality, and as bulking or lyo-cryo protective agents. Dextrose and sodium chloride are used to adjust tonicity in the majority of formulations. Some amino acids, glycine, alanine, histidine, imidazole, arginine, asparagine, aspartic acid, are used as bulking agents for lyophilization and may serve as stabilizers for proteins or peptides and as buffers. Monosaccharides (dextrose, glucose, lactose), disaccharide (sucrose), polyhydric alcohols (inositol, mannitol, sorbitol), glycol (PEG 3350), Povidone (polyvinylpyrrolidone), and proteins (albumin, gelatin) are commonly used as lyo-additives.

TABLE VII
Bulking Agents, Protectants, and Tonicity Adjustors

| Excipient | Example |
|--------------------------|--|
| Alanine | Thrombate III® (Bayer) |
| Albumin | Bioclote® (Arco) |
| Albumin human | Botax® (Allergan) |
| Amino acids | Havrix® (SmithKline Beecham) |
| L-Arginine | Activase® (Genentech) |
| Asparagine | Tice BCG® (Organon) |
| L-Aspartic acid | Pepcid® (Merck) |
| Calcium chloride | Phenergan Injection® (Wyeth-Ayerst) |
| Citric acid | Sensorcaine-MPF® (Astra) |
| Dextrose | Betaseron® (Berlex) |
| Gelatin hydrolyzed | Acthar® (Rhône-Poulanc Rorer) |
| Glucose | Iveegam® (Immuno-US) |
| Glycerin | Tice BCG® (Organon) |
| Glycine | Atgam Injection® (Upjohn) |
| Histidine | Antihemophilic Factor, human (Am. Red Cross) |
| Imidazole | Helixate® (Armour) |
| Inositol | OcraoScan® (Mallinckrodt) |
| Lactose | Caverject® (Upjohn) |
| Magnesium chloride | Terramycin Solution® (Roerig) |
| Magnesium sulfate | Tice BCG® (Organon) |
| Mannitol | Elspar® (Merck) |
| Polyethylene glycol 3350 | Bioclote® (Arco) |
| Polysorbate 80 | Helixate® (Armour) |
| Potassium chloride | Varivax® (Merck) |
| Povidone | Alkeran® (Glaxo-Wellcome) |
| Sodium chloride | WinRho SD® (Univax) |
| Sodium succinate | Actimmune® (Genentech) |
| Sodium sulfate | Depo-Provera® (Upjohn) |
| Sorbitol | Panhematin® (Abbott) |
| Sucrose | Prolastin® (Bayer) |

Special Additives

These additives have been included in pharmaceutical formulation to serve specific functions (Table VIII). Below is a summary of the special additives along with their intended use—

- (1) Calcium gluconate injection (American Regent) is a saturated solution of 10% w/v; calcium d-saccharate tetrahydrate 0.46% w/v is added to prevent crystallization during temperature fluctuations.
- (2) Cipro IV® (Ciprofloxacin, Bayer) contains lactic acid as a solubilizing agent for the antibiotic.
- (3) Premarin Injection® (Conjugated Estrogens, Wyeth-Ayerst Labs) is a lyophilized product that contains simethicone to prevent formation of foam during reconstitution.
- (4) Dexamethasone acetate (Dalaone DP, Forest, Decadron-LA, Merck, Dalaone DP Injection, UAD Labs) and Dexamethasone Na phosphate (Merck) are available as suspension or solution. These dexamethasone formulations contain creatine or creatinine as an additive.
- (5) Adriamycin RDE® (Doxorubicin hydrochloride, Pharmacia) contains methyl paraben, 0.2 mg/mL, to increase dissolution (10).
- (6) Ergotrate maleate (Ergonovine maleate, Lilly) contains 0.1% ethyl lactate as a solubilizing agent.
- (7) Estradurin Injection® (Polyestradiol phosphate, Wyeth-Ayerst Labs) uses Nicotinamide (12.5 mg/ml)

TABLE VIII
Special Additives

| Excipient | Example |
|--|--|
| Acetyl tryptophanate | Human Albumin (American Red Cross) |
| Aluminum hydroxide | Recombinant HB [®] (Merck) |
| Aluminum phosphate | Tetanus Toxoid Adsorbed [®] (Lederle) |
| Aluminum potassium sulfate | TD Adsorbed Adul [®] (Connaught) |
| E-Aminocaproic acid | Eminase [®] (Roberts) |
| Calcium d-saccharate | Calcium Gluconate (American Regent) |
| Caprylate sodium | Human Albumin (American Red Cross) |
| 8-Chlorotheophylline | Dimenhydrinate (Steris) |
| Creatine | Dalalone DP [®] (Forest) |
| Creatinine | Hydrocortone Phosphate (Merck) |
| Diatrizoic acid | Conray (Mallinckrodt) |
| Gamma Cyclodextrin | Cardiotec (Squibb) |
| Ethyl lactate | Ergostrate maleate [®] (Lilly) |
| Ethylenediamine | Aminophylline [®] (Abbott) |
| L-Glutamate sodium | Kabikinate [®] (Pharmacia) |
| Iron ammonium citrate | Tice BCG [®] (Oganon) |
| Lactic acid | Cipro IV [®] (Bayer) |
| D,L-Lactic and Glycolic acid copolymer | Zoladex [®] (Zeneca) |
| Maltose | Gamimune [®] (Bayer) |
| Meglumine | Magnevist [®] (Berlex) |
| Niacinamide | Estradurin [®] (Wyeth-Ayerst) |
| Paraben methyl | Adriamycin RDF [®] (Pharmacia) |
| Protamine | Insulatard NPH [®] (Novo Nordisk) |
| Simethicone | Premarin Injection [®] (Wyeth-Ayerst) |
| Sodium saccharin | Compazine Injection [®] (Smith-Kline Beecham) |
| Tri-n-butyl phosphate | Venoglobulin [®] (Apha Therapeutic) |
| von Willebrand factor | Bioclate [®] (Arco) |
| Zinc | Lente Insulin [®] (Novo Nordisk) |

as a solubilizing agent. Hydetrasol[®] (Merck) also contains niacinamide.

- (8) Aluminum in the form of aluminum hydroxide, aluminum phosphate or aluminum potassium sulfate is used as adjuvant in various vaccine formulations to elicit an increased immunogenic response.
- (9) Zoladex[®] (Goserelin acetate, Zeneca) is administered subcutaneously as microspheres. These spheres are made of D,L-lactic and glycolic acid copolymer. Lupron Depot Injection[®] (TAP) are lyophilized microspheres of gelatin and glycolic-lactic acid for intramuscular injection.
- (10) Gamma cyclodextrin is used as a stabilizer in Cardiotec[®] at a concentration of 50 mg/mL.
- (11) Sodium caprylate (sodium octoate) has antifungal properties, but it is also used to improve the stability of albumin solution against effects of heat. Albumin solution can be heat pasteurized by heating at 60°C for 10 hours in the presence of sodium caprylate. Acetyl tryptophanate sodium is also added to albumin formulations.
- (12) Meglumine (N-methylglucamine) is used as an ex-

TABLE IX

List of Excipient from 1996 FDA 'Inactive Ingredient Guide'

| | |
|---|--|
| Ammonium sulfate | Pentetate (DTPA) calcium trisodium |
| Benzyl chloride | Poloxamer 165 |
| Butyl paraben | PEG 4000 |
| Calcium sodium | PEG 600 |
| Calciferol calcium | Polyglactin |
| Castor oil | Poly lactide |
| Cellulose (microcrystalline) | Polyoxyethylene fatty acid esters |
| Cholesterol | Polyoxyethylene sorbitan monosterate |
| Deoxycholic acid | Polyoxyl 35 Castor oil |
| Diatrizoic acid | Polysorbate 40 |
| Dicyclohexyl carbodiimide | Polysorbate 85 |
| Diethyl amine | Potassium hydroxide |
| Dimyristoyl lecithin | Potassium phosphate, dibasic |
| Dimyristoyl phosphatidyl-glycerol | Sodium bisulfate |
| Disofenin | Sodium chloride |
| Docusate sodium | Sodium hypochlorite |
| Edamine | Sodium iodide |
| Exemetazine | Sodium pyrophosphate |
| Glucopate sodium | Sodium thiosulfate, anhydrous |
| Glucopate calcium | Sodium trimetaphosphate |
| Glucuronic acid | Sorbitan monopalmitate |
| Guanidine HCl | Stannous chloride |
| Iofetamine HCl | Stannous fluoride |
| Lactobionic acid | Stannous tartrate |
| Lecithin hydrogenated soy | Starch |
| Lidofenin | Succimer |
| Medrofenin | Succinic acid |
| Medronate disodium | Sulfurous acid |
| Medronic acid | Tetrakis (1-isocyno-2-methoxy-2,methyl-propante) copper (I) Te |
| Methyl boronic acid | Thiazosimic acid |
| Methyl cellulose | Trihiazosimic acid |
| Methylene blue | Urea |
| N-(carbamoyl-methoxy polyethylene-glycol 2000)-1,2-distearoyl | Zinc acetate |
| N-2-hydroxyethyl piperazine N'-2' ethane sulphonic acid | Zinc chloride |
| Nioxime | Zinc oxide |
| Nitric acid | 2-ethyl hexanoic acid |
| Oxyquinoline | PEG vegetable oil |

- (13) Surprisingly, sodium saccharine is used in Stelazine[®] and Compazine[®] formulations; our guess is that it serves as a stabilizer and tonicity adjuster.
- (14) Tri-n-butyl phosphate is present as an excipient in human immune globulin solution (Venoglobulin[®]). Its exact function in the formulation is not known, but it may serve as a scavenging agent.
- (15) von Willebrand factor is used to stabilize recombinant antihemophilic factor (Bioclate[®]).
- (16) Maltose serves as a tonicity adjuster and stabilizer in immune globulin formulation (Gamimune N[®]).
- (17) Epsilon amino caproic acid (6-amino hexanoic acid) is used as a stabilizer in anistreplase (Eminase injection[®]).
- (18) Zinc and protamine have been added to insulin to form complexes and control the duration of action.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.