PHYSICAL PHARMACY

ALFRED MARTIN



Physical Pharmacy

PHYSICAL CHEMICAL PRINCIPLES IN THE PHARMACEUTICAL SCIENCES

Alfred Martin, Ph.D.

Emeritus Coulter R. Sublett Professor Drug Dynamics Institute, College of Pharmacy, University of Texas

with the participation of PILAR BUSTAMANTE, Ph.D.

Titular Professor
Department of Pharmacy
and Pharmaceutical Technology,
University Alcala de Henares,
Madrid, Spain

and with illustrations by
A. H. C. CHUN, Ph.D.
Associate Research Fellow
Pharmaceutical Products Division,
Abbott Laboratories



B. I. Waverly Pvt Ltd New Delhi



| B. I. Waverly Pvt Ltd 54 Janpath, New Delbi - 110 001 |
|---|
| . Reprint authorised by Waverly International |
| Copyright © 1993 Waverly International, 428 East Preston Street, Baltimore, Maryland 2102-3993 USA |
| Indian Reprint 1994 Reprint 1995 |
| All rights reserved. This book is protected by copyright. No part of this book may be reproduced in any form or by any means, including photocopying or utilized by any information storage and retrieval system without written permission from the copyright owner. Violators will be prosecuted. |
| This edition is for sale in India, Bangladesh, Nepal, Bhutan and Maldives only. |
| |
| ISBN 81-7431-001-0 |
| Price Rs. 495.00 |
| Published in India by B.I. Waverly Pvt Ltd, 54 Janpath, New Delhi - 110 001 and printed at United India Press, New Delhi. |



2 States of Matter

Binding Forces Between Molecules States of Matter The Gaseous State The Liquid State Solids and the Crystalline State
The Liquid Crystalline State
Phase Equilibria and the Phase Rule
Thermal Analysis

BINDING FORCES BETWEEN MOLECULES

In order for molecules to exist in aggregates in gases, liquids, and solids, intermolecular forces must exist. An understanding of intermolecular forces is important in the study of pharmaceutical systems and follows logically from the previous discussion of intramolecular bonding energies. Cohesion, or the attraction of like molecules, and adhesion, or the attraction of unlike molecules, are manifestations of intermolecular forces. A knowledge of these forces is important for an understanding not only of the properties of gases, liquids, and solids, but also of interfacial phenomena, flocculation in suspensions, stabilization of emulsions, compaction of powders in capsules, and the compression of granules to form tablets.

Repulsive and Attractive Forces. When molecules interact, both repulsive and attractive forces operate. As two molecules are brought close together, the opposite charges in the two molecules are closer together than the like charges and cause the molecules to attract one another. When the molecules are brought so close that the outer charge clouds touch, the molecules repel each other like rigid elastic bodies.

Thus attractive forces are necessary in order that molecules cohere; repulsive forces are necessary in order that the molecules do not interpenetrate and annihilate one another. Moelwyn-Hughes¹ points to the analogy between human behavior and molecular phenomenon. Just as the actions of humans are often influenced by a conflict of loyalties, so molecular behavior is governed by attractive and repulsive forces.

Repulsion is due to the interpenetration of the electronic clouds of molecules and increases exponentially with a decrease in distance between the molecules. At a certain equilibrium distance, about 3 or 4×10^{-8} cm (3 or 4 angstroms), the repulsive and attractive

forces are equal. At this position, the potential energy of the two molecules is a minimum and the system is most stable (Fig. 2-1). This principle of minimum potential energy applies not only to molecules but to atoms and to large objects as well.

Under the following headings are discussed the various types of attractive intermolecular forces.

Van der Waals Forces. Dipolar molecules frequently tend to align themselves with their neighbors, so that the negative pole of one molecule points toward the positive pole of the next. Thus, large groups of molecules may be associated through weak attractions known as dipole-dipole or Keesom forces. Permanent dipoles are capable of inducing an electric dipole in nonpolar molecules (which are easily polarizable) in order to produce dipole-induced dipole, or Debye, interactions, and nonpolar molecules can induce polar-




Fig. 2-1. Repulsive and attractive energies and net energy as a function of the distance between molecules. Note that a minimum occurs in the net energy because of the different character of the attraction and repulsion curves.

ZZ



5

Solutions of Nonelectrolytes

Concentration Expressions Equivalent Weights Solutions of Nonelectrolytes Ideal and Real Solutions Colligative Properties Molecular Weight Determination

Materials may be mixed together to form a true solution, a colloidal solution, or a coarse dispersion. A true solution is defined as a mixture of two or more components that form a homogeneous molecular dispersion, in other words, a one-phase system, the composition of which can vary over a wide range. The terms in this definition warrant further comment, and an attempt at clarification is made in the following paragraphs.

A system is a bounded space or a definite quantity of substance that is under observation and experimentation. Under some circumstances, the system may consist only of radiant energy or an electric field, containing no material substances. The term phase has already been defined in Chapter 2 as a distinct homogeneous part of a system separated by definite boundaries from other parts of the system. Each phase may be consolidated into a contiguous mass or region, such as a single piece of ice floating in water, or it may be distributed as small particles throughout the system, such as oil droplets in an emulsion or solid particles in a pharmaceutical suspension.

These latter two are examples of coarse dispersions, the diameter of the particles in emulsions and suspensions for the most part being larger than 0.1 µm (100 Å or 10^{-5} cm). A colloidal dispersion represents a system having a particle size intermediate between that of a true solution and a coarse dispersion, roughly 10 to 5000 Å. A colloidal dispersion may be considered as a two-phase system (heterogeneous) under certain circumstances and as a one-phase system (homogeneous) under others. A colloidal dispersion of silver proteinate in water is heterogeneous since it consists of distinct particles constituting a separate phase. A colloidal dispersion of acacia or sodium carboxymethylcellulose in water, on the other hand, is homogeneous. It does not differ significantly from a solution of sucrose and

may be considered as a single-phase system or true solution 1

A solution composed of only two substances is known as a binary solution, and the components or constituents are referred to as the solvent and the solute. We use the terms component and constituent interchangeably here, as do other authors, to represent the pure chemical substances that make up a solution. The number of components has a definite significance in the phase rule, as explained on p. 37. The constituent present in the greater amount in a binary solution is arbitrarily designated as the solvent and the constituent in the lesser amount as the solute. When a solid is dissolved in a liquid, however, the liquid is usually taken as the solvent and the solid as the solute, irrespective of the relative amounts of the constituents.

When water is one of the constituents of a liquid mixture, it is usually considered the solvent. When dealing with mixtures of liquids that are miscible in all proportions, such as alcohol and water, it is less meaningful to classify the constituents as solute and solvent.

Properties of Solutions. The physical properties of substances may be classified as colligative, additive, and constitutive. Some of the constitutive and additive properties of molecules were considered in Chapter 4. In the field of thermodynamics, physical properties of systems are classified as extensive properties, depending on the quantity of the matter in the system (e.g., mass and volume) and intensive properties, which are independent of the amount of the substances in the system (e.g., temperature, pressure, density, surface tension, and viscosity of a pure liquid).

Colligative properties depend mainly on the number of particles in a solution. The colligative properties of solutions are osmotic pressure, vapor pressure lowering, freezing point depression, and boiling point eleva-



DOCKET

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

