

**ADVANCES IN  
EXPERIMENTAL MEDICINE AND BIOLOGY**

---

**VOLUME 206**

Teva – Fresenius

## ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY

Editorial Board:

NATHAN BACK, *State University of New York at Buffalo*

NICHOLAS R. DI LUZIO, *Tulane University School of Medicine*

EPHRAIM KATCHALSKI-KATZIR, *The Weizmann Institute of Science*

DAVID KRITCHEVSKY, *Wistar Institute*

ABEL LAJTHA, *Rockland Research Institute*

RODOLFO PAOLETTI, *University of Milan*

---

### Recent Volumes in this Series

Volume 201  
LIPOPROTEIN DEFICIENCY SYNDROMES  
Edited by Aubie Angel and Jiri Frohlich

Volume 202  
INFECTIONS IN THE IMMUNOCOMPROMISED HOST  
Laboratory Diagnosis and Treatment  
Edited by Paul Actor, Alan Evangelista, James Poupard, and Eileen Hinks

Volume 203  
EXCITATORY AMINO ACIDS AND EPILEPSY  
Edited by Robert Schwarcz and Yehezkel Ben-Ari

Volume 204  
NEUROBIOLOGY OF CENTRAL D<sub>1</sub>-DOPAMINE RECEPTORS  
Edited by George R. Breese and Ian Creese

Volume 205  
MOLECULAR AND CELLULAR ASPECTS OF REPRODUCTION  
Edited by Dharam S. Dhindsa and Om P. Bahl

Volume 206  
ESSENTIAL NUTRIENTS IN CARCINOGENESIS  
Edited by Lionel A. Poirier, Paul M. Newberne, and  
Michael W. Pariza

Volume 207  
THE MOLECULAR AND CELLULAR BIOLOGY OF FERTILIZATION  
Edited by Jerry L. Hedrick

Volume 208  
PHOSPHATE AND MINERAL HOMEOSTASIS  
Edited by Shaul G. Massry, Michel Olmer, and  
Eberhard Ritz

---

A Continuation Order Plan is available for this series. A continuation order will bring delivery of each new volume immediately upon publication. Volumes are billed only upon actual shipment. For further information please contact the publisher.

Teva – Fresenius

THE ROLE OF VITAMIN B<sub>12</sub> AND FOLATE IN CARCINOGENESIS\*

Victor Herbert

Department of Medicine  
 Mount Sinai School of Medicine  
 New York, New York, and  
 Hematology and Nutrition Laboratory  
 Bronx Veterans Administration Medical Center  
 Bronx, New York 10468

## ABSTRACT

The roles of vitamin B<sub>12</sub> and folate in carcinogenesis are largely extensions of and linked to their roles in normal metabolism, particularly 1-carbon unit metabolism. A possible key area may be hypomethylation to "switch on" genes and methylation to "switch them off." Some vitamin analogues may act as antivitamin in these reactions, as may some vitamin-binding proteins. Others may act as specific delivery proteins. Using appropriate radioactive substrates and suspensions of vitamin-dependent normal and malignant cells, it may be possible to work out their positive and negative control of DNA synthesis.

## INTRODUCTION

The roles of vitamin B<sub>12</sub> and folate in carcinogenesis are largely extensions of and linked to their roles in normal metabolism. One key area is the conversion of homocysteine to methionine (methyl homocysteine). This process is dependent on folate delivering its 1-carbon unit to vitamin B<sub>12</sub>, which then becomes methyl-B<sub>12</sub> and transfers that methyl unit to homocysteine (Fig. 1). Newberne et al.<sup>1</sup> recently reviewed the role of the lipotropes choline and methionine and related factors in oncogenesis, including the impaired hormonal and cell-mediated immunity in folate-deficient humans and animals, and they pointed out the synergism between high fat diets and methyl deprivation.<sup>2</sup> Poirier<sup>3</sup> reviewed the protective effect of methionine against hepatocarcinogenesis, and Farber<sup>4</sup> discussed the carcinogenesis promotion effect of the ethyl analogue of methionine, ethionine.

ABBREVIATIONS: azaC = 5-azacytidine; TC II = transcobalamin II; PGA = pteroglutamic acid; SAM = S-adenosylmethionine; dU = deoxyuridine; AIDS = acquired immunodeficiency syndrome; dThd = thymidine; PHA = phytohemagglutinin A.

\*Supported in part by the Research Service of the U.S. Veterans Administration and U.S. Public Health Service Grant AM35709.

AMe = S-adenosylmethionine.  
 1 = serine hydroxymethyltransferase  
 2 = methylene THF reductase  
 3 = homocysteine transmethylase (methyltransferase)  
 4 = thymidylate synthetase  
 (The numbers represent enzymes)

5 = formiminotransferase  
 THF = tetrahydrofolate  
 DHF = dihydrofolate  
 B<sub>12</sub> = reduced Vitamin B<sub>12</sub>




Fig. 1. Biochemical interrelationships between vitamin B<sub>12</sub> and folate in human metabolism.

In a series of painstaking studies, Poirier's group<sup>5</sup> determined that, over a 76-week period, dietary methyl deficiency markedly promoted liver carcinogenesis and exhibited complete carcinogenic activity in this organ in the rat. They showed this in rats fed methyl-deficient, amino acid-deficient diets. When the diets were also devoid of folic acid and vitamin B<sub>12</sub>, the diethylnitrosamine-initiated rats died within 23 experimental weeks, before developing hepatocellular carcinoma, but all had livers containing hepatocytes of atypical appearance and, particularly at the 2 higher dosages of diethylnitrosamine, a cirrhotic pseudonodular architecture. They also found neoplastic conversion of rat liver epithelial cells in culture by ethionine and S-adenosylethionine.<sup>6</sup>

Krumdieck<sup>7</sup> reviewed the literature pertaining to the role of folate deficiency in facilitating carcinogenesis through 1982 and Eto<sup>8</sup> has carried the subject through the beginning of 1985. Folate is essential in the biosynthesis of both purines and pyrimidines<sup>9,10</sup> and therefore is required by all dividing cells. The conversion of deoxyuridylylate to thymidylate (methyldeoxyuridylylate) is folate- and B<sub>12</sub>-dependent, involving these 2 vitamins in a key step in DNA synthesis (Fig. 1).<sup>9-11</sup> These biochemical facts underlie the chromosomal abnormalities that characterize human clinical deficiency of vitamin B<sub>12</sub> and/or folate.<sup>12,13</sup> A wide range of chemical carcinogens inhibit DNA methylation in vitro.<sup>14</sup> It has been suggested that deficiency of folate or vitamin B<sub>12</sub> or any cause of failure to methylate DNA and/or RNA can activate malignancy by hypomethylating oncogenes, leading to such gene expression and/or gene amplifications, and that methylating oncogenes can inhibit malignancy by making them

dormant.<sup>3,15</sup> This is similar to the concept of "relaxed control" of RNA synthesis, discussed 3 decades ago by Borek and co-workers.<sup>16</sup> They noted that when an organism auxotrophic for methionine is deprived of methionine, it loses its ability to suppress synthesis of RNA, which is then synthesized more rapidly; they tied that observation to methylation of RNA. We speculated that vitamin B<sub>12</sub> or folate deficiency could produce such "relaxed control,"<sup>17</sup> and we noted more recently<sup>15</sup> that folate, vitamin B<sub>12</sub>, and their antagonists could be involved in the control of normal gene expression if in fact hypomethylation of DNA "switches on" normal genes and methylation "switches them off."<sup>18</sup> Although the evidence of this process is significant but inconclusive, one would expect that hypomethylation of the DNA or RNA of oncogenes would activate them and methylation would inactivate them. Perhaps some of the second cancers that develop after successful antimetabolic chemotherapy are due to the same chemotherapy that directly destroys an active cancer, demethylating an oncogene of a dormant cancer.

Gene amplification is a mechanism for tumor resistance to anti-metabolites.<sup>19</sup> One can speculate that it may also be a mechanism to aid in tumor proliferation by, for example, producing gene amplification of the hepatic Phase I enzymes that activate carcinogens.<sup>20</sup>

Gautsch and Wilson<sup>21</sup> found that de novo methylation of the input provirus occurs in embryonal carcinoma cells but not in permissive, differentiated teratocarcinoma. Harrison et al.<sup>22</sup> demonstrated a 3-way correlation between tumorigenicity, trisomy for 3q, and specific demethylation, suggesting that decreased DNA methylation may be involved both in differentiation and in tumorigenicity and that the antileukemia drug azaC may induce chromosomal aberrations as well as altering DNA methylation. Altering DNA methylation is just one of the varied effects of azaC on cellular metabolism.<sup>23-25</sup> The drug reduces DNA methylation and induces theoretically therapeutically valuable differentiation of human promyelocytic leukemia cells (HL-60) in culture, although this induction is less effective than that brought about in these cells by dimethylsulfoxide and L-methionine.<sup>26</sup> Anderson and colleagues<sup>25</sup> found that azaC selectively hypomethylates fetal globin genes, supporting work by Ley et al.<sup>23</sup> and Charache and associates.<sup>27</sup>

Patients with neoplasms excrete elevated levels of certain methylated bases in their urine, and Borek's group<sup>28-30</sup> has been attempting to correlate the quantity of such excretion with the degree of tumor activity. Gross's group, in collaboration with our group,<sup>15,31,32</sup> were unable to show any reproducible inhibitory effect of 5-methylcytidine on the development of presumably RNA virus-induced transplanted L2C leukemia-lymphoma in guinea pigs; this appears to be an animal analogue to human leukemia-lymphoma of RNA virus etiology.<sup>33,34</sup> After we switched to 5-iodocytidine, which seemed more promising, both groups<sup>35</sup> unsuccessfully sought funding targeted to continue this work. Gross's<sup>35</sup> recent dramatic report of reduction in the incidence (i.e., the initial development) of radiation-induced tumors in rats after restriction of caloric food intake has been associated with renewed funding. His group previously noted that restriction of food intake will not significantly influence the growth or progress of established tumors in mice.<sup>32</sup> American Cancer Society statistics suggest an increased frequency of malignancy in obese persons (L. Gross, personal communication).

The roles of vitamin B<sub>12</sub> and folate in carcinogenesis are not at the simple level at which serum vitamin levels correlate with extent of disease. No correlation has been found between serum folate and vitamin B<sub>12</sub> levels and the extent of small cell lung cancer.<sup>36</sup> However, a correlation may exist between levels of certain naturally occurring folate and

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.