

---

# The prevalence of previously diagnosed and undiagnosed psoriasis in US adults: Results from NHANES 2003-2004

Shanu Kohli Kurd, MHS,<sup>a,b</sup> and Joel M. Gelfand, MD, MSCE<sup>a,b</sup>  
*Philadelphia, Pennsylvania*

**Background:** Psoriasis is a predictor of morbidity. It is important to determine the extent to which psoriasis remains undiagnosed.

**Objective:** To determine the prevalence of psoriasis.

**Methods:** We conducted a cross-sectional study using the National Health and Nutrition Examination Survey 2003-2004.

**Results:** The prevalence of diagnosed psoriasis was 3.15% (95% confidence interval [CI], 2.18-4.53), corresponding to 5 million adults. Approximately 17% of these patients have moderate to severe psoriasis based on body surface area report and 25% rate psoriasis a large problem in everyday life. The prevalence of undiagnosed active psoriasis by conservative estimate was 0.4% (95% CI, 0.19-0.82), corresponding to approximately 600,000 US adults, and 2.28% (95% CI, 1.47-3.50) by a broader definition, corresponding to 3.6 million US adults. Undiagnosed patients had a trend toward being more likely to be male, nonwhite, less educated, and unmarried compared with patients who had received a diagnosis.

**Limitations:** The method for determining the presence of psoriasis had limited ability to detect mild disease and only fair interrater agreement.

**Conclusion:** More than 5 million adults have been diagnosed with psoriasis. A large number have undiagnosed psoriasis and there are important disparities which may be associated with not receiving medical attention. (J Am Acad Dermatol 2008;60:218-24.)

## INTRODUCTION

Psoriasis is a chronic, inflammatory disease of the skin and joints that negatively impacts health-related quality of life. More recent data have also demonstrated that psoriasis, particularly when severe, is

### Abbreviations used:

BSA:	body surface area
CI:	confidence interval
MEC:	mobile examination center
NHANES:	National Health and Nutrition Examination Survey
OR:	odds ratio

---

From the Department of Dermatology<sup>a</sup> and Center for Clinical Epidemiology and Biostatistics,<sup>b</sup> University of Pennsylvania School of Medicine.

Supported in part by a National Research Service Award from the National Institute of Health (to S.K.K.) and a grant K23AR051125 from the National Institutes of Health/National Institute of Arthritis, Musculoskeletal, and Skin Diseases (to J.M.G.).

Conflicts of interest: None declared.

Accepted for publication September 17, 2008.

Reprints not available from the authors.

Correspondence to: Joel M. Gelfand, MD, MSCE, University of Pennsylvania, Department of Dermatology, 3600 Spruce St, 2 Maloney Building Suite 2M47, Philadelphia, PA 19104. E-mail: Joel.Gelfand@uphs.upenn.edu.

Published online November 21, 2008.

0190-9622/\$36.00

© 2008 by the American Academy of Dermatology, Inc.

doi:10.1016/j.jaad.2008.09.022

associated with metabolic disorders, obesity, excess mortality and may be an independent risk factor for developing atherosclerosis, myocardial infarction, and stroke.<sup>1-6</sup> The treatment paradigm of psoriasis is undergoing a revolution with the recent approval of multiple systemic psoriasis treatments and the development of consensus statements which have broadened recommendations for which patients may qualify for systemic therapy.<sup>7,8</sup> Given these recent advances, it is important to understand how many patients suffer from psoriasis and which patients have disease that has substantial severity and/or impact on quality of life to warrant systemic

therapy. Furthermore, since psoriasis is increasingly being recognized to be a predictor of current and future morbidities, it is important to determine the extent to which psoriasis remains undiagnosed in the general population.

Previous estimates of the prevalence of psoriasis in various locations throughout the world have ranged from 0.6% to 4.8%.<sup>9-26</sup> These studies have varied in source population studied (eg, various ages, general population-based vs clinic-based), definition of prevalence (point vs period vs lifetime), and definition of psoriasis (eg, self report vs physician diagnosed). Two population-based studies in the continental United States of adults have found a prevalence of psoriasis of 2.2%<sup>24,27</sup> and 2.6%<sup>9</sup> based on patient report of a physician diagnosis attained by telephone and mail questionnaire, respectively. This approach may underestimate the true prevalence of disease because a significant portion of patients with psoriasis may not seek medical care and therefore be unaware of their diagnosis. The 1971-1974 Health and Nutrition Examination Survey of persons 1-74 years of age found a point prevalence of psoriasis of 1.4% based on physician examination.<sup>10</sup> This study did not evaluate whether a patient had ever had psoriasis in the past and therefore may underestimate the prevalence of psoriasis as the disease may be in remission because of treatment, seasonal changes, or natural history.<sup>9</sup>

A more comprehensive method of assessing the prevalence of psoriasis is needed in order to capture psoriasis patients who are aware of their diagnosis as well as those who may remain undiagnosed. To further investigate the prevalence of psoriasis in the general US population, we examined data from the National Health and Nutrition Examination Survey (NHANES) (2003-2004), which is unique in that it contains both information ascertained by patient report and physician examination and therefore can be used to determine the prevalence of psoriasis in patients who are aware of their diagnosis as well as the prevalence of psoriasis in patients with active yet previously undiagnosed psoriasis.

## METHODS

### Study design

We investigated the prevalence of psoriasis by analyzing data from the NHANES in the United States from 2003 through 2004. The study was approved by the National Center for Health Statistics institutional review board and all subjects gave informed consent. NHANES is an ongoing population-based, cross-sectional study which is designed to assess the health and nutritional status of people living in the United States. It is unique in that it combines interviews and

physical examinations, and the purpose of the NHANES initiative is to determine the prevalence of major diseases and risk factors for diseases in the United States. NHANES data are collected in 2-year cycles using a continuous stratified sampling technique. The first step is for participants to complete an interviewer-administered questionnaire. Physical examinations are conducted within 1 to 2 weeks of the in-home interview in specially designed and equipped mobile examination centers (MECs). The survey team consists of a physician, medical and health technicians, and dietary and health interviewers who are extensively trained. The prevalence of psoriasis was determined by patient interview as well as examination of standardized clinical photographs by two dermatologists.

### Study population

This study uses a subset of the NHANES survey from 2003-2004 which is representative of the non-institutionalized US civilian population aged 20-59 years, which was selected using a complex, multi-stage, stratified sampling design. African Americans, Mexican Americans, and low-income white Americans have been oversampled to increase the accuracy and precision of estimates of health status indicators for these population subgroups. The sample weights for NHANES 2003-2004 reflect the unequal probabilities of selection, nonresponse adjustments and adjustments to independent population controls, and sampling adjustments are made during statistical analysis.

### Definition of psoriasis

The prevalence of psoriasis was defined by two methods. The prevalence of previously diagnosed psoriasis was determined by interviewer-assisted questionnaire. A response of "yes" when asked "Have you ever been told by a health care provider that you had psoriasis?" was classified in our study as having psoriasis. This self-report method of ascertaining the prevalence of psoriasis has been well accepted in numerous epidemiological studies of psoriasis.<sup>9,15,19,24,27</sup>

The presence of psoriasis was also determined by review of standardized clinical photographs by two dermatologists. The data collection method consisted of taking standardized images of potentially affected areas of the skin during the MEC examination using a digital camera. Sites included the back, lower extremities, hands, and arms as follows: one image of the back that includes the elbows; one image of the left upper inner arm; one image of the dorsal surface of both hands combined with an image of the front lower legs; and one image of the

palm of both hands combined with an image of the back and lower legs. Prior to concluding the examination, images were checked for quality. The images were then transferred to a digital video disc and given to two independent dermatologists for analysis.

A diagnosis of psoriasis by both dermatologists in a patient who reported no history of diagnosis of psoriasis by questionnaire was used as our conservative definition of "undiagnosed active" psoriasis. A diagnosis by at least one dermatologist in a patient who reported no history of diagnosis of psoriasis by questionnaire was used as our less conservative definition of "undiagnosed active" psoriasis.

### Definition of covariates

Those patients who reported a previous diagnosis of psoriasis by questionnaire were asked additional questions regarding the extent of their psoriasis as well as the impact that psoriasis has on their daily lives. Patients were asked, "Do you currently have little or no psoriasis, only a few patches that could be covered by one or two palms of your hand, scattered patches that could be covered between 3 and 10 palms of your hand, or extensive psoriasis covering large areas of the body that would require more than 10 palms of your hand?" By convention, the palm of the hand is estimated to be approximately 1% body surface area (BSA) and therefore we present these categories as little/no psoriasis, 1%-2% BSA, 3%-10% BSA, and >10% BSA, respectively.<sup>28</sup> For additional and logistic regression analyses, psoriasis severity categories were dichotomized into mild (<3%) and moderate to severe ( $\geq 3\%$ ). Patients were asked to answer the question "On a scale of 1 to 10 how much of a problem has your psoriasis been in your everyday life where 1 means no problem at all and 10 means a very large problem." Ordinal categories were created for additional analyses in a manner identical to previously published work (1-3: no or little impact, 4-7: a problem, 8-10: a large problem).<sup>24</sup>

Additional covariate information for all patients such as age, gender, race, education, income, health insurance and marital status were determined by questionnaire. Education was dichotomized into greater than high school education or not. Income was dichotomized into less than or equal to a median income level of \$35,000-\$44,999 or greater and marital status was dichotomized into married or not married. In descriptive statistics, the prevalence of race was reported based on questionnaire categories (Caucasian, black, Hispanic, other) but for logistic regression was dichotomized into white or non-white.

Other variables of interest such as frequency of physician visits, current smoking, alcohol use, depression and anxiety screening were not used because of a high proportion of missing data. To maintain quality analyses, only covariates for which data were recorded in at least 90% of patients were included.

### Statistical methods

All statistical analyses were performed using survey commands of STATA to incorporate sample weights and adjust for clusters and strata of the complex sample design (Version 10, STATA Corp, College Station, Tex). The prevalence of psoriasis was determined by responses to the dermatological questionnaire as well as dermatological clinical evaluation. Prevalence and covariate data were reported as percentages with 95% confidence intervals using the survey tabulate command in STATA. Odds ratios (ORs) with 95% confidence intervals were calculated using survey logistic regression for both unadjusted and adjusted (for age and gender) analyses. Quality of life data were analyzed using logistic regression and linear regression. Agreement about the presence of psoriasis on the clinical photographs was assessed via the kappa statistic. All *P* values reported are two sided.

### RESULTS

The dermatologic interview and examination were only administered to subjects aged 20-59 years and represent a subset of the total NHANES data which includes all ages. An unweighted sample size of 4,163 people aged 20-59 years were screened, 3140 of whom participated in questionnaire-based interviews. A total of 2984 of the interviewed subjects had nonmissing psoriasis questionnaire data and this was the cohort used to calculate prevalence estimates of psoriasis. Data from dermatologist review of clinical photographs were available for 90% of patients in this cohort. The raw data include 73 patients with psoriasis as determined by self-report of a healthcare provider diagnosis on questionnaire. Upon examination of standardized photographs, 84 patients were determined to have active psoriasis by at least one dermatologist, 66 of whom by self-report had never received a prior diagnosis of psoriasis by a healthcare provider. Among the 73 patients who reported a prior healthcare provider diagnosis of psoriasis, the presence of psoriasis, based on review of the standardized clinical photographs, could only be confirmed by at least one of two dermatologists in 18 patients. Confirmation of active psoriasis was strongly dependent upon patient self report of skin severity. For example, the odds of one or both

**Table I.** Summary statistics of psoriasis patients, 20-59 years of age, in the US population\*

	No psoriasis	Psoriasis		
		History of healthcare provider diagnosis	Undiagnosed active psoriasis: Conservative estimate <sup>†</sup>	Undiagnosed active psoriasis: Less conservative estimate <sup>‡</sup>
Psoriasis prevalence, % (95% CI)	NA			
Overall		3.15 (2.18-4.53)	0.40% (0.19-0.82)	2.28% (1.47-3.50)
Women		1.62% (1.02-2.55)		0.48% (0.22-1.03)
Men		1.53% (0.97-2.40)		1.8% (1.15-2.8)
Gender, % male (95% CI)	49.03% (47.69-50.36)	48.61% (35.54-61.86)	82.31% (25.62-98.43)	79.1% (63.99-88.96)
Mean age (95% CI)	38.91 (38.10-39.72)	41.11 (38.80-43.42)	44.06 (34.58-53.54)	41.69 (38.74-44.64)
Race, %				
White	68.79%	81.61%	79.94	70.01
Black	12.16	7.96	12.19	16.76
Hispanic	13.29	6.53	7.87	4.44
Other	5.76	3.9	0	8.79
Income ≤ median, <sup>§</sup> % (95% CI)	43.91 (39.49-48.43)	37.37 (28.24-47.49)	26.89 (4.78-72.94)	51.63 (36.83-66.16)
No advanced (post-high school) education, % (95% CI)	42.59 (40.38-44.83)	29.51 (19.58-41.86)	57.17 (25.50-83.89)	56.85 (45.58-67.45)
Not married, % (95% CI)	43.86 (40.45-47.32)	31.79 (23.01-42.09)	37.97 (11.62-74.03)	47.10 (29.19-65.79)
No health insurance, % (95% CI)	22.59 (20.44-24.89)	16.5 (9.19-26.83)	0	27.16 (14.63-44.79)

CI, Confidence interval.

\*Psoriasis patients with a history of healthcare provider diagnosis and patients with undiagnosed active psoriasis are compared with patients without psoriasis.

<sup>†</sup>The conservative estimate of undiagnosed active psoriasis is determined by agreement of psoriasis by both dermatologists upon review of standardized photographs in a patient who reported no previous diagnosis of psoriasis by a healthcare provider.

<sup>‡</sup>Undiagnosed active psoriasis by less conservative estimate is determined by at least one dermatologist upon review of standardized photographs in a patient who reported no previous diagnosis of psoriasis by a healthcare provider.

<sup>§</sup>Median income is \$35,000 to \$44,999.

evaluators confirming psoriasis was 11.71 (95% CI, 4.03-33.98;  $P < .001$ ) and 9.71 (95% CI, 1.77-53.34;  $P = .01$ ), respectively, for patients with self reported BSA  $\geq 3\%$  compared with patients with no to mild psoriasis by self-report (eg, BSA  $< 3\%$ ). Agreement between the two dermatologists on the presence of psoriasis for all subjects evaluated was only fair<sup>29</sup> ( $\kappa$  0.36,  $P < .001$ ) and both dermatologists agreed only 17% of the time in previously undiagnosed individuals. However, agreement between the two dermatologists among photographs of subjects who self-reported the presence of psoriasis via questionnaire was moderate<sup>29</sup> ( $\kappa$  0.54,  $P < .001$ ), and agreement among subjects with the most extensive disease ( $> 10\%$  BSA) was perfect ( $\kappa$  1.0,  $P = .02$ ).

To determine prevalence estimates and measures of association, the raw data were adjusted for the sampling technique (Table I). The prevalence of psoriasis based on patient report of a healthcare provider diagnosis of the disease was similar in men and women and is estimated to be 3.15% (95% CI, 2.18-4.53) of the US population aged 20-59 years.

The prevalence of undiagnosed active psoriasis is estimated to be 0.4% (95% CI, 0.19- 0.82) based on a conservative definition (eg, both evaluators needed to confirm the presence of psoriasis). The prevalence of undiagnosed active psoriasis based on a less conservative definition (eg, at least one of the two evaluators needed to confirm the presence of psoriasis) was 2.28% (95% CI, 1.47-3.50) of the US population aged 20-59 years.

Table I also summarizes characteristics of patients with previously diagnosed psoriasis compared with patients who have undiagnosed active psoriasis. Subsequent analyses were conducted to determine which factors were associated with not receiving a previous diagnosis of psoriasis. Although not statistically significant because of a small sample size, regression analysis adjusted for age and sex showed a trend toward a greater likelihood of being male (OR, 5.72; 95% CI, 0.49-67.35), non-white (OR, 1.34; 95% CI, 0.20-8.83), less educated (OR, 3.78; 95% CI, 0.64-22.37), and not married (OR, 1.70; 95% CI, 0.26-11.05) in patients with undiagnosed active psoriasis

**Table II.** Extent of psoriasis severity and impact on daily life in patients with history of a healthcare provider diagnosis of psoriasis\*

	Mild psoriasis		Moderate to severe psoriasis	
	Little/no psoriasis	1%-2% BSA psoriasis	3%-10% BSA psoriasis	>10% BSA psoriasis
Proportion of psoriasis patients	51.59%	31.68%	11.38%	5.25%
Mean psoriasis impact on daily life (95% CI)	3.22 (2.37-4.07)	4.11 (2.97-5.25)	7.32 (5.31-9.32)	10 (10.0-10.0)

BSA, Body surface area.

\*Of psoriasis patients with a history of physician diagnosis, 83.37% had mild psoriasis (little/no to 1%-2% BSA), whereas 16.63% had moderate to severe disease ( $\geq 3\%$  BSA). Mean psoriasis impact on daily life score increased with increasing psoriasis severity.

confirmed by both dermatologists compared to patients with a previous diagnosis of psoriasis. Patients with undiagnosed psoriasis based on the less conservative definition were more likely to be male (OR, 4.29; 95% CI, 2.34-7.87), non-white (OR, 2.12; 95% CI, 1.00-4.48), have a lower household income (OR, 2.46; 95% CI, 1.02-5.95), be less educated (OR, 3.51; 95% CI, 1.39-8.88), and not married (OR, 2.65; 95% CI, 1.39-5.07) compared to patients who had received a diagnosis. Health insurance status was not associated with having newly diagnosed psoriasis (OR, 1.83; 95% CI, 0.73-4.61).

Patients with a history of healthcare provider-diagnosed psoriasis were asked to provide information on the extent of their psoriasis as well as how psoriasis impacts their daily life. Psoriasis impact on daily life was scored from 1-10 where 1 indicates no impact and 10 indicates severe impact (Table II). In analysis of the extent of psoriasis on daily life, previously established categories were applied (1-3, little or no problem; 4-7, a problem; 8-10, a large problem).<sup>24</sup> More than half (56.66%) of all psoriasis patients with a previous diagnosis reported that psoriasis was a little or no problem in their daily lives, whereas 18.44% reported that it was a problem and 24.91% reported that it was a large problem. It is estimated that 83.37% of the previously diagnosed psoriasis population has limited skin disease ( $<3\%$  BSA) while 16.63% are estimated to suffer from more moderate to severe disease ( $\geq 3\%$  BSA). After adjustment for age and gender, the extent to which psoriasis impacts daily life was associated with psoriasis severity ( $P < .001$  for trend). For example, 16% of patients with mild psoriasis rated it as a large problem, whereas 72% of moderate to severe psoriasis patients rated it as a large problem.

## DISCUSSION

The unique design of the NHANES 2003-2004 in which the prevalence of psoriasis was measured by both patient self-report of a prior healthcare provider diagnosis as well as by review of clinical photographs provides important confirmation of previous epidemiological studies as well as novel findings.

First, the prevalence of self report of a healthcare provider diagnosis of psoriasis is statistically similar to previous estimates in the US adult population using comparable methods.<sup>9,24</sup> Second, this study confirms that a substantial number of psoriasis patients suffer from disease which is severe enough to warrant systemic therapy based on current consensus statements.<sup>7,8</sup> For example, about 17% have moderate to severe psoriasis based on BSA estimates, and 25% of patients rate psoriasis as a large problem in everyday life. Furthermore, we determined that by conservative estimates approximately 0.4% (2.28% based on a less strict definition) of the general population aged 20-59 years have undiagnosed, clinically active psoriasis. Importantly, our analyses demonstrate that significant disparities may exist among psoriasis patients in terms of the likelihood that they have been previously diagnosed by a healthcare provider. In particular, people with undiagnosed active psoriasis may be more likely to be male, unmarried, non-white, and have less education.

With application of these prevalence estimates to the population at the time these data were collected, it is estimated that approximately 5 million people 20-59 years of age have been previously diagnosed with psoriasis and that, despite having active disease, an additional 600,000 (conservative estimate) to 3.6 million people have psoriasis but remain undiagnosed. Furthermore, we estimate that 1.4 million patients aged 20-59 years have moderate to severe disease and that 2.1 million consider psoriasis to be a large problem in daily life.

A particular strength of this study is that NHANES participants are selected in a manner that is representative of the general US population. In addition, interviews and examinations are conducted on site at people's homes and participation rates are favorable to other published reports of the prevalence of psoriasis.<sup>9,30</sup> As with any study, there are important limitations to consider. As NHANES interviewers and examiners were unaware of our hypothesis to be tested, it is unlikely that any misclassification of covariates measured was directional (ie that

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.