



US005665772A

United States Patent [19]

[11] Patent Number: **5,665,772**

Cottens et al.

[45] Date of Patent: **Sep. 9, 1997**

[54] **O-ALKYLATED RAPAMYCIN DERIVATIVES AND THEIR USE, PARTICULARLY AS IMMUNOSUPPRESSANTS**

[51] Int. Cl.⁶ **A61K 31/395; C07D 498/16**

[52] U.S. Cl. **514/514; 540/456**

[58] Field of Search **540/456; 514/514**

[75] Inventors: **Sylvain Cottens, Witterswil; Richard Sedrani, Basel, both of Switzerland**

[56] **References Cited**

U.S. PATENT DOCUMENTS

5,120,842 6/1992 Failli et al. 540/542

5,151,413 9/1992 Caufield et al. 514/63

5,258,389 11/1993 Goulet et al. 514/291

Primary Examiner—Robert T. Bond

Attorney, Agent, or Firm—Robert S. Honor; Melvyn M. Kassenoff; Thomas O. McGovern

[57] **ABSTRACT**

Novel derivatives of rapamycin, particularly 9-deoxorapamycins, 26-dihydro-rapamycins, and 40-0-substituted and 28,40-0,0-di-substituted rapamycins, are found to have pharmaceutical utility, particularly as an immunosuppressants.

[73] Assignee: **Sandoz Ltd., Basel, Switzerland**

[21] Appl. No.: **416,673**

[22] PCT Filed: **Sep. 24, 1993**

[86] PCT No.: **PCT/EP93/02604**

§ 371 Date: **Apr. 7, 1995**

§ 102(e) Date: **Apr. 7, 1995**

[87] PCT Pub. No.: **WO94/09010**

PCT Pub. Date: **Apr. 28, 1994**

[30] **Foreign Application Priority Data**

Oct. 9, 1992 [GB] United Kingdom 9221220


10 Claims, No Drawings

**O-ALKYLATED RAPAMYCIN DERIVATIVES
AND THEIR USE, PARTICULARLY AS
IMMUNOSUPPRESSANTS**

This application is a 371 of PCT/EP93/02604, filed Sep. 24, 1993.


This invention comprises novel alkylated derivatives of rapamycin having pharmaceutical utility, especially as immunosuppressants.

Rapamycin is a known macrolide antibiotic produced by *Streptomyces hygroscopicus*, having the structure depicted in Formula A:



See, e.g., McAlpine, J. B., et al., *J. Antibiotics* (1991) 44: 688; Schreiber, S. L., et al., *J. Am. Chem. Soc.* (1991) 113: 7433; U.S. Pat. No. 3,929,992. Rapamycin is an extremely potent immunosuppressant and has also been shown to have antitumor and antifungal activity. Its utility as a pharmaceutical, however, is restricted by its very low and variable bioavailability as well as its high toxicity. Moreover, rapamycin is highly insoluble, making it difficult to formulate stable galenic compositions.

It has now surprisingly been discovered that certain novel derivatives of rapamycin (the Novel Compounds) have an improved pharmacologic profile over rapamycin, exhibit greater stability and bioavailability, and allow for greater ease in producing galenic formulations. The Novel Compounds are alkylated derivatives of rapamycin having the structure of Formula I:



X is (H,H) or O;

Y is (H,OH) or O;

R¹ and R² are independently selected from H, alkyl, thioalkyl, arylalkyl, hydroxyalkyl, dihydroxyalkyl, hydroxyalkylarylalkyl, dihydroxyalkylarylalkyl, alkoxyalkyl, acyloxyalkyl, aminoalkyl, alkylaminoalkyl, alkoxy-carbonylaminoalkyl, acylaminoalkyl, arylsulfonamidoalkyl, allyl, dihydroxyalkylallyl, dioxolanylallyl, carbalkoxyalkyl, and (R³)₃Si where each R³ is independently selected from H, methyl, ethyl, isopropyl, t-butyl, and phenyl; wherein "alk-" or "alkyl" refers to C₁₋₆ alkyl, branched or linear preferably C₁₋₃ alkyl, in which the carbon chain may be optionally interrupted by an ether (—O—) linkage; and

R⁴ is methyl, or R⁴ and R¹ together form C₂₋₅ alkylene; provided that R¹ and R² are not both H; and provided that where R¹ is (R³)₃Si or carbalkoxyalkyl, X and Y are not both O.

Preferred Novel Compounds include the following:

1. 40-O-Benzyl-rapamycin
2. 40-O-(4'-Hydroxymethyl)benzyl-rapamycin
3. 40-O-[4'-(1,2-Dihydroxyethyl)]benzyl-rapamycin
4. 40-O-Allyl-rapamycin
5. 40-O-[3'-(2,2-Dimethyl-1,3-dioxolan-4(S)-yl)-prop-2'-en-1'-yl]-rapamycin
6. (2'E,4'S)-40-O-(4',5'-Dihydroxypent-2'-en-1'-yl)-rapamycin
7. 40-O-(2-Hydroxy)ethoxycarbonylmethyl-rapamycin
8. 40-O-(2-Hydroxy)ethyl-rapamycin
9. 40-O-(3-Hydroxy)propyl-rapamycin
10. 40-O-(6-Hydroxy)hexyl-rapamycin
11. 40-O-[2-(2-Hydroxy)ethoxy]ethyl-rapamycin
12. 40-O-[(3S)-2,2-Dimethyldioxolan-3-yl]methyl-rapamycin
13. 40-O-[(2S)-2,3-Dihydroxyprop-1-yl]-rapamycin
14. 40-O-(2-Acetoxy)ethyl-rapamycin
15. 40-O-(2-Nicotinoyloxy)ethyl-rapamycin
16. 40-O-[2-(N-Morpholino)acetoxy]ethyl-rapamycin
17. 40-O-(2-N-Imidazolylacetoxy)ethyl-rapamycin
18. 40-O-[2-(N-Methyl-N'-piperazinyl)acetoxy]ethyl-rapamycin
19. 39-O-Desmethyl-39,40-O,O-ethylene-rapamycin
20. (26R)-26-Dihydro-40-O-(2-hydroxy)ethyl-rapamycin
21. 28-O-Methyl-rapamycin
22. 40-O-(2-Aminoethyl)-rapamycin
23. 40-O-(2-Acetaminoethyl)-rapamycin
24. 40-O-(2-Nicotinamidoethyl)-rapamycin
25. 40-O-(2-(N-Methyl-imidazo-2'-yl)carbethoxamido)ethyl)-rapamycin
26. 40-O-(2-Ethoxycarbonylaminoethyl)-rapamycin
27. 40-O-(2-Tolylsulfonamidoethyl)-rapamycin
28. 40-O-[2-(4',5'-Dicarboethoxy-1',2',3'-triazol-1'-yl)-ethyl]-rapamycin

The Novel Compounds for immunosuppressive use are preferably the 40-O-substituted rapamycins where X and Y are both O, R² is H, R⁴ is methyl, and R¹ is other than H; most preferably where R¹ is selected from hydroxyalkyl, hydroxyalkoxyalkyl, acylaminoalkyl, and aminoalkyl; especially 40-O-(2-hydroxy)ethyl-rapamycin, 40-O-(3-hydroxy)propyl-rapamycin, 40-O-[2-(2-hydroxy)ethoxy]ethyl-rapamycin, and 40-O-(2-acetaminoethyl)-rapamycin.

Preferably O-substitution at C40 or O,O-disubstitution at C28 and C40 is performed according to the following general process: Rapamycin (or dihydro or

e.g., an alkyl, allyl, or benzyl moiety, which is desired as the O-substituent, and X is the leaving group, e.g., $\text{CCl}_3\text{C}(\text{NH})\text{O}$ or CF_3SO_3 under suitable reaction conditions, preferably acidic or neutral conditions, e.g., in the presence of an acid like trifluoromethanesulfonic acid, camphorsulfonic acid, p-toluenesulfonic acid or their respective pyridinium or substituted pyridinium salts when X is $\text{CCl}_3(\text{NH})\text{O}$ or in the presence of a base like pyridine, a substituted pyridine, diisopropylethylamine or pentamethylpiperidine when X is CF_3SO_3 . O-substituents at C28 only are accomplished in the same manner, but with prior protection at C40. Further modifications are possible. For example, where the substituent is allyl, the isolated, monosubstituted double bond of the allyl moiety is highly amenable to further modification.

The 9-deoxorapamycin compounds are preferably produced by reducing a rapamycin using hydrogen sulfide, by reacting rapamycin with diphenyldiselenide and tributylphosphine or by other suitable reduction reaction.

The 26-dihydro-rapamycins are preferably produced by reducing rapamycins or 9-deoxorapamycins from keto to hydroxy at C26 by a mild reduction reaction, such as a borohydride reduction reaction.

The Novel Compounds are particularly useful for the following conditions:

- a) Treatment and prevention of organ or tissue transplant rejection, e.g. for the treatment of recipients of e.g. heart, lung, combined heart-lung, liver, kidney, pancreatic, skin or corneal transplants. They are also indicated for the prevention of graft-versus-host disease, such as following bone marrow transplantation.
- b) Treatment and prevention of autoimmune disease and of inflammatory conditions, in particular inflammatory conditions with an etiology including an autoimmune component such as arthritis (for example rheumatoid arthritis, arthritis chronica progrediente and arthritis deformans) and rheumatic diseases. Specific autoimmune diseases for which the compounds of the invention may be employed include, autoimmune hematological disorders (including e.g. hemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polycondritis, sclerodoma, Wegener granulomatosis, dermatomyositis, chronic active hepatitis, myasthenia gravis, psoriasis, Steven-Johnson syndrome, idiopathic sprue, autoimmune inflammatory bowel disease (including e.g. ulcerative colitis and Crohn's disease) endocrine ophthalmopathy, Graves disease, sarcoidosis, multiple sclerosis, primary billiary cirrhosis, juvenile diabetes (diabetes mellitus type I), uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis, glomerulonephritis (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy) and juvenile dermatomyositis.
- c) Treatment and prevention of asthma.
- d) Treatment of multi-drug resistance (MDR). The Novel Compounds suppress P-glycoproteins (Pgp), which are the membrane transport molecules associated with MDR. MDR is particularly problematic in cancer patients and AIDS patients who will not respond to conventional chemotherapy because the medication is pumped out of the cells by Pgp. The Novel Compounds are therefore useful for enhancing the efficacy of other chemotherapeutic agents in the treatment and control of multidrug resistant conditions such as multidrug resis-

- e) Treatment of proliferative disorders, e.g. tumors, hyperproliferative skin disorder and the like.
- f) Treatment of fungal infections.
- g) Treatment and prevention of inflammation, especially in potentiating the action of steroids.
- h) Treatment and prevention of infection, especially infection by pathogens having Mip or Mip-like factors.
- i) Treatment of overdoses of FK-506, rapamycin, immunosuppressive Novel Compounds, and other macrophilin binding immunosuppressants.

The invention thus provides the Novel Compounds described herein, for use as novel intermediates or as pharmaceuticals, methods of treating or preventing the above-described disorders by administering an effective amount of Novel Compound to a patient in need thereof, use of a Novel Compound in the manufacture of a medicament for treatment or prevention of the above-described disorders, and pharmaceutical compositions comprising a Novel Compound in combination or association with a pharmaceutically acceptable diluent or carrier.

Most of the Novel Compounds described herein are highly immunosuppressive, especially those Novel Compounds which are O-substituted at C40, and these Novel Compounds are particularly useful in indications a and b, but not in indication i. Those of the Novel Compounds which are less immunosuppressive, especially those which are O-substituted at C28 only, are particularly useful in indications h and i, but are less preferred in indications a or b.

The Novel Compounds are utilized by administration of a pharmaceutically effective dose in pharmaceutically acceptable form to a subject in need of treatment. Appropriate dosages of the Novel Compounds will of course vary, e.g. depending on the condition to be treated (for example the disease type or the nature of resistance), the effect desired and the mode of administration.

In general however satisfactory results are obtained on administration orally at dosages on the order of from 0.05 to 5 or up to 10 mg/kg/day, e.g. on the order of from 0.1 to 2 or up to 7.5 mg/kg/day administered once or, in divided doses 2 to 4x per day, or on administration parenterally, e.g. intravenously, for example by i.v. drip or infusion, at dosages on the order of from 0.01 to 2.5 up to 5 mg/kg/day, e.g. on the order of from 0.05 or 0.1 up to 1.0 mg/kg/day. Suitable daily dosages for patients are thus on the order of 500 mg p.o., e.g. on the order of from 5 to 100 mg p.o., or on the order of from 0.5 to 125 up to 250 mg i.v., e.g. on the order of from 2.5 to 50 mg i.v.

Alternatively and even preferably, dosaging is arranged in patient specific manner to provide pre-determined trough blood levels, e.g. as determined by RIA technique. Thus patient dosaging may be adjusted so as to achieve regular on-going trough blood levels as measured by RIA on the order of from 50 or 150 up to 500 or 1000 ng/ml, i.e. analogously to methods of dosaging currently employed for Cyclosporin immunosuppressive therapy.

The Novel Compounds may be administered as the sole active ingredient or together with other drugs. For example, in immunosuppressive applications such as prevention and treatment of graft vs. host disease, transplant rejection, or autoimmune disease, the Novel Compounds may be used in combination with Cyclosporin, FK-506, or their immunosuppressive derivatives; corticosteroids; azathioprene; immunosuppressive monoclonal antibodies, e.g., monoclonal antibodies to CD3, CD4, CD25, CD28, or CD45; and 7 or other immunomodulatory compounds. For anti-inflammatory applications the Novel Compounds can be

teroids. For anti-infective applications, the Novel Compounds can be used in combination with other anti-infective agents, e.g., anti-viral drugs or antibiotics.

The Novel Compounds are administered by any conventional route, in particular enterally, e.g. orally, for example in the form of solutions for drinking, tablets or capsules or parenterally, for example in the form of injectable solutions or suspensions. Suitable unit dosage forms for oral administration comprise, e.g. from 1 to 50 mg of a compound of the invention, usually 1 to 10 mg. Pharmaceutical compositions comprising the novel compounds may be prepared analogously to pharmaceutical compositions comprising rapamycin, e.g., as described in EPA 0 041 795, which would be evident to one skilled in the art.

The pharmacological activity of the Novel Compounds are demonstrated in, e.g., the following tests:

1. Mixed lymphocyte reaction (MLR)

The Mixed Lymphocyte Reaction was originally developed in connection with allografts, to assess the tissue compatibility between potential organ donors and recipients, and is one of the best established models of immune reaction in vitro. A murine model MLR, e.g., as described by T. Meo in "Immunological Methods", L. Lefkovits and B. Peris, Eds., Academic Press, N.Y. pp. 227-239 (1979), is used to demonstrate the immunosuppressive effect of the Novel Compounds. Spleen cells (0.5×10^6) from Balb/c mice (female, 8-10 weeks) are co-incubated for 5 days with 0.5×10^6 irradiated (2000 rads) or mitomycin C treated spleen cells from CBA mice (female, 8-10 weeks). The irradiated allogeneic cells induce a proliferative response in the Balb/c spleen cells which can be measured by labeled precursor incorporation into the DNA. Since the stimulator cells are irradiated (or mitomycin C treated) they do not respond to the Balb/c cells with proliferation but do retain their antigenicity. The antiproliferative effect of the Novel Compounds on the Balb/c cells is measured at various dilutions and the concentration resulting in 50% inhibition of cell proliferation (IC_{50}) is calculated. The inhibitory capacity of the test sample may be compared to rapamycin and expressed as a relative IC_{50} (i.e. IC_{50} test sample/ IC_{50} rapamycin).

2. IL-6 mediated proliferation

The capacity of the Novel Compounds to interfere with growth factor associated signalling pathways is assessed using an interleukin-6 (IL-6)-dependent mouse hybridoma cell line. The assay is performed in 96-well microtiter plates. 5000 cells/well are cultivated in serum-free medium (as described by M. H. Schreier and R. Tees in Immunological Methods, I. Lefkovits and B. Pernis, eds., Academic Press 1981, Vol. II, pp. 263-275), supplemented with 1 ng recombinant IL-6/ml. Following a 66 hour incubation in the absence or presence of a test sample, cells are pulsed with 1 μ Ci (3-H)-thymidine/well for another 6 hours, harvested and counted by liquid scintillation. (3-H)-thymidine incorporation into DNA correlates with the increase in cell number and is thus a measure of cell proliferation. A dilution series of the test sample allows the calculation of the concentration resulting in 50% inhibition of cell proliferation (IC_{50}). The inhibitory capacity of the test sample may be compared to rapamycin and expressed as a relative IC_{50} (i.e. IC_{50} test sample/ IC_{50} rapamycin).

3. Macrophilin binding assay

Rapamycin and the structurally related immunosuppressant, FK-506, are both known to bind in vivo to macrophilin-12 (also known as FK-506 binding

pounds. The Novel Compounds also bind strongly to macrophilin-12, as is demonstrated in a competitive binding assay.

In this assay, FK-506 coupled to BSA is used to coat microtiter wells. Biotinylated recombinant human macrophilin-12 (biot-MAP) is allowed to bind in the presence or absence of a test sample to the immobilized FK-506. After washing (to remove non-specifically bound macrophilin), bound biot-MAP is assessed by incubation with a streptavidin-alkaline phosphatase conjugate, followed by washing and subsequent addition of p-nitrophenyl phosphate as a substrate. The read-out is the OD at 405 nm. Binding of a test sample to biot-MAP results in a decrease in the amount of biot-MAP bound to the FK-506 and thus in a decrease in the OD405. A dilution series of the test sample allows determination of the concentration resulting in 50% inhibition of the biot-MAP binding to the immobilized FK-506 (IC_{50}). The inhibitory capacity of a test sample is compared to the IC_{50} of free FK-506 as a standard and expressed as a relative IC_{50} (i.e., IC_{50} -test sample/ IC_{50} -free FK-506).

4. Localized Graft-Versus-Host (GvH) Reaction

In vivo efficacy of the Novel Compounds is proved in a suitable animal model, as described, e.g., in Ford et al, TRANSPLANTATION 10 (1970) 258. Spleen cells (1×10^7) from 6 week old female Wistar/Furth (WF) rats are injected subcutaneously on day 0 into the left hind-paw of female (F344 \times WF) F_1 rats weighing about 100 g. Animals are treated for 4 consecutive days and the popliteal lymph nodes are removed and weighed on day 7. The difference in weight between the two lymph nodes is taken as the parameter for evaluating the reaction.

5. Kidney Allograft Reaction in Rat

One kidney from a female fisher 344 rat is transplanted onto the renal vessel of a unilaterally (left side) nephrectomized WF recipient rat using an end-to-end anastomosis. Ureteric anastomosis is also end-to-end. Treatment commences on the day of transplantation and is continued for 14 days. A contralateral nephrectomy is done seven days after transplantation, leaving the recipient relying on the performance of the donor kidney. Survival of the graft recipient is taken as the parameter for a functional graft.

6. Experimentally Induced Allergic Encephalomyelitis (EAE) in Rats

Efficacy of the Novel Compounds in EAE is measured, e.g., by the procedure described in Levine & Wenk, AMER J PATH 47 (1965) 61; McFarlin et al, J IMMUNOL 113 (1974) 712; Borel, TRANSPLANT. & CLIN. IMMUNOL 13 (1981) 3. EAE is a widely accepted model for multiple sclerosis. Male Wistar rats are injected in the hind paws with a mixture of bovine spinal cord and complete Freund's adjuvant. Symptoms of the disease (paralysis of the tail and both hind legs) usually develop within 16 days. The number of diseased animals as well as the time of onset of the disease are recorded.

7. Freund's Adjuvant Arthritis

Efficacy against experimentally induced arthritis is shown using the procedure described, e.g., in Winter & Nuss, ARTHRITIS & RHEUMATISM 9 (1966) 394; Billingham & Davies, HANDBOOK OF EXPERIMENTAL PHARMACOLOGY (Vane & Ferreira Eds, Springer-Verlag, Berlin) 50/II (1979) 108-144. OFA and Wistar rats (male or female, 150 g body weight) are injected i.e. at the base of the tail or in the hind paw with 0.1 ml of mineral oil containing 0.6 mg of lyophilized heat-killed Mycobacterium smegmatis. In the

established arthritis model treatment is started on day 14, when the secondary inflammation is well developed (days 14–20). At the end of the experiment, the swelling of the joints is measured by means of a micro-caliper. ED₅₀ is the oral dose in mg/kg which reduces the swelling (primary or secondary) to half of that of the controls.

8. Antitumor and MDR activity

The antitumor activity of the Novel Compounds and their ability to enhance the performance of antitumor agents by alleviating multidrug resistance is demonstrated, e.g., by administration of an anticancer agent, e.g., colchicine or etoposide, to multidrug resistant cells and drug sensitive cells in vitro or to animals having multidrug resistant or drug sensitive tumors or infections, with and without co-administration of the Novel Compounds to be tested, and by administration of the Novel Compound alone.

Such in vitro testing is performed employing any appropriate drug resistant cell line and control (parental) cell line, generated, e.g. as described by Ling et al., *J. Cell. Physiol.* 83, 103–116 (1974) and Bech-Hansen et al. *J. Cell. Physiol.* 88, 23–32 (1976). Particular clones chosen are the multidrug resistant (e.g. colchicine resistant) line CHR (subclone C5S3.2) and the parental, sensitive line AUX B1 (subclone AB1 S11).

In vivo anti-tumor and anti-MDR activity is shown, e.g., in mice injected with multidrug resistant and drug sensitive cancer cells. Ehrlich ascites carcinoma (EA) sub-lines resistant to drug substance DR, VC, AM, ET, TE or CC are developed by sequential transfer of EA cells to subsequent generations of BALB/c host mice in accordance with the methods described by Slater et al., *J. Clin. Invest.*, 70, 1131 (1982).

Equivalent results may be obtained employing the Novel Compounds test models of comparable design, e.g. in vitro, or employing test animals infected with drug-resistant and drug sensitive viral strains, antibiotic (e.g. penicillin) resistant and sensitive bacterial strains, anti-mycotic resistant and sensitive fungal strains as well as drug resistant protozoal strains, e.g. Plasmodial strains, for example naturally occurring sub-strains of *Plasmodium falciparum* exhibiting acquired chemotherapeutic, anti-malarial drug resistance.

9. FKBP binding

Certain of the Novel Compounds are not immunosuppressive, particularly those which are O-substituted at C28 only, such as 28-O-methyl-rapamycin. This can be shown in standard in vitro assays in comparison to FK506 and rapamycin. FK506, for example, is known to be a potent inhibitor of IL-2 transcription, as can be shown in an IL-2 reporter gene assay. Rapamycin, although not active in the IL-2 reporter gene assay, strongly inhibits IL-6 dependent T-cell proliferation. Both compounds are very potent inhibitors of the mixed lymphocyte reaction. Nonimmunosuppressivity can also be shown in the in vivo models 1–7 above. Even those Novel Compounds which are not immunosuppressive, however, bind to macrophilin, which confers certain utilities in which nonimmunosuppressivity is an advantage.

Those of the Novel Compounds which bind strongly to macrophilin and are not themselves immunosuppressive can be used in the treatment of overdoses of macrophilin-binding immunosuppressants, such as FK506, rapamycin, and the immunosuppressive Novel Compounds.

10. Steroid potentiation

The macrophilin binding activity of the Novel Compounds also makes them useful in enhancing or potentiating the action of corticosteroids. Combined treatment with the

dexamethasone, results in greatly enhanced steroidal activity. This can be shown, e.g., in the murine mammary tumor virus-chloramphenicol acetyltransferase (MMTV-CAT) reporter gene assay, e.g., as described in Ning, et al., *J. Biol. Chem.* (1993) 268: 6073. This synergistic effect allows reduced doses of corticosteroids, thereby reducing the risk of side effects in some cases.

11. Mip and Mip-like factor inhibition

Additionally, the Novel Compounds bind to and block a variety of Mip (macrophage infectivity potentiator) and Mip-like factors, which are structurally similar to macrophilin. Mip and Mip-like factors are virulence factors produced by a wide variety of pathogens, including those of the genera Chlamidia, e.g., *Chlamidia trachomatis*; Neisseria, e.g., *Neisseria meningitidis*; and Legionella, e.g., *Legionella pneumophila*; and also by the obligately parasitic members of the order Rickettsiales. These factors play a critical role in the establishment of intracellular infection. The efficacy of the Novel Compounds in reducing the infectivity of pathogens which produce Mip or Mip-like factors can be shown by comparing infectivity of the pathogens in cells culture in the presence and absence of the macrolides, e.g., using the methods described in Lundemose, et al., *Mol. Microbiol.* (1993) 7: 777. The nonimmunosuppressive compounds of the invention are preferred for use in this indication for the reason that they are not immunosuppressive, thus they do not compromise the body's natural immune defenses against the pathogens.

The Novel Compounds are also useful in assays to detect the presence or amount of macrophilin-binding compounds, e.g., in competitive assays for diagnostic or screening purposes. Thus, in another embodiment, the invention provides for use of the Novel Compounds as a screening tool to determine the presence of macrophilin-binding compounds in a test solution, e.g., blood, blood serum, or test broth to be screened. Preferably, a Novel Compound is immobilized in microtiter wells and then allowed to bind in the presence and absence of a test solution to labelled macrophilin-12 (FKBP-12). Alternatively, the FKBP-12 immobilized in microtiter wells and allowed to bind in the presence and absence of a test solution to a Novel Compound which has been labelled, e.g., fluoro-, enzymatically- or radio-labelled, e.g., a Novel Compound which has been O-substituted at C40 and/or C28 with a labelling group. The plates are washed and the amount of bound labelled compound is measured. The amount of macrophilin-binding substance in the test solution is roughly inversely proportional to the amount of bound labelled compound. For quantitative analysis, a standard binding curve is made using known concentrations of macrophilin bind compound.

EXAMPLES:

In the following examples, characteristic spectroscopic data is given to facilitate identification. Peaks which do not differ significantly from rapamycin are not included. Biological data is expressed as a relative IC₅₀, compared to rapamycin in the case of the mixed lymphocyte reaction (MLR) and IL-6 dependent proliferation (IL-6 dep. prol.) assays, and to FK-506 in the macrophilin binding assay (MBA). A higher IC₅₀ correlates with lower binding affinity. Example 1: 40-O-Benzyl-rapamycin

To a stirred solution of 183 mg (0.200 mmol) of rapamycin in 2.1 mL of 2:1 cyclo-hexane-methylene chloride is added 75 μ L (0.402 mmol) of benzyl-trichloroacetimidate, followed by 2.6 μ L (29 μ mol 15 mol %) of trifluoromethanesulfonic acid whereupon the mixture turned immediately

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.