

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use INSPRA safely and effectively. See full prescribing information for INSPRA.

INSPRA (eplerenone) tablets  
Initial U.S. Approval: 2002

### INDICATIONS AND USAGE

INSPRA is an aldosterone antagonist indicated for:

- Improving survival of stable patients with LV systolic dysfunction (LVEF  $\leq$ 40%) and CHF after an acute myocardial infarction. (1.2)
- Hypertension, alone or combined with other agents. (1.3)

### DOSAGE AND ADMINISTRATION

**CHF Post-MI:** Initiate treatment with 25 mg once daily. Titrate to maximum of 50 mg once daily within 4 weeks, as tolerated. Dose adjustments may be required based on potassium levels. (2.1)

**Hypertension:** 50 mg once daily, alone or combined with other antihypertensive agents. For inadequate response, increase to 50 mg twice daily. Higher dosages are not recommended. (2.2)

#### For all patients:

Measure serum potassium before starting INSPRA and periodically thereafter. (2.3)

### DOSAGE FORMS AND STRENGTHS

Tablets: 25 mg, 50 mg (3)

### CONTRAINDICATIONS

#### For all patients:

- Serum potassium  $>$ 5.5 mEq/L at initiation (4)
- Creatinine clearance  $\leq$ 30 mL/min (4)

- Concomitant use with strong CYP3A4 inhibitors (4, 7.1)  
*For the treatment of hypertension:*
- Type 2 diabetes with microalbuminuria (4)
- Serum creatinine  $>$ 2.0 mg/dL in males,  $>$ 1.8 mg/dL in females (4)
- Creatinine clearance  $<$ 50 mL/min (4)
- Concomitant use of potassium supplements or potassium-sparing diuretics (4)

### WARNINGS AND PRECAUTIONS

- Hyperkalemia:** Patients with decreased renal function and diabetics with proteinuria are at increased risk. Proper patient selection and monitoring and avoiding certain concomitant medications can minimize the risk. (5.1)

### ADVERSE REACTIONS

**CHF Post-MI:** Most common adverse reactions ( $>$ 2% and more frequent than with placebo): hyperkalemia and increased creatinine. (6.1)

**Hypertension:** Most common adverse reactions ( $\geq$ 2% and more frequent than with placebo): dizziness, diarrhea, coughing, fatigue and flu-like symptoms. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc at 1-800-438-1985 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- CYP3A4 Inhibitors: Reduce the starting dose for hypertension to 25 mg once daily when used with moderate CYP3A4 inhibitors (e.g., verapamil, erythromycin, saquinavir, fluconazole). (2.4, 7.1, 12.3)

See 17 for PATIENT COUNSELING INFORMATION

Revised: APR 2008

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- Patient Selection Considerations
- Congestive Heart Failure Post-Myocardial Infarction
- Hypertension

### 2 DOSAGE AND ADMINISTRATION

- Congestive Heart Failure Post-Myocardial Infarction
- Hypertension
- Recommended Monitoring
- Dose Modifications for Specific Populations

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- Hyperkalemia
- Impaired Hepatic Function
- Impaired Renal Function

### 6 ADVERSE REACTIONS

- Clinical Trials Experience
- Clinical Laboratory Test Findings
- Postmarketing Experience

### 7 DRUG INTERACTIONS

- CYP3A4 Inhibitors
- ACE Inhibitors and Angiotensin II Receptor Antagonists
- Lithium

7.4 Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

### 8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Nursing Mothers
- Pediatric Use
- Geriatric Use

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacokinetics

### 13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14 CLINICAL STUDIES

- Congestive Heart Failure Post-Myocardial Infarction
- Hypertension

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 Patient Selection Considerations

Serum potassium levels should be measured before initiating INSPRA therapy, and INSPRA should not be prescribed if serum potassium is  $>5.5$  mEq/L. [See *CONTRAINDICATIONS* (4).]

#### 1.2 Congestive Heart Failure Post-Myocardial Infarction

INSPRA is indicated to improve survival of stable patients with left ventricular (LV) systolic dysfunction (ejection fraction  $\leq 40\%$ ) and clinical evidence of congestive heart failure (CHF) after an acute myocardial infarction (MI).

#### 1.3 Hypertension

INSPRA is indicated for the treatment of hypertension. INSPRA may be used alone or in combination with other antihypertensive agents.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Congestive Heart Failure Post-Myocardial Infarction

Treatment should be initiated at 25 mg once daily and titrated to the recommended dose of 50 mg once daily, preferably within 4 weeks as tolerated by the patient. INSPRA may be administered with or without food.

Once treatment with INSPRA has begun, adjust the dose based on the serum potassium level as shown in Table 1.

**Table 1. Dose Adjustment in Congestive Heart Failure Post-MI**

| Serum Potassium (mEq/L) | Action   | Dose Adjustment  |
|-------------------------|----------|--|
| $< 5.0$                 | Increase | 25 mg every other day to 25 mg once daily<br>25 mg once daily to 50 mg once daily                                      |
| 5.0–5.4                 | Maintain | No adjustment  |
| 5.5–5.9                 | Decrease | 50 mg once daily to 25 mg once daily<br>25 mg once daily to 25 mg every other day<br>25 mg every other day to withhold |
| $\geq 6.0$              | Withhold | Restart at 25 mg every other day when potassium levels fall to $<5.5$ mEq/L  |

#### 2.2 Hypertension

The recommended starting dose of INSPRA is 50 mg administered once daily. The full therapeutic effect of INSPRA is apparent within 4 weeks. For patients with an inadequate blood pressure response to 50 mg once daily the dosage of INSPRA should be increased to 50 mg twice daily. Higher dosages of INSPRA are not recommended because they have no greater effect on blood pressure than 100 mg and are associated with an increased risk of hyperkalemia. [See *CLINICAL STUDIES* (14.2).]

#### 2.3 Recommended Monitoring

Serum potassium should be measured before initiating INSPRA therapy, within the first week, and at one month after the start of treatment or dose adjustment. Serum potassium should be assessed periodically thereafter. Patient characteristics and serum potassium levels may indicate that additional monitoring is appropriate. [See *WARNINGS AND PRECAUTIONS* (5.1), *ADVERSE REACTIONS* (6.2).] In the EPHEBUS study [See *CLINICAL STUDIES* (14.1)], the majority of hyperkalemia was observed within the first three months after randomization.

In all patients taking INSPRA who start taking a moderate CYP3A4 inhibitor, check serum potassium and serum creatinine in 3-7 days.

#### 2.4 Dose Modifications for Specific Populations

For hypertensive patients receiving moderate CYP3A4 inhibitors (e.g., erythromycin, saquinavir, verapamil, and fluconazole), the starting dose of INSPRA should be reduced to 25 mg once daily. [See *DRUG INTERACTIONS* (7.1).]

No adjustment of the starting dose is recommended for the elderly or for patients with mild-to-moderate hepatic impairment. [See *CLINICAL PHARMACOLOGY* (12.3).]

### 3 DOSAGE FORMS AND STRENGTHS

- 25 mg tablets: yellow diamond biconvex film-coated tablets debossed with *Pfizer* on one side and *NSR over 25* on the other
- 50 mg tablets: yellow diamond biconvex film-coated tablets debossed with *Pfizer* on one side and *NSR over 50* on the other

### 4 CONTRAINDICATIONS

#### *For All Patients*

INSPRA is contraindicated in all patients with:

- serum potassium  $>5.5$  mEq/L at initiation,
- creatinine clearance  $\leq 30$  mL/min, or
- concomitant administration of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, nefazodone, troleandomycin, clarithromycin, ritonavir, and nelfinavir). [See *DRUG INTERACTIONS* (7.1), *CLINICAL PHARMACOLOGY* (12.3).]

#### *For Patients Treated for Hypertension*

INSPRA is contraindicated for the treatment of hypertension in patients with:

- type 2 diabetes with microalbuminuria,
- serum creatinine  $>2.0$  mg/dL in males or  $>1.8$  mg/dL in females,
- creatinine clearance  $<50$  mL/min, or
- concomitant administration of potassium supplements or potassium-sparing diuretics (e.g., amiloride, spironolactone, or triamterene). [See *WARNINGS AND PRECAUTIONS* (5.1), *ADVERSE REACTIONS* (6.2), *DRUG INTERACTIONS* (7), and *CLINICAL PHARMACOLOGY* (12.3).]

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hyperkalemia

Minimize the risk of hyperkalemia with proper patient selection and monitoring, and avoidance of certain concomitant medications [See *CONTRAINDICATIONS* (4), *ADVERSE REACTIONS* (6.2), and *DRUG INTERACTIONS* (7)]. Monitor patients for the development of hyperkalemia until the effect of INSPRA is established. Patients who develop hyperkalemia ( $>5.5$  mEq/L) may continue INSPRA therapy with proper dose adjustment. Dose reduction decreases potassium levels. [See *DOSAGE AND ADMINISTRATION* (2.1).]

The rates of hyperkalemia increase with declining renal function. [See *ADVERSE REACTIONS* (6.2).] Patients with hypertension who have serum creatinine levels  $>2.0$  mg/dL (males) or  $>1.8$  mg/dL (females) or creatinine clearance  $\leq 50$  mL/min should not be treated with INSPRA. [See *CONTRAINDICATIONS* (4).] Patients with CHF post-MI who have serum creatinine levels  $>2.0$  mg/dL (males) or  $>1.8$  mg/dL (females) or creatinine clearance  $\leq 50$  mL/min should be treated with INSPRA with caution.

Diabetic patients with CHF post-MI should also be treated with caution, especially those with proteinuria. The subset of patients in the EPHEBUS study with both diabetes and proteinuria on the baseline urinalysis had increased rates of hyperkalemia compared to patients with either diabetes or proteinuria. [See *ADVERSE REACTIONS* (6.2).]

#### 5.2 Impaired Hepatic Function

Mild-to-moderate hepatic impairment did not increase the incidence of hyperkalemia. In 16 subjects with mild-to-moderate hepatic impairment who received 400 mg of eplerenone, no elevations of serum potassium above 5.5 mEq/L were observed. The mean increase in serum potassium was 0.12 mEq/L in patients with hepatic impairment and 0.13 mEq/L in normal controls. The use of INSPRA in patients with severe hepatic impairment has not been evaluated. [See *CLINICAL PHARMACOLOGY* (12.3).]

#### 5.3 Impaired Renal Function

Patients with decreased renal function are at increased risk of hyperkalemia. [See *CONTRAINDICATIONS* (4), *WARNINGS AND PRECAUTIONS* (5.1), *ADVERSE REACTIONS* (6.1).]

### 6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Hyperkalemia [See *WARNINGS AND PRECAUTIONS* (5.1)]

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

#### 6.1 Clinical Trials Experience

##### *Congestive Heart Failure Post-Myocardial Infarction*

In EPHEBUS, safety was evaluated in 3307 patients treated with INSPRA and 3301 placebo-treated patients. The overall incidence of adverse events reported with INSPRA (78.9%) was similar to placebo (79.5%). Adverse events occurred at a similar rate regardless of age, gender, or race. Patients discontinued treatment due to an adverse event at

similar rates in either treatment group (4.4% INSPRA vs. 4.3% placebo), with the most common reasons for discontinuation being hyperkalemia, myocardial infarction, and abnormal renal function.

Adverse reactions that occurred more frequently in patients treated with INSPRA than placebo were hyperkalemia (3.4% vs. 2.0%) and increased creatinine (2.4% vs. 1.5%). Discontinuations due to hyperkalemia or abnormal renal function were less than 1.0% in both groups. Hypokalemia occurred less frequently in patients treated with INSPRA (0.6% vs. 1.6%).

The rates of sex hormone-related adverse events are shown in Table 2.

**Table 2. Rates of Sex Hormone-Related Adverse Events in EPHEUS**

|         | Rates in Males |            |        | Rates in Females          |
|---------|----------------|------------|--------|---------------------------|
|         | Gynecomastia   | Mastodynia | Either | Abnormal Vaginal Bleeding |
| INSPRA  | 0.4%           | 0.1%       | 0.5%   | 0.4%                      |
| Placebo | 0.5%           | 0.1%       | 0.6%   | 0.4%                      |

**Hypertension**

INSPRA has been evaluated for safety in 3091 patients treated for hypertension. A total of 690 patients were treated for over 6 months and 106 patients were treated for over 1 year.

In placebo-controlled studies, the overall rates of adverse events were 47% with INSPRA and 45% with placebo. Adverse events occurred at a similar rate regardless of age, gender, or race. Therapy was discontinued due to an adverse event in 3% of patients treated with INSPRA and 3% of patients given placebo. The most common reasons for discontinuation of INSPRA were headache, dizziness, angina pectoris/myocardial infarction, and increased GGT. The adverse events that were reported at a rate of at least 1% of patients and at a higher rate in patients treated with INSPRA in daily doses of 25 to 400 mg versus placebo are shown in Table 3.

**Table 3. Rates (%) of Adverse Events Occurring in Placebo-Controlled Hypertension Studies in ≥1% of Patients Treated with INSPRA (25 to 400 mg) and at a More Frequent Rate than in Placebo-Treated Patients**

|  | INSPRA (n=945) | Placebo (n=372) |
|--|----------------|-----------------|
| <b>Metabolic</b>                         |                |                 |
| Hypercholesterolemia                     | 1              | 0               |
| Hypertriglyceridemia                     | 1              | 0               |
| <b>Digestive</b>                         |                |                 |
| Diarrhea                                 | 2              | 1               |
| Abdominal pain                           | 1              | 0               |
| <b>Urinary</b>                           |                |                 |
| Albuminuria                              | 1              | 0               |
| <b>Respiratory</b>                       |                |                 |
| Coughing                                 | 2              | 1               |
| <b>Central/Peripheral Nervous System</b> |                |                 |
| Dizziness                                | 3              | 2               |
| <b>Body as a Whole</b>                   |                |                 |
| Fatigue                                  | 2              | 1               |
| Influenza-like symptoms                  | 2              | 1               |

Note: Adverse events that are too general to be informative or are very common in the treated population are excluded.

Gynecomastia and abnormal vaginal bleeding were reported with INSPRA but not with placebo. The rates of these sex hormone-related adverse events are shown in Table 4. The rates increased slightly with increasing duration of therapy. In females, abnormal vaginal bleeding was also reported in 0.8% of patients on antihypertensive medications (other than spironolactone) in active control arms of the studies with INSPRA.

**Table 4. Rates of Sex Hormone-Related Adverse Events with INSPRA in Hypertension Clinical Studies**

|                                       | Rates in Males |            |        | Rates in Females          |
|---------------------------------------|----------------|------------|--------|---------------------------|
|                                       | Gynecomastia   | Mastodynia | Either | Abnormal Vaginal Bleeding |
| All controlled studies                | 0.5%           | 0.8%       | 1.0%   | 0.6%                      |
| Controlled studies lasting ≥ 6 months | 0.7%           | 1.3%       | 1.6%   | 0.8%                      |
| Open-label, long-term study           | 1.0%           | 0.3%       | 1.0%   | 2.1%                      |

**6.2 Clinical Laboratory Test Findings**

**Congestive Heart Failure Post-Myocardial Infarction**

**Creatinine:** Increases of more than 0.5 mg/dL were reported for 6.5% of patients administered INSPRA and for 4.9% of placebo-treated patients.

**Potassium:** In EPHEUS [see CLINICAL STUDIES (14.1)], the frequencies of patients with changes in potassium (<3.5 mEq/L or >5.5 mEq/L or ≥6.0 mEq/L) receiving INSPRA compared with placebo are displayed in Table 5.

**Table 5. Hypokalemia (<3.5 mEq/L) or Hyperkalemia (>5.5 or ≥6.0 mEq/L) in EPHEUS**

| Potassium (mEq/L) | INSPRA (N=3251)<br>n (%) | Placebo (N=3237)<br>n (%) |
|-------------------|--------------------------|---------------------------|
| < 3.5             | 273 (8.4)                | 424 (13.1)                |
| >5.5              | 508 (15.6)               | 363 (11.2)                |
| ≥ 6.0             | 180 (5.5)                | 126 (3.9)                 |

Table 6 shows the rates of hyperkalemia in EPHEUS as assessed by baseline renal function (creatinine clearance).

**Table 6. Rates of Hyperkalemia (>5.5 mEq/L) in EPHEUS by Baseline Creatinine Clearance\***

| Baseline Creatinine Clearance | INSPRA (N=508)<br>n (%) | Placebo (N=363)<br>n (%) |
|-------------------------------|-------------------------|--------------------------|
| ≤30 mL/min                    | 160 (32)                | 82 (23)                  |
| 31–50 mL/min                  | 122 (24)                | 46 (13)                  |
| 51–70 mL/min                  | 86 (17)                 | 48 (13)                  |
| >70 mL/min                    | 56 (11)                 | 32 (9)                   |

\* Estimated using the Cockcroft-Gault formula.

Table 7 shows the rates of hyperkalemia in EPHEUS as assessed by two baseline characteristics: presence/absence of proteinuria from baseline urinalysis and presence/absence of diabetes. [See WARNINGS AND PRECAUTIONS (5.1).]

**Table 7. Rates of Hyperkalemia (>5.5 mEq/L) in EPHEUS by Proteinuria and History of Diabetes\***

|                          | INSPRA (N=508)<br>n (%) | Placebo (N=363)<br>n (%) |
|--------------------------|-------------------------|--------------------------|
| Proteinuria, no Diabetes | 81 (16)                 | 40 (11)                  |
| Diabetes, no Proteinuria | 91 (18)                 | 47 (13)                  |
| Proteinuria and Diabetes | 132 (26)                | 58 (16)                  |

\* Diabetes assessed as positive medical history at baseline; proteinuria assessed by positive dipstick urinalysis at baseline.

**Hypertension**

**Potassium:** In placebo-controlled fixed-dose studies, the mean increases in serum potassium were dose-related and are shown in Table 8 along with the frequencies of values >5.5 mEq/L.

**Table 8. Increases in Serum Potassium in the Placebo-Controlled, Fixed-Dose Hypertension Studies of INSPRA**

| Daily Dosage | n   | Mean Increase mEq/L | % >5.5 mEq/L |
|--------------|-----|---------------------|--------------|
| Placebo      | 194 | 0                   | 1            |
| 25           | 97  | 0.08                | 0            |
| 50           | 245 | 0.14                | 0            |
| 100          | 193 | 0.09                | 1            |
| 200          | 139 | 0.19                | 1            |
| 400          | 104 | 0.36                | 8.7          |

Patients with both type 2 diabetes and microalbuminuria are at increased risk of developing persistent hyperkalemia. In a study of such patients taking INSPRA 200 mg, the frequencies of maximum serum potassium levels >5.5 mEq/L were 33% with INSPRA given alone and 38% when INSPRA was given with enalapril.

Rates of hyperkalemia increased with decreasing renal function. In all studies, serum potassium elevations >5.5 mEq/L were observed in 10.4% of patients treated with INSPRA with baseline calculated creatinine clearance <70 mL/min, 5.6% of patients with baseline creatinine clearance of 70 to 100 mL/min, and 2.6% of patients with baseline creatinine clearance of >100 mL/min. [See **WARNINGS AND PRECAUTIONS (5.1).**]

**Sodium:** Serum sodium decreased in a dose-related manner. Mean decreases ranged from 0.7 mEq/L at 50 mg daily to 1.7 mEq/L at 400 mg daily. Decreases in sodium (<135 mEq/L) were reported for 2.3% of patients administered INSPRA and 0.6% of placebo-treated patients.

**Triglycerides:** Serum triglycerides increased in a dose-related manner. Mean increases ranged from 7.1 mg/dL at 50 mg daily to 26.6 mg/dL at 400 mg daily. Increases in triglycerides (above 252 mg/dL) were reported for 15% of patients administered INSPRA and 12% of placebo-treated patients.

**Cholesterol:** Serum cholesterol increased in a dose-related manner. Mean changes ranged from a decrease of 0.4 mg/dL at 50 mg daily to an increase of 11.6 mg/dL at 400 mg daily. Increases in serum cholesterol values greater than 200 mg/dL were reported for 0.3% of patients administered INSPRA and 0% of placebo-treated patients.

**Liver Function Tests:** Serum alanine aminotransferase (ALT) and gamma glutamyl transpeptidase (GGT) increased in a dose-related manner. Mean increases ranged from 0.8 U/L at 50 mg daily to 4.8 U/L at 400 mg daily for ALT and 3.1 U/L at 50 mg daily to 11.3 U/L at 400 mg daily for GGT. Increases in ALT levels greater than 120 U/L (3 times upper limit of normal) were reported for 15/2259 patients administered INSPRA and 1/351 placebo-treated patients. Increases in ALT levels greater than 200 U/L (5 times upper limit of normal) were reported for 5/2259 of patients administered INSPRA and 1/351 placebo-treated patients. Increases of ALT greater than 120 U/L and bilirubin greater than 1.2 mg/dL were reported 1/2259 patients administered INSPRA and 0/351 placebo-treated patients. Hepatic failure was not reported in patients receiving INSPRA.

**BUN/Creatinine:** Serum creatinine increased in a dose-related manner. Mean increases ranged from 0.01 mg/dL at 50 mg daily to 0.03 mg/dL at 400 mg daily. Increases in blood urea nitrogen to greater than 30 mg/dL and serum creatinine to greater than 2 mg/dL were reported for 0.5% and 0.2%, respectively, of patients administered INSPRA and 0% of placebo-treated patients.

**Uric Acid:** Increases in uric acid to greater than 9 mg/dL were reported in 0.3% of patients administered INSPRA and 0% of placebo-treated patients.

### 6.3 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of INSPRA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Skin:* angioneurotic edema, rash

## 7 DRUG INTERACTIONS

### 7.1 CYP3A4 Inhibitors

Because eplerenone metabolism is predominantly mediated via CYP3A4, do not use INSPRA with drugs that are strong inhibitors of CYP3A4. [See **CONTRAINDICATIONS (4)** and **CLINICAL PHARMACOLOGY (12.3).**]

In patients with hypertension taking moderate CYP3A4 inhibitors, reduce the starting dose of INSPRA to 25 mg once daily. [See **DOSAGE**

AND **ADMINISTRATION (2.3, 2.4)** and **CLINICAL PHARMACOLOGY (12.3).**]

### 7.2 ACE Inhibitors and Angiotensin II Receptor Antagonists

#### *Congestive Heart Failure Post-Myocardial Infarction*

In EPHEBUS [see **CLINICAL STUDIES (14.1)**], 3020 (91%) patients receiving INSPRA 25 to 50 mg also received ACE inhibitors or angiotensin II receptor antagonists (ACEI/ARB). Rates of patients with maximum potassium levels >5.5 mEq/L were similar regardless of the use of ACEI/ARB.

#### *Hypertension*

In clinical studies of patients with hypertension, the addition of INSPRA 50 to 100 mg to ACE inhibitors and angiotensin II receptor antagonists increased mean serum potassium slightly (about 0.09–0.13 mEq/L). In a study in diabetics with microalbuminuria, INSPRA 200 mg combined with the ACE inhibitor enalapril 10 mg increased the frequency of hyperkalemia (serum potassium >5.5 mEq/L) from 17% on enalapril alone to 38%.

### 7.3 Lithium

A drug interaction study of eplerenone with lithium has not been conducted. Lithium toxicity has been reported in patients receiving lithium concomitantly with diuretics and ACE inhibitors. Serum lithium levels should be monitored frequently if INSPRA is administered concomitantly with lithium.

### 7.4 Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

A drug interaction study of eplerenone with an NSAID has not been conducted. The administration of other potassium-sparing antihypertensives with NSAIDs has been shown to reduce the antihypertensive effect in some patients and result in severe hyperkalemia in patients with impaired renal function. Therefore, when INSPRA and NSAIDs are used concomitantly, patients should be observed to determine whether the desired effect on blood pressure is obtained and monitored for changes in serum potassium levels.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B

There are no adequate and well-controlled studies in pregnant women. INSPRA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

#### *Teratogenic Effects*

Embryo-fetal development studies were conducted with doses up to 1000 mg/kg/day in rats and 300 mg/kg/day in rabbits (exposures up to 32 and 31 times the human AUC for the 100 mg/day therapeutic dose, respectively). No teratogenic effects were seen in rats or rabbits, although decreased body weight in maternal rabbits and increased rabbit fetal resorptions and post-implantation loss were observed at the highest administered dosage. Because animal reproduction studies are not always predictive of human response, INSPRA should be used during pregnancy only if clearly needed.

### 8.3 Nursing Mothers

The concentration of eplerenone in human breast milk after oral administration is unknown. However, preclinical data show that eplerenone and/or metabolites are present in rat breast milk (0.85:1 [milk:plasma] AUC ratio) obtained after a single oral dose. Peak concentrations in plasma and milk were obtained from 0.5 to 1 hour after dosing. Rat pups exposed by this route developed normally. Because many drugs are excreted in human milk and because of the unknown potential for adverse effects on the nursing infant, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

### 8.4 Pediatric Use

In a 10-week study of 304 hypertensive pediatric patients age 4 to 17 years treated with INSPRA up to 100 mg per day, doses that produced exposure similar to that in adults, INSPRA did not lower blood pressure effectively. In this study and in a 1-year pediatric safety study in 149 patients, the incidence of reported adverse events was similar to that of adults.

INSPRA has not been studied in hypertensive patients less than 4 years old because the study in older pediatric patients did not demonstrate effectiveness.

INSPRA has not been studied in pediatric patients with heart failure.

### 8.5 Geriatric Use

Of the total number of patients in EPHEBUS, 3340 (50%) were 65 and over, while 1326 (20%) were 75 and over. Patients greater than 75 years did not appear to benefit from the use of INSPRA. [See *CLINICAL STUDIES (14.1)*.]

No differences in overall incidence of adverse events were observed between elderly and younger patients. However, due to age-related decreases in creatinine clearance, the incidence of laboratory-documented hyperkalemia was increased in patients 65 and older. [See *WARNINGS AND PRECAUTIONS (5.1)*.]

#### Hypertension

Of the total number of subjects in clinical hypertension studies of INSPRA, 1123 (23%) were 65 and over, while 212 (4%) were 75 and over. No overall differences in safety or effectiveness were observed between elderly subjects and younger subjects.

## 10 OVERDOSAGE


No cases of human overdosage with eplerenone have been reported. Lethality was not observed in mice, rats, or dogs after single oral doses that provided  $C_{max}$  exposures at least 25 times higher than in humans receiving eplerenone 100 mg/day. Dogs showed emesis, salivation, and tremors at a  $C_{max}$  41 times the human therapeutic  $C_{max}$ , progressing to sedation and convulsions at higher exposures.

The most likely manifestation of human overdosage would be anticipated to be hypotension or hyperkalemia. Eplerenone cannot be removed by hemodialysis. Eplerenone has been shown to bind extensively to charcoal. If symptomatic hypotension should occur, supportive treatment should be instituted. If hyperkalemia develops, standard treatment should be initiated.

## 11 DESCRIPTION

INSPRA contains eplerenone, a blocker of aldosterone binding at the mineralocorticoid receptor.

Eplerenone is chemically described as Pregn-4-ene-7,21-dicarboxylic acid, 9,11-epoxy-17-hydroxy-3-oxo-,  $\gamma$ -lactone, methyl ester, (7 $\alpha$ ,11 $\alpha$ ,17 $\alpha$ )-. Its empirical formula is  $C_{24}H_{30}O_6$  and it has a molecular weight of 414.50. The structural formula of eplerenone is represented below:



Eplerenone is an odorless, white to off-white crystalline powder. It is very slightly soluble in water, with its solubility essentially pH-independent. The octanol/water partition coefficient of eplerenone is approximately 7.1 at pH 7.0.

INSPRA for oral administration contains 25 mg or 50 mg of eplerenone and the following inactive ingredients: lactose, microcrystalline cellulose, croscarmellose sodium, hypromellose, sodium lauryl sulfate, talc, magnesium stearate, titanium dioxide, polyethylene glycol, polysorbate 80, and iron oxide yellow and iron oxide red.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Eplerenone binds to the mineralocorticoid receptor and blocks the binding of aldosterone, a component of the renin-angiotensin-aldosterone system (RAAS). Aldosterone synthesis, which occurs primarily in the adrenal gland, is modulated by multiple factors, including angiotensin II and non-RAAS mediators such as adrenocorticotropic hormone (ACTH) and potassium. Aldosterone binds to mineralocorticoid receptors in both epithelial (e.g., kidney) and nonepithelial (e.g., heart, blood vessels, and

brain) tissues and increases blood pressure through induction of sodium reabsorption and possibly other mechanisms.

Eplerenone has been shown to produce sustained increases in plasma renin and serum aldosterone, consistent with inhibition of the negative regulatory feedback of aldosterone on renin secretion. The resulting increased plasma renin activity and aldosterone circulating levels do not overcome the effects of eplerenone.

Eplerenone selectively binds to recombinant human mineralocorticoid receptors relative to its binding to recombinant human glucocorticoid, progesterone, and androgen receptors.

### 12.3 Pharmacokinetics

Eplerenone is cleared predominantly by cytochrome P450 (CYP) 3A4 metabolism, with an elimination half-life of 4 to 6 hours. Steady state is reached within 2 days. Absorption is not affected by food. Inhibitors of CYP3A4 (e.g., ketoconazole, saquinavir) increase blood levels of eplerenone.

#### Absorption and Distribution

Mean peak plasma concentrations of eplerenone are reached approximately 1.5 hours following oral administration. The absolute bioavailability of eplerenone is 69% following administration of a 100 mg oral tablet. Both peak plasma levels ( $C_{max}$ ) and area under the curve (AUC) are dose proportional for doses of 25 to 100 mg and less than proportional at doses above 100 mg.

The plasma protein binding of eplerenone is about 50% and it is primarily bound to alpha 1-acid glycoproteins. The apparent volume of distribution at steady state ranged from 43 to 90 L. Eplerenone does not preferentially bind to red blood cells.

#### Metabolism and Excretion

Eplerenone metabolism is primarily mediated via CYP3A4. No active metabolites of eplerenone have been identified in human plasma.

Less than 5% of an eplerenone dose is recovered as unchanged drug in the urine and feces. Following a single oral dose of radiolabeled drug, approximately 32% of the dose was excreted in the feces and approximately 67% was excreted in the urine. The elimination half-life of eplerenone is approximately 4 to 6 hours. The apparent plasma clearance is approximately 10 L/hr.

#### Age, Gender, and Race

The pharmacokinetics of eplerenone at a dose of 100 mg once daily has been investigated in the elderly ( $\geq 65$  years), in males and females, and in Blacks. At steady state, elderly subjects had increases in  $C_{max}$  (22%) and AUC (45%) compared with younger subjects (18 to 45 years). The pharmacokinetics of eplerenone did not differ significantly between males and females. At steady state,  $C_{max}$  was 19% lower and AUC was 26% lower in Blacks. [See *DOSAGE AND ADMINISTRATION (2.4)* and *USE IN SPECIFIC POPULATIONS (8.5)*.]

#### Renal Impairment

The pharmacokinetics of eplerenone was evaluated in patients with varying degrees of renal impairment and in patients undergoing hemodialysis. Compared with control subjects, steady state AUC and  $C_{max}$  were increased by 38% and 24%, respectively, in patients with severe renal impairment and were decreased by 26% and 3%, respectively, in patients undergoing hemodialysis. No correlation was observed between plasma clearance of eplerenone and creatinine clearance. Eplerenone is not removed by hemodialysis. [See *WARNINGS AND PRECAUTIONS (5.3)*.]

#### Hepatic Impairment

The pharmacokinetics of eplerenone 400 mg has been investigated in patients with moderate (Child-Pugh Class B) hepatic impairment and compared with normal subjects. Steady state  $C_{max}$  and AUC of eplerenone were increased by 3.6% and 42%, respectively.

#### Heart Failure

The pharmacokinetics of eplerenone 50 mg was evaluated in 8 patients with heart failure (NYHA classification II-IV) and 8 matched (gender, age, weight) healthy controls. Compared with the controls, steady state AUC and  $C_{max}$  in patients with stable heart failure were 38% and 30% higher, respectively.

#### Drug-Drug Interactions [See *DRUG INTERACTIONS (7)*.]

Eplerenone is metabolized primarily by CYP3A4. Inhibitors of CYP3A4 cause increased exposure [see *DRUG INTERACTIONS (7.1)*].

Drug-drug interaction studies were conducted with a 100 mg dose of eplerenone.

A pharmacokinetic study evaluating the administration of a single dose of INSPRA 100 mg with ketoconazole 200 mg two times a day, a strong inhibitor of the CYP3A4 pathway, showed a 1.7-fold increase in  $C_{max}$  of eplerenone and a 5.4-fold increase in AUC of eplerenone.

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.