



Español



MENU



CONTACT



DICTIONARY



SEARCH

SECTION MENU

[Home](#) > [Cancer Types](#) > [Prostate Cancer](#)

Prostate-Specific Antigen (PSA) Test



ON THIS PAGE

- [What is the PSA test?](#)
- [Is the PSA test recommended for prostate cancer screening?](#)
- [What is a normal PSA test result?](#)
- [What if a screening test shows an elevated PSA level?](#)
- [What are some of the limitations and potential harms of the PSA test for prostate cancer screening?](#)
- [What research has been done to study prostate cancer screening?](#)
- [How is the PSA test used in men who have been treated for prostate cancer?](#)
- [What does an increase in PSA level mean for a man who has been treated for prostate cancer?](#)
- [How are researchers trying to improve the PSA test?](#)

What is the PSA test?

Prostate-specific antigen, or PSA, is a protein produced by cells of the prostate gland. The PSA test measures the level of PSA in a man's blood. For this test, a blood sample is sent to a laboratory for analysis. The results are usually reported as nanograms of PSA per milliliter (ng/mL) of blood.

The blood level of PSA is often elevated in men with prostate cancer, and the PSA test was originally approved by the FDA in 1986 to monitor the progression of prostate cancer in men who had already been diagnosed with the disease. In 1994, the FDA approved the use of the PSA test in conjunction with a digital rectal exam (DRE) to test asymptomatic men for prostate cancer. Men who report prostate symptoms often

In addition to prostate cancer, a number of benign (not cancerous) conditions can cause a man's PSA level to rise. The most frequent benign prostate conditions that cause an elevation in PSA level are prostatitis (inflammation of the prostate) and benign prostatic hyperplasia (BPH) (enlargement of the prostate). There is no evidence that prostatitis or BPH leads to prostate cancer, but it is possible for a man to have one or both of these conditions and to develop prostate cancer as well.

Is the PSA test recommended for prostate cancer screening?

Until recently, many doctors and professional organizations encouraged yearly PSA screening for men beginning at age 50. Some organizations recommended that men who are at higher risk of prostate cancer, including African American men and men whose father or brother had prostate cancer, begin screening at age 40 or 45. However, as more has been learned about both the benefits and harms of prostate cancer screening, a number of organizations have begun to caution against routine population screening. Although some organizations continue to recommend PSA screening, there is widespread agreement that any man who is considering getting tested should first be informed in detail about the potential harms and benefits.

Currently, [Medicare](#) provides coverage for an annual PSA test for all Medicare-eligible men age 50 and older. Many private insurers cover PSA screening as well.

What is a normal PSA test result?

There is no specific normal or abnormal level of PSA in the blood. In the past, most doctors considered PSA levels of 4.0 ng/mL and lower as normal. Therefore, if a man had a PSA level above 4.0 ng/mL, doctors would often recommend a prostate [biopsy](#) to determine whether prostate cancer was present.

However, more recent studies have shown that some men with PSA levels below 4.0 ng/mL have prostate cancer and that many men with higher levels do not have prostate cancer (1). In addition, various factors can cause a man's PSA level to fluctuate. For example, a man's PSA level often rises if he has [prostatitis](#) or a [urinary tract](#) infection. Prostate biopsies and prostate surgery also increase PSA level. Conversely, some drugs—including [finasteride](#) and [dutasteride](#), which are used to treat [BPH](#)—lower a man's PSA level. PSA level may also vary somewhat across testing laboratories.

Another complicating factor is that studies to establish the normal range of PSA levels have been conducted primarily in populations of white men. Although expert opinions vary, there is no clear consensus regarding the optimal PSA threshold for recommending a prostate biopsy for men of any racial or ethnic group.

In general, however, the higher a man's PSA level, the more likely it is that he has prostate cancer. Moreover, continuous rise in a man's PSA level over time may also be a sign of prostate cancer.

What if a screening test shows an elevated PSA level?

If a man who has no symptoms of prostate cancer chooses to undergo prostate cancer screening and is found to have an elevated PSA level, the doctor may recommend another PSA test to confirm the original finding. If the PSA level is still high, the doctor may recommend that the man continue with PSA tests and DREs at regular intervals to watch for any changes over time.

If a man's PSA level continues to rise or if a suspicious lump is detected during a DRE, the doctor may recommend additional tests to determine the nature of the problem. A urine test may be recommended to check for a urinary tract infection. The doctor may also recommend imaging tests, such as a transrectal ultrasound, x-rays, or cystoscopy.

If prostate cancer is suspected, the doctor will recommend a prostate biopsy. During this procedure, multiple samples of prostate tissue are collected by inserting hollow needles into the prostate and then withdrawing them. Most often, the needles are inserted through the wall of the rectum (transrectal biopsy); however, the needles may also be inserted through the skin between the scrotum and the anus (transperineal biopsy). A pathologist then examines the collected tissue under a microscope. The doctor may use ultrasound to view the prostate during the biopsy, but ultrasound cannot be used alone to diagnose prostate cancer.

What are some of the limitations and potential harms of the PSA test for prostate cancer screening?

Detecting prostate cancer early may not reduce the chance of dying from prostate cancer. When used in screening, the PSA test can help detect small tumors that do not cause symptoms. Finding a small tumor, however, may not necessarily reduce a man's chance of dying from prostate cancer. Some tumors found through PSA testing grow so slowly that they are unlikely to threaten a man's life. Detecting tumors that are not life threatening is called "overdiagnosis," and treating these tumors is called "overtreatment."

Overtreatment exposes men unnecessarily to the potential complications and harmful side effects of treatments for early prostate cancer, including surgery and radiation therapy. The side effects of these treatments include urinary incontinence (inability to control urine flow), problems with bowel function, erectile dysfunction (loss of erections, or having erections that are inadequate for sexual intercourse), and infection.

In addition, finding cancer early may not help a man who has a fast-growing or aggressive tumor that may have spread to other parts of the body before being detected.

The PSA test may give false-positive or false-negative results. A false-positive test result occurs when a man's PSA level is elevated but no cancer is actually present. A false-positive test result may create

can be harmful. Possible side effects of biopsies include serious infections, pain, and bleeding.

Most men with an elevated PSA level turn out not to have prostate cancer; only about 25 percent of men who have a prostate biopsy due to an elevated PSA level actually have prostate cancer (2).

A false-negative test result occurs when a man's PSA level is low even though he actually has prostate cancer. False-negative test results may give a man, his family, and his doctor false assurance that he does not have cancer, when he may in fact have a cancer that requires treatment.

What research has been done to study prostate cancer screening?


Several randomized trials of prostate cancer screening have been carried out. One of the largest is the Prostate, Lung, Colorectal, and Ovarian (PLCO) Cancer Screening Trial, which NCI conducted to determine whether certain screening tests can help reduce the numbers of deaths from several common cancers. In the prostate portion of the trial, the PSA test and DRE were evaluated for their ability to decrease a man's chances of dying from prostate cancer.

The PLCO investigators found that men who underwent annual prostate cancer screening had a higher incidence of prostate cancer than men in the control group but the same rate of deaths from the disease (3). Overall, the results suggest that many men were treated for prostate cancers that would not have been detected in their lifetime without screening. Consequently, these men were exposed unnecessarily to the potential harms of treatment.

A second large trial, the European Randomized Study of Screening for Prostate Cancer (ERSPC), compared prostate cancer deaths in men randomly assigned to PSA-based screening or no screening. As in the PLCO, men in ERSPC who were screened for prostate cancer had a higher incidence of the disease than control men. In contrast to the PLCO, however, men who were screened had a lower rate of death from prostate cancer (4).

The United States Preventive Services Task Force has analyzed the data from the PLCO, ERSPC, and other trials and estimated that, for every 1,000 men ages 55 to 69 years who are screened every 1 to 4 years for a decade (5):

- 0 to 1 death from prostate cancer would be avoided.
- 100 to 120 men would have a false-positive test result that leads to a biopsy, and about one-third of the men who get a biopsy would experience at least moderately bothersome symptoms from the biopsy.
- 110 men would be diagnosed with prostate cancer. About 50 of these



incontinence in 18 men, serious cardiovascular events in 2 men, deep vein thrombosis or pulmonary embolism in 1 man, and death due to the treatment in less than 1 man.

How is the PSA test used in men who have been treated for prostate cancer?

The PSA test is used to monitor patients who have a history of prostate cancer to see if their cancer has recurred (come back). If a man's PSA level begins to rise after prostate cancer treatment, it may be the first sign of a recurrence. Such a "biochemical relapse" typically appears months or years before other clinical signs and symptoms of prostate cancer recurrence.

However, a single elevated PSA measurement in a patient who has a history of prostate cancer does not always mean that the cancer has come back. A man who has been treated for prostate cancer should discuss an elevated PSA level with his doctor. The doctor may recommend repeating the PSA test or performing other tests to check for evidence of a recurrence. The doctor may look for a trend of rising PSA level over time rather than a single elevated PSA level.

What does an increase in PSA level mean for a man who has been treated for prostate cancer?

If a man's PSA level rises after prostate cancer treatment, his doctor will consider a number of factors before recommending further treatment. Additional treatment based on a single PSA test is not recommended. Instead, a rising trend in PSA level over time in combination with other findings, such as an abnormal result on imaging tests, may lead a man's doctor to recommend further treatment.

How are researchers trying to improve the PSA test?

Scientists are investigating ways to improve the PSA test to give doctors the ability to better distinguish cancerous from benign conditions and slow-growing cancers from fast-growing, potentially lethal cancers.

Some of the methods being studied include:

- **Free versus total PSA.** The amount of PSA in the blood that is "free" (not bound to other proteins) divided by the total amount of PSA (free plus bound). Some evidence suggests that a lower proportion of free PSA may be associated with more aggressive cancer.
- **PSA density of the transition zone.** The blood level of PSA divided by the volume of the transition zone of the prostate. The transition zone is the interior part of the prostate that surrounds the urethra. Some evidence suggests that this measure may be more accurate at detecting prostate cancer than the standard PSA test.
- **Age-specific PSA reference ranges.** Because a man's PSA level tends to increase with age, it has been suggested that the use of age-specific PSA reference ranges may increase the accuracy of PSA

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.