


Expert opinion on therapeutic patents.
Vol. 9, no. 5 (May 1999)
General Collection
772 C9765
Received: 06-22-2000

Expert Opinion on

Therapeutic Patents



Ashley Publications

Vol. 9 No. 5

May 1999

Patent Analysis

Patenting of combinatorial chemistry

Reviews & Updates

A₁ and A₂ adenosine receptor agonists: an overview

Low density lipoprotein receptor up-regulation

NOS inhibitors with cardiovascular therapeutic potential

The inhibition of isoforms of NOS: non-cardiovascular aspects

AMPA receptor antagonists

Neurokinin receptor antagonists

Patenting combinatorial libraries and associated technologies

Recent developments with oxazolidinone antibiotics

Patenting Perspective

Developing trends in pharmaceutical patent claims

Patent Evaluations

Somatostatin antagonists

Polymer-platinum compounds


Hepatitis C NS3 protease inhibitors



PROPERTY OF THE
NATIONAL

Expert Opinion on Therapeutic Patents

<http://www.ashley-pub.com>



Senior Advisory Panel

Abou-Gharbia M: *Wyeth-Ayerst, USA*
Arnaud J-C: *Servier, France*
Coombes J: *formerly Hoechst, UK*
Dalgleish A: *St George's Hospital, UK*
de Clercq E: *Rega Institut, Belgium*
Evans C: *Chiroscience, UK*
Fitzgerald JD: *Materia Medica, UK*
Floyd D: *Bristol-Myers Squibb, USA*
Glennon RA: *Virginia Commonwealth University, USA*
Hill CH: *Roche, UK*
Jacobson KA: *NIH, USA*
Krogsgaard-Larsen P: *Danish School of Pharmacy, Denmark*
Moos WH: *Chiron Corp., USA*
Mullen A: *Bayer AG, Germany*
Poste G: *SmithKline Beecham, USA*
Rees D: *Organon, UK*
Ross BC: *Glaxo Wellcome, UK*
Steele PR: *IP Consultant, UK*
Stevens RW: *Pfizer, Japan*
Taylor JB: *Rhône-Poulenc Rorer, UK*
Terrett N: *Pfizer, UK*
Timmermanns P: *DuPont Merck, USA*

Section Editorial Board

Pulmonary-Allergy, Dermatological, Gastrointestinal & Arthritis

Djuric SW, Kaminski JJ, Landry Y, Lipani J, Murthy S, Norman P, Sircar JC, Summers JB, Talley JJ, Trivedi B, Whittaker M

Anti-infectives

Cooper RDG, Croft S, Furman P, Hector RF, Highfield P, Hunter PA, Kirst H, Lee VJ, Ridley R, Rodriguez M, Smith PW

Biologicals, Immunologicals & Drug Delivery

Amos M, Deonarain MP, Epenetos AA, George A, Graf H, Gregoriadis G, Lerner A

Central & Peripheral Nervous Systems

Baudy RB, Bousquet P, Christos T, Cliffe I, Costentin J, Franco R, Halazy S, Hoban C, Howard H, Johnson G, McElroy AB, Spence RP, Williams M

Cardiovascular & Renal

Baldwin JJ, Chakravarty PK, Doherty AM, Grove R, Müller CE, Perone M, Spada AP, Suckling KE, Trybulski E

Oncologic, Endocrine & Metabolic

Baldwin RW, Beeley NRA, Connors T, Dale I, Davis P, Dueholm KL, Ecker G, Fields G, Gallagher JT, Hindley RM, Hughes L, Martini L, Miller WR, Remuzzi G, Rosenberg S, Skibo E, Williams T

Commissioning Editor: Liz Williams

Production Editor: Paula Rhodes

Publisher: James Drake

Aims and scope: *Expert Opinion on Therapeutic Patents* aims to provide an evaluated guide to developments in the recent patent literature. The journal is divided into six therapeutic sections. Each section is reviewed every six months, allowing systematic and comprehensive coverage of the most important topics:

- Pulmonary-Allergy, Dermatological, Gastrointestinal & Arthritis (January & July)
- Anti-infectives (February & August)
- Biologicals, Immunologicals & Drug Delivery (March & September)
- Central & Peripheral Nervous Systems (April & October)
- Cardiovascular & Renal (May & November)
- Oncologic, Endocrine & Metabolic (June & December)

Authors are encouraged to express their *Expert Opinion* of the status of the research under review, rather than to simply review the available data. Each issue of the journal contains annotated patent selections and short patent evaluations which highlight inventions and innovative ideas deemed by relevant experts to be of potential importance.

Editorial policy: Articles published in *Expert Opinion on Therapeutic Patents* are commissioned by the Editor in collaboration with the Section Editorial Board. Authors who wish to contribute should contact the Editor. All reviews are subject to peer-review.

Editorial boards: The Senior Advisory Panel and Section Editorial Board are composed of Senior Scientists involved in drug research and development. The Senior Advisory Panel is responsible for the development of the journal, while the Section Editors are responsible for selecting authors and topics for review to ensure comprehensive coverage of subjects in each therapeutic area.

Citations: *Expert Opinion on Therapeutic Patents* is indexed by *Current Contents®/Life Sciences*, *EMBASE/Excerpta Medica* and *Chemical Abstracts*.

Sources: Permission has been granted by the publisher of the following information source for Ashley Publications to provide its authors with relevant source material for use solely in the preparation of their reviews for *Expert Opinion on Therapeutic Patents*:

Patents Preview (© Derwent Publications)
Patent fast-alert (© Current Drugs Ltd.)

Copyright: Conditions of sale - *Expert Opinion on Therapeutic Patents* may be circulated only to those members of staff who are employed at the site at which the subscription is taken out. Readers are reminded that, under internationally agreed copyright legislation, photocopying of copyright materials is prohibited other than on a limited basis for personal use. This means that making copies of any article published in *Expert Opinion on Therapeutic Patents* is a breach of the law and can be prosecuted.

Whilst every effort is made by the publishers and advisory board to ensure that no inaccurate or misleading data, opinions or statements appear in this journal, they wish to make it clear that the data and opinions appearing herein are the responsibility of the contributor concerned. Accordingly, the publishers, advisory board, editors and their respective employees, officers and agents accept no liability whatsoever for the consequences of any inaccurate or misleading data, opinions or statements.

Subscription information: *Expert Opinion on Therapeutic Patents* is now available to print subscribers in searchable, electronic format as Adobe Acrobat PDF files. For more information contact the Subscriptions Department.

Europe and the rest of the world: Ashley Publications Ltd., 1st Floor The Library, 1 Shepherds Hill, Highgate, London, N6 5QJ, UK. Tel: +44 (0)181 347 5030, Fax: +44 (0)181 347 5040, Email: subscriptions@ashley-pub.com

Subscribers in North America make their payment in US dollars.

Japan: Technomics, Inc., Nihonbashi TM Bldg., 1-8-11 Nihonbashi Horidome-cho, Chuo-ku, Tokyo 103, Japan. Tel: +81 (03) 3666 2952, Fax: +81 (03) 3666 2730.

Subscribers in Japan make their payment in Yen.

Korea: Pharma Koreana Ltd., SL Kangnam, PO Box 99, Seoul 135-600, Korea. Tel: +82 (02) 554 9592, Fax: +82 (02) 557 1290.

Annual subscription rates: *Expert Opinion on Therapeutic Patents*, Volume 9, 12 monthly issues, £1750 / US\$3235 / ¥494,950.



<http://www.ashley-pub.com>

Review

1. Introduction
 2. N-Phenyl oxazolidinones
 3. N-Heteroaryl oxazolidinones
 4. Acyl group variations
 5. Oxazolidinone core unit modifications
 6. Non anti-infective activities
 7. Expert opinion
- Bibliography
Patents

Recent developments with oxazolidinone antibiotics

Bernd Riedl & Rainer Endermann

Bayer AG, Pharma Research Centre, D-42096 Wuppertal, Germany

The emerging problems with multiple antibiotic-resistant Gram-positive cocci led to the re-evaluation of an antibacterial class of compounds, the oxazolidinones. During the 1990s, many companies such as Upjohn, Bayer, Zeneca, Roussel Uclaf, Marion Merrell Dow and Glaxo published their work on antibacterial active oxazolidinones. The primary work in this area started in the 1980s at DuPont. The efforts of these scientists led to N-phenyl and N-heteroaryl oxazolidinones with strong antibacterial activity *in vitro* and *in vivo*. The most advanced oxazolidinone, linezolid, discovered by scientists at Upjohn, is currently undergoing Phase III clinical trials.

Keywords: *antibiotic, DuP-721, enterococci, eperezolid, Gram-positive, linezolid, methicillin-resistant Staphylococcus aureus, methicillin-resistant Streptococcus epidermidis, monoamine oxidase inhibitors, mycobacteria, N-phenyl oxazolidinone, N-pyridyl oxazolidinone, N-thienyl oxazolidinone, oxazolidinone, penicillin-resistant Streptococcus pneumoniae, staphylococci, vancomycin-resistant enterococci*


Exp. Opin. Ther. Patents (1999) 9(5):625-633

1. Introduction

The evolution and spread of resistance to currently available antimicrobial agents in important pathogens is a serious medical problem [1-3]. In particular, infections caused by multiresistant Gram-positive cocci are a threat in antibacterial therapy. Therefore, the primary focus of the worldwide antibiotic discovery programs is the identification of new compound classes with novel mechanisms of action.

The oxazolidinones represent a relatively new class of orally active, synthetic antimicrobial agents with a Gram-positive spectrum of activity. They inhibit an early event in the bacterial protein biosynthesis [4-6]. After sequencing the ribosomal RNA genes of oxazolidinone-resistant strains and comparing it to sensitive parental strains, mutations were found in the peptidyl transferase domain of the 23S rRNA [7]. This supports earlier results that demonstrated oxazolidinone binding to the 50S subunit of the ribosome [8-10]. A recent publication [11] describes the target for oxazolidinones in more detail. Drug binding sites were found on both ribosomal subunits. The 16S and the 23S rRNAs are directly involved in oxazolidinone binding. Furthermore, the inhibition of tRNA translocation is discussed as mechanism of action.

Oxazolidinones with antibacterial activity such as 5-(S)-acetamidomethyl-3-aryl-2-oxazolidinones were first described in the 1980s by scientists from DuPont [12-14]. In 1987 at the 27th ICAAC, DuPont presented two candidate compounds DuP-721 [15-17] and DuP-105 [15,16],



demonstrating high *in vitro* and *in vivo* activity *vs.* Gram-positive pathogens and *Mycobacterium tuberculosis* [18].

The increasing clinical problems with multiresistant pathogens along with the attractive properties of the oxazolidinones encouraged workers at Upjohn to start a discovery program [19]. These efforts led to two clinical candidates, eperezolid (**2**, U-100592) and linezolid (**1**, U-100766) [20,21]. Both compounds exhibited potent *in vitro* activity against Gram-positive bacteria, including multiresistant strains [22,23]. In various experimental infections, these oxazolidinones demonstrated excellent therapeutic efficacy [24]. Toxicological studies of the candidates in rats and dogs showed a favourable safety profile. In Phase I studies using either p.o. or iv. dosing, eperezolid (**2**) and linezolid (**1**) achieved plasma concentrations above the MIC values of relevant pathogens [25]. Both oxazolidinones are well-tolerated in humans at clinically relevant doses [26]. For a given dose, linezolid (**1**) produced higher plasma levels than eperezolid (**2**). Due to its advantageous pharmacokinetic profile, Upjohn continued the clinical development program with linezolid (**1**). During Phase II clinical trials, linezolid (**1**) was studied

in a dose range of 750 - 1000 mg/day for up to 21 days. The oxazolidinone showed good efficacy and excellent tolerability. Phase III clinical trials are underway.

Bayer has also taken interest in this compound class and new N-heteroaryl oxazolidinones with improved antimicrobial activity have been prepared. Modifications of the oxazolidinone core unit have been reported by Zeneca, Marion Merrell Dow [27], Roussel Uclaf [28], Glaxo [29] and Pharmacia & Upjohn.

In addition to the antimicrobial activity, other pharmacological activities of the oxazolidinones have been reported.

A comprehensive review from Brickner [19] discussed the structure-activity relationship (SAR) findings of DuPont and Upjohn and described in detail the oxazolidinones that entered clinical trials. Our review will focus on the recent activity selected from the patent literature since 1996.

2. N-Phenyl oxazolidinones

The N-phenyl oxazolidinones are the most advanced series of antibacterial oxazolidinones, first described by DuPont in the 1980s.

DuPont presented compounds bearing a (*S*)-5-acetamidomethyl side-chain, DuP-721 and DuP-105. Both oxazolidinones demonstrated good *in vitro* and *in vivo* activity against Gram-positive pathogens. These parentally and orally active candidates entered Phase I clinical trials, but the development was later discontinued for tolerability reasons.

Scientists at Upjohn focused on fluorine substituted N-phenyl oxazolidinones [30].

The most promising representative of the 3-fluoro-N-phenyl-oxazolidinone series is linezolid ((*S*)-N-[[3-[3-fluoro-4-(4-morpholinyl)phenyl]-2-oxo-5-oxazolidin-yl]methyl]-acetamide) (**1**) [116], which was identified in the early 1990s. It possesses good *in vitro* and *in vivo* potency *vs.* Gram-positive bacteria such as staphylococci [31], pneumococci [32] and enterococci [33,34], including resistant strains (methicillin-resistant *Staphylococcus aureus* [MRSA], methicillin-resistant *Streptococcus epidermidis* [MRSE], penicillin-resistant *Streptococcus pneumoniae* [PRSP], vancomycin-resistant enterococci [VRE]). Linezolid (**1**) is currently in clinical development

(Phase III) for severe hospital infections. The 3-fluoro-N-phenyl subclass seems to have both good *in vivo* potency and low toxicity.

Antibacterial therapy of intensive care unit (ICU) patients requires iv.-treatment. Most of the antibacterial oxazolidinones are insoluble in water and are therefore causing difficulties in finding a suitable iv. formulation. Upjohn described the water solubility and antibacterial activity of N-oxides [117] including the N-oxide of linezolid (1). This N-oxide retained most of linezolid's antibacterial activity (MIC = 4 µg/ml *vs.* *S. aureus* UC 9213) and had good water solubility (N-oxide 348 mg/ml, linezolid 3.7 mg/ml). This prodrug approach should help in finding suitable iv. formulations for oxazolidinones. This is not necessary for linezolid (1) since there is an appropriate iv. formulation already in use.






The main focus of the discovery program at Upjohn was the evaluation of the SAR around the 3-fluoro-N-phenyl oxazolidinones. The common feature of these 3-fluoro-N-phenyl oxazolidinones is a nitrogen containing heterocycle attached to the 4-position of the phenyl ring.

Small four- and five-membered heterocycles were described in 1996 [101], and exhibited reasonable *in vitro* potency (3, MIC *vs.* *S. aureus* or *Streptococcus pneumoniae* ~ 1 - 4 µg/ml).

N-Fluorophenyl oxazolidinones substituted with bicyclic nitrogen containing heterocycles were described [102,107] and exhibited good efficacy after oral dosing (ED₅₀ ~ 5 mg/kg) in murine infections caused by *S. aureus*.

Aromatic five-membered nitrogen containing heterocycles in the 4-position of the N-phenyl oxazolidinones increased the *in vitro* potency 2- to 4-fold compared to the corresponding non-aromatic analogues. These compounds were described by scientists from Upjohn [103] and Zeneca [104]. Formyl groups as substituents in the distal ring (4) increased the activity. The same effect was observed by Bayer scientists [105] in the N-heteroaryl oxazolidinone series. These formyl compounds cannot translate their excellent *in vitro* potency into *in vivo* efficacy (due to pharmacokinetic and metabolic problems).

The most promising compounds in the N-(3-fluorophenyl)oxazolidinone series, substituted with a five-membered aromatic heterocycle, are the thiazole substituted compounds [35]. (*S*)-N-(3-(3-Fluoro-4-(2-(5-cyanothiazolyl))phenyl)-2-oxo-5-



oxazolidinyl)methyl)acetamide (5) exhibited very strong antibacterial *in vitro* activity against Gram-positive cocci (MIC *vs.* *S. aureus* < 0.125 µg/ml) as well as against fastidious Gram-negative organisms (MIC *vs.* *Haemophilus influenzae* 30063 ~ 2 µg/ml).

The left side of the molecule allows for many variations. Besides the above mentioned small four- and five-membered heterocycles, large tricyclic systems connected to the oxazolidinone still showed reasonable *in vitro* potency [106] (6, MIC *vs.* *S. aureus* ~ 2 - 4 µg/ml).

Another field of intense research within the area of N-phenyl-oxazolidinones are the piperazine derivatives exemplified by (*S*)-N-(3-(3-fluoro-4-(4-(4-pyrimidinyl)-1-piperazinyl)phenyl)-2-oxo-5-oxazolidinyl)methyl)acetamide (7). Scientists from Upjohn [108] and Zeneca [109,110] reported compounds with good MIC values and reasonable *in vivo* potency (7, ED₅₀ = 4.4 mg/kg, *S. aureus* infection in mice). The heterocycles in 4-position of the piperazine unit are mainly pyrimidine, pyridine and other nitrogen containing five- and six-membered heterocycles [36]. Even large substituents like 2-(6-nitrobenzothiazole) on the piperazine unit [111] exhibited strong potency *vs.* Gram-positive bacteria (8, MIC *vs.*

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.