

# Chemistry

The Central Science

Eighth Revised Edition

Theodore L. Brown

University of Illinois at Urbana-Champaign

H. Eugene LeMay, Jr.

University of Nevada, Reno

Bruce E. Bursten

The Ohio State University

With contributions by Julia R. Burdge, University of Akron

PRENTICE HALL

Upper Saddle River, New Jersey 07458

Editor: John Challice  
Development Editor/Editor in Chief, Development: Carol Trueheart  
Associate Editor: Mary Hornby  
Editorial Assistants: Amanda K. Griffith, Gillian Buonanno  
Media Editor: Paul Draper  
Editorial/Production Supervision: Bob Walters  
Art Director: Joseph Sengotta  
Assistant Art Director: John Christiana  
Page layout: Richard Foster, Karen Noferi, Karen Stephens, Amy Peltier, Jeff Henn,  
Joanne Del Ben, Donna Marie Paukovits  
Art Studios: Academy Artworks/Michael Goodman/BioGrafx/Wellington  
Editor in Chief: Paul F. Corey  
Director of Marketing: John Tweeddale  
Assistant Vice President ESM Production and Manufacturing: David W. Riccardi  
Executive Managing Editor: Kathleen Schiaparelli  
Art Manager: Gus Vibal  
Art Editor: Karen Branson  
Assistant Art Editor: Adam Velthaus  
Senior Marketing Manager: Steve Sartori  
Marketing Assistant: Dorothy Marrero  
Director, Creative Services: Paul Belfanti  
Associate Creative Director: Amy Rosen  
Interior Design: Judith A. Matz-Coniglio  
Manufacturing Manager: Trudy Piscioti  
Photo Editor: Melinda Reo  
Photo Researcher: Yvonne Gerin  
Cover Illustration: © Kenneth Eward/BioGrafx, 1999  
Copy Editor: Fay Ahuja

© 2002, 2000, 1997, 1994, 1991, 1988, 1985, 1981, 1977 by Prentice-Hall, Inc.  
Upper Saddle River, NJ 07458

All rights reserved. No part of this book may be  
reproduced, in any form or by any means,  
without permission in writing from the publisher.  
Printed in the United States of America  
10 9 8 7 6 5 4 3 2 1

ISBN 0-13-061142-5

Prentice-Hall International (UK) Limited, *London*  
Prentice-Hall of Australia Pty. Limited, *Sydney*  
Prentice-Hall Canada Inc., *Toronto*  
Prentice-Hall Hispanoamericana, S.A., *Mexico*  
Prentice-Hall of India Private Limited, *New Delhi*  
Prentice-Hall of Japan, Inc., *Tokyo*  
Prentice-Hall (Singapore) Pte. Ltd., *Singapore*  
Editora Prentice-Hall do Brasil, Ltda., *Rio de Janeiro*

If we assume that  $x$  is small relative to 0.10 or 0.20  $M$ , this expression simplifies to give

$$\frac{(0.10)(x)}{0.20} = 6.8 \times 10^{-4}$$

$$x = \frac{0.20}{0.10} (6.8 \times 10^{-4}) = 1.4 \times 10^{-3} M = [F^-]$$

This  $F^-$  concentration is substantially smaller than it would be in a 0.20  $M$  solution of HF with no added HCl. The common ion,  $H^+$ , has repressed the dissociation of HF. The concentration of  $H^+(aq)$  is



$$[H^+] = (0.10 + x) M \approx 0.10 M$$

Thus,  $pH = 1.00$ . Notice that for all practical purposes,  $[H^+]$  is due entirely to the HCl; the HF makes a negligible contribution by comparison.

#### PRACTICE EXERCISE

Calculate the formate-ion concentration and pH of a solution that is 0.050  $M$  in formic acid,  $HCHO_2$  ( $K_a = 1.8 \times 10^{-4}$ ), and 0.10  $M$  in  $HNO_3$ . *Answer:*  $[CHO_2^-] = 9.0 \times 10^{-5}$ ;  $pH = 1.00$

Sample Exercises 17.1 and 17.2 both involve weak acids. The ionization of a weak base is also decreased by the addition of a common ion. For example, the addition of  $NH_4^+$  (as from the strong electrolyte  $NH_4Cl$ ) causes the base-dissociation equilibrium of  $NH_3$  to shift to the left, decreasing the equilibrium concentration of  $OH^-$  and lowering the pH:



## 17.2 Buffered Solutions

Solutions like those discussed in Section 17.1, which contain a weak conjugate acid–base pair, resist drastic changes in pH. Solutions that resist a change in pH upon addition of small amounts of acid or base are called **buffered solutions** (or merely **buffers**). Human blood is an important example of a complex aqueous medium with a pH buffered at about 7.4 (see the “Chemistry and Life” box near the end of this section). Much of the chemical behavior of seawater is determined by its pH, buffered at about 8.1 to 8.3 near the surface. Buffered solutions find many important applications in the laboratory and in medicine (Figure 17.1 ◀).

▼ **Figure 17.1** Prepackaged buffer solutions and ingredients for forming buffer solutions of predetermined pH.




### Composition and Action of Buffered Solutions

Buffers resist changes in pH because they contain both an acidic species to neutralize  $OH^-$  ions and a basic one to neutralize  $H^+$  ions. It is necessary that the acidic and basic species of the buffer do not consume each other through a neutralization reaction. These requirements are fulfilled by a weak acid–base conjugate pair such as  $HC_2H_3O_2-C_2H_3O_2^-$  or  $NH_4^+-NH_3$ . Thus, buffers are often prepared by mixing a weak acid or a weak base with a salt of that acid or base.

For example, the  $\text{HC}_2\text{H}_3\text{O}_2\text{-C}_2\text{H}_3\text{O}_2^-$  buffer can be prepared by adding  $\text{NaC}_2\text{H}_3\text{O}_2$  to a solution of  $\text{HC}_2\text{H}_3\text{O}_2$ ; the  $\text{NH}_4^+\text{-NH}_3$  buffer can be prepared by adding  $\text{NH}_4\text{Cl}$  to a solution of  $\text{NH}_3$ . By choosing appropriate components and adjusting their relative concentrations, we can buffer a solution at virtually any pH.

To understand better how a buffer works, let's consider a buffer composed of a weak acid (HX) and one of its salts (MX, where  $\text{M}^+$  could be  $\text{Na}^+$ ,  $\text{K}^+$ , or other cations). The acid-dissociation equilibrium in this buffered solution involves both the acid and its conjugate base:



The corresponding acid-dissociation-constant expression is


$$K_a = \frac{[\text{H}^+][\text{X}^-]}{[\text{HX}]} \quad [17.4]$$

Solving this expression for  $[\text{H}^+]$ , we have

$$[\text{H}^+] = K_a \frac{[\text{HX}]}{[\text{X}^-]} \quad [17.5]$$


We see from this expression that  $[\text{H}^+]$ , and thus the pH, is determined by two factors: the value of  $K_a$  for the weak-acid component of the buffer, and the ratio of the concentrations of the conjugate acid-base pair,  $[\text{HX}]/[\text{X}^-]$ .

If  $\text{OH}^-$  ions are added to the buffered solution, they react with the acid component of the buffer:





This reaction causes  $[\text{HX}]$  to decrease and  $[\text{X}^-]$  to increase. However, as long as the amounts of HX and  $\text{X}^-$  in the buffer are large compared to the amount of  $\text{OH}^-$  added, the ratio  $[\text{HX}]/[\text{X}^-]$  doesn't change much, and thus the change in pH is small (Figure 17.2 ▼).

If  $\text{H}^+$  ions are added, they react with the base component of the buffer:



This reaction can also be represented using  $\text{H}_3\text{O}^+$ :



◀ **Figure 17.2** A buffer consists of a mixture of a weak conjugate acid-base pair, here represented as HX and  $\text{X}^-$ . When a small portion of  $\text{OH}^-$  is added to the buffer (left), it reacts with HX, decreasing  $[\text{HX}]$  and increasing  $[\text{X}^-]$  in the buffer. When a small portion of  $\text{H}^+$  is added to the buffer (right), it reacts with  $\text{X}^-$ , decreasing  $[\text{X}^-]$  and increasing  $[\text{HX}]$  in the buffer.

Using either equation, we see that the reaction causes  $[X^-]$  to decrease and  $[HX]$  to increase. As long as the change in the ratio  $[HX]/[X^-]$  is small, the change in pH will be small (Figure 17.2).

Buffers most effectively resist a change in pH in *either* direction when the concentrations of HX and  $X^-$  are about the same. From Equation 17.5 we see that when  $[HX]$  equals  $[X^-]$ ,  $[H^+]$  equals  $K_a$ . For this reason, we usually try to select a buffer whose acid form has a  $pK_a$  close to the desired pH.

### Buffer Capacity and pH

Two important characteristics of a buffer are its capacity and its pH. **Buffer capacity** is the amount of acid or base the buffer can neutralize before the pH begins to change to an appreciable degree. This capacity depends on the amount of acid and base from which the buffer is made. The pH of the buffer depends on the  $K_a$  for the acid and on the relative concentrations of the acid and base that comprise the buffer. For example, we can see from Equation 17.5 that  $[H^+]$  for a 1-L solution that is 1 M in  $HC_2H_3O_2$  and 1 M in  $NaC_2H_3O_2$  will be the same as for a 1-L solution that is 0.1 M in  $HC_2H_3O_2$  and 0.1 M in  $NaC_2H_3O_2$ . However, the first solution has a greater buffering capacity because it contains more  $HC_2H_3O_2$  and  $C_2H_3O_2^-$ . The greater the amounts of the conjugate acid–base pair, the more resistant the ratio of their concentrations, and hence the pH, is to change.

Because conjugate acid–base pairs share a common ion, we can use the same procedures to calculate the pH of a buffer that we used in treating the common ion effect (see Sample Exercise 17.1). However, an alternate approach is sometimes taken that is based on an equation derived from Equation 17.5. Taking the negative log of both sides of Equation 17.5, we have

$$-\log [H^+] = -\log \left( K_a \frac{[HX]}{[X^-]} \right) = -\log K_a - \log \frac{[HX]}{[X^-]}$$

Because  $-\log [H^+] = \text{pH}$  and  $-\log K_a = \text{p}K_a$ , we have

$$\text{pH} = \text{p}K_a - \log \frac{[HX]}{[X^-]} = \text{p}K_a + \log \frac{[X^-]}{[HX]} \quad [17.8]$$

In general,

$$\text{pH} = \text{p}K_a + \log \frac{[\text{base}]}{[\text{acid}]} \quad [17.9]$$

where [acid] and [base] refer to the equilibrium concentrations of the conjugate acid–base pair. Note that when  $[\text{base}] = [\text{acid}]$ ,  $\text{pH} = \text{p}K_a$ .

Equation 17.9 is known as the **Henderson–Hasselbalch equation**. Biologists, biochemists, and others who work frequently with buffers often use this equation to calculate the pH of buffers. In doing equilibrium calculations, we have seen that we can normally neglect the amounts of the acid and base of the buffer that ionize. Therefore, we can usually use the starting concentrations of the acid and base components of the buffer directly in Equation 17.9.



Calculating pH  
Using the  
Henderson–Hasselbalch  
Equation activity

#### SAMPLE EXERCISE 17.3

What is the pH of a buffer that is 0.12 M in lactic acid,  $HC_3H_5O_3$ , and 0.10 M in sodium lactate? For lactic acid,  $K_a = 1.4 \times 10^{-4}$ .

**Solution** We will first determine the pH using the method described in Section 17.1. The major species in solution are  $HC_3H_5O_3$ ,  $Na^+$ ,  $C_3H_5O_3^-$ , and  $H_2O$ . The  $Na^+$  ion is

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.