

Research and Clinical Forums

VOLUME 20 NUMBER 1 1998
ISSN 0143-3083

Univ. of Minn.
Bio-Medical
Library

03 10 98

IBD AND SALICYLATES – 3



General Information

Research and Clinical Forums is an international journal devoted to publishing, in the English language, material presented and discussed at medico-scientific and medico-therapeutic congresses and symposia, either in the form of selected and collated papers, or as full reports and proceedings. Each issue is devoted to one subject area and may consist of the proceedings of, or selected papers from, one particular meeting or may be a compilation of material which, while relating to a particular subject, may have been presented at a number of meetings.

Material presented at congresses, symposia or meetings held anywhere in the world is considered for publication, the basic criterion for acceptance being the scientific and medical standard of the material. Each issue of *Research and Clinical Forums* is edited by an independent authority, or panel of assessors, expert in the relevant field. Views and factual claims expressed in individual contributions are personal to the respective contributor and are not necessarily endorsed by the Editors, Advisers, Assessors, Publishers, Distributors or Sponsors.

Research and Clinical Forums is published periodically to demand, is distributed to medical schools, medical libraries, hospitals and research establishments world-wide, and to all information and retrieval systems.

Where necessary, issues of *Research and Clinical Forums* will be translated into languages other than English and published as foreign language editions of those issues.

Copyright © 1998 by
Wells Medical Holdings Ltd
Speldhurst Place, Speldhurst Road
Royal Tunbridge Wells
Kent TN4 0JB
England

All rights reserved. This publication is copyright under the Berne Convention and the International Copyright Convention. Apart from any fair dealing under the UK Copyright Act 1956, Part 1, Section 7, no part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior permission of the Publisher.

Although great care has been taken in compiling and checking the information given in this publication to ensure that it is accurate, the Publisher shall not be held responsible for the continued currency of the information or for any errors, omissions or inaccuracies in this publication. Due to the rapid advances in medical science, it is recommended that independent verification of diagnoses and drug dosages should be made. In addition, no responsibility is assumed by the Publisher for any injury and/or damage to persons or property arising from any methods, products, instructions, ideas or otherwise contained in this publication.

Notes for Contributors see inside back cover.

The Publishing Editor
Research and Clinical Forums
Speldhurst Place, Speldhurst Road
Royal Tunbridge Wells
Kent TN4 0JB

British Library Cataloguing in Publications Data
Research and Clinical Forums
Volume 20, No. 1
1. Inflammatory Bowel Disease 2. Crohn's Disease 3. Ulcerative Colitis 4. Salicylates 5. Sulphasalazine
610'.5 R31
ISSN 0143-3083

Research and Clinical Forums is indexed in *Directory of Published Proceedings*, *Excerpta Medica/EMBASE*, *Current Contents Proceedings - Medical, Biological, Agricultural* and *Index to Scientific and Technical Proceedings*®.

New data on inflammatory bowel disease treatment with topical steroids

R. Löfberg, MD, PhD

Department of Gastroenterology
Huddinge University Hospital
Karolinska Institute
Stockholm
Sweden

SUMMARY

Glucocorticosteroids (GCSs) remain the mainstay primary treatment of active inflammatory bowel disease (IBD). These agents usually result in a rapid suppression of symptoms but due to the ubiquitous nature of the GCS receptor, they also cause systemic side-effects. Recently, new GCSs developed mainly for asthma (eg., budesonide) have been found to counter the problem of systemic side-effects due to a greater first-pass metabolism and an increased affinity for the GCS receptor, resulting in a predominantly topical mode of action. Budesonide seems particularly suitable for IBD conditions. It has been developed for both rectal and oral delivery to the gut and has been shown to have good efficacy in both Crohn's disease and ulcerative colitis with a mild side-effect profile.

INTRODUCTION

Effects and side-effects of glucocorticosteroids

Glucocorticosteroids (GCSs) remain the mainstay of medical treatment in active inflammatory bowel disease (IBD). The anti-inflammatory effects of GCSs in acute attacks of ulcerative colitis (UC) as well as in Crohn's disease (CD) are unsurpassed by any other type of drug and a rapid and substantial effect on clinical symptoms is usually seen during the first few days of treatment. Glucocorticosteroids exert their actions by binding to the intracellular GCS-receptor which is uniformly distributed in all types of human cells. Activation of the receptor will evoke a number of responses and effects, including inhibition of the production and action of certain key pro-inflammatory cytokines.

The anti-inflammatory effects of conventional GCSs in IBD are commonly offset by systemic side-effects: eg., cosmetic or psychological problems such as acne, moonface, striae and mood disturbances including insomnia. Others, like hypertension, dyspepsia and impaired glucose tolerance, may be of substantial clinical importance. Long-term treatment is precluded by the risk for hazardous and sometimes also irreversible complications such as osteoporosis and osteonecrosis, cataracts and diabetes mellitus. The suppression of endogenous cortisol levels and impairment of the hypothalamic-pituitary-adrenal gland function, with delayed recovery, are other issues of concern.

Development of GCSs for topical action

Recent steroid development, primarily aimed at refinement of topical inhalation therapy in asthma, has provided us with new GCS compounds such as budesonide and fluticasone, which have 100–200 times higher affinity for the GCS receptor compared with hydrocortisone, enabling treatment with much reduced doses. If the drug is administered in the gut, the major portion will be cleared through the portal vein to the liver where it is, to a large extent, inactivated during the hepatic first-pass metabolism. The systemic availability of the new GCSs will therefore be substantially reduced compared with that of older GCSs, and this improved ratio of efficacy at the target level *vs* decreased systemic load is the principle on which new topical steroids for IBD rely^{1,2}. Furthermore, other GCS prerequisites for optimal topical action in IBD conditions include sufficient water solubility for adequate distribution in the bowel lumen, a high mucosal uptake and deep penetration. These enhanced properties of new GCSs are dependent on the introduction of lipophilic substitutions in the 16- α and 17- α positions on the GCS skeleton.

Several new GCSs have been evaluated for use in IBD during the last years², but after review³, only a few compounds remain on the market for topical IBD applications.

BECLOMETHASONE

Beclomethasone, 17- α and 21-dipropionate, given in enema form for active distal UC, has previously been shown to have equal efficacy but reduced suppression of plasma cortisol levels in comparison with conventional GCS enemas⁴. Beclomethasone was recently compared with 5-aminosalicylic acid (5-ASA) in a randomised controlled trial (RCT); the two drugs were equally effective when given as monotherapy for active ulcerative proctitis, but significantly improved results were obtained when a combination of the two different compounds (3 mg GCS + 2 g 5-ASA in 100 ml) was used⁵. The results need confirmation, but are still of great clinical interest as they indicate that therapeutic synergism may be obtained by combining drugs utilising different modes of action, maintaining a low toxicity profile⁶. However, as beclomethasone is now out of patent, the commercial interest for a study using this particular drug is probably low and it is uncertain if further development for UC will be continued.

BUDESONIDE

To date, the best documented of the new generation of highly potent 17- α substituted GCSs for the treatment in IBD, is budesonide. It is now approved in several countries, both as an enema and in an oral preparation (Entocort[®], Astra, Sweden). Due to its relatively high water solubility, budesonide is readily dissolved, facilitating transport to the bowel wall. These hydrophilic properties ensure a high tissue uptake, resulting in high concentrations and high activity in the target tissues when the drug is applied topically. As shown in animal studies, budesonide has a greater topical uptake and remains within the mucosa for a longer period of time than prednisolone¹. The drug is biotransformed rapidly in the liver *via* the cytochrome P450 CYP3A4 enzymes, with 80–90% metabolised in the first pass resulting in low systemic availability.

BUDESONIDE EFFICACY IN THE TREATMENT OF ACTIVE DISTAL UC AND PROCTITIS

positive results⁷⁻¹¹. In an American RCT comprising 184 patients with distal UC, the budesonide enema was compared with standard hydrocortisone enema. Budesonide showed similar therapeutic efficacy, but had significantly less impact on the response to an adrenocorticotrophic hormone (ACTH) test after 6 weeks of treatment¹². Another large US study with 223 patients evaluated the budesonide enema in a dose-ranging RCT *vs* placebo, comparing doses of 0.5, 2 and 8 mg given at bedtime. The optimal dose was confirmed to be 2 mg/100 ml, although the 8 mg dose provided slightly higher remission rates¹³. In line with these results, a recent, Swedish, multicentre RCT investigating active distal UC and proctitis treatment using administration of the enema *bd.* could not demonstrate a significant advantage in remission rate over the standard, once-daily dose (62% *vs* 57% *bd.*) after 8 weeks¹⁴. Furthermore, in those patients achieving remission, a prolonged treatment with the budesonide enema given twice weekly for up to 6 months was not significantly superior to placebo in maintaining remission (38% *vs* 53% relapse rate)¹⁴. A clinical trial comparing budesonide enema 2 mg/100 ml to Pentasa[®] (mesalazine) 1 g/100 ml in acute distal UC or proctitis for 4 weeks¹⁵ showed equal efficacy in inducing endoscopic and histological remission; however, the Pentasa[®] enema was superior in obtaining clinical remission (60% *vs* 38%; $P = 0.03$).

BUDESONIDE FOR ORAL USE

A refined, slow-release delivery system developed for budesonide by Astra Draco AB, Sweden uses mm-sized enteric coated (Eudragit L) pellets with a rate-limiting polymer containing the active drug¹⁶. This time-dependent and pH-dependent delivery system (controlled ileal release, [CIR]) results in an absorption of around 70% of the given drug in the distal ileum and the caecum, making it appropriate for treatment of active distal ileal and/or right-sided colonic CD. Encouraging results in a small pilot trial were confirmed by two large multicentre RCTs. The optimal dose of the oral preparation for active ileal or ileocaecal CD was found to be 9 mg in a Canadian placebo-controlled, dose-ranging study of 258 patients¹⁷. Special attention was made to monitor GCS-associated side-effects, but interestingly, no significant differences were found between the active treatment groups compared with placebo in this respect. The 9 mg dose had the best clinical efficacy and a GCS-associated side-effect incidence similar to that of placebo (26%); however, significantly lower plasma cortisol was seen after 8 weeks treatment in the budesonide group compared to placebo (69% *vs* 14%; $P < 0.001$). In a European, multicentre, double-blind, double-dummy RCT, comparing 9 mg budesonide CIR given with a standard regimen of 40 mg/day of oral prednisolone tapering over a 10-week period¹⁸, no statistical difference in overall remission rate (as determined by a Crohn's disease activity index [CDAI]-score < 150) was seen (52% *vs* 65%; $P = 0.12$). While clinical efficacy was comparable, morning plasma cortisol levels were markedly suppressed in the prednisolone group after 8 weeks ($P = 0.02$). In line with this, GCS-associated side-effects were milder and significantly less frequent in the budesonide group (33% *vs* 55%; $P = 0.003$) during the study period.

These results have recently been confirmed and reinforced by an international comparative RCT¹⁹, involving 177 patients with active ileocaecal CD. The study evaluated two different dosing regimens for budesonide. A once daily *vs* a twice daily

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.