Handbook of PHARMACEUTICAL EXCIPIENTS

Published by

American Pharmaceutical Association 2215 Constitution Avenue, NW

The Pharmaceutical Society of Great Britain
1 Lambeth High Street



American Pharmaceutical Association Production Staff

James C. Boylan Jack Cooper Zak T. Chowhan

The Pharmaceutical Society of Great Britain Production Staff

Walter Lund Ainley Wade Robert F. Weir Bernard J. Yates

Publication Production Staff

Marlene Povich, Art and Production Manager Carol Niedziałek, Production Coordinator John Covert, Director of Publications

Library of Congress Data

The use of portions of the text of *USP XXI/NF XVI* is by permission of the USP Convention, Inc. The Convention is not responsible for any inaccuracy of quotation or for any false or misleading implication that may arise from separation of an excerpt from the original context or by obsolescence resulting from publication of a supplement.

Selected structural formulas from *The Merck Index*, Tenth Edition, 1983, published by Merck & Co., Rahway, New Jersey, are reprinted with permission.

© Copyright 1986 by the American Pharmaceutical Association, 2215 Constitution Avenue, N.W., Washington, D.C. 20037 USA, and The Pharmaceutical Society of Great Britain, 1 Lambeth High Street, London SE1 7JN, England. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage or retrieval system, without prior written permission from the joint publishers.

ISBN (U.S.): 0-917330-56-0 ISBN (U.K.): 0-85369-164-9



Lecithin

1. Nonproprietary Name

NF: Lecithin

2. Functional Categories

USP: Emulsifying and/or solubilizing agent.

Others: wetting and dispersing agent; emollient; digestible surfactant; moisturizer

3. Synonyms

Soybean phospholipids; mixed soybean phosphatides; soybean lecithin; vegetable lecithin; egg lecithin; ovolecithin.

4. Chemical Name and CAS Registry Number

Phosphatidylcholine—commercially, a mixture of phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phytoglycolipids, other phosphatides, carbohydrates, triglycerides and other components. [8002-43-5]

5. Empirical Formula

Molecular Weight

6. Structural Formula

$$\begin{array}{c} \operatorname{CH_2OCOR} \\ \vdash \\ \operatorname{CHOCOR} \\ \operatorname{CH_2O} - P - \operatorname{OCH_2CH_2N} (\operatorname{CH_3})_3 \end{array}$$

 α —phosphatidylcholine

 β —phosphatidylcholine

7. Commercial Availability

USA

American Lecithin Central Soya Co. Spencer-Kellogg

UK

Alembic (Lovelock) Products Steetley Chemicals, Ltd.

8. Method of Manufacture

Crude soybean lipids are obtained by the solvent extraction of soybeans. Lecithin is obtained as a water extract of these crude soybean lipids. This crude lecithin is then upgraded by various techniques. Lecithin can also be made from corn, peanut, cottonseed, castor seed and coffee beans. It has been made from egg, known as egg lecithin. Synthetic lecithins are also made.

Description

Lecithin is brown to light yellow, depending on whether it is unbleached or bleached. It has practically no odor and a bland to nut-like taste, similar to soybean oil. In consistency, it may vary from plastic to fluid depending on the free fatty acid content.

10. Pharmacopeial Specifications

Test	NF_
Water (method I)	≤1.5%

Test	NF
Hexane-insoluble matter	≤0.3%
Acetane-insoluble matter	≤50.0%

11. Typical Properties

Isoelectric point: Approximately 3.5

Miscibility: When mixed with water, lecithin hydrates to form an emulsion. It readily forms imbibitions (taking up liquid without a measurable increase in volume) with almost any solvent.

pH: Approx. 6.6 (soybean phosphatides)

Solubility: Soluble in aliphatic and aromatic hydrocarbons, halogenated hydrocarbons, mineral oil and fatty acid. Insoluble in polar solvents and water. Practically insoluble in cold vegetable and animal oils.

Physical form: Liquid or granule (oil-free lecithin)

Density: Liquid lecithin, 0.97 g/cm³; granule lecithin, 0.5 g/cm³

Iodine value: Liquid lecithin, 95-100; granule lecithin, 82-88 Saponification value: 196

12. Stability and Storage Conditions

Fluid lecithin grades should be stored at room temperature or above (22°-49°C) in well-closed containers. Temperatures of 160-180°C will cause degradation within 24 hours. Temperatures below 10°C may cause separation. Lecithins are hygroscopic; keep containers closed to prevent moisture pickup. Lecithin is subject to microbial degradation. Exposure to light should be avoided. A shelf life of several years can be expected when stored at ambient temperatures.

13. Incompatibilities

Lecithin decomposes at extreme pH. When heated, it oxidizes, darkens and decomposes. It is hydrolyzed by esterases.

14. Safety

Lecithin is recognized as a multiple purpose food substance by the FDA.

15. Handling Precautions

16. Regulatory Status

NF XVI

17. Applications in Pharmaceutical Formulation or Technology

Lecithins are used as dispersing and emulsifying agents for intramuscular injections. They are used in creams and ointments as dispersing, wetting, emulsifying and emollient agents.

18. Related Substances

19. Comments

Since lecithin contains a variety of unspecified materials, care should be exercised in the use of unpurified lecithin in injectable or topical dosage forms, as interaction with the active ingredient or formulation excipients may occur.

20. Specific References

 P. Sartoretto, Lecithin. In "Encyclopedia of Chemical Technology," 2nd Ed., Kirk & Othmer eds., Vol. 12, John Wiley and Sons, Inc., 1967, New York, p. 343.

