

12-1-1992

## Densities of Vegetable Oils and Fatty Acids

Hossein Nouredini

*Department of Chemical Engineering, University of Nebraska-Lincoln, hnoured@unlnotes.unl.edu*

B C. Teoh

*Department of Chemical Engineering, University of Nebraska-Lincoln*

L Davis Clements

*Department of Chemical Engineering, University of Nebraska-Lincoln*

Follow this and additional works at: [http://digitalcommons.unl.edu/chemeng\\_biomaterials](http://digitalcommons.unl.edu/chemeng_biomaterials)

 Part of the [Biomaterials Commons](#)

Nouredini, Hossein; Teoh, B C.; and Clements, L Davis, "Densities of Vegetable Oils and Fatty Acids" (1992). *Papers in Biomaterials*. Paper 14.

[http://digitalcommons.unl.edu/chemeng\\_biomaterials/14](http://digitalcommons.unl.edu/chemeng_biomaterials/14)

This Article is brought to you for free and open access by the Chemical and Biomolecular Engineering Research and Publications at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Biomaterials by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Densities of Vegetable Oils and Fatty Acids

H. Nouredini\*, B.C. Teoh and L. Davis Clements

Department of Chemical Engineering, University of Nebraska-Lincoln, Lincoln, Nebraska 68588-0126

Complete data for density as a function of temperature have been measured for a number of vegetable oils (crambe, rapeseed, corn, soybean, milkweed, coconut, lesquerella), as well as eight fatty acids in the range  $C_9$  to  $C_{22}$  at temperatures from above their melting points to  $110^\circ\text{C}$  ( $230^\circ\text{F}$ ). The specific gravity and density measurements were performed according to American Society for Testing and Materials (ASTM) standard test methods D 368, D 891 and D 1298 for hydrometers and a modified ASTM D 369 and D 891 for pycnometers. Correlation constants, based on the experimental data, are presented for calculating the density of fatty acids and vegetable oils in the range of temperature from  $24^\circ\text{C}$  ( $75^\circ\text{F}$ ) or the melting point of the substance, to  $110^\circ\text{C}$  ( $230^\circ\text{F}$ ). The constants are valuable for designing or evaluating such chemical process equipment as heat exchangers, reactors, process piping and storage tanks. Estimated density of fatty acids by a modified Rackett equation is also presented.

**KEY WORDS:** Density, fatty acids, specific gravity, vegetable oils.

Density or specific gravity data are important in numerous chemical engineering unit operations in the fatty acid industry. Representative examples include reactors for splitting of fatty acids or conversion of fatty acids to their derivatives, distillation for separation of fatty acids, or for designing storage tanks and process piping. In addition, the pure component data for fatty acids may be used to ascertain mixture property values and to estimate oil density.

The data were measured by both hydrometer and pycnometer procedures in a single constant-temperature bath. Pycnometer measurements provide the true density of the material and the hydrometer measures specific gravity. Due to the tedious procedure involved in pycnometer measurements, they were used only for some of the data to validate the hydrometer readings over the entire range of temperature. This procedure resulted in a correction factor for hydrometer measurements. In all measurements the correction never exceeded 0.2%.


### EXPERIMENTAL PROCEDURES

A Precision Scientific (Chicago, IL) constant-temperature bath with a Micro-set Thermoregulator, in accordance with American Society for Testing and Materials (ASTM) D 445 (1), was used in the procedures for specific gravity and density determinations. This assembly maintains a temperature uniformity of  $\pm 0.0033^\circ\text{C}$  ( $\pm 0.005^\circ\text{F}$ ) throughout the range of 20 to  $130^\circ\text{C}$  (68 to  $266^\circ\text{F}$ ), exceeding the ASTM D 445 (1) requirements. Fisher's Precision Specific Gravity Hydrometers (Fisher Scientific, Pittsburgh, PA), ASTM D 1298 (2) with subdivisions of 0.001, and Fisher thermometers, ASTM D 445 (1), with subdivisions of  $0.056^\circ\text{C}$  ( $0.1^\circ\text{F}$ ), were used for specific gravity and temperature readings, respectively.

\*To whom correspondence should be addressed at Department of

A glass cylinder was installed through one of the holes in the bath cover assembly. The sample was placed inside the glass cylinder for hydrometer measurements. The bath temperature was set at the desired temperature and was monitored with a glass thermometer during the experiment. The temperature of the sample inside the glass cylinder was monitored with a second thermometer. At higher temperatures, where a temperature gradient may exist, the bath temperature was increased to maintain the desired temperature inside the cylinder. The specific gravities were determined according to the procedures in ASTM methods D 1298, D 368, and D 891 (2-4).

The pycnometer measurements were performed along with the hydrometer procedures in the same bath. Three 10-mL Fisher pycnometers, in accordance with ASTM D 369 (5), were used in all measurements. Pycnometers were filled with the sample at a temperature below the bath temperature. Heat resistant tubing was slipped over the outside of the pycnometer neck to prevent direct contact between the bath fluid and the capillary opening of the bottle. Figure 1 shows the pycnometer assembly. This assembly was secured to the bath cover and clip and



## DENSITIES OF VEGETABLE OILS AND FATTY ACIDS

TABLE 1

## Fatty Acid Distributions and Sources of Vegetable Oils

Vegetable oils	% Composition <sup>a</sup>			Source or manufacturer
	Saturated	Monounsaturated	Polyunsaturated	
Crambe <sup>b</sup>	5	79	16	Texas A&M University, College Station, TX
Rapeseed <sup>b</sup>	7	72	21	University of Idaho, Moscow, ID
Corn	4.3	21.4	64.3	Hain Pure Food Co. Inc., Los Angeles, CA
Soybean	14.3	28.0	57.1	Hain Pure Food Co. Inc., Los Angeles, CA
Milkweed	NA <sup>c</sup>	NA	NA	Biological System Eng., U. of Nebraska, Lincoln, NE
Coconut	94.0	5.0	1.0	Vinton Pop Corn Co., Vinton, IA
Lesquerella	3.1	76.0	20.9	Jobba Growers & Processors Inc., Apache Junction, AZ

<sup>a</sup>Provided by the manufacturers. <sup>b</sup>Approximate values. <sup>c</sup>Not available.

TABLE 2

## Purity of Fatty Acids

Acids	% Purity <sup>a</sup>	Manufacturer
Nonanoic	98	Kodak, Rochester, NY
Capric	99-100	Sigma, St. Louis, MO
Lauric	99-100	Sigma
Myristic	99	Kodak
Palmitic	99	Sigma
Stearic	95	Aldrich, Milwaukee, WI
Oleic	98	J.T. Baker, Phillipsburg, PA
Erucic	90	Sigma

<sup>a</sup>Provided by the manufacturers.

immersed inside the bath. Except for the modifications for the pycnometer assembly, the procedures in the ASTM standard test methods D 891 and D 369 (4,5) were followed for the density determinations.

The fatty acid composition and source or manufacturer of each vegetable oil are shown in Table 1. Table 2 shows the purity and the manufacturers of the fatty acids used. The information on the fatty acid composition of vegetable oils and purity of the fatty acids were provided by the manufacturers.

## RESULTS

The scale readings for the hydrometer were corrected for the glass expansion by using a glass expansion coefficient of  $25 \times 10^{-6}$  per °C. The specific gravity data for crambe and milkweed seed oils and for oleic acid were measured with both hydrometers and pycnometers. The hydrometer scale readings were corrected according to the pycnometer measurements. The correction was equal to 0.0025. This correction was uniform and did not depend on temperature or type of oil or acid used. This correction was then applied to the rest of the samples, for which only the hydrometer measurements were performed. Densities were calculated by multiplying the specific gravities by water density (0.99904 g/mL) at 15.5°C (60.0°F). This is the calibration temperature of the hydrometers. The results are tabulated in Tables 3 and 4. Available literature values (6,7) at 25°C (77°F) are presented for comparison.

A linear correlation was fitted to the experimental data.

ments are also included. As expected, the deviations found were small. The mean deviation was less than 0.07% for all data. Figures 2 and 3 show the experimental and linear predictions for vegetable oils (lesquerella, milkweed, crambe) and for saturated fatty acids (nonanoic, capric, lauric, myristic, palmitic, stearic), respectively. Lines show the linear correlations compared to the experimental points.

The difference between successive test results by the same apparatus under constant operating conditions on identical test material never exceeded 0.0005 for hydrometer scale readings and 0.003 for pycnometer measurements. These are within the sensitivity limits imposed by ASTM test methods D 1298 and D 369 (2,5), respectively. Random examination of results for other samples, for which hydrometer measurements were principally performed, confirmed the correction applied to the scale readings. Corrected data were also within the repeatability specifications given by the ASTM methods, thus justifying the application of the correction factor for the hydrometer measurements.

The modified Rackett equation (8) also can be used for calculation of fatty acid and vegetable oil densities. This approach can be used directly in most computer-aided-process-design programs such as PRO/II<sup>TM</sup>, DESIGN II<sup>TM</sup> or HySim<sup>TM</sup>. The modified Rackett equation requires the critical properties and an empirical parameter,  $Z_{RA}$ , for each acid as the basis for computing density as a function of temperature. The required critical properties and Rackett parameters are usually available in the component library or may be easily incorporated into the program data file. The modified Rackett equation is as follows:

$$\rho = \frac{MW}{\frac{RT_c}{P_c} Z_{RA}^{[1+(1-T_r)^{2/7}]}} \quad [1]$$

$T_c$  and  $P_c$  are the critical temperature and pressure, respectively;  $\rho$  is the density;  $T_r$  is the reduced temperature ( $T_r = T/T_c$ );  $R$  is the universal gas constant;  $MW$  is the molecular weight; and  $Z_{RA}$  is the Rackett parameter, a correlating parameter unique to each compound. Critical properties for the fatty acids mentioned in this work are shown in Table 7. A more complete listing of fatty acid critical properties was found in earlier works

TABLE 3

## Density of Vegetable Oils

Temperature °C (°F)	Density (g/mL)						
	Crambe <sup>a</sup>	Rapeseed	Corn	Soybean	Milkweed <sup>a</sup>	Coconut	Lesquerella
23.9 (75.0)	0.9078	0.9073	0.9188	0.9193	0.9203		0.9398
37.8 (100.0)	0.8977	0.8987	0.9082	0.9082	0.9097	0.9107	0.9307
48.9 (120.0)	0.8898	0.8908	0.9028	0.9023	0.9018	0.9033	0.9229
60.0 (140.0)	0.8829	0.8839	0.8939	0.8939	0.8934	0.8949	0.9152
82.2 (180.0)	0.8681	0.8686	0.8800	0.8795	0.8790	0.8795	0.8999
100.0 (212.0)	0.8564	0.8579	0.8679	0.8674	0.8684	0.8669	0.8879
110.0 (230.0)	0.8501	0.8511	0.8610	0.8615	0.8605	0.8605	0.8830
Literature data		0.903-0.907	0.912-0.917	0.914-0.918		0.916-0.918	
Temp. (°C) <sup>b</sup>		25	25	25		25	

<sup>a</sup>By direct pycnometer determination. Other values by corrected hydrometer determination. <sup>b</sup>Ref. 6.

TABLE 4

## Density of Fatty Acids

Temperature °C (°F)	Density (g/mL)							
	Nonanoic 9:0	Capric 10:0	Lauric 12:0	Myristic 14:0	Palmitic 16:0	Stearic 18:0	Oleic <sup>a</sup> 18:1	Erucic 22:1
23.9 (75.0)	0.9018						0.8928	
37.8 (100.0)	0.8913	0.8863					0.8828	0.8763
48.9 (120.0)	0.8838	0.8788	0.8729				0.8753	0.8689
60.0 (140.0)	0.8744	0.8694	0.8634	0.8590			0.8674	0.8611
71.1 (160.0)	0.8665	0.8615	0.8560	0.8516	0.8496		0.8605	0.8532
82.2 (180.0)	0.8571	0.8526	0.8471	0.8432	0.8412	0.8387	0.8521	0.8459
98.9 (210.0)	0.8445	0.8405	0.8351	0.8321	0.8302	0.8276	0.8400	0.8348
110.0 (230.0)	0.8371	0.8332	0.8282	0.8257	0.8237	0.8217	0.8327	0.8284
121.1 (250.0)						0.8148		
Literature data	0.8570	0.8531	0.8477	0.8439	0.8414	0.8390	0.850	0.8699
Temp. (°C) <sup>b</sup>	80	80	80	80	80	80	80	55

<sup>a</sup>By direct pycnometer determination. Other values by corrected hydrometer determination. <sup>b</sup>Ref. 6, except for erucic acid (ref. 7).




FIG. 2. Density vs. temperature for vegetable oils. The lines are correlation derivatives and the points are experimental data. (+), Les-




FIG. 3. Density vs. temperature for fatty acids. The lines are correlation derivatives and the points are experimental data. (•), Nonanoic acid; (+), capric acid; (•), lauric acid; (□), myristic acid; (x)

## DENSITIES OF VEGETABLE OILS AND FATTY ACIDS

TABLE 5

Linear Correlations<sup>a</sup> for the Vegetable Oil Densities (g/mL)

Vegetable oils	Constants		Data points n	% Mean deviation r	Temperature range (°C)
	m	c			
Crambe	-6.665E-04	9.2300E-01	7	0.032	23.9-110.0
Rapeseed	-6.550E-04	9.2305E-01	7	0.029	23.9-110.0
Corn	-6.650E-04	9.3432E-01	7	0.056	23.9-110.0
Soybean	-6.674E-04	9.3441E-01	7	0.057	23.9-110.0
Milkweed	-6.820E-04	9.3552E-01	7	0.069	23.9-110.0
Coconut	-7.000E-04	9.3716E-01	6	0.026	37.8-110.0
Lesquerella	-6.709E-04	9.5569E-01	7	0.049	23.9-110.0

<sup>a</sup>Density = c + mT.

TABLE 6

Linear Correlations<sup>a</sup> for the Fatty Acid Densities (g/mL)

Fatty acids	Carbon	Constants		Data points n	% Mean deviation r	Temperature range (°C)
		m	c			
Nonanoic	9:0	-7.595E-04	9.2015E-01	8	0.048	23.9-110.0
Capric	10:0	-7.453E-04	9.1452E-01	7	0.047	37.8-110.0
Lauric	12:0	-7.319E-04	9.0792E-01	6	0.067	48.9-110.0
Myristic	14:0	-6.727E-04	8.9909E-01	5	0.055	60.0-110.0
Palmitic	16:0	-6.649E-04	8.9634E-01	4	0.060	71.1-110.0
Stearic	18:0	-6.081E-04	8.8838E-01	4	0.036	82.2-121.1
Oleic	18:1	-6.982E-04	9.0946E-01	8	0.023	23.9-110.0
Erucic	22:1	-6.694E-04	9.0131E-01	7	0.042	37.8-110.0

<sup>a</sup>Density = c + mT.

TABLE 7

Fatty Acid Densities and Properties for Use with the Rackett Equation

	Fatty acid (density, g/mL)							
	Nonanoic 9:0	Capric 10:0	Lauric 12:0	Myristic 14:0	Palmitic 16:0	Stearic 18:0	Oleic 18:1	Erucic 22:1
MW	158.24	172.27	200.32	228.37	256.43	284.48	284.48	340.59
T <sub>c</sub> (K)	717.03	730.84	756.21	779.07	799.89	819.00	819.41	853.41
P <sub>c</sub> (bar)	25.18	22.92	19.22	16.35	14.08	12.25	12.76	9.87
Z <sub>RA</sub>	0.24654	0.24419	0.23862	0.23263	0.22641	0.22041	0.22344	0.21177
Temperature [°C (°F)]								
23.9 (75.0)	0.9015						0.8910	
37.8 (100.0)	0.8912	0.8863					0.8820	0.8750
48.9 (120.0)	0.8829	0.8782	0.8717				0.8748	0.8680
60.0 (140.0)	0.8745	0.8700	0.8639	0.8593			0.8675	0.8609
71.1 (160.0)	0.8660	0.8617	0.8560	0.8517	0.8494		0.8602	0.8538
82.2 (180.0)	0.8573	0.8534	0.8480	0.8440	0.8419	0.8388	0.8528	0.8467
98.9 (210.0)	0.8442	0.8407	0.8358	0.8323	0.8305	0.8276	0.8415	0.8358
110.0 (230.0)	0.8353	0.8320	0.8276	0.8244	0.8228	0.8201	0.8340	0.8285
121.1 (250.0)						0.8125		
% Mean deviation, r	0.060	0.059	0.076	0.064	0.065	0.120	0.102	0.083

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.