



US007141249B2

(12) **United States Patent**
Ohashi et al.

(10) **Patent No.:** US 7,141,249 B2
(45) **Date of Patent:** *Nov. 28, 2006

(54) **RAPIDLY SOLUBLE DRUG COMPOSITION**

(75) Inventors: **Mamoru Ohashi**, Amagasaki (JP);
Kazuyoshi Ogasawara,
Kitakatsuragi-gun (JP); **Yoshimi Shirai**,
Suita (JP); **Hiroshi Fujioka**, Ibaraki
(JP)

(73) Assignee: **Dainippon Sumitomo Pharma Co.,
Ltd.**, Osaka-fu (JP)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **09/529,715**

(22) PCT Filed: **Oct. 15, 1998**

(86) PCT No.: **PCT/JP98/04658**

§ 371 (c)(1),
(2), (4) Date: **Apr. 19, 2000**

(87) PCT Pub. No.: **WO99/20277**

PCT Pub. Date: **Apr. 29, 1999**

(65) **Prior Publication Data**

US 2002/0197308 A1 Dec. 26, 2002

(30) **Foreign Application Priority Data**

Oct. 20, 1997 (JP) 9-306635

(51) **Int. Cl.**
A61K 9/20 (2006.01)
A61K 9/14 (2006.01)
A61K 31/495 (2006.01)

(52) **U.S. Cl.** **424/465**; 424/464; 424/489;
514/951; 514/960; 514/249

(58) **Field of Classification Search** 514/249,
514/951, 960; 424/94.4, 400, 464, 465, 489,
424/451, 452, 470
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,910,022 A * 3/1990 Bavitz et al. 424/465
5,258,382 A 11/1993 Negoro et al. 514/249
5,356,636 A * 10/1994 Schneider et al. 424/489
5,858,410 A * 1/1999 Muller et al. 424/489
5,952,356 A * 9/1999 Ikeda et al. 514/340
6,297,244 B1 * 10/2001 Ohashi et al. 514/249
6,458,811 B1 * 10/2002 Arbuthnot et al. 514/324

FOREIGN PATENT DOCUMENTS

JP 5-186472 4/1996

OTHER PUBLICATIONS

English Language Translation of the Document: "Design and Evaluation
of Peroral Pharmaceutical Preparations (In Japanese)", edited
by Mitsuru Hashida, Yakugyo Jihosya, Feb. 10, 1995, pp. 81-84,
168-171 (cited in the International Search Report of the original
PCT/JP98/04658) thereof.


(Continued)

Primary Examiner—Sharmilla S. Gollamudi
(74) *Attorney, Agent, or Firm*—Wenderoth, Lind & Ponack,
L.L.P.

(57) **ABSTRACT**

A fast-dissolving pharmaceutical composition comprising
micronized (R)-2-(4-bromo-2-fluorobenzyl)-1,2,3,4-tetra-
hydropyrrolo[1,2-a]pyrazine-4-spiro-3'-pyrrolidine-1,2',3,
5'-tetrone (hereinafter, referred to as AS-3201). The present
pharmaceutical composition has improved dissolution char-
acteristics as well as a good bioavailability.

5 Claims, 1 Drawing Sheet



OTHER PUBLICATIONS

English Language translation of the Document "Practical Drug Additives (In Japanese)", cited by Research Team Concerning Drug Additives, Mar. 5, 1974, pp. 215-217, 258-259 (cited in the International Search Report of the original PCT/JP98/04658) thereof. Chemical Abstracts, 122, 9860 (1995) (English Abstract of JP-A-6-192222).
Chemical Abstracts, 125, 221569 (1996) (English Abstract of JP-A-8-176105).


J. Lutomski et al., "Postep W Technice Farmaceutyusznej", p. 102, 1981 (English translation also enclosed).

"Design and Evaluation of Peroral Pharmaceutical Preparations (in Japanese)", edited by Mitsuru Hashida, Yakyogyo Jihosha, Feb. 10, 1995, pp. 81-84, 168-171, Trans. Sep. 20, 2000.

"Practical Drug Additives (in Japanese)", Research Team Concerning Drug Additives, Mar. 5, 1974, pp. 258-259, Trans Sep. 26, 2000.

* cited by examiner

FIG. 1




RAPIDLY SOLUBLE DRUG COMPOSITION

TECHNICAL FIELD

The present invention relates to a fast-dissolving pharmaceutical composition of (R)-2-(4-bromo-2-fluorobenzyl)-1,2,3,4-tetrahydropyrrolo[1,2-a]pyrazine-4-spiro-3'-pyrrolidine-1,2',3,5'-tetrone (hereinafter, referred to as "AS-3201") having a potent aldose reductase inhibitory activity.

BACKGROUND ART

AS-3201 is the compound of the following formula. Said compound is described in Example 22 of Japanese Patent No. 2516147 (U.S. Pat. No. 5,258,382), Reference Example 12 of JP-A-6-192222 (Chem. Abstr., 122, 9860 (1995)), and Experiment of JP-A-8-176105 (Chem. Abstr., 125, 221569 (1996)), and its potent aldose reductase inhibitory activities are disclosed therein.



Example 28 of Japanese Patent No. 2516147 (USP 5258382) describes a method for preparing specific tablets of AS-3201. That is, it is described therein that AS-3201 (1 g), corn starch (25 g), lactose (58 g), crystalline cellulose (11 g), hydroxypropylcellulose (3 g), light anhydrous silicic acid (1 g) and magnesium stearate (1 g) are blended, granulated and made into 1,000 tablets each weighing 100 mg by a conventional method.

During the studies on methods for preparing AS-3201-containing pharmaceutical compositions having an excellent bioavailability, the present inventors have found that the water-solubility of said substance is extremely low in the range of low pH to the extent of several $\mu\text{g}/\text{ml}$, and therefore, the plasma concentration of AS-3201 varies widely among the individuals to be administered.

Under such circumstances, the present inventors have further intensively studied, and have found that by using micronized AS-3201 in a composition, the dissolution characteristics of said substance from the composition are remarkably improved, and as a result, an AS-3201-containing fast-dissolving pharmaceutical composition having a good bioavailability can be obtained, and finally have accomplished the present invention.

DISCLOSURE OF INVENTION

The present invention provides a fast-dissolving pharmaceutical composition comprising micronized AS-3201.

The terms used in the present specification are explained below.

The "micronized AS-3201" means powders of AS-3201 having a mean particle size of less than about 20 μm . The "mean particle size" means a particle size of being at 50% in cumulative particle distribution on weight or volume basis (ref., HA Lieberman et al., "Pharmaceutical Dosage Forms:

Tablets", Marcel Dekker, Inc., New York, 1990, vol. 2, 174-186; Kouichi IINOYA (edit.) "Handbook of Powder and Particle Measurement (in Japanese)", The NIKKAN KOGYO SHINBUN LTD., 1981, 29-36). The "dissolution test" means a test in which the dissolution of AS-3201 from test pharmaceutical compositions in an amount corresponding to 20 mg of AS-3201 is evaluated according to Paddle method (50 rpm) specified in the Twelfth Edition of the Pharmacopoeia of Japan, using a 0.2 M phosphate buffer (pH 6.5, 900 ml) as a test solution, and assaying AS-3201 by spectrophotometry at 300 nm. The " pK_{a1} " means an acid dissociation exponent of an acidic substance at 25° C. in an infinitely diluted solution thereof. When an acidic substance is a polybasic acid, it means an acid dissociation exponent at the first step of dissociation. The "water-solubility" means a maximum amount of a solute being dissolved in 100 ml of water. The term "about" is used with the intention of including values following said term.

The mean particle size of the micronized AS-3201 is preferably less than about 10 μm , more preferably less than about 5 μm , and most preferably in the range of about 0.5 μm to about 3 μm .

According to the method disclosed in Japanese Patent No. 2516147 (U.S. Pat. No. 5,258,382), crystals of AS-3201 having a mean particle size of about 60 μm to about 120 μm can usually be obtained. The micronization of AS-3201 crystals is carried out using a mill that is conventionally used in the pharmaceutical field. Mills are, for example, a fluid energy mill such as Jet Mill (manufactured by SEISHIN ENTERPRISE Co., LTD., Japan), a high speed rotative impact mill such as Sample Mill (manufactured by Hosokawa Micron Corporation, Japan), Pin Mill (manufactured by ALPINE, Germany), or Angmill (manufactured by Hosokawa Micron Corporation, Japan), a wet form high speed tumbling trituration mill such as MICROS (manufactured by Nara Machinery Co., Ltd., Japan), and a tumbling mill such as a ball mill. In order to obtain micronized powders having a mean particle size of less than about 5 μm , a fluid energy mill is preferably used. The micronization can be carried out on AS-3201 crystals alone, or on a mixture of AS-3201 crystals and a part or whole of pharmaceutical excipients or carriers, which are used in the preparation of pharmaceutical compositions.

The AS-3201-containing fast-dissolving pharmaceutical composition of the present invention may be solid dosage forms, and includes, for example, tablets, capsules, granules, powders, etc. These pharmaceutical compositions can be prepared by mixing micronized AS-3201 with pharmaceutical excipients or carriers such as diluents, disintegrators, binders and lubricants by a conventional method. For example, the mixture is granulated by wet-granulation such as high-shear granulation, fluid bed granulation, agitation fluid bed granulation, centrifugal fluid bed granulation, or extrusion granulation, or by dry-granulation such as roller compaction or slugging, and then the resulting granules are put into capsules for capsule preparations, or compressed for tablet preparations. Alternatively, a mixture of micronized AS-3201 and pharmaceutical excipients or carriers can directly be put into capsules for capsule preparations, or compressed for tablet preparations. These pharmaceutical compositions may optionally be coated, or may additionally contain stabilizers, surfactants, coloring agents, flavoring agents, etc.

The pharmaceutical excipients or carriers may be any ones except for ones showing a bad compatibility with AS-3201. The diluents include, for example, lactose, starch, crystalline cellulose, D-mannitol, sucrose, glucose, erythritol, xylitol, D-sorbitol, anhydrous dibasic calcium phosphate, and calcium sulfate. The disintegrators are, for example, starch, crystalline cellulose, low substituted hydroxypropylcellulose, carmellose, carmellose calcium, sodium carboxymethyl starch, croscarmellose sodium, partly pregelatinized starch, and hydroxypropyl starch. The binders are, for example, acacia, starch, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinyl alcohol, pullulan, gelatin, ethylcellulose, methylcellulose, carmellose sodium, and dextrin. The lubricants are, for example, magnesium stearate, calcium stearate, stearic acid, sucrose esters of fatty acids, light anhydrous silicic acid, talc, hydrogenated oil, and macrogol.

The stabilizer may be any pharmaceutically acceptable acidic substances having an acidity more potent than that of AS-3201, i.e., $pK_a=5.6-5.8$, and preferable acidic substances are ones having a pK_{a1} of less than about 4.5 and a water-solubility of larger than about 10 g/100 ml at 15° C.–25° C. More preferable acidic substances are ones having a pK_{a1} of less than about 3.3 and a water-solubility of larger than about 50 g/100 ml at 15° C.–25° C. Especially preferable acidic substances are, for example, citric acid, tartaric acid, maleic acid, and phosphoric acid. Among these acidic substances, tartaric acid is most preferable. The content of the acidic substance is preferably in the range of about 0.5% by weight to about 2.5% by weight. It is preferable to add a stabilizer in the case of preparing a pharmaceutical composition containing AS-3201 in a ratio of less than about 5% by weight.

The surfactants to be used in the present pharmaceutical composition are, for example, sorbitan fatty acid esters and polysorbates. The coloring agents are, for example, tar color, caramel, and red iron oxide. The flavoring agents are, for example, sweeteners and perfumes.

The dissolution characteristics of the active substance from the composition can be remarkably improved by using micronized AS-3201, and by further controlling the combination ratio of pharmaceutical excipients or carriers, AS-3201-containing fast-dissolving pharmaceutical compositions having more improved dissolution characteristics as well as good bioavailability can be obtained. The combination ratio of the pharmaceutical excipients or carriers may vary depending on the content of AS-3201. The content of AS-3201 in the present fast-dissolving pharmaceutical composition is usually in the range of about 0.5% by weight to about 25% by weight, to the total weight of the pharmaceutical composition. When the content of AS-3201 is in the range of about 0.5% by weight to 5% by weight to the total weight of the pharmaceutical composition, then the pharmaceutical composition usually comprises a diluent in a ratio of about 51% by weight—about 93.8% by weight, a disintegrator in a ratio of about 5% by weight—about 35% by weight, a binder in a ratio of about 0.5% by weight—about 5% by weight, and a lubricant in a ratio of about 0.2% by weight—about 4% by weight. More preferably, the pharmaceutical composition comprises a diluent in a ratio of about 59% by weight—about 88% by weight, a disintegrator in a ratio of about 10% by weight—about 30% by weight, a binder in a ratio of about 1% by weight—about 3% by weight, and a lubricant in a ratio of about 0.5% by weight—about 3% by weight. When the content of AS-3201 is more than 5% by weight and less than about 25% by weight to the total weight of the pharmaceutical composition, then the

present composition usually comprises a diluent in a ratio of about 16% by weight—about 84.3% by weight, a disintegrator in a ratio of about 10% by weight—about 50% by weight, a binder in a ratio of about 0.5% by weight—about 5% by weight, and a lubricant in a ratio of about 0.2% by weight—about 4% by weight, and more preferably, a diluent in a ratio of about 29% by weight—about 73.5% by weight, a disintegrator in a ratio of about 20% by weight—about 40% by weight, a binder in a ratio of about 1% by weight—about 3% by weight, and a lubricant in a ratio of about 0.5% by weight—about 3% by weight.

Since AS-3201 has an extremely low water-solubility to the extent of several $\mu\text{g/ml}$ in the range of low pH, there is a correlation between the initial dissolution rate and the bioavailability of AS-3201-containing pharmaceutical compositions, and compositions having a better initial dissolution rate can show a better bioavailability. From the viewpoint of the above, preferable compositions are ones having a dissolution percentage of the active substance of 50% or more for 15 minutes after the start of the dissolution test, and more preferable pharmaceutical compositions are ones having a dissolution percentage of the active substance of 80% or more for 15 minutes after the start of the dissolution test.

The AS-3201-containing fast-dissolving pharmaceutical composition of the present invention may be packed in a bottle using materials of low moisture-permeability or in damp-proof packages such as heat-sealed packages, if necessary.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 is a graph showing a dissolution pattern of the tablets of Examples 1 and 2, and Comparative Example 1.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention is illustrated in more detail by Examples and Comparative Example, but the present invention should not be construed to be limited thereto. The mean particle size was measured using a laser diffraction particle size distribution analyzer (HELOS & RODOS (trademark), manufactured by SYMPATEC GmbH, Germany), and calculated from cumulative particle distribution on volume basis by dry air dispersion method (dispersion air pressure: 0.5 atm).

EXAMPLE 1

Preparation of tablets:

| | |
|--|--------|
| AS-3201 | 160 g |
| Tartaric acid | 8 g |
| Lactose | 492 g |
| Low substituted hydroxypropylcellulose | 300 g |
| Hydroxypropylcellulose | 20 g |
| Magnesium stearate | 20 g |
| Total | 1000 g |

AS-3201 crystals were micronized using Single Truck Jet Mill (manufactured by SEISHIN ENTERPRISE CO., LTD., hereinafter abbreviated as "Jet Mill") with compression air pressure of 6 kgf/cm^2 to give powders having a mean particle size of about 1.5 μm . The micronized AS-3201 powders thus obtained, lactose, and low substituted hydrox-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.