

BML Stacks
W1
NA772
V. 342
No. 6251

Nature

INTERNATIONAL WEEKLY JOURNAL OF SCIENCE

Volume 342 No.6251 14 December 1989 \$5.75

#00000262907#NA MAY90
UNIV OF CA SAN DIEGO
SERIALS LIFE SCIENCES
BIOMEDICAL LIB C-0758
LA JOLLA CA 92093
DIR 10

Nature
BML Stacks
UNIVERSITY OF CALIFORNIA,
SAN DIEGO - LIBRARIES
Received on: 12-29-89
v. 1- Nov. 4, 1869-

RESERVE COPY

RETURN TO
CIRCULATION DESK

LAKES OF LIQUID SULPHUR

RECOMBINANT DNA
product review



nature

14 December 1989
Vol. 342 Issue no. 6251

◀ Sulphur volcanoes grew on the desiccated lake floor of Poás crater, Costa Rica, earlier this year. This 2-metre-high cone later collapsed revealing a bubbling pool of liquid sulphur. Photo by David Stevenson. See page 790.

THIS WEEK

Setting the baseline

The greenhouse effect, its extent, extent and implications, has been the subject of hundreds of model and theoretical studies. Yet the physical basis of the effect — the infrared radiation energy trapped by atmospheric gases and clouds — has not previously been measured from observational data. Raval and Ramanathan (page 758) have now filled that gap, presenting an effective method for directly monitoring future changes in the greenhouse effect. See also News and Views, page 736.

Channel vision

The light-sensitivity of vertebrate photoreceptor cells depends on the closure of a cation channel which is gated directly by cyclic GMP, the internal messenger of visual transduction. Now, the cloning and sequencing of the complementary DNA for this channel from bovine retinal rod photoreceptors shows that it could belong to a new family of ligand-gated ion channels. Page 762.

Regal matters

Worker honeybees efficiently destroy worker-laid male eggs if a queen is present. This 'worker policing' ensures that the colony's male reproductives are sons of the queen, pages 796 and 741.

Positional cues

The *Hox-5* homoeogenes are activated in the developing limb bud of the chick in a temporal order that corresponds to their spatial ordering on the chromosome, and in a series of partially overlapping spatial domains such that the region in which all five genes are expressed coincides with the zone of polarizing activity that determines the antero-posterior limb axis. Pages 767 and 734.

Insulin mechanism

Changes in our understanding of how insulin regulates glucose uptake are likely to follow the discovery that the endothelial cells that line the capillaries of the insulin-regulated tissues, muscle and fat express high levels of the insulin-regulated glucose transporter. Page 798.

Beneath the rings

Infrared imaging has revealed several new features in the atmosphere of Saturn — including a hotspot at the north pole and an asymmetric structure at the equator. Page 777.

Yeast shows the way

The processes of differentiation by induction of different gene regulatory proteins under the influence of cell-cell signals, and of the establishment of stem cell lineages, fundamental to metazoan development, can be illustrated in the single-celled yeast. Recent work illuminates the molecular basis of some of these processes. Review Article, page 749. See also page 830.

Hairpin corner

Oligonucleotide encoding telomeric DNA sequences form stable dimers in solution, possibly by the formation of hydrogen bonds between two intramolecular hairpin loops to form antiparallel quadruplexes containing cyclic guanine base tetrads. Pages 825 and 737.

Alvin's finds

Samples of sediment taken by the research submersible *Alvin* at the Guaymas Basin hot vents (Gulf of California), contain a group of methanogenic archaeobacteria growing, quite unexpectedly, at a temperature of at least 110°C (page 833). From the same area, dense layers of large filamentous sulphur-oxidizing bacteria of the genus *Beggiatoa* have been found at a depth of 2010 m (page 834).

NATURE SAYS

Europe must define its objectives ■ An outstanding broadcasting system is endangered 721

NATURE REPORTS

India's new stance ■ Human genome matters ■ After the 'quake ■ Early reptile on the move ■ Drift-nets cause conflict ■ West German reactor ■ Support for East Germany ■ Aviation milestone ■ Mono Lake reprieved ■ French universities ■ MRC ■ AIDS epidemic ■ IBM awards 723

CORRESPONDENCE

Ranking universities ■ Reviewing policy ■ Carbon tax ■ Journal costs ■ Etcetera 729

COMMENTARY

Keeping up with the Russians
G W Fisher, P C Grew & B Yardley 731

NEWS AND VIEWS

Is the salami sliced too thin? John Maddox 733
Vertebrate limb development: A pattern emerges
Julian Lewis & Paul Martin 734
Diamond films: Low-pressure nucleation routes
Michael Pinneo 735
Greenhouse effect: Gauging water-vapour feedback
Robert D Cess 736
DNA structure: The turn of the quadruplex?
Maxim Frank-Kamenetskii 737
Palaeontology: Four legs to stand on for Devonian vertebrates
Henry Gee; Per Erik Ahlberg 738
Scanning atomic microscopy: Variations on an original theme
C F Quate 739
Mid-ocean ridges: Propagating rifts exposed
Ken C Macdonald 740
Social insects: Who are the drone police?
Jon Seger 741
Daedalus: A clean press 742

SCIENTIFIC CORRESPONDENCE

Sampling the lithosphere N W Rogers, C J Hawkesworth, D S Ormerod & P D Kempton: Reply —
W F McDonough, K P Jochum, H Palme & B Spittel 743
Cleaning up after Chernobyl K F Baverstock 744
Early warning for LTPR Anwyl 744

BOOK REVIEWS

Origins and Evolution of Planetary and Satellite Atmospheres S K Atreya, J B Pollack & M S Matthews eds
William B McKinnon 745
Debating Archaeology by L R Binford Warwick Bray ■
Diverse Divers: Physiology and Behavior
by G L Kooyman P J Butler 746
Boninites and Related Rocks A J Crawford ed,
Carbonatites; Genesis and Evolution K Bell ed
H S Yoder Jr 747
Speciation and its Consequences D Otte & J A Endler eds
Mark Ridley 748

REVIEW ARTICLE

A regulatory hierarchy for cell specialization in yeast
I Herskowitz 749▶

Nature® (ISSN 0028-0836) is published weekly on Thursday, except the last week in December, by Macmillan Magazines Ltd (4 Little Essex Street, London WC2R 3LF). Annual subscription for USA and Canada US\$295 (institutional/corporate), US\$125 (individual making personal payment). USA and Canadian orders to: Nature, Subscription Dept, PO Box 1733, Riverton, NJ 08077-7333, USA. Other orders to: Nature, Brunel Road, Basingstoke, Hants RG21 2XS, UK. Second class postage paid at New York, NY 10012 and additional mailing offices. Authorization to photocopy material for internal or personal use, or internal or personal use of specific clients, is granted by Nature to libraries and others registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided the base fee of \$1.00 a copy plus \$0.10 a page is paid direct to CCC, 21 Congress Street, Salem, MA 01970, USA. Identification code for Nature: 0028-0836/89 \$1.00 + \$0.10. US Postmaster send address changes to: Nature, 65 Bleeker Street, New York, NY 10012. Published in Japan by Nature Japan K.K., Shin-Mitsuke Bldg, 2F, 1-10-1, Shinjuku-ku, Tokyo 162, Japan. © 1989 Macmillan Magazines Ltd.

25. Ullier, C. D. *Tectonics and Landforms* (Longman, New York, 1981).
 26. Fitch, F. J. & Miller, J. A. *Spec. Publ. geol. Soc. S. Afr.* **13**, 247-266 (1984).
 27. Cox, K. G. *J. Petrol.* **21**, 629-650 (1980).
 28. England, P. C. & Molnar, P. *Geology* (in the press).
 29. Petri, S. & Furlanò, V. *J. Geologia do Brasil* (Editora da Universidade de São Paulo, 1983).
 30. Ankrand, A. J. et al. *Crustal Evolution of Southern Africa* (Springer, New York, 1982).
 31. Craddock, C. (ed.) *Antarctic Geoscience* (University of Wisconsin Press, Madison, 1982).
 32. King, L. C. *South African Scenery* 2nd edn (Oliver and Boyd, Edinburgh, 1951).
 33. King, L. C. *The Morphology of the Earth* (Oliver and Boyd, Edinburgh, 1962).
 34. Partridge, T. C. & Maud, R. R. *S. Afr. geol. J.* **90**, 179-208 (1987).
 35. Cahen, L., Snelling, N. J., Delhal, J. & Val, J. R. *The Geochronology and Evolution of Africa* (Clarendon, Oxford, 1984).
 36. Brown, R. W. *6th Int. Conf. Fission Track Dating Abstr.* Vol. Université de Franche-Comté, Besançon, 1988).

Conformations of immunoglobulin hypervariable regions

Cyrus Chothia^{††}, Arthur M. Lesk^{*‡}, Anna Tramontano[‡], Michael Levitt^{*§}, Sandra J. Smith-Gill^{||}, Gillian Air[¶], Steven Sheriff^{**}, Eduardo A. Padlan[#], David Davies[#], William R. Tulip^{††}, Peter M. Colman^{††}, Silvia Spinelli^{‡‡}, Pedro M. Alzari^{‡‡} & Roberto J. Poljak^{‡‡}

* MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, UK

‡ European Molecular Biology Laboratory, Meyerhofstrasse 1, Postfach 102209, D-6900 Heidelberg, FRG

§ Department of Cell Biology, Stanford University Medical School, Stanford, California 94305, USA

¶ Christopher Ingold Laboratory, University College London, 20 Gordon Street, London WC1H 0AJ, UK

|| National Institute of Cancer and # National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892, USA

† Department of Microbiology, University of Alabama, Birmingham, Alabama 35294, USA

†† CSIRO Division of Biotechnology, 343 Royal Parade, Parkville 3052, Australia

‡‡ Unité d'Immunologie Structurale, Département d'Immunologie, Institut Pasteur, 25 rue du Dr Roux, 75724 Paris, France

On the basis of comparative studies of known antibody structures and sequences it has been argued that there is a small repertoire of main-chain conformations for at least five of the six hypervariable regions of antibodies, and that the particular conformation adopted is determined by a few key conserved residues. These hypotheses are now supported by reasonably successful predictions of the structures of most hypervariable regions of various antibodies, as revealed by comparison with their subsequently determined structures.

THE relationships between the amino-acid sequences of immunoglobulins and the structures of their antigen-binding sites are important for understanding the molecular mechanisms of the generation and maturation of the immune response and for designing engineered antibodies. Antigen-binding sites are formed by six loops of polypeptide, the hypervariable regions; three from the variable domain of the light chain (VL) and three from the variable domain of the heavy chain (VH), denoted L1, L2, L3, and H1, H2, H3, respectively (Fig. 1a). Within the domains, the loops are connected to a β -sheet framework whose structure is conserved^{1,2}. The specificity and affinity of the binding sites are governed by the structures of the six hypervariable regions^{3,4}.

Two models can be proposed for the relationship between the amino-acid sequence and structure of the binding-site loops. In one model, different sequences produce different conformations for both the main chain and side chains of the loops. Because hypervariable regions have different sequences in different antibodies, this model implies that each region adopts a different conformation in different antibodies. In the other

model, antibodies have only a few main-chain conformations or 'canonical structures' for each hypervariable region. Most sequence variations would only modify the surface provided by the side chains on a canonical main-chain structure. Sequence changes at a few specific sets of positions would switch the main chain to a different canonical conformation.

Canonical structure model

Experimental evidence indicates that the canonical structure model describes the relationship between amino-acid sequence and structure for at least five of the six hypervariable regions⁵⁻⁹. Kabat *et al.*⁵ found conserved residues at sites within certain sets of hypervariable regions and suggested that they had a structural role. Padlan and Davies⁶, and more recently de la Paz *et al.*⁷, showed that some of the hypervariable regions in the immunoglobulins of known structure have the same main-chain conformation in spite of several differences in sequence.

Chothia and Lesk⁸ identified the residues that through packing, hydrogen bonding, or the ability to assume unusual values of the torsion angles ϕ , ψ or ω , are primarily responsible for the main-chain conformations of the hypervariable regions in the structures then known—the Fab fragments of NEW (ref. 10), McPC603 (ref. 11), KOL (ref. 12) and J539 (ref. 13) and the VL domains of REI (ref. 14) and RHE (ref. 15). The conformations are determined by the interactions of a few residues at specific sites in the hypervariable regions and, for certain loops, in the framework regions. Hypervariable regions that have the same conformations in different immunoglobulins have the same or very similar residues at these sites (Fig. 1 and Table 1). Examination of the amino-acid sequence of the antibody D1.3 showed that its hypervariable regions are the same size as those in known structures and contain the same or similar residues at the sites responsible for known conformations⁹. On the basis of these observations the atomic structure of the VL-VH dimer of D1.3 was predicted before its experimental determination. Comparison of this predicted structure with the preliminary crystal structure showed that the conformations of four of the hypervariable regions had been predicted correctly; the conformation of L3 was significantly different from that predicted, and

** Present address: The Squibb Institute for Medical Research, PO Box 4000, Princeton, New Jersey 08543-4000, USA.

ARTICLES

TABLE 1 Sequences and conformations of V_k and V_H hypervariable regions of known structure

L1 Regions†

Canonical Structure	Protein	26	27	28	29	30	31	a	b	c	d	e	f	32	2	25	33	71
1	J539	S	S	S	V	S	—	—	—	—	—	—	—	S	I	A	L	Y
	HyHEL-5	S	S	S	V	N	—	—	—	—	—	—	—	Y	I	A	M	Y
	NQ10	S	S	S	V	R	—	—	—	—	—	—	—	Y	I	A	M	Y
2	REI	S	Q	D	I	I	K	—	—	—	—	—	—	V	I	A	L	Y
	D1.3	S	G	N	I	H	N	—	—	—	—	—	—	Y	I	A	L	Y
	HyHEL-10	S	Q	S	I	G	N	—	—	—	—	—	—	N	I	A	L	F
	NC41	S	Q	D	V	S	T	—	—	—	—	—	—	A	I	A	L	Y
3	McPC603	S	E	S	L	L	N	S	G	N	E	K	N	F	I	S	L	F
4	4.4-20	S	Q	S	L	V	H	S	—	N	G	N	T	Y	V	S	L	F

Total no. of sequences known for L1 regions: human, 95; mouse, 299.

Canonical structure	1	2	3	4
Human sequences that fit (%)	—	60	5	5
Mouse sequences that fit (%)	15	25	20	10

L2 Regions

Canonical Structure	Protein	50	51	52	48	64
1	REI	E	A	S	I	G
	McPC603	G	A	S	I	G
	J539	E	I	S	I	G
	D1.3	Y	T	T	I	G
	HyHEL-5	D	T	S	I	G
	HyHEL-10	Y	A	S	I	G
	NC41	W	A	S	I	G
	NQ10	D	T	S	I	G
	4.4-20	K	V	S	I	G

Total no. of sequences known for L2 regions: human, 69; mouse, 183.

Canonical structure	1
Human sequences that fit (%)	95
Mouse sequences that fit (%)	95

L3 Regions

Canonical Structure	Protein	91	92	93	94	95	96	90
1	REI	Y	Q	S	L	P	Y	Q
	McPC603	D	H	S	Y	P	L	N
	D1.3	F	W	S	T	P	R	H
	HyHEL-10	S	N	S	W	P	Y	Q
	NC41	H	Y	S	P	P	W	Q
	4.4-20	S	T	H	V	P	W	Q
	NQ10	W	S	S	N	P	L	Q
2	J539	W	T	Y	P	L	I	Q
3	HyHEL-5	W	G	R	N	P	—	Q

Total no. of sequences known for L3 regions: human, 52; mouse, 152.

Canonical structure	1	2	3
Human sequences that fit (%)	90	—	2
Mouse sequences that fit (%)	80	10	1

H1 had a very different fold from that predicted⁹. (We report below that the refined conformation of D1.3 corresponds more closely to the predicted structure.)

An examination of the library of the known immunoglobulin sequences shows that many immunoglobulins have hypervariable regions that are the same size as those in the known structures and contain the same or closely related residues at the sites responsible for the known conformations⁸. These observations indicate that for at least five of the hypervariable regions there is only a small repertoire of canonical main-chain conformations and that the conformation actually present can often be predicted from the sequence by the presence of specific residues.

The accuracy of the canonical structure model for immunoglobulin binding sites depends on (1) the correct deter-

mination of the sets of residues responsible for the observed conformations and (2) changes in the identity of residues at other sites not significantly affecting the conformations of the canonical structures. The model can be tested, refined and extended by using it to predict the atomic structures of binding sites in immunoglobulins before their structures have been determined by X-ray crystallography.

We have now tested the canonical structure model by using it to predict the structures of four immunoglobulins before their structures had been experimentally determined. These immunoglobulins are HyHEL-5 (ref. 16), HyHEL-10 (ref. 17), NC41 (ref. 18) and NQ10 (S.S., P.M.A. and R.J.P., manuscript in preparation). The analysis of the amino-acid sequences of these immunoglobulins indicated that 19 of their 24 hypervariable regions should have conformations close to known canoni-

H1 Regions‡

Canonical Structure	Protein	26	27	28	29	30	31	32	34	94
1	McPC603	*	*		*				*	*
	KOL	G	F	T	F	S	D	F	M	R
	J539	G	F	I	F	S	S	Y	M	R
	D1.3	G	F	D	F	S	K	Y	M	R
	HyHEL-5	G	F	S	L	T	G	Y	V	R
	NC41	G	Y	T	F	S	D	Y	I	R
	NQ10	G	Y	T	F	T	N	Y	M	R
	4-4-20	G	F	T	F	S	S	F	M	R
		G	F	T	F	S	D	Y	M	G
1'	NEW	G	S	T	F	S	N	D	Y	R
	HyHEL-10	G	D	S	I	T	D	D	W	N

Total no. of sequences known for H1 regions: human, 50; mouse, 321.

Canonical structure	1
Human sequences that fit (%)	50
Mouse sequences that fit (%)	80

H2 Regions§

Canonical Structure	Protein	52a	b	c	53	54	55	71
1	NEW *	—	—	—	Y	H	G	
	D1.3				G	D	G	
	HyHEL-10	—	—	—	Y	S	G	
2	HyHEL-5	*	—	—	G	S	G	(*)
	NC41	T	—	—	N	T	G	A
3	KOL	D	—	—	D	G	S	R
	J539	P	—	—	D	S	G	R
	NQ10	S	—	—	G	S	S	R
4	McPC603	N	K	G	N	K	Y	(*)
	4-4-20	N	K	P	Y	N	Y	R

Total no. of sequences known for H2 regions: human, 54; mouse, 248.

Canonical structure	1	2	3	4
Human sequences that fit (%)	15	1	40	15
Mouse sequences that fit (%)	15	40	5	20

The residues listed here (single-letter code) are those that form the hypervariable regions and those in the framework regions that are important for the observed conformations of these regions⁸. The hypervariable regions are taken as those outside the framework β -sheet⁸. Except for H2, they are similar to, but not identical with the regions that show high sequence variations and which Kabat *et al.*²⁶ use to define hypervariable regions. The sequences are grouped so that those that have the same main-chain conformation, or canonical structure, are adjacent. The canonical structure numbers used below refer to the conformations shown in Fig. 1. The residues in the hypervariable and framework regions that are mainly responsible for these conformations⁸ are indicated by an asterisk. The classification and sequence requirements of the H2 conformations have been revised in the light of work described here and elsewhere²⁸. For each hypervariable region the number of human and mouse sequences listed by Kabat *et al.*²⁶ are given. We also give the percentage of these sequences that are the same size as the known canonical structures and have the same residues at the positions marked by an asterisk.

† Canonical structure 4 is illustrated in Fig. 4. Although the size of the known L1 structures varies between 6 and 13 residues, they have closely related folds with residues 26-19 and 32 packed against the framework in the same conformation⁸. The remaining residues form a turn or loop on the surface (Figs 1 and 4). The ends of the long loops have some flexibility. There are another 25% of the human sequences and 20% of the mouse sequences that have one more residue than structure 2, or one fewer than structure 4, and whose sequences satisfy the requirements listed above. It is expected that these differ only in the conformations of the tips of the surface loops.

‡ The H1 hypervariable regions with canonical structure 1 have very similar conformations: the r.m.s. differences in the coordinates of their main-chain atoms are 0.3-0.8 Å. The H1 regions in NEW and HyHEL-10 only partly satisfy the sequence requirements for structure 1 and have a distorted version of its conformation.

§ The H2 region here comprises residues 52a-55. The region with high sequence variation is 50-65 (ref. 26). In the known structures the main-chain conformation of 50-52 and 56-63 do not differ significantly⁸ (Fig. 1b). (*), The residues at positions 55 or 54 in the canonical structures 2, 3 and 4 have residues with positive values for ϕ and ψ and usually, but not in all cases, Gly, Asn or Asp is found at these sites. For a sequence to match that of canonical structure 2, 3 or 4 the presence of these residues at sites 54 or 55 is required.

cal structures. We then compared the predicted structures of these hypervariable regions with the subsequently determined structures. Another immunoglobulin structure, 4-4-20 (ref. 19) has recently been reported. We did not have the opportunity to predict the structure of 4-4-20 before its experimental determination, and we discuss here only how its hypervariable regions have the conformations expected from the known canonical structures. Also, we report that the refined conformation of D1.3 (ref. 20) corresponds more closely to the predicted structure.

Model building procedure

The main-chain conformations of the hypervariable regions in the V κ and V λ domains of known structure are shown in Fig. 1. The residues responsible for these conformations are listed

in Table 1. Each hypervariable region in the immunoglobulins of unknown structure was examined to determine (1) whether it has the same size as any homologous hypervariable region of known structure and (2) whether its sequence contains the set of residues responsible for a known conformation. Except for L3 in HyHEL-5, all the light-chain regions correspond to a known canonical structure, as do all the H1 regions and the H2 region in HyHEL-10 (Table 1). The conformation predicted for the H2 region in NC41 and HyHEL-5 was based on the analysis of the H2 region in the preliminary structure of J539 (ref. 8). In all three of these antibodies the H2 region is a four-residue turn with Gly at the fourth position and the predicted conformation is that almost always found for such turns²¹. (Below we present a more accurate analysis of H2 regions.) For H3 regions in HyHEL-5, HyHEL-10, NC41 and NQ10, no prediction of

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.