



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12P 21/00, C12N 5/10, 7/01, 15/00 *	A1	(11) International Publication Number: WO 90/07861 (43) International Publication Date: 26 July 1990 (26.07.90)
(21) International Application Number: PCT/US89/05857 (22) International Filing Date: 28 December 1989 (28.12.89) (30) Priority data: 290,975 28 December 1988 (28.12.88) US 310,252 13 February 1989 (13.02.89) US (71) Applicant: PROTEIN DESIGN LABS, INC. [US/US]; 3181 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors: QUEEN, Cary, L. ; 1300 Oak Creek Drive, Palo Alto, CA 94304 (US). SELICK, Harold, Edwin ; 1673 Sunnyslope Avenue, Belmont, CA 94002 (US). (74) Agent: SMITH, William, M.; Townsend and Townsend, One Market Plaza, 2000 Steuart Tower, San Francisco, CA 94105 (US).		(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (Euro- pean patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI pa- tent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: CHIMERIC IMMUNOGLOBULINS SPECIFIC FOR p55 TAC PROTEIN OF THE IL-2 RECEPTOR		
(57) Abstract <p>Novel methods for designing humanized immunoglobulins having one or more complementary determining regions (CDR's) from a donor immunoglobulin and a framework region from a human immunoglobulin comprising first comparing the framework or variable region amino acid sequence of the donor immunoglobulin to corresponding sequences in a collection of human immunoglobulin chains, and selecting as the human immunoglobulin one of the more homologous sequences from the collection. Each humanized immunoglobulin chain may comprise about 3 or more amino acids from the donor immunoglobulin in addition to the CDR's, usually at least one of which is immediately adjacent to a CDR in the donor immunoglobulin. The heavy and light chains may each be designed by using any one or all three additional position criteria. When combined into an intact antibody, the humanized immunoglobulins of the present invention will be substantially non-immunogenic in humans and retain substantially the same affinity as the donor immunoglobulin to the antigen, such as a protein or other compound containing an epitope.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LJ	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

CHIMERIC IMMUNOGLOBULINS SPECIFIC FOR p55 TAC PROTEIN
OF THE IL-2 RECEPTOR

Field of the Invention

5 The present invention relates generally to the combination of recombinant DNA and monoclonal antibody technologies for developing novel therapeutic agents and, more particularly, to the production of non-immunogenic antibodies and their uses.

10 Background of the Invention

In mammals, the immune response is mediated by two types of cells that interact specifically with foreign material, i.e., antigens. One of these cell types, B-cells, are responsible for the production of antibodies. The second cell class, T-cells, include a wide variety of cellular subsets controlling the in vivo function of both B-cells and a wide variety of other hematopoietic cells, including T-cells.

20 One way in which T-cells exert this control is through the production of a lymphokine known as interleukin-2 (IL-2), originally named T-cell growth factor. IL-2's prime function appears to be the stimulation and maintenance of T-cells. Indeed, some immunologists believe that IL-2 may be at the center of the entire immune response (see, Farrar, J., et al., Immunol. Rev. 63:129-166 (1982), which is incorporated herein by reference).

30 To exert its biological effects, IL-2 interacts with a specific high-affinity membrane receptor (Greene, W., et al., Progress in Hematology XIV, E. Brown, Ed., Grune and Statton, New York (1986), at pgs. 283 ff). The human IL-2 receptor is a complex multichain glycoprotein, with one chain, known as the Tac peptide, being about 55kD in size (see, Leonard, W., et al., J. Biol. Chem. 260:1872 (1985), which is incorporated herein by reference). A gene encoding this protein has been isolated, and predicts a 272 amino acid peptide, including a 21 amino acid signal peptide (see, Leonard, W., et al., Nature 311: 626 (1984)). The 219 NH₂-

terminal amino acids of the p55 Tac protein apparently comprise an extracellular domain (see, Leonard, W., et al., Science, 230:633-639 (1985), which is incorporated herein by reference).

5 Much of the elucidation of the human IL-2
receptor's structure and function is due to the development
of specifically reactive monoclonal antibodies. In
particular, one mouse monoclonal antibody, known as anti-Tac
(Uchiyama, et al., J. Immunol. 126:1393 (1981)) has shown
10 that IL-2 receptors can be detected on T-cells, but also on
cells of the monocyte-macrophage family, Kupffer cells of the
liver, Langerhans' cells of the skin and, of course,
activated T-cells. Importantly, resting T-cells, B-cells or
circulating macrophages typically do not display the IL-2
15 receptor (Herrmann, et al., J. Exp. Med. 162:1111 (1985)).

 The anti-Tac monoclonal antibody has also been used
to define lymphocyte functions that require IL-2 interaction,
and has been shown to inhibit various T-cell functions,
including the generation of cytotoxic and suppressor T
20 lymphocytes in cell culture. Also, based on studies with
anti-Tac and other antibodies, a variety of disorders are now
associated with improper IL-2 receptor expression by T-cells,
in particular adult T-cell leukemia.

 More recently, the IL-2 receptor has been shown to
25 be an ideal target for novel therapeutic approaches to T-cell
mediated diseases. It has been proposed that IL-2 receptor
specific antibodies, such as the anti-Tac monoclonal
antibody, can be used either alone or as an immunoconjugate
(e.g., with Ricin A, isotopes and the like) to effectively
30 remove cells bearing the IL-2 receptor. These agents can,
for example, theoretically eliminate IL-2 receptor-expressing
leukemic cells, certain B-cells, or activated T-cells
involved in a disease state, yet allow the retention of
mature normal T-cells and their precursors to ensure the
35 capability of mounting a normal T-cell immune response as
needed. In general, most other T-cell specific agents can
destroy essentially all peripheral T-cells, which limits the
agents' therapeutic efficacy. Overall, the use of

appropriate monoclonal antibodies specific for the IL-2 receptor may have therapeutic utility in autoimmune diseases, organ transplantation and any unwanted response by activated T-cells. Indeed, clinical trials have been initiated using, 5 e.g., anti-Tac antibodies (see, generally, Waldman, T., et al., Cancer Res. 45:625 (1985) and Waldman, T., Science 232:727-732 (1986), both of which are incorporated herein by reference).

10 Unfortunately, the use of the anti-Tac and other non-human monoclonal antibodies have certain drawbacks, particularly in repeated therapeutic regimens as explained below. Mouse monoclonal antibodies, for example, do not fix human complement well, and lack other important immunoglobulin functional characteristics when used in 15 humans.

Perhaps more importantly, anti-Tac and other non-human monoclonal antibodies contain substantial stretches of amino acid sequences that will be immunogenic when injected into a human patient. Numerous studies have shown that, 20 after injection of a foreign antibody, the immune response elicited by a patient against an antibody can be quite strong, essentially eliminating the antibody's therapeutic utility after an initial treatment. Moreover, as increasing numbers of different mouse or other antigenic (to humans) 25 monoclonal antibodies can be expected to be developed to treat various diseases, after the first and second treatments with any different non-human antibodies, subsequent treatments even for unrelated therapies can be ineffective or even dangerous in themselves.

30 While the production of so-called "chimeric antibodies" (e.g., mouse variable regions joined to human constant regions) has proven somewhat successful, a significant immunogenicity problem remains. In general, the production of human immunoglobulins reactive with the human 35 IL-2 receptor, as with many human antigens, has been extremely difficult using typical human monoclonal antibody production techniques. Similarly, utilizing recombinant DNA technology to produce so-called "humanized" antibodies (see,

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.