V 17	715	272		
,PRESS MAIL	LABEL M	NO, 85	993	7585
DATE N	IAILED:	June	14,	1991

29 1991 H

GENENTECH, INC. 460 Point San Bruno Boulevard, South San Francisco, CA 94080 (415) 266-1000

Docket No. 709

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

NEW APPLICATION TRANSMITTAL

SIR:

Transmitted herewith for filing is the patent application of Inventor(s): PAUL J. CARTER ET AL.

Title: IMMUNOGLOBULIN VARIANTS

CERTIFICATION UNDER 37 CFR §1.10

I hereby certify that this New Application and the documents referred to as enclosed herein are being deposited with the United States Postal Service on this date June 14, 1991, in an envelope bearing "Express Mail Post Office To Addressee" Mailing Label Number 859937585 addressed to: Patent Application, Honorable Commissioner of Batents, and Trademarks, Mashington, D.C. 20231.

Carolyn	R.	Adler
		Hatter

(Name of person mailing paper)

Enclosed are:

- 1. The papers required for filing date under CFR §1.53(b):
- 106 Pages of specification (including claims); _5_ Sheets of drawings (_ formal / _x_ informal)
- a. x Declaration/Oath/Power of Attorney
- 3. ____ Assignment of the invention to GENENTECH, INC.
- 4. Fee Calculation

	CLAINS AS FILED	••		
	/ Number Filed	Number Extra	Rate	Basic Fee \$630
Total Claims	16 - 20 =	· •	x \$20.00	630.
Indep. Claims	.8 - 3 ∝ .	* 5	× \$60.00	300.
_	Multiple dependent claim(s), if any	\$200.00	

*If less than zero, enter "0".

7. Recording Assignment [\$8.00]

8. Payment of Fees

9.

. .

x. Charge Account No. 07-0630 In the amount of \$_. A duplicate of this transmittal is attached.

1

<u>x</u> Authorization to Charge Additional Fees

- The Commissioner is hereby authorized to charge any additional fees (or credit any overpayment) associated with this communication and which may be required under 37 CFR §1.16 or §1.17 to Account No. 07-0630. <u>A duplicate sheet is attached.</u>
- 10. Information Disclosure Statement

11. x Return Receipt Postcard

C.All

Name: Carolyn R/Adler Registration No. 32,324

Dated <u>June 14, 1991</u>

Pfizer v. Genentech IPR2017-01489 Genentech Exhibit 2032



₩ 715272

FIGURE 1A: VL DOMAIN

50 40 30 10 20 DIVMTQSHKFMSTSVGDRVSITCKASQDVNTAVAWYQQKPGHSPKLLIYSASFRYT 4D5 DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLL IYSASFLES HU4D5 DIQMTQSPSSLSASVGDRVTITCRASQDVSSYLAWYQQKPGKAPKLLIYAASSLES HUVLKI VL-CDR1 V_L-CDR2

 60
 70
 80
 90
 100

 4D5
 GVPDRFTGNRSGTDFTFTISSVQAEDLAVYYCQQHYTTPPTFGGGTKLEIKRA

 1
 1
 1
 1

 HU4D5
 GVPSRFSGSRSGTDFTLTISSLQPEDFATYYCQQHYTTPPTFGQGTKVEIKRT


 HUVL×I
 GVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYNSLPYTFGQGTKVEIKRT

2

VL-CDR3

∅ 715272


FIGURE 1B: V_H DOMAIN




	60	70	80	ABC	90	100ABC
4D5					EDTAVYYCSR	WGGDGFYAMDYW
HU4D5	GYTRYADSVK	RFTISADT	SKNTAYLQ	MNSLRAI	SDTAVYYCSR	WGGDGFYAMDVW
		i_i	i		i	
HUV _H III	SDTYYADSVKG	RFTISRDDS	SKNTLYLQN	INSLRAE	DTAVYYCAR	DRGGAVSYFDVW
		•				

V_H-CDR3

4D5	110 ^P GQGASVTVSS
HU4D5	GQGTLVTVSS
HUV _H III	GQGTLVTVSS




Anneal huV_L or huV_H oligomers to pAK1 template





1 715272

FIGURE 3



FIGURE



15272

DOCKET 709 EXPRESS MAIL NO. B59937585 MAILED 14 JUNE 1991

IMMUNOGLOBULIN VARIANTS

Field of the Invention

This invention relates to methods for the preparation and use of variant antibodies and finds application particularly in the fields of immunology and cancer diagnosis and therapy.

Background of the Invention

Naturally occurring antibodies (immunoglobulins) comprise two heavy chains linked together by disulfide bonds and two light chains, one light chain being linked to each of the heavy chains by disulfide bonds. Each heavy chain has at one end a variable domain $\langle V_H \rangle$ followed by a number of constant domains. Each light chain has a variable domain $\langle V_L \rangle$ at one end and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains, see e.g. Chothia *et al.*, *J. Mol. Biol.* 186:651-663 (1985); Novotny and Haber, *Proc. Natl. Acad. Sci.*

1

7

15

20

25

30

10

USA 82:4592-4596 (1985).

The constant domains are not involved directly in binding the antibody to an antigen, but are involved in various effector functions, such as participation of the antibody in antibody-dependent cellular cytotoxicity. The variable domains of each pair of light and heavy chains are involved directly in binding the antibody to the antigen. The domains of natural light and heavy chains have the same general structure, and each domain comprises four framework (FR) regions, whose sequences are somewhat conserved, connected by three hyper-variable or complementarity determining regions (CDRs) (see Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda, MD, (1987)). The four framework regions largely adopt a β -sheet conformation and the CDRs form loops connecting, and in some cases forming part of, the framework regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site.

Widespread use has been made of monoclonal antibodies, particularly those derived from rodents including mice, however they are frequently antigenic in human clinical use. For example, a major limitation in the clinical use of rodent monoclonal antibodies is an anti-globulin response during therapy (Miller, R. A. *et al.*, *Blood* 62:988-995 (1983); Schroff, R. W. *et al.*, *Cancer Res.* 45:879-885 (1985)).

The art has attempted to overcome this problem by constructing "chimeric" antibodies in which an animal antigen-binding variable domain is coupled to a human constant domain (Cabilly *et al.*, U.S. patent No. 4,816,567; Morrison, S. L. *et al.*, *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984); Boulianne, G. L. *et al.*, *Nature* 312:643-646 (1984); Neuberger, M. S. *et al.*, *Nature* 314:268-270 (1985)). The term "chimeric" antibody is used herein to describe a polypeptide comprising at least the antigen binding portion of an antibody molecule linked to at least part of another protein (typically an immunoglobulin constant domain).

The isotype of the human constant domain may be selected to tailor

10

15

20

30

25

2

the chimeric antibody for participation in antibody-dependent cellular cytotoxicity (ADCC) and complement-dependent cytotoxicity (see e.g. Brüggemann, M. *et al.*, *J. Exp. Med.* **166**:1351-1361 (1987); Riechmann, L. *et al.*, *Nature* **332**:323-327 (1988); Love *et al.*, *Methods in Enzymology* 178:515-527 (1989); Bindon *et al.*, *J. Exp. Med.* **168**:127-142 (1988).

In the typical embodiment, such chimeric antibodies contain about one third rodent (or other non-human species) sequence and thus are capable of eliciting a significant anti-globulin response in humans. For example, in the case of the murine anti-CD3 antibody, OKT3, much of the resulting anti-globulin response is directed against the variable region rather than the constant region (Jaffers, G. J. *et al.*, *Transplantation* **41**:572-578 (1986)).

In a further effort to resolve the antigen binding functions of antibodies and to minimize the use of heterologous sequences in human antibodies, Winter and colleagues (Jones, P. T. *et al.*, *Nature* **321**:522-525 (1986); Riechmann, L. *et al.*, *Nature* **332**:323-327 (1988); Verhoeyen, M. *et al.*, *Science* **239**:1534-1536 (1988)) have substituted rodent CDRs or CDR sequences for the corresponding segments of a human antibody. As used herein, the term "humanized" antibody is an embodiment of chimeric antibodies wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The therapeutic promise of this approach is supported by the clinical efficacy of a humanized antibody specific for the CAMPATH-1 antigen with two non-Hodgkin lymphoma patients, one of whom had previously developed an anti-globulin response to the parental rat antibody (Riechmann, L. *et al.*, *Nature* 332:323-327 (1988); Hale, G. *et al.*, *Lancet* i:1394-1399 (1988)). A murine antibody to the interleukin 2 receptor has also recently been humanized (Queen, C. *et al.*, *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989)) as a potential immunosuppressive reagent. Additional references related to humanization of antibodies include Co *et al.*, *Proc. Natl. Acad. Sci.*

25

20

5

10

15

30

3

USA 88:2869-2873 (1991); Gorman *et al.*, *Proc. Natl. Acad. Sci. USA* 88:4181-4185 (1991); Daugherty *et al.*, *Nucleic Acids Research* 19(9):2471-2476 (1991); Brown *et al.*, *Proc. Natl. Acad. Sci. USA* 88:2663-2667 (1991); Junghans *et al.*, *Cancer Research* 50:1495-1502 (1990).

In some cases, substituting CDRs from rodent antibodies for the human CDRs in human frameworks is sufficient to transfer high antigen binding affinity (Jones, P. T. *et al.*, *Nature* **321**:522-525 (1986); Verhoeyen, M. *et al.*, *Science* **239**:1534-1536 (1988)), whereas in other cases it has been necessary to additionally replace one (Riechmann, L. *et al.*, *Nature* **332**:323-327 (1988)) or several (Queen, C. *et al.*, *Proc. Natl. Acad. Sci. USA* **86**:10029-10033 (1989)) framework region (FR) residues. See also Co *et al.*, *supra*.

For a given antibody a small number of FR residues are anticipated to be important for antigen binding. Firstly for example, certain antibodies have been shown to contain a few FR residues which directly contact antigen in crystal structures of antibody-antigen complexes (e.g., reviewed in Davies, D. R. et al., Ann. Rev. Biochem. 59:439-473 (1990)). Secondly, a number of FR residues have been proposed by Chothia, Lesk and colleagues (Chothia, C. & Lesk, A. M., J. Mol. Biol. 196:901-917 (1987); Chothia, C. et al., Nature 342:877-883 (1989); Tramontano, A. et al., J. Mol. Biol. 215:175-182 (1990)) as critically affecting the conformation of particular CDRs and thus their contribution to antigen binding. See also Margolies et al., Proc. Natl. Acad. Sci. USA 72:2180-2184 (1975).

It is also known that, in a few instances, an antibody variable domain (either V_H or V_L) may contain glycosylation sites, and that this glycosylation may improve or abolish antigen binding, Pluckthun, *Biotechnology* 9:545-51 (1991); Spiegelberg *et al.*, *Biochemistry* 9:4217-4223 (1970); Wallic *et al.*, *J. Exp. Med.* 168:1099-1109 (1988); Sox *et al.*, *Proc. Natl. Acad. Sci. USA* 66:975-982 (1970); Margni *et al.*, *Ann. Rev. Immunol.* 6:535-554 (1988). Ordinarily, however, glycosylation has no influence on the antigen-binding properties of an antibody, Pluckthun, *supra*, (1991).

10

15

20

5

25

The three-dimensional structure of immunoglobulin chains has been studied, and crystal structures for intact immunoglobulins, for a variety of immunoglobulin fragments, and for antibody-antigen complexes have been published (see e.g., Saul et al., Journal of Biological Chemistry 25:585-97 (1978); Sheriff et al., Proc. Natl. Acad. Sci. USA 84:8075-79 (1987); Segal et al., Proc. Natl. Acad. Sci. USA 71:4298-4302 (1974); Epp et al., Biochemistry 14(22):4943-4952 (1975); Marguart et al., J. Mol. Biol. 141:369-391 (1980); Furey et al., J. Mol. Biol. 167:661-692 (1983); Snow and Amzel, Protein: Structure, Function, and Genetics 1:267-279, Alan R. Liss, Inc. pubs. (1986); Chothia and Lesk, J. Mol. Biol. 196:901-917 (1987); Chothia et al., Nature 342:877-883 (1989); Chothia et al., Science 233:755-58 (1986); Huber et al., Nature 264:415-420 (1976); Bruccoleri et al., Nature 335:564-568 (1988) and Nature 336:266 (1988); Sherman et al., Journal of Biological Chemistry 263:4064-4074 (1988); Amzel and Poljak, Ann. Rev. Biochem. 48:961-67 (1979); Silverton et al., Proc. Natl. Acad. Sci. USA 74:5140-5144 (1977); and Gregory et al., Molecular Immunology 24:821-829 (1987). It is known that the function of an antibody is dependent on its three dimensional structure, and that amino acid substitutions can change the three-dimensional structure of an antibody, Snow and Amzel, supra. It has previously been shown that the antigen binding affinity of a humanized antibody can be increased by mutagenesis based upon molecular modelling (Riechmann, L. et al., Nature 332:323-327 (1988); Queen, C. et al., Proc. Natl. Acad. Sci. USA 86:10029-10033 (1989)).

Humanizing an antibody with retention of high affinity for antigen and other desired biological activities is at present difficult to achieve using currently available procedures. Methods are needed for rationalizing the selection of sites for substitution in preparing such antibodies and thereby increasing the efficiency of antibody humanization.

The proto-oncogene *HER2* (human epidermal growth factor receptor 2) encodes a protein tyrosine kinase (p185^{HER2}) that is related to and somewhat homologous to the human epidermal growth factor receptor

5

10

15

20

30

(see Coussens, L. *et al.*, *Science* 230:1132-1139 (1985); Yamamoto, T. *et al.*, *Nature* 319:230-234 (1986); King, C. R. *et al.*, *Science* 229:974-976 (1985)). *HER2* is also known in the field as *c-erbB-2*, and sometimes by the name of the rat homolog, *neu*. Amplification and/or overexpression of *HER2* is associated with multiple human malignancies and appears to be integrally involved in progression of 25-30% of human breast and ovarian cancers (Slamon, D. J. *et al.*, *Science* 235:177-182 (1987), Slamon, D. J. *et al.*, *Science* 244:707-712 (1989)). Furthermore, the extent of amplification is inversely correlated with the observed median patient survival time (Slamon, *supra*, Science 1989).

The murine monoclonal antibody known as muMAb4D5 (Fendly, B. M. et al., Cancer Res. 50:1550-1558 (1990)), directed against the extracellular domain (ECD) of p185HER2, specifically inhibits the growth of tumor cell lines overexpressing p185^{HER2} in monolayer culture or in soft agar (Hudziak, R. M. et al., Molec. Cell. Biol. 9:1165-1172 (1989); Lupu, R. et al., Science 249:1552-1555 (1990)). MuMAb4D5 also has the potential of enhancing tumor cell sensitivity to tumor necrosis factor, an important effector molecule in macrophage-mediated tumor cell cytotoxicity (Hudziak, supra, 1989; Shepard, H. M. and Lewis, G. D. J. Clinical Immunology 8:333-395 (1988)). Thus muMAb4D5 has potential for clinical intervention in and imaging of carcinomas in which p185HER2 is overexpressed. The muMAb4D5 and its uses are described in copending U.S. patent applications 07/143,912 and 07/147,461, and in corresponding PCT application WO 89/06692 published 27 July 1989. This murine antibody was deposited with the ATCC and designated ATCC CRL 10463. However, this antibody may be immunogenic in humans.

It is therefore an object of this invention to provide methods for the preparation of antibodies which are less antigenic in humans than non-human antibodies but have desired antigen binding and other characteristics and activities.

It is a further object of this invention to provide methods for the efficient humanization of antibodies, i.e. selecting non-human amino acid

6

10

Ņ

15

20

25

residues for importation into a human antibody background sequence in such a fashion as to retain or improve the affinity of the non-human donor antibody for a given antigen.

It is another object of this invention to provide humanized antibodies capable of binding p185^{HER2}.

Other objects, features, and characteristics of the present invention will become apparent upon consideration of the following description and the appended claims.

Summary of the Invention

a.

b.

c.

d.

e.

f.

The objects of this invention are accomplished by a method for making a humanized antibody comprising amino acid sequence of an import, non-human antibody and a human antibody, comprising the steps of:

> obtaining the amino acid sequences of at least a portion of an import antibody variable domain and of a consensus human variable domain;

identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human variable domain sequences;

substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;

identifying import antibody FR residues in the aligned FR sequences that are non-homologous to the corresponding consensus antibody residues;

determining if the non-homologous import amino acid residue is reasonably expected to have at least one of the following effects:

1. non-covalently binds antigen directly,

10

15

20

25

30

2. interacts with a CDR; or

g.

3. participates in the $V_L - V_H$ interface; and for any non-homologous import antibody amino acid residue which is reasonably expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence.

Optionally, the method of this invention comprises the additional steps of determining if any non-homologous residues identified in step (e) are exposed on the surface of the domain or buried within it, and if the residue is exposed but has none of the effects identified in step (f), retaining the consensus residue.

Additionally, in certain embodiments the method of this invention comprises the feature wherein the corresponding consensus antibody residues identified in step (e) above are selected from the group consisting of 4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H (utilizing the numbering system set forth in Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)).

In certain embodiments, the method of this invention comprises the additional steps of searching either or both of the import, non-human and the consensus variable domain sequences for glycosylation sites, determining if the glycosylation is reasonably expected to be important for the desired antigen binding and biological activity of the antibody (i.e., determining if the glycosylation site binds to antigen or changes a side chain of an amino acid residue that binds to antigen, or if the glycosylation enhances or weakens antigen binding, or is important for maintaining antibody affinity). If the import sequence bears the glycosylation site, it is preferred to substitute that site for the corresponding residues in the consensus human sequence if the glycosylation site is reasonably expected to be important. If only the

10

15

20

25

30

R

consensus sequence, and not the import, bears the glycosylation site, it is preferred to eliminate that glycosylation site or substitute therefor the corresponding amino acid residues from the import sequence.

Another embodiment of this invention comprises aligning import antibody and the consensus antibody FR sequences, identifying import antibody FR residues which are non-homologous with the aligned consensus FR sequence, and for each such non-homologous import antibody FR residue, determining if the corresponding consensus antibody residue represents a residue which is highly conserved across all species at that site, and if it is so conserved, preparing a humanized antibody which comprises the consensus antibody amino acid residue at that site.

Certain alternate embodiments of the methods of this invention comprise obtaining the amino acid sequence of at least a portion of an import, non-human antibody variable domain having a CDR and a FR, obtaining the amino acid sequence of at least a portion of a consensus human antibody variable domain having a CDR and a FR, substituting the non-human CDR for the human CDR in the consensus human antibody variable domain, and then substituting an amino acid residue for the consensus amino acid residue at at least one of the following sites:

> (in the FR of the variable domain of the light chain) 4L, 35L, 36L, 38L, 43L, 44L, 58L, 46L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, or (in the FR of the variable domain of the heavy chain) 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

In preferred embodiments, the non-CDR residue substituted at the consensus FR site is the residue found at the corresponding location of the non-human antibody.

Optionally, this just-recited embodiment comprises the additional steps of following the method steps appearing at the beginning of this summary and determining whether a particular amino acid residue can

9

10

15

20

а.

b.

25

30

reasonably be expected to have undesirable effects.

This invention also relates to a humanized antibody comprising the CDR sequence of an import, non-human antibody and the FR sequence of a human antibody, wherein an amino acid residue within the human FR sequence located at any one of the sites 4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H has been substituted by another residue. In preferred embodiments, the residue substituted at the human FR site is the residue found at the corresponding location of the non-human antibody from which the non-human CDR was obtained. In other embodiments, no human FR residue other than those set forth in this group has been substituted.

This invention also encompasses specific humanized antibody variable domains, and isolated polypeptides having homology with the following sequences.

1. SEQ. ID NO. 1, which is the light chain variable domain of a humanized version of muMAb4D5:

DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAP KLLIYSASFLESGVPSRFSGSRSGTDFTLTISSLQPEDFATYYCQQHY TTPPTFGQGTKVEIKRT

2. SEQ. ID NO. 2, which is the heavy chain variable domain of a humanized version of muMAb4D5):

EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLE WVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAEDT AVYYCSRWGGDGFYAMDVWGQGTLVTVSS

In another aspect, this invention provides a consensus human antibody variable domain amino acid sequence for use in the preparation of humanized antibodies, methods for obtaining, using, and storing a computer representation of such a consensus sequence, and computers comprising the

10

10

15

25

20

sequence data of such a sequence. In one embodiment, the following consensus human antibody variable domain amino acid sequences are provided:

SEQ. ID NO. 3 (light chain):

DIQMTQSPSSLSASVGDRVTITCRASQDVSSYLAWYQQKPGKAPK LLIYAASSLESGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYN SLPYTFGQGTKVEIKRT, and

SEQ. ID NO. 4 (heavy chain): EVQLVESGGGLVQPGGSLRLSCAASGFTFSDYAMSWVRQAPGKG LEWVAVISENGGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAE DTAVYYCSRWGGDGFYAMDVWGQGTLVTVSS

Brief Description of the Drawings

FIGURE 1A shows the comparison of the V_L domain amino acid residues of muMAb4D5, huMAb4D5, and a consensus human sequence (Fig. 1A, SEQ.ID NO. 5, SEQ. ID NO. 1 and SEQ. ID NO. 3, respectively). FIGURE 1B shows the comparison between the V_H domain amino acid residues of the muMAb4d5, huMAb4D5, and a consensus human sequence (Fig. 1B, SEQ. ID NO. 6, SEQ. ID NO. 2 and SEQ. ID NO. 4, respectively). Both Figs 1A and 1B use the generally accepted numbering scheme from Kabat, E. A., *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD (1987)). In both Fig. 1A and Fig. 1B, the CDR residues determined according to a standard sequence definition (as in Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)) are indicated by the first underlining beneath the sequences, and the CDR residues determined according to a structural definition (as in Chothia, C. & Lesk, A. M., *J. Mol. Biol.* 196:901-917 (1987)) are indicated by the second, lower underlines.

10

15



20

30

sequences

The mismatches between genes are shown by the vertical lines.

FIGURE 2 shows a scheme for humanization of muMAb4D5 $\rm V_L$ and $\rm V_H$ by gene conversion mutagenesis.

FIGURE 3 shows the inhibition of SK-BR-3 proliferation by MAb4D5 variants. Relative cell proliferation was determined as described (Hudziak, R. M. *et al.*, *Molec. Cell. Biol.* 9:1165-1172 (1989)) and data (average of triplicate determinations) are presented as a percentage of results with untreated cultures for muMAb4D5 (I), huMAb4D5-8 (n) and huMAb4D5-1 (I).

FIGURE 4 shows a stereo view of *a*-carbon tracing for model of huMAb4D5-8 V_L and V_H. The CDR residues (Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)) are shown in bold and side chains of V_H residues A71, T73, A78, S93, Y102 and V_L residues Y55 plus R66 (see Table 1) are shown.

Detailed Description of the Invention

Definitions

In general, the following words or phrases have the indicated definitions when used in the description, examples, and claims:

The murine monoclonal antibody known as muMAb4D5 (Fendly, B. M. *et al.*, *Cancer Res.* 50:1550-1558 (1990)) is directed against the extracellular domain (ECD) of p185^{HER2}. The muMAb4D5 and its uses are described in copending U.S. patent applications 07/143,912 and 07/147,461, and in corresponding PCT application WO 89/06692 published 27 July 1989. This murine antibody was deposited with the ATCC and designated ATCC CRL 10463. In this description and claims, the terms muMAb4D5, chMAb4D5 and huMAb4D5 represent murine, chimerized and humanized versions of the monoclonal antibody 4D5, respectively.

A humanized antibody for the purposes herein is an immunoglobulin amino acid sequence variant or fragment thereof which is capable of binding to a predetermined antigen and which comprises a FR region having

12

18

10

15

20

25

substantially the amino acid sequence of a human immunoglobulin and a CDR having substantially the amino acid sequence of a non-human immunoglobulin.

In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains (Fab) in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Ordinarily, the antibody will contain both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include the CH1, hinge, CH2, CH3, and CH4 regions of the heavy chain.

The humanized antibody will be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG3 and IgG4. Usually the constant domain is a complement fixing constant domain where it is desired that the humanized antibody exhibit cytotoxic activity, and the class is typically IgG_1 . Where such cytotoxic activity is not desirable, the constant domain may be of the IgG_2 class. The humanized antibody may comprise sequences from more than one class or isotype, and selecting particular constant domains to optimize desired effector functions is within the ordinary skill in the art.

The FR and CDR regions of the humanized antibody need not correspond precisely to the parental sequences, e.g., the import CDR or the consensus FR may be mutagenized by substitution, insertion or deletion of a residue so that the CDR or FR residue at that site does not correspond to either the consensus or the import antibody. Such mutations, however, will not be extensive. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental FR and CDR sequences, more often 90%, and most preferably greater than 95%.

In general, humanized antibodies prepared by the method of this invention are produced by a process of analysis of the parental sequences

10

15

5

25

20

30

and various conceptual humanized products using three dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen.

Residues that influence antigen binding are defined to be residues that are substantially responsible for the antigen affinity or antigen specificity of a candidate immunoglobulin, in a positive or a negative sense. The object here is to select FR residues from the consensus and import sequence so that the desired immunoglobulin characteristic is achieved. Such desired characteristics include increases in affinity and greater specificity for the target antigen, although it is conceivable that in some circumstances the opposite effects might be desired. In general, the CDR residues are directly and most substantially involved in influencing antigen binding (although not all CDR residues are so involved and therefore need not be substituted into the consensus sequence). However, FR residues also have a significant effect and can exert their influence in at least three ways: They may noncovalently directly bind to antigen, they may interact with CDR residues and they may affect the interface between the heavy and light chains.

A residue that noncovalently directly binds to antigen is one that, by three dimensional analysis, is reasonably expected to noncovalently directly bind to antigen. Typically, it is necessary to impute the position of antigen from the spatial location of neighboring CDRs and the dimensions and structure of the target antigen. In general, only those humanized antibody residues that are capable of forming salt bridges, hydrogen bonds, or hydrophobic interactions are likely to be involved in non-covalent antigen binding, however residues which are separated spatially by 3.2 Angstroms or less may also non-covalently interact. Such residues typically are the

10

15

20

7

25

30

 $\mathbf{20}$

relatively larger amino acids, such as tyrosine, arginine, and lysine. Antigenbinding FR residues also typically will have side chains that are oriented into an envelope surrounding the solvent oriented face of a CDR which extends about 7 Angstroms into the solvent from the CDR domain and about 7 Angstroms on either side of the CDR domain, again as visualized by three dimensional modeling.

5

10

15

20

25

30

A residue that interacts with a CDR generally is a residue that either affects the conformation of the CDR polypeptide backbone or forms a noncovalent bond with a CDR residue side chain. Conformation-affecting residues ordinarily are those that change the spatial position of any CDR backbone atom (N, Ca, C, O, C β) by more than about 0.2 Angstroms. Backbone atoms of CDR sequences are displaced for example by residues that interrupt or modify organized structures such as beta sheets, helices or loops. Residues that can exert a profound affect on the conformation of neighboring sequences include proline and glycine, both of which are capable of introducing bends into the backbone. Other residues that can displace backbone atoms are those that are capable of participating in salt bridges and hydrogen bonds.

A residue that interacts with a CDR side chain is one that is reasonably expected to form a noncovalent bond with a CDR side chain, generally either a salt bridge or hydrogen bond. Such residues are identified by three dimensional positioning of their side chains. A salt or ion bridge could be expected to form between two side chains positioned within about 2.5 - 3.2 Angstroms of one another that bear opposite charges, for example a lysinyl and a glutarryl pairing. A hydrogen bond could be expected to form between the side chains of residue pairs such as seryl or threonyl with aspartyl or glutarryl (or other hydrogen accepting residues). Such pairings are well known in the protein chemistry art and will be apparent to the artisan upon three dimensional modeling of the candidate immunoglobulin.

Immunoglobulin residues that affect the interface between heavy and light chain variable regions ("the $V_L - V_H$ interface") are those that affect the proximity or orientation of the two chains with respect to one another.

15

Ŏ

Certain residues involved in interchain interactions are already known and include V_L residues 34, 36, 38, 44, 46, 87, 89, 91, 96, and 98 and V_H residues 35, 37, 39, 45, 47, 91, 93, 95, 100, and 103 (utilizing the nomenclature set forth in Kabat *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)). Additional residues are newly identified by the inventors herein, and include 43L, 85L, 43H and 60H. While these residues are indicated for IgG only, they are applicable across species. In the practice of this invention, import antibody residues that are reasonably expected to be involved in interchain interactions are selected for substitution into the consensus human sequence. It is believed that heretofore no humanized antibody has been prepared with an intrachain affecting residue selected from an import antibody sequence.

Since it is not entirely possible to predict in advance what the exact impact of a given substitution will be it may be necessary to make the substitution and assay the candidate antibody for the desired characteristic. These steps, however, are *per se* routine and well within the ordinary skill of the art.

CDR and FR residues are determined according to a standard sequence definition (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda MD (1987), hereby specifically incorporated by reference), and a structural definition (as in Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987), hereby specifically incorporated by reference). Where these two methods result in slightly different identifications of a CDR, the structural definition is preferred, but the residues identified by the alternate method are considered important FR residues for determination of which framework residues to import into a consensus sequence.

The terms "consensus sequence" and "consensus antibody" as used herein refers to an amino acid sequence which comprises the most frequently occurring amino acid residues at each location in all human immunoglobulins of any particular subclass. In preferred embodiments, the

16

10

15

20

25

30

consensus human variable domain sequences are derived from the most abundant subclasses in the sequence compilation of Kabat *et al.*, Sequences, of Proteins of Immunological Interest, National Institutes of Health, Bethesda MD (1987), namely $V_L \kappa$ subgroup I and V_H group III. In such preferred embodiments, the V_L consensus domain has the amino acid sequence: DIQMTQSPSSLSASVGDRVTITCRASQDVSSYLAWYQQKPGKAPKLLIYAAS SLESGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYNSLPYTFGQGTKVEI KRT (SEQ. ID NO. 3);

the V_H consensus domain has the amino acid sequence: EVQLVESGGGLVQPGGSLRLSCAASGFTFSDYAMSWVRQAPGKGLEWVAVI SENGGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGD GFYAMDVWGQGTLVTVSS (SEQ. ID NO. 4).

While not wishing to be limited to any particular theories, it may be that these preferred embodiments are less likely to be immunogenic in an individual than less abundant subclasses. However, in other embodiments, the consensus sequence is derived from human constant domains, or from other subclasses of human immunoglobulin variable domains.

Identity or homology with respect to a specified amino acid sequence of this invention is defined herein as the percentage of amino acid residues in a candidate sequence that are identical with the specified residues, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology, and not considering any conservative substitutions as part of the sequence identity. None of Nterminal, C-terminal or internal extensions, deletions, or insertions into the specified sequence shall be construed as affecting homology. All sequence alignments called for in this invention are such maximal homology alignments. fruction of

"Non-homologous" import antibody residues are those residues which are not identical to the amino acid residue at the analogous or corresponding location in a consensus sequence, after the import and consensus sequences are aligned.

The term "computer representation" refers to information which is

10

15

5

25

20

ر**30**

17

in a form that can be manipulated by a computer. The act of storing a computer representation refers to the act of placing the information in a form suitable for manipulation by a computer.

This invention is also directed to novel polypeptides, and in certain aspects, isolated novel humanized anti-p185^{HER2} antibodies are provided. These novel anti-p185^{HER2} antibodies are sometimes collectively referred to herein as huMAb4D5, and also sometimes as the light or heavy chain variable domains of huMAb4D5, and are defined herein to be any polypeptide sequence which possesses a biological property of a polypeptide comprising the following polypeptide sequence:

DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAP KLLIYSASFLESGVPSRFSGSRSGTDFTLTISSLQPEDFATYYCQQHY TTPPTFGQGTKVEIKRT (SEQ. ID NO. 1, which is the light chain variable domain of huMAb4D5); or

EVOLVESGGGLVOPGGSLRLSCAASGFNIKDTYIHWVROAPGKGLE WVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLOMNSLRAEDT AVYYCSRWGGDGFYAMDVWGOGTLVTVSS (SEQ. ID NO. 2, which is the heavy chain variable domain of huMAb4D5).

"Biological property" for the purposes herein means an *in vivo* effector or antigenic function or activity that is directly or indirectly performed by huMAb4D5 (whether in its native or denatured conformation). Effector functions include receptor binding, any enzyme activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any mitogenic or angiogenic activity, any cytotoxic activity, any activity in promoting or inhibiting adhesion of cells to extracellular matrix or cell surface molecules, or any structural role. However, effector functions do not include possession of an epitope or antigenic site that is capable of cross-reacting with antibodies raised against the polypeptide sequence of huMAb4D5.

10

15

20



30

Biologically active huMAb4D5 is defined herein as a polypeptide that shares an effector function of huMAb4D5 and which may (but need not) in addition possess an antigenic function. A principal known effect or function of huMAb4D5 is its ability to bind to p185^{HER2}.

Antigenically active huMAb4D5 is defined as a polypeptide that possesses an antigenic function of huMAb4D5 and which may (but need not) in addition possess an effector function.

In preferred embodiments, antigenically active huMAb4D5 is a polypeptide that binds with an affinity of at least about 10-9 l/mole to an antibody capable of binding huMAb4D5. Ordinarily the polypeptide binds with an affinity of at least about 10-8 l/mole. Isolated antibody capable of binding huMAb4D5 is an antibody which is identified and separated from a component of the natural environment in which it may be present. Most preferably, antigenically active huMAb4D5 is a polypeptide that binds to an antibody capable of binding huMAb4D5 in its native conformation. HuMAb4D5 in its native conformation is huMAb4D5 as recovered according to the methods described in Example 1 below, which has not been denatured by chaotropic agents, heat or other treatment that substantially modifies the three dimensional structure of huMAb4D5 as determined for example by migration on nonreducing, nondenaturing sizing gels. Antibody used in this determination is rabbit polyclonal antibody raised by formulating native huMAb4D5 in Freund's complete adjuvant, subcutaneously injecting the formulation, and boosting the immune response by intraperitoneal injection of the formulation until the titer of anti-huMAb4D5 antibody plateaus.

Ordinarily, biologically or antigenically active huMAb4D5 will have an amino acid sequence having at least 75% amino acid sequence identity with the huMAb4D5 amino acid sequence, more preferably at least 80%, more preferably at least 90%, and most preferably at least 95%. Identity or homology with respect to this sequence is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the huMAb4D5 residues, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology, and not considering

10

5

15

20

25

19

any conservative substitutions as part of the sequence identity. None of Nterminal, C-terminal or internal extensions, deletions, or insertions into the huMAb4D5 sequence shall be construed as affecting homology.

Thus, the biologically active and antigenically active huMAb4D5 polypeptides that are the subject of certain embodiments of this invention include the sequence of the entire translated nucleotide sequence of huMAb4D5; mature huMAb4D5; fragments thereof having a consecutive sequence of at least 5, 10, 15, 20, 25, 30 or 40 amino acid residues from huMAb4D5; amino acid sequence variants of huMAb4D5 wherein an amino acid residue has been inserted N- or C-terminal to, or within, huMAb4D5 or its fragment as defined above; amino acid sequence variants of huMAb4D5 or its fragment as defined above wherein an amino acid residue of huMAb4D5 or its fragment as defined above has been substituted by another residue, including predetermined mutations by, e.g., site-directed or PCR mutagenesis; derivatives of huMAb4D5 or its fragments as defined above wherein huMAb4D5 or its fragments have been covalent modified, by substitution, chemical, enzymatic, or other appropriate means, with a moiety other than a naturally occurring amino acid; and glycosylation variants of huMAb4D5 (insertion of a glycosylation site of deletion of any glycosylation site by deletion, insertion or substitution of suitable residues). Such fragments and variants exclude any polypeptide heretofore identified, including muMAb4D5 or any known polypeptide fragment, which are anticipatory order 35 U.S.C.102 as well as polypeptides obvious thereover under 35 U.S.C. 103.

"Isolated" huMAb4D5 means huMAb4D5 which has been identified and separated and/or recovered from a component of its natural cell culture /environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for huMAb4D5, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, huMAb4D5 will be purified (1) to greater than 95% by weight of protein as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a

.20

26

10

15

20

25

degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated huMAb4D5 includes huMAb4D5 in situ within recombinant cells since at least one component of the huMAb4D5 natural environment will not be present. Ordinarily, however, isolated huMAb4D5 will be prepared by at least one purification step.

In accordance with this invention, huMAb4D5 nucleic acid is RNA or DNA containing greater than ten bases that encodes a biologically or antigenically active huMAb4D5, is complementary to nucleic acid sequence encoding such huMAb4D5, or hybridizes to nucleic acid sequence encoding such huMAb4D5 and remains stably bound to it under stringent conditions.

Preferably, the huMAb4D5 nucleic acid encodes a polypeptide sharing at least 75% sequence identity, more preferably at least 80%, still more preferably at least 85%, even more preferably at 90%, and most preferably 95%, with the huMAb4D5 amino acid sequence. Preferably, a nucleic acid molecule that hybridizes to the huMAb4D5 nucleic acid contains at least 20, more preferably 40, and most preferably 90 bases. Such hybridizing or complementary nucleic acid, however, is further defined as being novel under 35 U.S.C. 102 and unobvious under 35 U.S.C. 103 over any prior art nucleic acid.

Stringent conditions are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate/0/1% NaDodSO₄ at 50° C; (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0/1% Ficoll/0/1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42° C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 g/ml), 0.1% SDS, and 10% dextran sulfate at 42 C, with washes at 42 C in 0.2 x SSC and 0.1% SDS.

10

5

15

20

25

30

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, a ribosome binding site, and possibly, other as yet poorly understood sequences. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

5

10

15

20

25

30

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

An "exogenous" element is defined herein to mean nucleic acid sequence that is foreign to the cell, or homologous to the cell but in a position within the host cell nucleic acid in which the element is ordinarily not found.

As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are

22

intended, it will be clear from the context.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are commercially available, are publicly available on an unrestricted basis, or can be constructed from such available plasmids in accord with published procedures. In addition, other equivalent plasmids are known in the art and will be apparent to the ordinary artisan.

"Restriction Enzyme Digestion" of DNA refers to catalytic cleavage of the DNA with an enzyme that acts only at certain locations in the DNA. Such enzymes are called restriction endonucleases, and the sites for which each is specific is called a restriction site. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors, and other requirements as established by the enzyme suppliers are Restriction enzymes commonly are designated by abbreviations used. composed of a capital letter followed by other letters representing the microorganism from which each restriction enzyme originally was obtained and then a number designating the particular enzyme. In general, about 1 μ g of plasmid or DNA fragment is used with about 1-2 units of enzyme in about 20 Appropriate buffers and substrate amounts for μ of buffer solution. particular restriction enzymes are specified by the manufacturer. Incubation of about 1 hour at 37°C is ordinarily used, but may vary in accordance with the supplier's instructions. After incubation, protein or polypeptide is removed by extraction with phenol and chloroform, and the digested nucleic acid is recovered from the aqueous fraction by precipitation with ethanol. Digestion with a restriction enzyme may be followed with bacterial alkaline phosphatase hydrolysis of the terminal 5' phosphates to prevent the two restriction cleaved ends of a DNA fragment from "circularizing" or forming a closed loop that would impede insertion of another DNA fragment at the restriction site. Unless otherwise stated, digestion of plasmids is not followed by 5' terminal dephosphorylation. Procedures and reagents for dephosphorylation are conventional as described in sections 1.56-1.61 of Sambrook et al. (Molecular Cloning: A Laboratory Manual New York: Cold

10

15

20

25

30

23

Spring Harbor Laboratory Press, 1989).

"Recovery" or "isolation" of a given fragment of DNA from a restriction digest means separation of the digest on polyacrylamide or agarose gel by electrophoresis, identification of the fragment of interest by comparison of its mobility versus that of marker DNA fragments of known molecular weight, removal of the gel section containing the desired fragment, and separation of the gel from DNA. This procedure is known generally. For example, see Lawn *et al.*, <u>Nucleic Acids Res.</u>, <u>9</u>: 6103-6114 (1981), and Goeddel *et al.*, <u>Nucleic Acids Res.</u> <u>8</u>: 4057 (1980).

"Southern blot analysis" is a method by which the presence of DNA sequences in a restriction endonuclease digest of DNA or DNA-containing composition is confirmed by hybridization to a known, labeled oligonucleotide or DNA fragment. Southern analysis typically comprises electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer of the DNA to nitrocellulose, nylon, or another suitable membrane supports for analysis with a radiolabeled, biotinylated or enzyme-labeled probe as described in sections 9.37-9.52 of Sambrook et al, *supra*.

"Northern analysis" is a method used to identify RNA sequences that hybridize to a known probe such as an oligonucleotide, DNA fragment, cDNA or fragment thereof, or RNA fragment. The probe is labeled with a radioisotope such as 32-P, or by biotinylation, or with an enzyme. The RNA to be analyzed is usually electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe, using standard techniques well known in the art such as those described in sections 7.39-7.52 of Sambrook et al., *supra*.

"Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments. To ligate the DNA fragments together, the ends of the DNA fragments must be compatible with each other. In some cases, the ends will be directly compatible after endonuclease digestion. However, it may be necessary to first convert the staggered ends

24

15

20



commonly produced after endonuclease digestion to blunt ends to make them compatible for ligation. To blunt the ends, the DNA is treated in a suitable buffer for at least 15 minutes at 15°C with about 10 units of the Klenow fragment of DNA polymerase I or T4 DNA polymerase in the presence of the four deoxyribonucleotide triphosphates. The DNA is then purified by phenolchloroform extraction and ethanol precipitation. The DNA fragments that are to be ligated together are put in solution in about equimolar amounts. The solution will also contain ATP, ligase buffer, and a ligase such as T4 DNA ligase at about 10 units per 0.5 μ g of DNA. If the DNA is to be ligated into a vector, the vector is first linearized by digestion with the appropriate restriction endonuclease(s). The linearized fragment is then treated with bacterial alkaline phosphatase, or calf intestinal phosphatase to prevent selfligation during the ligation step.

"Preparation" of DNA from cells means isolating the plasmid DNA from a culture of the host cells. Commonly used methods for DNA preparation are the large and small scale plasmid preparations described in sections 1.25-1.33 of Sambrook *et al., supra*. After preparation of the DNA, it can be purified by methods well known in the art such as that described in section 1.40 of Sambrook *et al., supra*.

"Oligonucleotides" are short-length, single- or double-stranded polydeoxynucleotides that are chemically synthesized by known methods (such as phosphotriester, phosphite, or phosphoramidite chemistry, using solid phase techniques such as described in EP 266,032 published 4 May 1988, or via deoxynucleoside H-phosphonate intermediates as described by Froehler *et al.*, <u>Nucl. Acids Res.</u>, <u>14</u>: 5399-5407 [1986]). They are then purified on polyacrylamide gels.

The technique of "polymerase chain reaction," or "PCR," as used herein generally refers to a procedure wherein minute amounts of a specific piece of nucleic acid, RNA and/or DNA, are amplified as described in U.S. Pat. No. 4,683,195 issued 28 July 1987. Generally, sequence information from the ends of the region of interest or beyond needs to be available, such that oligonucleotide primers can be designed; these primers will be identical

10

•

20

15

30

25

25

or similar in sequence to opposite strands of the template to be amplified. The 5' terminal nucleotides of the two primers may coincide with the ends of the amplified material. PCR can be used to amplify specific RNA sequences, specific DNA sequences from total genomic DNA, and cDNA transcribed from total cellular RNA, bacteriophage or plasmid sequences, etc. See generally Mullis *et al.*, <u>Cold Spring Harbor Symp. Quant. Biol.</u>, <u>51</u>: 263 (1987); Erlich, ed., <u>PCR Technology</u>, (Stockton Press, NY, 1989). As used herein, PCR is considered to be one, but not the only, example of a nucleic acid polymerase reaction method for amplifying a nucleic acid test sample, comprising the use of a known nucleic acid (DNA or RNA) as a primer and utilizes a nucleic acid polymerase to amplify or generate a specific piece of nucleic acid or to amplify or generate a specific piece of nucleic acid which is complementary to a particular nucleic acid.

Suitable Methods for Practicing the Invention

Some aspects of this invention include obtaining an import, nonhuman antibody variable domain, humanizing the antibody sequence, and producing the humanized antibody. Methods for determining a desired humanized antibody sequence and for humanizing an antibody gene sequence are described below. A particularly preferred method of gene conversion from a non-human or consensus sequence into a humanized nucleic acid sequence is described in Example 1. Additionally, methods are given for obtaining and producing antibodies generally, which apply equally to native non-human antibodies as well as to humanized antibodies.

Generally, the antibodies and antibody variable domains of this invention are conventionally prepared in recombinant cell culture, as described in more detail below. Recombinant synthesis is preferred for reasons of safety and economy, but it is known to prepare peptides by chemical synthesis and to purify them from natural sources; such preparations are included within the definition of antibodies herein.

10 --

15

20

30

Molecular Modeling

Where it is desired to prepare molecular models for the antibodies of this invention, one may utilize any of the commercially available modeling programs described in the literature cited in the Background above.

Generally, models for a particular antibody domains, for example non-human, import antibody variable V_H and V_L domains, are constructed separately from consensus coordinates based upon FAb structures which have similar sequences. Models of consensus human antibody sequences are similarly created.

For example, in modeling the muMAb4d5, the models were constructed based upon seven Fab structures from the Brookhaven protein data bank (entries 1FB4, 2RHE, 2MCP, 3FAB, 1FBJ, 2HFL and 1REI). The Fab fragment KOL (Marquart, M. *et al.*, *J. Mol. Biol.* 141:369-391 (1980)) was first chosen as a template for V_L and V_H domains and additional structures were then superimposed upon this structure using their main chain atom coordinates (INSIGHT program, Biosym Technologies). Similar programs and techniques are utilized for modeling the desired antibody.

The distance from the template Ca to the analogous Ca in each of the superimposed structures is calculated for each residue position. Generally, if all (or nearly all) Ca-Ca distances for a given residue are ≤ 1 Å, then that position is included in the consensus structure. In some cases the β -sheet framework residues will satisfy these criteria whereas the CDR loops may not. For each of these selected residues the average coordinates for individual N, Ca, C, O and C β atoms are calculated and then corrected for resultant deviations from non-standard bond geometry by 50 cycles of energy minimization using a commercially available program such as the DISCOVER program (Biosym Technologies) with the AMBER forcefield (Weiner, S. J. *et al.*, *J. Amer. Chem. Soc.* 106:765-784 (1984)), and the Ca coordinates are fixed. The side chains of highly conserved residues, such as the disulfide-bridged cysteine residues, are then incorporated into the resultant consensus structure. Next the sequences of the particular antibody V_L and V_H domains are incorporated starting with the CDR residues and

.

15

10

20

25

30

33 ~

using the tabulations of CDR conformations from Chothia et al. (Chothia, C. et al., Nature 342:877-883 (1989)) as a guide. Side-chain conformations are chosen on the basis of Fab crystal structures, rotamer libraries (Ponder, J. W. & Richards, F. M., J. Mol. Biol. 193:775-791 (1987)) and packing considerations. Since V_{H} -CDR3 typically cannot be assigned a definite backbone conformation from these criteria, models may be created from a search of similar sized loops using the INSIGHT program, derived using packing and solvent exposure considerations, or created using other routine and commercially available techniques. It is preferable to subject the model to 5000 cycles of energy minimization.

Methods for Obtaining a Humanized Antibody Sequence

In humanizing muMAb4D5, consensus human sequences are first derived, and then a molecular model is generated for these sequences using the methods described above. In certain embodiments of this invention, the consensus human sequences are derived from the most abundant subclasses in the sequence compilation of Kabat et al. (Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)), namely $V_L \kappa$ subgroup 1 and V_H group III, and have the sequences indicated in the definitions above.

While these steps may be taken in different order, typically a structure for the candidate humanized antibody is created by transferring the CDRs from the non-human, import sequence into the consensus human structure. The humanized antibody may contain human replacements of the non-human import residues at positions within CDRs as defined by sequence variability (Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)) or as defined by structural variability (Chothia, C. & Lesk, A. M., *J. Mol. Biol.* 196:901-917 (1987)). For example, huMAb4D5 contains human replacements of the muMAb4D5 residues at three positions within CDRs as defined by sequence variability (Kabat, E. A. *et al.*, Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)) or as defined by sequence variability (Kabat, E. A. *et al.*, Sequences of Proteins within CDRs as defined by sequence variability (Kabat, E. A. *et al.*, Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987))

10

15

20

25

30

but not as defined by structural variability (Chothia, C. & Lesk, A. M., *J. Mol. Biol.* **196**:901-917 (1987)): V_L -CDR1 K24R, V_L -CDR2 R54L and V_L -CDR2 T56S.

Differences between the non-human import and the human consensus framework residues are individually investigated to determine their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is desirably performed through modeling, by examination of the characteristics of the amino acids at particular locations, or determined experimentally through evaluating the effects of substitution or mutagenesis of particular amino acids.

In certain preferred embodiments of this invention, a humanized antibody is made comprising amino acid sequence of an import, non-human antibody and a human antibody, utilizing the steps of:

a. obtaining the amino acid sequences of at least a portion
 of an import antibody variable domain and of a consensus
 human variable domain;

identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human variable domain sequences;

substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;

identifying import antibody FR residues in the aligned FR sequences that are non-homologous to the corresponding consensus antibody residues;

determining if the non-homologous import amino acid residue is reasonably expected to have at least one of the following effects:

1. non-covalently binds antigen directly,

interacts with a CDR; or

29

10

15

20

b.

c.

d.

e.

f.

2.

25

30

3. participates in the $V_L - V_H$ interface; and for any non-homologous import antibody amino acid residue which is reasonably expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence.

Optionally, one determines if any non-homologous residues identified in step (e) are exposed on the surface of the domain or buried within it, and if the residue is exposed but has none of the effects identified in step (f), one may retain the consensus residue.

Additionally, in certain embodiments the corresponding consensus antibody residues identified in step (e) above are selected from the group consisting of 4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H (utilizing the numbering system set forth in Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)).

In preferred embodiments, the method of this invention comprises the additional steps of searching either or both of the import, non-human and the consensus variable domain sequences for glycosylation sites, determining if the glycosylation is reasonably expected to be important for the desired antigen binding and biological activity of the antibody (i.e., determining if the glycosylation site binds to antigen or changes a side chain of an amino acid residue that binds to antigen, or if the glycosylation enhances or weakens antigen binding, or is important for maintaining antibody affinity). If the import sequence bears the glycosylation site, it is preferred to substitute that site for the corresponding residues in the consensus human sequence if the glycosylation site is reasonably expected to be important. If only the consensus sequence, and not the import, bears the glycosylation site, it is preferred to eliminate that glycosylation site or substitute therefor the corresponding amino acid residues from the import sequence.

g.

20

10

25

30

30

Another preferred embodiment of the methods of this invention comprises aligning import antibody and the consensus antibody FR sequences, identifying import antibody FR residues which are nonhomologous with the aligned consensus FR sequence, and for each such nonhomologous import antibody FR residue, determining if the corresponding consensus antibody residue represents a residue which is highly conserved across all species at that site, and if it is so conserved, preparing a humanized antibody which comprises the consensus antibody amino acid residue at that site.

In certain alternate embodiments, one need not utilize the modeling and evaluation steps described above, and may instead proceed with the steps of obtaining the amino acid sequence of at least a portion of an import, non-human antibody variable domain having a CDR and a FR, obtaining the amino acid sequence of at least a portion of a consensus human antibody variable domain having a CDR and a FR, substituting the non-human CDR for the human CDR in the consensus human antibody variable domain, and then substituting an amino acid residue for the consensus amino acid residue at at least one of the following sites:

> a. (in the FR of the variable domain of the light chain) 4L, 35L, 36L, 38L, 43L, 44L, 58L, 46L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, or
> b. (in the FR of the variable domain of the heavy chain) 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

Preferably, the non-CDR residue substituted at the consensus FR site is the residue found at the corresponding location of the non-human antibody. If desired, one may utilize the other method steps described above for determining whether a particular amino acid residue can reasonably be expected to have undesirable effects, and remedying those effects.

If after making a humanized antibody according to the steps above and testing its activity one is not satisfied with the humanized antibody, one

10

20

15

30

25

31

preferably reexamines the potential effects of the amino acids at the specific locations recited above. Additionally, it is desirable to reinvestigate any buried residues which are reasonably expected to affect the $V_L - V_H$ interface but may not directly affect CDR conformation. It is also desirable to reevaluate the humanized antibody utilizing the steps of the methods claimed herein.

In certain embodiments of this invention, amino acid residues in the consensus human sequence are substituted for by other amino acid residues. In preferred embodiments, residues from a particular non-human import sequence are substituted, however there are circumstances where it is desired to evaluate the effects of other amino acids. For example, if after making a humanized antibody according to the steps above and testing its activity one is not satisfied with the humanized antibody, one may compare the sequences of other classes or subgroups of human antibodies, or classes or subgroups of antibodies from the particular non-human species, and determine which other amino acid side chains and amino acid residues are found at particular locations and substituting such other residues.

Antibodies

Certain aspects of this invention are directed to natural antibodies and to monoclonal antibodies, as illustrated in the Examples below and by antibody hybridomas deposited with the ATCC (as described below). Thus, the references throughout this description to the use of monoclonal antibodies are intended to include the use of natural or native antibodies as well as humanized and chimeric antibodies. As used herein, the term "antibody" includes the antibody variable domain and other separable antibody domains unless specifically excluded.

In accordance with certain aspects of this invention, antibodies to be humanized (import antibodies) are isolated from continuous hybrid cell lines formed by the fusion of antigen-primed immune lymphocytes with myeloma cells.

In certain embodiments, the antibodies of this invention are

20

15

25

30

32

obtained by routine screening. Polyclonal antibodies to an antigen generally are raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the antigen and an adjuvant. It may be useful to conjugate the antigen or a fragment containing the target amino acid sequence to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCI₂, or R¹N = C = NR, where R and R¹ are different alkyl groups.

The route and schedule of the host animal or cultured antibodyproducing cells therefrom are generally in keeping with established and conventional techniques for antibody stimulation and production. While mice are frequently employed as the test model, it is contemplated that any mammalian subject including human subjects or antibody-producing cells obtained therefrom can be manipulated according to the processes of this invention to serve as the basis for production of mammalian, including human, hybrid cell lines.

Animals are typically immunized against the immunogenic conjugates or derivatives by combining 1 mg or 1 μ g of conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of conjugate in Freund's complete adjuvant (or other suitable adjuvant) by subcutaneous injection at multiple sites. 7 to 14 days later animals are bled and the serum is assayed for antigen titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking agent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are used to enhance the immune response.

10

15

5

20[°]

30

25

39

After immunization, monoclonal antibodies are prepared by recovering immune lymphoid cells--typically spleen cells or lymphocytes from lymph node tissue--from immunized animals and immortalizing the cells in conventional fashion, e.g. by fusion with myeloma cells or by Epstein-Barr (EB)-virus transformation and screening for clones expressing the desired antibody. The hybridoma technique described originally by Kohler and Milstein, *Eur. J. Immunol.* 6:511 (1976) has been widely applied to produce hybrid cell lines that secrete high levels of monoclonal antibodies against many specific antigens.

It is possible to fuse cells of one species with another. However, it is preferable that the source of the immunized antibody producing cells and the myeloma be from the same species.

The hybrid cell lines can be maintained in culture in vitro in cell culture media. The cell lines of this invention can be selected and/or maintained in a composition comprising the continuous cell line in hypoxanthine-aminopterin thymidine (HAT) medium. In fact, once the hybridoma cell line is established, it can be maintained on a variety of nutritionally adequate media. Moreover, the hybrid cell lines can be stored and preserved in any number of conventional ways, including freezing and storage under liquid nitrogen. Frozen cell lines can be revived and cultured indefinitely with resumed synthesis and secretion of monoclonal antibody. The secreted antibody is recovered from tissue culture supernatant by conventional methods such as precipitation, Ion exchange chromatography, affinity chromatography, or the like. The antibodies described herein are also recovered from hybridoma cell cultures by conventional methods for purification of IgG or IgM as the case may be that heretofore have been used to purify these immunoglobulins from pooled plasma, e.g. ethanol or polyethylene glycol precipitation procedures. The purified antibodies are sterile filtered, and optionally are conjugated to a detectable marker such as an enzyme or spin label for use in diagnostic assays of the antigen in test samples.

While routinely rodent monoclonal antibodies are used as the source

10

15

20

5

25 ·

30

34

of the import antibody, the invention is not limited to any species. Additionally, techniques developed for the production of chimeric antibodies (Morrison *et al.*, *Proc. Natl. Acad. Sci.*, 81:6851 (1984); Neuberger *et al.*, *Nature* 312:604 (1984); Takeda *et al.*, *Nature* 314:452 (1985)) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity (such as ability to activate human complement and mediate ADCC) can be used; such antibodies are within the scope of this invention.

Techniques for creating recombinant DNA versions of the antigenbinding regions of antibody molecules (known as Fab fragments) which bypass the generation of monoclonal antibodies are encompassed within the practice of this invention. One extracts antibody-specific messenger RNA molecules from immune system cells taken from an immunized animal, transcribes these into complementary DNA (cDNA), and clones the cDNA into a bacterial expressions system. One example of such a technique suitable for the practice of this invention was developed by researchers at Scripps/Stratagene, and incorporates a proprietary bacteriophage lambda vector system which contains a leader sequence that causes the expressed Fab protein to migrate to the periplasmic space (between the bacterial cell membrane and the cell wall) or to be secreted. One can rapidly generate and screen great numbers of functional FAb fragments for those which bind the antigen. Such FAb fragments with specificity for the antigen are specifically encompassed within the term "antibody" as it is defined, discussed, and claimed herein.

Amino Acid Sequence Variants

Amino acid sequence variants of the antibodies and polypeptides of this invention (referred to in herein as the target polypeptide) are prepared by introducing appropriate nucleotide changes into the DNA encoding the target polypeptide, or by *in vitro* synthesis of the desired target polypeptide. Such variants include, for example, humanized variants of non-human antibodies, as well as deletions from, or insertions or substitutions of,

10

.

20

15

25

30

residues within particular amino acid sequences. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the target polypeptide, such as changing the number or position of glycosylation sites, altering any membrane anchoring characteristics, and/or altering the intra-cellular location of the target polypeptide by inserting, deleting, or otherwise affecting any leader sequence of the native target polypeptide.

In designing amino acid sequence variants of target polypeptides, the location of the mutation site and the nature of the mutation will depend on the target polypeptide characteristic(s) to be modified. The sites for mutation can be modified individually or in series, e.g., by (1) substituting first with conservative amino acid choices and then with more radical selections depending upon the results achieved, (2) deleting the target residue, or (3) inserting residues of the same or a different class adjacent to the located site, or combinations of options 1-3. In certain embodiments, these choices are guided by the methods for creating humanized sequences set forth abové.

10

15

20

25

30

A useful method for identification of certain residues or regions of the target polypeptide that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells (<u>Science, 244</u>: 1081-1085 [1989]). Here, a residue or group of target residues are identified (e.g., charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Those domains demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at or for the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to optimize the performance of a mutation at

36

a given site, ala scanning or random mutagenesis may be conducted at the target codon or region and the expressed target polypeptide variants are screened for the optimal combination of desired activity.

There are two principal variables in the construction of amino acid sequence variants: the location of the mutation site and the nature of the mutation. In general, the location and nature of the mutation chosen will depend upon the target polypeptide characteristic to be modified.

Amino acid sequence deletions of antibodies are generally not preferred, as maintaining the generally configuration of an antibody is believed to be necessary for its activity. Any deletions will be selected so as to preserve the structure of the target antibody.

10

15

20

25

30

Amino acid sequence insertions include amino- and/or carboxylterminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions (i.e., insertions within the target polypeptide sequence) may range generally from about 1 to 10 residues, more preferably 1 to 5, most preferably 1 to 3. Examples of terminal insertions include the target polypeptide with an N-terminal methionyl residue, an artifact of the direct expression of target polypeptide in bacterial recombinant cell culture, and fusion of a heterologous N-terminal signal sequence to the N-terminus of the target polypeptide molecule to facilitate the secretion of the mature target polypeptide from recombinant host cells. Such signal sequences generally will be obtained from, and thus homologous to, the intended host cell species. Suitable sequences include STII or Ipp for *E. coli*, alpha factor for yeast, and viral signals such as herpes aD for mammalian cells.

Other insertional variants of the target polypeptide include the fusion to the N- or C-terminus of the target polypeptide of immunogenic polypeptides, e.g., bacterial polypeptides such as beta-lactamase or an enzyme encoded by the *E. coli trp* locus, or yeast protein, and C-terminal fusions with proteins having a long half-life such as immunoglobulin constant regions (or other immunoglobulin regions), albumin, or ferritin, as described

37

in WO 89/02922 published 6 April 1989.

Another group of variants are amino acid substitution variants. These variants have at least one amino acid residue in the target polypeptide molecule removed and a different residue inserted in its place. The sites of greatest interest for substitutional mutagenesis include sites identified as the active site(s) of the target polypeptide, and sites where the amino acids found in the target polypeptide from various species are substantially different in terms of side-chain bulk, charge, and/or hydrophobicity. Other sites for substitution are described infra, considering the effect of the substitution of the antigen binding, affinity and other characteristics of a particular target antibody.

Other sites of interest are those in which particular residues of the target polypeptides obtained from various species are identical. These positions may be important for the biological activity of the target polypeptide. These sites, especially those falling within a sequence of at least three other identically conserved sites, are substituted in a relatively conservative manner. If such substitutions result in a change in biological activity, then other changes are introduced and the products screened until the desired effect is obtained.

Substantial modifications in function or immunological identity of the target polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side chain properties:

(1) hydrophobic: norleucine, met, ala, val, leu, ile;

(2) neutral hydrophilic: cys, ser, thr;

(3) acidic: asp, glu;

(4) basic: asn, gln, his, lys, arg;

(5) residues that influence chain orientation: gly, pro; and

(6) aromatic: trp, tyr, phe.

10

15

25

20

30

38

Non-conservative substitutions will entail exchanging a member of one of these classes for another. Such substituted residues may be introduced into regions of the target polypeptide that are homologous with other antibodies of the same class or subclass, or, more preferably, into the non-homologous regions of the molecule.

Any cysteine residues not involved in maintaining the proper conformation of target polypeptide also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking.

DNA encoding amino acid sequence variants of the target polypeptide is prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the target polypeptide. A particularly preferred method of gene conversion mutagenesis is described below in Example 1. These techniques may utilized target polypeptide nucleic acid (DNA or RNA), or nucleic acid complementary to the target polypeptide nucleic acid.

Oligonucleotide-mediated mutagenesis is a preferred method for preparing substitution, deletion, and insertion variants of target polypeptide DNA. This technique is well known in the art as described by Adelman *et al.*, <u>DNA, 2</u>: 183 (1983). Briefly, the target polypeptide DNA is altered by hybridizing an oligonucleotide encoding the desired mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of the target polypeptide. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the target polypeptide DNA.

Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are

10

15

20

.

30

25

completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea *et al.* (Proc. Natl. Acad. Sci. USA, 75: 5765 [1978]).

Single-stranded DNA template may also be generated by denaturing double-stranded plasmid (or other) DNA using standard techniques.

For alteration of the native DNA sequence (to generate amino acid sequence variants, for example), the oligonucleotide is hybridized to the single-stranded template under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of DNA polymerase I, is then added to synthesize the complementary strand of the template using the oligonucleotide as a primer for synthesis. A heteroduplex molecule is thus formed such that one strand of DNA encodes the mutated form of the target polypeptide, and the other strand (the original template) encodes the native, unaltered sequence of the target polypeptide. This heteroduplex molecule is then transformed into a suitable host cell, usually a prokaryote such as E. coli JM101. After the cells are grown, they are plated onto agarose plates and screened using the oligonucleotide primer radiolabeled with 32-phosphate to identify the bacterial colonies that contain the mutated DNA. The mutated region is then removed and placed in an appropriate vector for protein production, generally an expression vector of the type typically employed for transformation of an appropriate host.

The method described immediately above may be modified such that a homoduplex molecule is created wherein both strands of the plasmid The modifications are as follows: The contain the mutation(s). single-stranded oligonucleotide is annealed to the single-stranded template A mixture of three deoxyribonucleotides, as described above. deoxyriboadenosine (dATP), deoxyriboguanosine (dGTP), and (dTTP), is combined with modified deoxyribothymidine а thio-deoxyribocytosine called dCTP-(aS) (which can be obtained from is added the Amersham Corporation). This mixture to

10

15

20

25

30

46

template-oligonucleotide complex. Upon addition of DNA polymerase to this mixture, a strand of DNA identical to the template except for the mutated bases is generated. In addition, this new strand of DNA will contain dCTP-(aS) instead of dCTP, which serves to protect it from restriction endonuclease digestion.

After the template strand of the double-stranded heteroduplex is nicked with an appropriate restriction enzyme, the template strand can be digested with ExoIII nuclease or another appropriate nuclease past the region that contains the site(s) to be mutagenized. The reaction is then stopped to leave a molecule that is only partially single-stranded. A complete double-stranded DNA homoduplex is then formed using DNA polymerase in the presence of all four deoxyribonucleotide triphosphates, ATP, and DNA ligase. This homoduplex molecule can then be transformed into a suitable host cell such as *E. coli* JM101, as described above.

DNA encoding target polypeptide variants with more than one amino acid to be substituted may be generated in one of several ways. If the amino acids are located close together in the polypeptide chain, they may be mutated simultaneously using one oligonucleotide that codes for all of the desired amino acid substitutions. If, however, the amino acids are located some distance from each other (separated by more than about ten amino acids), it is more difficult to generate a single oligonucleotide that encodes all of the desired changes. Instead, one of two alternative methods may be employed.

In the first method, a separate oligonucleotide is generated for each amino acid to be substituted. The oligonucleotides are then annealed to the single-stranded template DNA simultaneously, and the second strand of DNA that is synthesized from the template will encode all of the desired amino acid substitutions.

The alternative method involves two or more rounds of mutagenesis to produce the desired mutant. The first round is as described for the single mutants: wild-type DNA is used for the template, an oligonucleotide encoding the first desired amino acid substitution(s) is annealed to this template, and

10

20

15

30

25

the heteroduplex DNA molecule is then generated. The second round of mutagenesis utilizes the mutated DNA produced in the first round of mutagenesis as the template. Thus, this template already contains one or more mutations. The oligonucleotide encoding the additional desired amino acid substitution(s) is then annealed to this template, and the resulting strand of DNA now encodes mutations from both the first and second rounds of mutagenesis. This resultant DNA can be used as a template in a third round of mutagenesis, and so on.

PCR mutagenesis is also suitable for making amino acid variants of While the following discussion refers to DNA, it is target polypeptide. understood that the technique also finds application with RNA. The PCR technique generally refers to the following procedure (see Erlich, supra, the chapter by R. Higuchi, p. 61-70): When small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs from the template sequence only at the positions where the primers differ from the template. For introduction of a mutation into a plasmid DNA, one of the primers is designed to overlap the position of the mutation and to contain the mutation; the sequence of the other primer must be identical to a stretch of sequence of the opposite strand of the plasmid, but this sequence can be located anywhere along the plasmid DNA. It is preferred, however, that the sequence of the second primer is located within 200 nucleotides from that of the first, such that in the end the entire amplified region of DNA bounded by the primers can be easily sequenced. PCR amplification using a primer pair like the one just described results in a population of DNA fragments that differ at the position of the mutation specified by the primer, and possibly at other positions, as template copying is somewhat error-prone.

If the ratio of template to product material is extremely low, the vast majority of product DNA fragments incorporate the desired mutation(s). This product material is used to replace the corresponding region in the plasmid that served as PCR template using standard DNA technology.

42

10

15

20

25

30

. 48 Mutations at separate positions can be introduced simultaneously by either using a mutant second primer, or performing a second PCR with different mutant primers and ligating the two resulting PCR fragments simultaneously to the vector fragment in a three (or more)-part ligation.

In a specific example of PCR mutagenesis, template plasmid DNA (1 μ g) is linearized by digestion with a restriction endonuclease that has a unique recognition site in the plasmid DNA outside of the region to be amplified. Of this material, 100 ng is added to a PCR mixture containing PCR buffer, which contains the four deoxynucleotide tri-phosphates and is included in the GeneAmp^Φ kits (obtained from Perkin-Elmer Cetus, Norwalk, CT and Emeryville, CA), and 25 pmole of each oligonucleotide primer, to a final volume of 50 μ l. The reaction mixture is overlayed with 35 μ l mineral oil. The reaction is denatured for 5 minutes at 100°C, placed briefly on ice, and then 1 μ l *Thermus aquaticus (Taq)* DNA polymerase (5 units/ μ l, purchased from Perkin-Elmer Cetus, Norwalk, CT and Emeryville, CA) is added below the mineral oil layer. The reaction mixture is then inserted into a DNA Thermal Cycler (purchased from Perkin-Elmer Cetus) programmed as follows: 2 min. at 55°C, then 30 sec. at 72°C, then 19 cycles of the following: 30 sec. at 94°C, 30 sec. at 55°C, and 30 sec. at 72°C.

At the end of the program, the reaction vial is removed from the thermal cycler and the aqueous phase transferred to a new vial, extracted with phenol/chloroform (50:50:vol), and ethanol precipitated, and the DNA is recovered by standard procedures. This material is subsequently subjected to the appropriate treatments for insertion into a vector.

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells *et al.* (Gene, 34: 315 [1985]). The starting material is the plasmid (or other vector) comprising the target polypeptide DNA to be mutated. The codon(s) in the target polypeptide DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at

10

15

5

25

20

30

appropriate locations in the target polypeptide DNA. After the restriction sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures. The two strands are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated target polypeptide DNA sequence.

Insertion of DNA into a Cloning Vehicle

The cDNA or genomic DNA encoding the target polypeptide is inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Many vectors are available, and selection of the appropriate vector will depend on 1) whether it is to be used for DNA amplification or for DNA expression, 2) the size of the DNA to be inserted into the vector, and 3) the host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

(a) Signal Sequence Component

In general, the signal sequence may be a component of the vector, or it may be a part of the target polypeptide DNA that is inserted into the vector.

The target polypeptides of this invention may be expressed not only directly, but also as a fusion with a heterologous polypeptide, preferably a

15

20

5

25

signal sequence or other polypeptide having a specific cleavage site at the Nterminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the target polypeptide DNA that is inserted into the vector. Included within the scope of this invention are target polypeptides with any native signal sequence deleted and replaced with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed (i.e. cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the native target polypeptide signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the native target polypeptide signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

(b) Origin of Replication Component

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

Most expression vectors are "shuttle" vectors, i.e. they are capable

10

15

25

20

30

51

of replication in at least one class of organisms but can be transfected into another organism for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells for expression even though it is not capable of replicating independently of the host cell chromosome.

DNA may also be amplified by insertion into the host genome. This is readily accomplished using *Bacillus* species as hosts, for example, by including in the vector a DNA sequence that is complementary to a sequence found in *Bacillus* genomic DNA. Transfection of *Bacillus* with this vector results in homologous recombination with the genome and insertion of the target polypeptide DNA. However, the recovery of genomic DNA encoding the target polypeptide is more complex than that of an exogenously replicated vector because restriction enzyme digestion is required to excise the target polypeptide DNA.

(c) Selection Gene Component

Expression and cloning vectors should contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g. ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g. the gene encoding D-alanine racemase for *Bacilli*.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene express a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin (Southern *et al.*, <u>J. Molec. Appl. Genet.</u>, <u>1</u>: 327 [1982]), mycophenolic acid (Mulligan *et al.*, <u>Science</u>, <u>209</u>: 1422 [1980]) or

25

20

10

15

30

52

hygromycin (Sugden *et al.*, <u>Mol. Cell. Biol.</u>, <u>5</u>: 410-413 [1985]). The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid), or hygromycin, respectively.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the target polypeptide nucleic acid, such as dihydrofolate reductase (DHFR) or thymidine kinase. The mammalian cell transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of having taken up the marker. Selection pressure is imposed by culturing the transformants under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes the target polypeptide. Amplification is the process by which genes in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Increased quantities of the target polypeptide are synthesized from the amplified DNA.

10

15

20

25

30

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, <u>Proc. Natl. Acad. Sci. USA</u>, <u>77</u>: 4216 [1980]. The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA comprising the expression vectors, such as the DNA encoding the target polypeptide. This amplification technique can be used with any otherwise suitable host, e.g., ATCC No. CCL61 CHO-K1, notwithstanding the presence of endogenous DHFR if, for example, a mutant DHFR gene that is highly resistant to Mtx is employed (EP 117,060). Alternatively, host cells (particularly wild-type hosts

47

that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding the target polypeptide, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3' phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See U.S. Pat. No. 4,965,199.

A suitable selection gene for use in yeast is the *trp*1 gene present in the yeast plasmid YRp7 (Stinchcomb *et al.*, <u>Nature</u>, <u>282</u>: 39 [1979]; Kingsman *et al.*, <u>Gene</u>, <u>7</u>: 141 [1979]; or Tschemper *et al.*, <u>Gene</u>, <u>10</u>: 157 [1980]). The *trp*1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 (Jones, <u>Genetics</u>, <u>85</u>: 12 [1977]). The presence of the <u>trp</u>1 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu*2-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu*2 gene.

(d) Promoter Component

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the target polypeptide nucleic acid. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular nucleic acid sequence, such as that encoding the target polypeptide, to which they are operably linked. Such promoters typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g. the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known. These promoters are operably linked to DNA encoding the target polypeptide

3

15

20

10

25

by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the vector. Both the native target polypeptide promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of the target polypeptide DNA. However, heterologous promoters are preferred, as they generally permit greater transcription and higher yields of expressed target polypeptide as compared to the native target polypeptide promoter.

Promoters suitable for use with prokaryotic hosts include the β lactamase and lactose promoter systems (Chang *et al.*, <u>Nature</u>, <u>275</u>: 615 [1978]; and Goeddel *et al.*, <u>Nature</u>, <u>281</u>: 544 [1979]), alkaline phosphatase, a tryptophan (trp) promoter system (Goeddel, <u>Nucleic Acids Res.</u>, <u>8</u>: 4057 [1980] and EP 36,776) and hybrid promoters such as the tac promoter (deBoer *et al.*, <u>Proc. Natl. Acad. Sci. USA</u>, <u>80</u>: 21-25 [1983]). However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to DNA encoding the target polypeptide (Siebenlist *et al.*, <u>Cell</u>, <u>20</u>: 269 [1980]) using linkers or adaptors to supply any required restriction sites. Promoters for use in bacterial systems also generally will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the target polypeptide.

Suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem., 255: 2073 [1980]) or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg., 7: 149 [1968]; and Holland, Biochemistry, 17: 4900 [1978]), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism,

10

15

5

-

25

20

30

49

metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in Hitzeman *et al.*, EP 73,657A. Yeast enhancers also are advantageously used with yeast promoters.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into mammalian expression vectors.

Target polypeptide transcription from vectors in mammalian host cells is controlled by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g. the actin promoter or an immunoglobulin promoter, from heat-shock promoters, and from the promoter normally associated with the target polypeptide sequence, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. Fiers *et al.*, <u>Nature</u>, <u>273</u>:113 (1978); Mulligan and Berg, <u>Science</u>, <u>209</u>: 1422-1427 (1980); Pavlakis *et al.*, <u>Proc. Natl. Acad. Sci. USA</u>, <u>78</u>: 7398-7402 (1981). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a <u>HindIII E restriction fragment</u>. Greenaway *et al.*, <u>Gene</u>, <u>18</u>: 355-360 (1982). A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed

10

5

15

20

25

30

50

in U.S. 4,419,446. A modification of this system is described in U.S. 4,601,978. See also Gray *et al.*, <u>Nature</u>, <u>295</u>: 503-508 (1982) on expressing cDNA encoding immune interferon in monkey cells; , Reyes *et al.*, <u>Nature</u>, <u>297</u>: 598-601 (1982) on expression of human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus, Canaani and Berg, <u>Proc. Natl. Acad. Sci. USA</u>, <u>79</u>: 5166-5170 (1982) on expression of the human interferon β 1 gene in cultured mouse and rabbit cells, and Gorman *et al.*, <u>Proc. Natl. Acad. Sci. USA</u>, <u>79</u>: 6777-6781 (1982) on expression of bacterial CAT sequences in CV-1 monkey kidney cells, chicken embryo fibroblasts, Chinese hamster ovary cells, HeLa cells, and mouse NIH-3T3 cells using the Rous sarcoma virus long terminal repeat as a promoter.

(e) Enhancer Element Component

Transcription of DNA encoding the target polypeptide of this invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10-300 bp, that act on a promoter to increase its transcription. Enhancers are relatively orientation and position independent having been found 5' (Laimins et al., Proc. Natl. Acad. Sci. USA, 78: 993 [1981]) and 3' (Lusky et al., Mol. Cell Bio., 3: 1108 [1983]) to the transcription unit, within an intron (Banerji et al., Cell, 33: 729 [1983]) as well as within the coding sequence itself (Osborne et al., Mol. Cell Bio., 4: 1293 [1984]). Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, σ -fetoprotein and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature, 297: 17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a

10

5

15

20

25

30

51

position 5' or 3' to the target polypeptide DNA, but is preferably located at a site 5' from the promoter.

(f) Transcription Termination Component

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the target polypeptide. The 3' untranslated regions also include transcription termination sites.

Construction of suitable vectors containing one or more of the above listed components the desired coding and control sequences employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced by the method of Messing *et al.*, <u>Nucleic Acids Res.</u>, <u>9</u>: 309 (1981) or by the method of Maxam *et al.*, <u>Methods in Enzymology</u>, <u>65</u>: 499 (1980).

Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding the target polypeptide. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the

10

-20

15

30

25

52

expression vector. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of polypeptides encoded by cloned DNAs, as well as for the rapid screening of such polypeptides for desired biological or physiological properties. Thus, transient expression systems are particularly useful in the invention for purposes of identifying analogs and variants of the target polypeptide that have target polypeptide-like activity.

Other methods, vectors, and host cells suitable for adaptation to the synthesis of the target polypeptide in recombinant vertebrate cell culture are described in Gething *et al.*, <u>Nature</u>, <u>293</u>: 620-625 [1981]; Mantei *et al.*, <u>Nature</u>, <u>281</u>: 40-46 [1979]; Levinson *et al.*; EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of the target polypeptide is pRK5 (EP pub. no. 307,247) or pSVI6B (U.S. Ser. No. 07/441,574 filed 22 November 1989, the disclosure of which is incorporated herein by reference).

Selection and Transformation of Host Cells

Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes include eubacteria, such as Gram-negative or Gram-positive organisms, for example, *E. coli*, *Bacilli* such as *B. subtilis*, *Pseudomonas* species such as *P. aeruginosa*, *Salmonella typhimurium*, or *Serratia marcescans*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* χ 1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting. Preferably the host cell should secrete minimal amounts of proteolytic enzymes. Alternatively, *in vitro* methods of cloning, e.g. PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable hosts for target polypeptide-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly

10

15

20

25

30

Suitable host cells for the expression of glycosylated target polypeptide are derived from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture. Examples of invertebrate cells include plant and insect Numerous baculoviral strains and variants and corresponding cells. permissive insect host cells from hosts such as Spodoptera frugiperda (caterpillar), Aedes aegypti (mosquito), Aedes albopictus (mosquito), Drosophila melanogaster (fruitfly), and Bombyx mori host cells have been identified. See, e.g., Luckow et al., Bio/Technology, 6: 47-55 (1988); Miller et al., in Genetic Engineering, Setlow, J.K. et al., eds., Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda et al., Nature, 315: 592-594 (1985). A variety of such viral strains are publicly available, e.g., the L-1 variant of Autographa californica NPV and the Bm-5 strain of Bombyx mori NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of Spodoptera frugiperda cells. Plant cell cultures of cotton, corn, potato, soybean, petunia,

10

15

20

25

30

54

tomato, and tobacco can be utilized as hosts. Typically, plant cells are transfected by incubation with certain strains of the bacterium *Agrobacterium tumefaciens*, which has been previously manipulated to contain the target polypeptide DNA. During incubation of the plant cell culture with *A. tumefaciens*, the DNA encoding target polypeptide is transferred to the plant cell host such that it is transfected, and will, under appropriate conditions, express the target polypeptide DNA. In addition, regulatory and signal sequences compatible with plant cells are available, such as the nopaline synthase promoter and polyadenylation signal sequences. Depicker *et al.*, <u>J.</u> <u>Mol. Appl. Gen.</u>, <u>1</u>: 561 (1982). In addition, DNA segments isolated from the upstream region of the T-DNA *780* gene are capable of activating or increasing transcription levels of plant-expressible genes in recombinant DNAcontaining plant tissue. See EP 321,196 published 21 June 1989.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure in recent years [Tissue Culture, Academic Press, Kruse and Patterson, editors (1973)]. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol., 36: 59 [1977]); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77: 4216 [1980]); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23: 243-251 [1980]); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad, Sci., 383: 44-68 [1982]); MRC 5 cells; FS4 cells; and a human hepatoma cell line (Hep G2). Preferred host cells are human embryonic kidney 293 and Chinese hamster ovary cells.

10

5

15

20

25

30

55

Host cells are transfected and preferably transformed with the above-described expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Transfection refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, $CaPO_4$ and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook et al., supra, is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with Agrobacterium tumefaciens is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23: 315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method described in sections 16.30-16.37 of Sambrook et al, supra, is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. 4,399,216, issued 16 August 1983. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130: 946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76: 3829 (1979). However, other methods for introducing DNA into cells such as by nuclear injection, electroporation, or protoplast fusion may also be used.

Culturing the Host Cells

10

15

20

-5

25

30

Prokaryotic cells used to produce the target polypeptide of this invention are cultured in suitable media as described generally in Sambrook *et al., supra.*

The mammalian host cells used to produce the target polypeptide of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ([MEM], Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ([DMEM], Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, Meth. Enz., 58: 44 (1979), Barnes and Sato, Anal. Biochem., 102: 255 (1980), U.S. 4,767,704; 4,657,866; 4,927,762; or 4,560,655; WO 90/03430; WO 87/00195; U.S. Pat. Re. 30,985; or copending U.S.S.N. 07/592,107 or 07/592,141, both filed in 3 October 1990, the disclosures of all of which are incorporated herein by reference, may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as Gentamycin[™] drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The host cells referred to in this disclosure encompass cells in *in vitro* culture as well as cells that are within a host animal.

It is further envisioned that the target polypeptides of this invention may be produced by homologous recombination, or with recombinant production methods utilizing control elements introduced into cells already containing DNA encoding the target polypeptide currently in use in the field. For example, a powerful promoter/enhancer element, a suppressor, or an

57

125

15

10

5

20

25

30

exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired target polypeptide. The control element does not encode the target polypeptide of this invention, but the DNA is present in the host cell genome. One next screens for cells making the target polypeptide of this invention, or increased or decreased levels of expression, as desired.

Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77: 5201-5205 [1980]), dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Various labels may be employed, most commonly radioisotopes, particularly ³²P. However, other techniques may also be employed, such as using biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. With immunohistochemical staining techniques, a cell sample is prepared, typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product coupled, where the labels are

10

15

20



30

usually visually detectable, such as enzymatic labels, fluorescent labels, luminescent labels, and the like. A particularly sensitive staining technique suitable for use in the present invention is described by Hsu *et al.*, <u>Am. J.</u> <u>Clin. Path.</u>, <u>75</u>: 734-738 (1980).

Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native target polypeptide or against a synthetic peptide based on the DNA sequences provided herein as described further in Section 4 below. Purification of The Target polypeptide

The target polypeptide preferably is recovered from the culture medium as a secreted polypeptide, although it also may be recovered from host cell lysates when directly expressed without a secretory signal.

When the target polypeptide is expressed in a recombinant cell other than one of human origin, the target polypeptide is completely free of proteins or polypeptides of human origin. However, it is necessary to purify the target polypeptide from recombinant cell proteins or polypeptides to obtain preparations that are substantially homogeneous as to the target polypeptide. As a first step, the culture medium or lysate is centrifuged to remove particulate cell debris. The membrane and soluble protein fractions are then separated. The target polypeptide may then be purified from the soluble protein fraction and from the membrane fraction of the culture lysate, depending on whether the target polypeptide is membrane bound. The following procedures are exemplary of suitable purification procedures: or ion-exchange columns; ethanol fractionation on immunoaffinity precipitation; reverse phase HPLC; chromatography on silica or on a cation exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; and protein A Sepharose columns to remove contaminants such as IgG.

Target polypeptide variants in which residues have been deleted, inserted or substituted are recovered in the same fashion, taking account of

10

5

20

15

25

30

any substantial changes in properties occasioned by the variation. For example, preparation of a target polypeptide fusion with another protein or polypeptide, e.g. a bacterial or viral antigen, facilitates purification; an immunoaffinity column containing antibody to the antigen (or containing antigen, where the target polypeptide is an antibody) can be used to adsorb the fusion. Immunoaffinity columns such as a rabbit polyclonal anti-target polypeptide column can be employed to absorb the target polypeptide variant by binding it to at least one remaining immune epitope. A protease inhibitor such as phenyl methyl sulfonyl fluoride (PMSF) also may be useful to inhibit proteolytic degradation during purification, and antibiotics may be included to prevent the growth of adventitious contaminants. One skilled in the art will appreciate that purification methods suitable for native target polypeptide may require modification to account for changes in the character of the target polypeptide or its variants upon expression in recombinant cell culture.

Covalent Modifications of Target Polypeptides

Covalent modifications of target polypeptides are included within the scope of this invention. One type of covalent modification included within the scope of this invention is a target polypeptide fragment. Target polypeptide fragments having up to about 40 amino acid residues may be conveniently prepared by chemical synthesis, or by enzymatic or chemical cleavage of the full-length target polypeptide or variant target polypeptide. Other types of covalent modifications of the target polypeptide or fragments thereof are introduced into the molecule by reacting specific amino acid residues of the target polypeptide or fragments thereof with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteinyl residues most commonly are reacted with *a*-haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, *a*-

10

15

20

.

25

30

60

bromo-*B*-(5-imidozoyl)propionic acid, chloroacetyl phosphate, Nalkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, pchloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing *a*-amino-containing residues include imidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK, of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidizole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ¹²⁵I or ¹³¹I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R'-N = C = N-R'), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl

15

20

25

30

and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking target polypeptide to a water-insoluble support matrix or surface for use in the method for purifying anti-target polypeptide antibodies, and vice versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing such as methyl-3-[(p-azidophenyl)dithio]propioimidate yield agents photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the *a*-amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, <u>Proteins: Structure and Molecular Properties</u>, W.H. Freeman & Co., San Francisco, pp. 79-86 [1983]), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the target polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. By altering is meant deleting one or more carbohydrate moieties found in the native target polypeptide, and/or adding one or more glycosylation sites that are not present in the native target polypeptide.

62

10

15

20

25

Glycosylation of polypeptides is typically either N-linked or Olinked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-Xserine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars Nacetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

5

10

15

20

, ‡

25

30

Addition of glycosylation sites to the target polypeptide is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for Nlinked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the native target polypeptide sequence (for O-linked glycosylation sites). For ease, the target polypeptide amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids. The DNA mutation(s) may be made using methods described above under the heading of "Amino Acid Sequence Variants of Target Polypeptide".

Another means of increasing the number of carbohydrate moleties on the target polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. These procedures are advantageous in that they do not require production of the polypeptide in a host cell that has glycosylation capabilities for N- and O- linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan,

63

or (f) the amide group of glutamine. These methods are described in WO 87/05330 published 11 September 1987, and in Aplin and Wriston (<u>CRC</u> <u>Crit. Rev. Biochem.</u>, pp. 259-306 [1981]).

5

10

15

20

25

30

Removal of carbohydrate moieties present on the native target polypeptide may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the polypeptide to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (Nacetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin *et al.* (Arch. <u>Biochem. Biophys.</u>, 259:52 [1987]) and by Edge *et al.* (Anal. Biochem., 118:131 [1981]). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exoglycosidases as described by Thotakura *et al.* (Meth. Enzymol., 138:350 [1987]).

Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin *et al.* (<u>J. Biol.</u> <u>Chem.</u>, <u>257</u>:3105 [1982]). Tunicamycin blocks the formation of protein-Nglycoside linkages.

Another type of covalent modification of the target polypeptide comprises linking the target polypeptide to various nonproteinaceous polymers, e.g. polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in U.S. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The target polypeptide also may be entrapped in microcapsules, prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-[methylmethacylate] microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in <u>Remington's Pharmaceutical Sciences</u>, 16th edition, Osol, A., Ed., (1980).

64

Target polypeptide preparations are also useful in generating antibodies, for screening for binding partners, as standards in assays for the target polypeptide (e.g. by labeling the target polypeptide for use as a standard in a radioimmunoassay, enzyme-linked immunoassay, or radioreceptor assay), in affinity purification techniques, and in competitivetype receptor binding assays when labeled with radioiodine, enzymes, fluorophores, spin labels, and the like.

Since it is often difficult to predict in advance the characteristics of a variant target polypeptide, it will be appreciated that some screening of the recovered variant will be needed to select the optimal variant. For example, a change in the immunological character of the target polypeptide molecule, such as affinity for a given antigen or antibody, is measured by a competitive-type immunoassay. The variant is assayed for changes in the suppression or enhancement of its activity by comparison to the activity observed for the target polypeptide in the same assay. Other potential modifications of protein or polypeptide properties such as redox or thermal stability, hydrophobicity, susceptibility to proteolytic degradation, stability in recombinant cell culture or in plasma, or the tendency to aggregate with carriers or into multimers are assayed by methods well known in the art.

Diagnostic and Related Uses of the Antibodies

10

15

20

25

30

The antibodies of this invention are useful in diagnostic assays for antigen expression in specific cells or tissues. The antibodies are detectably labeled and/or are immobilized on an insoluble matrix.

The antibodies of this invention find further use for the affinity purification of the antigen from recombinant cell culture or natural sources.

Suitable diagnostic assays for the antigen and its antibodies depend on the particular antigen or antibody. Generally, such assays include competitive and sandwich assays, and steric inhibition assays. Competitive and sandwich methods employ a phase-separation step as an integral part of the method while steric inhibition assays are conducted in a single reaction mixture. Fundamentally, the same procedures are used for the assay of the

65

antigen and for substances that bind the antigen, although certain methods will be favored depending upon the molecular weight of the substance being assayed. Therefore, the substance to be tested is referred to herein as an analyte, irrespective of its status otherwise as an antigen or antibody, and proteins that bind to the analyte are denominated binding partners, whether they be antibodies, cell surface receptors, or antigens.

Analytical methods for the antigen or its antibodies all use one or more of the following reagents: labeled analyte analogue, immobilized analyte analogue, labeled binding partner, immobilized binding partner and steric conjugates. The labeled reagents also are known as "tracers."

The label used (and this is also useful to label antigen nucleic acid for use as a probe) is any detectable functionality that does not interfere with the binding of analyte and its binding partner. Numerous labels are known for use in immunoassay, examples including moieties that may be detected directly, such as fluorochrome, chemiluminescent, and radioactive labels, as well as moieties, such as enzymes, that must be reacted or derivatized to be detected. Examples of such labels include the radioisotopes ³²P, ¹⁴C, ¹²⁵I, ³H, and ¹³¹I, fluorophores such as rare earth chelates or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, luceriferases, e.g., firefly luciferase and bacterial luciferase (U.S. Pat. No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase (HRP), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases, e.g., glucose oxidase, galactose oxidase, and glucose-6phosphate dehydrogenase, heterocyclic oxidases such as uricase and xanthine oxidase, coupled with an enzyme that employs hydrogen peroxide to oxidize a dye precursor such as HRP, lactoperoxidase, or microperoxidase, biotin/avidin, spin labels, bacteriophage labels, stable free radicals, and the like.

Conventional methods are available to bind these labels covalently to proteins or polypeptides. For instance, coupling agents such as dialdehydes, carbodiimides, dimaleimides, bis-imidates, bis-diazotized benzidine, and the like may be used to tag the antibodies with the above-

10

-

20

15

25

30

described fluorescent, chemiluminescent, and enzyme labels. See, for example, U.S. Pat. Nos. 3,940,475 (fluorimetry) and 3,645,090 (enzymes); Hunter *et al.*, <u>Nature</u>, <u>144</u>: 945 (1962); David *et al.*, <u>Biochemistry</u>, <u>13</u>: 1014-1021 (1974); Pain *et al.*, <u>J. Immunol. Methods</u>, <u>40</u>: 219-230 (1981); and Nygren, <u>J. Histochem. and Cytochem.</u>, <u>30</u>: 407-412 (1982). Preferred labels herein are enzymes such as horseradish peroxidase and alkaline phosphatase.

The conjugation of such label, including the enzymes, to the antibody is a standard manipulative procedure for one of ordinary skill in immunoassay techniques. See, for example, O'Sullivan *et al.*, "Methods for the Preparation of Enzyme-antibody Conjugates for Use in Enzyme Immunoassay," in <u>Methods in Enzymology</u>, ed. J.J. Langone and H. Van Vunakis, Vol. 73 (Academic Press, New York, New York, 1981), pp. 147-166. Such bonding methods are suitable for use with the antibodies and polypeptides of this invention.

Immobilization of reagents is required for certain assay methods. Immobilization entails separating the binding partner from any analyte that remains free in solution. This conventionally is accomplished by either insolubilizing the binding partner or analyte analogue before the assay procedure, as by adsorption to a water-insoluble matrix or surface (Bennich *et al..*, U.S. 3,720,760), by covalent coupling (for example, using glutaraldehyde cross-linking), or by insolubilizing the partner or analogue afterward, e.g., by immunoprecipitation.

Other assay methods, known as competitive or sandwich assays, are well established and widely used in the commercial diagnostics industry.

Competitive assays rely on the ability of a tracer analogue to compete with the test sample analyte for a limited number of binding sites on a common binding partner. The binding partner generally is insolubilized before or after the competition and then the tracer and analyte bound to the binding partner are separated from the unbound tracer and analyte. This separation is accomplished by decanting (where the binding partner was preinsolubilized) or by centrifuging (where the binding partner was precipitated after the competitive reaction). The amount of test sample

67

10

5

15

20

25

analyte is inversely proportional to the amount of bound tracer as measured by the amount of marker substance. Dose-response curves with known amounts of analyte are prepared and compared with the test results to quantitatively determine the amount of analyte present in the test sample. These assays are called ELISA systems when enzymes are used as the detectable markers.

Another species of competitive assay, called a "homogeneous" assay, does not require a phase separation. Here, a conjugate of an enzyme with the analyte is prepared and used such that when anti-analyte binds to the analyte the presence of the anti-analyte modifies the enzyme activity. In this case, the antigen or its immunologically active fragments are conjugated with a bifunctional organic bridge to an enzyme such as peroxidase. Conjugates are selected for use with antibody so that binding of the antibody inhibits or potentiates the enzyme activity of the label. This method *per se* is widely practiced under the name of EMIT.

Steric conjugates are used in steric hindrance methods for homogeneous assay. These conjugates are synthesized by covalently linking a low-molecular-weight hapten to a small analyte so that antibody to hapten substantially is unable to bind the conjugate at the same time as anti-analyte. Under this assay procedure the analyte present in the test sample will bind anti-analyte, thereby allowing anti-hapten to bind the conjugate, resulting in a change in the character of the conjugate hapten, e.g., a change in fluorescence when the hapten is a fluorophore.

Sandwich assays particularly are useful for the determination of antigen or antibodies. In sequential sandwich assays an immobilized binding partner is used to adsorb test sample analyte, the test sample is removed as by washing, the bound analyte is used to adsorb labeled binding partner, and bound material is then separated from residual tracer. The amount of bound tracer is directly proportional to test sample analyte. In "simultaneous" sandwich assays the test sample is not separated before adding the labeled binding partner. A sequential sandwich assay using an anti-antigen monoclonal antibody as one antibody and a polyclonal anti-antigen antibody

15

20

25

30

5

as the other is useful in testing samples for particular antigen activity.

The foregoing are merely exemplary diagnostic assays for the import and humanized antibodies of this invention. Other methods now or hereafter developed for the determination of these analytes are included within the scope hereof, including the bioassays described above.

<u>Immunotoxins</u>

This invention is also directed to immunochemical derivatives of the antibodies of this invention such as immunotoxins (conjugates of the antibody and a cytotoxic moiety). Antibodies which carry the appropriate effector functions, such as with their constant domains, are also used to induce lysis through the natural complement process, and to interact with antibody dependent cytotoxic cells normally present.

For example, purified, sterile filtered antibodies are optionally conjugated to a cytotoxin such as ricin for use in AIDS therapy. US Patent Application Serial No. 07/350,895 illustrates methods for making and using immunotoxins for the treatment of HIV infection, and its teachings are specifically incorporated by reference herein. The methods of this invention, for example, are suitable for obtaining humanized antibodies for use as immunotoxins for use in AIDS therapy.

The cytotoxic moiety of the immunotoxin may be a cytotoxic drug or an enzymatically active toxin of bacterial, fungal, plant or animal origin, or an enzymatically active fragment of such a toxin. Enzymatically active toxins and fragments thereof used are diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. In another embodiment, the antibodies are conjugated to small molecule anticancer drugs such as cis-platin or 5FU. Conjugates of the monoclonal antibody and such cytotoxic moieties are made

10

5

..

20

15

25

69

using a variety of bifunctional protein coupling agents. Examples of such reagents are SPDP, IT, bifunctional derivatives of imidoesters such as dimethyl adipimidate HCl, active esters such as disuccinimidyl suberate, aldehydes such as glutaraldehyde, bis-azido compounds such as bis (p-azidobenzoyl) hexanediamine, bis-diazonium derivatives such as bis- (pdiazoniumbenzoyl)- -ethylenediamine, diisocyanates such as tolylene 2,6-diisocyanate and bis-active fluorine compounds such as 1,5-difluoro-2,4-dinitrobenzene. The lysing portion of a toxin may be joined to the Fab fragment of the antibodies.

Immunotoxins can be made in a variety of ways, as discussed herein. Commonly known crosslinking reagents can be used to yield stable conjugates.

Advantageously, monoclonal antibodies specifically binding the domain of the antigen which is exposed on the infected cell surface, are conjugated to ricin A chain. Most advantageously the ricin A chain is deglycosylated and produced through recombinant means. An advantageous method of making the ricin immunotoxin is described in Vitetta *et al.*, *Science* 238:1098 (1987) hereby incorporated by reference.

When used to kill infected human cells *in vitro* for diagnostic purposes, the conjugates will typically be added to the cell culture medium at a concentration of at least about 10 nM. The formulation and mode of administration for *in vitro* use are not critical. Aqueous formulations that are compatible with the culture or perfusion medium will normally be used. Cytotoxicity may be read by conventional techniques.

Cytotoxic radiopharmaceuticals for treating infected cells may be made by conjugating radioactive isotopes (e.g. I, Y, Pr) to the antibodies. Advantageously alpha particle-emitting isotopes are used. The term 'cytotoxic moiety" as used herein is intended to include such isotopes.

In a preferred embodiment, ricin A chain is deglycosylated or produced without oligosaccharides, to decrease its clearance by irrelevant clearance mechanisms (e.g., the liver). In another embodiment, whole ricin (A chain plus B chain) is conjugated to antibody if the galactose binding

70

10

15

25

20

property of B-chain can be blocked ("blocked ricin").

In a further embodiment toxin-conjugates are made with Fab or $F(ab')_2$ fragments. Because of their relatively small size these fragments can better penetrate tissue to reach infected cells.

In another embodiment, fusogenic liposomes are filled with a cytotoxic drug and the liposomes are coated with antibodies specifically binding the particular antigen.

Antibody Dependent Cellular Cytotoxicity

Certain aspects of this invention involve antibodies which are (a) directed against a particular antigen and (b) belong to a subclass or isotype that is capable of mediating the lysis of cells to which the antibody molecule binds. More specifically, these antibodies should belong to a subclass or isotype that, upon complexing with cell surface proteins, activates serum complement and/or mediates antibody dependent cellular cytotoxicity (ADCC) by activating effector cells such as natural killer cells or macrophages.

Biological activity of antibodies is known to be determined, to a large extent, by the constant domains or Fc region of the antibody molecule (Uananue and Benacerraf, Textbook of Immunology, 2nd Edition, Williams & Wilkins, p. 218 (1984)). This includes their ability to activate complement and to mediate antibody-dependent cellular cytotoxicity (ADCC) as effected by leukocytes. Antibodies of different classes and subclasses differ in this respect, as do antibodies from the same subclass but different species; according to the present invention, antibodies of those classes having the desired biological activity are prepared. Preparation of these antibodies involves the selection of antibody constant domains are their incorporation in the humanized antibody by known technique. For example, mouse immunoglobulins of the IgG3 and IgG2a class are capable of activating serum complement upon binding to the target cells which express the cognate antigen, and therefore humanized antibodies which incorporate IgG3 and IgG2a effector functions are desirable for certain therapeutic applications. In general, mouse antibodies of the IgG2a and IgG3 subclass and

10

15

25

30

20

occasionally IgG1 can mediate ADCC, and antibodies of the IgG3, IgG2a, and IgM subclasses bind and activate serum complement. Complement activation generally requires the binding of at least two IgG molecules in close proximity on the target cell. However, the binding of only one IgM molecule activates serum complement.

The ability of any particular antibody to mediate lysis of the target cell by complement activation and/or ADCC can be assayed. The cells of interest are grown and labeled *in vitro*; the antibody is added to the cell culture in combination with either serum complement or immune cells which may be activated by the antigen antibody complexes. Cytolysis of the target cells is detected by the release of label from the lysed cells. In fact, antibodies can be screened using the patient's own serum as a source of complement and/or immune cells. The antibody that is capable of activating complement or mediating ADCC in the *in vitro* test can then be used therapeutically in that particular patient.

This invention specifically encompasses consensus Fc antibody domains prepared and used according to the teachings of this invention.

Therapeutic and Other Uses of the Antibodies

When used *in vivo* for therapy, the antibodies of the subject invention are administered to the patient in therapeutically effective amounts (i.e. amounts that have desired therapeutic effect). They will normally be administered parenterally. The dose and dosage regimen will depend upon the degree of the infection, the characteristics of the particular antibody or immunotoxin used, e.g., its therapeutic index, the patient, and the patient's history. Advantageously the antibody or immunotoxin is administered continuously over a period of 1-2 weeks, intravenously to treat cells in the vasculature and subcutaneously and intraperitoneally to treat regional lymph nodes. Optionally, the administration is made during the course of adjunct therapy such as combined cycles of radiation, chemotherapeutic treatment, or administration of tumor necrosis factor, interferon or other cytoprotective or immunomodulatory agent.

10

20

25

30

For parenteral administration the antibodies will be formulated in a unit dosage injectable form (solution, suspension, emulsion) in association with a pharmaceutically acceptable parenteral vehicle. Such vehicles are inherently nontoxic, and non-therapeutic. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils and ethyl oleate can also be used. Liposomes may be used as carriers. The vehicle may contain minor amounts of additives such as substances that enhance isotonicity and chemical stability, e.g., buffers and preservatives. The antibodies will typically be formulated in such vehicles at concentrations of about 1 mg/ml to 10 mg/ml.

Use of IgM antibodies may be preferred for certain applications, however IgG molecules by being smaller may be more able than IgM molecules to localize to certain types of infected cells.

There is evidence that complement activation in vivo leads to a variety of biological effects, including the induction of an inflammatory response and the activation of macrophages (Uananue and Benecerraf, Textbook of Immunology, 2nd Edition, Williams & Wilkins, p. 218 (1984)). The increased vasodilation accompanying inflammation may increase the ability of various agents to localize in infected cells. Therefore, antigen-antibody combinations of the type specified by this invention can be used therapeutically in many ways. Additionally, purified antigens (Hakomori, Ann. Rev. Immunol. 2:103 (1984)) or anti-idiotypic antibodies (Nepom et al., Proc. Natl. Acad. Sci. 81:2864 (1985); Koprowski et al., Proc. Natl. Acad. Sci. 81:216 (1984)) relating to such antigens could be used to induce an active immune response in human patients. Such a response includes the formation of antibodies capable of activating human complement and mediating ADCC and by such mechanisms cause infected cell destruction.

Optionally, the antibodies of this invention are useful in passively immunizing patients, as exemplified by the administration of humanized anti-HIV antibodies.

10

5

15

20

25

30

The antibody compositions used in therapy are formulated and dosages established in a fashion consistent with good medical practice taking into account the disorder to be treated, the condition of the individual patient, the site of delivery of the composition, the method of administration and other factors known to practitioners. The antibody compositions are prepared for administration according to the description of preparation of polypeptides for administration, *infra*.

Deposit of Materials

As described above, cultures of the muMAb4D5 have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, USA (ATCC).

This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of viable cultures for 30 years from the date of the deposit. The organisms will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the cultures to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.12 with particular reference to 886 OG 638).

In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC)

10

15

20

25

30

The assignee of the present application has agreed that if the cultures on deposit should die or be lost or destroyed when cultivated under suitable conditions, they will be promptly replaced on notification with a viable specimen of the same culture. Availability of the deposited strain is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the constructs deposited, since the deposited embodiments are intended to illustrate only certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that they represent. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

It is understood that the application of the teachings of the present invention to a specific problem or situation will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein. Examples of the products of the present invention and representative processes for their isolation, use, and manufacture appear below, but should not be construed to limit the invention. All literature citations herein are expressly incorporated by reference.

EXAMPLES

EXAMPLE 1. HUMANIZATION OF muMAb4D5

75

81

10

5

15

20

25

Here we report the chimerization of muMAb4D5 (chMAb4D5) and the rapid and simultaneous humanization of heavy (V_H) and light (V_L) chain variable region genes using a novel "gene conversion mutagenesis" strategy. Eight humanized variants (huMAb4D5) were constructed to probe the importance of several FR residues identified by our molecular modeling or previously proposed to be critical to the conformation of particular CDRs (see Chothia, C. & Lesk, A. M., *J. Mol. Biol.* **196**:901-917 (1987); Chothia, C. *et al.*, *Nature* **342**:877-883 (1989); Tramontano, A. *et al.*, *J. Mol. Biol.* **215**:175-182 (1990)). Efficient transient expression of humanized variants in non-myeloma cells allowed us to rapidly investigate the relationship between binding affinity for p185^{HER2} ECD and anti-proliferative activity against p185^{HER2} overexpressing carcinoma cells.

MATERIALS and METHODS

Cloning of Variable Region Genes. The muMAb4D5 V_H and V_L genes were isolated by polymerase chain reaction (PCR) amplification of mRNA from the corresponding hybridoma (Fendly, B. M. et al., Cancer Res. 50:1550-1558 (1990)) as described by Orlandi et al. (Orlandi, R. et al., Proc. Natl. Acad. Sci. USA 86:3833-3837 (1989)). Amino terminal sequencing of muMAb4D5 V_L and V_H was used to design the sense strand PCR primers, whereas the anti-sense PCR primers were based upon consensus sequences of murine framework residues (Orlandi, R. et al., Proc. Natl. Acad. Sci. USA 86:3833-3837 (1989); Kabat, E. A. et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)) incorporating restriction sites for directional cloning shown by underlining and V₁ sense, 5'listed after the sequences: TCCGATATCCAGCTGACCCAGTCTCCA-3' (SEQ. ID NO. 7), EcoRV; VI anti-sense, 5'-GTTTGATCTCCAGCTTGGTACCHSCDCCGAA-3' (SEQ. ID NO. 8), Asp718; V_H sense, 5'-AGGTSMARCTGCAGSAGTCWGG-3' (SEQ. ID NO. anti-sense, Pstl 5 🔶 -9), a n d νн TGAGGAGACGGTGACCGTGGTCCCTTGGCCCCAG-3' (SEQ. ID NO. 10), BstEll; where H = A or C or T, S = C or G, D = A or G or T, M = A or C, R = A or G and W = A or T. The PCR products were cloned into pUC119

10

15

20

25

30 ·

(Vieira, J. & Messing, J., *Methods Enzymol.* **153**:3-11 (1987)) and five clones for each variable domain sequenced by the dideoxy method (Sanger, F. *et al.*, *Proc. Natl. Acad. Sci. USA* **74**:5463-5467 (1977)).

Molecular Modelling. Models for muMAb4D5 V_H and V_L domains were constructed separately from consensus coordinates based upon seven Fab structures from the Brookhaven protein data bank (entries 1FB4, 2RHE, 2MCP, 3FAB, 1FBJ, 2HFL and 1REI). The Fab fragment KOL (Marquart, M. et al., J. Mol. Biol. 141:369-391 (1980)) was first chosen as a template for V_{L} and V_{H} domains and additional structures were then superimposed upon this structure using their main chain atom coordinates (INSIGHT program, Biosym Technologies). The distance from the template $C\alpha$ to the analogous Ca in each of the superimposed structures was calculated for each residue position. If all (or nearly all) Ca-Ca distances for a given residue were $\leq 1\text{\AA}$, then that position was included in the consensus structure. In most cases the β -sheet framework residues satisfied these criteria whereas the CDR loops did not. For each of these selected residues the average coordinates for individual N, Ca, C, O and Cß atoms were calculated and then corrected for resultant deviations from non-standard bond geometry by 50 cycles of energy minimization using the DISCOVER program (Biosym Technologies) with the AMBER forcefield (Weiner, S. J. et al., J. Amer. Chem. Soc. 106:765-784 (1984)) and Ca coordinates fixed. The side chains of highly conserved residues, such as the disulfide-bridged cysteine residues, were then incorporated into the resultant consensus structure. Next the sequences of muMAb4D5 V_L and V_H were incorporated starting with the CDR residues and using the tabulations of CDR conformations from Chothia et al. (Chothia, C. et al., Nature 342:877-883 (1989)) as a guide. Side-chain conformations were chosen on the basis of Fab crystal structures, rotamer libraries (Ponder, J. W. & Richards, F. M., J. Mol. Biol. 193:775-791 (1987)) and packing considerations. Since V_H-CDR3 could not be assigned a definite backbone conformation from these criteria, two models were created from a search of similar sized loops using the INSIGHT program. A third model was derived using packing and solvent exposure considerations. Each model

10

15

20

25

30

77

was then subjected to 5000 cycles of energy minimization.

In humanizing muMAb4D5, consensus human sequences were first derived from the most abundant subclasses in the sequence compilation of Kabat et al. (Kabat, E. A. et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)), namely V₁ κ subgroup I and $V_{\mbox{H}}$ group III, and a molecular model generated for these sequences using the methods described above. A structure for huMAb4D5 was created by transferring the CDRs from the muMAb4D5 model into the All huMAb4D5 variants contain human consensus human structure. replacements of muMAb4D5 residues at three positions within CDRs as defined by sequence variability (Kabat, E. A. et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)) but not as defined by structural variability (Chothia, C. & Lesk, A. M., J. Mol. Biol. 196:901-917 (1987)): VL-CDR1 K24R, VL-CDR2 R54L and VL-CDR2 Differences between muMAb4D5 and the human consensus T56S. framework residues (Fig. 1) were individually modeled to investigate their possible influence on CDR conformation and/or binding to the p185^{HER2} ECD.

Construction of Chimeric Genes. Genes encoding chMAb4D5 light and heavy chains were separately assembled in previously described phagemid vectors containing the human cytomegalovirus enhancer and promoter, a 5' intron and SV40 polyadenylation signal (Gorman, C. M. *et al.*, *DNA & Prot. Engin. Tech.* 2:3-10 (1990)). Briefly, gene segments encoding muMAb4D5 V_L (Fig. 1A) and REI human κ_1 light chain C_L (Palm, W. & Hilschmann, N., *Z. Physiol. Chem.* 356:167-191 (1975)) were precisely joined as were genes for muMAb4D5 V_H (Fig. 1B) and human y1 constant region (Capon, D. J. *et al.*, *Nature* 337:525-531 (1989)) by simple subcloning (Boyle, A., in *Current Protocols in Molecular Biology*, Chapter 3 (F. A. Ausubel *et al.*, eds., Greene Publishing & Wiley-Interscience, New York, 1990)) and site-directed mutagenesis (Carter, P., in *Mutagenesis: A Practical Approach*, Chapter 1 (IRL Press, Oxford, UK 1991)). The y1 isotype was chosen as it has been found to be the preferred human isotype for

10

15

20



30

supporting ADCC and complement dependent cytotoxicity using matched sets of chimeric (Brüggemann, M. et al., J. Exp. Med. 166:1351-1361 (1987)) or humanized antibodies (Riechmann, L. et al., Nature 332:323-327 The PCR-generated V_L and V_H fragments (Fig. 1) were (1988)). subsequently mutagenized so that they faithfully represent the sequence of muMAb4D5 determined at the protein level: $V_H Q1E$, $V_L V104L$ and T109A (variants are denoted by the amino acid residue and number followed by the replacement amino acid). The human y1 constant regions are identical to those reported by Ellison et al. (Ellison, J. W. et al., Nucleic Acids Res. 13:4071-4079 (1982)) except for the mutations E359D and M361L (Eu numbering, as in Kabat, E. A. et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)) which we installed to convert the antibody from the naturally rare A allotype to the much more common non-A allotype (Tramontano, A. et al., J. Mol. Biol. 215:175-182 (1990)). This was an attempt to reduce the risk of anti-allotype antibodies interfering with therapy.

Construction of Humanized Genes. Genes encoding chMAb4D5 light chain and heavy chain Fd fragment (VH and CH1 domains) were subcloned together into pUC119 (Vieira, J. & Messing, J., Methods Enzymol. 153:3-11 (1987)) to create pAK1 and simultaneously humanized in a single . step (Fig. 2). Briefly, sets of 6 contiguous oligonucleotides were designed to humanize V_H and V_L (Fig. 1). These oligonucleotides are 28 to 83 nucleotides in length, contain zero to 19 mismatches to the murine antibody template and are constrained to have 8 or 9 perfectly matched residues at promote efficient annealing and ligation of adjacent each end to oligonucleotides. The sets of V_H and V_L humanization oligonucleotides (5 pmol each) were phosphorylated with either ATP or y-³²P-ATP (Carter, P. Methods Enzymol. 154:382-403 (1987)) and separately annealed with 3.7 pmol of pAK1 template in 40 µl 10 mM Tris-HCl (pH 8.0) and 10 mM MgCl₂ by cooling from 100 °C to room temperature over ~30 min. The annealed oligonucleotides were joined by incubation with T4 DNA ligase (12 units; New England Biolabs) in the presence of $2 \mu I 5$ mM ATP and $2 \mu I 0.1$ M DTT

10

15

20



30

for 10 min at 14 °C. After electrophoresis on a 6% acrylamide sequencing gel the assembled oligonucleotides were located by autoradiography and recovered by electroelution. The assembled oligonucleotides (~0.3 pmol each) were simultaneously annealed to 0.15 pmol single-stranded deoxyuridine-containing pAK1 prepared according to Kunkel et al. (Kunkel, T. A. et al., Methods Enzymol. 154:367-382 (1987)) in 10 μ l 40 mM Tris-HCI (pH 7.5) and 16 mM MgCl₂ as above. Heteroduplex DNA was constructed by extending the primers with T7 DNA polymerase and transformed into E. coli BMH 71-18 mut as previously described (Carter, P., in Mutagenesis: A Practical Approach, Chapter 1 (IRL Press, Oxford, UK 1991)). The resultant phagemid DNA pool was enriched first for huV₁ by restriction purification using Xhol and then for huV_H by restriction selection using Stul as described in Carter, P., in Mutagenesis: A Practical Approach, Chapter 1 (IRL Press, Oxford, UK 1991); and in Wells, J. A. et al., Phil. Trans. R. Soc. Lond. A 317:415-423 (1986). Resultant clones containing both huV₁ and huV_H genes were identified by nucleotide sequencing (Sanger, F. et al., Proc. Natl. Acad. Sci. USA 74:5463-5467 (1977)) and designated pAK2. Additional humanized variants were generated by site-directed mutagenesis (Carter, P., in Mutagenesis: A Practical Approach, Chapter 1 (IRL Press, Oxford, UK 1991)). The muMAb4D5 V_L and V_H gene segments in the transient expression vectors described above were then precisely replaced with their humanized versions.

Expression and Purification of MAb4D5 Variants. Appropriate MAb4D5 light and heavy chain cDNA expression vectors were co-transfected into an adenovirus transformed human embryonic kidney cell line, 293 (Graham, F. L. *et al.*, *J. Gen. Virol.* 36:59-72 (1977)) using a high efficiency procedure (Gorman, C. M. *et al.*, *DNA & Prot. Engin. Tech.* 2:3-10 (1990); Gorman, C., in *DNA Cloning*, vol II, pp 143-190 (D. M. Glover, ed., IRL Press, Oxford, UK 1985)). Media were harvested daily for up to 5 days and the cells re-fed with serum free media. Antibodies were recovered from the media and affinity purified on protein A sepharose CL-4B (Pharmacia) as described by the manufacturer. The eluted antibody was buffer-exchanged

10

15

20

5

25

30

80

into phosphate-buffered saline by G25 gel filtration, concentrated by ultrafiltration (Centriprep-30 or Centricon-100, Amicon), sterile-filtered (Millex-GV, Millipore) and stored at 4 °C. The concentration of antibody was determined by using both total immunoglobulin and antigen binding ELISAs. The standard used was huMAb4D5-5, whose concentration had been determined by amino acid composition analysis.

Cell Proliferation Assay. The effect of MAb4D5 variants upon proliferation of the human mammary adenocarcinoma cell line, SK-BR-3, was investigated as previously described (Fendly, B. M. *et al.*, *Cancer Res.* 50:1550-1558 (1990)) using saturating MAb4D5 concentrations.

Affinity Measurements. The antigen binding affinity of MAb4D5 variants was determined using a secreted form of the p185^{HER2} ECD prepared as described in Fendly, B. M. *et al.*, *J. Biol. Resp. Mod.* 9:449-455 (1990). Briefly, antibody and p185^{HER2} ECD were incubated in solution until equilibrium was found to be reached. The concentration of free antibody was then determined by ELISA using immobilized p185^{HER2} ECD and used to calculate affinity (K_d) according to Friguet et al. (Friguet, B. *et al.*, *J. Immunol. Methods* 77:305-319 (1985)).

RESULTS

Humanization of muMAb4D5. The muMAb4D5 V_L and V_H gene segments were first cloned by PCR and sequenced (Fig. 1). The variable genes were then simultaneously humanized by gene conversion mutagenesis using preassembled oligonucleotides (Fig. 2). A 311-mer oligonucleotide containing 39 mismatches to the template directed 24 simultaneous amino acid changes required to humanize muMAb4D5 V_L . Humanization of muMAb4D5 V_H required 32 amino acid changes which were installed with a 361-mer containing 59 mismatches to the muMAb4D5 template. Two out of 8 clones sequenced precisely encode huMAb4D5-5, although one of these clones contained a single nucleotide imperfection. The 6 other clones were essentially humanized but contained a small number of errors: < 3 nucleotide changes and < 1 single nucleotide deletion per kilobase.

10

5

-

20

15

25

30

81

Additional humanized variants (Table 1) were constructed by site-directed mutagenesis of huMAb4D5-5.

Expression levels of huMAb4D5 variants were in the range of 7 to $15 \,\mu$ g/ml as judged by ELISA using immobilized p185^{HER2} ECD. Successive harvests of five 10 cm plates allowed 200 μ g to 500 mg of each variant to be produced in a week. Antibodies affinity purified on protein A gave a single band on a Coomassie blue stained SDS polyacrylamide gel of mobility consistent with the expected M_r of ~150 kDa. Electrophoresis under reducing conditions gave 2 bands consistent with the expected M_r of free heavy (48 kDa) and light (23 kDa) chains (not shown). Amino terminal sequence analysis (10-cycles) gave the mixed sequence expected (see Fig. 1) from an equimolar combination of light and heavy chains (not shown).

huMAb4D5 Variants. In general, the FR residues were chosen from consensus human sequences (Kabat, E. A. et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, MD, 1987)) and CDR residues from muMAb4D5. Additional variants were constructed by replacing selected human residues in huMAb4D5-1 with their muMAb4D5 counterparts. These are V_H residues 71, 73, 78, 93 plus 102 and V_L residues 55 plus 66 identified by our molecular modeling. V_H residue 71 has previously been proposed by others (Tramontano, A. et al., J. Mol. Biol. 215:175-182 (1990)) to be critical to the conformation of V_H-CDR2. Amino acid sequence differences between huMAb4D5 variant molecules are shown in Table 1, together with their p185^{HER2} ECD binding affinity and maximal anti-proliferative activities against SK-BR-3 cells. Very similar K_d values were obtained for binding of MAb4D5 variants to either SK-BR-3 cells (unpublished data) or to p185^{HER2} ECD (Table 1). However, K_d estimates derived from binding of MAb4D5 variants to p185^{HER2} ECD were more reproducible with smaller standard errors and consumed much smaller quantities of antibody than binding measurements with whole cells.

The most potent humanized variant designed by molecular modeling, huMAb4D5-8, contains 5 FR residues from muMAb4D5. This

82

10

15

20

antibody binds the p185^{HER2} ECD 3-fold *more* tightly than does muMAb4D5 itself (Table 1) and has comparable anti-proliferative activity with SK-BR-3 cells (Fig. 3). In contrast, huMAb4D5-1 is the most humanized but least potent muMAb4D5 variant, created by simply installing the muMAb4D5 CDRs into the consensus human sequences. huMAb4D5-1 binds the p185^{HER2} ECD 80-fold *less* tightly than does the murine antibody and has no detectable anti-proliferative activity at the highest antibody concentration investigated (16 μ g/ml).

The anti-proliferative activity of huMAb4D5 variants against p185^{HER2} overexpressing SK-BR-3 cells is not simply correlated with their binding affinity for the p185^{HER2} ECD. For example, installation of three murine residues into the V_H domain of huMAb4D5-2 (D73T, L78A and A93S) to create huMAb4D5-3 does not change the antigen binding affinity but does confer significant anti-proliferative activity (Table 1).

The importance of V_H residue 71 (Tramontano, A. *et al.*, *J. Mol. Biol.* 215:175-182 (1990)) is supported by the observed 5-fold increase in affinity for p185^{HER2} ECD on replacement of R71 in huMAb4D5-1 with the corresponding murine residue, alanine (huMAb4D5-2). In contrast, replacing V_H L78 in huMAb4D5-4 with the murine residue, alanine (huMAb4D5-5), does not significantly change the affinity for the p185^{HER2} ECD or change anti-proliferative activity, suggesting that residue 78 is not of critical functional significance to huMAb4D5 and its ability to interact properly with ° the extracellular domain of p185^{HER2}.

 V_L residue 66 is usually a glycine in human and murine κ chain sequences (Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)) but an arginine occupies this position in the muMAb4D5 k light chain. The side chain of residue 66 is likely to affect the conformation of V_L -CDR1 and V_L -CDR2 and the hairpin turn at 68-69 (Fig. 4). Consistent with the importance of this residue, the mutation V_L G66R (huMAb4D5-3 \rightarrow huMAb4D5-5) increases the affinity for the p185^{HER2} ECD by 4-fold with a concomitant increase in anti-proliferative activity.

· 10

15

20

25

30

From molecular modeling it appears that the tyrosyl side chain of muMAb4D5 V_L residue 55 may either stabilize the conformation of V_H-CDR3 or provide an interaction at the V_L-V_H interface. The latter function may be dependent upon the presence of V_H Y102. In the context of huMAb4D5-5 the mutations V_L E55Y (huMAb4D5-6) and V_H V102Y (huMAb4D5-7) individually increase the affinity for p185^{HER2} ECD by 5-fold and 2-fold respectively, whereas together (huMAb4D5-8) they increase the affinity by 11-fold. This is consistent with either proposed role of V_L Y55 and V_H Y102.

10

15

20

25

30

Secondary Immune Function of huMAb4D5-8. MuMAb4D5 inhibits the growth of human breast tumor cells which overexpress p185^{HER2} (Hudziak, R. M. *et al.*, *Molec. Cell. Biol.* 9:1165-1172 (1989)). The antibody, however, does not offer the possibility of direct tumor cytotoxic effects. This possibility does arise in huMAb4D5-8 as a result of its high affinity (K_d = 0.1 μ M) and its human IgG₁ subtype. Table 2 compares the ADCC mediated by huMAb4D5-8 with muMAb4D5 on a normal lung epithelial cell line, WI-38, which expresses a low level of p185^{HER2} and on SK-BR-3, which expresses a high level of p185^{HER2}. The results demonstrate that: (1) huMAb4D5 has a greatly enhanced ability to carry out ADCC as compared with its murine parent; and (2) that this activity may be selective for cell types which overexpress p185^{HER2}.

DISCUSSION

MuMAb4D5 is potentially useful for human therapy since it is cytostatic towards human breast and ovarian tumor lines overexpressing the *HER2*-encoded p185^{HER2} receptor-like tyrosine kinase. Since both breast and ovarian carcinomas are chronic diseases it is anticipated that the optimal MAb4D5 variant molecule for therapy will have low immunogenicity and will be cytotoxic rather than solely cytostatic in effect. Humanization of muMAb4D5 should accomplish these goals. We have identified 5 different huMAb4D5 variants which bind tightly to p185^{HER2} ECD ($K_d \leq 1$ nM) and which have significant anti-proliferative activity (Table 1). Furthermore

huMAb4D5-8 but not muMAb4D5 mediates ADCC against human tumor cell lines overexpressing p185^{HER2} in the presence of human effector cells (Table 2) as anticipated for a human y1 isotype (Brüggemann, M. *et al.*, *J. Exp. Med.* **166**:1351-1361 (1987); Riechmann, L. *et al.*, *Nature* **332**:323-327 (1988)).

Rapid humanization of huMAb4D5 was facilitated by the gene conversion mutagenesis strategy developed here using long preassembled oligonucleotides. This method requires less than half the amount of synthetic DNA as does total gene synthesis and does not require convenient restriction sites in the target DNA. Our method appears to be simpler and more reliable than a variant protocol recently reported (Rostapshov, V. M. *et al.*, *FEBS Lett.* 249:379-382 (1989)). Transient expression of huMAb4D5 in human embryonic kidney 293 cells permitted the isolation of a few hundred micrograms of huMAb4D5 variants for rapid characterization by growth inhibition and antigen binding affinity assays. Furthermore, different combinations of light and heavy chain were readily tested by co-transfection of corresponding cDNA expression vectors.

The crucial role of molecular modeling in the humanization of muMAb4D5 is illustrated by the designed variant huMAb4D5-8 which binds the p185^{HER2} ECD 250-fold more tightly than the simple CDR loop swap variant, huMAb4D5-1. It has previously been shown that the antigen binding affinity of a humanized antibody can be increased by mutagenesis based upon molecular modelling (Riechmann, L. *et al.*, *Nature* 332:323-327 (1988); Queen, C. *et al.*, *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989)). Here we have extended this earlier work by others with a designed humanized antibody which binds its antigen 3-fold *more* tightly than the-parent-rodent-antibody. While this result is gratifying, assessment of the success of the molecular modeling must await the outcome of X-ray structure determination. From analysis of huMAb4D5 variants (Table 1) it is apparent that their anti-proliferative activity is not a simple function of their binding affinity for p185^{HER2} ECD. For example the huMAb4D5-8 variant binds p185^{HER2} 3-fold more tightly than muMAb4D5 but the humanized variant is

10

5

20

15

25

30

91

slightly less potent in blocking the proliferation of SK-BR-3 cells. Additional huMAb4D5 variants are currently being constructed in an attempt to identify residues triggering the anti-proliferative activity and in an attempt to enhance this activity.

In addition to retaining tight receptor binding and the ability to inhibit cell growth, the huMAb4D5-8 also confers a secondary immune function (ADCC). This allows for direct cytotoxic activity of the humanized molecule in the presence of human effector cells. The apparent selectivity of the cytotoxic activity for cell types which overexpress p185^{HER2} allows for the evolution of a straightforward clinic approach to those human cancers characterized by overexpression of the HER2 protooncogene.

Table 1. p185^{HER2} ECD binding affinity and anti-proliferative activities of MAb4D5 variants

		. V ₁	_i Resi	due*		V _L Res	idue'	: :	
MAb4D5	71	73	78	93	102	. 55	- 66	Rel a ztve	cell
Variant	FR3	FR3	FR3	FR3	CDR3	CDR2	FR3	nM	
proliferatio	n‡								
huMAb4D5-1	R	D	L		v	E .	G	103	 - -
huMAb4D5-2	Ala	D	Ľ	A	v	Ē	G	4.7	10
huMAb4D5-3	Ala	Thr	Ala	Ser	v	E	G	4.4	6
huMAb4D5-4	Ala	Thr	L	Ser	V	E	Arg	0.82	5
nuMAb4D5-5	Ala	Thr	Ala	Ser	V	E	Arg	1.1	4
nuMAb4D5-6	Ala	Thr	Ala	Ser	v	Tyr	Arg	0.22	5
nuMAb4D5-7	Ala	Thr	Ala	Ser	Tyr	E	Arg	0.62	5
nuMAb4D5-8	Ala	Thr	Ala	Ser	Tyr	Tyr	Arg	0.10	5
nuMAb4D5	Ala	Thr	Ala	Ser	Tyr	Tyr	Arg	0.30	3

* Human and murine residues are shown in one letter and three letter amino acid code respectively.

[†] K_d values for the p185^{HER2} ECD were determined using the method of Friguet *et al.* (43) and the standard error of each estimate is $\leq \pm 10\%$.

[‡] Proliferation of SK-BR-3 cells incubated for 96 hr with MAb4D5 variants shown as a percentage of the untreated control as described (Hudziak, R. M. *et al., Molec. Cell. Biol.* 9:1165-1172 (1989)). Data represent the maximal anti-proliferative effect for each variant (see Fig. 3A) calculated as the mean of triplicate determinations at a MAb4D5 concentration

87

10

5

-

15

20

25





of 8 μ g/ml. Data are all taken from the same experiment with an estimated standard error of

≤ ± 15%.

88

	WI	-38*	SK-BR-3		
Effector	:Target				
ratio [†]	muMAb4D5	huMAb4D5-8	muMAb4D5	huMAb4D5-8	
		1			
A.‡	25:1	<1.0	9.3	7.5 40.6	
	12.5:1	<1.0	11,1	4.7 36.8	
	6.25:1	<1.0	8.9	0.9 35.2	
	3.13:1	<1.0	8.5	4.6 19.6	
в.	25:1	<1.0	3.1	6.1 33.4	
	12.5:1	<1.0	1.7	5.5 26.2	
	6.25:1	1.3	2.2	2.0 21.0	
	3,13:1	<1.0	0.8	2.4 13.4	

Table 2. Selectivity of antibody dependent tumor cell cytotoxicity mediated by huMAb4D5-8

* Sensitivity to ADCC of two human cell lines (WI-38, normal lung epithelium; and SK-BR-3, human breast tumor cell line) are compared. WI-38 expresses a low level of p185^{HER2} (0.6 pg per μ g cell protein) and SK-BR-3 expresses a high level of p185^{HER2} (64 pg p185^{HER2} per μ g cell protein), as determined by ELISA (Fendly *et al.*, *J. Biol. Resp. Mod.* 9:449-455 (1990)). † ADCC assays were carried out as described in Brüggemann *et al.*, *J. Exp. Med.* **166**:1351-1361 (1987). Effector to target ratios were of IL-2 activated human peripheral blood lymphocytes to either WI-38 fibroblasts or SK-BR-3 tumor cells in 96-well microtiter plates for 4 hours at 37 °C. Values given represent percent specific cell lysis as determined by ⁵¹Cr release. Estimated standard error in these quadruplicate determinations was $\leq \pm 10\%$.

⁺ Monoclonal antibody concentrations used were 0.1 μ g/ml (A) and 0.1 μ g/ml (B).

15.

10

25

20

89

<u>95</u>

EXAMPLE 2, Schematic Method for Humanizing an Antibody Sequence

This example illustrates one stepwise elaboration of the methods for creating a humanized sequence described above. It will be understood that not all of these steps are essential to the claimed invention, and that steps may be taken in different order.

- ascertain a consensus human variable domain amino acid sequence and prepare from it a consensus structural model.
- prepare model of import (the non-human domain to be humanized) variable domain sequences and note structural differences with respect to consensus human model.
- 3. identify CDR sequences in human and in import, both by using Kabat (supra, 1987) and crystal structure criteria. If there is any difference in CDR identity from the different criteria, use of crystal structure definition of the CDR, but retain the Kabat residues as important framework residues to import.

 substitute import CDR sequences for human CDR sequences to obtain initial "humanized" sequence.

 compare import non-CDR variable domain sequence to the humanized sequence and note divergences.

 Proceed through the following analysis for each amino acid residue where the import diverges from the humanized.

- a. If the humanized residue represents a residue which is generally highly conserved across all species, use the residue in the humanized sequence. If the residue is not conserved across all species, proceed with the analysis described in 6b.
- b. If the residue is not generally conserved across all species, ask if the residue is generally conserved in humans.
 - i. If the residue is generally conserved in humans but the import residue differs, examine the structural models of the

10

15

20

25

30

90

import and human sequences and determine if the import residue would be likely to affect the binding or biological activity of the CDRs by considering 1) could it bind antigen directly and 2) could it affect the conformation of the CDR. If the conclusion is that an affect on the CDRs is likely, substitute the import residue. If the conclusion is that a CDR affect is unlikely, leave the humanized residue unchanged.

5

10

15

20

25

30

ii.

a)

If the residue is also not generally conserved in humans, examine the structural models of the import and human sequences and determine if the import residue would be likely to affect the binding or biological activity of the CDRs be considering 1) could it bind antigen directly and 2) could it affect the conformation of the CDR. If the conclusion is that an affect on the CDRs is likely, substitute the import residue. If the conclusion is that a CDR affect is unlikely, proceed to the next step.

Examine the structural models of the import and human sequences and determine if the residue is exposed on the surface of the domain or is buried within. If the residue is exposed, use the residue in the humanized sequence. If the residue is buried, proceed to the next step.

(i) Examine the structural models of the import and human sequences and determine if the residue is likely to affect the V_L - V_H interface. Residues involved with the interface include: 34L, 36L, 38L, 43L, 33L, 36L, 85L, 87L, 89L, 91L, 96L, 98L, 35H, 37H, 39H, 43H, 45H, 47H, 60H, 91H, 93H, 95H, 100H, and 103H. If no effect is likely, use the residue in the humanized sequence. If some affect is likely, substitute the

91

import residue...

7. Search the import sequence, the consensus sequence and the humanized sequence for glycosylation sites outside the CDRs, and determine if this glycosylation site is likely to have any affect on antigen binding and/or biological activity. If no effect is likely, use the human sequence at that site; if some affect is likely, eliminate the glycosylation site or use the import sequence at that site.

5

10

15

20

25

30

b.

8. After completing the above analysis, determine the planned humanized sequence and prepare and test a sample. If the sample does not bind well to the target antigen, examine the particular residues listed below, regardless of the question of residue identity between the import and humanized residues.

 Examine particular peripheral (non-CDR) variable domain residues that may, due to their position, possibly interact directly with a macromolecular antigen, including the following residues (where the * indicates residues which have been found to interact with antigen based on crystal structures):

i. Variable light domain: 36, 46, 49°, 63-70

ii. Variable heavy domain: 2, 47°, 68, 70, 73-76.

Examine particular variable domain residues which could interact with, or otherwise affect, the conformation of variable domain CDRs, including the following (not including CDR residues themselves, since it is assumed that, because the CDRs interact with one another, any residue in one CDR could potentially affect the conformation of another CDR residue) (L = LIGHT, H = HEAVY, residues appearing in **bold** are indicated to be structurally important according the Chothia *et al.*, Nature 342:877 (1989), and residues appearing in *italic* were altered during humanization by Queen *et al.* (PDL), Proc. Natl. Acad. Sci. USA 86:10029 (1989) and Proc. Natl. Acad. Sci. USA 88:2869 (1991).):

i. Variable light domain:

92

- a) CDR-1 (residues 24L-34L): 2L, 4L, 66L-69L, 71L
- b) CDR-2 (residues 50L-56L): 35L, 46L, 47L, 48L, 49L,
 - 58L, 62L, 64L-66L, 71L, 73L
- c) CDR-3 (residues 89L-97L): 2L, 4L, 36L, 98L, 37H, 45H, 47H, 58H, 60H

ii. Variable heavy domain:

- a) CDR-1 (residues 26H-35H): 2H, 4H, 24H, 36H, 71H, 73H, 76H, 78H, 92H, **94H**
- b) CDR-2 (residues 50H-55H): 49H, 69H, 69H, 71H, 73H, 78H
- c) CDR-3 (residues 95H-102H): examine all residues as possible interaction partners with this loop, because this loop varies in size and conformation much more than the other CDRs.
- If after step 8 the humanized variable domain still is lacking in desired binding, repeat step 8. In addition, re-investigate any buried residues which might affect the $V_L V_H$ interface (but which would not directly affect CDR conformation). Additionally, evaluate the accessibility of non-CDR residues to solvent.

15

20

9.

5

10

	SEQUENCE LISTING
	(1) GENERAL INFORMATION:
	(i) APPLICANT: Carter, Paul J. Presta, Leonard G.
	(ii) TITLE OF INVENTION: Immunoglobulin Variants
	(iii) NUMBER OF SEQUENCES: 10
· ·	 (iv) CORRESPONDENCE ADDRESS: (A) ADDRESSEE: Genentech, Inc. (B) STREET: 460 Point San Bruno Blvd (C) CITY: South San Francisco (D) STATE: California (E) COUNTRY: USA (F) ZIP: 94080
	 (v) COMPUTER READABLE FORM: (A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy di (B) COMPUTER: IBM PC compatible (C) OPERATING SYSTEM: PC-DOS/MS-DOS (D) SOFTWARE: patin (Genentech)
	 (vi) CURRENT APPLICATION DATA: (A) APPLICATION NUMBER: (B) FILING DATE: 14-June-1991 (C) CLASSIFICATION:
	(vii) PRIOR APPLICATION DATA: (A) APPLICATION NUMBER: (B) FILING DATE:
	(viii) ATTORNEY/AGENT INFORMATION: (A) NAME: Adler, Carolyn R. (B) REGISTRATION NUMBER: 32,324 (C) REFERENCE/DOCKET NUMBER: 709
	(ix) TELECOMMUNICATION INFORMATION: (A) TELEPHONE: 415/266-2614 (B) TELEFAX: 415/952-9881 (C) TELEX: 910/371-7168
	(2) INFORMATION FOR SEQ ID NO:1:
	(i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids

. 20

.

disk

(B) TYPE: amino acid (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Asp lie Gin Met Thr Gin Ser Pro Ser Ser Leu Ser Ala Ser Val 1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gin Asp Val Asn 20 25 30

Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys354045

Leu Leu Ile Tyr Ser Ala Ser Phe Leu Glu Ser Gly Val Pro Ser 50 55 60

Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile 65 70 75

Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln 80 85 90

His Tyr Thr Thr Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu 95 100 105

lle Lys Arg Thr 109

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly151015

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys 20 25 30

45

5

. 10

15

20

25

30

35

40

Asp Thr Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu 35 40 45

Glu Trp Val Ala Arg lle Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr

50	55	60	
Ala Asp Ser Val Lys Gl	y Arg Phe T	hr lle Ser Ala Asp Thr Se	r
65	70	75	
Lys Asn Thr Ala Tyr Le	eu Gln Met A	Asn Ser Leu Arg Ala Glu /	Asp
80	85	90	

Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr95100105

Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser 110 115 120

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 109 amino acids
 (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

5

10

15

20

25

30

35

40

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Asp lie Gin Met Thr Gin Ser Pro Ser Ser Leu Ser Ala Ser Val 1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser 20 25 30

Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys 35 40 45

Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser 50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile 65 70 75

Ser Ser Leu Gin Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gin Gin 80 85 90

Tyr Asn Ser Leu Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu 95 100 105

Ile Lys Arg Thr 109

	-	(2) INFORMATION FOR SEQ ID NO:4:		
5		 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear 		
	•	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:		
10		Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly 1 5 10 15	• .	
· .		Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser 20 25 30	· .	
15	. •	Asp Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu 35 40 45		
20	• · · ·	Glu Trp Val Ala Val IIe Ser Glu Asn Gly Gly Tyr Thr Arg Tyr 50 55 60		. ·
		Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser 65 70 75		
25		Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp 80 85 90		
-		Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr 95 100 105		
30		Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser 110 115 120		
35		(2) INFORMATION FOR SEQ ID NO:5:		
40		 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear 		
		(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:		
45		Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val 1 5 10 15	·	
		Gly Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn 20 25 30		, ,

,	
	Thr Ala Val Ala Trp Tyr Gin Gin Lys Pro Gly His Ser Pro Lys 35 40 45
	Leu Leu IIe Tyr Ser Ala Ser Phe Arg Tyr Thr Gly Val Pro Asp 50 55 60
•	Arg Phe Thr Gly Asn Arg Ser Gly Thr Asp Phe Thr Phe Thr Ile 65 70 75
	Ser Ser Val Gin Ala Giu Asp Leu Ala Val Tyr Tyr Cys Gin Gin 80 85 90
	His Tyr Thr Thr Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu 95 100 105
	lle Lys Arg Ala 109
	(2) INFORMATION FOR SEQ ID NO:6:
	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
	Glu Val Gin Leu Gin Gin Ser Giy Pro Giu Leu Val Lys Pro Giy 1 5 10 15
	Ala Ser Leu Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn lie Lys 20 25 30
	Asp Thr Tyr Ile His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu 35 40 45
	Glu Trp Ile Gly Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr 50 55 60
	Asp Pro Lys Phe Gin Asp Lys Ala Thr Ile Thr Ala Asp Thr Ser 65 70 75
	Ser Asn Thr Ala Tyr Leu Gln Val Ser Arg Leu Thr Ser Glu Asp 80 85 90
	Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr 95 100 105

-

	Ala Met Asp Tyr Trp Gly Gln Gly Ala Ser Val Thr Val Ser Ser 110 115 120
5	(2) INFORMATION FOR SEQ ID NO:7:
10	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
15	TCCGATATCC AGCTGACCCA GTCTCCA 27
20	(2) INFORMATION FOR SEQ ID NO:8:
· .	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 31 bases (B) TYPE: nucleic acid
25	(C) STRANDEDNESS: single (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
30	GTTTGATCTC CAGCTTGGTA CCXXCDCCGA A 31
35	(2) INFORMATION FOR SEQ ID NO:9:
	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 22 bases (B) TYPE: nucleic acid (C) CTRANDEDNESS: sincle
40	(C) STRANDEDNESS: single (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
.45	AGGTXXAXCT GCAGXAGTCX GG 22

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 bases
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

10

15

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TGAGGAGACG GTGACCGTGG TCCCTTGGCC CCAG 34

100

CLAIMS

WE CLAIM:

a.

b.

с.

d. .

e.

f.

g.

2.

1.

5

10

15

20

25

A method for making a humanized antibody comprising amino acid sequence of a non-human, import antibody and a human antibody, comprising the steps of:

> obtaining the amino acid sequences of at least a portion of an import variable domain and of a consensus human variable domain;

identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human amino variable domain sequences;

substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;

identifying import antibody FR residues in the aligned FR sequences that are non-homologous to the corresponding consensus antibody residues;

determining if the non-homologous import amino acid residue is reasonably expected to have at least one of the following effects:

non-covalently binds antigen directly, 1.

interacts with a CDR; or 2.

participates in the $V_{\rm L}$ - $V_{\rm H}$ interface; and 3. for any non-homologous import antibody amino acid residue which is reasonably expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence.

30

The method of claim 1, having an additional step of determining if N Carter da 1715,272 114/91

101

any such non-homologous residues are exposed on the surface of the domain or buried within it, and if the residue is exposed, retaining the consensus residue.

The method of claim 1, having the additional steps of searching the import variable domain sequence for glycosylation sites, determining if any such glycosylation site is reasonably expected to affect the antigen binding or affinity of the antibody, and if so, substituting the glycosylation site into the consensus sequence.

The method of claim 1, having the additional steps of searching the consensus variable domain sequence for glycosylation sites which are not present at the corresponding amino acid in the import sequence, and if the glycosylation site is not present in the import sequence, substituting the import amino acid residues for the amino acid residues comprising the consensus glycosylation site.

The method of claim 1, having an additional step which comprises aligning import antibody and consensus antibody FR sequences, identifying import antibody FR residues which are non-homologous with the aligned consensus FR sequence, and for each such nonhomologous import antibody FR residue, determining if the corresponding consensus antibody residue represents a residue which is highly conserved across all species at that site, and if it is so conserved, preparing a humanized antibody which comprises the consensus antibody amino acid residue at that site.

6.

4.

5.

5

10

15

20

25

30

The method of claim 1, wherein the corresponding consensus antibody residues are selected from the group consisting of 4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H,

76H, 78H, 91H, 92H, 93H, and 103H.

A method comprising providing at least a portion of an import, nonhuman antibody variable domain amino acid sequence having a CDR and a FR, obtaining the amino acid sequence of at least a portion of a consensus human antibody variable domain having a CDR and a FR, substituting the non-human CDR for the human CDR in the consensus human antibody variable domain, and then substituting an amino acid residue for the consensus amino acid residue at at least one of the following sites:

4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

The method of claim 7, wherein the substituted residue is the residue found at the corresponding location of the non-human antibody.

A humanized antibody variable domain having a non-human CDR incorporated into a human antibody variable domain, wherein the improvement comprises substituting an amino acid residue for the human residue at a site selected from the group consisting of: 4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

10.

7.

8.

9.

5-

10

15

20

25

30

The humanized antibody variable domain of claim 9, wherein the substituted residue is the residue found at the corresponding location of the non-human antibody from which the non-human CDR was obtained.



11.

13.

15.

b.



The humanized antibody variable domain of claim 9, wherein no human FR residue other than those set forth in the group has been substituted.

12. A polypeptide comprising the amino acid sequence: DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAP KLLIYSASFLESGVPSRFSGSRSGTDFTLTISSLQPEDFATYYCQQHY TTPPTFGQGTKVEIKRT

A polypeptide comprising the sequence: EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLE WVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAEDT AVYYCSRWGGDGFYAMDVWGQGTLVTVSS

A computer comprising the sequence data of the following amino acid sequence:

a. DIQMTQSPSSLSASVGDRVTITCRASQDVSSYLAWYQQ KPGKAPKLLIYAASSLESGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQQYNSLPYTFGQGTKVEIKRT, or

EVOLVESGGGLVOPGGSLRLSCAASGFTFSDYAMSWVR OAPGKGLEWVAVISENGGYTRYADSVKGRFTISADTSKN TAYLOMNSLRAEDTAVYYCSRWGGDGFYAMDVWGOG TLVTVSS

A computer representation of the following amino acid sequence: a. DIOMTOSPSSLSASVGDRVTITCRASODVSSYLAWYOO KPGKAPKLLIYAASSLESGVPSRFSGSGSGTDFTLTISSLO PEDFATYYCOOYNSLPYTFGOGTKVEIKRT, or b. EVOLVESGGGLVOPGGSLRLSCAASGFTFSDYAMSWVR

QAPGKGLEWVAVISENGGYTRYADSVKGRFTISADTSKN TAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDVWGQG

104

.

30

25

5-

10

15

20

TLVTVSS

- A method comprising storing a computer representation of the following amino acid sequence:
 - a. DIQMTQSPSSLSASVGDRVTITCRASQDVSSYLAWYQQ KPGKAPKLLIYAASSLESGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQQYNSLPYTFGQGTKVEIKRT, or
 - b.

10

16.

5

10

PEDFATYYCQQYNSLPYTFGQGTKVEIKRT, or EVQLVESGGGLVQPGGSLRLSCAASGFTFSDYAMSWVR QAPGKGLEWVAVISENGGYTRYADSVKGRFTISADTSKN TAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDVWGQG TLVTVSS



Abstract

Variant immunoglobulins, particularly humanized antibody polypeptides are provided, along with methods for their preparation and use. Consensus immunoglobulin sequences and structural models are also provided.

5

e

EXPRESS MAIL NO. B59937585 MAILED: 14 JUNE 1991

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

Docket No. 709

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

IMMUNOGLOBULIN VARIANTS

the specification of which (check one) \underline{x} is attached hereto or _ was filed on as Application Serial No. and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I hereby state that any Sequence Listing submitted with this application is submitted in paper copy and a computerreadable diskette, and that the content of the paper and computer readable copies are the same.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate have a filing date before that of the application on which priority is claimed:

Prior Foreign	Application(s)		Priority	Claimed	
,			Yes	No	
Number	Country	Day/Month/Year Filed	1		

I hereby claim the benefit under Title 35, United States Code, §120 of any United States applications(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Ser. No.Filing DateStatus: Patented, Pending, Abandoned

Filing Date

Status: Patented, Pending, Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

30/ Carolyn R. Adler - Reg. No. 32,324 Robert H. Benson - Reg. No. 30,446 Walter E. Buting - Reg. No. 23,092 Ginger R. Dreger - Reg. No. 33,055 Debbie Glaister - Reg. No. 33,888 Janet E. Hasak - Reg. No. 28,616

Application Ser. No.

Max D. Hensley- Reg. No. 27,043Dennis G. Kleid- Reg. No. 32,037Nancy Olseki- Reg. No. 34,688Stephen Raines- Reg. No. 25,912Daryl B. Winter- Reg. No. 32,637





Send correspondence to 401

(40) Genentech, Inc.
(402 Attn: Carolyn R. Adler
(70) 460 Point San Bruno Boulevard
(702 South San Francisco, CA 94080 Telephone: (415) 266-2614

I hereby declare that all statements made herein of my own knowledge and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that willful false statements may jeopardize the validity of the application or any patent issued thereon.

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from his foreign patent agent as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

Full name of sole or first inventor	
Paul J/Carter 40100	
Inventor's signature	Date
Residence 2074 18th Avenue San Francisco, CA 94116	· · ·
Citizenship United Kingdom	
Fost Office Address 460 Point San Bruno Boulevard South San Francisco, CA 94080	
Full name of second joint inventor, if any	
Leonard G Presta 40200	
Second Inventor's signature	Date
Residence 1900 Gough Street, #206 San Francisco, CA 94109	
Citizenship United States of America	
Post Office Address 460 Point San Bruno Boulevard South San Francisco, CA 94080	
Full name of third joint inventor, if any	
Third Inventor's signature	Date
Residence	
Citizenship	•
Post Office Address 460 Point San Bruno Boulevard South San Francisco, CA 94080	

Page 2 of 2

				U.S.	PATENT A	PPLICATION	
	SERVAL NUP	BER		ILING DATE	CLASS	GROUP ART UN	IT
	07/7	5,272		06/14/91	530	183	
	PAUL	J. CARTER, S	AN FRANCISCO	, CA; LEONARD	G. PRESTA, SA	N FRANCISCO, CA.	
	APPL					· · · ·	
		ITINUING DATA FIED	****	****			
	·					· · · ·	
(• •		·		,	
	FOR VERI	EIGN/PCT APP	LICATIONS*	*****			
	VERI						У. С. К.
	FORFI	GN FILING LI	CENSE GRANTE	D 08/03/91			
	STATE OR	SHEETS	TOTAL	INDEPENDENT	FILING FEE	ATTURNEY DOCKET N	
	COUNTRY	DRAWING	CLAIMS 16	CLAIMS	RECEIVED	709	
		5 NTECH, INC.			\$1,050.00		
	ATTN 460	: CAROLYN R. POINT SAN BR H SAN FRANCI	UNO BLVD.	0			
		NOGLOBULIN V	ARIANTS	<u></u>			
	F.						
	This is to	certify that	annexed hereto	is a true copy	y from the record	is of the United St	ates /
	By authori	tv of the	•		iginally filed w	nich is identified a	LOOVE. /
	Date	R OF PATENTS	AND TRADEMARKS	Certifying Off	icer		
· .			-				
	•						

1715272

930.00CH

 $(\cdot)_{ij}$

7

PATENT APPLICATION SERIAL NO.

.

4

L.

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE FEE RECORD SHEET

07-0630 030 101

116

06/27/91 07715272

P 30192

PTO-1556 (5/87)

	AIMS AS FILED - (Column 1) UMBER FILED	PARTI	uma 2)		715. L entity	<u>Э7</u> ОЯ	ン OTHER T SMALL E	
FOR N BASIC FEE	(Column 1) UMBER FILED						OTHER T	
BASIC FEE		NUMBER		RATE	FEE	1		
	/ ¥	s 20 = *			ļ		RATE	FEE
TOTAL CLAIMS	/ ¥	s 20 = *			\$ 315.00	OR		\$ 630.00
	8 minu	· · · ·		x \$10) ==	OR	x \$20 ≖	
INDEPENDENT CLAIMS	0	us 3 = * 🕹	5	x 30		OR	x 60 ≖	300
MULTIPLE DEPENDENT CLA	IM PRESENT	· .		+ 100) 📼	OR	+ 200 =	
 If the difference in column 1 is less t 	hen zero, enter "0" in c	olumn 2	·	TOTA	L	OR	TOTAL	930
CI (Column	LAIMS AS AME	NDED - PART	(Column 3)	SMAL		OR	OTHER T SMALL E	
CLAIM REMAIN AFTEF AMENDM	ING R	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE	ADDI- TIONAL FEE		RATE	ADDI- TIONAL FEE
Total	Minus	**	=	x \$10	-	OR	x \$20 🛥	
AFTEF AMENDM Total Independent	Minus	***	-	x 30	- .	OR	x 60=	-
FIRST PRESENTATION	OF MULTIPLE DE	PENDENT CLAIM		+ 100	-	ÖR	+ 200 =	
(Column	i)	(Column 2)	(Column 3)	TOTA ADDIT. FEI		OR	TOTAL DDIT: FEE	
CLAIM REMAIN AFTER AMENDM Total Independent	S ING R	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE	ADDI- TIONAL FEE		RATE	ADDI- TIONAL FEE
Total *	Minus	**		x \$10	a	OR OR	. x \$20 =	
Independent *	Minus	***		x 30	=	OR	x 60 ≖	
FIRST PRESENTATION	N OF MULTIPLE DE	PENDENT CLAIM	· •	+ 100	-	OR	+ 200 =	
(Column	1)	(Column 2)	(Column 3)	TOTA ADDIT. FE		OR	TOTAL DDIT. FEE	
CLAIM REMAIN AFTEF AMENDM	ING R	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE	ADDI- TIONAL FEE		RATE	ADDI- TIONAL FEE
Total *	Minus	*	æ .	x \$10	Ē	OR OR	x \$20 -	
AFTEF AMENDM Total * Independent *	Minus	***	Ξ	x 30	3	OR OR	x 60 =	
FIRST PRESENTATION				+ 100	K24	OR	+ 200 =	
If the entry in column 1 is less If the "Highest Number Previou If the "Highest Number Previou The "Highest Number Previou FORM PTO-875	usly Paid For" IN TH Isly Paid For" IN TH	IS SPACE is less the IS SPACE is less the IS Independent) is t	han 20, enter *20* han 3, enter *3*. he highest numbe	er found in th	E	ox in c		ONGAEDCE

INTERNATIONAL SEARCH REPORT

PCT

Applicant's or agent's file reference 709P1		of Transmittal of International Search Report (220) as well as, where applicable, item 5 below.
International application No.	International filing date(day/month/year)	(Earliest) Priority Date (day/month/year)
PCT/US 92/05126	15/06/92	14/06/91
Applicant		
GENENTECH, INC. et al.		
This international search report has been according to Article 18. A copy is being t	prepared by this International Searching Auth ransmitted to the International Bureau.	nority and is transmitted to the applicant
This international search report consists $\overline{\mathbf{X}}$ It is also accompanied by a cop	of a total of <u>4</u> sheets. y of each prior art document cited in this repo	or t .
1. X Certain claims were found unsea	urchable (see Box I).	
2. Unity of invention is lacking (see	e Box II).	
3. The international application or international search was carried	ntains disclosure of a nucleotide and/or amino l out on the basis of the sequence listing	acid sequence listing and the
file	d with the international application.	
fur	nished by the applicant separately from the in	
	but not accompanied by a statement to matter going beyond the disclosure in th	the effect that it did not include ie international application as filed.
	anscribed by this Authority	
4. With regard to the title, the	text is approved as submitted by the applicar	nt.
X the	text has been established by this Authority to	o read as follows:
METHOD FOR MAKING HUM	ANIZED ANTIBODIES.	
5. With regard to the abstract,	,	
	e text is approved as submitted by the applicat	nt.
	e text has been established, according to Rule ox III. The applicant may, within one month f arch report, submit comments to this Authori	38.2(b), by this Authority as it appears in rom the date of mailing of this international
	an a	
6. The figure of the drawings to be put	blished with the abstract is: suggested by the applicant.	None of the figures.
	cause the applicant failed to suggest a figure.	
	cause this figure better characterizes the inver	uion.
	,	

Form PCT/ISA/210 (first sheet) (July 1992)

ار بله دسر د

	International application No.
INTERNATIONAL STOCH REPORT	PCT 92/ 05126
ox I Observations, where certain claims were found unsearchable (Contin	nuation of item 1 of first sheet)
his international search report has not been established in respect of certain claims	under Arneie 1 (12 As) for the following reasons
X Claims Nos.: 17-18	uthority namely:
Claims Nos: 17-18 because they relate to subject matter not required to be searched by this A see PCT-Rule 39.1(iv)	
	,
Claims Nos.: because they relate to parts of the international application that do not con an extent that no meaningful international search can be carried out, speci	mply with the prescribed requirements to such fically:
3. Claims Nos.:	the second and third sentences of Rule 6.4(a).
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with	ale actoria and and a serientees of the origin
Box II Observations where unity of invention is lacking (Continuation of i	item 2 of first sheet)
This International Searching Authority found multiple inventions in this internation	nal application, as follows:
	-
, ,	· · · · ·
1. As all required additional search fees were timely paid by the applicant, t searchable claims.	his international search report covers all
searchable claims.	
 As all required additional search fees were timely paid by the applicant, the searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. 	
2. As all searchable claims could be searches without effort justifying an ad	
2. As all searchable claims could be searches without effort justifying an ad	lditional fee, this Authority did not invite payment the applicant, this international search report
 2. As all searchable claims could be searches without effort justifying an ad of any additional fee. 	lditional fee, this Authority did not invite payment the applicant, this international search report
 2. As all searchable claims could be searches without effort justifying an ad of any additional fee. 	lditional fee, this Authority did not invite payment the applicant, this international search report
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. As only some of the required additional search fees were timely paid by covers only those claims for which fees were paid, specifically claims No 	lditional fee, this Authority did not invite payment the applicant, this international search report 25.:
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. 	lditional fee, this Authority did not invite payment the applicant, this international search report 25.: onsequently, this international search report is
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. As only some of the required additional search fees were timely paid by covers only those claims for which fees were paid, specifically claims No 	lditional fee, this Authority did not invite payment the applicant, this international search report 25.: onsequently, this international search report is
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. As only some of the required additional search fees were timely paid by covers only those claims for which fees were paid, specifically claims No 	lditional fee, this Authority did not invite payment the applicant, this international search report 25.: onsequently, this international search report is
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. As only some of the required additional search fees were timely paid by covers only those claims for which fees were paid, specifically claims No No required additional search fees were timely paid by the applicant. Correstricted to the invention first mentioned in the claims; it is covered by 	lditional fee, this Authority did not invite payment the applicant, this international search report 25.: onsequently, this international search report is
 searchable claims. As all searchable claims could be searches without effort justifying an ad of any additional fee. As only some of the required additional search fees were timely paid by covers only those claims for which fees were paid, specifically claims No No required additional search fees were timely paid by the applicant. Correstricted to the invention first mentioned in the claims, it is covered by Remark on Protest 	Iditional fee, this Authority did not invite payment the applicant, this international search report ss.: onsequently, this international search report is claims Nos.:

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)

	`	INTERN	SEARCH REPORT	US 92/05126
	FICATION OF SUBJE	CT MATTER (if several classification	symbols apply, indicate all) ⁶	¥7 -
		Classification (IPC) or to both National		
· ·	. 5 C12N15/1 G06F15/0	3; C12P21/08;		2N5/10
D. FTELDS	SEARCHED			
		Minimum Docus	mentation Searched?	
Classificat	tion System		Classification Symbols	
Int.Cl	. 5	CO7K ; C12N ;	G06F	
		Documentation Searched oth to the Extent that such Document	er than Minimum Documentation s are included in the Fields Searched ⁸	
III. DOCU		ED TO BE RELEVANT ⁹		
Category °	Citation of D	ocument, 11 with indication, where approp	priate, of the relevant passages 12	Relevant to Claim No.13
Y	vol. 21	OF MOLECULAR BIOLOGY 5, 1990, ACADEMIC PRES 75 - 182	S	1-12,15
	Arthur determi conform region immunog cited i	ano, Anna; Chothia, Cy M. 'Framework residue nant of the position a ation of the second hy in the VH domains of lobulins' n the application whole document, espec ph 7	71 is a major and ypervariable	
Y	26 July	007 861 (PROTEIN DESIG 1990 es 1-6; 9-25	GN LABS, INC.)	1-12,15
		225 Atto wak		
1			- / ,	
		· · ·		
"A" do re "E" es fi "L" do wh ch "O" do 0 "P" do	ponsidered to be of parti partier document but pub- ling date ocument which may thrushich is cited to establisi- tation or other special in ocument referring to an ther means	eneral state of the art which is not cular relevance ilshed on or after the international ow doubts on priority claim(s) or h the publication date of another reason (as specified) a oral disclosure, use, exhibition or r to the international filling date but	 "T" later document published after the inter- or priority date and not in conflict with cited to understand the principle or the invention "X" document of particular relevance; the cl cannot be considered novel or cannot be involve an inventive step "Y" document of particular relevance; the cl cannot be considered to involve an inve document is combined with one or more ments, such combination being obvious in the art. "&" document member of the same patent from 	the application but ry underlying the aimed invention considered to aimed invention tive step when the other such docu- to a person skilled
1	IFICATION			
Date of the		the International Search BER 1992	Date of Malling of this International Se 0 2. 11. 92	arch Report
Internation	EUROPH	EAN PATENT OFFICE	Signature of Authorized Officer NAUCHE S.A.	for for
Form PCT/LS	A/210 (second sheet) (Junn	ury 1945)		"See notes on accompanying shor

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO. US SA

9205126

61838

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 07/10/92

Patent document cited in search report	Publication date	I	Patent family member(s)	Publication date
WO-A-9007861	26-07-90	AU-A- CA-A- EP-A-	5153290 2006865 0451216	13-08-90 28-06-90 16-10-91

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

	International Application No	
III. DOCUME	NTS CONSIDERED TO BE BELEVANT (CONTINUED FROM THE SECOND SHEET)	Relevant to Claim No.
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	
Y	NATURE. vol. 342, December 1989, LONDON GB pages 877 - 883 Chothia, Cyrus; Lesk, Arthur M.; Tramontano, Anna; Levitt, Michael;	1-12,15
	Smith-Gill, Sandra J.; Air, Gillian; Sheriff, Steven; Padlan, 'Conformations of immunoglobulin hypervariable region' cited in the application See the whole document, especially 'Discussion'	
P ,X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA. vol. 89, May 1992, WASHINGTON US pages 4285 - 4289 Carter, Paul et al. 'Humanization of an	1-15
	anti-p185HER2 antibody for human cancer therapy. ¹ see the whole document	
	: :	
		•
,		

SCORE Placeholder Sheet for IFW Content

Application Number: 07715272

Document Date: 06/14/1991

The presence of this form in the IFW record indicates that the following document type was received in paper and is scanned and stored in the SCORE database.

Drawings

Images of the original documents are scanned in gray scale or color and stored in SCORE. Bi-tonal images are also stored in IFW. Defects visible in both IFW and SCORE are indicative of defects in the original paper documents.

To access the documents in the SCORE database, refer to instructions developed by SIRA.

At the time of document entry (noted above):

- Examiners may access SCORE content via the eDAN interface.
- Other USPTO employees can bookmark the current SCORE URL (http://es/ScoreAccessWeb/).
- External customers may access SCORE content via the Public and Private PAIR interfaces.


Form Revision Date: December 8, 2006

Best Available Copy

· .					
			The second s	- A. 63	
	• •		11000000000000000000000000000000000000	- Channel	
		NUMB			
TILING DAT	5 33	1901aas' 387	aroup ary unit	Feise	2
TER. SAN F	ANCISCO, 'CA; L	EONARD G. PF	ESTA- SAN FR	ANCISCO- CA.	
	• • ·	1) 		inter State internationalista	
VERIFIED	****	***			
Noute	· .	4	16-1 		
	1			i	
			•		
FOREIGN/PCT APPLICA	TIONS******	***			
VERIFIED					
NEI4 0	• •	· · ·		· •	
		2/01		1. 1.	
FOREIGN FILING LICENS	E GRANIED USIO	SHEETS TOTAL DRWGS. CLAIMS	INDEP. PILING PE	E ATTORNEY'S	
Poreign priority'scalmed U yes Ar 36 USC 319 conditions met U yes Vir Vertilied and Acknowledged Exeminar's In	FILED	5 16	3 51.050		<u></u>
ATTN: CAROLYN R. ADI	LER				
SOUTH SAN FRANCISCO	, CA 94080				
IMMUNOGLOBULIN VARI	ANTS		<u>.</u>	· · ·	
	1	U.S. DI	EPT. OF COMM. PAL & T	A Office PTD-438L (rev.	10-78)
	and the second second second				
IRTS OF APPLICATION		the state of the			
AED SEPARATELY		FOR ISSUE		LAIMS ALLOWED	
	Assistant Examiner	Dockat Clerk	Total Claims	Print Claim	
IBSUE FEE			Bheets	DRAWING Figs, Drwg. Print P	
Amount Due Date Peld		Primary E		The state of the	
	and the state of the	LASSIFICATION	ISSUE		
	Class		NUMBER		
Label		n disclosed berein n	nay be restricted. Una Title 35, Sections 122, 1	uthorized claciosure mi	iy 154

21.02 33. K Foralle SEARCHED 34 in Class Sub. Exmr. Date . On 9/29/42 1) 580 530 381.3 350 UF 102 J -/02 the W ipd 13 Ŵ Dol. 50 6 Эð 59/ 10/15/14 part 10/54 wid INTERFERENCE SEARCHED Request for allesu 4-15-5à. 5 Sub. Class Date Exmr. Lin Ne pl

Best Available Copy



•

Page: 1

Raw Sequence Listing

06/25/91 10:32:10

#1/

Patent Application US/07/715,272

SEQUENCE LISTING 1 2 3 (1) GENERAL INFORMATION: 4 (i) APPLICANT: Carter, Paul J. 5 Presta, Leonard G. 6 7 (ii) TITLE OF INVENTION: Immunoglobulin Variants 8 9 (iii) NUMBER OF SEQUENCES: 10 10 11 (iv) CORRESPONDENCE ADDRESS: 12 (A) ADDRESSEE: Genentech, Inc. 13 (B) STREET: 460 Point San Bruno Blvd 14 (C) CITY: South San Francisco 15 16 (D) STATE: California (E) COUNTRY: USA 17 (F) ZIP: 94080 18 19 (v) COMPUTER READABLE FORM: 20 (A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy disk 21 (B) COMPUTER: IBM PC compatible 22 (C) OPERATING SYSTEM: PC-DOS/MS-DOS 23 (D) SOFTWARE: patin (Genentech) 24 25 (vi) CURRENT APPLICATION DATA: 26 27 (A) APPLICATION NUMBER: (B) FILING DATE: 14-June-1991 28 29 (C) CLASSIFICATION: 30 31 (vii) PRIOR APPLICATION DATA: 32 (A) APPLICATION NUMBER: 33 (B) FILING DATE: 34 (viii) ATTORNEY/AGENT INFORMATION: 35 (A) NAME: Adler, Carolyn R. 36 (B) REGISTRATION NUMBER: 32,324 37 (C) REFERENCE/DOCKET NUMBER: 709 38 39 (ix) TELECOMMUNICATION INFORMATION: 40 41 (A) TELEPHONE: 415/266-2614 (B) TELEFAX: 415/952-9881 42 (C) TELEX: 910/371-7168 43 44 (2) INFORMATION FOR SEQ ID NO:1: 45 46 (i) SEQUENCE CHARACTERISTICS: 47 48 (A) LENGTH: 109 amino acids (B) TYPE: amino acid 49 50 (D) TOPOLOGY: linear 51 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1: 52 53

and the second second

· · · · · · · · · · · · · /

Raw Sequence Listing

Patent Application US/07/715,272

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val . 15 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ser Ala Ser Phe Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr . 109 (2) INFORMATION FOR SEQ ID NO:2: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2: Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys Asp Thr Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu 35 4 Glu Trp Val Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp) Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr

06/25/91 10:32:11

06/25/91 10:32:13

Patent Application US/07/715,272

107	and an all all mar ton Val Thr Val Ser Ser	
108	Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser 110 115 120	
109	110 115 120	
110		
111		
112	(2) INFORMATION FOR SEQ ID NO:3:	
113		
114	(i) SEQUENCE CHARACTERISTICS:	
115	(A) LENGTH: 109 amino acids	
116	(B) TYPE: amino acid	
117	(D) TOPOLOGY: linear	
118		
119	(XI) SEQUENCE DESCRIPTION: SEQ ID NO:3:	
120		
121	Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val	
122	10^{-15}	
123		
124	Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser	
125	20 25 30	
126		
127	Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys	
128	35 40 45	
129		
130	Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser	
131	50 55 60	
131		
132	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile	
134	65 70 75	
	•••	
135	Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln	•
136	80 85 90	
137		
138	Tyr Asn Ser Leu Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu	
139	95 100 105	
140		
141	man manage milese	
142	Ile Lys Arg Thr	
143	109	
144	THE THE TON FOR SEC IN NO.4.	
145	(2) INFORMATION FOR SEQ ID NO:4:	
146		
147	(i) SEQUENCE CHARACTERISTICS:	
148	(A) LENGTH: 120 amino acids	
149	(B) TYPE: amino acid	
150	(D) TOPOLOGY: linear	
151	The second secon	
152	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:	
153	dia the state of the dia ten Hel dia Bao dia	
154	Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly 10 15	
155	1 5 10 15	
156		
157	Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser	•
158	20 25 30	
159.		

Raw Sequence Listing

Patent Application US/07/715,272

Asp Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Val Ile Ser Glu Asn Gly Gly Tyr Thr Arg Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr ·95 Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser (2) INFORMATION FOR SEQ ID NO:5: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5: Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly His Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Phe Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly Asn Arg Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala (2) INFORMATION FOR SEQ ID NO:6:

06/25/91 10:32:15

Page: 5

Raw Sequence Listing

Patent Application US/07/715,272

(i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6: Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser Leu Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Thr Tyr Ile His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile Gly Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Asp Pro Lys Phe Gln Asp Lys Ala Thr Ile Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr Leu Gln Val Ser Arg Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr Ala Met Asp Tyr Trp Gly Gln Gly Ala Ser Val Thr Val Ser Ser (2) INFORMATION FOR SEQ ID NO:7: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear (x1) SEQUENCE DESCRIPTION: SEQ ID NO:7: TCCGATATCC AGCTGACCCA GTCTCCA 27 (2) INFORMATION FOR SEQ ID NO:8: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 31 bases (B) TYPE: nucleic acid

06/25/91 10:32:17 Raw Sequence Listing

Page:

6

Patent Application US/07/715,272

06/25/91 10:32:19

266 267 268		(C) STRANDEDNESS: single (D) TOPOLOGY: linear
269	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:8:
270		
271		a not not not according
272 273		GTTTGATCTC CAGCTTGGTA COXXCDCCGA A 31 to the rule.
274		
275		
276	(2) IN	FORMATION FOR SEQ ID NO:9:
277		
278 279	(i)	SEQUENCE CHARACTERISTICS: (A) LENGTH: 22 bases
2/9		(A) LENGTA: 22 DASES (B) TYPE: nucleic acid
281		(C) STRANDEDNESS: single
282		(D) TOPOLOGY: linear
283		
284	(Xi)	SEQUENCE DESCRIPTION: SEQ ID NO:9:
285 286		
287	,	AGGTXXAXCT GCAQXAGTCX)GG 22
288		
289		
290 291	(3) Th	FORMATION FOR SEQ ID NO:10:
292	(2) 11	FORMATION FOR SEQ 1D NO. 10.
293	(i)	SEQUENCE CHARACTERISTICS:
294		(A) LENGTH: 34 bases
295		(B) TYPE: nucleic acid
296 297		(C) STRANDEDNESS: single (D) TOPOLOGY: linear
298	•	(<i>D</i>) 10100001, 11001
299	(Xi)	SEQUENCE DESCRIPTION: SEQ ID NO:10:
300		
301		
302	· · · ·	TGAGGAGACG GTGACCGTGG TCCCTTGGCC CCAG 34

PAGE: 1

SEQUENCE VERIFICATION REPORT PATENT APPLICATION US/07/715,272 DATE: 06/25/91 TIME: 10:32:20

LINE ERROR

ORIGINAL TEXT

272	Wrong Nucleic Acid Designator	GTTTGATCTC CAGCTTGGTA COXXCDCCGA A 31
269	Entered and Calc. Seq. Length difference we to	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
287	Wrong Nucleic Acid Designator	AGGÍXYAXCT GCAGXAGTCX GG 22 (xi) sequence description: seq id NO:9:

PAGE: 1

SEQUENCE MISSING ITEM REPORT PATENT APPLICATION US/07/715,272 DATE: 06/25/91 TIME: 10:32:20

MANDATORY IDENTIFIER THAT WAS NOT FOUND

PAGE: 1

SEQUENCE CORRECTION REPORT PATENT APPLICATION US/07/715,272 DATE: 06/25/91 TIME: 10:32:20

15-07

LINE ORIGINAL TEXT

CORRECTED TEXT

Genentech, Inc. Attn: Carolyn R. Adler 460 Point San Bruno Blvd. South San Francisco, CA 94080 Paul J. Carter 07/715,272 June 14, 1991

NOTICE TO COMPLY WITH REQUIREMENTS FOR PATENT APPLICATIONS CONTAINING NUCLEOTIDE SEQUENCE AND/OR AMINO ACID SEQUENCE DISCLOSURES

This application contains sequence disclosures that are encompassed by the definitions for nucleotide and/or amino acid sequences set forth in 37 CFR § 1.821(a)(1) and (a)(2). However, this application fails to comply with one or more of the requirements of 37 CFR §§ 1.821 through 1.825 as follows:

1. This application clearly fails to comply with the collective requirements of §§ 1.821 through 1.825. Applicant's attention is directed to these regulations, a copy of which is attached.

2. This application does not conform exclusively to the requirements of §§ 1.821 through 1.825. The non-conforming material should be deleted. § 1.821(b).

3. This application does not contain, as a separate part of the disclosure on paper copy, a "Sequence Listing." § 1.821(c).

4. This application does contain, as a separate part of the disclosure on paper copy, a "Sequence Listing." However, the "Sequence Listing" does not comply with the requirements of §§ 1.821 through 1.825 as follows:

a. The sequence data does not comply with the symbol and format requirements of paragraphs (b) through (p) of § 1.822. Specifically:

b. The "Sequence Listing" does not comply with the location and page requirements of paragraph (a) of § 1.823.

c. The "Sequence Listing" does not comply with the information requirements of paragraph (b) of § 1.823. Specifically:

6. A copy of the "Sequence Listing" in computer readable form has not been submitted as required by § 1.821(c).

 7 A copy of the "Sequence Listing" in computer readable form has been submitted. However, the computer readable form does not comply with the requirements of § 1.824. Specifically:

8. A statement that the content of the paper and computer readable copies are the same has not been submitted as required by § 1.821(f).

9. The amendment to or replacement of the paper and/or computer readable copies of the "Sequence Listing" does not comply with the requirements of § 1.825(a) through (c).

10. The computer readable form that has been filed with this application has been found to be damaged and/or unreadable. Applicant must provide a substitute copy of the data in computer readable form accompanied by a statement that the substitute data is identical to that originally
 § 1.825(d). Specifically:

Other:

TVEN ONE MONTH FROM THE DATE OF THIS LETTER WITHIN WHICH THE ABOVE REQUIREMENTS. Failure to comply with the above require-SANDONMENT of the application under 37 CFR 1.821(g). Extensions of v filing a petition accompanied by the extension fee under the provisions of the response to, and any questions about, this notice to the undersigned. A "b returned with your response.

				Patent and Trademark	PATENTS AND TRADEMARKS
, ,	APPLICATION NUM	ADER FILING DATE	FIRST NAMED API	LICANT	ATTY DOCKET NO/TITLE
	07/7	15,272 06/14/91	CARTER	l ₂	709
	ATTN 460	NTECH, INC. : CAROLYN R. ADLER PUINT SAN BRUNG BLV H SAN FRANCISCO, CA	7D. 1 94080	០០០ DATE MAILED:	07/03/91
		NOTICE TO I		RTS OF APPLICATIO	
		, ,	FILING DATE G		
		ng date has been granted to this			
		missing parts are filed within th		otal amount owed by applica イコム・のり	nt as a
	-	ge entity, 🗆 small entity (verified		120-	
	' 1.0	The statutory basic filing fee is			
	•	small entity, must submit SUBMIT THE SURCHARGE.		nplote the basic filing fee V.	and MUST ALSO
	2. 🗆	Additional claim fees of \$ dependent claim fee, are requir claims for which fees are due.	as a 🗇 large ent	ity € small entity, including ait the additional claim fees o	r cancel the additional
	3. []	The eath or declaration: is missing. does not cover items emitted	d at time of execution.		· .
:		An oath or declaration in compli Number and Filing Date is req BELOW.	once with 37 CFR 1.63. i	dentifying the application by MUST ALSO BE SUBMITI	the above Application TED AS INDICATED
	4. 🗅	The oath or declaration doos a compliance with 37 CFR 1.63, i is required. A SURCHARGE	dentifying the application	m by the above Application N	umber and Filing Date
	5. 🗗	The signature to the calh or do inventor or a person qualified compliance with 37 CFR 1.63, i is required. A SURCHARGE 1	under 37 CFR 1.42, 1.43 dentificing the applicatio	3, or 1.47. A properly signed on by the above Application N	umber and Filing Date
	6. 🟳	The signature of the following	joint inventor(s) is miss	ing from the oath or declarat	ion:
		the omitted inventor(s), identi- required. A SURCHARGE MI	iving this application by	on listing the names of all in the above Application Numl TED AS INDICATED BELO	er and Receipt Date is
	7. 🗆	The application was filed in a translation of the application of paid. NO SURCHARCE IS R	nd a fee of \$30.00 unde	r 37 CFR 1.17(k), unless thi	e a verified English s fee has already been
	8. 🗂	A \$50.00 processing fee is requ			
	9. 🗀	Your filing receipt was mailed	in error because check	was returned without payme	nt.
	10. 🗔	Other.			
2 2	·	An Application Number and Fi identified above in items 1 a SURCHARGE of \$120.00 for la claiming such status. The surc THE DATE OF THIS LETTE WHICHEVER IS LATER, will abandomment. Extensions of 1 under the provisions of 37 CFI	nd 3-6 must be timely rge entities or \$60.00 ft hago is set forth in 37 Cl CR, OR TWO MONTHS in which to file all miss ime may be obtained by	provided ALONG WITH T or small entities who have file FR 1.16(e). Applicant is giver S FROM THE FILING DAT ing marts and pay any fees r	He PAYMENT OF A ed a verified statement ONE MONTH FROM E of this application, equired above to avoid

Direct the response to, and any questions about, this notice to ATTENTION: Application Division, Special Handling Unit.

137

A copy of this notice <u>MUST</u> be returned with response. (Maywad) For: Manager, Application Division (703) 557-348-1242

OFFICE COPY

FORM PTO 1533 (REV. 6-90)

• .			
	, , , ,		1
	· · ·	UNITED STA DEPARTMENT OF COMM	V
	1	NULL 12 12 12 12 12 12 12 12 12 12	
		1 191 Washington, D.C. 20231	₩S
· ·	<u> </u>	FR. NO DATE FIRST NAMED APPLICANT ATTY DOCKET NO./ITLE	
		- 07/715, 272 06/14/21 CARTER F 702	,
		5	1
		GENERITECH, INC.	
		ATTNE CAROLYN R. ADLER 460 Politi'san brund biyd.	
		SOUTH SAM ERANCISCO, CA 94080 000	
		••• DATE MAILED: 07/03/91	•
•		TOTICE TO FILE MISSING PARTS OF APPLICATION FILING DATE GRANTED	
		2.	
· .		A filing date has been granted to this application. However, the following parts are missing. If all missing parts are filed within the period set below, the total amount owed by applicant as a	
		I an integrating party are ined within the period set below, the total amount owed by applicant as a \Box barge entity, \Box small entity (verified statement filed), is $\frac{1}{2} \frac{1}{2} \frac{1}{$	
	1	1. The statutory basic filing fee is: missing insufficient. Applicant as a large entity	
•	1.	I small entity, must submit \$ to complete the basic filing fee and MUST ALSO	
•		SUBMIT THE SURCHARGE AS INDICATED BELOW. 2. Additional claim fees of \$as a □ large ontity □ small entity, including any required multiple	
		dependent claim fee, are required. Applicant must aubmit the additional claim fees or cancel the additional claims for which fees are due. NO SURCHARGE IS REQUIRED FOR THIS ITEM,	
<u>.</u>		3. [] The eath or declaration:	
		 is missing. idees not cover items omitted at time of execution. 	
		An each or declaration in compliance with 37 CFR 1.63, identifying the application by the above Application Number and Filing Date is required. A SURCHARGE MUST ALSO BE SUBMITTED AS INDICATED BELOW.	•
		4.	
	•	5. C-The signature to the oath or declaration is: Comissing; C a reproduction; C by a person other than the	
		inventor or a person qualified under 37 CFR 1.42, 1.43, or 1.47. A properly signed eath or declaration in compliance with 37 CFR 1.63, identifying the application by the above Application Number and Filing Date is required. A SURCHARGE MUST ALSO BE SUBMITTED AS INDICATED BELOW	
` .		6. The signature of the following joint inventor(e) is missing from the oath or declaration:	
		An oath or declaration listing the names of all inventors and signed by the omitted inventor(s), identifying this application by the above Application Number and Receipt Date is required. A SURCHARGE MUST ALSO BE SUBMITTED AS INDICATED BELOW.	
•	•	7. The application was filed in a language other than English. Applicant must file a verified English translation of the application and a fee of \$30.00 under 37 CFR 1.17(k), unless this fee has already been paid. NO SURCHARGE IS RERQUIRED FOR THIS ITEM.	
		 8. A \$50.00 processing fee is required for returned thecks. (37 CFR 1.21(m)). 	
		9. 🔲 Your filing receipt was mailed in error because check was returned without payment.	
· •		10. 🖸 Other.	
		An Application Number and Filing Date have been assigned to this application. The missing parts and fees identified above in items 1 and 3-6 must be timely provided ALONG WITH THE PAYMENT OF A SURCHARCE of \$120.00 for large entities or \$60.00 for small entities who have filed a verified statement claiming such status. The surchage is set forth in \$7 CFR 1.16(o). Applicant is given ONE MONTH FROM	
		THE DATE OF THIS LETTER, OR TWO-MONTHS FROM THE FILING DATE of this application, WHICHEVER IS LATER, within which to file all missing parts and pay any fees required above to avoid abandonment. Extensions of time may be obtained by filing a polition accompanied by the extension fee under the provisions of 37 CFR 1. 136(a).	
		Direct the response to, and any questions about, this notice to ATTENTION: Application Division, Special Handling Unit.	
		A copy of this notice \underline{MUST} be returned with response.	
	لي الح	For: Manager, Application Division	
<u>[</u>		D520157 07/17/91 715272 138	

A	L RO	
61	JUL	15
PA	1. 1	E)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

intre Application of

Paul J. Carter et al.

Serial No. 07/715,272

Filed: 14 June 1991

For: IMMUNOGLOBULIN VARIANTS

Examiner:

Group Art Unit:

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 266-2614

PATENT DOCKET 709

¥5

HEGEIVED

JUL 1 8 1991

APPLICATION DIVISION-401

TRANSMITTAL LETTER

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231 Attn: Application Branch

Sir:

Transmitted herewith are the following documents:

1. Declaration duly executed.

2 Copy of PTO-1553.

The Commissioner is hereby authorized to deduct the appropriate surcharge fee of \$120 associated with this communication or credit any overpayment to Deposit Account No. 07-0630. A duplicate of this sheet is enclosed.

Respectfully submitted,

GENENTECH, INC. Aler

Carolyn R. Ádler Reg. No. 32,324

9 July 1991

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Parol Moulu Carol Koehler

Date: 9 July 1991

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

Docket No. 709

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

IMMUNOGLOBULIN VARIANTS

the specification of which (check one) __ is attached hereto or \underline{x} was filed on <u>14 June 1991</u> as Application Serial No. 07/715,272 and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I hereby state that any Sequence Listing submitted with this application is submitted in paper copy and a computerreadable diskette, and that the content of the paper and computer readable copies are the same.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate have a filing date before that of the application on which priority is claimed:

Prior Foreign App		Priority Claimed		
			Yes	No
Number	Country	Day/Month/Year Filed		

I hereby claim the benefit under Title 35, United States Code, §120 of any United States applications(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Ser. No.	Filing Date	Status: Patented, Pending, Abandoned
		· · · · · · · · · · · · · · · · · · ·
Application Ser. No.	Filing Date	Status: Patented, Pending, Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Carolyn R. Adler - Reg. No. 32,324 301 Robert H. Benson - Reg. No. 30,446 Walter E. Buting - Reg. No. 23,092 Ginger R. Dreger - Reg. No. 33,055 Debbie Glaister - Reg. No. 33,888 Janet E. Hasak - Reg. No. 27,043

 Max D. Hensley
 - Reg. No. 27,043

 Dennis G. Kleid
 - Reg. No. 32,037

 Nancy Olseki
 - Reg. No. 34,688

 Stephen Raines
 - Reg. No. 25,912

 Daryl B. Winter
 - Reg. No. 32,637





Senid correspondence to Lol

Genentech, Inc.
Attn: Carolyn R. Adler
460 Point San Bruno Boulevard
South San Francisco, CA 94080

Telephone: (415) 266-2614

I hereby declare that all statements made herein of my own knowledge and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or Imprisonment or both, under Section 1001 of Title 18 of the United States Code and that willful false statements may jeopardize the validity of the application or any patent issued thereon.

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from his foreign patent agent as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

Full name of sole or first inventor Paul J. Carter 4000, Paul John Cotes	18th June 1991
Paul J/Carter 40100 / 1000 000000000	Date
Inventor's signature	
Residence 2074 18th Avenue San Francisco, CA 94116	
Citizenship United Kingdom	
Post Office Address 460 Point San Brumo Boulevard South San Francisco, CA 94080	
Full name of second joint inventor, if any	· · ·
Leonard G Presta 40208	
Second Inventor's signature & Presta	Date 6-19-91
Residence # 1900 Gough Street, #206 San Francisco, CA-94109	
Citizenship United States of America	
Post Office Address 460 Point San Bruno Boulevard South San Francisco, CA 94080	
Full name of third joint inventor, if any	
Third Inventor's signature	Date
Residence	
Citizenship	
Post Office Address 460 Point San Bruno Boulevard South San Francisco, CA 94080	

Best Available Copy

WIL ROOM Paul J. Carter CALENDARED RECEIVED at 191nc E JUL 0 8 1991 07/715,272 3 +4991 Attur Carolynes Adler 460 Pole Adler Bruno Bl June 14, 1991 Bruno Blvd. Genentech, Inc. Legal Dept. South San Francisco, CA 94080 NOTICE TO COMPLY WITH REQUIREMENTS FOR PATENT APPLICATIONS CONTAINING NUCLEOTIDE SEQUENCE AND/OR AMINO ACID SEQUENCE DISCLOSURES This application contains sequence disclosures that are encompassed by the definitions for nucleo-tide and/or amino acid sequences set forth in 37 CFR § 1.821(a)(1) and (a)(2). However, this application fails to comply with one or more of the requirements of 37 CFR §§ 1.821 through 1.8251. This application clearly fails to comply with the collective requirements of §§ 1.821 through 1.825. Applicant's attention is directed to these regulations, a copy of which is attached. as follows: 2. This application does not conform exclusively to the requirements of §§ 1.821 through 1.825. The non-conforming material should be deleted. § 1.821(b). 3: This application does not contain, as a separate part of the disclosure on paper copy, a "Sequence Listing." § 1.821(c). 4. This application does contain, as a separate part of the disclosure on paper copy, a "Sequence Listing." However, the "Sequence Listing" does not comply with the requirements of \$5 1.821 through 1.825 as follows: a. The sequence data does not comply with the symbol and format requirements of paragraphs (b) through (p) of § 1.822. Specifically: b. The "Sequence Listing" does not comply with the location and page requirements of paragraph (a) of § 1.823. c. The "Sequence Listing" does not comply with the information requirements of paragraph (b) of § 1.823. Specifically: _ d. Other: _ 5. The description and/or claims of the patent application mention a sequence that is set forth in the "Sequence Listing" but reference is not properly made to the sequence by use of a sequence identifier as required by § 1.821(d). 6. A copy of the "Sequence Listing" in computer readable form has not been submitted as required by § 1.821(c). L 1. A copy of the "Sequence Listing" in computer readable form has been submitted. How-ever, the computer readable form does not comply with the requirements of § 1.824. Specifically: 8. A statement that the content of the paper and computer readable copies are the same has not been submitted as required by \$ 1.821(f). 9. The amendment to or replacement of the paper and/or computer readable copies of the "Sequence Listing" does not comply with the requirements of \$ 1.825(e) through (c). 10. The computer readable form that has been filed with this application has been found to be damaged and/or unreadable. Applicant must provide a substitute copy of the data in computer readable form accompanied by a statement that the substitute data is identical to that originally filed. § 1.825(d). Specifically: _ APPLICANT IS GIVEN ONE MONTH FROM THE DATE OF THIS LETTER WITHIN WHICH TO COMPLY WITH THE ABOVE REQUIREMENTS. Failure to comply with the above require-ments will result in ABANDONMENT of the application under 37 CFR 1.821(g). Extensions of time may be obtained by filing a petition accompanied by the extension fee under the provisions of 37 CFR §-1.136. Direct the response to, and any questions about, this notice to the undersigned. A copy of this notice MUST be returned with your response. 11. Other. _

J C Mylayword 142

	— • • •
ROOM.	SEQUENCE LISTING
Le jui	(1) GENERAL INFORMATION:
	(i) APPLICANT: Carter, Paul J. Presta, Leonard G.
PRADEN	(ii) TITLE OF INVENTION: Immunoglobulin Variants
10	(iii) NUMBER OF SEQUENCES: 10
15	 (iv) CORRESPONDENCE ADDRESS: (A) ADDRESSEE: Genentech, Inc. (B) STREET: 460 Point San Bruno Blvd (C) CITY: South San Francisco (D) STATE: California (E) COUNTRY: USA (F) ZIP: 94080
20	 (v) COMPUTER READABLE FORM: (A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy disk (B) COMPUTER: IBM PC compatible (C) OPERATING SYSTEM: PC-DOS/MS-DOS (D) SOFTWARE: patin (Genentech)
25	 (vi) CURRENT APPLICATION DATA: (A) APPLICATION NUMBER: 07/715,272 (B) FILING DATE: 14-June-1991 (C) CLASSIFICATION:
30	<pre>(vii) PRIOR APPLICATION DATA: (A) APPLICATION NUMBER: (B) FILING DATE:</pre>
35	 (viii) ATTORNEY/AGENT INFORMATION: (A) NAME: Adler, Carolyn R. (B) REGISTRATION NUMBER: 32,324 (C) REFERENCE/DOCKET NUMBER: 709
40	<pre>(ix) TELECOMMUNICATION INFORMATION: (A) TELEPHONE: 415/266-2614 (B) TELEFAX: 415/952-9881 (C) TELEX: 910/371-7168</pre>
45	(2) INFORMATION FOR SEQ ID NO:1:
50	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
55	Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val 1 5 10 15
	Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Asn 20 25 30
60	Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys 35 40 45
4 	Leu Leu Ile Tyr Ser Ala Ser Phe Leu Glu Ser Gly Val Pro Ser 50 55 60

Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln ÷2 His Tyr Thr Thr Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr (2) INFORMATION FOR SEQ ID NO:2: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2: Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys Asp Thr Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser (2) INFORMATION FOR SEQ ID NO:3: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3: Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys

	1 · · · · · · · · · · · · · · · · · · ·
· ·	Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser 50 55 60
-5	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile 65 70 75
	Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln 80 85 90
10	Tyr Asn Ser Leu Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu 95 100 105
15	Ile Lys Arg Thr 109
	(2) INFORMATION FOR SEQ ID NO:4:
20	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
25	Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly 1 5 10 15
20	Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser 20 25 30
30	Asp Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu 35 40 45
35	Glu Trp Val Ala Val Ile Ser Glu Asn Gly Gly Tyr Thr Arg Tyr 50 55 60
	Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser 65 70 75
40	Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp 80 85 90
45	Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr 95 100 105
	Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser 110 1)5 120
50	(2) INFORMATION FOR SEQ ID NO:5:
55	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
60	Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val 1 5 10 15
	Gly Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn 20 25 30

Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly His Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Phe Arg Tyr Thr Gly Val Pro Asp -5 . . Arg Phe Thr Gly Asn Arg Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala (2) INFORMATION FOR SEQ ID NO:6: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 120 amino acids (B) TYPE: amino acid(D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6: Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser Leu Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Thr Tyr Ile His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile Gly Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Asp Pro Lys Phe Gln Asp Lys Ala Thr Ile Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr Leu Gln Val Ser Arg Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr 95 100 105 Ala Met Asp Tyr Trp Gly Gln Gly Ala Ser Val Thr Val Ser Ser (2) INFORMATION FOR SEQ ID NO:7: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TCCGATATCC AGCTGACCCA GTCTCCA 27

(2) INFORMATION FOR SEQ ID NO:8: (i) SEQUENCE CHARACTERISTICS: -5 . . (A) LENGTH: 31 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single
(D) TOPOLOGY: linear 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8: GTTTGATCTC CAGCTTGGTA CCHSCDCCGA A 31 15 (2) INFORMATION FOR SEQ ID NO:9: (i) SEQUENCE CHARACTERISTICS: 20 (A) LENGTH: 22 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9: AGGTSMARCT GCAGSAGTCW GG 22 30 (2) INFORMATION FOR SEQ ID NO:10: (i) SEQUENCE CHARACTERISTICS: 35 (A) LENGTH: 34 bases (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

45

147

TGAGGAGACG GTGACCGTGG TCCCTTGGCC CCAG 34

	CED AN T
61 1991 5	PATENT DOCKET 709
THE UNITED STATES PATEN	IT AND TRADEMARK OFFICE
C IRADE	
In re Application of)	
PAUL J. CARTER ET AL.	Art Unit: to be assigned
Serial No. 07/715,272	
) Filed: June 14, 1991)	Examiner: to be assigned Thereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Com- first class mail in an envelope addressed to: Com-
For: IMMUNOGLOBULIN VARIANTS)	first class mail in an anyologic during the second
)	LOUISE STRASBAUCH
RESPONSE AND PRELIN	
	Signature of Depositing Party JULY 12, 1991
Honorable Commissioner of Patents and Tradem Washington, D.C. 20231	

Sir:

This is responsive to the Notice to Comply with Requirements for Patent Applications Containing Nucleotide and/or Amino Acid Sequence Disclosures, mailed June 25, 1991. The inventors also take this opportunity to correct two minor grammatical errors in the application, and add no new matter.

Enclosed is an amended sequence listing submitted with a paper copy and a computerreadable diskette. The sequence listing has been corrected to conform exactly to the sequences as recited in the specification as originally filed. I hereby state that the content of this paper and computer readable copies are the same, and that this amendment corrects errors in the previous sequence listing submission without adding new matter.

IN THE SPECIFICATION:

Please make the following amendments:

On page 12, line 1, delete genes and insort --sequences--.

On page 16, line 12, delete intrachain-affecting and insert --interchain--affecting.

Respectfully Submitted, GENENTECH, INC.

Carolyn R. Ádler Reg. No. 32,324

July 12, 1991 460 Point San Bruno Blvd South San Francisco, CA 94080 Page: 1

Ţ

Raw Sequence Listing



Patent Application US/07/715,272A

1		SEQUENCE LISTING
2		
3 4	(1) GE1	VERAL INFORMATION:
* 5	(i)	APPLICANT: Carter, Paul J.
6	()	Presta, Leonard G.
. 7		
8	(ii)	TITLE OF INVENTION: Immunoglobulin Variants
9 10	(111)	NUMBER OF SEQUENCES: 10
11	(111)	
12	(iv)	CORRESPONDENCE ADDRESS:
13		(A) ADDRESSEE: Genentech, Inc.
14		(B) STREET: 460 Point San Bruno Blvd
15	*	(C) CITY: South San Francisco
16		(D) STATE: California (E) COUNTRY: USA
17 18	,	(F) ZIP: 94080
19		(1) 111. 74000
20	(v)	COMPUTER READABLE FORM:
21		(A) MEDIUM TYPE: 5.25 inch, 360 Kb floppy disk
22		(B) COMPUTER: IBM PC compatible
23		(C) OPERATING SYSTEM: PC-DOS/MS-DOS
24	`	(D) SOFTWARE: patin (Genentech)
25		AND A DET TOTOT DAMA
26	(V1)	CURRENT APPLICATION DATA: (A) APPLICATION NUMBER: 07/715,272
27 28		(B) FILING DATE: 14-June-1991
29		(C) CLASSIFICATION:
30		
31	(vii)	PRIOR APPLICATION DATA:
32		(A) APPLICATION NUMBER:
33		(B) FILING DATE:
34		· · · · · · · · · · · · · · · · · · ·
35	(viii)	ATTORNEY/AGENT INFORMATION:
36		(A) NAME: Adler, Carolyn R. (B) REGISTRATION NUMBER: 32,324
37		(C) REFERENCE/DOCKET NUMBER: 709
38 39		(C) REFERENCE/DOCKET NOMBER. 703
40	(ix)	TELECOMMUNICATION INFORMATION:
41	()	(A) TELEPHONE: 415/266-2614
42	· •	(B) TELEFAX: 415/952-9881
43	,	(C) TELEX: 910/371-7168
44		
45	(2) IN	FORMATION FOR SEQ ID NO:1:
46	. .	
47	(i)	SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids
48	,	(A) LENGTH: 109 amino acids (B) TYPE: amino acid
49 50		(B) TOPOLOGY: linear
51		
52	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:1:
53		

Page: 2

Raw Sequence Listing

Patent Application U8/07/715,272A

07/19/91 16:16:26

54 55 56	Asp 1	Ile	Gln	Met	Thr 5	Gln	Ser	Pro	Ser	Ser 10	Leu	Ser	Ala	Ser	Val 15	
57 58 59	Gly	Asp	Arg	Val	Thr 20	Ile	Thr	Cys	Arg	Ala 25	Ser	Gln	Asp	Val	Asn 30	
60 61 62	Thr	Ala	Val	Ala	Trp 35	Tyr	Gln	Gln	Lys	Pro 40	Gly	Lys	Ala	Pro	Lув 45	
63 64 65	Leu	Leu	Ile	Tyr	Ser 50	Ala	Ser	Phe	Leu	Glu 55	Ser	Gly	Val	Pro	Ser 60	
66 67 68	-		Ser	-	65	_				70					75	
69 70 71	Ser	Ser	Leu	Gln	Pro 80	Glu	Asp	Phe	Ala	Thr 85	Tyr	Tyr	Сув	Gln	Gln 90	
72 73 74	Bis	Tyr	Thr	Thr	Pro 95	Pro	Thr	Phe	Gly	Gln 100	Gly	Thr	Lys	Val	Glu 105	
75 76 77	Ile	Lys	Arg	Thr 109												
78 79	(2) 1															
80	· (2	L) SI	EQUEN	ICE (CHARJ	ACTEI	RIST	ICS:								
80 81 82 83	(2	(1 (1	EQUEN A) LI B) TY D) T(engti (Pe:	H: 12 amin	20 ar 10 ac	nino cid		ls			•				
81 82 83 84 85		(1 (1 (1	A) LH B) TY	ENGTH (PE: OPOL(H: 12 amin DGY:	20 an no ad line	nino cid ar	acio		10:2 :	;					
81 82 83 84 85 86 87 88	(x:	(1 (1 (1 (1	A) LI B) T) D) T(ENGTI (PE:)POL(VCE I	H: 12 amin DGY: DESCI	20 ar no ac line RIPTI	nino cid æar [ON:	acio SEQ	IDI			Val	Gln	Pro	Gly 15	
81 82 83 84 85 86 87	(x: Glu 1	(1 (1 (1 (1 (1 Val	A) LH B) T) D) T(EQUEN	ENGTI (PE:)POL(VCE I Leu	H: 12 Amin DGY: DESCI Val 5	20 ar no ao line RIPTI Glu	aino cid æar ION: Sær	acio SEQ Gly	ID 1 Gly	Gly 10	Leu		· •		15	
81 82 83 84 85 86 87 88 89 90 91	(xi Glu 1 Gly	(1 (1 (1 (1 (1) SI Ser	A) LH B) TY D) TC EQUEN Gln	ENGTI (PE:)POLO NCE I Leu Arg	H: 12 amin DGY: DESCH Val 5 Leu 20	20 an 10 ac line RIPTI Glu Ser	nino cid æar ION: Sær Cys	acio SEQ Gly Ala	ID I Gly Ala	Gly 10 Ser 25	Leu Gly	Phe	Asn	Ile	15 Lys 30	•
81 82 83 84 85 86 87 88 89 90 91 92 93 94	(x: Glu 1 Gly Asp Glu	(i (i) SI Val Ser Thr Trp	A) LH B) TY D) TO EQUEN Gln Leu Tyr Val	ENGTH (PE:)POLO NCE I Leu Arg Ile Ala	H: 12 Amin DGY: DESCI Val 5 Leu 20 His 35 Arg 50	20 ar no ad line RIPTI Glu Ser Trp Ile	aino cid æar [ON: Ser Cys Val Tyr	acio SEQ Gly Ala Arg Pro	ID F Gly Ala Gln Thr	Gly 10 Ser 25 Ala 40 Asn 55	Leu Gly Pro Gly	Phe Gly Tyr	Asn Lys Thr	Ile Gly Arg	15 Lys 30 Leu 45 Tyr 60	
81 82 84 85 86 87 88 90 91 92 93 94 95 96 97	(x: Glu 1 Gly Asp Glu	(i (i) SI Val Ser Thr Trp	A) LH B) TY D) TO EQUEN Gln Leu Tyr	ENGTH (PE:)POLO NCE I Leu Arg Ile Ala	H: 12 Amin DGY: DESCI Val 5 Leu 20 His 35 Arg 50	20 ar no ad line RIPTI Glu Ser Trp Ile	aino cid æar [ON: Ser Cys Val Tyr	acio SEQ Gly Ala Arg Pro	ID F Gly Ala Gln Thr	Gly 10 Ser 25 Ala 40 Asn 55	Leu Gly Pro Gly	Phe Gly Tyr	Asn Lys Thr	Ile Gly Arg	15 Lys 30 Leu 45 Tyr 60	
81 82 83 84 85 86 87 88 90 91 92 93 94 95 96 97 98 99 100	(x: Glu 1 Gly Asp Glu Ala	(i (1 (1) Val Ser Thr Trp Asp	A) LH B) TY D) TO EQUEN Gln Leu Tyr Val	NGTI (PE:)POLO NCE I Leu Arg Ile Ala Val	H: 12 Amin DGY: DESCI Val 5 Leu 20 His 35 Arg 50 Lys 65	20 ar no ad line RIPTI Glu Ser Trp Ile Gly	aino cid æar ION: Sær Cys Val Tyr Arg	acio SEQ Gly Ala Arg Pro Phe	ID R Gly Ala Gln Thr Thr	Gly 10 Ser 25 Ala 40 Asn 55 Ile 70	Leu Gly Pro Gly Ser	Phe Gly Tyr Ala	Asn Lys Thr Asp	Ile Gly Arg Thr	15 Lys 30 Leu 45 Tyr 60 Ser 75	

Raw Sequence Listing

07/19/91 16:16:28

Patent Application US/07/715,272A

107	Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser	•
108	110 115 120	
109	110	
110		
111	(2) INFORMATION FOR SEQ ID NO:3:	
112	(2) INFORMATION FOR DBg 15 HOLD.	
113	(i) SEQUENCE CHARACTERISTICS:	
114	(1) SEQUENCE CHARACEERCEERCEERCEERCEERCEERCEERCEERCEERC	
115	(B) TYPE: amino acid	
116	(D) TOPOLOGY: linear	
117	(b) ideologi. insu	
118	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:	
119	(XI) SEQUENCE DEDONITIZION. DEE EL CONTE	
120	Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val	
121	- 10 15	
122	1 5 10 22	
123	Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser	
124	$\begin{array}{c} \text{Gly Asp Arg var int file int cys my int cost of 30 \\ 20 & 25 & 30 \end{array}$	
125	20 20	
126	Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys	
127	$\begin{array}{c} \text{Ser tyr Let All lip tyl old old ljo 1001 -1 -40} \\ 35 & 40 & 45 \end{array}$	
128 129		
130	Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser	
130	50 55 60	
131	.	
	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile	
133 134	75	•
135 136	Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln	
130	80 85 90	•
	55	
138	Tyr Asn Ser Leu Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu	
139	95 100 105	
140		
141	Ile Lys Arg Thr	
142	109	
143	103	
144	(2) INFORMATION FOR SEQ ID NO:4:	
145		
146.	(i) SEQUENCE CHARACTERISTICS:	
147	(A) LENGTH: 120 amino acids	
148	(B) TYPE: amino acid	
149		
150	(D) TOPOLOGY: linear	
151	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:	
152	(XI) READENCE DESCRIPTION, SEX ID NO.4.	
153	Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly	
154	- 10 15	
155	1 5 10 13	•
156	The des due ble ble Con dly Dhe Thy Dhe Cor	
157	Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser 30 30	
158	20 25 30	
159		

Raw Sequence Listing

Patent Application US/07/715,272A

Asp Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Val Ile Ser Glu Asn Gly Gly Tyr Thr Arg Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr 100 . Ala Met Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser (2) INFORMATION FOR SEQ ID NO:5: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 109 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5: Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val 10 -· 49 Gly Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly His Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Phe Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly Asn Arg Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala (2) INFORMATION FOR SEQ ID NO:6:

07/19/91 16:16:30

Page: 5

Raw Sequence Listing

07/19/91 16:16:32

Patent Application US/07/715,272Å

13	
14	(i) SEQUENCE CHARACTERISTICS:
15	(A) LENGTH: 120 amino acids
16	(B) TYPE: amino acid
17	(D) TOPOLOGY: linear
18	
19	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
20 °	
21	Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gl
22	1 5 10 1
23	
24	Ala Ser Leu Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Ly
25	20 25 3
26	
27	Asp Thr Tyr Ile His Trp Val Lys Gln Arg Pro Glu Gln Gly Le
28	35 40 4
29	
30	Glu Trp Ile Gly Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Ty
31	50 55 6
32	
33	Asp Pro Lys Phe Gln Asp Lys Ala Thr Ile Thr Ala Asp Thr Se
34	65 70 7
35	
36	Ser Asn Thr Ala Tyr Leu Gln Val Ser Arg Leu Thr Ser Glu As
37	80 85 9
38	
39	Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Ty
40	
41	
42	Ala Met Asp Tyr Trp Gly Gln Gly Ala Ser Val Thr Val Ser Se
43	110 115 12
44	
45	· · · · · · · · · · · · · · · · · · ·
46	(2) INFORMATION FOR SEQ ID NO:7:
47	
48	(i) SEQUENCE CHARACTERISTICS:
49	(A) LENGTH: 27 bases
50	(B) TYPE: nucleic acid
51	(C) STRANDEDNESS: single
52	(D) TOPOLOGY: linear
53	(~)
54	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
55	(<i>m</i>)
56	
57	TCCGATATCC AGCTGACCCA GTCTCCA 27
58	TARRITINGAN WARTONGAN, MARTANI WI
59	
60 ·	
61	(2) INFORMATION FOR SEQ ID NO:8:
	(2) INFORMATION FOR SEX ID NO:0:
67	
62	
63	(i) SEQUENCE CHARACTERISTICS:
	 (1) SEQUENCE CHARACTERISTICS: (A) LENGTH: 31 bases (B) TYPE: nucleic acid

Raw Sequence Listing

07/19/91 16:16:34

Patent Application US/07/715,272A

266	(C) STRANDEDNESS: single
267	(D) TOPOLOGY: linear
268	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
269	(XI) SEQUENCE DEBONIT 2
270	
271	GTTTGATCTC CAGCTTGGTA CCHSCDCCGA A 31
272	GTITGAICIC CROCIIGGIA COMPLETE
273	
274	· ·
275	TOT TE NO. 91
276	(2) INFORMATION FOR SEQ ID NO:9:
277	
278	(i) SEQUENCE CHARACTERISTICS:
279	(A) LENGTH: 22 bases
280	(B) TYPE: nucleic acid
281	(C) STRANDEDNESS: single
282	(D) TOPOLOGY: linear
283	
284	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
285	
286	
287	AGGTSMARCT GCAGSAGTCW GG 22
288	
289	
290	
291	(2) INFORMATION FOR SEQ ID NO:10:
292	
293	(i) SEQUENCE CHARACTERISTICS:
294	(A) LENGTH: 34 bases
295	(B) TYPE: nucleic acid
296	(C) STRANDEDNESS: single
297	(D) TOPOLOGY: linear
298	
299	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:
300	
301	
302	TGAGGAGACG GTGACCGTGG TCCCTTGGCC CCAG 34
303	
303	

Page:

SEQUENCE VERIFICATION REPORT PATENT APPLICATION US/07/715,272A

DATE: 07/19/91 TIME: 16:16:36

LINE ERROR

PAGE: 1

ORIGINAL TEXT

(A) APPLICATION NUMBER: 07/715,272

Wrong application Serial Number 27

SEQUENCE MISSING ITEM REPORT PATENT APPLICATION US/07/715,272A

DATE: 07/19/91 TIME: 16:16:36

MANDATORY IDENTIFIER THAT WAS NOT FOUND

ŝ

A

PAGE: 1

c .

SEQUENCE CORRECTION REPORT PATENT APPLICATION US/07/715,272A DATE: 07/19/91 TIME: 16:16:36

LINE ORIGINAL TEXT

CORRECTED TEXT

8M Feisee PATENT DOCKET IN THE UNITED STATES PATENT AND TRADEMARK OFFICE Group Art Unit: In re Application of Examiner: MAY U 8 1992 Paul J. Carter et al. **GROUP 180** Serial No. 07/715272 Filed: June 14, 1991 460 Point San Bruno Boulevard Immunoglobulin Variants For: South San Francisco, CA 94080 (415) 266-2614 INFORMATION DISCLOSURE STATEMENT by certify that this correspondences is being coposited with the United States Postal Service as iirst class mail in an envelope addressed to: Com-Honorable Commissioner of Patents missioner of Potents and Trademarks, Washington, and Trademarks D.C., 20231 on Dril 30, 1992 Washington, D.C. 20231 (Date of Deposit) RASBAUGH Sir: The following items are supplied to the United States Patent and Tradema k Officere advance Deuso Strau the prosecution of the subject application. Signature of Depositing Party tor 0,199) Chothia et al., J. Mol. Biol. 186:651-663 (1985) Date of Signature Novotny and Haber, Proc. Natl. Acad. Sci. USA 82:4592-4596 (1985) Cabilly et al., U.S. patent No. 4,816,567 Morrison, S. L. et al., Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984) Boulianne, G. L. et al., Nature 312:643-646 (1984) Neuberger, M. S. et al., Nature 314:268-270 (1985) Brüggemann, M. et al., J. Exp. Med. 166:1351-1361 (1987) Riechmann, L. et al., Nature 332:323-327 (1988) Love et al., Methods in Enzymology 178:515-527 (1989) Bindon et al., J. Exp. Med. 168:127-142 (1988) Jones, P. T. et al., Nature 321:522-525 (1986) Verhoeyen, M. et al., Science 239:1534-1536 (1988) Hale, G. et al., Lancet i:1394-1399 (1988) Queen, C. et al., Proc. Natl. Acad. Sci. USA 86:10029-10033 (1989) Co et al., Proc. Natl. Acad. Sci. USA 88:2869-2873 (1991) Gorman et al., Proc. Natl. Acad. Sci. USA 88:4181-4185 (1991) Daugherty et al., Nucleic Acids Research 19(9):2471-2476 (1991)

07/715272

Page No. 2

Brown et al., Proc. Natl. Acad. Sci. USA 88:2663-2667 (1991)
Junghans et al., Cancer Research 50:1495-1502 (1990)
Davies, D. R. et al., Ann. Rev. Biochem. 59:439-473 (1990)
Chothia, C. & Lesk, A. M., J. Mol. Biol. 196:901-917 (1987)
Chothia, C. et al., Nature 342:877-883 (1989)
Tramontano, A. et al., J. Mol. Biol. 215:175-182 (1990)
Margolies et al., Proc. Natl. Acad. Sci. USA 72:2180-2184 (1975)
Pluckthun, Biotechnology 9:545-51 (1991)
Spiegelberg et al., Biochemistry 9:4217-4223 (1970)
Wallick et al., J. Exp. Med. 168:1099-1109 (1988)
Sox et al., Proc. Natl. Acad. Sci. USA 66:975-982 (1970)
Margni et al., Ann. Rev. Immunol.6:535-554 (1988)
Fendly, B. M. et al., Cancer Res. 50:1550-1558 (1990)

Neuberger et al., Nature 312:604-608 (1984)

Takeda et al., Nature 314:452-454 (1985)

Snow and Amzel, Protein: Structure, Function, and Genetics 1:267-279, Alan R. Liss, Inc. pubs. (1986)

Cheetham, J., Protein Engineering, 2(3): 170-172 (1988)

WO 91/09967, pub. 07/11/91, Adair et al.

One copy of each item cited above is supplied, along with a completed Form PTO-1449. The Examiner is requested to make the citations of record.

This submission is understood to complement the results of the Examiner's own independent search. The submission of this Disclosure Statement should not be construed as a representation that a search was made, or that the cited itms are inclusive of all the relevant and amterial citations that may be available publicly.

The citation of any item is not an admission that the item is prior art. The right is reserved to antedate any item in adherence with standard procedures.

Respectfully submitted, GENENTECH: INC.

Carolyn R. Adler Reg. No. 32,324

Dated: April 30, 1992

	UNITED STATES DEPARTMENT OF COMMERCE Patent and Trademark Office Address* COMMIBBIONER OF PATENTS AND TRADEMARKS Washington, D.C. 20231
SERIAL NUMBER FILING DATE F	RET HAMED INVENTOR AT LORING DOCKET NO.
	FEISEE-I
GENENTECH, INC. Attn. Carolyn R. Adler 460 Point San Brund Blvd, South San Francisco, CA 94080	1806 95/12/92
This is a communication harm the exeminar in charge of your application COMMISSIONER OF PATIENTS AND TRADEMARKS	
YPGWIGOD This application test been examined Arsponsive to oc	
A shortened statutory period for response to the action is set to e Failure to respond within the period for response will cause the a	pplication to become abandoned: S5 U.S.C. 133
S. Notice of Art Cited by Applicant, PTO-1449. S. Information on How to Effect Drawing Changes, PTO Part II SUMMARY OF ACTION 1. Claims	ere pending in the applicat
Of the above, daims	are withdrawn from consideral
2. Claims	•
3, 🔲 Claims	
4. Ciałms	ero rejocted.
	am abjacted to
5. Ckelma	are objected to
e.X. cialma1-16	are subject to restriction or election requirement.
e Claims	are subject to restriction or election requirement.
Claims Chaims This application has been filed with informal drawing 8. Formal drawings are required in response to this Off	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. Nos action.
Claims Claims This application has been filed with internal drawing	are subject to restriction or election requirement. Is under 37 C.F.A. 1.65 which are acceptable for examination purposes. Ice solon. elved on Under 37 C.F.R. 1.64 these draw vation or Notice re Patent Drawing, PTO-948).
 Claims	are subject to restriction or election requirement. is under 37 C.F.R. 1.85 which are acceptable for examination purposes. fice action. elved on Under 37 C.F.R. 4.84 these draw sation or Notice re Patent Drawing, PTO-948). awings, field on thas (have) been [] approved by the planation).
Claims	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. Ilce action. elved on Under 37 C.F.R. 4.64 these draw sation or Notice re Patent Drawing, PTO-948). swings, filed on thas (have) been [] approved by the planation). has been [] approved; [] disapproved (see explanation).
Claims	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. fice action. elved on Under 37 C.F.R. 1.84 these draw astion or Notice re Patent Drawing, PTO-948). awings, field onhas (heve) been a approved by the planation). has been approved; acceptable core explanation). nder U.S.C. 119. The certified core has a been received and here received ; field on; field on;
6. Claims	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. Ites action. elved on Under 37 C.F.R. 1.84 these draw astion or Notice re Patant Drawing, PTO-948). awings, filed on thas (have) been [] approved by the planation). has been [] approved; [] disapproved (see explanation). nder U.S.C. 119. The contified copy has [] been received [] not been received ; filed on; thousand the merits is closed in r slowance except for formal matters, prosocution as to the merits is closed in
Claims	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. Ites action. elved on Under 37 C.F.R. 1.84 these draw astion or Notice re Patant Drawing, PTO-948). awings, filed on thas (have) been [] approved by the planation). has been [] approved; [] disapproved (see explanation). nder U.S.C. 119. The contified copy has [] been received [] not been received ; filed on; there is the merits is closed in r slowance except for formal matters, prosocution as to the merits is closed in
6. Claims	are subject to restriction or election requirement. Is under 37 C.F.R. 1.85 which are acceptable for examination purposes. Ites action. elved on Under 37 C.F.R. 1.84 these draw astion or Notice re Patant Drawing, PTO-948). awings, filed on thas (have) been [] approved by the planation). has been [] approved; [] disapproved (see explanation). nder U.S.C. 119. The contified copy has [] been received [] not been received ; filed on; thousand the merits is closed in r slowance except for formal matters, prosocution as to the merits is closed in

Serial No. 715272

Art Unit 1806

Restriction to one of the following inventions is required under 35 U.S.C. 121:

I. Claims 1-13, drawn to a method of making an antibody and an antibody comprising a polypeptide, classified in Class 435, 530 subclass 69.1, 350.

II. Claims 14-16, drawn to computer

representations, classified in Class 364, subclass 282.1+.

The inventions are distinct, each from the other because of the following reasons:

The two Groups are drawn to two different products, Group I being a biological molecule and Group II being a machine. These constitute two different statutory classes of invention and are 15 therefore patentably distinct one from the other.

Because these inventions are distinct for the reasons given above and have acquired a separate status in the art as shown by their different classification and divergent subject matter, and because the searches for the individual Groups are not

20 coextensive, restriction for examination purposes as indicated is proper.

Applicant is reminded that upon the cancellation of claims to a non-elected invention, the inventorship must be amended in compliance with 37 C.F.R. 1.48(b) if one or more of the currently

162

10

Serial No. 715272

Art Unit 1806

5

named inventors is no longer an inventor of at least one claim remaining in the application. Any amendment of inventorship must be accompanied by a diligently-filed petition under 37 C.F.R. 1.48(b) and by the fee required under 37 C.F.R. 1.17(h).

A telephone call was made to Carolyn Adler, on 12/9/91, to request an oral election to the above restriction requirement, but did not result in an election being made and a written restriction was requested.

Applicant is advised that the response to this requirement 10 to be complete must include an election of the invention to be examined even though the requirement be traversed. (37 C.F.R. 1.143).

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Lila 15 Feisee whose telephone number is (703) 308-2731.

Any inquiry of a general nature or relating to the status of this application should be directed to the Group receptionist whose telephone number is (703) 308-0196.

Papers related to this application may be submitted to Group 20 180 by facsimile transmission. Papers should be faxed to Group 180 via the PTO FAX Center located in Crystal Mall 1. The faxing of such papers must conform with the notice published in the Official Gazette, 1096 OG 30 (November 15, 1989). The CM1 FAX

Serial No. 715272

Art Unit 1806

. . . V

Center number is (703) 308-4227. The hours of operation of the Center are 8:45 am - 4:45 pm, Monday - Friday.

Feisee/lf May 11, 1992

JOHN J. DOLL SUPERVISORY PATENT EXAMINER GROUP 180

FORM 948 . 5-90)	U.S. DEPARTME Patent a	
		APPLICATION NUMBER
		S PATENT DRAWING REVIEW
The drawings filed	6/14/91	checked below. The examiner will require submission of new, drawings must be submitted according to the instructions listed
Margin 8 1/2 by 8 1/2 by Top 2 inches 1 3 inche Top 2 inches 1 inch Left 1/4 inch 1/4 inch Right 1/4 inch 1/4 inch Bottom 1/4 inch 1/4 inch Proper Size Paper All Sheets Must be Sheet(s) Sheet(s) Proper Margins Rest Sheet(s) 1/2 TOP LEFT LEFT	Poor. Poor. Poor. DIN size A4 DIN size A4 DIN size A4 21 by 29.7 cm. A 2.5 cm. A 2.5 cm. A 1.5 cm. A 1.0 cm. Required. Same Size. - 4 equired. BIGHT BOTTOM	 4. Hatching and Shading. 37 CFR 1.84(d) Shade Lines are Required. Fig(s) Criss-Cross Hatching Not Allowed. Fig(s) Double Line Hatching Not Allowed. Fig(s) Parts in Section Must be Hatched. Fig(s) Feference Characters. 37 CFR 1.84(f) Reference Characters Poor or Incorrectly Sized. Fig(s) Reference Characters Placed Incorrectly. Fig(s) Reference Must be Numbered Properly.
 3. Character of Lines. 37 CFR Lines Pale or Rough a Fig(s) Solid Black Shading N Fig(s) Telephone inquires cond number (703) 557-6404 	nd Blurred. ot Allowed. cerning this review sho	Figures Must Not be Connected. Fig(s) 7. Photographs Not Approved. 5.5 4 8. Other. Other. ould be directed to the Chief Draftsman at telephone 8.6



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Paul J. Carter et al.

Serial No. 07/715,272

Filed: 14 June 1991

Immunoglobulin Variants For:

Group Art Unit: 1806

Examiner: L. Feisee

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

PATENT DOCKET 709

JUL 2 2 1992

Response

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

This is responseive to the Restriction Requirement mailed 12 May 1992. A request for a onemonth extension of time to respond is submitted herewith, bringing the due date for this response to

11 July 1992. This response is timely filed.

The inventors hereby elect to prosecute Group 1, claims 1-13.

Respectfully submitted, GENENTECH, INC.

10 July 1992

Carolyn R. Adler Reg. No. 32,324

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, on 10 July 1992.

Dated: 10 July 1992

Carolyn R. Adler



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of	
Paul J. Carter et al.	J
Serial No. 07/715,272	
Filed: 14 June 1991	
For: Immunoglobulin Variants	

Group Art Unit: 1806 Examiner: L. Feisee

Sciend

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

PETITION AND FEE FOR EXTENSION OF TIME (37 CFR 1.136(a))

JUL 2 2 1992

PATENT DOCKET 709

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Applicant petitions the Commissioner of Patents and Trademarks to extend the time for response to the Office action dated 12 May 1992 for one month(s) from 11 June 1992 to 11 July 1992. The extended time for response does not exceed the statutory period.

Please charge Deposit Account Number 07-0630 in the amount of \$110 to cover the cost of the extension. Any deficiency or overpayment should be charged or credited to this deposit account. A duplicate of this sheet is enclosed.

Respectfully submitted,

GENENTECH, INC.

Carolyn Ř. Adler Reg. No. 32,324

Date: 10 July 1992

CERTIFICATE OF MAILING (37 CFR 1.88)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and 20231. DY.C.

Jashington. Adler 26 Carolyn

LA10099 07/17/92 07715272

010 07-0630 167

110.00CH

Date: <u>10 July 1992</u>

U.S. DEPARTMENT OF COMMEN ONLINE SEARCH REQUEST-FORM USER _ ABBOLD FEISCE SERIAL NUMBER _ 715272 ART UNIT 1806 PHONE 273 9 DATE Please give a detailed statement of requirements. Describe as specifically as possible the subject matter to be searched. Define any terms that may have special meaning. Give examples or relevant citations, authors, or keywords, if known-You may include a copy of the broadest and or relevant claim(s). Please search. Making Humanezed Ant kodies by . CDR: Grafting . and al agric man and and See claims 1-13 ispécially! 1.2 53 10 FI 3 01 U.S. PÅT 2 TK OFF. / CHELICAL FORECTO STAFF USE ONLY 168

Fersie 715272 9/15/92 show files File 155:MEDLINE_1966-1992/NOV (9211W1) File 5: BIOSIS PREVIEWS_69-92/OCT BA9407: BARRM4307 (C. BIOSIS 1992) 73: EMBASE (EXCERPTA MEDICA) 74-92/ISS37 File (COPR. ESP BV/EM 1992) File 399:CA SEARCH_1967-1992 UD=11710 (Copr. 1992 by the Amer. Chem. Soc.) ?ds Description Items Set HUMANIZED()ANTIBODIES/TI SYNONYN for CDR. 16 S1 ANTIBODIES! FROM 155 332298 IMMUNOGLOBULIN VARIABLE REGION! FROM 155 **S**2 2253 S3 S2 AND S3 2253 S4 HUMANIZ? 862 S5 HUMANIS? 2005 S6 S4 AND (HUMANIZ? OR HUMANIS?) 16 **S**7 ANTIBOD? FROM 5,73,399 636823 S8 IMMUNOGLOBULIN 165469 S9 TG 41830 S10 VARIABLE 113462 S11 REGION 392448 (IMMUNOGLOBULIN OR IG) (W) VARIABLE (W) REGION S12 862 S13 604 CDR S14 COMPLEMENTARY 67991 S15 DETERMINING 112646 S16 COMPLEMENTARY (W) DETERMINING 63 S17 HYPERVARIABLE 1904 S18 (COMPLEMENTARY (W) DETERMINING OR HYPERVARIABLE) (W) REGION REGION 392448 S19 747 S20 ANTIBODY 428778 S21 RELATED 1469126 S22 BINDING 623755 S23 SITE? ? 544344 ANTIBODY (W) RELATED (W) BINDING (W) SITE? ? S24 (IMMUNOGLOBULIN OR IG) () VARIABLE() REGION OR CDR OR (COMPLE-0 S25 MENTARY() DETERMINING OR HYPERVARIABLE) () REGION OR ANTIBODY() R-S26 2161 ELATED () BINDING () SITE? ? FROM 5,73,399 SYNONYMS for COR 8 AND 26 897 S27 27 AND (5 OR 6) 18 S28 28 OR 7 34 S29 RD (unique items) 21 S30 21 Sort S30/ALL/PY,D a S31 ?t31/7/1-21 (Item 1 from file: 5) 31/7/1 BIOSIS Number: 94073885 *HUMANIZED* OKT3 *ANTIBODIES* SUCCESSFUL TRANSFER OF IMMUNE MODULATING 9568885 PROPERTIES AND IDIOTYPE EXPRESSION WOODLE E S; THISTLEWAITE J R; JOLLAFFE L K; ZIVIN R A; COLLINS A; ADAIR J A; BODMER M; ATHWAL D; ALEGRE M-L; BLUESTONE J A SECT. ORGAN TRANSPLANTATION, DEP. SURGERY, WASH. UNIV. SCH. MED., ONE BARNES HOSP. PLAZA, QUEENY TOWER, SUITE 6107, ST. LOUIS, MO. 63110. J IMMUNOL 148 (9). 1992. 2756-2763. Full Journal Title: Journal of Immunology

Language: ENGLISH

, *Antibodies* that possess the Ag-binding regions of OKT3 within the context of a human framework (Hu-OKT3 Ab) offer distinct advantages for optimizing anti-CD3 mAb therapy. First, manipulation of Ab genes to produce *humanized*. Ab that retain Ag-binding activity may circumvent antigenicity problems. Second, Ab gene engineering provides a means for modifying functional properties, including T cell activation and immune suppression. The purpose of this study was to determine the functional properties of Hu-OKT3 Ab and to compare the functional properties and idiotypes of Hu-OKT3 Ab to those of maurine OKT3. Three Hu-OKT3 IgG4 aAb, a chimeric OKT3 *antibody* (cOKT3-1) (grafted sequences comprising all OKT3 VH and VL and two complementarity determining region (*CDR*)-grafted regions) gOKT3-5 and gOKT3-6 (grafted sequences comprising only OKT3 *antibodies* , VH and VL *CDR* and some framework amino acids, were analyzed. Initial studies demonstrated that the cOKT3 and gOKT3-5 Ab bound selectively to T cells and competitively inhibited OKT3-FITC binding with avidities similar to that of murine OKT3. binding avidity of the gOKT3-6 Ab was markedly less than that of the other Hu-OKT3 Ab. Serologic analysis suggested that COKT3 and gOKT3-5 Ab possess idiotypes (combining sites) similar to murine OKT3. C cell activation potency of all three Hu-OKT3 Ab was assessed by proliferation, induction of activation marker expression (IL-2R and Leu 23), and lymphokine production (TNF-.alpha. and IFN-.gamma.). The COKT3 and gOKT3-5 Ab demonstrated T cell activation potencies similar to murine OKT3 as assessed by each parameter. CD3 coating and modulation by these two Ab was effective but somewhat less potent than that observed with OKT3. Finally, COKT3 and gOKT3-5 Ab both inhibited CTL activity comparably to murine OKT3. In conclusion, these studies indicate that gOKT3-5 and COKT3 Ab possess immune modulating properties similar to murine OKT3 and thus offer attractive alternatives to murine OKT3 for in vivo therapy.

(Item 2 from file: 155) 31/7/2

08124424 92262424

Humanization of an anti-p185HER2 antibody for human cancer therapy. Carter P; Presta L; Gorman CM; Ridgway JB; Henner D; Wong WL; Rowland AM;

Kotts C; Carver ME; Shepard HM Department of Protein Engineering, Genentech Inc., South San Francisco, CA 94080.

May 15 1992, 89 (10) p4285-9, Proc Natl Acad Sci U S A (UNITED STATES) Journal Code: PV3 ISSN 0027-8424

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The murine monoclonal antibody mumAb4D5, directed against human epidermal growth factor receptor 2 (p185HER2), specifically inhibits proliferation of human tumor cells overexpressing p185HER2. However, the efficacy of in human cancer therapy is likely to be limited by a human mumAb4D5 anti-mouse antibody response and lack of effector functions. A "*humanized* antibody, humAb4D5-1, containing only the antigen binding loops from mumAb4D5 and human variable region framework residues plus IgG1 constant domains was constructed. Light- and heavy-chain variable regions were simultaneously *humanized* in one step by "gene conversion mutagenesis" using 311-mer and 361-mer preassembled oligonucleotides, respectively. The humAb4D5-1 variant does not block the proliferation of human breast. carcinoma SK-BR-3 cells, which overexpress p185HER2, despite tight antigen binding (Kd = 25 nM). One of seven additional *humanized* variants designed by molecular modeling (humAb4D5-8) binds the p185HER2 antigen 250-fold and 3-fold more tightly than humAb4D5-1 and mumAb4D5, respectively. In addition, humAb4D5-8 has potency comparable to the murine antibody in blocking SK-BR-3 cell proliferation. Furthermore, humAb4D5-8 is much more efficient in supporting antibody-dependent cellular cytotoxicity against SK-BR-3 cells than mumAb4D5, but it does not efficiently kill WI-38 cells, which express p185HER2 at lower levels.

(Item 3 from file: 155) 31/7/3

08081267 92219267

conformation of the the residues affecting framework Antibody hypervariable loops.

Foote J; Winter G

MRC Laboratory of Molecular Biology, Cambridge, England. J Mol Biol (ENGLAND) Mar 20 1992, 224 (2) p487-99, ISSN 0022-2836

Journal Code: J6V

Languages: ENGLISH Document type: JOURNAL ARTICLE

Rodent monoclonal antibodies have been "*humanized*" or "reshaped" for therapy by transplanting the antigen-binding loops from their variable domains onto the beta-sheet framework regions of human antibodies. However, additional substitutions in the human framework regions are sometimes required for high affinity antigen binding. Here we describe antigen binding by a reshaped antibody derived from the mouse anti-lysozyme antibody D1.3, and several variants in which point mutations had been introduced into framework positions to improve its affinity. The affinities were determined from the relaxation kinetics of reactant mixtures using the formation of upon that occurs fluorescence of quenching antibody-antigen complex. The dissociation constant of lysozyme ranged from 3.7 nM (for D1.3) to 260 nM. Measurement of antibody-antigen association kinetics using stopped-flow showed that D1.3 and most of the reshaped antibodies had bimolecular rate constants of 1.4 x 10(6) s-1 M-1, indicating that differences in equilibrium constant were predominantly due to different rates of dissociation of lysozyme from immune complexes. Mutations in a triad of heavy chain residues, 27, 29 and 71, contributed 0.9 kcal/mol in antigen binding free energy, and a Phe to Tyr substitution of light chain residue 71 contributed an additional 0.8 kcal/mol. The combined effect of all these mutations brought the affinity of the reshaped antibody to within a factor of 4 of D1.3. All of these substitutions were the underlying closely beta-sheet framework the complementarity-determining regions, and do not participate in a direct interaction with antigen. The informed selection of residues in such positions may prove essential for the success of loop transplants in antibodies. Variation of these sites may also have a role in shaping the diversity of structures found in the primary repertoire, and in affinity maturation.

(Item 4 from file: 155) 31/7/4 92148135 08010135

Chimeric and *humanized* antibodies with specificity for the CD33 antigen.

Co MS; Avdalovic NM; Caron PC; Avdalovic MV; Scheinberg DA; Queen C

Protein Design Labs, Inc., Mountain View, CA 94043. J Immunol (UNITED STATES) Feb 15 1992, 148 (4) p1149-54, ISSN Journal Code: IFB

0022-1767 Contract/Grant No.: NIH CA55349

Languages: ENGLISH

Document type: JOURNAL ARTICLE

L and H chain cDNAs of M195, a murine mAb that binds to the CD33 Ag on normal and leukemic myeloid cells, were cloned. The cDNAs were used in the construction of mouse/human IgG1 and IgG3 chimeric antibodies. In addition, combined the were constructed which antibodies complementarity-determining regions of the M195 antibody with human *humanized* framework and constant regions. The human framework was chosen to maximize homology with the M195 V domain sequence. Moreover, a computer model of M195 was used to identify several framework amino acids that are likely to interact with the complementarity-determining regions, and these residues

were also retained in the *humanized* antibodies. Unexpectedly, the *humanized* IgG1 and IgG3 M195 antibodies, which have reshaped V regions, have higher apparent binding affinity for the CD33 Ag than the chimeric or mouse antibodies.

(Item 5 from file: 155) 31/7/5

92134790 07996790 conversion of immunoglobulin variable regions in mutagenesis Gene cassettes by replacement PCR mutagenesis.

Near RI.

Molecular Research Laboratory, Massachusetts General Cellular and Hospital, Boston 02144.

Biotechniques (UNITED STATES) Jan 1992, 12 (1) p88-97, ISSN 0736-6205 Journal Code: AN3

Contract/Grant No.: HL-19259

Languages: ENGLISH

Document type: JOURNAL ARTICLE

A technique, Replacement PCR Mutagenesis, was developed to replace one immunoglobulin variable region (V) in a M13 phage cassette with a different, homologous V. This allows the use of the same mutagenesis and subsequent expression vectors for many V regions or V segments. The method combines PCR of V fragments and in vitro mutagenesis. Primers homologous to and 5' ends of both V regions initiate PCR synthesis of the V DNA fragment (donor) that will replace the V region (recipient) in M13. Donor V PCR DNA may originate from mRNA, cloned V genes or genomic templates. The donor V PCR DNA is denatured and annealed to the Mi3 cassette containing the recipient V to be supplanted. The second strand is synthesized, transfected into bacteria and mutant plaques selected by hybridization. Since restriction sites in primers are not required, altered primer-encoded amino acids are avoided. Further, the PCR donor piece can be of any length if it shares homology with the recipient gene. This allows construction and expression of complete gene replacements and chimeras. This method is also applicable to V "*humanization* " and studying sets of homologous genes containing polymorphic or evolutionary disparities. The potential uses of the technique are discussed.

(Item 6 from file: 5) 31/7/6

BIOSIS Number: 42004979 8779979 IMMUNOHISTOCHEMICAL CHARACTERIZATION OF THE *CDR*-GRAFTED *HUMANIZED*

MONOCLONAL *ANTIBODY* BW 431-26 HUMAB PRECLINICAL STUDY

MASCHEK W; BOSSLET K

INST. NUCLEARMED., LINZ BEHRING RES. LABS, MARBURG, FRG.

EUROPEAN ASSOCIATION OF NUCLEAR MEDICINE CONGRESS, VIENNA, AUSTRIA, SEPTEMBER 1-5, 1991. EUR J NUCL MED 18 (8). 1991. 546. CODEN: EJNMD Language: ENGLISH

(Item 7 from file: 5) 31/7/7

BIOSIS Number: 92028624 8563624

POLYMERASE CHAIN REACTION FACILITATES THE CLONING *CDR*-GRAFTING AND RAPID EXPRESSION OF A MURINE MONOCLONAL *ANTIBODY* DIRECTED AGAINST THE CD18 COMPONENT OF LEUKOCYTE INTEGRINS

DAUGHERTY B L; DEMARTINO J A; LAW M-F; KAWKA D W; SINGER I I; MARK G E DEP. CELL. MOL. BIOL., MERCK SHARP DOHME RES. LAB., RAHWAY, N.J. 07065, USA.

NUCLEIC ACIDS RES 19 (9). (1991.) 2471-2476. CODEN: NARHA Full Journal Title: Nucleic Acids Research

Language: ENGLISH

Two novel approaches of recombinant PCR technology were employed to graft the complementarity determining regions from a murine monoclonal *antibody* (mAb) onto human *antibody* frameworks. One approach relied on the availability of cloned human variable region templates, whereas the other strategy was dependent only on human variable region protein sequence data. The transient expression of recombinant *humanized* *antibody* was driven by the adenovirus major late promoter and was detected 48 hrs post-transfection into non-lymphoid mammalian cells. The application of these new approaches enables the expression of a recombinant *humanized* *antibody* just 6 weeks after initiating the cDNA cloning of the murine mAB.

(Item 8 from file: 155) 31/7/8

92187594 08049594

Humanization of a mouse monoclonal antibody by CDR-grafting: the importance of framework residues on loop conformation.

Kettleborough CA; Saldanha J; Heath VJ; Morrison CJ; Bendig MM Medical Research Council Collaborative Centre, London, UK.

Oct 1991, 4 (7) p773-83, ISSN 0269-2139 Protein Eng (ENGLAND) Journal Code: PR1

Languages: ENGLISH

Document type: JOURNAL ARTICLE

A mouse monoclonal antibody (mAb 425) with therapeutic potential was ' *humanized* ' in two ways. Firstly the mouse variable regions from mAb 425 were spliced onto human constant regions to create a chimeric 425 antibody. Secondly, the mouse complementarity-determining regions (CDRs) from mAb 425 were grafted into human variable regions, which were then joined to human constant regions, to create a reshaped human 425 antibody. Using a molecular model of the mouse mAb 425 variable regions, framework residues that might be critical for antigen-binding were identified. To test (FRs) importance of these residues, nine versions of the reshaped human 425 the (VH) regions and two versions of the reshaped human heavy chain variable light chain variable (VL) regions were designed and constructed. The 425 recombinant DNAs coding for the chimeric and reshaped human light and heavy chains were co-expressed transiently in COS cells. In antigen-binding assays and competition-binding assays, the reshaped human antibodies were compared with mouse 425 antibody and to chimeric 425 antibody. The different versions of 425-reshaped human antibody showed a wide range of avidities for antigen, indicating that substitutions at certain positions in the human FRs significantly influenced binding to antigen. Why certain individual FR residues influence antigen-binding is discussed. One version of reshaped human 425 antibody bound to antigen with an avidity approaching that of the mouse 425 antibody.

(Item 9 from file: 155) 31/7/9 07969093 92107093 *Humanization* of monoclonal antibodies. Gussow D; Seemann G 203 p99-121, ISSN 0076-6879 Methods Enzymol (UNITED STATES) 1991, Journal Code: MVA Languages: ENGLISH Document type: JOURNAL ARTICLE (Item 10 from file: 155) 31/7/10 92091750 07953750 expression and characterization of *humanized* antibodies Construction, directed against the human alpha/beta T cell receptor. Shearman CW; Pollock D; White G; Hehir K; Moore GP; Kanzy EJ; Kurrle R Genzyme Corporation, Framingham, MA 01701. Dec 15 1991, 147 (12) p4366-73, ISSN Immunol (UNITED STATES) J Journal Code: IFB 0022-1767 Languages: ENGLISH

Document type: JOURNAL ARTICLE

antibodies with specificity for the human alpha/beta TCR have been produced by genetic engineering. The L and H chain *humanized* V region exons encoding the murine mAb BMA 031 CD regions and human EU framework regions were synthesized and replaced into previously isolated genomic fragments. These fragments were inserted into mammalian expression vectors containing the human kappa and gamma 1 C region exons. Two variants were constructed each containing selected BMA 031 amino acids within the human frameworks. The *humanized* genes were transfected into Sp2/0 hybridoma cells by electroporation and transfectomas secreting *humanized* antibody were isolated. Levels of antibody expression up to 7 pg/cell/24 h were obtained. The *humanized* antibody, BMA 031-EUCIV2, competed poorly with murine BMA 031 for binding to T cells. BMA 031-EUCIV3, however, bound specifically to T cells and competed effectively with both the murine BMA 031 antibody and a previously constructed chimeric BMA 031 antibody for binding to these cells. The relative affinity of BMA 031-EUCIV3 was about 2.5 times lower than BMA 031. The ability to promote antibody dependent cell-mediated cytolysis was significantly enhanced with the engineered antibodies as compared to murine BMA 031. *Humanized* BMA 031 is a clinically relevant, genetically engineered antibody with potential uses in transplantation, graft vs host disease, and autoimmunity.

(Item 11 from file: 155) 31/7/11

Antigenicity of mouse monoclonal antibodies. A study on the variable 07909485

region of the heavy chain. Olsson PG; Hammarstrom L; Smith CI

Karolinska Institute, Huddinge of Clinical Immunology, Department University Hospital, Sweden.

ISSN 0022-5193 151 (x) p111-22, Jul 7 1991, J Theor Biol (ENGLAND)

Journal Code: K8N

Languages: ENGLISH

Mouse monoclonal antibodies (Mabs) against human tumour antigens are currently used in therapy, but up to 50% of the patients receiving treatment form anti-Mab antibodies thus reducing the efficiency of the Document type: JOURNAL ARTICLE One attempt to minimize the immunogenicity of the mouse Mabs is "*humanize* " them by replacing the constant part of the molecule with treatment. the human equivalent by genetic engineering. However, this does not reduce the immunogenicity of the variable part of the antibody. Some variable regions may be expected to be less antigenic than others. We therefore compared consensus sequences for the 11 mouse VH families with the human VH antigenicity predictions sequences published so far. Theoretical and relative accessibility surface flexibility, antigenicity) were made and two families; VH I(J558) and VH XI (CP5 B5-3) were predicted to be immunogenic by all four methods. One family, VH X (MRL-DNA4), was not predicted to be immunogenic by any of the four methods. The residues predicted to form antigenic epitopes in the two families VH II (Q52) and VH III (36-60) are predicted not to be exposed on the surface of the antibody molecule and may therefore not be immunogenic.

31/7/12

1/7/12 (Item 12 from file: 5) 05670 BIOSIS Number: 40106670 CHIMERIC MOUSE-HUMAN AND *CDR*-GRAFTED *ANTIBODIES* TO HUMAN IL2 RECEPTOR 7905670

WEIDLE U H; RUSSMANN E; LENZ H; KALUZA B BOEHRINGER MANNHEIM GMBH, NONNENWALD 2, D-8122 PENZBERG, FRG. MEETING ON MOLECULAR BIOLOGY AND THE IMMUNOPATHOGENESIS OF RHEUMATOID ARTHRITIS HELD AT THE 20TH ANNUAL MEETING OF THE KEYSTONE SYMPOSIA ON MOLECULAR AND CELLULAR BIOLOGY, LAKE TAHOE, CALIFORNIA, USA, MARCH 15-2+, 1991. J CELL BIOCHEM SUPPL 15 (PART E). 1991. 186.

Language: ENGLISH

(Item 13 from file: 155) 31/7/13

07899816 92037816 A *humanized* monovalent CD3 antibody which can activate homologous complement.

Routledge EG; Lloyd I; Gorman SD; Clark M; Waldmann H Department of Pathology, Cambridge University.

Eur J Immunol (GERMANY) Nov 1991, 21 (11) p2/17-25, ISSN 0014-2980

Journal Code: EN5

Languages: ENGLISH Document type: JOURNAL ARTICLE

The rat monoclonal antibody (mAb) YTH12.5, specific for the CD3 antigen complex on human T cells has been modified in order to improve its efficacy in human therapy. With the aim of rendering it less immunogenic, it has been *humanized* using the method of framework grafting. During this process sequence analysis of the YTM12.5 VL gene indicated that it was of lambda subclass, however, it was markedly dissimilar from previously the published rat and mouse V lambda gene sequences and may represent a new V lambda gene family. The *humanization* of this light chain represents the first successful reshaping of a lambda light chain V region. To improve the effector function of the antibødy we have created a monovalent form (1 Fab, 1 Fc) using a novel method/involving the introduction of an N-terminally truncated human IgG1 heavy chain gene into cells producing the *humanized* CD3 mAb. Comparison of the mono- and bivalent *humanized* mAb in a complement-mediated cell/lysis assay revealed that the monovalent antibody mediated lysis of human T cell blasts whereas the bivalent form did not. The availability of a *humanized*, complement-fixing CD3 mAb may improve opportunities for human therapy, in the management of organ rejection, autoimmunity and the treatment of T cell lymphoma.

(Item 14 from file: 155) 31/7/14 91287736 07768736

A possible procedure for reducing the immunogenicity of antibody variable domains while preserving their ligand-binding properties. Padlan EA

Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892.

Apr-May 1991, 28 (4-5) p489-98, ISSN 0161-5890 Mol Immunol QR180+152. Journal Code: NG1

Languages: ENGLISH

Document type: JOURNAL ARTICLE

proposed to reduce the immunogenicity of allogeneic antibody It is domains, while preserving ligand-binding properties, by reducing their antigenicity through replacement of the exposed residues in the variable framework regions which differ from those usually found in host antibodies. The results of a comparison of representative murine antibody sequences with those of human origin suggest that the number of residues that need to be replaced to "*humanize*" those antibodies could be small.

(Item 15 from file: 155) 31/7/15 07757287 91276287

Immunoglobulin complementarity-determining region grafting by recombinant polymerase chain reaction to generate *humanised* monoclonal antibodies.

Lewis AP; Crowe JS

Department of Cell Biology, Wellcome Research Laboratories, Beckenham, Kent, U.K.

ISSN 0378-1119 Journal Code: $QH 442 \cdot \cancel{B}43$ 30 1991, 101 (2) p297-302, Gene May FOP

Languages: ENGLISH

Document type: JOURNAL ARTICLE

We describe an approach to rapidly generate *humanised* monoclonal antibodies by grafting rodent complementarity-determining regions onto human immunoglobulin frameworks using recombinant polymerase chain reaction (PCR) methodology. The approach was applied to grafting a rat complementarily-determining region onto a human framework and amplifying the entire *humanised* heavy chain. The terminal oligodeoxyribonucleotide primers incorporated restriction sites to allow forced cloning into plasmid vectors for sequencing and expression. No nucleotide errors were introduced into the 1463-bp sequence even after sequential applications of PCR.

31/7/16 (Item 16 from file: 155)

07668893 91187893

Journal Code: PV3

Humanized antibodies for antiviral <u>therapy</u>. Co MS; Deschamps M; Whitley RJ; Queen C Protein Design Labs, Inc., Mountain View, CA 94043. Proc Natl Acad Sci <u>U S A</u> Apr 1 1991, 88 (7) p2869-73,

ISSN 0027-8424

Languages: ENGLISH Document type: JOURNAL ARTICLE

Antibody therapy holds great promise for the treatment of cancer, autoimmune disorders, and viral infections. Murine monoclonal antibodies are relatively easy to produce but are severely restricted for therapeutic use by their immunogenicity in humans. Production of human monoclonal antibodies has been problematic. *Humanized* antibodies can be generated by introducing the six hypervariable regions from the heavy and light chains of a murine antibody into a human framework sequence and combining it with human constant regions. We *humanized*, with the aid of computer modeling, two murine monoclonal antibodies against herpes simplex virus gB and gD glycoproteins. The binding, virus neutralization, and cell protection results all indicate that both *humanized* antibodies have retained the binding activities and the biological properties of the murine monoclonal antibodies.

31/7/17 (Item 17 from file: 399)

117024688 CA: 117(3)24688r PATENT

Humanized complementarily-determing region (CDR)-grafted antibodies to intercellular adhesion molecule-1 (ICAM-1), methods of preparation and usage thereof

INVENTOR (AUTHOR): Adair, John Robert; Athwal, Diljeet Singh; Rothlein, Robert A.

LOCATION: UK,

ASSIGNEE: Celltech Ltd.; Boehringer Ingelheim Pharmaceuticals, Inc. PATENT: PCT International ; WO 9116927 A1 DATE: 911114

APPLICATION: WO 91US2942 (910429) *GB 909549 (900427)

PAGES: 81 pp. CODEN: PIXXD2 LANGUAGE: English CLASS: A61K-039/395A; C07K-015/28B DESIGNATED COUNTRIES: AT; AU; BB; BG; BR; CA; CH; DE; DK; ES; FI; GB; HU; JP; KP; KR; LK; LU; MC; MG; MW; NL; NO; PL; RO; SD; SE; SU; US DESIGNATED REGIONAL: AT; BE; BF; BJ; CF; CG; CH; CM; DE; DK; ES; FR; GA; GB; GR; IT; LU; ML; MR; NL; SE; SN; TD; TG

SECTION:

CA215003 Immunochemistry

CA201XXX Pharmacology

CA203XXX Biochemical Genetics

IDENTIFIERS: humanized antibody intercellular adhesion mol 1, inflammation inhibitor humanized antibody ICAM1, asthma inhibitor humanized antibody ICAM1, AIDS virus humanized antibody ICAM1, virucide humanized antibody ICAM1, diagnosis humanized antibody ICAM1 DESCRIPTORS:

Dermatitis...

acute, treatment of, with humanized antibody to intercellular adhesion mol.-1Immunosuppressants... and humanized antibody to intercellular adhesion mol.-1, pharmaceutical compn. contq. anti-intercellular adhesion mol.-1 antibody variable region Rodent.. complementary detg. region of, in humanized antibody prodn. Integrins, antigens LFA-1... antibody to, and humanized antibody to intercellular adhesion mol.-1, for inflammation treatment Neoplasm inhibitors, metastasis... chimeric antibody to intercellular adhesion mol.-1, for hemopoietic cell tumors cytokine-induced, inflammation of, treatment of, humanized antibody to Toxicity... intercellular adhesion mol.-1 for Inflammation... diagnosis of, with chimeric antibody binding to cell expressing intercellular adhesion mol.-1 Deoxyribonucleic acids... for antibody heavy and light chains, in humanized antibody to intercellular adhesion mol.-1 prodn. Deoxyribonucleic acid sequences... for monoclonal antibody R6-5-D6 heavy and light chain components for humanized antiintercellular adhesion mol.-1 antibody human immunodificiency virus infection of, inhibition of, with Leukocyte... humanized antibody to intercellular adhesion mol.-1 Bronchodilators, antiasthmatics... Inflammation inhibitors... Inflammation inhibitors, antirheumatics... Therapeutics... Virucides and Virustats... humanized antibody to intercellular adhesion mol.-1 humanized antibody to intercellular adhesion mol.-1 derivatized with, Toxins... for inhibition of intercellular adhesion mol.-1-expressing tumor cell Diagnosis... humanized antibody to intercellular adhesion mol.-1 for Inflammation inhibitors, antiarthritics... humanized antibody to intercellular adhesion mol.-1, for reaction Glycoproteins, specific or class, ICAM-1 (intercellular adhesion mol. 1)... humanized recombinant antibody to Antibodies... humanized recombinant, to intercellular adhesion mol.-1 Thyroid gland, disease, autoimmune thyroiditis... inflammation in, treatment of, with humanized antibody to intercellular adhesion mol.-1 Nervous system, central... inflammation of, treatment of, humanized antibody to intercellular adhesion mol.-1 for Autoimmune disease... Blood vessel, disease, Raynaud's phenomenon... Brain, disease, stroke... Dialysis, hemo-... Encephalomyelitis... Intestine, disease, Crohn's... Intestine, disease, pseudomembranous enterocolitis... Intestine, disease, ulcerative colitis... Kidney, disease, acute glomerulonephritis... Leukapheresis... Lupus erythematosus... Multiple sclerosis... Psoriasis... Respiratory distress syndrome, adult... inflammation of, treatment of, with humanized antibody to intercellular adhesion mol.-1 Neoplasm, composition... intercellular adhesion mol.-1-expressing, diagnosis of, with humanized

antibody to intercellular adhesion mol.-1 Mouse... monoclonal antibody R6-5-D6 of, in humanized antibody to intercellular adhesion mol.-1 prodn. Sepsis and Septicemia... multiple organ injury syndrome secondary to, inflammation of, treatment of, humanized antibody to intercellular adhesion mol.-1 for Protein sequences... of monoclonal antibody R6-5-D6 heavy and light chain components for humanized antiintercellular adhesion mol.-1 antibody Plasmid and Episome... pAL5, in grafted humanized antibody to intercellular adhesion mol.-1 prodn. Plasmid and Episome... pAL6, in grafted humanized antibody to intercellular adhesion mol.-1 prodn. Plasmid and Episome... pBJ1, in grafted humanized antibody to intercellular adhesion mol.-1 prodn. Kidney, transplant... Organ, transplant... Transplant and Transplantation... rejection of, inflammation of, treatment of, humanized antibody to intercellular adhesion mol.-1 for Antibodies, monoclonal... R6-5-D6, of mouse, in humanized antibody to intercellular adhesion mol.-1 prodn. Organ, disease, multiple organ failure... secondary to septicemia or trauma, treatment of, humanized antibody to intercellular adhesion mol.-1 for Temperature effects, biological... thermal injury, inflammation of, treatment of, humanized antibody to intercellular adhesion mol.-1 for Perfusion, re-... tissue injury from, treatment of, humanized antibody to intercellular adhesion mol.-1 for Lymphokines and Cytokines... toxicity induced by, inflammation of, treatment of, humanized antibody to intercellular adhesion mol.-1 for Neoplasm inhibitors... toxin-derivatized humanized antibody to intercellular adhesion mol.-1, for intercellular adhesion mol.-1-expressing tumor cell Leukocyte, granuločyte... transfusion-assocd. syndrome, treatment of, humanized antibody to intercellular adhesion mol.-1 for Allergy, delayed hypersensitivity... treatment of, humanized antibody to intercellular adhesion mol.-1 for Picornaviridae... Virus, animal, Coxsackie A... Virus, animal, human immunodeficiency... Virus, animal, human immunodeficiency 1... Virus, animal, Mengo... Virus, animal, rhino-... treatment of infection with, with humanized antibody to intercellular adhesion mol.-1 Hematopoietic precursor cell... tumorous, metastasis of, inhibition of, chimeric antibody to intercellular adhesion mol.-1 Genetic vectors ... with DNA for antibody heavy and light chains, in humanized antibody to intercellular adhesion mol.-1 prodn. CAS REGISTRY NUMBERS: 142007-78-1 142007-79-2 142007-80-5 142007-81-6 142007-82-7 142007-83-8 142007-85-0 amino acid sequence of 142007-84-9 amino acid sequence of, humanized antibody to intercellular

adhesion mol.-1 in relation to 140876-28-4 140876-29-5 142007-86-1 142007-87-2 amino acid sequence of, humanized antibody to intercellular adhesion mol.-1 prodn. in relation to 140857-88-1 142008-94-4 nucleotide sequence of, humanized antibody to intercellular adhesion mol.-1 prodn. in relation to 140857-89-2 142008-93-3 nucleotide sequence of, humanized antibody to intercellular adhesion mol.01 prodn. in relation to Copyright 1992 by the American Chemical Society (Item 18 from file: 155) 31/7/18 90356972 07449972 Immunoglobulin V regions of a bactericidal anti-Neisseria meningitidis outer membrane protein monoclonal antibody. J; Fernandez ME; Fry KE; Valle JW; Coloma MJ; del Larrick Gavilondo-Cowley JV Genelabs Inc., Redwood City, California. Aug 1990, 32 (2) p121-8, ISSN 0300-9475 Scand J Immunol Journal Code: UCW Languages: ENGLISH Document type: JOURNAL ARTICLE a potentially therapeutic murine monoclonal antibody that recognizes the class 1 outer membrane protein of Neisseria meningitidis. C6 C6 is specifically immunoblots this antigen and augments in vitro killing of N. meningitidis bacteria. We describe a general method of obtaining the heavy light chain variable-region sequence from immunoglobulin-secreting The method uses mixed polymerase chain reaction (PCR) primers and designed from the 5' end of the framework 1 (FR1) sequences of the heavy chains, and 3'-end primers for constant-region conserved sequences. The method has been applied to the cloning and sequencing of the variable region of C6 to construct a *humanized* monoclonal antibody. Rapid amplification and sequencing of variable regions by this general method have multiple applications in the study of the immune response to infectious diseases. (Item 19 from file: 155) 31/7/19 90199738 07292738 Cloning of the genes for T84.66, an antibody that has a high specificity and affinity for carcinoembryonic antigen, and expression of chimeric human/mouse T84.66 genes in myeloma and Chinese hamster ovary cells. Neumaier M; Shively L; Chen FS; Gaida FJ; Ilgen C; Paxton RJ; Shively JE; Division of Biology, Beckman Research Institute of the City of Hope, Riggs AD Duarte, California 91010. Apr 1 1990, 50 (7) p2128-34, ISSN 0008-5472 Cancer Res Journal Code: CNF Contract/Grant No.: CA 43904 Languages: ENGLISH Document type: JOURNAL ARTICLE antigen (CEA) is one of the best characterized Carcinoembryonic antigens and is extensively used in the in vitro tumor-associated immunodiagnosis of human colon adenocarcinomas. Among a number of anti-CEA monoclonal antibodies, the murine monoclonal antibody T84.66 shows the highest specificity and affinity for CEA and has been used successfully for in vivo tumor imaging in mice and humans. We report here the cloning and sequencing of the genes coding for monoclonal antibody T84.66 and the amino acid sequence of the variable regions for the heavy and light chains. We also report the construction of mouse/human chimeric IgG1 antibody genes using T84.66 variable region genes and human constant region genes. The resulting chimeric gene constructs were transfected into murine myeloma 179

cells (Sp2/0) by electroporation and into Chinese hamster ovary cells by chimeric antibodies obtained exhibited the same specificity and affinity for CEA as that of the T84.66 immunoglobulin produced by the murine hybridoma cell line. Antibody concentrations in culture medium supernatants were clonally variable but similar (15-480 ng/ml) for both Sp2/0 and Chinese hamster ovary transfectants; the average production by Chinese hamster ovary transfectants was only 3-5-fold less than Sp2/0 transfectants. Ascites production of Sp2/0 transfectants is sufficiently high (900 micrograms/ml) for initial in vivo studies with *humanized* T84.66.

(Item 20 from file: 155) 31/7/20 07192290 90099290

A *humanized* antibody that binds to the interleukin 2 receptor. Queen C; Schneider WP; Selick HE; Payne PW; Landolfi NF; Duncan JF; Avdalovic NM; Levitt M; Junghans RP; Waldmann TA

Protein Design Labs, Palo Alto, CA 94304. ISSN 0027-8424 Dec 1989, 86 (24) p10029-33, Proc Natl Acad Sci U S A sot this. Journal Code: PV3

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The anti-Tac monoclonal antibody is known to bind to the p55 chain of the human interleukin 2 receptor and to inhibit proliferation of T cells by blocking interleukin 2 binding. However, use of anti-Tac as an immunosuppressant drug would be impaired by the human immune response against this murine antibody. We have therefore constructed a "*humanized*" antibody by combining the complementarity-determining regions (CDRs) of the anti-Tac antibody with human framework and constant regions. The human framework regions were chosen to maximize homology with the anti-Tac antibody sequence. In addition, a computer model of murine anti-Tac was used to identify several amino acids which, while outside the CDRs, are likely to interact with the CDRs or antigen. These mouse amino acids were also retained in the *humanized* antibody. The *humanized* anti-Tac antibody has an affinity for p55 of 3 x 10(9) M-1, about 1/3 that of murine anti-Tac.

(Item 21 from file: 155) 31/7/21 88178056 06533056

Reshaping human antibodies: grafting an antilysozyme activity.

Verhoeyen M; Milstein C; Winter G Medical Research Council Laboratory of Molecular Biology, Cambridge,

England. ISSN 0036-8075 Mar 25 1988, 239 (4847) p1534-6, Science

Journal Code: UJ7

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The production of therapeutic human monoclonal antibodies by hybridoma technology has proved difficult, and this has prompted the "*humanizing*" of mouse monoclonal antibodies by recombinant DNA techniques. It was shown previously that the binding site for a small hapten could be grafted from the heavy-chain variable domain of a mouse antibody to that of a human myeloma protein by transplanting the hypervariable loops. It is now shown that a large binding site for a protein antigen (lysozyme) can also be transplanted from mouse to human heavy chain. The success of such constructions may be facilitated by an induced-fit mechanism. ?save temp

Temp SearchSave "TD101" stored ?b351,350

15sep92 09:23:49 User209197 Session D126.3 SYSTEM: OS - DIALOG OneSearch File 351: Derwent World Patents Index Latest 1981+;DW=9227,UA=9214,UM=9143 **FILE351: Formats 32,33,35,37 & 39 display the new 'Expanded' Patent For more info. type ?NEWS351 Family table for UD=9216 and greater. File 350: Derwent World Patents Index 1963-1980, EQUIVALENTS THRU DW=9227 **FLLE350:-Formats 32,33,35,37 & 39 display the new 'Expanded' Patent Family table for UD=9219 and greater. For more info. type ?NEWS350 Description Items Set ?exs Executing TD101 HILIGHT set on as '*' 0 HUMANIZED/TI 2945 ANTIBODIES/TI 0 HUMANIZED()ANTIBODIES/TI **S1** >>>File 155 is not open >>>No valid files specified in FROM >>>File 155 is not open >>>No valid files specified in FROM >>>Set "S2" does not exist >>>"S4" does not exist 0 S4 S2 1 HUMANIZ? **S**3 26 HUMANIS? **S**4 S4 AND (HUMANIZ? OR HUMANIS?) 0 **S**5 HILIGHT set on as '*' >>>File 5 is not open >>>File 73 is not open >>>File 399 is not open >>>No valid files specified in FROM >>>File 5 is not open >>>File 73 is not open >>>File 399 is not open >>>No valid files specified in FROM >>>Set "S8" does not exist >>>Set "S27" does not exist >>>Set "S28" does not exist >>>Duplicate detection is not supported for File 351. >>>Duplicate detection is not supported for File 350. >>>All specified files are unsupported, command ignored. >>>Set '30' has not yet been created. = OFF. COST ?ss antibod? and (s3 or s4) 13936 ANTIBOD? S6 **S**3 1 S4 26 22 ANTIBOD? AND (S3 OR S4) ?ss cdr or (ig or immunoglobulin)()variable()region or (complementary()determing Processing Processing CDR 31 S8 S9 786 IG

١.

•	,	•
S10	1576	IMMUNOGLOBULIN
S11	108404	VARIABLE
S12	108131	REGION
S13	4	(IG OR IMMUNOGLOBULIN) (W) VARIABLE (W) REGION
S14	23564	COMPLEMENTARY
S15	501	DETERMING
S16	0	COMPLEMENTARY (W) DETERMING
S17	23	HYPERVARIABLE
S18	108131	REGION
S19	12	(COMPLEMENTARY (W) DETERMING OR HYPERVARIABLE) (W) REGION
S20	11218	ANTIBODY
S21	43127	RELATED
S22	28329	BINDING
S23	29492	SITE? ?
S24	0	ANTIBODY (W) RELATED (W) BINDING (W) SITE? ?
S25	45	CDR OR (IG OR IMMUNOGLOBULIN) ()VARIABLE()REGION OR (COMPLEMENTARY()DETERMING OR HYPERVARIABLE)()REGION OR ANTIBODY()RELATED()BINDING()SITE? ?

?c 7 and 25

?t261

22 7 45 25 87 77 AND 25

(Item 1 from file: 351) 26/7/1 009040436 WPI Acc No: 92-167794/21 XRAM Acc No: C92-077239 New *humanised* *antibody* specific for interleukin-2 receptor - with complementarity determn. regions and framework from different immunoglobulin(s), is non immunogenic and used to treat T-cell Patent Assignee: (PROT-) PROTEIN DESIGN LABS INC Author (Inventor): QUEEN C L; SELICK H E Number of Patents: 001 Number of Countries: 001 Patent Family: Date Week Kind CC Number (Basic) 9221 911219 A5 DD 296964

Priority Data (CC No Date): DD 337159 (900117) Abstract (Basic): DD 296964 A

Compsn. comprises a practically pure human-type immunoglobulin (Ig) that reacts specifically with p55-Tac protein and/or inhibits binding of human interleukin-2 (I1-2) to its specific receptor.

Also new are (1) human-type Ig having 2 pairs of light chain/heavy chain dimers and able to react specifically with an epitope of human IL-2 receptor with affinity at least 10 power 8 M-1, in which the complementarity determining regions (*CDR*) and human-type frame work regions are from different Ig molecules; (2) *humanised* Ig able to bind to IL-2 receptors with one or more *CDR* from anti-Tac *antibody* in a human framework, where the framework includes includes at least one amino acid (AA) from anti-Tac; (3) nucleic acid encoding a human Ig framework and murine *CDR* which, when expressed, produces an Ig specifically reactive with p55-Tac protein and can block binding of IL-2 to its receptor; (4) cells transformed with this nucleic acid.

USE/ADVANTAGES - These Ig are used to treat humans with T-cell related diseases (e.g. transplant rejection; T-cell leukaemia or autoimmune diseases such as diabetes, multiple sclerosis, etc.). They are specific for the IL-2 receptors; are engineered to be

non-immunising and can be produced by recombinant DNA method. The new Ig are admin. in usual parenteral formulation e.g. in doses of 150 mg for therapy or 0.5-2.5 mg for prophylaxis. Ig can also be used, opt. labelled, for diagnosis; T-cell typing; specific receptor isolation or vaccine prodn. 0/10 Derwent Class: B04; D16; Int Pat Class: A61K-039/395; C12N-015/13 (Item 2 from file: 351) 26/7/2 009039793 WPI Acc No: 92-167155/20 Prepn. of chimeric *humanised* *antibodies* - using a new polymerase XRAM Acc No: C92-076891 chain reaction technique; PCR Patent Assignee: (WELL) WELLCOME FOUND LTD Author (Inventor): CROWE J S; LEWIS A P Number of Patents: 001 Number of Countries: 015 Patent Family: Week Date Kind CC Number (Basic) 9220 920430 A1 WO 9207075 Priority Data (CC No Date): GB 9022011 (901010) Applications (CC, No, Date): WO 91GB1744 (911008) Language: English EP and/or WO Cited Patents: 4.Jnl.Ref; WO 9007861 Designated States (Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; SE Abstract (Basic): WO 9207075 Prodn. of ds or ss DNA of formula: 5' F1-M-F2 3' encoding an *antibody* (Ab) chain or fragment in which at least one of the complementarily determining regions (CDRs) of the variable region is derived from a first mammalian Ab and the framework of the variable region is derived from a second different mammalian Ab, where M is DNA encoding a *CDR* of the second Ab and F1 and F2 resp. encode 5' and 3' sequences flanking M, by: (a) prepg. a ss or ds DNA template of formula: 5' f1-H-f2 3' where H is DNA encoding a *CDR* of a different specificity from M, and f1 and f2 are homologous to F1 and F2, resp.; (b) obtaining DNA oligonucleotide primers A, B, C and D, where: A comprises the sequence al with a 5' end corresp. to the 5' and of F1 and which is identical to the corresp. length of F1 and is oriented in a 5' to 3' direction towards H; B has of the sequence 5' b1-b2 3', where bl comprises a sequence complementary to a corresp. length of M and has a 3' end complementary to the 5' end of M, and b2 is complementary to a sequence of corresp. length in F1 and has a 5' end which starts at the nucleotide complementary to the 3' end of F1, C has of the sequence 5' c1-c2 3' where c1 comprises a sequence identical to the corresp. length of M and has a 3'end corresp. to the 3' end of M, and c2 is identical to a sequence of corresp. length in F2 and has a 5' end which starts at the nucleotide corresp. to the 5' end of F2, and D comprises a sequence d1 which has a 5' end complementary to the 3' end of F2 and which is complementary to a corresp. length of F2 and is oriented in a 5' to 3' direction towards H, where b1 and c1 overlap by a sufficient length to permit annealing of their 5' ends under conditions which allow PCR to be performed; (c) performing, in any desired order, PCR reactions with primer pairs A, B and C, D on the template prepd. in (a), and (d) mixing the prods. of (c) and performing PCR using primers A and D. USE/ADVANTAGE - The method allows the prepn. of chimeric,

esp. *humanised* Abs. The resulting Ab retains the antigen binding

capability of the non-human Ab from which the *CDR*(s) are derived. 0/4 Derwent Class: B04; D16; Int Pat Class: C12N-005/10; C12N-015/12; C12N-015/69; C12P-021/08 (Item 3 from file: 351) 26/7/3 008937440 WPI Acc No: 92-064709/08 XRAM Acc No: C92-029621 New multivalent anti-cytokine immunoglobulins - for treating disorders associated with elevated cytokine levels, e.g. septic and endotoxic shock, AIDS, allergies, etc.; ACQUIRE IMMUNE DEFICIENT SYNDROME Patent Assignee: (CLLT) CELLTECH LTD; (CELL-) CELLTECH LTD Author (Inventor): ALLEN R A; MORGAN S A Number of Patents: 002 Number of Countries: 035 Patent Family: Week Kind Date CC Number (Basic) 9208 920206 Α WO 9201472 9222 920218 Α AU 9182381 Priority Data (CC No Date): GB 9015908 (900719) Applications (CC, No, Date): AU 9182381 (910719); WO 91GB1216 (910719) EP and/or WO Cited Patents: 2.Jnl.Ref; EP 347057; EP 355067; WO 9006371; WO 9007118; WO 9106305 (National): AT; AU; BB; BG; BR; CA; CH; CS; DE; DK; ES; FI; GB; HU; JP; KP Designated States ; KR; LK; LU; MC; MG; MW; NL; NO; PL; RO; SD; SE; SU; US (Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; OA; SE Filing Details: AU9182381 Based on WO 9201472 Abstract (Basic): WO 9201472 New multivalent immunoglobulin (I) has at least 3 linked antigen-binding domains (ABD's) each being specific for a complementary site on a cytokine. The combining interactions between ABD and cytokine sites are neutralising. (I) is specific for tumour necrosis factor (TNF) alpha or beta; an interleukin, an interferon or a colony-stimulating factor, and it contains 4-20 ABD. ABD are all of class IgG (most pref.) or all of class IgM (but must be different from a native IgM molecule) and can be linked by covalent crosslinking (e.g. 2-iminothiolane/ maleimide system) or by non-covalent interaction (e.g. using an *antibody* reactive with sites on Ig other than those involved in antigen binding; or the biotin-avidin system). (I) are made by joining together appropriate immunoglobulin molecules or fragments esp *CDR*-grafted or *humanised* chimaeric Ig. USE/ADVANTAGE- (I) are used to treat or prevent diseases assciated with elevated cytokine levels, e.g. immuno regulatory and inflammatory disease, sepsis, endotoxic or cardiovascular shock, AIDS, psoriasis, organ transplant rejection or excessive TNF generation induced cancer therapy etc., Compared with monomeric Ig, (I) have much greater neutralising activity. @(43pp)@ Derwent Class: B04; D16; Int Pat Class: A61K-039/39; A61K-039/395; C07K-015/28; C12P-021/08 (Item 4 from file: 351) 26/7/4 008929605 WPI Acc No: 92-056874/07 Related WPI Accession(s): 91-222915 New *cdr*-grafted anti carcinoembryonic antigen *antibodies* - useful XRAM Acc No: C92-025713 in therapy and diagnosis of carcinoma

Patent Assignee: (CELL-) CELLTECH LTD Author (Inventor): ADAIR J R; BODMER M W; MOUNTAIN A; OWENS R J Number of Patents: 001 Patent Family: Week Date Kind CC Number 9207 v (Basic) 920123 Priority Data (CC No Date): WO 91GB1108 (910705); GB 9014932 (900705); WO 90GB2017 (901221) EP and/or WO Cited Patents: WO 8910140; WO 8901783; EP 323806; 6.Jnl.REF Designated States (National): AT; AU; BB; BG; BR; CA; CH; CS; DE; DK; ES; FI; GB; HU; JP; KP ; KR; LK; LU; MC; MG; MW; NL; NO; PL; RO; SD; SE; SU; US (Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; SE; OA Abstract (Basic): WO 9201059 New *humanised* *antibody* molecule (HAM) is specific for carcino-embryonic antigen (CEA) and has an antigen binding site in which at least one of the complementarity determining regions (*CDR*'s) of the variable domain is derived from the mouse monoclonal *antibody* (MAb) A5B7. The remaining Ig-derived parts of HAM are of human origin. HAM is a chimeric or *CDR*-grafted *humanised* *antibody*, prepd. by recombinant DNA techniques. It can be a complete *antibody* or an Fab, Fab', (Fab')2 or Fv fragment, or a single-chain fragment. It may have a reporter or effector molecule attached to it. USE/ADVANTAGE - HAM are useful in therapy or diagnosis (including imaging) of carcinomas which produce CEA, e.g., when coupled to a toxin such as ricin. @(70pp Dwg.No.0/19 Derwent Class: B04; D16; Int Pat Class: A61K-039/39; C07K-015/28; C12N-015/13; C12P-021/08 (Item 5 from file: 351) 26/7/5 008849515 WPI Acc No: 91-353533/48 XRAM Acc No: C91-152448 New *humanised* *CDR*-grafted anti-ICAM *antibodies* - used to treat and prevent inflammation (e.g. psoriasis) tumours, viral infections and asthma and in diagnosis; INTER CELLULAR ADHESIVE MOLECULAR Patent Assignee: (CELL-) CELLTECH LTD; (BOEH) BOEHRINGER INGELHEIM PHA Author (Inventor): ADAIR J R; ATHWAL D S; ROTHLEIN R A Number of Patents: 002 Patent Family: Week Date CC Number Kind (Basic) 9148 911114 WO 9116927 А 9210 911127 AU 9179001 Α Priority Data (CC No Date): GB 909549 (900427) Applications (CC, No, Date): WO 91US2942 (910429) Language: English EP and/or WO Cited Patents: US 4816567; WO 8901783; 7.Jnl.REF (National): AT; AU; BB; BG; BR; CA; CH; DE; DK; ES; FI; GB; HU; JP; KP; KR Designated States ; LK; MC; MG; MW; NL; NO; PL; RO; SD; SE; SU; US (Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; SE; OA Abstract (Basic): WO 9116927 A recombinant *antibody* molecule comprising antigen binding regions derived from the heavy and/or light chain variable regions of an anti-intracellular adhesion molecule-1 (anti-ICAM-1) *antibody* is claimed. The Ab is *CDR*-grafted and comprises several non-human residues. Also claimed are DNA encoding an Ab heavy or light chain, a vector comprising the DNA, host cells transformed with the vector and a method for producing the anti-ICAM-1 grafted Ab.

USE/ADVANTAGE - The Abs are used to treat - and prevent

inflammation in e.g. delayed type hypersensitivity, psoriasis, an autoimmune disease e.g. Reynaud7s syndrome, autoimmune thyroiditis, EAE, multiple sclerosis, rheumatoid arthritis and lupus erythematosus, tissue or organ transplant or graft rejection. They are also used to treat and prevent tumours, viral infections (e.g. rhinoviruses of the major serotype within the genus Picornavididae, group A coxsackievirus, a Mengo virus and HIV); asthma and non-specific defence system response, e.g. adult respiratory distress syndrome, CNS inflammatory disorder, multiple organ injury syndrome secondary to septicaemia or trauma, ulcerative colitis and Crohn's disease. Administration can be enteral, parenteral, topical, intranasal or by inhalation. The are also used to diagnose an ICAM-1-expressing tumour cell and

inflammation. @(68pp Dwg.No.0/4 Derwent Class: B04; D16; Int Pat Class: A61K-039/39; C07K-015/28

(Item 6 from file: 351) 26/7/6 008718897 WPI Acc No: 91-222916/30

XRAM Acc No: C91-096865 CD3 specific *humanised* recombinant *antibody* - is chimeric or *cdr* grafted for immunotherapy and diagnosis; COMPLEMENTARY DETERMINE REGION

Week

9130 9143 (Basic)

Patent Assignee: (CELL-) CELLTECH LTD Author (Inventor): JOLLIFFE L K; ZIVIN R A; ADAIR J R; ATHWAL D S Number of Patents: 003

Patent Family:

	ramral.		
CC	Number	Kind	Date
	9109968	А	910711
	9170330	A	910724
	22/6201	Δ	920212

9207 Priority Data (CC No Date): WO 90GB2018 (901221); GB 8928874 (891221); GB 9117611 (910815)

Applications (CC,No,Date): GB 9017611 (901221)

Language: English

EP and/or WO Cited Patents: EP 403156; EP 328404 Designated States

(National): AT; AU; BB; BG; BR; CA; CH; DE; DK; ES; FI; GB; GR; HU; JP; KR ; LK; LU; MC; MG; MW; NL; NO; RO; SD; SE; SU; US

(Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; SE; OA Filing Details: GB2246781 Based on W09109968 (E) (1251CH)

Abstract (Basic): WO 9109968

A recombinant *antibody* (RAM) comprising antigen binding regions derived from the heavy and or light chain variable regions of a donor anti- CD3 *antibody*. The *antibody* preferably has binding affinity similar to that of OKT3. The RAM comprises antigen binding regions from suitable anti-CD3 *antibodies* such as rodent e.g. mouse or rat anti-CD3 MAb. The RAM may comprises only the variable region (VH and/or VL) or one or more CDRs of such a MAb.

The RAM is preferably a *humanised* *antibody* molecule specific for CD3 having an antigen binding site where at least one of the CDRs of the variable domain and usually two more of the CDRs are derviced from non human anti-CD3 *antibody*. The RAM may be a chimeric or *CDR* grafted *antibody*. Usually, the donor and acceptor *antibodies* are derived from different species. Typically the donor anti CD3 *antibody* is non-human (e.g. rodent) and the acceptor *antibody* is human. A *CDR* grafted *antibody* heavy chain comprising variable region with acceptor and donor CD3 binding comprising donor residues at one or more of positions 6, 37, 48 and 94. The *CDR* grafted light chain is also claimed.

DNA coding these *antibodies* and their production by recombinant DNA technology is claimed.

USE/ADVANTAGE - The *antibodies* may be used for treatment or diagnosis of human or veterinary conditions. The *humanised* *antibodies* do not have the immunologic complications associated with administration of non human *antibodies* to human subjects. @(81pp Dwg.No.0/13)@ Derwent Class: B04; D16; Int Pat Class: A61K-039/39; A61K-049/00; C07K-015/06; C12N-005/10; C12N-015/13; C12P-021/08 (Item 7 from file: 351) 26/7/7 WPI Acc No: 91-222915/30 008718896 Related WPI Accession(s): 92-056874 XRAM Acc No: C92-025713 New *humanised* *antibodies* comprising *CDR* grafted *antibody* - with heavy and light chains, for use in vivo therapy and diagnosis; COMPLEMENTARY DETERMINE REGION Patent Assignee: (CLLT) CELLTECH LTD; (CELL-) CELLTECH LTD Author (Inventor): ADAIR J R; BODMER M W; MOUNTAIN A; OWENS R J; ATHWAL D S ; EMTAGE J S Number of Patents: 005 Number of Countries: 035 Patent Family: Week Date X CC Number Kind (Basic) 9130 91.07.11 WO 9109967. Α 9143 910724) AU 9169740 Α 920205 9206 А GB 2246570 9207 920123 Ά WO 9201059 9220 920204 AU 9182005 Α WO 90GB20174 (901221); GB 891221) Priority Data (CC No Date): GB 8928874 9014932 (900705) Applications (CC, No, Date): AU 9182005 (910705); WO 91GB1108 (910705); GB 9017612 (901221) Language: English EP and/or WO Cited Patents: EP 239400; EP 323806; EP 328404; EP 403156; 6.Jnl.Ref; WO 8901783; WO 8910140 Designated States (National): AT; AU; BB; BG; BR; CH; DE; DK; FI; GB; HU; JP; KP; KR; LK; LU ; MC; MG; MW; NL; NO; RO; SD; SE; SU; US; CA; CS; ES; PL (Regional): AT; BE; CH; DE; FR; GB; GR; IT; LU; NL; OA; SE; DK; ES Filing Details: AU9182005 Based on WO 9201059 Abst/ract (Basic): WO 9109967 A *CDR* grafted *antibody* heavy chain is claimed having a variable region comprising acceptor frame-work and donor antigen binding regions in at least one of positions 6, 23 and/or 24, 48 and/or 49, 71 and/or 73, 75 and/or 76 and/or 78 and 88 and/or 91. Preferably the heavy chain framework also comprises donor residues at positions 37, 48 and 94. Also claimed is a *CDR*-grafted *antibody* light chain having a variable region domain comprising acceptor framework and donor antigen binding regions comprising donor residues in at least one of positions 1 and/or 3 and preferably at positions 46 and/or 47. A *CDR* grafted *antibody* molecule is also claimed comprising at least one *CDR* grafted heavy chain and light chain. DNA encoding the *CDR* grafted heavy and light chains is also claimed. The heavy or light chains may have an effector or reporter molecule attached e.g. a macrocycle for chelating a metal atom or a toxin such as ricin. The *CDR* grafted *antibodies* preferably have non-human e.g. rodent donor and human acceptor frameworkers. USE/ADVANTAGE - For use in treatment and diagnosis of human and veterinary conditions. @(91pp Dwg.No.0/13

Derwent Class: B04; D16; Int Pat Class: A61K-039/39; A61K-039/395; C07K-015/06; C07K-015/28; C12N-005/10; C12N-015/13; C12P-021/08; C12R-001/91 (Item 8 from file: 351) 26/7/8 WPI Acc No: 90-253800/33 008366799 XRAM Acc No: C90-109897 Chimaeric immunoglobulin(s) blocking IL-2 binding to receptors comprising human framework and murine complementary determining regions, less immunogenic than murine *antibodies* Patent Assignee: (PROT-) PROTEIN DESIGN LABS INC; (PROT-) PROTEIN DESIGN LABS Author (Inventor): QUEEN C L; SELICK H E Number of Patents: 010 Number of Countries: 034 Patent Family: Week Kind Date CC Number 9033 (Basic) 900726 Α WO 9007861 9033 900629 PT 92758 А 9037 900628 CA 2006865 А 9044 900813 AU 9051532 А 9048 901031 ZA 8909956 А 9115 900718 Α CN 1043875 9133 910520 Α FI 9102436 9142 910619 Α NO 9102385 9143 Α 910619 DK 9101191 9225 920507 W JP 4502408 Priority Data (CC No Date): US 290975 (881228); US 310252 (890213) Applications (CC, No, Date): WO 89US5857 (891228); JP 90503677 (891228); ZA 899956 (891228) Language: English; German EP and/or WO Cited Patents: 7.Jnl.Ref; EP 239400; GB 2188941; US 4816567; WO 8901783 Designated States (National): AT; AU; BB; BG; BR; CH; DE; DK; FI; GB; HU; JP; KP; KR; LK; LU ; MC; MG; MW; NL; NO; RO; SD; SE; SU (Regional): AT; BE; CH; DE; ES; FR; GB; IT; LU; NL; OA; SE Filing Details: JP04502408 Based on WO 9007861 Abstract (Basic): WO 9007861 Compsn. comprises a pure human-like immunoglobulin (Ig) which (a) reacts specifically with p55 Tac protein and/or (b) inhibits binding of human interleukin-2 (IL-2) to its receptor. Also new are (1) human-like Ig having 2 pairs of light/heavy chains and able to react specifically with an epitope of a human IL-2 receptor with affinity at least 10 power 8 per mole, the chains including complementarily determg. regions (*CDR*'s) and human-like framework regions (FR's), the *CDR*'s being from different Ig molecules than FR's; (2) *humanised* Ig (hIg) which can bind to IL-2 receptors and contain at least one *CDR* from anti-Tac *antibody* in a human-like FR contg. at least one amino acid from the anti-Tac *antibody*; (3) nucleic acid encoding for human-like FR and at least one murine *CDR*, and (4) cells transfected with nucleic acid. USE/ADVANTAGE - hIG are not significantly immunogenic in humans; are easily and economically produced, and have a longer half-life in vivo than mouse *antibodies*. They are useful (opt. when attached to a cytotoxic agent, for treatment of T-cell mediated disorders, e.g. graft or transplant rejection, and autoimmune diseases. LIG can also be used in vitro for T-cell typing; isolation of IL-2 receptor bearing cells, vaccine prodn., etc. @(52pp Dwg.No.0/10)@ Abstract (EP): 9142 EP 451216

Compsn. comprises a pure human-like immunoglobulin (Ig) which (a) reacts specifically with p55 Tac protein and/or (b) inhibits binding of human interleukin-2 (IL-2) to its receptor. Also new are (1) human like Ig having 2 pairs of light/heavy chains and able to react specifically with an epitope of a human IL-2 receptor with affinity at least 10 power 8 per mole, the chains including complementarily determg. regions (*CDR*'s) and human-like framework regions (FR's) the *CDR*'s being from different Ig molecules than FR's. (2) *humanised* IG (hIg) which can bind to IL-2 receptors and contain at least one *CDR* from anti-Tac *antibody* in a numan-like FR contg. at lesdt one amino acid from the anti-Tac *antibody*, (3) nucleic acid encoding for human-like FR and at least one murine *CDR*, and (4) cells transfected with nucleic acid.

USE/ADVANTAGE - hIG are not significantly immunogenic in humans, are easily and economically produced, and have a longer half-life in vivo than mouse *antibodies*. They are useful (opt. when attached to a cytotoxic agent, for treatment of T-cell mediated disorders, e.g. graft or transplant rejection, and autoimmune diseases, LIG can also be used in vitro for T-cell typing, isolation of IL-2 receptor bearing cells, vaccine prodn etc.

189

Derwent Class: B04; D16;

?

Int Pat Class: A61K-039/39; C07K-007/10; C07K-013/00; C07K-015/14; C12N-005/10; C12N-007/01; C12N-015/00; C12P-021/08

: • set cost off COST = OFF. ?set hi * HILIGHT set on as '*' Hilight option is not available in file(s) 399. ?show files File-155:MEDLINE-1966-1992/NOV (9211W1) 5:BIOSIS PREVIEWS 69-92/OCT BA9407:BARRM4307 File (C. BIOSIS 1992) File 73: EMBASE (EXCERPTA MEDICA) 74-92/ISS37 (COPR. ESP BV/EM 1992) File 3.99: CA SEARCH 1967-1992 UD=11710 (Copr. 1992 by the Amer. Chem. Soc.) ?ds Set Description Items HUMANIZED()ANTIBODIES/TI **S1** 16 ANTIBODIES! FROM 155 S2 332298 IMMUNOGLOBULIN VARIABLE REGION! FROM 155 2253 **S**3 2253 S2 AND S3 **S**4 HUMANIZ? 862 **S**5 HUMANIS? 2005 S6 S4 AND (HUMANIZ? OR HUMANIS?) 16 **S**7 ANTIBOD? FROM 5,73,399 636823 **S**8 IMMUNOGLOBULIN S9 165469 41830 IG S10 VARIABLE S11 113462 REGION S12 392448 (IMMUNOGLOBULIN OR IG) (W) VARIABLE (W) REGION 862 S13 604 CDR S14 67991 COMPLEMENTARY S15 DETERMINING 112646 S16 COMPLEMENTARY (W) DETERMINING S17 63 HYPERVARIABLE 1904 S18 392448 REGION S19 (COMPLEMENTARY (W) DETERMINING OR HYPERVARIABLE) (W) REGION S20 747 428778 ANTIBODY S21 1469126 RELATED S22 S23 623755 BINDING SITE? ? S24 544344 ANTIBODY (W) RELATED (W) BINDING (W) SITE? ? S25 n (IMMUNOGLOBULIN OR IG) () VARIABLE () REGION OR CDR OR (COMPLE-S26 2161 MENTARY()DETERMINING OR HYPERVARIABLE)()REGION OR ANTIBODY()R-ELATED()BINDING()SITE? ? FROM 5,73,399 S8 AND S26 897 S27 sort S30/ALL/PY, D COMPLEMENTARITY DETERMIN? REGION COMPLEMENTARITY(W) DETERMIN? (W) REGION COMPLEMENTARITY() DETERMIN? (W) REGION COMPLEMENTARITY() DETERMIN? (W) REGION COMPLEMENTARITY() DETERMIN? (W) REGION COMPLEMENTARITY() DETERMIN? (W) REGION COMPLEMENTARITY (W) DETERMIN? (W) REGION AND (S5 OR S6) AND S8 7 OR 36 (37 OR 29) NOT 29 190 HOULD HOUL S27 AND (S5 OR S6) S28 18 S29 34 21 \$30 S31 21 3165 S32 2005813 S33 S34 524927 358 ____COMPLEMENTARITY (W) DETERMIN? (W) REGION----- 9 S35 S36 12 28 S37 8 S38

RD (unique items) \$39 6 6 Sort S39/ALL/PX,D ?t40/7/1-6

(Item 1 from file: 5) 40/7/1

BIOSIS Number: 93066780 9081780

DEVELOPMENT OF *HUMANIZED* BISPECIFIC *ANTIBODIES* REACTIVE WITH CYTOTOXIC LYMPHOCYTES AND TUMOR CELLS OVEREXPRESSING THE HER2 PROTOONCOGENE SHALABY M R; SHEPARD H M; PRESTA L; RODRIGUES M L; BEVERLEY P C L;

FELDMANN M; CARTER P

DEP. CELL BIOL., GENENTECH, INC., 460 POINT SAN BRUNO BOULEVARD, SOUTH SAN FRANCISCO, CALIF. 94080.

J EXP MED 175 (1). 1992. 217-226. CODEN: JEMEA

Full Journal Title: Journal of Experimental Medicine Language: ENGLISH

transmembrane 185-kD protooncogene encodes а HER2 phosphoglycoprotein, human epidermal growth factor receptor 2 (p185HER2), The whose amplified expression on the cell surface can lead to malignant transformation. Overexpression of HER2/p185HER2 is strongly correlated with progression of human ovarian and breast carcinomas. Recent studies have shown that human T cells can be targeted with bispecific *antibody* to against human tumor cells in vitro. We have developed a bispecific react F(ab')2 *antibody* molecule consisting of a *humanized* arm with a specificity to 185HER2 linked to another arm derived from a murine anti-CD3 monoclonal *antibody* that we have cloned from UCHT1 hybridoma. The antigen-binding loops for the anti-CD3 were installed in the context of human variable region framework residues, thus forming a fully *humanized* BsF(ab')2 fragment. Additional variants were produced by replacement of amino acid residues located in light chain *complementarity* *determining* *region* 2 and heavy chain framework region 3 of the *humanized* anti-CD3 Flow cytometry analysis showed that the bispecific F(ab')2 molecules arm. can bind specifically to cells overexpressing p185HER2 and to normal human peripheral blood mononuclear cells bearing the CD3 surface marker. In experiments, the presence of bispecific F(ab')2 caused up to additional enhancement in the cytotoxic activities of human T cells against fourfold cells overexpressing p185HER2 as determined by a 51Cr release assay. tumor These bispecific molecules have a potential use as therapeutic agents for the treatment of cancer.

(Item 2 from file: 399) 40/7/2

CA: 117(7)68366p PATENT 117068366

Chimeric and complementarity-determining region-grafted anti-carcingembryonic antigen antibodies and their production

INVENTOR (AUTHOR): Adair, John Robert; Bodmer, Mark William; Mountain, Andrew; Owens, Raymond John

LOCATION: UK,

ASSIGNEE: Celltech Ltd.

PATENT: PCT International ; WO 9201059 A1 DATE: 920123 APPLICATION: WO 91GB1108 (910705) *GB 9014932 (900705) *WO 90GB2017 (901221)

PAGES: 70 pp. CODEN: PIXXD2 LANGUAGE: English CLASS: C12P-021/08A; A61K-039/395B; C12N-015/13B; C07K-015/28B DESIGNATED COUNTRIES: AT; AU; BB ; BG; BR; CA; CH; CS; DE; DK; ES; FI; GB; HU; JP; KP; KR; LK; LU; MC; MG; MN; MW; NL; NO; PL; RO; SD; SE; SU; US DESIGNATED REGIONAL: AT; BE; BF; BJ ; CF; CG; CH; CI; CM; DE; DK; ES; FR; GA; GB; GN; GR; IT; LU; ML; MR; NL; SE; SN; TD; TG

SECTION:

1

CA215003 Immunochemistry

IDENTIFIERS: carcinoembryonic antigen humanized chimeric antibody,

complementarity detg region grafted antibody CEA, cloning DNA humanized antibody CEA **DESCRIPTORS:** Antibodies, monoclonal... A5B7 murine, to carcinoembryonic antigen, in humanized antibody prodn. Animal cell line... CHO L761 h, humanized anti-carcinoembryonic antigen antibody recombinant prodn. in Deoxyribonucleic acid sequences... for antibody variable regions in humanized anti-carcinoembryonic antigen antibody prodn. Genetic vectors... Molecular cloning... for humanized anti-carcinoembryonic antigen antibody prodn. Diagnosis... Therapeutics... humanized anti-carcinoembryonic antigen antibodies for Escherichia coli... humanized anti-carcinoembryonic antigen antibody fragment recombinant prodn. in Animal cell line,CHO-K1... Animal cell line,COS-1... Bacteria... humanized anti-carcinoembryonic antigen antibody recombinant prodn. in Mammal... humanized anti-carcinoembryonic antigen antibody recombinant prodn. in cells of Immunoglobulins, fusion products... humanized, prodn. of Antibodies... humanized, to carcinoembryonic antigen Immunoglobulins... in humanized anti-carcinoembryonic antigen antibody prodn. Protein sequences... of antibody variable regions in humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Eiisme... pAL43, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pAL44, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pAL45, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pAL46, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pAL53, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pAL54, for humanized anti-carcinoembryonic antigen antibody prodn. Genetic vectors... pEE6hCMV gpt, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pHMC19, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pHMC30, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pHMC31, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pHMC43, for humanized anti-carcinoembryonic antigen antibody prodn. Plasmid and Episome... pHMC44, for humanized anti-carcinoembryonic antigen antibody prodn. Genetic vectors... pMRR028, for humanized anti-carcinoembryonic antigen antibody fragment prodn.

Genetic vectors...

pMRR045, for humanized anti-carcinoembryonic antigen antibody fragment prodn.

CAS REGISTRY NUMBERS:

142661-53-8 142661-54-9 142661-55-0 142661-56-1 142661-57-2 142661-58-3 amino acid sequence of, humanized anti-carcinoembryonic antigen antibody prodn. in relation to

142662-72-4 142662-81-5 142662-71-3 142662-70-2 142662-69-9 142662-82-6 nucleotide sequence of, humanized anti-carcinoembryonic antigen antibody prodn. in relation to

Copyright 1992 by the American Chemical Society

(Item 3 from file: 5) 40/7/3

BIOSIS Number: 92064131 8599131

IMMUNOGLOBULIN *COMPLEMENTARITY*-*DETERMINING* *REGION* GRAFTING BY RECOMBINANT POLYMERASE CHAIN REACTION TO GENERATE *HUMANIZED* MONOCLONAL *ANTIBODIES*

LEWIS A P; CROWE J S

DEP. CELL BIOLOGY, WELLCOME RES. LAB., LANGLEY COURT, BECKENHAM, KENT, BR3 3BS UK.

GENE (AMST) 101 (2). 1991. 297-302. CODEN: GENED Full Journal Title: GENE (Amsterdam)

Language: ENGLISH

describe an approach to rapidly generate *humanised* monoclonal *antibodies* by grafting rodent complementarity-determining regions into human immunoglobulin frameworks using recombinant polymerase chain reaction applied to grafting a rat was approach The methodology. (PCR) onto a human framework and *region* *complementarity*-*determining* terminal chain. The *humanised* heavy entire amplifying the oligodeoxyribonucleotide primers incorporated restriction sites to allow forced clonign into plasmid vectors for sequencing and expression. No nucleotide errors were introduced into the 1463-bp sequence even after sequential applications of PCR.

(Item 4 from file: 5) 40/7/4

BIOSIS Number: 40113269 7912269 CONSTRUCTION OF *HUMANIZED* *ANTIBODIES* AND_TESTING IN PRIMATES QUEEN C; CO M S; DESCHAMPS M; WHITLEY R; BENJAMIN W; HAKIMI J PROTEIN DESIGN LAB. INC., 2375 GARCIA AVE., MOUNTAIN VIEW, CALIF. 94043. MEETING ON MONOCLONAL ANTIBODIES HELD AT THE 20TH ANNUAL MEETING OF THE KEYSTONE SYMPOSIA ON MOLECULAR AND CELLULAR BIOLOGY, DENVER, COLORADO, USA, MARCH 10-16, 1991. J CELL BIOCHEM SUPPL 15 (PART E) - 1991. 137. CODEN: JCBSD

Language: ENGLISH

(Item 5 from file: 5) 40/7/5

BIOSIS Number: 89052006 7400987 A *HUMANIZED* *ANTIBODY* THAT BINDS TO THE INTERLEUKIN 2 RECEPTOR QUEEN C; SCHNEIDER W P; SELICK H E; PAYNE P W; LANDOLFI N F; DUNCAN J F; AVDALOVIC N M; LEVITT M; JUNGHANS R P; WALDMANN T A

PROTEIN DESIGN LABS., 3181 PORTER DRIVE, PALO ALTO, CALIF. 94304. PROC NATL ACAD SCI U S A 86 (24). 1989. 10029-10033. CODEN: PNASA Full Journal Title: Proceedings of the National Academy of Sciences of

the United States of America

Language: ENGLISH The anti-Tac monoclonal *antibody* is known to bind to the p55 chain of the human interleukin 2 receptor and to inhibit proliferation of T cells by 2 binding. However, use of anti-Tac as an interleukin blocking immunosuppressant drug would be impaired by the human immune response

against this murine *antibody*. We have therefore constructed a " *humanized*" *antibody* by combining the complementarity-determining regions (CDRs) of the anti-Tac *antibody* with human framework and constant regions. The human framework regions were chosen to maximize homology with the anti-Tac *antibody* sequence. In addition, a computer model of murine anti-Tac was used to identify several amino acids which, while outside the CDRs, are likely to interact with the CDRs or antigen. These mouse amino acids were also retained in the *humanized* *antibody*. The *humanized* anti-Tac *antibody* has an affinity for p55 of 3 .times. 109 M-1, about 1/3 that of murine anti-Tac.

40/7/6 (Item 6 from file: 399)

113170316 CA: 113(19)170316b PATENT Recombinant antibodies to Campath-1 antigen, containing foreign complementarity determining region(s), and their use in immunosuppression and cancer therapy INVENTOR(AUTHOR): Waldmann, Herman; Clark, Michael Ronald; Winter, Gregory Paul; Riechmann, Lutz

LOCATION: UK,

ASSIGNEE: Medical Research Council

PATENT: PCT International ; WO 8907452 A1 DATE: 890824 APPLICATION: WO 89GB113 (890210) *GB 883228 (880212) *GB 884464 (880225) PAGES: 61 pp. CODEN: PIXXD2 LANGUAGE: English CLASS: A61K-039/395A; C12N-015/00B DESIGNATED COUNTRIES: AU; DK; JP; US

SECTION:

CA215003 Immunochemistry

CA201XXX Pharmacology

CA203XXX Biochemical Genetics

IDENTIFIERS: chimeric antibody Campath 1 antigen, lymphoma neoplasm inhibitor Campath 1H antibody DESCRIPTORS:

Rat...

complementarity detg. regions of, in recombinant antibody to Campath-1 antigen

Immunoglobulins, G2... Immunoglobulins, G3... Immunoglobulins, G4...

const. domains of human, in recombinant antibody contg. complementarity detg. regions to Campath-1 antigen

Lymphocyte...

depletion of, in human, by recombinant human antibody contg. foreign complementarity detg. regions to Campath-1 antigen

Gene and Genetic element, animal, synthetic...

for humanized light chain variable region, construction of, in prodn. of recombinant human antibody contg. rat complementarity detg. regions to Campath-1 antigen

Protein sequences...

of IgG2a YTH 34.5 HL heavy and light chain variable domains, of rat Deoxyribonucleic acid sequences, IgG2a-specifying...

of rat

Antigens, CAMPATH-1...

recombinant antibodies to, foreign complementarity detg. regions in Immunosuppressants... Neoplasm inhibitors... Neoplasm inhibitors,lymphoma

recombinant antibody contg. foreign complementarity detg. regions to Campath-1 antigen as

Gene and Genetic element, animal...

recombinant, for anti-Campath-1 antigen antibody of human, sequences encoding rat complementary detg. regions in

Immunoqlobulins, G2a...

recombinant human antibody to Campath-1 antigen contg. complementary detg. regions of rat

Leukemia, B-cell... recombinant human antibody to Campath-1 antigen killing leukemia cells of Antibodies... recombinant, to Campath-1 antigen, foreign complementarity detg. regions in Immunoglobulins, G1... Immunoglobulins, G... Immunoglobulins, M... recombinant, to Campath-1 antigen, foreign complementary detg. regions in CAS REGISTRY NUMBERS: 129711-40-6 amino acid sequence encoded by HuVLLYS gene 129711-41-7 amino acid sequence encoded by synthetic HuVLLYS.degree. gene 129711-01-9 129711-02-0 cloning and nucleotide sequence of, of human and 129711-19-9 129711-20-2 cloning and nucleotide sequence of, of rate 128096-06-0 128096-07-1 128096-08-2 128096-09-3 128096-10-6 complementarity detg. region of rat YTH 34.5 HL, human 128096-11-7 recombinant antibody contg., Campath-1 antigen binding by 129711-56-4 heavy chain variable region of human contg. rat complementarity detg. regions, recombinant antibody contg., Campath-1 antigen binding by 129711-60-0 heavy chain variable region of rat YTH 34.5 HL, recombinant antibody contg., Campath-1 antigen binding by 129710-86-7P HuVLLYS gene, prepn. of, in prepn. of recombinant human antibody contg. rat complementarity detg. regions to Campath-1 antigen 129711-59-7 light chain variable region of human contg. rat complementarity detg. regions, recombinant antibody contg., Campath-1 antigen binding by light chain variable region of rat YTH 34.5 HL, recombinant 129711-61-1 antibody contg., Campath-1 antigen binding by 127859-21-6P 127859-23-8P 127859-24-9P 127859-26-1P 127859-62-5P 127859-79-4P 127859-82-9P 127859-92-1P 127859-70-5P 127859-72-7P 127859-99-8P 127860-01-9P 127860-02-0P 127859-94-3P 127859-93-2P 129924-59-0P prepn. of, in 129924-57-8P 127860-03-1P 127860-04-2P gene synthesis for recombinant human antibody contg. rat complementarity detg. regions to Campath-1 antigen 129711-58-6 recombinant human antibody contg., Campath-1 129711-57-5 antigen binding by 129710-91-4P synthetic gene HuVLLYS.degree., prepn. of, in prepn. of recombinant human antibody contg. rat complementary detg. regions to Campath-1 antigen Copyright 1992 by the American Chemical Society ?b351,350 15sep92 10:26:26 User209197 Session D127.2 SYSTEM:OS - DIALOG OneSearch File 351: Derwent World Patents Index Latest 1981+;DW=9227,UA=9214,UM=9143 **FILE351: Formats 32,33,35,37 & 39 display the new 'Expanded' Patent Family table for UD=9216 and greater. For more info. type ?NEWS351 File 350:Derwent World Patents Index 1963-1980, EQUIVALENTS THRU DW=9227 **FILE350: Formats 32,33,35,37 & 39 display the new 'Expanded' Patent Family table for UD=9219 and greater. For more info. type ?NEWS350 Set Items Description ?ds

Description Items Set ANTIBOD? AND (HUMANIS? OR HUMANIZ?) 22 S1 S1 AND (CDR OR (IG OR IMMUNOGLOBULIN) () VARIABLE() REGION OR S2 8 HYPERVARIABLE() REGION) S1 AND COMPLEMENTARITY()DETERMIN?()REGION 0 \$3 S1 AND COMPLEMENT? () DETERMIN? () REGION 3 **S**4 (2 OR 4) NOT 2 1 s5_ ?t5/7/1 (Item 1 from file: 351) 5/7/1 WPI Acc No: 89-085403/11 007820291 XRAM Acc No: C89-037905 Recombinant *humanised* *antibody* specific for TAG-72 - having complementarity determining regions of variable domains from mouse *antibody* and the remainder from human immunoglobulin Patent Assignee: (CELL-) CELLTECH LTD Author (Inventor): BODMER M W; ADAIR J R; WHITTLE N R Number of Patents: 001 Patent Family: Week CC Number Kind Date (Basic) 8911 890309 А WO 8901783 Priority Data (CC No Date): WO 88GB731 (880905); GB 8720833 (870904) Language: English EP and/or WO Cited Patents: No.SR.Pub; 4.Jnl.REF Designated States (National): AU; DK; FI; HU; JP; KR; NO; RO; SU; US (Regional): AT; BE; CH; DE; FR; GB; IT; LU; NL; SE Abstract (Basic): WO 8901783 A *humanised* *antibody* molecule (HAM) is claimed having specificity for the TAG-72 antigen and having an antigen binding site in which at least the *complementary* *determining* *region* (CDRs) of the variable domains are derived from the mouse monoclonal *antibodies* (MAb) B72.3 and the remaining immunoglobulin-derived parts of the HAM are derived from a human immunoglobulin. USE/ADVANTAGE - *Humanising* the B72.3 MAb does not adversely affect its binding activity and this produces a HAM which is useful in both therapy and diagnosis of certain carcinomas, e.g. solid tumours expressing TAG-72. @(49pp Dwg.No.0/13)@ Derwent Class: B04; D16; Int Pat Class: A61K-039/39; C12N-015/00; C12P-021/00 ?s complement?()determin?(w)region? ? Processing Processing Processing COMPLEMENT? 27431 DETERMIN? 234285 REGION? ? 124968 COMPLEMENT?()DETERMIN?(W)REGION? ? 23 **S6** ?c 1 and 6 1 22 23 6 1 AND 6 10 **S**7 ? ?c 7 not (2 or 4) 196

10 7 8 2 3 4 3 7 NOT (2 OR 4) S8 ?t8/7/1-3 (Item 1 from file: 351) 8/7/1 WPI Acc No: 92-132139/16 009004842 XRAM Acc No: C92-061892 *Humanisation* of *antibodies* binding to human CD4 antigen - by mutation of framework-encoding regions of DNA encoding variable domain of rat or mouse *antibody* chain Patent Assignee: (GORM/) GORMAN S D Author (Inventor): CLARK M R; COBBOLD S P; GORMAN S D; WALDMANN H Number of Patents: 001 Number of Countries: 018 Patent Family: Date Week Kind CC Number 9216 (Basic) 920402 WO 9205274 А Priority Data (CC No Date): GB 9020282 (900917) Applications (CC, No, Date): WO 91GB1578 (910916) Language: English EP and/or WO Cited Patents: 7.Jnl.Ref; EP 328404; EP 365209; EP 403156; WO 9007861; WO 9107492; WO 9109966; WO 9109967 Designated States (National): AU; CA; JP; KR; US (Regional): AT; BE; CH; DE; DK; ES; FR; GB; GR; IT; LU; NL; SE Abstract (Basic): WO 9205274 A *Complementarity* *determining* *regions* (CDRs) of the variable domain of the *antibody* chain are derived from a first mammalian species and the framework of the variable domain and any constant domains of the Ab chain are derived from a second different mammalian species; comprising (a) mutating the framework-encoding regions of DNA encoding a variable domain of the first mammalian Ab chain such that it encodes the framework derived from the second species; and (b) expressing the Ab chain using this mutated DNA. The process specifically comprises: (i) determining nucleotide and predicted aminoacid sequence of a variable domain of a selected Ab chain of the first species; (ii) determining the Ab framework to which the framework of this domain is to be altered; (iii) mutating framework-encoding regions of DNA encoding this variable domain such that the mutated region encodes the framework determined in (ii); (iv) linking mutated DNA to DNA encoding a constant domain of the second species and cloning the DNA into an expression vector; and (v) introducing expression vector into a compatible host cell and culturing it to express Ab chain. USE/ADVANTAGE - Altered Abs is prepd., used to *humanise* an Ab, typically a monoclonal Ab and, e.g. a rat or mouse Ab. The resulting Ab retains the antigen binding capabilities of the Ab from which it is derived. Reshaped CD4 Ab is used to induce tolerance against an antigen. Used to alleviate autoimmune diseases e.g. rheumatoid arthritis, and to prevent graft rejection. 0/13Derwent Class: B04; D16; Int Pat Class: A61K-039/39; C12N-015/13; C12P-021/08 (Item 2 from file: 351) 8/7/2 WPI Acc No: 91-216983/30 008712964

XRAM Acc No: C91-094177

Prodn. of *humanised* recombinant immunoglobulin - including polymerase chain reaction amplification of murine *antibody* light and heavy chain variable portions

Week

(Basic)

Patent Assignee: (MERI) MERCK & CO INC

Author (Inventor): LAW M F; MARK G E; WILLIAMSON A R Number of Patents: 002 Patent Family:

Date

CC Number Kind

EP 438310A9107249130CA 2034553A9107209139

Priority Data (CC No Date): US 627423 (901220); US 467700 (900119) Applications (CC,No,Date): EP 91300362 (910117) Language: English

EP and/or WO Cited Patents: EP 239400; WO 8901783; 1.Jnl.REF Designated States

(Regional): CH; DE; FR; GB; IT; LI; NL

Abstract (Basic): EP 438310

Method for producing a *humanised* recombinant immunoglobulin comprises: (a) prepg. polymerase chain reaction (PCR) primers to amplify the variable portion of the light and heavy chain of a murine *antibody* which binds to a predefined antigen; (b) using the primers to amplify the variable portions of both heavy and light chains and sequencing the resulting nucleotide chains; (c) determining the murine *complementary* *determining* *regions* of the heavy and light chains; (d) selecting human variable heavy and light chain frameworks which show a high degree of amino acid similarity with the variable heavy and light chain framework of the murine immunoglobulin; (e) selecting human constant heavy and light chain frameworks; (f) grafting the murine *complementary* *determining* *regions* of (c) to the human framework regions of (e); (g) incorporating the complete DNA sequence for the *humanised* recombinant immunoglobulin into an appropriate expression vector; (h) transfecting host cells with the vector; (i) growing the transfected cells in an environment in which the *humanised* recombinant immunoglobulin is expressed; and (j) collecting the immunoqlobulin.

A PCR method for the simultaneous synthesis and assembly of at least 4 deoxyoligonucleotides is also claimed.

USE/ADVANTAGE - The *humanised* recombinant immunoglobulins are weakly immunogenic or non-immunogenic when admin. to humans, and may be used as therapeutic agents. Recombinant human anti-CD18 *antibodies* or active fragments which bind to the CD18 antigen of leukocytes can be used to inhibit influx of the leukocytes into a site of inflammation or tissue liable to become inflamed following influx. @(78pp Dwg.No.0/38)@ Derwent Class: B04; D16;

```
Int Pat Class: C12N-015/13; C12P-021/08; C12Q-001/68
```

8/7/3 (Item 3 fro	m file: 351)				
007275804 WPI Acc No:	87-272811/3	9			
YRAM ACC No: C87-11582	5				
Pecombinant altere	d *antibodie	s* - hay	ving *compl	ementarity	*
determining *reg	ions* replac	ed with	those from	*antibody	* of
different specific	ity		·		
Patent Assignee: (WINT	/) WINTER G	Р	•		
Author (Inventor): WIN	TER G P				
Number of Patents: 004			٠		
Patent Family:					
CC Number Kind	Date	Week	·.		
EP 239400 A	870930	8739	(Basic)		
GB 2188638 A	871007	8740			

GB 2188638 B 900523 9021
 Priority Data (CC No Date): GB 867679 (860327); GB 877252 (870326)
 Applications (CC,No,Date): EP 87302620 (870326); JP 8773980 (870327)
 Language: English
 EP and/or WO Cited Patents: A3...8914; 3.Jnl.REF

EP and/or WO Cited Patents: A3...0914, 5.0011.NB1 Designated States

Α

JP 62296890

(Regional): AT; BE; CH; DE; ES; FR; GB; GR; IT; LI; LU; NL; SE Abstract (Basic): EP 239400

871224

An altered *antibody* in which at least parts of the *complementary* *determining* *regions* (CDRs) in the light or heavy chain variable domains have been replaced by analogous parts of CDRs from an *antibody* of different specificity is new.

8806

The altered *antibody* can be produced by (a) prepg. a first replicable expression vector including a suitable promoter operably linked to a DNA sequence which encodes at least a variable domain of an Ig heavy or light chain, the variable domain comprising framework regions from a first *antibody* and CDRs comprising at least parts of the CDRs from a second *antibody* of different specificity, (b) if necessary, prepg. a second replicable expression vector including a suitable promoter operably linked to a DNA sequence which encodes at least the variable domain of a complementary Ig light or heavy chain, (c) transforming a cell line with the first or both prepd. vectors and (d) culturing the transformed cell line to produce the altered *antibody*.

USE/ADVANTAGE - The method is used for ''*humanising*'' non-human monoclonal *antibodies* (MAbs) e.g. CDRs from mouse MAb can be partially or totally grafted into the framework regions of a human MAb, which is then produced in quantity by a suitable cell line. Only the CDRs of the *antibody* will be foreign to the body and this should minimise side effects if used for human therapy. @(41pp Dwg.No.0/8)@

Derwent Class: B04; D16; Int Pat Class: C12N-015/00; C12P-021/02; C07K-015/00; A61K-039/39; C12N-005/00; C12R-001/91

?ds

Set S1 S2	Items 22 8	Description ANTIBOD? AND (HUMANIS? OR HUMANIZ?) S1 AND (CDR OR (IG OR IMMUNOGLOBULIN)()VARIABLE()REGION OR
	НУ	(PERVARIABLE() REGION)
S3	· 0	S1 AND COMPLEMENTARITY()DETERMIN?()REGION
S4	3	S1 AND COMPLEMENT? () DETERMIN? () REGION
S5	1	(2 OR 4) NOT 2
S6	23	COMPLEMENT? () DETERMIN? (W) REGION? ?
S7	10	1 AND 6
S8	3	7 NOT (2 OR 4)
S9	5	S1 AND CDRS
SHO.		(9 OR 7 OR 2 OR 4) NOT (7 OR 2 OR 4)
·>		

	077715.27?	05/11/91		Patent and T	redomerk Off	MENT OF COMME Ice TENTS AND TRADEMAR
[SERIAL NUMBER	FILING DATE	, FIRST	NAMED INVENTOR		ATTORNEY DOCKET
_	GENENTECH.				1.1.1.25.1	
		AN RE ADLER AN ERUND BL				EVANUED.
	SOUTH GAN F	TANCI CO, C	.201 26 - 94086	•	1866	EXAMINER
	•					. •
					ART UNIT	I PAPER NUMBER
						13
		•				1-
	t				DATE MAILED:	
This i COM	 a communication from the MISSIONER OF PATENTS 	examiner in charge of yes AND TRADEMAINS	ur application.		-	
			l			
	· .					
х ты	is application has been t	exemined	Perponsive to comn	nunication filed on	113/92] This action is made final
				3	ab	
	tened statutory pariod fi to respond within the p			monti		ays from the date of this let 3
		,	,.			•
Part I	THE FOLLOWING	TTACHMENT(S) ARE	E PART OF THIS AC	TION:		
1.	Notice of Reference	e Cited by Examiner, i	PTO-892.		Patent Drawing, PTC	
1	Notice of Art Cited	by Applicant, PTO-144	19. 3page	4. D Notice of I	nformal Patent Appl	lication, Form PTO-152.
8.	Information on How	10 Effect Drawing Chi	anges, PTO-1474.	مستحمد السا 6.		
Part II	SUMMARY OF AC	TION				
Part II			1600			
Part II 1.>	SUMMARY OF AC	тюн (6			are pending in the applic
Part II 1>	Claima	<u> </u>	16 4 - 16			
ı)	Claims	<u> </u>	16 4 - 16			are pending in the applic withdrawn from coneidera
ı)	Claima	<u> </u>	1 E 4 - 16			
1) 2	Claims	<u> </u>	1 6 4 - 16			withdrawn from considere
1) 2 3	Claims	(((4	1 6 4 - 16			withdrawn from considera _ have been cancelled.
1) 2 3	Claima Ol the above	(((4	1 6			withdrawn from considera _ have been cancelled.
1>	Claima Of the above Claima Claima Claima	(((4	1 6 4 - 16 			withdrawn from considere have been cancelled, ere ellowed. ere rejected.
1>	Claims	(((4	1 & 4 - 16			withdrawn from considera have been cancelled. are allowed.
1) 2 3 4 8	Claima Of the above Claima Claima Claima	(((4	169 4 - 16	er	0/0	withdrawn from considere have been cancelled, ere ellowed. ere rejected.
1) 2 3 4 3 8	Claime OI the above OI the above Claime Claime Claime Claime Claime Claime Claime Claime Claime	 , cleires[4 1_13	4 - 16		ore exubject to restriction	withdrawn from considera
1	Claima Of the above Claima Claima Claima Claima Claima Claima	1 14 , cleires 14 1 1.3 s been filed with intervi	4 - 16 a) d/sewinge under 33	7 C.F.R. 1.85 which are	ore exubject to restriction	withdrawn from considera
1	Claima Of the above Claima Claima Claima Claima Claima Claima	1 14 , cleires 14 1 1.3 s been filed with intervi	4 - 16 a) d/sewinge under 33	7 C.F.R. 1.85 which are	ore exubject to restriction	withdrawn from considera
1) 2 3 4 3 5 7. 5	Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims	 , cleines (4 13 a been flied with into ⁻ⁿⁱ e required	4 - 16 al drawinge under 37 to this Office ection.	7 C.F.R. 1.85 which are	e subject to restriction	withdrawn from considera
1) 2 3 4 3 8. 4 7. 5	Claims Claims Claims Claims Claims Claims Claims Claims Claims This application hat Formal drawings ar The corrected or au	1 14 a been filed with intervi e required 1 - feaponee batterie drewings have	4 - 16 al drawings under 3; to this Office action.	7 C.F.R. 1.85 which are	e subject to restrictly acceptable for example	withdrawn from considera
1.) 2. 3. 4. 3. 8. 8. 7. 7. 8.	Claims Claims Claims Claims Claims Claims Claims Claims This application had Formal drawings ar The corrected or au are ecospitable	, cleines (4 , cleines (4 ,) 13 s been filed with inter- s cleines(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4	4 — 16 al drawinge under 33 to this Office action. been received on _ ee explanation or No	7 C.F.R. 1.85 which are	e subject to restrictly acceptable for example	withdrawn from considera
1.) 2. 3. 4. 3. 8. 7. 8. 8. 8.	Claims Claims Claims Claims Claims Claims Claims Claims This application has Formal drawings er The corrected or su are acceptable The proposed additioned	I	4 — 16 al drawinge under 37 to this Office action. been received on _ bee explanation or No bet(e) of drawinge, file	7 C.F.R. 1.85 which are	ore o	withdrawn from considera
1.) 2. 3. 4. 3. 8. 7. 8. 8. 8.	Claims Claims Claims Claims Claims Claims Claims Claims This application has Formal drawings er The corrected or su are acceptable The proposed additioned	, cleines (4 , cleines (4 ,) 13 s been filed with inter- s cleines(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4)(4	4 — 16 al drawinge under 37 to this Office action. been received on _ bee explanation or No bet(e) of drawinge, file	7 C.F.R. 1.85 which are	ore o	withdrawn from considera
1.) 2. 3. 4. 3. 8. 7. 8. 8. 8.	Claima Of the above Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima Claima This application has are acceptable The corrected or au are acceptable	I , claims I I a been filed with inter- a required response ballwie drawings have ballwie drawings have ballwie drawings have ballwie drawings have ballwie drawings have proved by the expension	A - 16 al drawings under 3: to this Office action. been received on - we explanation or Ne est(e) of drawings, file er (ese explanation).	7 C.F.R. 1.85 which are otice re Patent Drawing d on	ore o	withdrawn from considera - have been cancelled, - are allowed. - are rejected, - are objected to. on or election requirement mination purposes. - R. 1.84 these drawings - approved by the
2 2 3 4 3 8 3 8 5 5 10. [11. [Claims Claims Claims Claims Claims Claims Claims Claims Claims This application hat Formal drawings ar The corrected or au are acceptable The proposed addifi- examiner, diag	, cleines (4 , cleines))))))))))))))))))))))))))))))))	4 — 16 al drawinge under 37 to this Office action. been received on _ bee explanation or No et(e) of drawinge, file er (eee explanation).	7 C.F.R. 1.85 which are otice re Patent Drawing d on 	ore o	withdrawn from considera
2 2 3 4 3 8 3 8 5 5 10. [11. [Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims This application hat Formal drawings ar The corrected or au are ecceptable Construction additionation The proposed additionation Claims The proposed drawing	, cleines (4 , cleines))))))))))))))))))))))))))))))))	4 - 16 al drawinge under 37 to this Office action. been received on _ bee explanation or No et(e) of drawinge, file er (eee explanation).	7 C.F.R. 1.85 which are otice re Patent Drawing d on 	ore o	withdrawn from considera - have been cancelled, - are allowed. - are rejected, - are objected to. on or election requirement mination purposes. - R. 1.84 these drawings - approved by the
2 2 3 4 3 8 3 8 5 5 10. [11. [Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims This application hat Formal drawings ar The corrected or au are ecceptable Construction additionation The proposed additionation Claims The proposed drawing	, cleines (4 , cleines))))))))))))))))))))))))))))))))	4 - 16 al drawinge under 37 to this Office action. been received on _ bee explanation or No et(e) of drawinge, file er (eee explanation).	7 C.F.R. 1.85 which are otice re Patent Drawing d on 	ore o	withdrawn from considera have been cancelled. are allowed. are relacted. are objected to. on or election requirement mination purposes. C.R. 1.84 these drawings approved by the ved (see explanation).
3. 4. 7. 9. 10. 11. [12. [13. [Claims Of the above Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims	1 12 a been filed with internities a beginned fragming bave beginned fragming bave beginned fragming bave beginned or substitute sheep over a by the examination of the claim for pent application, serial results to be in concerning appears to be in concerning a beginned to beginned to be in concerning a beginned to beginned	4 - 16 al drawinge under 37 to this Office action. been received on to explanation or No been splanation or No been received on to this Office action. been received on been received on been received on to this Office action. been received on to this Office action. been received on been receive	7 C.F.R. 1.85 which are oilce re Patent Drawing d on	ore o	withdrawn from considera have been cancelled. are allowed. are relacted. are objected to. on or election requirement mination purposes. C.R. 1.84 these drawings approved by the ved (see explanation).
1) 2. 3. 4. 7. 8. 7. 8. 10. 11. [11. [11. [11.]	Claims Claims Claims Claims Claims Claims Claims Claims Claims This application hat Formal drawings ar The corrected or au are acceptable The proposed additi examiner dear The proposed drawing Acknowledgment is 	1 12 a been filed with internities a beginned fragming bave beginned fragming bave beginned fragming bave beginned or substitute sheep over a by the examination of the claim for pent application, serial results to be in concerning appears to be in concerning a beginned to beginned to be in concerning a beginned to beginned	4 - 16 al drawinge under 37 to this Office action. been received on to explanation or No been splanation or No been received on to this Office action. been received on been received on been received on to this Office action. been received on to this Office action. been received on been receive	7 C.F.R. 1.85 which are oilce re Patent Drawing d on	ore o	withdrawn from considera have been cancelled. c are allowed. are relected. are objected to. on or election requirement mination purposes. E.R. 1.84 these drawings approved by the ved (see explanation). sived [
1) 2. 3. 4. 7. 9. 10. 11. (12. [13. [Claims Of the above Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims Claims	1 12 a been filed with internities a beginned fragming bave beginned fragming bave beginned fragming bave beginned or substitute sheep over a by the examination of the claim for pent application, serial results to be in concerning appears to be in concerning a beginned to beginned to be in concerning a beginned to beginned	4 - 16 al drawinge under 37 to this Office action. been received on to explanation or No been splanation or No been received on to this Office action. been received on been received on been received on to this Office action. been received on to this Office action. been received on been receive	7 C.F.R. 1.85 which are oilce re Patent Drawing d on	ore o	withdrawn from considera have been cancelled. c are allowed. are relected. are objected to. on or election requirement mination purposes. E.R. 1.84 these drawings approved by the ved (see explanation). sived [

PTOL-326 (Rev. 9-89)

24.

Applicant's election of Group 1, in Paper No. 12, is acknowledged. Because applicant did not distinctly and specifically point out the supposed errors in the restriction requirement, the election has been treated as an election without traverse. See M.P.E.P. 818.03(a).

Claims 1-10 are rejected under 35 U.S.C. § 112, second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject, matter which applicant regards as the invention. Claims 1, 3, 4, 5 and 7 are indefinite in the use of the language "import antibody" in that it is not 10 clear what constitutes an important antibody, ie. what the determines what is to be an import antibody. Claim 1 step a) is indefinite in that it is not clear what is meant by a "consensus" human variable domain". Claim 1 step d) is indefinite in that it 15 is not clear what is actually taking place when one aligns the amino acid sequences of the FR, ie. is this a physical or mental step? Claim 1 step e) is unclear in what type of homology is indicated, ie. are conservative amino acids considered as homologs or should their be identical amino acid residues at the indicated portion of the framework. Claim 1 step f), 3 is indefinite in the 20 language "participates" in that the use of the nature of participation is unclear. Claim 1 step f) is indefinite in that it is not clear how one of ordinary skill can determine the effects which are listed in steps 1-3, ie. through antigen binding, through 25 hybridization? Claim 1 step g) is indefinite in that it is not clear what effects are reasonably expected to occur. Claim 2 is indefinite in that the antecedent basis for "the domain" is unclear. Claim 3 is indefinite in that it is not clear when in the antibody one would search for the of making the process glycosylation sites. Claim 4 is indefinite for the same reason 30 that claim \mathscr{X} is indefinite. Claim 5 is indefinite in that it is believed that the claims up to this point were directed to making a



Serial No. 715272

Art Unit 1806

"humanized antibody", and it is unclear how "preparing a humanized antibody" in claim 5 differs from the preparation of the antibody up to this point. Furthermore, it is not clear what is intended in the preparation of the antibody of claim 5. Claim 6 is vague in 5 that it is not clear what the numbers are meant to designate. lt is suggested that applicant clarify the nature of the numbers or Claim 7 is indefinite in that it is not clear point to a figure. what the method is drawn to. it is suggested that the language "a method of making a humanized antibody" be inserted within the claim.

The following is a quotation of the first paragraph of 35 U.S.C. § 112:

specification shall contain written The а description of the invention, and of the manner and process of making and using it, in such full, clear, concise, and exact terms as to enable any person skilled in the art to which it pertains, or with which it is most nearly connected, to make and use the same and shall set forth the best mode contemplated by the inventor of carrying out his invention.

specification is objected to under 35 U.S.C. 5 112, first The paragraph, as failing to adequately describe the invention and failing to adequately teach how to make and or use the invention, ie. failing to provide an enabling disclosure. The following terms lack enablement in the specification:.

Claims 1 and 7 lack enablement in the language "at least a portion of an import variable domain". Applicant has only indicated specific residues which may be transferred, but they are claiming an antibody wherein the a portion of the import antibody are to be transferred. There is no guidance in the specification which would enable one of skill in the art to make antibodies with

15

10

20

25

transferred variable domains other than CDRs. Applicant is aware that a portion of the variable domain can be any one of the CDRs as well as the framework regions. However, this language also reads on small amino acid sequences which are incomplete regions of the 5 variable region of the antibody. There is no support in the specification for linking the variable region of the antibody to any or all of the myriad "portions" which are encompassed within this language. One of skill in the art would neither expect nor predict the appropriate functioning of the antibody as broadly as 10 is claimed. It is suggested that the specific portion of the human variable region which is described in the specification be recited within the claim or this language be removed completely in order to obviate this rejection.

Claim 1 step c) lacks enablement in that it is not clear how 15 one would determine which amino acids are to be substituted. There is no specific recitation of what characteristics of the amino acids are necessary for deciding whether it is to be replaced or not. Without this description one of skill in the art would not be able to choose the appropriate amino acid residues without 20 hindering the function of the antibody.

Claim 1 step f), lacks enablement in that the protocol for determining whether the amino acid residues in the import amino acid sequence are reasonably expected to interact with the antigen is not described anywhere in the specification. There is no

10

15

20

explicit step which enables one of ordinary skill in the art to determine the effects which are recited. It would require undue experimentation of one of ordinary skill in the art to make the variations which may be made in order to test the effects of the mutant antibodies.

Claim 2 lacks enablement in that there is no description in the specification of how to determine which residues are exposed on the surface or which residues are buried within the domain, is this through computer modeling or through x-ray crystallography or other methods?

Claim 3 lacks enablement in that there is no guidance in the specification on how one would determine which glycosylation site affects antigen binding, or what comprises "reasonable expectation".

Claims 5, 7 and 9 lack enablement in that it would appear that these amino acids are relevant to IgG and not to other isotypes. There is no indication that one of skill in the art would extrapolate the use of these amino acids to all or other isotypes of immunoglobulins. Furthermore, there is insufficient description and guidance in the specification with regards to the properties of these amino acids which would enable one of ordinary skill in the art to make humanized antibodies with other isotypes using these amino acid sequences.

Applicant has not shown that antibodies which have been modified as that which is claimed are capable of functioning as that which is being disclosed, ie. maintaining the binding affinity Protein chemistry is probably one of the of the parent antibody. most unpredictable areas of biotechnology. For example, 5 replacement of a single lysine residue at position 118 of acidic fibroblast growth factor by glutamic acid led to the substantial loss of heparin binding, receptor binding and biological activity Burgess et. al. Journal of Cell biology, 111: of the protein. 10 2129-2138 (1990). In transforming growth factor alpha, replacement of aspartic acid at position 47 with alanine or asparagine did not affect biological activity while replacement with serine or glutamic acid sharply reduced the biological activity of the mitogen. Lazar et. al. Molecular and Cellular Biology, 8:1247-1252 Similarly it has been shown that aglycosylation of 15 (1988). antibodies reduces the resistance of the antibodies to proteolytic degradation, while CH2 deletions increase the binding affinity of the antibodies. See Tao et. al. The Journal of Immunology, Vol. 143. No. 8. 2595-2601 (1989) and Gillies et. al. Human Antibodies 47-54 (1990). and Hybridomas, Vol 1, no. 1, These references 20 amino acid substitution or what demonstrate that even a single appears to be an inconsequential chemical modification, will often dramatically affect the biological activity and characteristic of a Therefore, without sufficient guidance in the protein.

6

Serial No. 715272

Art Unit 1806

10

specification to support the use of the above terms and for the reasons mentioned above one of ordinary skill in the art would forced into undue experimentation in order to practice the invention as is claimed.

5 Claims 1-11 are rejected under 35 U.S.C. § 112, first paragraph, for the reasons set forth in the objection to the specification.

35 U.S.C. 5 101 reads as follows:

Whoever invents or discovers any new and useful process, machine, manufacture, or composition of matter or any new and useful improvement thereof, may obtain a patent therefore, subject to the conditions and requirements of this title.

15 Claims 1-4, 6-8 are rejected under 35 U.S.C. § 101 because the claimed invention is directed to non-statutory subject matter. The above claims are drawn to a method of preparing an antibody, however, there is no indication within the claims that actual physical steps are taking place. For example, there is no step 20 which includes isolating an antibody, rather obtaining an amino acid sequence. All of the steps which are listed in the claims can be done on paper as mental steps or on a computer terminal.

The specification is objected to under 35 U.S.C. § 112, first paragraph, and claims 9-13 are rejected under 35 U.S.C. § 112, 25 first paragraph and 35 U.S.C. § 101 as the specification fails to adequately teach how to use the claimed monoclonal antibodies in the manner in which they are disclosed ie. for the therapeutic

7

Ŋ

Applicants claims are supported only by in vitro data purposes. showing the ability of muMab4D5, which is a humanized anti-p185 antibody which reacts with breast and ovarian cancers, to react with different cell lines (see page 88-90 of the specification). Applicant has made no showing that these data correlate with utility for in vivo therapy in humans of the complex array of diseases encompassed by the claims. ln general, effective treatment of human cancers has not been routinely achieved in the art using monoclonal antibodies. Further, in vitro data such as that reported in the specification and animal model studies 10 frequently do not correlate with clinical utility in in vivo trials Based on the evidence of record, the alleged utility in patients. of the claimed composition for the treatment of cancer would not be believable on its face to the person of skill in the art in view of the contemporary knowledge in the art. Applicant has not provided 15 any showing of therapeutic utility of the subject monoclonal antibodies which would lead one of skill in the art to believe that the antibodies are broadly applicable for the treatment of all types of autoimmune diseases. Applicant is required to provide evidence commensurate with the scope of the claims, which would be 20 art that the claimed convincing to those skilled in the compositions have utility for the treatment of malignant and autoimmune diseases in humans. See MPEP 608.01(p).

8



Waldmann, in a recent review of the literature pertaining to clinical applications of monoclonal antibodies for diagnosis and therapy of human disease, teaches that effective therapy using monoclonal antibodies has been elusive and indicates that hopes for 5 antibody-based treatment methods engendered by in vitro studies have not correlated well with in vivo clinical trial results in patients with cancer. It does not appear that the exemplary material provided in the specification in support of the assertions that the claimed antibodies have therapeutic utility would be 10 viewed by those skilled in the art as being predictive of their utility for treating humans. Applicant has not exemplified how to use the claimed antibodies in vivo and has not shown that the antibodies would be effective in vivo. It appears that undue experimentation would be required of one skilled in the art to practice the claimed invention for the single utility disclosed in 15 the specification.

The following is a quotation of the appropriate paragraphs of 35 U.S.C. 5 102 that form the basis for the rejections under this section made in this Office Action:

20

25

A person shall be entitled to a patent unless--

- (a) the invention was known or used by others in this country, or patented or described in a printed publication in this or a foreign country, before the invention thereof by the applicant for a patent.
- (b) the invention was patented or described in a printed publication in this country or a

Serial No. 715272

Art Unit 1806

5

10

15

20

foreign country or in public use or on sale in this country, more than one year prior to the date of application for patent in the United States.

The following is a quotation of 35 U.S.C. § 103 which forms the basis for all obviousness rejections set forth in this Office action:

A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negatived by the manner in which the invention was made.

Subject matter developed by another person, which qualifies as prior art only under subsection (f) or (g) of section 102 of this title, shall not preclude patentability under this section where the subject matter and the claimed invention were, at the time the invention was made, owned by the same person or subject to an obligation of assignment to the same person.

Claims 1, 2, 5-10 are rejected under 35 U.S.C. § 102(b) as being anticipated by Queen et. al.. The above claims are drawn to 25 a method of producing a humanized antibody wherein the amino acid import antibody and a consensus antibody are an sequences of compared, wherein the CDRs of the import antibody are substituted for the antibody of the consensus antibody, and wherein certain framework residues which are responsible for the binding of 30 antigen, interaction with CDR, or participating in the VI-Vh interaction are also imported to the consensus antibody. In essence, residues of the framework region are also transferred with

10

Serial No. 715272

Art Unit 1806

15

the CDRs in order to retain the antigen binding affinity of the parent antibody.

Queen et. al. describe the production of humanized antibodies wherein the murine antibody is compared to human antibodies and the 5 most homologous human antibody is chosen as the acceptor molecule. The CDRs of the murine antibody are then substituted for the CDRs of the human antibody and certain framework residues are also changed. Queen et. al. describe computer modeling and sequence comparison in order to determine the amino acid residues which are 10 to be substituted (see page 10031-10033). Although the steps of the methods are not in exactly the same order, all of the claimed elements are present with in the reference.

Claims 1,2 and 5-10 are rejected under 35 U.S.C. § 102(a) as being anticipated by Co et. al.. See above discussion.

Co et. al. show the production of humanized anti-HSV

using the general concept of Queen et. al. (see Results and Table 1).

Claims 3 and 4 are rejected under 35 U.S.C. § 103 as being unpatentable over Queen et. al. or Co et. al. in view of Wallick 20 et. al.

The above claims are drawn to a method of making a humanized antibody wherein the CDRs of an import antibody are transferred to a consensus human antibody along with certain residues of the framework. Furthermore, the claims require that the glycosylation

11

sites, if any, of the import amino acid also be imported with the CDRs and framework regions if these sites have an affect on the binding of antigen.

Queen et. al. and Co et. al. both describe the production of humanized antibodies by transferring the CDRs and certain framework 5 regions of the donor antibody to the human consensus antibody (see Queen et. al. pages 10031-10033 and Co et. al. page 2871). They further state that any residue which might have an affect on the antigen binding of the antibody should be changed substituted in 10 order to maintain the binding affinity of the parent antibody (see page 10033 of Queen et. al. at the last paragraph on the page). They do not however, specifically discuss the glycosylation sites as potential targets for transfer. Wallick et. al. teach the importance of carbohydrate interaction with antigen for maintaining 15 or increasing antigen binding affinity (see pages 1107-1108). ït would have been prima facie obvious to one of ordinary skill in the art at the time the invention was made to make humanized antibodies using the method of Queen et. al. or Co et. al. and further incorporating the concept taught by Wallick et. al.. One of ordinary skill in the art would have been motivated to combine the 20 teachings of the two references in view of the teaching of Queen that retaining high antigen binding affinity is desirable in the production of humanized antibodies. Knowing the role of carbohydrates in antigen antibody interaction as was pointed out by

12

Wallick et. al. one of ordinary skill would have had the means and the motivation to make humanized antibodies using both of the teachings of the primary and secondary references.

Claim 11 is rejected under 35 U.S.C. § 103 as being 5 unpatentable over Queen et. al. or Co et. al. in view of Reichmann et. al.

The above claim is drawn to a humanized antibody wherein only one amino acid (listed in claim 9) in the framework and the CDRs have been substituted in the consensus antibody.

Queen et. al. and Co et. al. both teach the production of 10 humanized antibodies by transferring the CDRs of a murine antibody along with specific residues of the framework region to the They do not however teach only acceptor antibody molecule. substituting one of the framework residues among those listed in claim 9. Queen et. al. introduce the general concept of a scaffold 15 wherein certain amino acid residues of the framework must be present and certain are dispensable. Reichmann et. al. teach that a single amino acid substitution in an antibody is sufficient to retain the antigen binding specificity of the parent antibody (see 20 final paragraph). It would have been prima facie obvious to one of ordinary skill in the art at the time the invention was made to make only a single substitution in the antibody of Queen et. al. or Co et. al. in positions among those listed in claim 9. It would have been obvious to one of ordinary skill to complete the

13

invention in light of the success of Reichmann et. al. in only mutating one amino acid of the framework. Knowing that each antibody varies slightly in the non-conserved region, and given the computer modelling protocol set forth by Queen et. al. one of 5 ordinary skill would have been motivated to make a single mutation in the variable region with the expectation of obtaining a functional antibody.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Lila Feisee 10 whose telephone number is (703) 308-2731.

Any inquiry of a general nature or relating to the status of this application should be directed to the Group receptionist whose telephone number is (703) 308-0196.

Feisee/lf () 15 September 29, 1992

SUPERVISORY PATENT EXAMINER GROUP 180 9/30/92

--- ' ATTACHMENT TO PAPER NUMBER SERIAL NO. CROUPART UNIT FORM PTO-892 (REV. 3-78) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE 801 13 \$ 272 NOTICE OF REFERENCES CITED Have Carter et 2 U.S. PATENT DOCUMENTS FILING DATE IF SUB. CLASS CLASS DATE NAME DOCUMENT NO. 8 С D G н FOREIGN PATENT DOCUMENTS PERTINENT SUB-CLASS DATE COUNTRY NAME CLASS • DOCUMENT NO. м -N 0 a OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.) alli Ø an Ê She Journal ull C 21 990 2 29 $\bar{\mathcal{A}}$ he mme 0 $\frac{2}{8}$ Doursa Maloen 198 250 XAMI .. J 12 1/101 A copy of this reference is not being furnished with this office action. (See Manual of Patent Examining Procedure, section 707.05 (a).) 214

M PTO-1449	· ·	U.S. Dept of Commerce Patern and Hadelpark Office	Atty Docket No. 709	Serial No. 07/715,272
ST OF PRIOR	ART CITED BY APPLICANT	A MAY	Applicant Paul J. Carter et	al. # htt
(Use sever	al sheets if necessary)	86 1992 5	Filing Date June 14, 1991	Group 1806
xaminer nitials	OTHER PRIOR ART (Includ	ling Author, Ditte, Date, Pertinent P	ages, Etc.)	
Ac. A	Chothia <u>et al., J. Mol. Biol.</u> 186:6	51-663 (1985)		,
B	Novotny and Haber, Proc. Natl. Acad	. <u>Sci. USA</u> 82:4592-4596 (1985)		
A ·	Morrison, S. L. <u>et al.</u> , <u>Proc. Natl.</u>	Acad. Sci. USA 81:6851-6855 (1984)	~ .	
Ro	Boulianne, G. L. <u>et al.</u> , <u>Nature</u> 312	:643-646 (1984)		
E	Neuberger, M. S. <u>et al.</u> , <u>Nature</u> 314	:268-270 (1985)	********	
F	Brüggemann, M. <u>et al.</u> , <u>J. Exp. Med</u> .	166:1351-1361 (1987)		
G	Riechmann, L. <u>et al.</u> , <u>Nature</u> 332:3 2	3-327 (1988)		
(A) H	Love <u>et al.</u> , <u>Methods in Enzymology</u>	178:515-527 (1989)		
IK I	Bindon et al., J. Exp. Med. 168:127	-142 (1988)		
(b) ·	Jones, P. T. <u>et al.</u> , <u>Nature</u> 321:522	-525 (1986)		
K	Verhoeyen, M. <u>et al.</u> , <u>Science</u> 239:1	534-1536 (1988)	t 1	
L L	Hale, G. <u>et al.</u> , <u>Lancet</u> i:1394-1399	(1988))	
	Queen, C. <u>et al.</u> , <u>Proc. Natl. Acad.</u>	<u>Sci. USA</u> 86:10029-10033 (1989)		
N	Co <u>et al.</u> , <u>Proc. Natl. Acad. Sci. U</u>	<u>sa</u> 88:2869-2873 (1991)		
F º	Gorman <u>et al., Proc. Natl. Acad. Sc</u>	i <u>. USA</u> 88:4181-4185 (1991)	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	#####
NEP	Daugherty <u>et al.</u> , <u>Nucleic Acids Res</u>	earch 19(9):2471-2476 (1991)		
p °	Brown <u>et al.</u> , <u>Proc. Natl. Acad. Sci</u>	. USA 88:2663-2667 (1991)		
R	Junghans <u>et al., Cancer Research</u> 50	:1495-1502 (1990)		
S S	Davies, D. R. <u>et al.</u> , <u>Ann. Rev. Bio</u>	chem, 59:439-473 (1990)		
IA T	Chothia, C. & Lesk, A. M., J. Mol. 1	Biol. 196:901-917 (1987)		
xaminer 🔀		Date Consider	a/32	

USCOMM-DC 80-398

FORM PTO-1449 U.S. Dept. of Com Patent and Trademark LIST OF PRIOR ART CITED BY APPLICANT				U.S. Dept. of Competence Patent and Trademark Office	Atty 709	Docket	No. Se (erial No.)7/715272	
				NULAX	Applicant Paul J. Carter et al.				
(Use s	evera	l sheets if neces	sary)	86 1992	Filing June 1	Date 4, 1991	Gre	oup (ROB	
				U.S. PATENT DOCUMENTS & TRADERNIN					
Examiner Initials		Document Number	Date	Name		Class	Subclass	Filing Date	
JA	AA	4,816,567	3/28/89	Cabilly et al.		630	387.1		
40	AB								
	AC								
	AD		,				-		
	AE								
	AF AG								
	AH								
	AI								
	AJ								
	AK			· .					
<u>`</u>	•	·····		FOREIGN PATENT DOCUMENTS		1		Translatio	
		Document Number	Date	Country		Class	Subclass	Yes No	
н,	AL								
2.	AM,								
. 4.	AN								
	AO			· · · ·					
	AP	, , , ,	HER PRIOR	ART (Including Author, Title, Date, Pertinent	Pages, E	 tc.)			
- RI	AR								
0			Chothia, C. <u>et al.</u> , <u>Nature</u> 342:877-883 (1989)						
W.		Tramontano, A. <u>et al.</u> , <u>J. Mol. Biol.</u> 215:175-182 (1990)							
h	· AT	Margolies <u>et al., Proc. Natl. Acad. Sci. USA</u> 72:2180-2184 (1975)							
	AU	Pluckthun, <u>Biotechnology</u> 9:545-51 (1991)							
W									
P		Spiegelberg <u>et al.</u> , <u>Biochemistry</u> 9:4217-4223 (1970)							
10/	AW	Wallick <u>et al.</u> , <u>J. Exp. Med.</u> 168:1099-1109 (1988)							
IN	AX	Sox <u>et al.</u> , <u>Proc</u>	. Natl. A	cad. Sci. USA 66:975-982 (1970)					
1111/	1	1		Date Consider	nord C	~ 1-	20100	`	

USCOMM-DC 80-398.

FORM PTO-1449				U.S. Dept. of C Patent and Tradema		Atty Docket No. Serial No. 709 07/715272		
IST OF P	RIOR	ART CITED BY APPLI	CANT		ROOM	Applicant Paul J. Car	ter et al.	
(Use	sever	al sheets if neces	sary)		I MAY	Fling Date		806
				U.S. PATENT DOCUMEN	85 42	June 14, 199	1 1	80 P
xaminer	.]	1		U.S. PATENT DOCUMEN	Ver.	/		
nitials		Document Number	Date	Name	Tha	Class	Subclass	Filing Date
	BA							
	BB				· ·			
	BC BD							
	BE							
	BF	-						
	BG							
	BH							
·	B1							
	ВJ							
· · · ·	вк			1				
	.	·····	· · · · ·	FOREIGN PATENT DOCUME	INTS	· · · · · · · · · · · · · · · · · · ·	T	
~		Document Number	Date	Count	ъ	Class	Subclass	Translation Yes No
Ø	BL	WO 91/09967	7/11/91	PCT				
Ŭ	BM							
	BN			i i	·			
	BO	-		· · ·				
	BP			ART (Including Author, Title, Da	ta Dontinant Dago			l
N7 -	BR	T		munol.6:535-554 (1988)	ite, rentinent rage	5, 210.7		<u>,</u>
18								
()	BS	Fendly, B. M. <u>et a</u>	al., <u>Canc</u>	er Res. 50:1550-1558 (1990)				· .
R/	ВТ	Neuberger <u>et al.</u> ,	Nature 3	12:604-608 (1984)				
JY.		Tabada at al Na		50 / F/ (1095)				
Jr_	BU	Takeda <u>et al.</u> , <u>Nature</u> 314 :452-454 (1985)						
R	вv	Snow and Amzel, <u>Protein: Structure, Function, and Genetics</u> 1:267-279, Alan R. Liss, Inc. pubs. (1986)						
ÜP	BW	Cheetham, J., Prot	tein Engir	neering, 2(3): 170-172 (1988)		• • • • • • • • • • • • • • • • • • •	·····	
	вх							
ami <u>ner</u>	L	10-			Date Considered	ali	Alr.)
	X	~~X	L			- Id	U 190	

USCOMM-DC 80-398.

, 20, ^{1,16} PATENT DOCKET 709

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Paul J. Carter et al. Serial No. 07/715,272 Filed: 14 JUNE 1991 For: IMMUNOGLOBULIN VARIANTS

SUPPLEMENTAL INFORMATION DISCLOSURE STATEMENT

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

The attached materials were received in connection with the prosecution of a foreign patent application corresponding to the captioned case. These materials contain at least two reference citations, the relevance of which is apparent from the communication from the foreign patent office that is also enclosed.

A PTO Form 1449 is submitted herewith to facilitate citation to the record of all references contained in these materials.

Respectfully submitted, GENENTECH, INC Carolyn R. Adler Reg. No. 32,324

December 30, 1992 460 Point San Bruno Boulevard South San Francisco, CA 94080 415-225-2614

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231.

astraug Strasbaugh

Date: December 30, 1992

PATENT DOCKET 709

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Paul J. Carter et al.

Serial No. 07/715,272

Filed: June 14, 1991

For: Immunoglobulin Variants



Group Art Unit: 1806

Examiner: L. FEISEE

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

CERTIFICATION UNDER 37 C.F.R. § 1.97(e)

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

I hereby certify that each item of information contained in this information disclosure statement was cited in a communication from a foreign patent office in a counterpart foreign application not more than three months prior to the filing of this statement. Pursuant to §1.97, this information disclosure has been filed in a timely fashion and no fees are required.

Respectfully submitted,

GENENTECH, INC. Carolvn R. Adler

Reg. No. 32,324

Dated: December 30, 1992

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D_C. 20231.

trasbaug

Date: December 30, 1992

PR: 1806



PATENT DOCKET 709

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Paul J. Carter et al.

Serial No. 07/715,272

Filed: 14 June 1991

For: Immunoglobulin Variants

Group Art Unit: 1806

Examiner: L. Feisee

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

Amendment and Response

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

b.

d.

e.

f.

Responsive to the Office Action mailed 5 October 1992, please amend the claims as follows:

(Amended) A method for making at least a portion of a humanized antibody variable domain comprising amino-acid sequence of a non-human[, import] antibody which is desired to be humanized (import antibody) and a human antibody, comprising the steps of:

- a. obtaining the amino acid sequences of [at least a portion of] an import variable domain and of a consensus human variable domain;
 - identifying Complementarity_Determining Region (CDR) amino acid sequences in the import and the human amino variable domain sequences;
- c. substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;
 - aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;
 - identifying import antibody FR residues in the aligned FR sequences that are nonhomologous to the corresponding consensus antibody residues;
 - determining if the non-homologous import amino acid residue is reasonably expected to have at least one of the following effects:

1. non-covalently binds antigen directly,

c124.u

SC13194 02/17/93 07715272

07-0630 130 102

148.00CH

Page No. 2

07/715,272

g,

2.

3.

4.

2. interacts with a CDR; or

- 3. participates in the $V_L V_H$ interface by affecting the proximity or orientation of the V_L and V_H regions with respect to one another; [and]
- for any non-homologous import antibody amino acid residue which is [reasonably] expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence; and

<u>preparing a humanized antibody variable domain having amino acid sequences</u>
 <u>determined in steps a-g</u>.

(Amended) The method of claim 1, having an additional step of determining if any such nonhomologous residues are exposed on the surface of the <u>consensus human antibody variable</u> domain or buried within it, and if the residue is exposed, retaining the consensus residue.

(Amended) The method of claim 1, having the additional steps, which may be taken between any two steps in the method of claim 1, of searching the import antibody variable domain amino acid sequence for glycosylation sites, determining if any such glycosylation site is reasonably expected to affect the antigen binding-or-affinity of the antibody, and if so, substituting the glycosylation site into the consensus amino acid sequence.

(Amended) The method of claim 1, having the additional steps, which may be taken between any two steps in the method of claim 1, of searching the consensus variable domain amino acid sequence for glycosylation sites which are not present at the corresponding amino acid in the import antibody amino acid sequence, and if the glycosylation site is not present in the import sequence, substituting the import amino acid residues for the amino acid residues comprising the consensus glycosylation site.

5. (Amended) The method of claim 1, having an additional step which comprises aligning import antibody and consensus antibody FR <u>amino acid</u> sequences, identifying import antibody FR <u>amino acid</u> residues which are non-homologous with the aligned consensus FR sequence, and for each such non-homologous import antibody FR <u>amino acid</u> residue, determining if the corresponding consensus antibody <u>amino acid</u> residue represents a residue which is highly conserved across all species at that site, and if it is so conserved, preparing a humanized antibody which comprises the consensus antibody amino acid residue at that site.

7.

--17.

--19.

Page No. 3

(Amended) A method <u>for making a humanized antibody</u> comprising providing [at least a portion of] a non-human antibody variable domain amino acid sequence <u>which is desired to be</u> <u>humanized (import antibody)</u> having a CDR and a FR, obtaining the amino acid sequence of at least a portion of a consensus human antibody variable domain having a CDR and a FR, substituting the non-human CDR for the human CDR in the consensus human antibody variable domain, and then substituting an amino acid residue for the consensus amino acid residue at at least one of the following sites:

4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

Please add the following new claims 17-21:

A method of using a consensus human antibody variable domain amino acid sequence in the preparation of a humanized antibody.-- $||2|2^{-2}$ where $||2|2^{-2}$ is more than $||2|2^{-2}$ where $||2|2^{-2}$ is the preparation of a humanized antibody.--

--18. In a method for making a humanized antibody variable domain, the improvement consisting of using consensus human antibody variable domain amino acid sequence.--

A method for making an improved antibody, comprising amino acid sequence from a nonhuman (import) antibody and a human antibody, comprising the steps of:

- a. obtaining the amino acid sequences of at least a portion of an import antibody variable domain and of a consensus human antibody variable domain;
- b. identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human amino variable domain sequences;

c. substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

d. aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;

e. identifying import antibody FR residues in the aligned FR sequences that are nonhomologous to the corresponding consensus antibody residues;

determining if the non-homologous import amino acid residue is reasonably expected to have at least one of the following effects:

1. non-covalently binds antigen directly,

c124.u

f.

g.

i.

2. interacts with a CDR; or

participates in the V_L - V_H interface by affecting the proximity or orientation of 3. the V_L and V_H regions with respect to one another;

for any non-homologous import antibody amino acid residue which is reasonably expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence; and

 preparing an improved, humanized antibody having amino acid sequences determined in steps a-g; and

evaluating the antigen binding or immunogenicity of the improved, humanized antibody with respect to the parental antibody.--

--20. A method comprising, following the identification of an antibody by the method of any one of claims 1, 7, or 17-19, the manufacture of the antibody.--

--21. A method comprising, following the identification of an antibody by the method of any one of claims 1, 7, or 17-19, the expression of nucleic acid encoding the antibody.--

Remarks

Claims 1-13, and 17-21 are presented herein for examination. Reconsideration of the outstanding rejections is respectfully requested for the reasons that follow. A request for a one-month extension of time to respond is submitted herewith, bringing the due date for this response to 5 February 1993. This response is timely filed.

Amendments

Claims 1, 3, 4, 5 and 7 have been amended to indicate that an import antibody is a non-human antibody which is desired to be humanized. Support for this language is found in the specification at page 6, line 27 to page 7, line 3.

Claim 1, step (f) has been amended to clarify that the word "participates" in the $V_L - V_H$ interface means to affect the proximity or orientation of the V_L and V_H regions with respect to one another. Support for this amendment is found on page 15, lines 30-32. New step (h) has been added to claim 1, directed to the physical step of preparation of a humanized antibody variable domain. Support for this step appear throughout the specification.

Claims 3 and 4 have been amended to provide that the additional steps may be taken between

Page No. 5

any two steps in the method of claim 1. Claims 2-7 have been amended to clarify that the residues or sequences referred to relate to amino acids.

New claims 17 - 21 have been added. These claims are alternate approaches to claiming the subject matter claimed in claim 1. Additional support for claims 20-21 is found in Example 1.

It is believed that these amendments introduce no new matter. The inventors respectfully request entry of these amendments.

The rejection under 35 U.S.C. § 112, second paragraph

Claims 1-10 were rejected under 35 U.S.C. § 112, second paragraph for indefiniteness. Claims 1, 3, 4, 5 and 7 were rejected for use of the term "import antibody". These claims have been amended to indicate that the import antibody is a non-human antibody which is desired to be humanized.

Claim 1 step (a) was rejected because of the term "consensus human variable domain". The terms "consensus sequence", "consensus antibody" and "consensus human variable domain" are defined at specification page 16, line 29 to page 17, line 17:

The terms "consensus sequence" and "consensus antibody" as used herein refers to an amino acid sequence which comprises the most frequently occurring amino acid residues at each location in all human immunoglobulins of any particular subclass. In preferred embodiments, the consensus human variable domain sequences are derived from the most abundant subclasses in the sequence compilation of Kabat *et al., Sequences of Proteins of Immunological Interest,* National Institutes of Health, Bethesda MD (1987), namely $V_L \kappa$ subgroup I and V_H group III....

As described in the specification, a "consensus human variable domain" would have an amino acid sequence comprising, amino acid residue by residue, the most frequently occurring amino acid residue gathered from a group of human immunoglobulins. The identity of each amino acid residue making up the consensus sequence is determined separately, requiring merely routine tabulation of the amino acids present in each member of a particular immunoglobulin subclass. To expedite the routine tabulation of the most commonly occurring amino acids, workers in the field are referred to the Kabat *et al.*, publication cited in the quoted material above, which presents such tabulations.

Claim 1 step (d) was rejected as indefinite as to whether the alignment of the amino acid sequences is a physical or mental step. This rejection is somewhat confusing. The inventors intend claim 1, step (d) to refer to a maximal homology alignment of representations of amino acid sequences, as described in the specification at page 17, lines 18-27. Preparing such a homology alignment typically <u>combines</u> physical and mental actions. This connotation for the phrase "alignment of sequences" is common in the art to which this invention pertains. Step (d) of claim 1 does not require

Page No. 6

manipulation of the actual, tangible amino acids, merely manipulation of symbolic representations of the actual amino acids.

Claim 1 step (e) was rejected because of the term "homology"; the Examiner questioned whether conservative amino acids are to be considered as homologs. Identity or homology with respect to a specified amino acid sequence of this invention is defined on page 17, lines 18-27. At lines 22-23, the specification indicates that this invention does "not consider[ing] any conservative substitutions as part of the sequence identity". Conservative substitutions are therefore not considered as homologs.

Claim 1 step (f) was rejected for use of the language "participates". Step (f) of claim 1 refers to an amino acid residue which "participates in the in the $V_L - V_H$ interface". This step has been amended to clarify that immunoglobulin residues which so participate are those that affect the proximity or orientation of the V_L and V_H regions with respect to one another.

Claim 1 step (f) was also rejected as indefinite as to how one of ordinary skill can determine the effects listed in steps 1-3. Steps 1-3 presently list the following effects an import amino acid residue might have:

1. non-covalently binds antigen directly,

- 2. interacts with a CDR; or
- 3. participates in the $V_L V_H$ interface by affecting the proximity or orientation of the V_L and V_H regions with respect to one another.

The specification discusses, at pages 13-16, the interactions of amino acid residues within an immunoglobulin and describes at least two methods for evaluating the role of any particular amino acid residue: three dimensional models and assays. As stated at page 14, lines 2-9:

"Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen."

The specification provides detailed information how to evaluate the three-dimensional models to determine the various potential effects of amino acid residue changes.

The specification also suggests an alternate method for evaluating the effect of an amino acid

residue change. On page 16, lines 14-18, the specification teaches:

"Since it is not entirely possible to predict in advance what the exact impact of a given substitution will be it may be necessary to make the substitution and assay the candidate antibody for the desired characteristic. These steps, however, are *per se*

Page No. 7

routine and well within the ordinary skill of the art."

The inventors submit that methods for determining the effects of amino acid changes are known in the art, and that those skilled in the art would understand what is claimed in step (f). Claim 1 step (n) was rejected as held in the step of the s

Claim 1 step (g) was rejected as being indefinite as to what effects are reasonably expected to occur. The word "reasonably" has been deleted from the claim.

Claim 2 was rejected as lacking antecedent basis for "the domain". This claim has been amended to clarify that the intended domain is the consensus human antibody variable domain.

Claims 3 and 4 were rejected as indefinite for not specifying when in the process one would search for the glycosylation sites. These claims have been amended to indicate that one would search for glycosylation sites between any two steps in the method of claim 1.

Claim 5 was rejected as unclear in the use of the phrase "preparing a humanized antibody"; this phrase has now been added by amendment as the last step of claim 1. The phrase is intended to mean the physical making of a humanized antibody, methods for which are described in the specification, including *in vitro* mutagenesis and recombinant engineering. The Examiner also seems to be questioning how claim 5 differs from the previous claims. Claim 5 adds an additional step of determining if a particular amino acid residue in the consensus human variable domain--which differs from the import antibody amino acid residue at that site--also appears at that site in antibodies of other species at that particular site (is conserved). If the particular amino acid residue is conserved across species at that site, than that residue is retained in the humanized antibody, and not substituted by the import antibody amino acid residue at that site, and without requiring evaluation of the impact of such a change on the antibody's characteristics.

Claim 6 was rejected as vague for unclear use of numbers. These numbers refer to particular amino acids in the light (L) and heavy (H) chains of immunoglobulins. By convention, workers in this field generally utilize the immunoglobulin numbering system set forth in Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987)), as described in the specification at page 8, lines 19-21. The Examiner's attention is drawn to Queen *et al.*, already of record in this case, especially at page 10032 column 1 first paragraph (and reference 38 therein) where antibody amino acid residues are referred to with numbers representing certain positions. It is submitted that workers in the field will understand clearly what is claimed in claim 5.

Claim 7 was rejected as indefinite as to what the method is drawn, and has been amended according to the Examiner's suggestion.

According to the CAFC, a decision as to whether a claim is invalid for indefiniteness "requires a determination whether those skilled in the art would understand what is claimed", <u>Amgen v. Chugai</u>,

Page No. 8

18 USPQ2d 1116, 1030 (CAFC 1991). The presently pending claims use terminology with clear meanings in the field, especially in light of the definitions provided in the specification. The wordings of the claims comply with the requirements of 35 USC § 112, and this rejection should be reconsidered and withdrawn.

The rejection under 35 U.S.C. § 112, first paragraph

Claims 1-11 were rejected under 35 U.S.C. § 112, first paragraph as lacking enablement.

Claims 1 and 7 were rejected as lacking enablement in the language "at least a portion of an import variable domain". These terms have been deleted from the claims.

Claim 1 step (c) was rejected for being unclear as to how one would determine which amino acids are to be substituted. This step recites "substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence". The identification of the Complementarity Determining Region (CDR) amino acid sequence of the import and the human amino variable domain sequences is made in previous step (b). To accomplish step (c), therefore, one substitutes the amino acids identified in step (b).

Methods for identifying CDRs and distinguishing them from Framework Residues (FRs) are known in the art. As the specification describes on page two, antibody variable domains of natural light and heavy chains have the same general structure, and each domain comprises four framework (FR) regions, whose sequences are somewhat conserved, connected by three hyper-variable or complementarity determining regions (CDRs) (see Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda, MD, (1987)). The four framework regions largely adopt a β -sheet conformation and the CDRs form loops connecting, and in some cases forming part of, the β -sheet structure, The CDRs in each chain are held in close proximity by the framework regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site. The CDR may be identified following three-dimensional modeling of the antibody. The CDRs may also be identified based on comparison of the an antibody amino acid sequence with a known antibody.

Attached as Exhibit A for the Examiner's convenience are pages from the Introduction to Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest, Fifth Edition*, National Institutes of Health, Bethesda, MD, (1991). This work, along with the earlier Kabat compendiums referred to in the specification and other references, guide the practitioner in the numbering of antibody amino acid sequences, and the assignment of particular amino acids to one of the FR or CDR regions. The Examiner's attention is drawn the sections beginning on page xv, the section entitled "Variable Region Sequence" and especially to Table I, page xvi. Table I presents the amino acid residues associated

Page No. 9

with FRs and CDRs of the variable domains of immunoglobulin light and heavy chains. See also Figure 1. page xviii, which shows a schematic view of an immunoglobulin; please note the mention in that figure description to the use of a maximum homology alignment to determine the proper numbering of the amino acids (as referred to in the response to the previous § 112 rejection). The inventors submit that the identification of immunoglobulin amino acid residues as belonging to a CDR or to the framework is routine in the art, requiring no undue experimentation.

The specification teaches, in detail, several ways to substitute amino acid residues, including mutagenesis and the construction of nucleic acid encoding the desired sequence. Alanine scanning mutagenesis is described at page 36, line 20 to page 37, line 3. Oligonucleotide-mediated mutagenesis, PCT mutagenesis and cassette mutagenesis are described in the specification at page 39, line 10 through page 44, line 10. The inventors submit that steps (b) and (c) of claim 1 are fully enabled by the specification.

Claim 1 step (f) was rejected as lacking enablement for determining which amino acid residues may be expected to interact with the antigen. At page 29, lines 4-10, the specification teaches that:

"Differences between the non-human import and the human consensus framework residues are individually investigated to determine their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is desirably performed through modeling, by examination of the characteristics of the amino acids at particular locations, or determined experimentally through evaluating the effects of substitution or mutagenesis of particular amino acids."

Techniques for molecular modeling are described on pages 27-28. Experimental evaluation of the role of particular amino acids will utilize assays tailored to the activities of the antibody to be humanized.

More detailed teaching on identifying residues that influence antigen binding is contained in the specification at page 14, line 10 through page 15, line 6, where it is stated:

"A residue that noncovalently directly binds to antigen is one that, by three dimensional, analysis, is reasonably expected to noncovalently directly bind to antigen. Typically, it is necessary to impute the position of antigen from the spatial location of neighboring CDRs and the dimensions and structure of the target antigen. In general, only those humanized antibody residues that are capable of forming salt bridges, hydrogen bonds, or hydrophobic interactions are likely to be involved in non-covalent antigen binding, however residues which are separated spatially by 3.2 Angstroms or less may also non-covalently interact. Such residues typically are the relatively larger amino acids, such as tyrosine, arginine, and lysine. Antigen-binding FR residues also typically will

Page No. 10

have side chains that are oriented into an envelope surrounding the solvent oriented face of a CDR which extends about 7 Angstroms into the solvent from the CDR domain and about 7 Angstroms on either side of the CDR domain, again as visualized by three dimensional modeling.

The inventors submit that determining whether a residue may be expected to influence antigen binding is routine in the art, in light of the detailed teachings of the specification.

Claim 2 was rejected as lacking enablement for determining which residues are exposed on the surface or buried within the domain. As indicated in the specification, for example at page 91, lines 18-21, the worker in this field would examine the structural models of the import and human sequences to determine if an amino acid residue is exposed on the surface of the domain or is buried within. Evaluation of structural models, preparation of which are described in the specification, to determine whether a residue is exposed or buried is routine and within the ordinary skill in the art.

Claim 3 was rejected as lacking enablement for how one would determine which glycosylation site affects antigen binding, or what comprises "reasonable expectation". The specification teaches, at page 8, lines 22-32, teaches that determining if the glycosylation is reasonably expected to be important for the desired antigen binding and biological activity of the antibody involves determining if the glycosylation site binds to antigen or changes a side chain of an amino acid residue that binds to antigen, or if the glycosylation enhances or weakens antigen binding, or is important for maintaining antibody affinity. As with other aspects of this invention, evaluation of the impact of glycosylation typically is performed by evaluation of molecular models, or experimental evaluation of a modified polypeptide. Such evaluation is routine within the field.

Claims 6, 7 and 9 were rejected as being enabled only with respect to IgG and not other antibody isotypes. The specification, at page 13 lines 14-22, states:

"The humanized antibody will be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG3 and IgG4. Usually the constant domain is a complement fixing constant domain where it is desired that the humanized antibody exhibit cytotoxic activity, and the class is typically IgG_1 . Where such cytotoxic activity is not desirable, the constant domain may be of the IgG_2 class. The humanized antibody may comprise sequences from more than one class or isotype, and selecting particular constant domains to optimize desired effector functions is within the ordinary skill in the art."

The Examples presented in the specification involve the use of a IgG₁ constant domain. As noted in the specification, specific method steps and illustrative reagents for the use of IgG₁ are taught, as well

Page No. 11

as their applicability to other antibody isotypes. The inventors respectfully reminds the Examiner that working examples are not a required component of a patent application. As stated in MPEP § 608.01(h), "There is no statutory requirement for the disclosure of a specific example." Thus, the absence of a working example describing particular embodiments of the invention cannot negative the patentability of the invention. The examples included in the specification, which illustrate the preparation of IgG₁ antibodies, are representative of the manner in which the invention may be practiced. From reading these examples and the detailed description of the invention, the ordinarily skilled artisan would immediately deduce the applicability of the methods described in the specification to other immunoglobulin isotypes.

The Examiner has not made a prima facie case for the § 112, first paragraph rejections, supplying no basis for her skepticism about the scope of the claims. The burden is on the Examiner to provide evidence to support rejections of this sort. "Mere broad generalizations and allegations are insufficient for holding of non-enablement," <u>Ex parte Goeddel</u>, 5 U.S.P.Q. 1449, 1450 (TTAB 1987).

If the Examiner is only prepared to allow claims to exemplified embodiments, what incentive exists for an inventor to disclose the invention to the public? Trade secret protection obviously would be superior to the following circumstances foreseen by the CCPA in <u>In re Goffe</u>, 191 USPQ 429, 431 (CCPA 1976):

For all practical purposes, the board would limit appellant to claims involving the specific materials disclosed in the examples, so that a competitor seeking to avoid [literally] infringing the claims would merely have to follow the disclosure in the subsequently-issued patent to find a substitute. However, to provide effective incentives, claims must adequately protect inventors. To demand that the first to disclose shall limit his claims to what he has found will work or to materials which meet the guidelines specified for 'preferred' materials in a process such as the one herein involved would not serve the constitutional purpose of promoting progress in the useful arts. See In re Fuetterer, 50 CCPA 1453, 1462, 319 F.2d 259, 265, 138 USPQ 217, 223 (1963).

For a similar case, see <u>In re Strahilevitz</u>, 212 USPQ 561 (P.O.B.A. 1982), where the Board was reversed for rejecting as non-enabling an application that was devoid of even a single working example.

The first paragraph of 35 U.S.C. § 112 requires nothing more than objective enablement. Whether this is achieved by the use of illustrative examples or by broad terminology is of no importance, <u>In re Marzocchi et al.</u>, 169 USPQ 267 (CCPA 1971). Further, an assertion by the Examiner that the enabling disclosure is not commensurate with the protection being sought must be supported by reasons for doubting the truth or accuracy of any statement in the presumptively accurate supporting disclosure. It is also incumbent upon the Examiner to back up such assertions with acceptable evidence or reasoning to substantiate the doubts so expressed, <u>In re Armbruster</u>, 185

Page No. 12

USPQ 152 (CCPA 1975), In re Strahilevitz, op cit.

Se also <u>In re Smith</u>, *supra*, wherein the CCPA reversed an Office ruling that the description in the specification of two categories of prepolymers was not sufficient to support the broad claim for all polymers having a certain desired property. In this case, the court even acknowledged that the specification did not contain language that was precisely identical to the language of the claims. However, the tenor of the specification was that the applicant had made a generic invention rather than one limited to two categories of polymers.

In the present situation, the Examiner has provided no evidence to support the assertion that the invention is not enabled for the preparation of humanized antibodies. Broad claims should be allowed if there is adequate disclosure and where, as in the present situation, there is no pertinent art to prevent such claims. As stated in <u>In re Sus and Schaefer</u>, 134 USPQ 301, 304 (CCPA 1962) (emphasis added):

The public purpose on which the patent law rests required the granting of claims commensurate in scope with the invention disclosed. This requires as much the granting of broad claims on <u>broad inventions</u> as it does the granting of more specific claims on more specific inventions. It is neither contemplated by the public purpose of the patent laws nor required by the statute that an inventor shall be forced to accept claims narrower than his invention in order to secure allowance of his patent.

The inventors submit that in view of the detailed information provided in the specification as discussed above, the specification adequately teaches how to practice the claimed invention. The rejections under 35 USC § 112, first paragraph, should be reconsidered and withdrawn, as they are not statutorily based, are inconsistent with court and Patent Office decisions on the subject, and are contrary to public policy.

The rejection under 35.U.S.C. § 101

Claims 1-4, and 6-8 were rejected under 35 U.S.C. § 101 as being directed to non-statutory subject matter. It is believed that the amendments to the claims made above render most this ground of rejection.

The rejection under 35 U.S.C. § 112, first paragraph and under 35 U.S.C. § 101

Claims 9-13 were rejected under 35 U.S.C. § 112, first paragraph and under 35 U.S.C. § 101 as lacking utility for the treatment of malignant and autoimmune diseases in humans. The inventors request clarification of this rejection, because none of claims 9-13 are directed to methods of treatment. These claims are directed to humanized antibody variable domains and the polypeptides

c124.u

Cuar

Page No. 13

of recited sequences. These polypeptides are useful as probes, and in diagnostic assays, as described in the specification at pages 65-66, and need not rely on therapeutic utility.

The rejections under 35 U.S.C. § 102(b) and § 102(a)

Claims 1, 2, and 5-10 were rejected under 35 U.S.C. § 102(b) as being anticipated by Queen *et al.*, and under 35 U.S.C. § 102(a) as being anticipated by Co *et al.*. The inventors respectfully traverse these rejections.

To constitute anticipation, all material elements of a claim must be found in one prior art source. In re Marshall, 198 USPQ 344 (CCPA 1978); In re Kalm, 154 USPQ 10 (CCPA 1967). The inventors will show that neither Queen nor Coe contains all the material elements of these claims, particularly the limitation regarding the use of a consensus sequence.

The rejected claims are directed to the humanization an antibody, namely the combination of amino acid sequence from a non-human antibody desired to be humanized, and from a <u>consensus</u> <u>human variable domain</u>. Methods for preparing such a consensus sequence are fully described in the specification and are discussed above. The inventors believe that the use of a such a consensus sequence achieve a superior result, or a "better" humanized antibody.

The cited prior art utilizes a different approach, which approach had apparently been taken by all other workers in the field prior to the present invention. These workers did not prepare a consensus human antibody to combine with their non-human antibody. Instead, they selected only one human antibody for use, based on the similarity of that human antibody to their non-human antibody. Queen *et al.* state this objective explicitly, at page 10031, column 2 of their paper:

"In selecting a human antibody to provide the variable region framework for the humanized anti-Tac antibody, we reasoned that the more homologous the human antibody was to the original anti-Tac antibody, the less likely would combining the anti-

Tac CDRs with the human framework be to introduce distortions into the CDRs." Queen continues to describe selecting a human heavy chain V region which was 57% identical to their non-human antibody, after dismissing all other candidate as between 30-52% identical to their nonhuman. They selected the human light chain V region from the same human antibody for their use.

Co et al. are equally explicit describing their similar reasoning. At page 2871, column 1 they state:

"First, a human antibody variable region with maximal homology to the mouse antibody is selected to provide the framework sequence for humanization of the mouse antibody. Normally the heavy chain and light chain from the same human antibody are

Page No. 14

07/715,272

chosen so as to reduce the possibility of incompatibility in the assembly of the two chains. Based on a sequence homology search against the NBRF protein sequence data base, the antibody Pom was chosen to provide the framework sequences for humanization of Fd79."

The approach of the present invention is quite distinct, in its use of a consensus human variable domain sequence. This consensus sequence might or might not have a high degree of homology with the non-human antibody. Neither Queen *et al.* or Coe *et al.* supply this teaching, and therefore do not anticipate the claimed invention. This rejection should be reconsidered and withdrawn.

The rejection under 35 U.S.C. § 103

Claims 3 and 4 were rejected under 35 U.S.C. § 103 as being obvious over Queen *et al.* or Co *et al.* in view of Wallick *et al.*. Claim 11 was rejected under 35 U.S.C. § 103 as being unpatentable over Queen *et al.* or Co *et al.* in view of Reichmann *et al.*

None of the cited references teaches or suggests the claimed invention, which involves the preparation humanized antibodies using a consensus human antibody variable domain. Such a method is not suggested in any of the prior references, and absent such a teaching there was no motivation to try the methods described in the present specification.

The Obviousness Rejections Do Not Meet the Test of Graham v. Deere

The proper context for determining the issue of obviousness is provided in the seminal decision of <u>Graham v. John Deere</u>, 383 U.S. 1, 148 U.S.P.Q. 459 (1966). In that case, the U.S. Supreme Court set forth the following considerations for deciding this issue:

(1) The scope and the content of the prior art;

(2) The difference between the prior art and the claims at issue;

(3) The level of ordinary skill in the pertinent art; and

(4) Secondary considerations such as commercial success, long-felt and unresolved needs, failure of others, etc.

a. Scope and Content of the Prior Art.

1. Queen *et al.* teach the humanization of an anti-Tac antibody. They do not teach the use of a human consensus variable domain to provide the framework for their non-human CDRs.

Page No. 15

Co et al. teach the humanization of an anti-HSV antibody. They do not teach the use of a human consensus variable domain to provide the framework for their non-human CDRs.
 Wallick et al. teach the importance of glycosylation for maintaining the affinity of a monoclonal antibody for its antigen. They do not teach methods for humanization of antibodies,

nor teach the creation of a human antibody variable domain consensus sequence.

4. Reichmann *et al.* teach the humanization of an anti-CAMPATH-1 antibody. They do not teach the creation of a human antibody variable domain consensus sequence, or suggest that such might be desirable to provide the framework for their non-human antibody CDRs.

b. The Differences Between the Prior Art and the Claims at Issue

The Examiner has chosen various pieces of prior art and concludes that the combination of these references would have rendered the invention obvious.

The prior art shows that it was known as of the filing date to produce antibody fragments comprising sequence from a non-human antibody and from a human antibody. Prior to the present filing date, however, methods were not known which included the use of a consensus human variable domain for mounting the non-human CDRs. There would have been no impetus on the part of the skilled artisan at the filing date to attempt to produce such a consensus sequence or use it in antibody humanization, in view of the teachings of the prior art literature. The cited references do not teach or suggest the claimed invention, alone or in any combination, nor would there have been any reason from these references to practice the claimed methods. The absence of a suggestion of the claimed invention in the art of record precludes the Patent Office from satisfying its initial burden of showing prima facie obviousness.

c. Level of Ordinary Skill in the Art.

The <u>Graham</u> inquiries point to a conclusion of non-obviousness of the present claims regardless of the presumed level of skill in the art. However, absent evidence to the contrary, a person of ordinary skill in the art is presumed to be one who essentially follows conventional wisdom and does not undertake to innovate. As stated by the Federal Circuit in <u>Standard Co. v. American Cyanamid</u> <u>Co.</u>, 227 U.S.P.O. 293, 298 (Fed. Cir. 1985):

A person of ordinary skill in the art is also presumed to be one who thinks along the line of convention wisdom in the art and is not one who undertakes to innovate, whether by patient, and often expensive, systematic research or by extraordinary insights, it makes no difference which.

The inventors submit that one who followed the conventional wisdom would not have

Page No. 16

extrapolated from the teachings of the cited references methods for using a consensus human antibody variable domain for humanizing a non-human antibody. Such an extension of the prior art teachings is based entirely upon hindsight analysis of the inventors' methods. The teachings of this invention should not be considered sufficient to support a conclusion of obviousness in this regard.

The inventors submit that in light of the foregoing amendments and remarks the subject matter defined by the pending claims is useful, enabled, and patentable over the references relied upon by the Examiner, which in no way teach or suggest the invention. The inventors believe the claims are now in condition for allowance and earnestly solicit a Notice to that effect. If the Examiner has any questions, she should feel free to contact the undersigned attorney at the telephone number indicated above.

Respectfully Submitted, GENENTECH, INC.

Carolyn R. Adler Reg. No. 32,324

29 January 1993


CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, on 29 January 1993.

Dated: 29 January 1993

c124.u

\$~



EXHIBITA

SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST

FIFTH EDITION

 $\begin{array}{c} \mbox{Tabulation and Analysis of} \\ \mbox{Amino Acid and Nucleic Acid Sequences of Precursors,} \\ \mbox{V-Regions, C-Regions, J-Chain, T-Cell Receptors for Antigen,} \\ \mbox{T-Cell Surface Antigens, } \beta_2\mbox{-Microglobulins,} \\ \mbox{Major Histocompatibility Antigens, Thy-1, Complement,} \\ \mbox{C-Reactive Protein, Thymopoietin, Integrins, Post-gamma Globulin,} \\ \mbox{α_2-Macroglobulins, and Other Related Proteins} \end{array}$

1991

Elvin A. Kabat*, Tai Te Wu*, Harold M. Perry[†], Kay S. Gottesman[†], and Carl Foeller[†]

*Depts. of Microbiology, Genetics and Development, and Neurology, Cancer Center/Institute of Cancer Research, College of Physicians and Surgeons, Columbia University, New York, NY 10032 and the National Institute of Allergy and Infectious Diseases, and the Office of the Director National Institutes of Health, Bethesda, MD 20892.

*Depts. of Biochemistry, Molecular Biology, and Cell Biology, and Engineering Sciences and Applied Mathematics and Biomedical Engineering, Northwestern University, Evanston, IL 60208 and the Cancer Center, Northwestern University Medical School, Chicago, IL 60611

⁷BBN Systems and Technologies, 10 Moulton Street, Cambridge, MA 02138

*Formerly with BBN. Present address Laboratory for Applied Research in Academic Information, William H. Welch Medical Library, The Johns Hopkins University, Baltimore, Md 21205

The collection and maintenance of this data base is sponsored under grant 5R01 AI-125616 to E.A. Kabat of Columbia University by the following components of the National Institutes of Health, Bethesda, MD 20892:

Office of the Director

- National Center for Research Resources
- National Cancer Institute
- National Institute of Allergy and Infectious Diseases
- National Institute of Diabetes, Digestive and Kidney Diseases
- National Institute of General Medical Sciences
- National Library of Medicine

Work with the PROPHET software package is supported by a subcontract from Columbia University to BBN Systems and Techologies, Cambridge, MA 02138

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

Public Health Service National Institutes of Health

NIH Publication No. 91-3242

INTRODUCTION

Our earlier "Variable Regions of Immunoglobulin Chains" (1), the second edition "Sequences of Immunoglobulin Chains" (2) and the third edition "Sequences of Proteins of Immunological Interest" (3) have been further exixpanded in the Fourth Edition (4) and now in the Fifth Edition to include amino acid and nucleotide sequences of precursors, variable regions, constant regions, J-chains of immunoglobulins, β^2 -microglobulins, antigens of the major histocompatibility complex (HLA, H-2, Ia, DR) as well as of Thy-1, complement, T-lymphocyte receptors for antigens, other T-cell antigens of the immunoglobulin superfamily, interleukins, integrins and various other proteins related to immune functions. The identification and sequencing of clones obtained using recombinant DNA techniques has yielded nucleotide sequences of signal, variable, and constant regions of immunoglobulins (5,6), and these nucleotide sequences have been translated into amino acid sequences with those determined earlier directly by amino acid sequencing and are indicated by an apostrophe followed by CL after the name of the clone. We have continued to use the PROPHET Software Package of the National Center for Research Resources, National Institutes of Health (7,8) to tabulate the sequences.

In compiling the data for this Fifth Edition we have tried to be as up-to-date as possible and have included only sequences which have been published or which have been accepted for publication. Residues which have not been definitely determined have been excluded. It should be remembered that sequences are often published in review articles without detailed documentary evidence. These have often been revised. We have listed such revisions in the notes in many instances; others can readily be found by comparison with sequences in previous editions. We have compiled sequences determined directly as amino acids and have merged with them those translated from the nucleotide sequences thus making all comparable data available. When antibody activities were known, they have been listed after the amino acid and nucleotide sequence tables and are included in the indexes.

When doubts arise as to the validity of any residue in a sequence, the original reference should be examined to ascertain whether editions, we have sent the amino acid and nucleotide sequences as stored in the computer to the original authors for verification. If so verified, this was denoted by "checked by author" at the end of each reference and except for the earliest sequences, the date on which the checked sequences from GenBank (9) have been used. Programs for converting a GenBank sequence to the codon format of our tables have been developed. The correctness of the table sequence has been verified by converting back into the linear form and comparing with GenBank. When this has been done the sequence is listed as "from GenBank". Recently we have developed newer programs that automatically process a GenBank entry completely e.g.: extract the relevant feature, determine the appropriate table, and perform alignment. In such cases, the reference will of the GenBank accession numbers from Which the data was obtained. Some nucleotide sequences were transmitted to us by electronic mail, and they are indicated by "received from authors through email." If the sequences were entered by us from the literature and then checked with GenBank. In general, we have not included stretches of sequence such as enhancers, switch regions and introns. Much information about such sequences may be found in references (10-13). We have also had access to the Protein Information Resource (14) and to the European Molecular Biology Laboratories Data Base (15).

It is also possible, by examining the numbers of sequences at the

xiv

end of each table and the summary tables, to evaluate the probability that a given amino acid at a given position may not be correct. This is most readily done for the framework residues of the V-region and for the C-region; in the complementarity-determining regions this is more difficult because of the high variability. variability.

AMINO ACID SEQUENCES

The first column in each table gives the residue number. Except for complement, T-cell surface antigens, integrins and miscellaneous proteins, the second column is a tabulation of invariant residues. Since exceptions to invariance are found, the frequency, if less than 1.0 and greater than or equal to 0.95, is indicated alongside the residue listed as invariant; when only a single sequence is available, this is not given. These rows are shaded in grey. Each sequence is tabulated in each subsequent column. Three dashes (--) indicate that no amino acid is present at that position and that the sequence continues. In all instances residues considered uncertain by the authors have not been included in the table. In some instances the symbol # is used to indicate that several amino acid residues were found in one position, and these residues are listed in the notes. The four columns at the end of each table give: The first column in each table gives the residue number. Except for give:

2.	the number of residues sequenced at that position, the number of different amino acids found at that position, the number of times the most common amino acid occurred and that amino acid in parentheses, and
4.	the variability.

These columns are included only in tables with more than five sequences. Miscellaneous tables have only columns corresponding to the first two above.

Variability is calculated (16) as:

Number of different amino acids occurring at a given position

Variability =

Frequency of the most common amino acid at that position

An invariant position would have a variability of one; if 20 amino acids occurred with equal frequency, the variability would be 20 divided by 0.05 equals 400. If, for example, four different amino acids Ser, Asp, Pro, and Thr occurred at a given position, and of 100 sequences available at that position, Ser occurred 80 times, the variability would be 4/0.8 = 5. When any of the amino acid residues, sequenced directly as amino acids, were not identified completely and are listed as Glx (or Asx), two values, separated by a comma, are given in the last three columns. The first value in each of these columns is calculated assuming that only one of the two possibilities, e.g., Glu or Gln (or Asp or Asn) occurred, while the second considers that both were present and maximizes variability. In the variability plots, the horizontal bars indicate the two values. the two values.

When two or more amino acids are most common and occur with equal When two or more amino acids are most common and occur with equal frequency, they are tabulated as a note, and the symbol + is used in the next to last column. If no sequence data have been reported for any position, there are no entries in the last four columns. Variability is not calculated for insertions or if only a single sequence is known. When the translated sequence of a clone corresponds to a previously listed sequence of a plasmacytoma from which it was prepa variability computa If a given sequence is indicated by an antibody specificit constants if avail rabbit heavy chain domain of the rabb sequence is given; usually the most ne included, especial) Notes are of two t the symbol #, and s

Signal Sequences The signal (precu chains are listed light chains, for total of nine precu sequencing of sign: sequences from DNA acid residues in Genomic DNA clones the coding sequenc -4, and in rare ca: leader peptide to for positions -4 t

The signal amino antigens, $\beta 2$ -mici proteins, complem amino proteins are liste

By conformational Leu-Leu-Leu-Trp-Va alpha helical conformations in t four amino termin (20).

Variable Region So The variable regi contain hypervari. (27-30) chains, labeled with hapt segments of ligh examination of se chains aligned These and the thre were hypothesized regions or segme contact with vari high resolution x been verified by antibodies all hypervariable req antibody combinin the framework (] framework segme complementarity-(the three CDRs 5 Figures 3-47 have comments are giv bibliography. The Table I.



which it was prepared, only one sequence is listed so that the variability computations are not affected, and a note is included. variability computations are not affected, and a note is included. If a given sequence is associated with any antibody activity, this is indicated by an asterisk alongside the protein heading, and the antibody specificities are given in a separate list with binding constants if available. The notes list the a-allotypes for the rabbit heavy chain V-region and the b-allotypes for the constant domain of the rabbit kappa light chain. A key reference to the sequence is given; generally the most recent references since it is usually the most nearly complete, but often several references are included, especially when revisions of a sequence have been made. included, especially when revisions of a sequence have been made. Notes are of two types: general notes about a table indicated by the symbol #, and specific notes indicated by the sequence number.

xν

972

Signal Sequences The signal (precursor) amino acid sequences of immunoglobulin chains are listed as human, mouse, and miscellaneous for kappa light chains, for lambda light chains, and for heavy chains for a total of nine precursor tables. They were obtained either by direct sequencing of signal proteins (17-19) or by translating nucleotide sequences from DNA clones. Signal segments range from 17-29 amino acid residues in length and are thus numbered from -29 to -1. Genomic DNA clones contain introns of varving length that interrupt Genomic DNA clones contain introns of varying length that interrupt the coding sequence of the precursor within the codon for position -4, and in rare cases for position -6. Thus, the L-gene encodes the leader peptide to position -4 and the 5' end of the V-gene codes for positions -4 to -1.

The signal amino acid sequences of the T-cell receptors for antigens, $\beta_{2-microglobulins}$, major histocompatibility complex proteins, complement components, integrins, and other related proteins are listed in separate tables.

By conformational energy calculations, the core V, hydrophobic Leu-Leu-Leu-Trp-Val-Leu-Leu (MOPC321, MOPC63) exists in an alpha helical conformation, terminated by chain reversal conformations in the four C-terminal residues Trp-Val-Pro-Gly; the four amino terminal residues are compatible with the alpha helix (20):

Variable Region Sequences The variable regions (21) of immunoglobulins have been shown to contain hypervariable segments in their light (16,22-26) and heavy (27-30) chains, of which certain residues have been affinity labeled with haptenic determinants (31-44). Three hypervariable segments of light chain were delineated from a statistical examination of sequences of human V_x , human V_y , and mouse V_z light chains aligned for maximum sequence similarity (16,23,24,27). These and the three corresponding segments of the heavy chains (27) were hypothesized (16,27) to be the complementarity-determining regions or segments (CDR) containing the residues which make contact with various antigenic determinants, several years before high resolution x-ray structures were determined, and this has now been verified by X-ray diffraction studies at high resolution for all antibodies examined Figures 3-47. The proposed fourth hypervariable region (cf. 30) of heavy chains is not part of the antibody combining site (27). The rest of the V-region constitutes the framework (16,27,45-54). It is convenient to identify the framework segments (FR1, FR2, FR3, and FR4) and the complementarity-determining segments (CDR1, CDR2, and CDR3) with the three CDRs separating the four FRs. The CDRs in the stereo Figures 3-47 have solid circles for each residue. References and comments are given with each figure and are not listed in the bibliography. The residue numbers for these segments are given in Table I. Table I.

xvi

Amino Acid Residues Associated with Framework(FR) and Complementarity Determining Regions (CDR) of the Variable Domains of Immunoglobulin Light (V_L) and Heavy (V_H) Chains

Segment	Light Chain	Heavy Chain		
FRI	1-23 (with an occasional	1-30 (with an occasional		
	residue at 0, and a	residue at 0)		
	deletion at 10 in V_{λ} chains)			
CDR1	24-34 (with possible	31-35 (with possible		
	insertions numbered	insertions numbered		
	as 27A, B, C, D, E, F)	as 35A,B)		
FR2°	35-49°	36-49		
CDR2	50~56	50-65 (with possible		
CDRZ		insertions numbered		
		as 52A, B, C) ^b		
	57-88	66-94 (with possible		
FR3		insertions numbered		
		as 82A, B, C)		
0000	89-97 (with possible	95-102 (with possible		
CDR3	insertions numbered as	insertions numbered as		
	95A, B, C, D, E, F)	100A, B, C, D, E, F, G, H, I, J, K)		
201	98-107 (with a possible			
FR4	insertion numbered as 106A)	103-113		

⁴ Five Basilea rabbits (λ) immunized with type II pneumococci and which produced anti-type II pneumococcal polysaccharide had Met at position 48 and an insertion of four amino acid residues between positions 48 and 49; in four of the five the sequence was Glu, Leu, Lys, Ser and the fifth was Trp, Leu, Arg, Lys (53,54,63,64); the others were not sequenced at these positions (for references see table of rabbit λ amino acid sequences.)

 $^{\rm b}$ In the rabbit, Mage et al. (65) consider position 65 in V₈ to be in FR3, since it is allotype related.

241

The V-genes for the and the J-minigenes f kappa light chains. I by recombination and by the J-minigene. I occur at different pr residues may result a of the inserted resi for better alignment the V-gene region. In times more frequentl:

The V-genes for the ł and are followed b extensive variation ability to be read boundary between D a acid position. In add sequences vary by a i of D-J joining appea between V and D and and correlates with t B cells (60). The or has therefore been re evidence suggesting perhaps a minigene nucleotides. Light (V₁-J₂ junction (62), probably results from in fetal and neonata and 17/146 RNA sequi lower than in adults regulated both in T diversity but are te

1

In the tables of V horizontal lines for chain, MPC 11, has between position 1 have internal deleti xvii.

The V-genes for the light chains code to amino acid position 95, and the J-minigenes from position 97 to 107 for lambda and 108 for kappa light chains. Position 96 is usually the site of V-J joining by recombination and may be coded partly by the V-gene and partly by the J-minigene. Because the site of V-J recombination could occur at different positions within a codon, different amino acid residues may result at this position. We have changed the location of the inserted residues from 97A-F (2) to 95A-F, since it makes for better alignment by confining chains of different lengths to the V-gene region. In mouse V_x chains, J1 and J2 were used 5 to 10 times more frequently than J4 and J5 (55).

The V-genes for the heavy chains code up to amino acid position 94 and are followed by the D- and J-minigenes. Because of the extensive variation in the lengths of D-minigenes, and their ability to be read in different reading frames (56), the exact boundary between D and J is not always located at the same amino acid position. In addition, the lengths of the J encoded amino acid sequences vary by a few amino acid residues. Moreover, the process of D-J joining appears to involve insertions of extra nucleotides between V and D and between D and J, termed the N region (57-61) and correlates with the appearance of terminal deoxytransferase in B cells (60). The original numbering system for the heavy chains has therefore been retained. Wysocki et al. (61) have provided some evidence suggesting a non-random origin for the V_R-D_R junction, perhaps a minigene, rather than random addition of the N nucleotides. Light chains do not appear to have N sequences at the V-J_j junction (62), but show an additional residue 95A which probably results from V_J_j joining. N sequences are generally rare in fetal and neonatal mouse V_R-D-J_R junctions (62), only 1/87 DNA and 17/146 RNA sequences contained N regions, an incidence much lower than in adults indicating that N insertion is developmentally regulated both in T and B cells. P elements also contribute to diversity but are templated (62a).

In the tables of V-regions, the FR and CDR are separated by horizontal lines for convenience in reading. One mouse kappa light chain, MPC 11, has an extra segment of 12 amino acid residues between position 1 and the signal sequence (66). Several chains have internal deletions.

xviii

S.

Figure 1 (50) shows the domain structure for IgGl protein EU. Numbering on the left half indicates the CDR for the light and heavy chains (50), while that on the right half gives the EU numbering (67).




FIG. 1. Schematic view of four-chain structure of human IgG1, molecule. Numbers on right side: actual residue numbers in protein EU [Edelman et al. (67)]; Numbers of Fab fragment on left side aligned for maximum homology: light chains numbered as in Wu and Kabat (16) and heavy chains as in Kabat and Wu (27). Heavy chains of EU have residue 52A, three residues 82A, B, C, and lack residues termed 100A, B, C, D, E, F, G, H, I, J, K, and 35A, B. Thus residue 110 (end of variable region) is 114 in actual sequence. Hypervariable regions, complementarity-determining segments or regions (CDR): heavier lines. V, and V_x: light and heavy chain variable region of light chain. Hinge region in which two heavy chains are linked by disulfide bonds is indicated approximately. Attachment of carbohydrate is at residue 297. Arrows at residues 107 and 110 denote transition from variable to constant regions. Sites of action of papain and pessin and locations of a number of genetic factors are given. Modified from 50. Critical understanding sites and the genetic antibody complementar evaluation of a large and especially of the c and heavy chains of im to locate residues i determinants (68,69) a combining sites will d and scope V_A and V_L chai must be resolved. ? immunochemical data in in addition to other m

and the second second

Through the generous C_i been provided with the Fab molecules, $V_{\rm H}$ dime. Drs. Eduardo Padlan $\hat{\epsilon}$ shown. Legends and k model.

Critical understanding of the architecture of antibody combining sites and the genetics of the generation of diversity and of antibody complementarity depends to a great extent on the evaluation of a large number of sequences of the variable regions and especially of the complementarity-determining segments of light and heavy chains of immunoglobulins of different species. Ability to locate residues in the site making contact with antigenic determinants (68,69) and to predict (70) the structures of antibody combining sites will depend heavily upon such sequences. The role and scope V_n and V_n chains in contributing to binding of the epitope must be resolved. This can be often accomplished by use of immunochemical data in defining antibody combining sites (68,70-73) in addition to other methodologies such as 2D-NMR (71,51), or high resolution X-ray crystallography. resolution X-ray crystallography.

xix

Through the generous cooperation of X-ray crystallographers we have been provided with the α -carbon coordinates of almost all available Fab molecules, V_h dimers and antigen-antibody complexes from which Drs. Eduardo Padlan and Chantal Abergal made the stereo models shown. Legends and key references for each are listed with the model. model.

Best Available Copy

: <u>3</u> 21 Long, R. Mage, C'. WiB and L. Steiner for Hier editions.

大学の大学

х

bn.

貼り

asia-carbon coordinates an and Chantal Abergel Ristereodiagrams from Drindebted to them. Dr. Steed to Drs. Rose Mage tions to organizing the $K_{\rm A}$ families as $V_{\rm A}$ and $V_{\rm K}$ families as $V_{\rm M}$ and $V_{\rm K}$ sequences into lar.

lxxiii

REFERENCES TO INTRODUCTION

Kabat, E.A., Wu, T.T., and Bilofsky, H. (1976) Variable Regions of Immunoglobulin Chains. Medical Computer Systems, Bolt Beranek and Newman Inc., Cambridge, MA.
 Kabat, E.A., Wu, T.T., and Bilofsky, H. (1979) Sequences of Immunoglobulin Chains. National Institutes of Health, NIH Publication 80-2008.

1.1 3. Kabat, E.A., Wu, T.T., Bilofsky, H., Reid-Miller, M., and Perry, H. (1983) Sequences of Proteins of Immunological Interest. National Institutes of Health.

Institutes of Health.
Institutes of Health.
Institutes of Health.
Institutes of Health.
Institutes of Health.
Institutes of Proteins of Immunological Interest.
Institutes of Health and Human Services, Public Health Service National Institutes of Health.
5. Tonegawa, S., Maxam, A.M., Tizard; R., Bernard, O., and Gilbert,
W. (1978) Sequence of a mouse germ-line gene for a variable region of an immunoglobulin light chain. Proc. Natl. Acad. Sci. U.S.A. 5:1485-1489.

6. Seidman, J.G., Leder, A., Edgell, M.H., Plosky, F., Tilghman, S.M., Tiemeier, D.C., and Leder, P. (1978) Multiple related immunoglobulin variable region genes identified by cloning and sequence analysis. Proc. Natl. Acad. Sci. U.S.A. 75:3881-3885.

7. Raub, W.F. (1974) The PROPHET system and resource sharing. Federation Proceedings 33:2390-2392.

8. Hollister, C. (1988) A national computational resource for life science research. Nucleic Acids Research 16:1873.

9. Burks, C., Fickett, J.W., Goad, W.B., Kanehisa, M., Lewitter, F.I., Rindone, W.P., Swindell, C.D., Tung, C.-S., and Bilofsky, H (1985) The GenBank nucleic acid sequence database. CABIO Lewitter, CABTOS 1:225-233.

10. Trifonov, E.N. and Brendel, V. (1986) GNOMIC, A Dictionary of Genetic Codes. Balaban Publishers; Rehovot, Philadelphia.

11. Nussinov, R. (1986) Some guidelines for identification of recognition sequences: regulatory sequences frequently contain (T)GTG/CAC(A), TGA/TCA and (T)CTC/GAG(A). Biochimica et Biophysica Acta 866:93-108.

12. Ghosh, D. (1990) A relational data base of transcription factors. Nucleic Acids Res. 18:1749-1756.

13. Wu, T.T., Reidmiller, M., Perry, H.M., and Kabat, E.A. (1984) Long identical repeats in the mouse $\gamma 2b$ switch region and their implications for the mechanism of class switching. EMBO J. 3:2033-2040.

14. Protein Information Resource. National Biomedical Research Foundation, Georgetown University Medical Center, Washington, DC 20007.

15. European Molecular Biology Organization (1990) EMBL File Server. See issues of Nucleic Acid Research for recent sequences added to their data bank.

1. 16. Wu, T.T. and Kabat, E.A. (1970) An analysis of the sequences of the variable regions of Bence Jones proteins and myeloma light the variable regions for antibody complementarity. J.

lxxiv

17. Milstein, C., Brownlee, G.G., Harrison, T.M., and Mathews, M.B. (1972) A possible precursor of immunoglobulin light chains. Nature New Biol. 239:117-120.

18. Schechter, I. and Burstein, Y. (1976) Partial evidence of the precursors of immunoglobulin light chains of different subgroups: Evidence that the immunoglobulin variable-region gene is larger than hitherto known. Biochem. Biophys. Res. Comm. 68:489-496.

19. Rose, S.M., Kuehl, W.M., and Smith, G.P. (1977) Cloned MPC11 myeloma cells express two kappa genes: a gene for a complete light chain and a gene for a constant region polypeptide. Cell 12:453-462.

20. Pincus, M.R. and Klausner, R.D. (1982) Prediction of the three dimensional structure of the leader sequence of the pre-light chain, a hexadecapeptide. Proc. Natl. Acad. Sci. U.S.A. 79:3413-3417.

21. Hilschmann, N. and Craig, L.C. (1965) Amino acid sequence studies with Bence Jones proteins. Proc. Natl. Acad. Sci. U.S.A. 53:1403-1409.

22. Milstein, C. (1967) Linked groups of residues in immunoglobulin chains. Nature 216:330-332.

23. Kabat, E.A. (1967) Unique features of the variable regions of Bence Jones proteins and their possible relation to antibody complementarity. Proc. Natl. Acad. Sci. U.S.A. 59:613-619.

24. Kabat, E.A. (1970) Heterogeneity and structure of antibody combining sites. Landsteiner Centennial, Dec. 5,6,7, 1968. Ann. N.Y. Acad. Sci. 169:43-54.

25. Franêk, F. (1969) The character of variable sequences in immunoglobulins and its evolutionary origin. In Developmental Aspects of Antibody Formation and Structure. Academia, Czechoslovak Academy of Sciences, Prague, pp. 311-313.

E.A. (1969) Discussion in Developmental Aspects of Antibody Formation and Structure. Academia, Czechoslovak Academy of Sciences, Prague, pp. 391-393.

27. Kabat, E.A. and Wu, T.T. (1971) Attempts to locate complementarity determining residues in the variable positions of light and heavy chains. Ann. N.Y. Acad. Sci. 190:382-393.

28. Milstein, C. and Pink, J.L.R. (1970) Structure and evolution of immunoglobulins. Prog. Biophys. Mol. Biol. 21:209-263.

29. Capra, J.D., Kehoe, J.M., Winchester, R.J., and Kunkel, H.G. (1971) Structure-function relationships among anti-gamma globulin antibodies. Ann. N.Y. Acad. Sci. 190:371-381.

30. Capra, J.D. and Kehoe, J.M. (1975) Hypervariable regions, idiotypy and the antibody combining site. Advances in Immunology. Academic Press, New York 20:1-40.

31. Thorpe, N.O. and Singer, S.J. (1969) The affinity-labeled residues in antibody active sites. II. Nearest-neighbor analyses. Biochemistry 8:4523-4534. (1969) The affinity-labeled

E.J. and Metzger, H. (1970) Affinity labeling of a mouse myeloma protein which binds nitrophenyl ligands. Kinetics of labeling and isolation of a labeled peptide. Biochemistry 9:1267-1278.

33. Franêk, F. (1971) Affinity labeling by m-nitrobenzenediazonium fluoroborate of porcine anti-dinitrophenyl antibodies. Position of labeled tyrosine in the λ -chains. Eur. J. Biochem. 19:176-183.

34. Fleet, G.W.J., Kn antibody binding site. photoprecursor of an a Kn

35. Haimovich. J. Eis Bocalization of affini chains of two myelo Biochemistry 11:2389-2.

36 Ray, A. and Cebra, C fesidues in the primar faised in strain 13 gu:

M. L. Kor 37: Yoshioka, M., Li Armstrong, M.Y.K., Kor Studies on the combin immunoglobulin which bi of two types of photo. 12:4679-4685. Buge .

2:4679-4685. Bulge J 38.4Fisher, C.E. and Pr Binding site of rabbit the hypervariable rec 139:135-149. 39:55 Koo, P.H. and Cek distinctive lysyl residi 72 chain of guinea pic Blochemistry 13:184-195 Section 40:2 Chesebro, B., Hadl

Brochemistry 13:184-195 Gerniss 40:3 Chesebro, B., Hadl Pabeling studies on 1 phosphorylcholine. In. andgCells. 3rd Int. Cc Karger, Basel, pp. 205-41. Givol, D. (1974) Aff Combining site. Essays Agrwight. 42.3Roholt, O.A., Frieder (1973) A light chain ty Fabbit antibenzoate anti Slavia. 43T Cebra, J.J., Koo, 1

A3T Cebra, J.J., Koo, 1

AGT Cebra, J.J., Koo, i antibodies: primary stru 1786:263-265. 11B, MA4.eRichards, F.F., Li: Skonigsberg, W.H. (1974) Frégion of myeloma prot Pabeling patterns. Bioc

ASSEPadlan, E.A. (1977) antigen-antibody reacti diversification of anti Brophysics 10:35-65.

6999 Davies; D.R. and I antibody function. Ann.

Kabat, E.A. (1983) I munology V, 67-85. Fi Muramamura and T. Tad

GNOVOTNY, J., Bruccole deKarplus, M. (1983) M GenzulaBiol. Chem. 23: eretion The section

Nat:

34. Fleet, G.W.J., Knowles, J.R., and Porter, R.R. (1972) The antibody binding site. Labeling of a specific antibody against the photoprecursor of an aryl nitrene. Biochem. J. 128:499-508.

lxxv

35. Haimovich, J., Eisen, H.N., Hurwitz, E., and Givol, D. (1972) Localization of affinity-labeled residues on the heavy and light chains of two myeloma proteins with anti-hapten activity. Biochemistry 11:2389-2398.

36. Ray, A. and Cebra, J.J. (1972) Localization of affinity-labeled residues in the primary structure of antidinitrophenyl antibody raised in strain 13 guinea pigs. Biochemistry 11:3647-3656.

Converse, C.A., F.F. (1973) 37. Yoshioka, M., Lifter, J., Hew, C.-L., Converse, C.A., Armstrong, M.Y.K., Konigsberg, W.H., and Richards, F.F. (1973) Studies on the combining region of protein 460, a mouse YA immunoglobulin which binds several haptens. Binding and reactivity of two types of photoaffinity labeling reagents. Biochemistry 12:4679-4685.

38. Fisher, C.E. and Press, E.M. (1974) Affinity labeling of the binding site of rabbit antibody. Evidence for the involvement of the hypervariable regions of the heavy chain. Biochem J. the hypervariable regions 139:135-149.

ĥi n

đν

dý

in

al àk

n F ôf

íťe of 11

Milin

tóns, lógy. Ě. nel`éd

zses. . . lof» a

Cs c Vstry of

£1 e

nium òn of

P:

÷ V 39. Koo, P.H. and Cebra, J.J. (1974) Affinity labeling of a distinctive lysyl residue within the second hypervariable region of γ^2 chain of guinea pig anti-p-azobenzenearsonate antibody. γ2 chain of guinea pig Biochemistry 13:184-195.

40. Chesebro, B., Hadler, N., and Metzger, H. (1973) Affinity labeling studies on five mouse myeloma proteins which bind phosphorylcholine. In Specific Receptors of Antibodies, Antigens and Cells. 3rd Int. Convoc. Immunol., 1972, Buffalo, N.Y., S. Karger, Basel, pp. 205-217.

. ...

41. Givol, D. (1974) Affinity labeling and topology of the antibody combining site. Essays in Biochemistry 10:1-31.

42. Roholt, O.A., Friedenson, B., Radzimski, G., and Pressman, D. (1973) A light chain tyrosyl sequence in the antibody site of a rabbit antibenzoate antibody. J. Immunol. 111:1367-1375.

43. Cebra, J.J., Koo, P.H., and Ray, A. (1974) Specificity of antibodies: primary structural basis of antibody binding. Science 186:263-265.

44. Richards, F.F., Lifter, J., Hew, C.-L., Yoshioka, M., and Konigsberg, W.H. (1974) Photo- affinity labeling of the combining region of myeloma protein 460. II. An interpretation of the labeling patterns. Biochemistry 13:3572-3574.

45. Padlan, E.A. (1977) Structural basis for the specificity of antigen-antibody reactions and structural mechanisms for the diversification of antigen-binding specificities. Quart. Rev.

46? Davies, D.R. and Metzger, H. (1983) Structural basis of antibody function. Ann. Rev. Immunol. 1:87-117.

47. Kabat, E.A. (1983) The antibody combining site. Progress in Immunology V, 67-85. Fifth International Congress of Immunology. Eds. Yamamura and T. Tada. Academic Press, Tokyo.

48. Novotny, J., Bruccoleri, R., Newell, J., Murphy, D., Haber, E., and Karplus, M. (1983) Molecular anatomy of the antibody binding site. J. Biol. Chem. 23: 14433-14437. ЪЦ.,

a formation of the second

247

station after a

lxxvi

.

49. Kabat, E.A., Wu, T.T., and Bilofsky, H. (1977) Unusual distributions of amino acids in complementarity-determining (hypervariable) segments of heavy and light chains of immunoglobulins and their possible roles in specificity of antibody-combining sites. J. Biol. Chem. 252:6609-6616.

The basis of antibody (1978) structural 50. Kabat, E:A. (1978) The structural complementarity. Adv. Protein Chem. 32:1-75.

51. Anglister, J. and Zilber, B. (1990) Antibodies against a peptide of cholera toxin differing in cross reactivity with the toxin differ in their specific interactions with the peptide as observed by 1H NMR spectroscopy. Biochemistry 29:921-928.

52. Kabat, E.A., Wu, T.T., and Bilofsky, H. (1978) Variable region genes for the immunoglobulin framework are assembled from small segments of DNA -- A hypothesis. Proc. Natl. Acad. Sci. U.S.A. 75:2429-2433.

53. Hayzer, D.J. and Jaton, J.-C. (1987) Nucleotide Sequence of a cDNA clone encoding a rabbit immunoglobulin- λ light chain: The V, region differs markedly from that of other species. J. Immunol. 138:2316-2322.

54. Hayzer, D.J. and Jaton, J.-C. (1989) Cloning and sequencing of two functional rabbit germ-line immunoglobulin V_{λ} genes. Gene 80:185-191.

55. Wood, D. and Coleclough, C. (1984) Different joining region J elements of the murine K immunoglobulin light chain locus are used at markedly different frequencies. Proc. Natl. Acad. Sci. U.S.A. 81:4756-4762.

56. Kaartinen, M. and Mäkela, O. (1985) Reading of D genes in variable frames as a source of antibody diversity. Immunology Today 6:324-327.

57. Alt, F.W. and Baltimore, D. (1982) Joining of immunoglobulin heavy chain gene segments: Implications from a chromosome with evidence of three $D-J_{\rm H}$ fusions. Proc. Natl. Acad. Sci. U.S.A. 79:4118-4122.

58. Perlmutter, R.M., Crews, J.T., Douglas, R., Sorensen, G., Johnson, N., Nivera, N., Gearhart, P.J., and Hood, L. (1984) The generation of diversity in phoshorylcholine-binding antibodies. Adv. in Immunol. Academic Press, New York 35:1-37.

59. Manser, T., Huang, S.Y., and Gefter, M. (1984) Influence of clonal selection on the expression of variable region genes. Science 226:1283-1288.

60. Desiderio, S.V., Yancopoulos, G.D., Paskind, M., Thomas, E., Boss, M.A., Landau, N., Alt, F.W. and Baltimore, D. (1984) Insertion of N regions into heavy-chain genes is correlated with expression of terminal deoxytransferase in B cells. Nature 311:752-755.

61. Wysocki, L., Manser, T., Gridley, T. and Gefter, M.L. (1986) Molecular limitations on variable-gene junctional diversity. J. Immunol. 137:3699-3701.

62. Feeney, A.J. (1990) Lack of N regions in fetal and neonatal mouse immunoglobulin V-D-J junctional sequences. J. Exp. Med. mouse 172:1377-1390.

62a. Lafaille, J.J., DeCloux, A., Bonneville, M., Takagaki, Y., and Tonegawa, S. (1989) Junctional sequences of T cell receptor $\gamma\delta$ T cell lineage and for a novel intermediate of VDJ joining. Cell 59:859-870.

63:CGarcia, I. and Jaton J.I. and type VII pneumoco Hight chains. Immunoc studies of the light cha 1851. Ko

1.1

64. Duvoisin, R.M., Ko-Jaton, J.C. (1984) Nuc constant region of Eur. J. Immunol. 14:379-

Mage, R.G., Bernstei CliB:, Young-Cooper, G.O. structural and genetic | allotypes of the rabbit.

14

66arSmith, G.P. (1978) S

 κ 67. Edelman, G.M. Cunni Rutishauser, U., and Wax-cancentire γ G immunoglobu

ancentire γG immunoglobu 63:78-85. 683. Kabat, E.A., Wu, T. Constraints in complet Completing sites which m Proc. Natl. Acad. Sci. U official 693 Padlan, E.A., Davies

693 Padlan, E.A., Davies Othm (1976) Model buildin Chard(1976) Model bulldin hapten-binding site of M Biol. 41:627-637. 7054:Kabat, E.A. (1982) Complementarity. Pharma

711: Dwek, R.A., Wain-Hot Disserving, S.J., and C Combining site by magnet $\mathcal{O}_{i,i}$

22:4)Feldman, R.J., Pott Hypothetical space-filli: Binding myeloma J539. M 23:3-Givol, D. (1979) The Recognition II (E. Lenno 11B, 23:71-125.

7418, 23:71-125. 7449,Padlan, E.A. and Ki Combining Sites. in Mole Applications. Mether Appli

yelw. 25 gKabat, E.A. and Wu, 1 and segments of sequences Relative contributions o Dindinghof antibody comb:

20u (Gibbs, M.R., Moody, I Structure of the asp chloramphenicol acetyltra Consequences of disruptic 29:11261-11265.

ເຄັບກຸດ ! SKriegler; M. (1990) BesydNew: York, pp.1-24(

500 L 71 B Horwitz, A.H., Chang, ecretion of functional a roc. Natl. Acad. Sci. US lxxvii

63. Garcia, I. and Jaton, J.C. (1979) The Immune response to type II and type VII pneumococcal vaccines in Basilea rabbits lacking K light chains. Immunochemical and partial amino acid sequence studies of the light chains. Mol. Immunol. 16:1063-1071. studies of the light chains.

Inina i. of

httpody

the ide, as

region Small ILSI A .

Manof a The V_λ

inol.

Mat- Say 18 m ing of

Gion J Reused DS.A.

meslin 1003odA

. Milin 1 with

Ú₽́S.A. 56 G.,

dies.

énce.of

nes. gene Di iffε.

(1984) dowith Nature alor:

值986) 腹 J. 165

what Med.

and

NGYδ T VCell

• • • • • • • • • •

100 tatal.

of

64. Duvoisin, R.M., Kocher, H.P., Garcia, I., Rougeon, F. and Jaton, J.C. (1984) Nucleotide sequence of a cDNA encoding the constant region of a rabbit immunoglobulin light chain of λ type. Eur. J. Immunol. 14:379-382. F. and

65. Mage, R.G., Bernstein, K.E., McCartney-Francis, N., Alexander, C.B., Young-Cooper, G.O., Padlan, E.A., and Cohen, G.H. (1984) The structural and genetic basis for expression of normal and latent allotypes of the rabbit. Mol. Immunol. 21:1067-1081.

66. Smith, G.P. (1978) Sequence of the full-length immunoglobulin kappa-chain of mouse myeloma MPC 11. Biochem. J. 171:337-347.

67. Edelman, G.M. Cunningham, B.A. Gall, W.E., Gottlieb, P.D., Rutishauser, U., and Waxdal, M.J. (1969) The covalent structure of an entire YG immunoglobulin molecule. Proc. Natl. Acad. Sci. USA 63:78-85.

68. Kabat, E.A., Wu, T.T., and Bilofsky, H. (1976) Attempts to locate residues in complementarity determining regions of antibody combining sites which make contact with antigenic determinants. Proc. Natl. Acad. Sci. U.S.A. 73:617-619.

69. Padlan, E.A., Davies, D.R., Pecht, I., Givol. D., and Wright, C. (1976) Model building studies of antigen-binding sites: the hapten-binding site of MOPC 315. Cold Spring Harbor Symp. Quant. Biol. 41:627-637.

70.: Kabat, E.A. (1982) Antibody diversity versus antibody complementarity. Pharmacol. Reviews 34:23-38.

171. Dwek, R.A., Wain-Hobson, S., Dower, S., Gettins, P., Sutton, B., Perkins, S.J., and Givol, D. (1977) Structure of an antibody combining site by magnetic resonance. Nature 266:31-37.

#2. Feldman, R.J., Potter, M., and Glaudemans, C.P.J. (1981) A hypothetical space-filling model of the V-regions of the galactan binding myeloma J539. Mol. Immunol. 18:683-698.

73. Givol, D. (1979) The antibody combining site. In Defense and Recognition II (E. Lennox, Ed.) Univ. Park Press, Baltimore, MD, 11B, 23:71-125.

74. Padlan, E.A. and Kabat, E.A. (1991) Modelling of antibody combining sites. <u>in</u> Molecular Design and Modelling. <u>in</u> Concepts and Applications. Methods in Enzymology Academic Press. J.N. Abelson and M.I. Simon, Editors in chief. .74.

p: [75. Kabat, E.A. and Wu, T.T. (1991) Identical V-region sequences and segments of sequences in antibodies of different specificities-Relative contributions of $V_{\rm H}$ and $V_{\rm L}$ genes, minigenes and CDRs to binding of antibody combining sites. J. Immunol. (in press).

176. Gibbs, M.R., Moody, P.C.E., and Leslie, A.G.W. (1990) Crystal structure of the aspartic acid-199 asparagine mutant of chloramphenicol acetyltransferase to 2.35-A resolution: structural consequences of disruption of a buried salt bridge. Biochemistry 29:11261-11265.

77. Kriegler, M. (1990) Gene Transfer and Expression. Stockton Press, New York, pp.1-240.

0.3 78. Horwitz, A.H., Chang, C.P., Better, M. and Hellstrom, K. (1988) Secretion of functional antibody and Fab fragment from yeast cells. Proc. Natl. Acad. Sci. USA 8678-8682.





IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Paul J. Carter et al. Serial No. 07/715,272 Filed: 14 June 1991 For:Immunoglobulin Variants Group Art Unit: 1806

Examiner: L. Feisee

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

PETITION AND FEE FOR EXTENSION OF TIME (37 CFR 1.136(a))

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Applicant petitions the Commissioner of Patents and Trademarks to extend the time for response to the Office action dated 05 October 1992 for one month(s) from 5 January 1993 to 5 February 1993. The extended time for response does not exceed the statutory period.

Please charge Deposit Account Number 07-0630 in the amount of \$110. to cover the cost of the extension. Any deficiency or overpayment should be charged or credited to this deposit account. A duplicate of this sheet is enclosed.

Respectfully submitted,

GENENTECH, INC.

Carolyn R.^gAdler Reg. No. 32,324

Date: 29 January 1993

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington (D. C. / 20231.

Date: 29 January 1993

Carolyn R. Adler



GENENTECH, INC. 460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-2614

Attorney Docket No. 709 Examiner:L. Feisee Group Art Unit 1806

SCEIVED

FEB 1 9 1993

GROUP 180

In re Application of: Paul J. Carter et al.

Serial No.: 07/715,272

Filed: 14 June 1991

Immunoglobulin Variants For:

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Transmitted herewith is an amendment in the above-identified application.

The fee has been calculated as shown below.

		(Col. 1)		(Col. 2)	(Col. 3)	-
	- -	Claims Remaining After Amendment		Highest No. Previously Paid For	Present Extra	Rat
•	Total	* 21	Minus	** 21	= 0	x
	Indep.	* 10	Ninus	*** 8	= 2	x
		_ First f	Presentat	ion of Multiple	Dep. Claim	+ 2

Raté	Addit. Fee
x 20≍	\$ 0
x 72=	\$ 144
+ 220=	\$0
TOTAL	, \$ 144.

*If the entry in Col. 1 is less than the entry in Col. 2, write "O" in Col. 3. **If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, write "20" in this space. ***If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, write "3" in this space. The "Highest Number Previously Paid For" (Total or Independent is the highest number found from the equivalent box in col. 1 of a prior amendment or the number of claims originally filed.)

No additional fee is required. 1.

Please charge any additional fees, including any fees necessary for extensions of time, or credit overpayment to Deposit Account No. 07-0630. A duplicate copy of this X sheet is enclosed.

Any additional filing fees required under 37 CFR 1.16. Any patent application processing fees under 37 CFR 1.17. х X

Dated: 29 January 1993

(Attorney of Record)

02/17/93 07715272 SC13193

Carolyn R. Adler Registration No. 32,324 07-0630 130 115 CERTIFICATE OF MAILING 110.00CH

I hereby certify that this correspondence is being deposited with the U.S. Postal Service on the date below as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Dated: 29 January 1993

Carolyn R. Adler

c124.u

2.

	, ×.	سەر. بر		UNITED BT Patent and Address : CDA	ATEL DEPARTMENT OF COMMERCE Trademark Office IMISBIONER OF PATENTS AND TRADEMARKS MINISDIO, D.C. 20231
SEF	IAL NUMBER	FILING DATE	FIRST N	AMED INVENTOR	ATTORNEY DOCKET NO.
ΰ	7/715,272	06/14/91	CARTER		P 709 EXAMINER PETISEE -L
•			18	MZ	ABT UNIT PAPER NUMBER
A	40 POINT 3	ING. LYN R. ADLER SAN BRUND BL FRANCISCO, C	VD.		1806 17
Teis la a d		e examinor in charge of you	IT REOFFERSON.		05/19/93
COMMISS	IONER OF PATENTS	AND TRADEMARKS	1		
7. N A shorter	od statutory period	d for response to this as	Responsive to comm ction is set to expire	<u> </u>	213193 (This action is mode final.
Fallure to	respond within the	period for response w	il cause the applicatio	n to become aband	oned. 35 U.S.C. 133
Part I		D ATTACHMENT(B) A			
	Notice of Art Cite	noes Cited by Examiner ad by Applicant, PTO-1 low to Effect Drawing C	449.		re Patent Drawing, PTO-648. of Informal Patent Application, Form PTO-152.
Part II 1. JK	Ciaima	Dive, claims	14-16		ere pending in the application.
۹ 🗆	Claims		20 1/2		have been cancelled.
* (x	Claima	/	a mars		are allowed.
• 🕅	Ciaime	1-13	, 11-3	21	Bre rejected.
s. []	Claims				are objected to.
e. C	Glaima	······			are subject to restriction or election requirement.
7.	This application	has been filed with info	rmal drawinga under S	17 C.F.A. 1.85 which	are acceptable for examination purposes.
a. []	Formal drawings	are required in respon	se to this Office action	ı.	
9.	The corrected or are accepti	r substitute drawings he able. 🗋 not scosptabl	eve been received on a (see explanation or f	iotice re Petent Dre	, Under 37 C.F.R. 1.84 these drawings wing, PTO-948).
· 10.	The proposed ac examiner. 🔲 d	iditional or substitute s isapproved by the exer	heet(s) of drawings, fil ninar (see explanation	od on	has (have) been 🔲 approved by the
13. 🗖	The proposed di	rawing correction, filed	on		pproved. D disapproved (see explanation).
12. E	Acknowledgmen	it is made of the claim f	or priority under U.S.C	. 119, The certified	copy has D been received D not been received
		parent application, ser		; tiled	•
N. [ation appears to be in (the prectice under Ex)			nattera, prosecution as to the merits is closed in
14. 🗆	Other				
					4

PTOL-326 (Rev. 9-89)

EXAMINER'S ACTION

Serial No. 715272 Art Unit 1806

The text of those sections of Title 35, U.S. Code not included in this action can be found in a prior Office action.

-2-

Some of the rejections under 35 USC 112 second paragraph have been obviated in view of the amendments to the claims. However, the following rejection still remain. The language "consensus human variable domain" is still unclear despite the description in the specification. It is unclear whether the consensus human variable domain is a culmination of different variable domains or a single universal variable domain which is homologous to other human variable domains.

With regards to the langauge "import amino acid", it is suggested the import amino acid be described in the following manner: "an import antibody comprising the amino acid sequence of a non-human antibody which binds to ...". The language "reasonably expected" is unclear since it is not known what criteria determines what is "reasonable".

Claim 1 remains rejected and new claims 19-21 are rejected under 35 USC 112 first paragraph as lacking enablement for the language "at least a portion" for the same reasons as set forth in pages 3 and 4 of paper #13.

Applicant states that this language has been deleted from claim 1, but, this is not the case. This language has been moved to the beginning of the claim and the claim contains the same objectionalble language, therefore, the rejection set forth Serial No. 715272 Art Unit 1806

previously still applies.

The rejection of claims 1-4, 6-8 under 35 USC 101 is withdrawn in view of the amendment to the claims.

The rejection of claims 9-13 as lacking utility is withdrawn in view of the argument set forth in the letter of 2/3/93.

- 7 -

The objection to the specification and the rejection of claims 1-11 under 35 USC 112 first paragraph is maintained and newly added claims 17-21 are rejected for the reasons of record.

The language "at least a portion" still remains in claim 1 and newly added claims 19-21. Therefore, the rejection set forth previously on pages 3-4 of paper #13 still applies. With regards to substituting an import CDR in place of the human CDR, the rejection still applies, since there is no clear guidance in the specification to enable one of ordinary skill in the art to make the human "consensus variable region" which is to contain the claimed substitution. It is true that once the amino acid sequences are known, it is routine to determine the CDRs according to Kabat, and substitute the rodent CDRs in place of the human CDRs. However, the only guidance presented in the specification with regards to the substitutions is the amino acid sequences of SEQ ID NO: 3 and 4, which are specific variable regions. The specification vaguely alludes to variable domain sequences which are derived from the most abundant subclasses but shows no way of making such variable domains. The fact remains

Serial No. 715272 Art Unit 1806

that applicant has not clearly taught how to determine which amino acids are the ones to be substituted since there is only a single example of the appropriate variable region which is to support the substitutions.

-4-

The rejection of claim 2 with regards to determining which residues are surface or buried residues is withdrawn in view of the argument presented explaining that computer modeling is well known in the art to determine the position of various amino acid residues.

The rejection of claims 1 and 3 with regards to the language "reasonably" and newlymadded claim 19 is maintained, since there is no set standard for determining what is reasonable interaction, or interfacing or what amount of glycosylation reasonably affects binding.

The rejection of claims 6,7 and 9 based on the specific amino acids sequences which are only relevant to IgG is maintained. Applicant argues that he is not required to exemplify every embodiment, however, if the claim requires the presence of a certain sequence which does not exist in a particular isotype, than clearly there is a lack of enablement for making that particular embodiment of the claim.

The rejections of claims 1,2,5-10 under 35 USC 102(a) and 102(b) is maintained and newly added claims 17-21 are rejected under 35 USC 102(a) and 35 USC 102(b) as being anticipated by

Serial No. . Art Unit

Queen et. al. or Co et. al. for the same reasons as set forth in the previous Office action.

-5-

Applicant argues that the distinction between the prior art and the instant invention is that the framework amino acids are chosen from a consensus human variable region. However, as previously mentioned there is no clear indication of what is meant by consensus variable regions and as it is stated by applicant on page 14 of the response the chosen amino acids in the references may indeed be the same as what applicant calls consensus variable domain sequences.

The rejection of claims 3 and 4 under 35 USC 103 is maintained for the same reasons as set forth in the previous Office action. Applicant again argues that the use of "consensus region variable domains" is different from the prior art methods, however, as previously mentioned, the consensus amino acids may be the same as the most homologous murine antibodies of the references. The lack of clarity of the language "consensus" amino acid region" is what allows this particular interpretation of the claims.

Claims 17,18, 20 and 21 are rejected under 35 U.S.C. § 112, second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject, matter which applicant regards as the invention. New claims 17,18,20 and 21 are indefinite in that there are no discrete method steps.

Serial No.

Art Unit

THIS ACTION IS MADE FINAL. Applicant is reminded of the extension of time policy as set forth in 37 C.F.R. § 1.136(a).

A SHORTENED STATUTORY PERIOD FOR RESPONSE TO THIS FINAL ACTION IS SET TO EXPIRE THREE MONTHS FROM THE DATE OF THIS ACTION. IN THE EVENT A FIRST RESPONSE IS FILED WITHIN TWO MONTHS OF THE MAILING DATE OF THIS FINAL ACTION AND THE ADVISORY ACTION IS NOT MAILED UNTIL AFTER THE END OF THE THREE-MONTH SHORTENED STATUTORY PERIOD, THEN THE SHORTENED STATUTORY PERIOD WILL EXPIRE ON THE DATE THE ADVISORY ACTION IS MAILED, AND ANY EXTENSION FEE PURSUANT TO 37 C.F.R. § 1.136(a) WILL BE CALCULATED FROM THE MAILING DATE OF THE ADVISORY ACTION. IN NO EVENT WILL THE STATUTORY PERIOD FOR RESPONSE EXPIRE LATER THAN SIX MONTHS FROM THE DATE OF THIS FINAL ACTION.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Lila Feisee whose telephone number is (703) 308-2731.

Any inquiry of a general nature or relating to the status of this application should be directed to the Group receptionist whose telephone number is (703) 308-0196.

Feisee/em May 18, 1993

SUPERVISORY PATENT EXAMINER GROUP 180

6-

ORM PTO-1	449		,	U.S. Dept. of Commerce Patent and Trademark Office	Atty Do 709	cket	No.	Serial No. 07/715,2	
IST OF DISC	LOSUI	RES CITED BY APPLICA	NT		Applican Paul J.		er et al.	Ŧ	14
(Use s	(Use several sheets if necessary) (JAN) 66 JUNE 14, 1991			G	roup	18			
				U.S. PATENT DOCUMENTS		-			
Examiner		· · · · · · · · · · · · · · · · · · ·		TRAVE		lass	Subclass	Filing	Date
Initials		Document Number	Date	Name		,1455		(i (i / i / j	
	AA								
	AB								
	AC								
	AD								
	AE			· · · ·				\ \	
-	AF	· ·							
	AG								
	AH			· · ·					
	AI								
	· AJ			. •					
	AK								
				FOREIGN PATENT DOCUMENTS					
		Document Number	Date	Country		Class	Subclass	Trans s Yes	lation No
A	AL	WO 90/07861	7/26/90	РСТ		• • • • • • • • • • • • • • • • • • • •			
U.	AM								
	AN								
	AO								
	AP								
	I	011	IER DISCLO	SURES (Including Author, Title, Date, Pertinent	: Pages, Et	c.)		<u></u>	
Q	AR			SURES (Including Author, Title, Date, Pertinent Acad. Sci.,89: 4285-4289 (1992)	Pages, Et	c.)			
UP.	I				: Pages, Et	c.)	<u> </u>		
WP.	AR				: Pages, Et	c.)			
UP.	AR				: Pages, Et	c.)		· · · ·	
ÛR.	AR AS AT				: Pages, Et	c.)		· · · · · · · · · · · · · · · · · · ·	
ÛR.	AR AS AT AU				: Pages, Et	c.)			
UP .	AR AS AT AU AV	Carter <u>et al., P</u>			: Pages, Et	c.)			

*Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.



GENENTECH, INC. 460 Point San Bruno Boulevard South San Francisco, CA 94080 **6) 225-2614**

> Attorney Docket No.709 Examiner:L. FEISEE Group Art Unit 1806

SEP 2 5 1993

المعربة التوجه والمروي

rres. and Mail

re Application of: Paul J. Carter

Serial No.: 07/715272

Filed: June 14, 1991

Immunoglobulin Variants For:

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

2.

Transmitted herewith is an amendment in the above-identified application.

The fee has been calculated as shown below.

	(Col. 1)		(Col. 2)	(Col. 3)	
	Claims Remaining After Amendment		Highest No. Previously Paid For	Present Extra	Rate
Total	* 17	Minus	** 21	- 0	x 22=
Indep.	* 6	Minus	*** 10	= 0	x 74=
	First P	resentat	ion of Multiple	Dep. Claim	+ 230=
Conversion of the local division of the loca					70741

Rate .	Addit. Fee
x 22≖	\$ 0
x 74≖	.\$0
+ 230=	\$
TOTAL	\$ 0

*If the entry in Col. 1 is less than the entry in Col. 2, write "O" in Col. 3.

If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, write "20" in this space. *If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, write "3" in this space. The "Highest Number Previously Paid For" (Total or Independent is the highest number found from the equivalent box in col. 1 of a prior amendment or the number of claims originally filed.)

x_ No additional fee is required. 1.

Please charge any additional fees, including any fees necessary for extensions of time, <u>_X</u>___ or credit overpayment to Deposit Account No. 07-0630. A duplicate copy of this sheet is enclosed.

> Any additional filing fees required under 37 CFR 1.16. х Any patent application processing fees under 37 CFR 1.17. <u>_X</u>

Dated: September 20, 1993

Janut E. Husak Attorney of Record

Janet E. Hasak Registration No. 28,616

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the U.S. Postal Service on the date below as inereby certify that this correspondence is being deposited with the 0.5. Poster service on the date below as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Mashington, D.C. 20231. Dated: <u>20 Sept 1993</u>

Louise Strasbaugh

806

PATENT DOCKET 709

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Corres. and Mail

)



Group Art Unit: 1806

Examiner: L. FEISEE

- Filed: June 14, 1991

For: Immunoglobulin Variants

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-1896

PETITION AND FEE FOR EXTENSION OF TIME (37 CFR 1.136(a))

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Applicant petitions the Commissioner of Patents and Trademarks to extend the time for response to the Office action dated 19 May 1993 for one month(s) from 19 August 1993 to 19 September 1993. The extended time for response does not exceed the statutory period.

Please charge Deposit Account Number 07-0630 in the amount of \$110 to cover the cost of the extension. Any deficiency or overpayment should be charged or credited to this deposit account. A duplicate of this sheet is enclosed.

Respectfully submitted,

GENENTECH, INC.

net E. Hozak

Janet E. Hasak Reg. No. 28,616

Date: September 20, 1993 CS14005 09/24/93 07715272 07-0630 140 115

110.00CH

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231.

straslauc muse (Strasbaugh

Date: September 20, 1993

THE UNITED STATES PATENT AND TRADEMARK OFFICE

Examining Group

Filed: June 14, 1991

For: Immunoglobulin Variants

Group Art Unit: 1806

Amendmentinder 37 CFR 1.116 Expedited Procedure

PATENT DOCKET 709

Examiner: L. FEISEE

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-1896

AMENDMENT AFTER FINAL REJECTION PURSUANT TO 37 CFR § 1.116

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Do not enter

This is responsive to the Office Action mailed May 19, 1993, which is a final rejection of claims 1 to 11 and 17 to 21. Claims 12 & 13 have been allowed. A request for a one-month extension of time to respond is submitted herewith, bringing the due date for this response to September 20, 1993 as September 19 is a Sunday. This response is timely filed.

IN THE SPECIFICATION:

On page 19, line 3, please delete "effect or" and insert --effector--. On page 87, please amend Table 1 as follows:

line 6, please amend the heading of the second to last column to read

---K⁺---;

lines 6 & 7, please amend the heading of the last column to read --Relative

cell proliferation[‡]--;

line 8, please delete "proliferation[‡]";

line 10, please amend the second to last column to read --25-- and the last column to read --102--.

IN THE CLAIMS:

1:

Please cancel claims 14-16 and 18 from the application, without prejudice. Please amend claims 1, 3, 7, 17, 19, 20 and 21 as follows:

(Twice amended) A method for making [at least a portion of] a humanized antibody variable domain comprising amino acid sequences of <u>an import antibody</u> <u>comprising</u> a non-human antibody which is desired to be humanized [(import antibody)] and a human antibody, comprising the steps of:

obtaining the amino acid sequences of an import variable domain and of a consensus human variable domain <u>of a human immunoglobulin</u> subgroup;) New

b. identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human amino variable domain sequences;
c. substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

d. aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;

e. identifying import antibody FR residues in the aligned FR sequences that are non-homologous to the corresponding consensus antibody residues;

determining if the non-homologous import amino acid residue is [reasonably] expected to have at least one of the following effects:

1. non-covalently binds antigen directly,

2. interacts with a CDR; or

3. participates in the $V_L - V_H$ interface by affecting the proximity or orientation of the V_L and V_H regions with respect to one another;

w014.u

f:

- 07/715272
 - g. for any non-homologous import antibody amino acid residue which is expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence; and
 - h. preparing a humanized antibody variable domain having amino acid sequences determined in steps a-g.

In claim 3, line 4, please delete "reasonably".

7. (Twice amended) A method for making a humanized antibody comprising providing an import antibody comprising a non-human antibody variable domain amino acid sequence which is desired to be humanized [(import antibody)] having a CDR and a FR, obtaining the amino acid sequence of [at least a portion of] a consensus human antibody variable domain <u>of a human immunoglobulin</u> <u>subgroup</u>, having a CDR and a FR, substituting the non-human CDR for the human CDR in the consensus human antibody variable domain, and then substituting an amino acid residue for the consensus amino acid residue <u>at</u> at least one of the following sites:

4L, 35L, 36L, 38L, 43L, 44L, 46L, 58L, 62L, 63L, 64L, 65L, 66L, 67L, 68L, 69L, 70L, 71L, 73L, 85L, 87L, 98L, 2H, 4H, 24H, 36H, 37H, 39H, 43H, 45H, 49H, 58H, 60H, 68H, 69H, 70H, 73H, 74H, 75H, 76H, 78H, 91H, 92H, 93H, and 103H.

17. (Amended) A method of <u>making a humanized antibody variable domain</u> <u>comprising the step of substituting Complementary Determining Region (CDR)</u> <u>amino acid residues of a variable domain of a non-human antibody for the</u> <u>corresponding CDR amino acid residues of</u> [using] a consensus human antibody variable domain amino acid sequence <u>of a human immunoglobulin subgroup</u> [in the preparation of a humanized antibody].

c.

e.

f.

Page No. 4

- 19. (Amended) A method for making an improved antibody, comprising amino acid sequences from <u>an import antibody comprising</u> a non-human [(import)] antibody and a human antibody, comprising the steps of:
 - a. obtaining the amino acid sequences of [at least a portion of] an import antibody variable domain and of a consensus human antibody variable domain <u>of a human immunoglobulin subgroup</u>;
 - identifying Complementarity Determining Region (CDR) amino acid sequences in the import and the human [amino] variable domain sequences;

substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence;

- d. aligning the amino acid sequences of a Framework Region (FR) of the import antibody and the corresponding FR of the consensus antibody;
 - identifying import antibody FR residues in the aligned FR sequences that are non-homologous to the corresponding consensus antibody residues; determining if the non-homologous import amino acid residue is
 - [reasonably] expected to have at least one of the following effects:
 - 1. non-covalently binds antigen directly,
 - 2. interacts with a CDR; or
 - 3. participates in the $V_L V_H$ interface by affecting the proximity or orientation of the V_L and V_H regions with respect to one another;
- g. for any non-homologous import antibody amino acid residue which is [reasonably] expected to have at least one of these effects, substituting that residue for the corresponding amino acid residue in the consensus antibody FR sequence[; and] ; and
- h. preparing an improved, humanized antibody having amino acid sequences determined in steps a-g; and
- i.
- evaluating the antigen binding or immunogenicity of the improved,

humanized antibody with respect to the parental antibody.

20. A method <u>of making a humanized antibody</u> comprising <u>the step of making the</u> <u>antibody identified</u> [, following the identification of an antibody] by the method of any one of claims [1,] 7[,] or 17 [-19, the manufacture of the antibody].

21. A method <u>of making a humanized antibody</u> comprising <u>the step of expressing</u> <u>nucleic acid encoding the antibody identified</u> [, following the identification of an antibody] by the method of any one of claims 1, 7, [or] 17, [-] <u>or</u> 19 [, the expression of nucleic acid encoding the antibody].

<u>REMARKS</u>

The claims pending in this application are claims 1 to 13, 17 and 19 to 21. Applicants have canceled claims 14 to 16 and 18, without prejudice to file divisional applications directed thereto.

The proposed amendments to the claims are purely in response to the rejections of the Final Action. No new matter has been introduced by the claim amendments. These amendments should be considered under Rule 116 because they do not introduce issues not already fully joined in this case and because they are believed to place the claims in better condition for appeal. Further, they are offered in a good faith effort to place this case in condition for allowance.

I. Amendments

The specification has been amended to correct obvious typographical errors. With respect to the amendment to Table 1 on page 87, a copy of Carter *et al.*, *Proc. Natl. Acad. Sci.*, **89**, (1992) is attached, which is a publication of the experimental data disclosed in the above application, and was published after the filing date thereof. It is clear that the

Page No. 6

last two column headings of Table 1 were inadvertently superimposed and the amendment to the specification serves merely to correct these errors. It would have been obvious from the information provided on page 87 of the specification, that the last two headings were intended to be "Kd nM", and "Relative cell proliferation", respectively, as the key under Table 1 discloses what the headings indicated by ⁺ and [‡] are. Also, it is clear that the figures in the last two columns of the first line of data in Table 1 were intended to be 25 and 102 respectively, and were inadvertently superimposed. Applicants respectfully request that the specification be amended to correct the obvious typographical errors discussed above.

Claims 1, 7, 17 and 19 have been amended to refer to the consensus human variable domain "of a human immunoglobulin subgroup", with support for the amendment found on at least page 16, lines 29-32 and page 17, line 4. Claim 17, 19, and 20 have been amended to recite a preamble and a positive step, which steps are clear from at least the original set of claims filed.

II. Rejections under 35 U.S.C. § 112, second paragraph

Most of the rejections under 35 U.S.C. § 112, second paragraph, which were raised in the earlier Office Action dated October 5, 1992 have been withdrawn. Applicants thank the Examiner for withdrawing these rejections.

The Examiner has, however, maintained some of the rejections under 35 U.S.C. § 112, second paragraph, which relate to claims 1, 3-5 and 7. The separate sets of rejections are addressed separately below.

A. The Examiner has maintained the rejection of claim 1 with respect to the phrase "consensus human variable domain" because it is allegedly not clear whether the consensus domain is a culmination of different variable domains or a single universal variable domain which is homologous to other human variable domains.

In the interests of expediting examination, claims 1, 7, 17 and 19 have been amended to recite that the consensus human variable domain is "of a human

Page No. 7

07/715272

immunoglobulin subgroup". Information concerning the amino acid sequences of the variable domains of antibodies belonging to various human immunoglobulin subgroups was compiled by Kabat et al., Sequences of Proteins of Immunological Interest, Fourth Edition, U.S. Dept. of Health & Human Services, pubs., (1987), a copy of which is attached to the enclosed Kelley Declaration as Exhibit "B". Kabat et al. grouped various heavy and light chain variable domains according to their amino acid sequence identity to form several human immunoglobulin "subgroups" i.e. human kappa light chains subgroups I to IV, human lambda light chains subgroups I to VI and human heavy chains subgroups I to III (see pages 41-76 and 160-167 of Kabat et al.). The "occurrences of most common amino acids" at each position of the variable domain are provided in the second to last column for each immunoglobulin subgroup in Kabat et al. The consensus human variable domain claimed in the above application is an amino acid sequence comprising the most commonly occurring amino acid residues at each position of the variable domain for a particular human immunoglobulin subgroup as defined by Kabat et al. It would have been readily apparent, to the ordinarily skilled biochemist, what constitutes a consensus human variable domain of a human immunoglobulin subgroup upon reading the above application.

Applicants respectfully request the withdrawal of the rejection of claim 1 as indefinite in light of the above submissions.

B. The Examiner has suggested that the "import amino acid" be described as "an import antibody comprising the amino acid sequence of a non-human antibody which binds to ...". Applicants understand that the Examiner considers that inclusion of the wording "import antibody" in parentheses is unclear and that the rejection relates to claims 1, 3, 4, 5 and 7. In order to overcome the rejection, claims 1, 7 and 19 have been amended to recite "an import antibody comprising a non-human antibody...". The non-human, import antibody may be the muMAb4D5 disclosed in Example 1 of the application, for example. Claims 3-5 depend on claim 1 and because there is clear antecedence basis for the phrases "import antibody variable domain amino acid

sequence", "import sequence" and "import antibody" in claim 1, the rejection of these claims is also rendered moot.

C. The Examiner has maintained the rejection of claim 1 under 35 U.S.C. §112, second paragraph, with respect to the wording "reasonably expected" on the grounds that it is not known what criteria determines what is "reasonable". In order to obviate the rejection, Applicants have deleted the word "reasonably" from claims 1, 3 and 19. Applicants respectfully submit that the amendment to the claims renders the rejection moot.

Applicants respectfully request that the maintained rejections of claims 1, 3-5 and 7 under 35 U.S.C. § 112, second paragraph, be withdrawn in light of the amendments to the claims and the submissions under paragraphs A to C above.

III. Objection and Rejections under 35 U.S.C. § 112, first paragraph

The Examiner has maintained the objection to the specification and the rejection of claims 1 to 11 under 35 U.S.C. § 112, first paragraph as lacking enablement. New claims 17 to 21 have also been rejected under 35 U.S.C. § 112, first paragraph as lacking enablement. The various sets of rejections are addressed separately below.

A. The Examiner has maintained the rejection of claim 1 and has rejected claims 19 to 21 for including the language "at least a portion". In the interests of expediting examination, claims 1, 7 and 19 have been amended by deleting the wording "at least a portion of" therefrom. Applicants submit that the amendment of the claims renders the rejection of claims 1 and 19-20 under 35 U.S.C. § 112, first paragraph, moot and respectfully request the withdrawal thereof.

B. The Examiner has maintained the rejection that step c) of claim 1 (i.e. the step of substituting an import CDR amino acid sequence for the corresponding human CDR amino acid sequence) is not enabled by the specification. The Examiner asserts that there is no clear guidance in the specification to enable one of ordinary skill in the art to make the human "consensus variable domain". The Examiner further asserts that the

Page No. 9

07/715272

only guidance presented in the specification with regards to the substitutions is the amino acid sequences of SEQ ID NO: 3 and 4. Applicants understand that the basis for the Examiner's rejection is that the information provided in the specification would not have enabled the ordinarily skilled biochemist to carry out the methods claimed in order to produce a humanized antibody.

Applicants respectfully traverse this rejection on the grounds that the specification is enabling for the method claimed. In support of the above position, a Declaration pursuant to 37 C.F.R. § 1.132 by Robert Kelley is attached. See specifically his opinion in paragraph 3 and the bases for this opinion set forth in paragraphs 4 to 7.

This Declaration was not earlier submitted because it was believed, in good faith, that the rejection would be overcome without the need for a Declaration. Applicants respectfully request the entry of this Declaration in the above application pursuant to Rule 116, because it does not introduce issues not already fully joined in this case. The Declaration is offered in a good faith effort to place this case in condition for allowance.

As discussed under section II (A) above and in paragraph 4 of the Kelley Declaration, the consensus human variable domain constitutes an amino acid sequence comprising the most commonly occurring amino acids at each position in the variable domain of a particular human immunoglobulin subgroup as defined by Kabat *et al.* The immunoglobulin subgroups referred to in Kabat *et al.* were grouped according to the amino acid sequence homology between human immunoglobulin *variable* domains, and the most commonly occurring amino acids at each position in the variable domain for each subgroup were identified (i.e. the "consensus human variable domain"). The skilled biochemist could have used the consensus human variable domains of the light chain and heavy chain subgroups having the greatest number of sequences therein (i.e. light chains kappa subgroup I and heavy chains subgroup III) as disclosed in Kabat *et al.* (see page 17, first paragraph of the specification) to humanize the non-human antibody of interest. Alternatively, the skilled biochemist could have chosen the consensus human variable domain and variable domain of another human immunoglobulin subgroup as defined in Kabat *et al.*

07/715272 ·

Page No. 10

i.e. the consensus human variable domain for human kappa light chains subgroups II to IV, human lambda light chains subgroups I to VI, or human heavy chains subgroups I or II (see pages 41-76 and 160-167 of Kabat et al.). Therefore, the skilled biochemist could have elected to use a consensus human variable domain other than those defined as SEQ ID NO: 3 & 4 on page 17 of the above application, as the consensus human variable domains for other subgroups were compiled in Kabat et al. Page ix of Kabat et al. identifies the residues forming the CDR regions of heavy and light chain variable domains tabulated from human and mouse variable domains. Kabat et al. have adopted standardized numbering for each of the residue locations. Accordingly, the skilled biochemist could have identified the CDR regions of the consensus human variable domain and the import variable domain using the teachings of Kabat et al. Alternatively, the structural definition of Chothia et al., J. Mol. Biol., 196: 901-917 (1987) (see page 16, third paragraph of the specification) could have been adopted to identify the CDR regions of the consensus and import variable domains. See paragraph 4 of the Kelley Declaration. The above submissions show that steps a & b of claim 1 were enabled by the specification as filed.

Also, step c of claim 1 could have been carried out by the ordinarily skilled biochemist using the information provided in the specification and techniques such as manual tabulation of amino acid sequences or a computer program which was known in the art prior to June 14, 1991. See paragraph 5 of the Kelley Declaration.

Steps d to g of claim 1 would similarly have been straightforward to perform. These steps of claim 1 relate to the identification of Framework Region (FR) residues in the consensus human variable domain which are non-homologous to the corresponding import FR residues and replacement of such non-homologous human residues with corresponding import residues, if the residues are expected to have any one of the effects specified in step f. The locations of FR residues in human and mouse variable domains are indicated in Kabat *et al.* (see page ix) and the structural definition of the FR's was available (see Chothia *et al.*) Hence, it would have been straightforward for the skilled

Page No. 11

07/715272

immunologist to identify the FR residues in the consensus human variable domain and the import sequence. Using computer programs available before June 14, 1991, the skilled biochemist would have been able to study the 3-dimensional structure of the antibody in order to establish whether a particular non-homologous import amino acid residue is likely to have one of the effects discussed in section f of claim 1. Information is provided on pages 14 to 16 of the specification which would have enabled the skilled biochemist to determine whether any non-homologous residue(s) would be expected to have the effects claimed. The techniques claimed in steps d to g of claim 1 could have been carried out routinely by a person versed in the relevant art, prior to June 14, 1991. See paragraph 6 of the Declaration.

As discussed in paragraph 7 of the Declaration, once the primary amino acid sequence of the antibody had been characterized, it would have been routine to make the protein using recombinant techniques or a peptide synthesizer, which techniques were well known in the art prior to the filing date of the above application.

Applicants conclude that, contrary to the Examiner's assertions, the ordinarily skilled biochemist would have been able to carry out the method claimed in the above application, using the information provided in the specification and techniques which were well known in the relevant art, prior to June 14, 1991.

Accordingly, Applicants request that the rejection of claim 1 under 35 U.S.C. § 112, first paragraph, be reconsidered and withdrawn in light of the above submissions and the Declaration.

C. The Examiner has maintained the rejection of claims 1 and 3, and has rejected claim 19 under 35 U.S.C. § 112, first paragraph, with respect to the wording "reasonably" therein. In order to obviate the rejection, the wording "reasonably" has been deleted from claims 1, 3 and 19.

Accordingly, Applicants request that the rejection of claims 1, 3 and 19 under 35 U.S.C. § 112, first paragraph, be withdrawn.

D. The Examiner has maintained the rejection of claims 6, 7 and 9 as lacking

Page No. 12

enablement under 35 U.S.C. § 112, first paragraph, the Examiner's position being that the amino acids are relevant to IgG only and not to other isotypes. Applicants respectfully traverse this rejection on the basis that the immunoglobulin sites claimed would have been relevant with respect to antibodies, other than IgG antibodies. Applicants refer the Examiner to paragraphs 8 & 9 of the Kelley Declaration which support this position. The Examiner appears to suggest that the rejected claims cover sequences which would not be found in immunoglobulin isotypes, other than IgG isotypes. However, as pointed out in paragraph 9 of the Kelley Declaration, the claims refer to positions or sites of the variable domain, not specific amino acid residues. These sites relate to the position of a residue in the 3-D structure of the variable domain. Kabat et al. have used universal numbering for the amino acid residue locations of the variable domains for each of the immunoglobulin subgroups mentioned therein. The FR residue sites indicated may be occupied by an amino acid residue which is non-homologous to the corresponding consensus human variable domain residue, and which residue is likely to have at least one of the effects discussed in step f of claim 1. The residue at the particular site can be any amino acid residue, depending on the antibody in which it is located. These residue locations or sites are applicable across species (see page 16, line 8). Accordingly, it is likely that an amino acid residue located at one of the sites indicated in claims 6, 7 and 9 will have one of the effects of claim 1 (step f) regardless of the antibody in which it is located. It is apparent that the particular sites claimed are applicable to immunoglobulins other than IgG.

Accordingly, Applicants submit that the rejection of claims 6, 7 & 9 under 35 U.S.C. § 112, first paragraph, should be reconsidered and withdrawn in light of the above submissions and Declaration.

In light of the submissions presented in paragraphs A to D above, Applicants respectfully request that the objection to the specification and the rejection of claims 1-11 and 17-21 under 35 U.S.C. §112, first paragraph, be withdrawn.

Applicants thank the Examiner for withdrawing the rejections which were raised

under 35 U.S.C. § 101 in the earlier Office Action dated October 5, 1992 .

IV. Rejection of claims 1, 2 and 5-10 under 35 U.S.C. 102 (a) and 102(b)

The rejection of claims 1, 2 and 5-10 under 35 U.S.C. § 102(a) and 102(b) has been maintained and newly added claims 17-21 have been rejected under 35 U.S.C. § 102(a) and 102(b) as being anticipated by Queen *et al.*, *Proc. Natl. Acad. Sci.*, 86:10029-10033 (1989) and Co *et al.*, *Proc. Natl. Acad. Sci.*, 88:2869-2873 (1991). The basis for the rejection is that there is allegedly no clear indication as to what is meant by the consensus human variable domain claimed in the above application.

To constitute anticipation, all material elements of a claim must be found in one prior art source. *In re Marshall*, 198 USPQ 344 (CCPA 1978), *In re Kalm*, 154 USPQ 10 (CCPA 1967). Applicants will show that Queen *et al.* and Co *et al.* do not contain all material elements of claims 1, 2, 5-10 and 17-21.

The nature of the "consensus human variable domain of a human immunoglobulin subgroup" as defined in the claims as amended has been discussed above under Section II(A) of this response and in paragraph 4 of the Kelley Declaration, those discussions being incorporated herein. Applicants submit that the meaning of the phrase consensus human variable domain of a human immunoglobulin subgroup would have been clearly understood by those skilled in the art upon reading the specification. The prior art relied upon in the Office Action fails to disclose a method of making a humanized antibody using a consensus human variable domain to "humanize" a nonhuman antibody. The Declaration by Kelley supports this position. In particular, Applicants direct the Office's attention to paragraphs 11-13 of the attached Declaration. It is apparent from the information given in Table 1 of Exhibit C and in the Figures of Exhibits D and E of the Kelley Declaration (see paragraphs 12 & 13 thereof), that the variable domains of the human immunoglobulin sequences used by Queen *et al.* and Co *et al.* are not a consensus human variable domain of any human immunoglobulin subgroup as set forth in the claims of the above application.

Page No. 14

Since, as shown above, Queen *et al.* and Co *et al.* do not teach all the material elements of the instant claims as required under *Marshall* and *Kalm, supra*, Applicants respectfully submit that the rejection of claims 1, 2, 5-10 and 17-21 under 35 U.S.C. § 102(a) and (b) can not be upheld and therefore request that the rejections be withdrawn.

V. Rejection of claims 3 and 4 under 35 U.S.C. § 103

The rejection of claims 3 and 4 as unpatentable under 35 U.S.C. § 103 over Queen *et al.*, or Co *et al.*, *supra*, in view of Wallick *et al.*, *J. Exp. Med.*, **168** (1988) has been maintained. The basis for the rejection relates to the alleged lack of clarity of the language "consensus human variable domain" in the claims of the above application. The consensus human variable domain as defined in the above application would have been readily understood by the ordinarily skilled biochemist (see paragraph 4 of the Kelley Declaration). Claim 1 of the above application relates to a method of using a consensus human variable domain to "humanize" a non-human antibody (e.g. muMAb4D5). As established in section IV above, use of a consensus human variable domain from a human immunoglobulin subgroup is not disclosed in Queen *et al.* or Co *et al.*

The publication by Wallick *et al.* does not compensate for the deficiencies in the primary references. Wallick *et al.* refer to the importance of glycosylation for maintaining antigen binding affinity of monoclonal antibodies. Wallick *et al.* fail to disclose or suggest a method of humanizing a non-human antibody, much less a method of humanizing a non-human antibody, much less a method of humanizing a non-human antibody. The skilled biochemist would have had no motivation to use a consensus human variable domain based on the prior art referred to in the Office Action, because the prior art techniques had all relied upon using a human variable domain sequence which has the closest sequence homology to the non-human variable sequence (to be humanized) in order to reduce the likelihood of introducing distortions into the CDR's (see column 2 on page 10031 of Queen *et al.*) and "to retain high binding affinity in the humanized antibody" (see column 1 on page 2871 of Co *et al.*). The method claimed in

Page No. 15

07/715272

the above application does not rely on a high degree of homology between the variable domain of the non-human sequence and the consensus variable domain which is used to humanize the non-human sequence.

Also, as supported by paragraph 15 of the Kelley Declaration, the invention claimed in the above application resulted in an unexpected result which could not have been reasonably predicted from the prior art. It was surprising that a consensus variable domain of a selected immunoglobulin subgroup could be used to humanize a non-human antibody, regardless of the degree of homology between the human and non-human amino acid sequences. It was also surprising that the humanized antibody so formed retained, and in some instances, had increased antigen binding affinity compared to the non-human antibody from which it was derived. The above application shows that the huMAb4D5-8 variant actually binds the p185^{HER2} ECD 3-fold more tightly than muMAb4D5 (see page 82 lines 31 & 32 to page 83, line 1 of the specification) which could not have been predicted by the ordinarily skilled biochemist. See paragraph 15 of the Kelley Declaration. The evidence of unexpected results in Applicants' application is sufficient to support a conclusion of nonobviousness. *Ralston Purina Co. Far-Mar-Co., Inc.,* 222 USPQ 863 (DC KS, 1984).

It is apparent that the invention claimed in claim 1 was novel and nonobvious over the citations because the combination of the prior art failed to disclose, or suggest, the invention claimed in claim 1 and, moreover, the method resulted in a new and unexpected result which could not have been reasonably predicted from the art.

Claims 3 & 4 depend on claim 1 which, as established above, is novel and nonobvious over the citations. Claim 3 refers to the step of finding any glycosylation site which is likely to affect the antigen binding or affinity in the import antibody and substituting the glycosylation site *into* the *consensus* amino acid sequence. Claim 4 refers to the step of *replacing* glycosylation sites of the consensus domain with the corresponding import amino acid residues if such glycosylation sites are not present in the import sequence. These claims would not have been obvious over the prior art of

Page No. 16

07/715272

record because the prior art failed to disclose the use of a human consensus variable domain to humanize the non-human antibody. Accordingly, the skilled biochemist would have had no motivation to replace or insert glycosylation sites into a consensus amino acid sequence, as claimed in claims 3 and 4 of the application. See paragraph 15 of the Kelley Declaration.

The law is clear that obviousness cannot be established by combining the teachings of the references to produce the claimed invention, absent some teaching, suggestion, or incentive supporting the combination. *ACS Hospital Systems, Inc. v. Montefiore Hospital*, 221 USPQ 929, 933 (Fed. Cir. 1984). The above discussion shows that the cited references, alone or in combination, lack the requisite teaching of the use of a consensus human variable domain to humanize a non-human antibody. In this case, the combined art would not have reasonably enabled or motivated the skilled practitioner to use a human consensus variable domain in this manner, which provides a method of making improved humanized antibodies. Accordingly, it is clear that the invention claimed in claims 3 & 4 is novel and nonobvious over the prior art of record.

Applicants submit that the rejection of claims 3 and 4 under 35 U.S.C. § 103 should be reconsidered and withdrawn in light of the above submissions and the Declaration.

VI. Rejection of claims 17,18, 20 and 21 under 35 U.S.C. §112, second paragraph.

Claims 17, 18, 20 and 21 have been rejected under 35 U.S.C. § 112, second paragraph, as being indefinite in that there are allegedly no discrete method steps. In order to obviate the rejection, claims 17, 20 and 21 have been amended to each recite a definite method step and claim 18 has been deleted.

Applicants respectfully request the withdrawal of the rejection of claims 17, 20, and 21 under 35 U.S.C. § 112, second paragraph, in light of the amendments to the claims.

As all objections and rejections have been addressed and overcome, Applicants

w014.u

Page No. 17

believe that the claims are now in condition for allowance. Notice to that effect is respectfully requested. If the Examiner has any questions concerning the response, she should feel free to call the undersigned attorney at the number indicated above.

Respectfully submitted, GENENTECH, INC.

Janel E. Hasek

Date: September 20, 1993

Janet E. Hasak Reg. No. 28,616

Date: September 20, 1993

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231.



PATENT DOCKET 709

THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Paul J. Carter et al.

Serial No. 07/715272

Filed: June 14, 1991

For: Immunoglobulin Variants

Group Art Unit: 1806

Examiner: L. FEISEE

460 Point San Bruno Boulevard South San Francisco, CA 94080

DECLARATION OF ROBERT F. KELLEY PURSUANT TO 37 CFR §1.132

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

I, ROBERT F. KELLEY, do hereby declare as follows:

1. I received my Ph.D. in Biochemistry in 1984 from the University of Iowa. Following my Ph.D, I was a NIH postdoctoral fellow in the Department of Molecular Biophysics & Biochemistry at Yale University from July 1984 to December 1985. In 1986, I joined the Biocatalysis Department at Genentech, Inc. as an Associate Scientist. In September 1988, I was promoted to Scientist and I am employed in that capacity at present. (The Biocatalysis Department has been renamed "Protein Engineering"). I am the author or co-author of 22 publications relating to the 3-D structures and folding of various proteins. A copy of my curriculum vitae is attached as Exhibit "A".

2. I understand that the Patent Office has rejected the above application on the basis that the application as filed does not provide sufficient disclosure to enable a skilled biochemist to carry out the method of claim 1 because the Examiner believes no clear guidance exists in the specification to allow a skilled biochemist to make the "consensus human variable domain" and substitute an import (i.e. non-human) Complementary Determining Region (CDR) amino acid sequence for the corresponding human CDR amino acid sequence, as set forth in claim 1. I further understand that the Office considers that

Page No. 2

07/715272

the only guidance in the specification with regards to the substitutions is the amino acid sequences of SEQ ID NO: 3 and 4.

3. I have read the above application, the Office Action date May 19, 1992 (Paper # 17) rejecting the claims of the application, and the proposed amendment of the claims in response to the rejection. In my opinion, the skilled biochemist could have readily carried out the method of claim 1 in order to make a humanized antibody, using the general knowledge available in the field on and before June 14, 1991, and the information given in the above application. The bases for my opinion are given in paragraphs 4 to 7 below.

4. Claim 1 relates to a method of making a humanized antibody. Steps a and b of claim 1, as amended, discuss identification of the CDR amino acid sequences of a non-human import antibody (to be humanized) and a consensus human variable domain of a human immunoglobulin subgroup. The consensus human variable domain constitutes an amino acid sequence comprising the most commonly occurring amino acids at each position in the variable domain of a particular human immunoglobulin subgroup as defined by Kabat et al., Sequences of Proteins of Immunological Interest, Fourth Edition, U.S. Dept. of Health & Human Services, pubs., (1987), a copy of which is attached as Exhibit "B". The immunoglobulin subgroups referred to in Kabat et al. were grouped according to the amino acid sequence homology between human immunoglobulin variable domains, and the most commonly occurring amino acids at each position in the variable domain for each subgroup were identified (i.e. the "consensus human variable domain"). The skilled biochemist could have used the consensus human variable domains of the light chain and heavy chain subgroups having the greatest number of sequences (i.e. light chains kappa subgroup I and heavy chains subgroup III) as disclosed in Kabat et al. (see page 17, first paragraph of the specification) to humanize the non-human antibody of interest. Alternatively, the skilled biochemist could have chosen the consensus human variable domain of another human immunoglobulin subgroup as defined in Kabat et al. (i.e. the consensus human variable domain for human kappa light chains subgroups II to IV, human lambda light chains subgroups I to VI, or human

Page No. 3

heavy chains subgroups I or II [see pages 41-76 and 160-167 of Kabat *et al.*]). Therefore, the skilled biochemist could have elected to use a consensus human variable domain other than those defined as SEQ ID NO: 3 & 4 on page 17 of the above application, as the consensus human variable domains for other subgroups were compiled in Kabat *et al.* Page ix of Kabat *et al.* identifies the residues forming the CDR regions of heavy and light chain variable domains tabulated from human and mouse variable domains. Kabat *et al.* have adopted standardized numbering for each of the residue locations. Accordingly, the skilled biochemist could have identified the CDR regions of the consensus human variable domain and the import variable domain using the teachings of Kabat *et al.* Alternatively, the structural definition of Chothia *et al., J. Mol. Biol.,* **196**: 901-917 (1987) (see page 16, third paragraph of the specification) could have been adopted to identify the CDR regions of the consensus and import variable domains. Hence, it would have been straightforward for the skilled biochemist to carry out steps a and b of claim 1 using the information provided in the specification.

5. Step c of claim 1 discloses the step of replacing the corresponding human CDR sequence with the import CDR amino acid sequence. This step could have been carried out routinely by the skilled biochemist by manual tabulation or using a computer program such as the ALIGN program, (Dayhoff *et al., Meth. Enzymol.*, 91:524-545 [1983]) which was available prior to June 14, 1991. Steps a to c of claim 1 would have resulted in the characterization of a primary amino acid sequence encoding a humanized variable domain with import (non-human) CDR regions.

6. Steps d to g of claim 1 relate to the identification of Framework Région (FR) residues in the consensus human variable domain which are non-homologous to the corresponding import FR residues and replacement of such non-homologous human residues with corresponding import residues, if the residues are expected to have any one of the effects specified in step f. The locations of FR residues in human and mouse variable domains are indicated in Kabat *et al.* (see page ix) and the structural definition of the FR's was available (see Chothia *et al.*) Hence, it would have been straightforward for the skilled immunologist to identify the FR residues in the consensus human variable domain and the

Page No. 4

import sequence. Using computer programs (such as the INSIGHT program [Biosym Technologies], available before June 14, 1991), the skilled biochemist would have been able to study the 3dimensional structure of an antibody in order to establish whether a particular non-homologous import amino acid residue is likely to have one of the effects discussed in section f of claim 1. Information is provided on pages 14 to 16 of the specification which would have enabled the skilled biochemist to determine whether any non-homologous residue(s) would be expected to have the effects claimed. The techniques claimed in steps d to g of claim 1 could have been carried out routinely by a person versed in the relevant art, prior to June 14, 1991.

7. Steps a to g of claim 1 would have lead to the characterization of an amino acid sequence of a humanized antibody having non-human CDR amino acid residues and, optionally, having one or more non-human FR residues. In order to prepare the humanized antibody as claimed in claim 1, step h, the skilled biochemist could have synthesized the antibody using a peptide synthesizer which was commercially available before June 14, 1991. Alternatively, the antibody could have been made in recombinant cell culture (see page 26, last paragraph of the specification). Preparation of the antibody would have been straightforward to perform by the person skilled in the art, once the amino acid sequence of the humanized antibody had been characterized.

8. I understand that the Patent Office has rejected the above application on the basis that the sites in the variable domain referred to in claims 6, 7, and 9 are relevant to IgG antibodies only. It is my opinion that the sites referred to in claims 6, 7, and 9 would be relevant to other immunoglobulins. The basis for my opinion is given in paragraph 9 below.

9. The sites referred to in claims 6, 7, and 9 are the residue locations, or sites, of the FR residues in the heavy or light chain forming the variable domain of immunoglobulins. The residue sites referred to in claims 6, 7 & 9 relate to the position of a residue in the 3-D structure of the variable domain. Kabat *et al.* have used universal numbering for the amino acid residue locations of the variable domains for each of the immunoglobulin subgroups mentioned in the reference. The FR residue sites

Page No. 5

indicated may be occupied by an amino acid residue which is non-homologous to the corresponding consensus human variable domain residue, and which is likely to have at least one of the effects discussed in step f of claim 1. These residue locations or sites are applicable *across species* (see page 16, line 8 of the specification). Accordingly, it is likely that an amino acid residue located at one of the sites indicated in claims 6, 7 and 9 will have one of the effects of claim 1 (step f), regardless of the antibody in which it is located, because it will be in the same position in the 3-D structure of the antibody variable domain as the residue sites referred to in the rejected claims. Accordingly, the examples of residue locations to be substituted in the variable domains would be applicable to antibodies, other than IgG antibodies.

10. I understand that the Patent Office has rejected the above application on the grounds that the invention as claimed is disclosed in Queen *et al.*, *Proc. Natl. Acad. Sci.*, 86:10029-10033 (1989) or Co *et al.*, *Proc. Natl. Acad. Sci.*, 88:2869-2873 (1991) and that the Office has suggested that the human variable domains disclosed in these references may have the same amino acid sequences as one of the consensus human variable domains disclosed in Kabat *et al.*

11. The above statements regarding the state of knowledge as of June 14, 1991, do not establish that the invention claimed in this application was known, or would have been obvious, to the skilled biochemist at the time the invention was made. To the contrary, after having read the citations relied upon by the Patent Office, it is my judgement that these documents would not have disclosed, nor suggested, the methods claimed. The basis for my opinion is given below.

12. The invention of the above application can be distinguished on the basis that a *consensus human variable domain* is used to "humanize" a non-human antibody of interest. The Queen *et al.* and Co *et al.* publications fail to disclose a consensus human variable domain. Instead, these publications refer to the use of a human variable domain having the closest sequence homology to the variable domain of the non-human antibody to be humanized. Queen *et al.* used the Eu human variable domain sequence (see Fig 2 thereof) and Co *et al.* used the variable domains of the Pom or Eu human

Page No. 6

antibodies (see Fig 1 thereof). The sequences used in Queen *et al.* and Co *et al.* do not constitute a consensus human variable domain of a human immunoglobulin subgroup. The sequence identity between the amino acid sequences of the FR residues of the variable domains of the Pom or Eu heavy or light chains compared to the FR residues of the consensus human variable domains of each of the human immunoglobulin subgroups as defined by Kabat *et al.* is illustrated in Table 1 (see Exhibit "C", attached hereto). The CDR residues were not used in the comparison because of the large number of differences between these residues for variable domains of different antibodies. The Pom and Eu variable domain sequences were taken from Kabat *et al.* The consensus human variable domains of the V_L lambda subgroups IV and V were not compared, as these subgroups have too few members. While the variable domain of Eu is classified in subgroups V_L kappa I and V_HII, and the variable domain of Pom is classified in subgroups V_L kappa III and V_H III, it is apparent that the Eu and Pom variable domain amino acid sequences are not consensus human variable domains of any immunoglobulin subgroup. This is further demonstrated in the following paragraph.

13. Exhibits "D" and "C" attached hereto, show the differences in the amino acid sequences of the Pom and Eu heavy and light chain variable domains compared to the consensus human variable domain of the subgroup in which they are classified. Exhibit D illustrates an alignment of the amino acid sequences of the light chain variable domains of Eu, Pom and the consensus variable domain of the V_L kappa subgroup I (in which the light chain variable domain of Eu is classified). Exhibit E illustrates an alignment of the amino acid sequences of the heavy chain variable domains of Eu, Pom and the consensus variable domain of the V_H subgroup I (in which the light chain variable domain of Eu is classified). Exhibit E illustrates an alignment of the amino acid sequences of the heavy chain variable domains of Eu, Pom and the consensus variable domain of the V_H subgroup III (in which the heavy chain variable domain of Pom is classified). Even though Eu is classified in V_L kappa I, it has seven framework residues which are different from the framework residues of the kappa I consensus sequence. Furthermore, while Pom is classified in the V_H III subgroup, eight of its framework residues differ from the corresponding framework residues of the V_H III consensus sequence. There are, of course, many differences between the CDR residues of the consensus sequences and the corresponding CDR residues of Pom and Eu.

Page No. 7

07/715272

It is clear from the information in Exhibits C, D, & E that the Queen *et al.* and Co *et al.* publications fail to disclose a method wherein a non-human import antibody is humanized using a consensus human variable domain of an immunoglobulin subgroup.

14. I understand the Patent Office has rejected the above application on the basis that the invention claimed in claims 3 & 4 would have been obvious in light of Queen *et al.*, or Co *et al.*, when read in conjunction with Wallick *et al.*, *J. Exp. Med.*, **168** (1988). After reading these references, it is my opinion that the invention claimed in claims 3 and 4 is novel and would not have been obvious in light of the citations. The basis for my opinion is given in the following paragraph.

15. Claim 1 of the above application relates to a method of using a consensus human variable domain to "humanize" a non-human antibody (e.g. muMAb4D5). Use of a consensus human variable domain from a human immunoglobulin subgroup to humanize a non-human antibody is not disclosed in Queen et al., Co et al. or Wallick et al. Wallick et al. does not relate to a method of humanizing a non-human antibody, much less a method of humanizing a non-human antibody using a consensus human variable domain of a human immunoglobulin subgroup. The skilled biochemist would have had no motivation at the filing date of this application to use a consensus human variable domain to humanize a non-human antibody, because the techniques in the prior literature had all relied upon using a human variable domain sequence which has the closest sequence homology to the non-human variable sequence (to be humanized) in order to reduce the likelihood of introducing distortions into the CDR's (see column 2 on page 10031 of Queen et al.) or to "retain high binding affinity in the humanized antibodies" (see column 1 on page 2871 of Co et al.). The method claimed in the above application does not rely on a high degree of homology between the variable domain of the non-human sequence and the consensus variable domain which is used to humanize the non-human sequence. It was surprising that a consensus variable domain of a selected immunoglobulin subgroup could be used to humanize a non-human antibody, regardless of the degree of homology between the human and nonhuman amino acid sequences. It was also surprising that the humanized antibody so formed retained,

Page No. 8

and in some instances, had increased antigen binding affinity compared to the non-human antibody from which it was derived. The above application shows that the huMAb4D5-8 variant actually binds the p185^{HER2} ECD 3-fold more tightly than muMAb4D5 (see page 82 lines 31 & 32 to page 83, line 1 of the specification), which could not have been predicted by the ordinarily skilled biochemist at the time the specification was filed. Claim 3 refers to the step of finding any glycosylation site which is likely to affect the antigen binding or affinity in the import antibody and substituting the glycosylation site into the consensus amino acid sequence. Claim 4 refers to the step of replacing glycosylation sites of the consensus domain with the corresponding import amino acid residues if such glycosylation sites are not present in the import sequence. In my opinion, these claims would not have been obvious over the prior literature because the reference failed to disclose the use of a human consensus variable domain to humanize the non-human antibody. Accordingly, the skilled biochemist would have had no motivation to replace or insert glycosylation sites into a consensus amino acid sequence, as claimed in claims 3 and 4 of the application.

16. I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Dated: <u>9/20/93</u> Signed: <u>Motert F. Kelley</u> ROBERT F. KELLEY

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, on September 20, 1993.

Dated: September 20, 1993

Louise Strasbaugh

XHIBIT A

Robert F. Kelley

September, 1993

Work address:

Genentech, Inc.

Protein Engineering Dept.

South San Francisco, CA 94080

Home address:

1029 San Felipe Ave.

San Bruno, CA 94066

Personal information:

Birthdate: November 7, 1957

Married, Wife's name: Wendy L. Kelley

One child: Brian F. Kelley

Education:

B.S., Biochemistry, Michigan State University, 1979

Ph. D., Biochemistry, University of Iowa, 1984

Employment positions:

Teaching assistant, Chemistry Dept., Michigan State Univ., 1978-1979 Teaching assistant, Biochemistry Dept., University of Iowa, 1979-1981 Graduate research assistant, Biochemistry Dept., University of Iowa, 1982-

April 1984

Postdoctoral associate, Biochemistry Dept., University of Iowa, May 1984-June 1984

NIH postdoctoral fellow, Dept. of Molecular Biophysics & Biochemistry, Yale University, July 1984-Dec. 1985

Associate Scientist, Biocatalysis Dept., Genentech, Inc., January 1986-September 1988

Scientist, Biomolecular Chemistry Dept., Genentech, Inc., September 1988-present

Awards and membership in professional organizations:

Biophysical Society, 1983-present

American Chemical Society, 1991-present

Scientific publications

- Kelley, R.F., & Stellwagen, E. (1984) "Conformational transitions of thioredoxin in guanidine hydrochloride", Biochemistry 23, 5095-5102.
- Kelley, R.F., Wilson, J., Bryant, C., & Stellwagen, E. (1986)"Effects of guanidine hydrochioride on the refolding kinetics of denatured thioredoxin", Biochemistry 25, 728-732.
- Wilson, J., Kelley, R.F., Shalongo, W., Lowery, D., & Stellwagen, E. (1986) "Equilibrium and kinetic measurements of the conformational transition of thioredoxin in urea", Biochemistry 25, 7560-7566.
- Kelley, R., Wilson, J., Bryant, C., Shalongo, W., Ledger, R., Lowery, D., & Stellwagen, E. (1986) "Folding of E. coli thioredoxin into its native conformation" in Thioredoxin and Glutaredoxin Systems: Structure & Function (Holmgren, A., Branden, C.-L., Jornvall, H., & Sjoberg, B.-M., Eds.) pp 77-85, Raven Press, New York.
- Kelley, R., Ledger, R., Shalongo, W., & Stellwagen, E. (1986)" Spectral and hydrodynamic methods for identification of intermediates in protein folding", UCLA Symp. Mol. Chem. Biol., New Ser. 39, 269-281.
- Kelley, R.F., Shalongo, W., Jagannadham, M.V., & Stellwagen, E. (1987) "Equilibrium and kinetic measurements of the conformational transition of reduced thioredoxin", Biochemistry 26, 1406-1411.
- Kelley, R.F., & Richards, F.M. (1987) "Replacement of proline 76 with alanine eliminates the slowest kinetic phase in thioredoxin folding", Biochemistry 26, 6765-6774.
- Cleary, S., Mulkerrin, M.G., & Kelley, R.F. (1989) "Purification and characterization of tissue plasminogen activator kringle-2 domain expressed in E. coli", Biochemistry, 28, 1884-1891.


Kelley, R.F., & Cleary, S. (1989) "Effect of residue 65 substitutions on thermal stability of t-PA kringle-2 domain", Biochemistry 28, 4047-4054.

EXHIBIT C

TABLE 1 SEQUENCE IDENTITY - (%)

CONSENSUS VARIABLE DOMAIN SUBGROUP	EU	POM		
V, kappa I	92	76		,
V _i kappa II	61	71		
V _L kappa III	72	85		•
V, kappa IV	73	78	100 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 - 100 -	
V, lambda l	61 .	59		
V, lambda II	57	54		
V _L lambda III	59	56		
V, lambda VI	52	49		
V _H I	83	64		
V _H II	53	62		
V _H III	61	91		

EXHIBIT D



Kappa-1	AASSLESGVPSRFSGSGSGTDFTLTISSLOPEDFA 0 0000 * * GASTRATGIPARFSGSGSGSGTEFTLTISSLOSEDFA	QQ Q VYYCQQYNNWPPTFGQ
POM		CDR-L3
	CDR-L2	

EU	GTKVEVKGT
Kappa-I	GTKVEIKRT
POM	GTRVEIKR

KEY: • = differences in FR residues

@ = differences in CDR residues



EXHIBIT E

Variable Heavy Domain

human-III	EVOLVESGGGI	LVOPGGSLRL	SCAASGFT	FSSYA Q	MSWVRQA	PGKGLEWVS
POM	EVOLLESGGG	LVOPGGSLRL	SCAASGFT	FSSSA	MSWVRQA	PGKGLEWVZ
		•		CDR	-H1	
	50 a.	60	70	80	abc	90
EU	GIVPMFGPPN @ @@@@ @@@	★ 66660	** * *	* **	** *	R A
human-III	VIŠČDČGŠTY COCCO COCC	YADŠVKGRFI A	**	*	*	*
POM	WKYENGNDKH	YADSVNGRFI	ISRNDSKN	TLYLI	MNSLQAE	DTALYYCA
	CDR-H2					
		11				
		EYNGGLV	TVSS			
EU	GYGIYSPE	*** *				
EU human-III	GYGIYSPE GRCGGSDY GRCGGSDY QQ QQQQ	WGQGTLV	TVSS			

KEY: • = differences in FR residues

@ = differences in CDR residues

RODert F. Kelley

- Byeon, I.-J.L., Kelley, R.F., & Llinas, M. (1989) "¹H-NMR structural characterization of a recombinant kringle-2 domain from human tissue-type plasminogen activator", Biochemistry 28, 9350-9360.
- Kelley, R.F., & Winkler, M.E. (1990) "Folding of eukaryotic proteins produced in Escherichia coli" Genetic Engineering (ed. J.K. Setlow) 12, 1-19.
- Byeon, I.-J., Kelley, R.F., & Llinas, M. (1991) "Kringle-2 domain of the tissuctype plasminogen activator: ¹H-NMR assignments and secondary structure" Eur. J. Biochem. 197, 155-165.
- Kelley, R.F., DeVos, A.M., & Cleary, S. (1991) "Thermodynamics of ligand binding and denaturation for His64 mutants of tissue plasminogen activator kringle-2 domain" Proteins: Structure, Function and Genetics 11, 35-44.
- Menhart, N., Sehl, L.C., Kelley, R.F., & Castellino, F.J. (1991) "Construction, expression and purification of recombinant kringle 1 of human plasminogen and analysis of its interaction with ω -amino acids" Biochemistry 30, 1948-1957.
- deVos, A.M., Ultsch, M.H., Kelley, R.F., Padmanabhan, K., Tulinsky, A., Westbrook, M.L., & Kossiakoff, A.A. (1991) "Crystal structure of the kringle-2 domain of tissue plasminogen activator at 2.4 Å resolution" Biochemistry, 31, 270-279.
- Garrard, L.J., Yang, M., O'Connell, M.P., Kelley, R.F., & Henner, D.J. (1991) "Fab assembly and enrichment in a monvalent phage display system" Bio/Technology, 9, 1373-1377.
- Carter, P., Kelley, R.F., Rodrigues, M.L., Snedecor, B.R., Covarrubias, M., Velligan, M.D., Wong, W.-L. T, Rowland, A.M., Kotts, C.E., Carver, M.E., Yang, M., Bourell, J.H., Shepard, H.M., & Henner, D.J. (1992)
 "High level Escherichia coli expression and production of a bivalent humanized antibody fragment" Bio/Technology, 10, 163-167.

3

- Kelley, R.F., O'Connell, M.P., Carter, P., Presta L., Eigenbrot, C., Covarrubias, M., Snedecor, B., Bourell, J.H., & Vetterlein, D. (1992) "Antigen binding thermodynamics and anti-proliferative effects of chimeric and humanized antip185^{HER2} antibody Fab fragments" Biochemistry, 31, 5434-5441.
- Kelley, R.F., O'Connell, M.P., Carter, P., Presta, L., Eigenbrot, C.,
 Covarrubias, M., Snedecor, B., Speckart, R., Blank, G., Vetterlein, D., &
 Kotts, C. (1993) "Characterization of humanized anti-p185^{HER2} antibody Fab fragments produced in *E. coli*" ACS Symposium Series No. 526, *Protein Folding: In Vivo and In Vitro*, J.L. Cleland & J. King, eds., pp. 218-239.

Kelley, R.F., & O'Connell, M.P. (1993) "Thermodynamic analysis of an antibody functional epitope" Biochemistry, 32, 6828-6835.

Huber, A.H., Kelley, R.F., Gastinel, L.N, & Bjorkman, P.J. (1993) "Crystallization and stoichiometry of binding of a complex between a rat intestinal Fc receptor and Fc" J. Mol. Biol., 230, 1077-1083.

292

Kelley, R.F. (1993) "Engineering Therapeutic Antibodies", in Protein Engineering: Principles & Practice, eds. J.L. Cleland & C. Craik, Hanser, in press.

GLU GLU GLU VAL VAL VAL GLN GLN GLN LEU LEU VAL VAL VAL SEP TRP TRF FOURTH EDUTION VAL ARG

VAL QUU thr GLY GLY

SER GLY PHE THR Val

SER

RF

ARG

ÚLY THR

ŝ

NOR AD A POST AT ALL OF AD

H

G: SE

PRO(.9

120 00 40 69 ŠER 1.60 RESIDUE

hable Cally

GLN

GLN

GLN

Bet

GLU VAL GLN

5

LEU(.99)

ΥS

ALA(.95) SER(.97) GLY(.95) PHE(.97)

BG(.95

APC

iLE-SÈR

MET1.951

CYS

PHF

LEU

THR

THF

91

109

212224

30

38

38 39 40

4344546

SERU96

1987

50

DERARTMENT OF HEALTH

NO HUMAN SERVICES Public Health Service ional Institutes of Health CYS ALA ARG PHE GLY LYS GLY THR THR

VAL THR PRO GLU ASP

λAL

SER

TRE

GL)

SEF

ANG

SER

PRO

PHE -

PHE

TAP

EU

GLU ALA GLU

тня

ALA VAL TYR TYR CYS ALA ARG

THR PR

GL

SEA

ASP

THE

GLY GLN GLY THR LEU

VÁL SER

AS SF

GLL VATA SE



ÜÞ 190 1

1:12

< 1,

FOURTH EDITION

Tabulation and Analysis of Amino Acid and Nucleic Acid Sequences of Precursors, V-Regions, C-Regions, J-Chain, T-Cell Receptor for Antigen, T-Cell Surface Antigens, β_2 -Microglobulins, Major Histocompatibility Antigens, Thy-1, Complement, C-Reactive Protein, Thymopoletin, Post-gamma Globulin, and α_2 -Macroglobulin

1987

Elvin A. Kabat*, Tai Te Wu[†], Margaret Reid-Miller[‡], Harold M. Perry[‡], and Kay S. Gottesman[‡]

* Depts. of Microbiology, Genetics and Development, and Neurology, Cancer Center/ Institute of Cancer Research, College of Physicians and Surgeons, Columbia University, New York, NY 10032 and the National Institute of Allergy and Infectious Diseases, Bethesda, MD 20892.

† Depts. of Biochemistry, Molecular Biology, and Cell Biology, and Engineering Sciences and Applied Mathematics and Biomedical Engineering, Northwestern University, Evanston, IL 60201 and the Cancer Center, Northwestern University Medical School, Chicago, IL 60611

‡ Bolt Beranek and Newman Inc., Cambridge, MA 02238

0057/0

The collection and maintenance of this data base is sponsored through Contract N01-RR-8-2158 by the following components of the National Institutes of Health, Bethesda, MD 20892:

Division of Research Resources

National Cancer Institute National Institute of Allergy and Infectious Diseases National Institute of Arthritis, Diabetes, Digestive and Kidney Diseases National Institute of General Medical Sciences

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

Public Health Service National Institutes of Health

(1987)

GENENTECH, INC. 460 Pt. San Brune Blvd. South San Francisco, CA 94080

iii TABLE OF CONTENTS

INTRO	DUCTION
SIGNA	L SEQUENCES
210144	SIGNAL PEPTIDES OF KAPPA LIGHT CHAINS 1 SIGNAL PEPTIDES OF LAMBDA LIGHT CHAINS 8 SIGNAL PEPTIDES OF LAMBDA LIGHT CHAINS 9 SIGNAL PEPTIDES OF HEAVY CHAINS 11 SIGNAL PEPTIDES OF BETA-2-MICROGLOBULINS 19 SIGNAL PEPTIDES OF MACROGLOBULINS 21 SIGNAL PEPTIDES OF I REGION ANTIGENS CLASS I 21 SIGNAL PEPTIDES OF I REGION ANTIGENS CLASS I 24 SIGNAL PEPTIDES OF COMPLEMENT 30 SIGNAL PEPTIDES OF T-CLYMPHOCYTE RECEPTOR FOR ANTIGEN 30 SIGNAL PEPTIDES OF T-CELL SUFFACE ANTIGENS 36 SIGNAL PEPTIDES OF OTHER PROTEINS 36 SIGNAL PEPTIDES OF T-CELLS SUFFACE ANTIGENS 36 SIGNAL PEPTIDES OF T-CELLS SUFFACE ANTIGENS 36 SIGNAL PEPTIDES OF OTHER PROTEINS 36
VARIA	BLE REGION LIGHT CHAIN SEQUENCES 41 HUMAN KAPPA LIGHT CHAINS SUBGROUP 1
	HUMAN KAPPA LIGHT CHAINS SUBGROUP II
	HUMAN LAMBDA LIGHT CHAINS SUBGROUP II
	MOUSE KAPPA LIGHT CHAINS I
	DAT KARRA LIGHT CHAINS
	RABBIT KAPPA LIGHT CHAINS
	OTHER KAPPA LIGHT CHAINS
VARIA	BLE REGION HEAVY CHAIN SEQUENCES
•	HUMAN HEAVY CHAINS SUBGROUP I
	MOUSE HEAVY CHAINS SUBGROUP I (A) 176 MOUSE HEAVY CHAINS SUBGROUP I (B) 181 MOUSE HEAVY CHAINS SUBGROUP II (A) 186 MOUSE HEAVY CHAINS SUBGROUP II (A) 182 MOUSE HEAVY CHAINS SUBGROUP II (A) 182 MOUSE HEAVY CHAINS SUBGROUP II (B) 200 MOUSE HEAVY CHAINS SUBGROUP II (C) 200 MOUSE HEAVY CHAINS SUBGROUP II (C) 204 MOUSE HEAVY CHAINS SUBGROUP II (C) 219 MOUSE HEAVY CHAINS SUBGROUP II (D) 222 MOUSE HEAVY CHAINS SUBGROUP V (A) 226 MOUSE HEAVY CHAINS SUBGROUP V (A) 226 MOUSE HEAVY CHAINS SUBGROUP V (A) 232 MOUSE HEAVY CHAINS SUBGROUP MISCELLANEOUS 232
	241 RAT HEAVY CHAINS
·	PAR BEIT HEAVY CHAINS 243 GUINEA PIG HEAVY CHAINS 248 CAT HEAVY CHAINS 250 DOG HEAVY CHAINS 252 CHICKEN HEAVY CHAINS 252 CHICKEN HEAVY CHAINS 254 SHARK HEAVY CHAINS 255 MISCELLANEOUS HEAVY CHAINS 259 T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (ALPHA CHAINS) 264
	T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (RETA CHAINS) T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (BETA CHAINS SUBGROUP I)
CONST	ANT REGION SEQUENCES
	KAPPA LIGHT CONSTANT CHAINS
	HEAVY CONSTANT CHAINS CH1 REGION
SEQUE	NCES OF RELATED PROTEINS
	1 CHAIN
	334 334
	MAJOR HISTOCOMPATIBILITY ANTIĞENS CLASS I N REGION
·	I REGION GENE PRODUCTS CLASS II A-CHAIN AT REGION
	TREGIUN GENE PRODUCTS CLASS II D'CHAIN D' NEGICIA

パ

	I REGION GENE PRODUCTS CLASS II B-CHAIN B2 REGION	5
	I REGION GENE PRODUCTS CLASS II B-CHAIN MEMBRANE BOUND REGION	17
	3	20
	HUMAN COMPLEMENT COMPONENTS	21
	T-LYMPHOCYTE RECEPTOR FOR ANTIGEN CONSTANT REGION	29
	T OF L CUREACE ANTIGENS	34
	MISCELLANEOUS PROTEINS OF THE IMMUNOGLOBULITS OF LIMITATION OF LIMITATIONO OF LIMI	
CODO	DNS OF NUCLEIC ACID SEQUENCES CODONS OF SIGNAL PEPTIDES OF KAPPA LIGHT CHAINS	52 58
	CODONS OF SIGNAL PEPTIDES OF HEAVY CHAINS. CODONS OF SIGNAL PEPTIDES OF BETA-2MICROGLOBULINS. CODONS OF SIGNAL PEPTIDES OF BAJOR HISTOCOMPATIBILITY ANTIGENS CLASS I. CODONS OF SIGNAL PEPTIDES OF I REGION ANTIGENS CLASS II. CODONS OF SIGNAL PEPTIDES OF COMPLEMENT. CODONS OF SIGNAL PEPTIDES OF COMPLEMENT. CODONS OF SIGNAL PEPTIDES OF T-LYMPHOCYTE RECEPTOR FOR ANTIGEN.	67 69 71 76 76 78
		508
	CODONS OF VARIABLE REGION HEAVY CHAINS CODONS OF KAPPA LIGHT CONSTANT CHAINS CODONS OF KAPPA LIGHT CONSTANT CHAINS CODONS OF LAMBDA LIGHT CONSTANT CHAINS	528 530
	CODONS OF HEAVY CONSTANT CHAINS CHI HEGION CODONS OF HEAVY CONSTANT CHAINS HINGE REGION CODONS OF HEAVY CONSTANT CHAINS CH2 REGION CODONS OF HEAVY CONSTANT CHAINS CH3 REGION CODONS OF HEAVY CONSTANT EXTRA LONG CH3 REGION CODONS OF HEAVY CONSTANT CHAINS CH4 REGION CODONS OF HEAVY CONSTANT CHAINS CH4 REGION CODONS OF HEAVY CONSTANT CHAINS CH4 AND MEMBRANE BOUND REGIONS	536 539 542 545 548 553
	CODONS OF BETA-2-MICHOGLOBULING	555
	CODONS OF MAJOR HISTOCOMPATIBILITY ANTIGENS CLASS I C2 REGION CODONS OF MAJOR HISTOCOMPATIBILITY ANTIGENS CLASS I MEMBRANE REGION CODONS OF MAJOR HISTOCOMPATIBILITY ANTIGENS CLASS I CYTOPLASMIC REGION	561 563 565
	CODONS OF I REGION GENE PRODUCTS CLASS II A-CHAIN & REGION. CODONS OF I REGION GENE PRODUCTS CLASS II A-CHAIN A2 REGION.	570
	CODONS OF I REGION GENE PRODUCTS CLASS II B-CHAIN B2 REGION	577
	CODONS OF HUMAN COMPLEMENT COMPONENTS	610
	CODONS OF T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (BETA CHAINS SUBGROUP I). CODONS OF T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (BETA CHAINS SUBGROUP II). CODONS OF T-LYMPHOCYTE RECEPTOR FOR ANTIGEN VARIABLE REGION (GAMMA CHAINS).	623 628 630
÷	CODONS OF T-LYMPHOCYTE RECEPTOR FOR ANTIGEN CONSTANT ALCION	. 637
	CODONS OF T-CELL SURFACE ANTIGENS CODONS OF MISCELLANEOUS PROTEINS OF THE IMMUNOGLOBULIN SUPERFAMILY CODONS OF MISCELLANEOUS PROTEINS ASSOCIATED WITH THE IMMUNE SYSTEM	. 641 . 644
	CODONS OF HIGELL SOLUTIONS OF THE IMMUNOGLOBULIN SUPERFAMILY CODONS OF MISCELLANEOUS PROTEINS ASSOCIATED WITH THE IMMUNE SYSTEM CODONS OF HUMAN D-MINIGENES	. 656
	CODONS OF HUMAN D-MINIGENES	663
	A REAL OF VEMANIT LYMPHOCYTE RECEPTOR FOR ANTIOCH O THE STORE	.004
	A THE AND A LANDRA LANDRES	. 000
	CODONS OF MOUSE KAPPA J-MINIGENES	
		. 671
	COCOUR OF BARRIT KAPPA J-MINIGENES	674
	CODONS OF HUMAN HEAVY JMINIGENES	675
	CODONS OF HUMAN T-LYMPHOCYTE RECEPTOR FOR ANTIGEN ALPHA CHAIN J-MINIGENES	680 683
	CODONS OF MOUSE T-LYMPHOCYTE RECEPTOR FOR ANTIGEN BETA CHAIN J-MINIGENES	687
	PSEUDOGENES OF IMMUNOGLOBULIN VARIABLE HEGIONS PSEUDOGENES OF T-LYMPHOCTYE RECEPTOR FOR ANTIGEN VARIABLE REGIONS PSEUDOGENES OF IMMUNOGLOBULIN CONSTANT REGIONS PSEUDOGENES OF OTHER PROTEINS	
S	UMMARY TABLES AMINO ACID DISTRIBUTION IN ALL VARIABLE REGION LIGHT CHAINS AMINO ACID DISTRIBUTION IN ALL VARIABLE REGION HEAVY CHAINS	716 718
	AMINO ACID DISTRIBUTION IN ALL VARIABLE REGION HEAVY CHAINS	
		730
v	JARIABILITY PLOTS LIGHT CHAINS HEAVY CHAINS	
	ACAVI CHAINS	
	LAMEDA LIGHT CHAINS	732
	HUMAN LIGHT CHAINS	

iv

I

296

1

	HUMAN KAPPA LIGHT CHAINS	733 733
	HUMAN KAPPA LIGHT CHAINS SUBGROUP I. HUMAN KAPPA LIGHT CHAINS SUBGROUP II. HUMAN KAPPA LIGHT CHAINS SUBGROUP III.	734
	HUMAN HEAVY CHAINS SUBGROUP I. HUMAN HEAVY CHAINS SUBGROUP II. HUMAN HEAVY CHAINS SUBGROUP III.	735
	HUMAN LAMBDA LIGHT CHAINS SUBGROUP I. HUMAN LAMBDA LIGHT CHAINS SUBGROUP II. HUMAN LAMBDA LIGHT CHAINS SUBGROUP II.	737
	MOUSE KAPPA LIGHT CHAINS	738
-	RABBIT KAPPA LIGHT CHAINS	738
	MOUSE KAPPA LIGHT CHAINS SUBGROUP II	739 739
	MOUSE HEAVY CHAINS SUBGROUP IA 7 MOUSE HEAVY CHAINS SUBGROUP IB 7 MOUSE HEAVY CHAINS SUBGROUP IIA 7 MOUSE HEAVY CHAINS SUBGROUP IIB 7 MOUSE HEAVY CHAINS SUBGROUP IIB 7 MOUSE HEAVY CHAINS SUBGROUP IIC 7 MOUSE HEAVY CHAINS SUBGROUP IIIA 7	740 741 741 741 742
	T-LYMPHOCYTE RECEPTOR FOR ANTIGEN ALPHA CHAINS	43 -
	CYTOCHROMES C	45
	HUMAN HLA DR BETA CHAIN CLASS II	
	MOUSE MHC IA ALPHA CHAIN CLASS II	
INDICES	S	
	NDEX OF PROTEINS	47 95

viii

considered uncertain by the authors have not been included in the table. In some instances the symbol # is used to indicate that several amino acid residues were found in one position, and these residues are listed in the notes. The four columns at the end of each table give:

- 1. the number of residues sequenced at that position,
- 2. the number of different amino acids found at that position,
- 3. the number of times the most common amino acid occurred and that amino acid in parentheses, and
- -4. the variability.

Variability is calculated (11) as:

Number of different amino acids occurring at a given position

Variability =

Frequency of the most common amino acid at that position

An invariant position would have a variability of one; if 20 amino acids occurred with equal frequency, the variability would be 20 divided by 0.05 equals 400. If, for example, four different amino acids Ser, Asp, Pro, and Thr occurred at a given position, and of 100 sequences available at that position, Ser occurred 80 times, the variability would be 4/0.8 = 5. When any of the amino acid residues sequenced were not identified completely and are listed as GIx (or Asx), two values, separated by a colon, are given in the last three columns. The first value in each of these columns is calculated assuming that only one of the two possibilities, e.g., Glu or Gln (or Asp or Asn) occurred, while the second considers that both were present and maximizes variability. In the variability plots, the horizontal bars indicate the two values.

When two or more amino acids are most common and occur with equal frequency, they are tabulated as a note, and the symbol + is used in the next to last column. If no sequence data have been reported for any position, there are no entries in the last four columns. Variability is not calculated for insertions or if only a single sequence is known. When the translated sequence of a clone corresponds to a previously listed sequence of a plasmacytoma from which it was prepared, only one sequence is listed so that the variability computations are not affected, and a note is included.

If a given sequence is associated with any antibody activity, this is indicated by an asterisk alongside the protein heading, and the antibody specificities are given in a separate list with binding constants if available. The notes list the a-allotypes for the rabbit heavy chain V-region and the b-allotypes for the constant domain of the rabbit kappa light chain. A key reference to the sequence is given; generally the most recent reference since it is usually the most nearly complete, but often several references are included, especially when revisions of a sequence have been made. Notes are now of two types; general notes about a table indicated by the symbol #, and specific notes indicated by the sequence number.

Signal Sequences

The signal (precursor) amino acid sequences of immunoglobulin chains are listed in three tables: one for kappa light chains, one for lambda light chains, and one for heavy chains. They were obtained either by direct sequencing of signal proteins (12-14) or by translating nucleotide sequences from DNA clones. Signal segments range from 17-29 amino acid residues in length and are thus numbered from -29 to -1. Genomic DNA clones contain introns of varying length that interrupt the coding sequence of the precursor within the codon for position -4, and in rare cases for position -6. Thus, the L-gene encodes the leader peptide to position -4 and the 5' end of the V-gene codes for positions -4 to -1.

The signal amino acid sequences of the T-cell receptors for antigens, β_2 -microglobulins, major histocompatibility complex proteins, and complement components are listed in separate tables.

Leu (MOPC321, MOPC63) exists in an alpha helical conformation, terminated by chain reversal conformations in the four C-terminal residues Trp-Val-Pro-Gly; the four amino terminal residues are compatible with the alpha helix (15).

Variable Region Sequences

The variable regions (16) of immunoglobulins have been shown to contain hypervariable segments in their light (11, 17-23) and heavy (22, 24-27) chains, of which certain residues have been affinity labeled (28-41). Three hypervariable segments of light chain were delineated from a statistical examination

of sequences of human V_x , human V_λ , and mouse V_x light chains aligned for maximum homology (11,22). These and the three corresponding segments of the heavy chains (22,26,27) were hypothesized (11,22) to be the complementarity-determining regions or segments (CDR) containing the residues which make contact with various antigenic determinants, and this has been verified by X-ray diffraction studies at high resolution (42-67). The rest of the V-region constitutes the framework (11,22,66-68). It is convenient to identify the framework segments (FR1, FR2, FR3, and FR4) and the complementarity-determining segments (CDR1, CDR2, and CDR3) with the three CDRs separating the four FRs. The residue numbers for these segments are as follows:

Segment	Light Chain	Heavy Chain
FR1	1-23 (with an occasional residue at 0, and a deletion at 10 in V_{λ} chains)	1-30 (with an occasional residue at 0)
CDR1	24-34 (with possible insertions numbered as 27A,B,C,D,E,F)	31-35 (with possible insertions numbered as 35A,B)
FR2	35-49	36-49
CDR2	50-56	50-65 (with possible insertions numbered as 52A,B,C) ^a
FR3	57-88	66-94 (with possible insertions numbered as 82A,B,C)
CDR3	89-97 (with possible insertions numbered as 95A,B,C,D,E,F)	95-102 (with possible insertions numbered as 100A,B,C,D,E,F,G,H,I,J,K)
FR4	98-107 (with a possible insertion numbered as 106A)	103-113

^a In the rabbit, Mage et al. (69) consider position 65 in V_H to be in FR3, since it is allotype related.

In the tables of V-regions, the FR and CDR are separated by horizontal lines for convenience in reading. One mouse kappa light chain, MPC11, has an extra segment of 12 amino acid residues between position 1 and the signal sequence (70). Several chains have internal deletions.

In the tables, the V-genes for the light chains code to amino acid position 95, and the J-minigenes from position 97 to 107 for lambda and 108 for kappa light chains. Position 96 is usually the site of V-J joining by recombination and may be coded partly by the V-gene and partly by the J-minigene. Because the site of V-J recombination could occur at different positions within a codon, different amino acid residues may result at this position. We have changed the location of the inserted residues from 97A-F (2) to 95A-F, since it makes for better alignment by confining chains of different lengths to the V-gene region. In V_x chains, J1 and J2 were used 5 to 10 times more frequently than J4 and J5 (71).

The V-genes for the heavy chains code up to amino acid position 94 and are followed by the Dand J-minigenes. Because of the extensive variation in the lengths of D-minigenes, the exact boundary between D and J is not always located at the same amino acid position. In addition, the lengths of the J encoded amino acid sequences vary by a few amino acid residues. Moreover, the process of D-J joining appears to involve insertions of extra nucleotides between V and D and between D and J, termed the N region (72-76) and correlates with the appearance of terminal deoxytransferase in B cells (75). The original numbering system for the heavy chains has therefore been retained. Wysocki *et al.* (76) have provided some evidence suggesting a non-random origin for the V_H-D_H junction, perhaps a minigene, rather than random addition of the N nucleotides.

It has become evident that a critical understanding of the architecture of antibody combining sites and the genetics of the generation of diversity and of antibody complementarity will depend to a great extent on the evaluation of a large number of sequences of the variable regions and especially of the complementarity-determining segments of light and heavy chains of immunoglobulins of different species. Ability to locate residues in the site making contact with antigenic determinants (77) and to predict (67,78-82) the structures of antibody combining sites will depend heavily upon such sequences.

Figures 1 and 2 are stereoviews of the α -carbon skeletons of the four Fv regions for which high resolution X-ray structures have been determined, NEWM (44), KOL (62), MCPC603 (47, 48, 63), and J539 (64). The residues in the CDRs are shown as solid circles. In Fig. 1 the combining site is at the

	нυ	MAN	KAPPA LIGHT	CHAI	vs s u	BGRO	UPI																			
	•			, POA	2 AU	3 REI	HÂU	нк101 °С∟ #	รตั้พ	7 AG #	WEA	9 HK 137 'CL #	10 HK 134 CL	DAUDI CL	12 WALKER CL	13 HF3- 16/6	14 HF2- 1/13B	HF2-	HF2-	17 BJ 26	18 RFZ	19 PSM	ном 20	21 ESM IGG	22 ESM IGM	23 WAT
		0 1 2 3			ILE	ASP ILE GLN	ILE	ASP ILE GLN	IL E	GLN	ASP ILE GLN	ILE	ASP ILE	ASP ILE GLN	ASP ILE GLN	ASP ILE GLN	N.E	ASP ILE GLN	ILE GLN	GLN	ASP ILE GLN	ILE GLN	ILE. GLN	ILE GLN	GLN	ASP ILE GLN
		4 50	THR(.98)	MET	MET	MET	MET THR	мет тна	мет тня	THR	THR	MET THR	THB	THB	тня	THR	тня		THR	THR	тна	THR	THR	THR	THA	
		7 8 9-	SER(.95) PRO(.98)	SEA PRO	SER PRO	SEA PRO	SER PRO	SER PRO	SER PRO	SEM	SER	SER PRO	SER	PRO	SER	PRO	SER	SER	SER PRO	SER PRO	PRO SER	SER PRO	SER PRO SER	PRO	PRO SER	SEF PRC SEP
	F	11	SEB(96)	LEU	SEA LEU	SER LEU SER	SER LEU	SER LEU	SER LEU SER	SER LEU SER	SER	LEU SER	SEA LEU SER	SER	SEA LEU SEA	SER	SEA LEU SEA	SER LEU SER	LEU SER	LEU SER	SER LEU SER	SER	SER LEU SER	SER LEU SER	SER	SEF LEL SEF
	1	13 14	-	ALA SER	ALA SER	ALA SEA	ALA SER	ALA SER	SER	ALA SER VAL	SER	ALA SER	SER	ALA SER	SER		ALA SER	SER	SER VAL	SEF						
		16		GLY ASP ARG	ASP	GLY ASP ARG	GLY ASP ARG	ARG	GLY ASP ARG	GLY ASP ARG	ARG	ARG	GLY ASP ARG	ASP ARG	ASP	GLY ASP ARG	GLÝ ASP ARG	GLY ASP ARG	GLY ASP ARG	ASP	GLY ASP ARG	ASP	ASP	ARG	ASP ARG	GL1 ASF ARC
B CYA		20 21	THR(.96)	THR	THR	THR	тня	THR		THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THB	THR	тня		THA ILE	THF
		22 23		CYS		CYS	CYS	CYS	CYS	CYS	CYS	ÇYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	CYS	GLX
3550		25 26 27	ALA(.95)	ALA SER GLN	ALA SER	ALA SER	ALA SER GLN	ALA ARG GLN	ALA SER GLN	ALA SER GLN	ALA SER GLN	ALA SER GLN	GLN	ALA GLY HIS	SER GLN	SER GLN	ALA SER GLN	ALA SER GLN		ALA SER GLN	ALA ASN GLN		HIS	GLX	SER	ALA SEF GLN
	r	27B 27C				 			•••							 										
Image: Section of the sectio	C D A 1	27E 27F																								
33 LEU LE		29 30		ILE	ILE	ILE ILE	ILE	ILE	ILE	ILE ASN	ILE ARG	11 5	N.E SER	ILE THR	ILE SER	ARG	ILE ARG	ILE ABG	ARG	ILE ASN	ILE SER	ILE ARG	ILE	ILE SER SER	ILE SER SER	ILE
3 The		32 33		PHELEU	LEU	LÉU	LEU	LEU	LEU		LEU	LEO	LEU	PHE LEU	LEU	ASP LEU	LEU		LEU	ALA	LEU	TYR LEU	LEU	LEU	LEU	TYR VAL ASN
38 PHRO(199) LYG		35 36	тяр	TRP	TBP	TAP	TAP	THP TYR	TAP	TAP TYR GLN	TRP TYR	TRP	TAP TYR	TRP TYR	TAP	TRP	THP	TAP	TRP TYR	TAP	TRP	TRP TYR GLN	TRP PHE GLN	TRP	TRP TYR	TRP
43 PhO ALA AL		38 39		GLN		GLN	GLN	GLN	GLN		GLN	GLN	GLN	GLN LYS	GLN LYS	GLN			GLN			LYS	LYS			
44 PRO	FR	41 42	FR0(.95)	LYS	GLY LYS	LYS	LYS	GLU LYS	LYS	LÝS	GLY THR ALA	LYS	GLU LYS	GLY LYS	GLY LYS					LYS	ALA	LYS	GLY			
49 TYM		45 46		LYS	PRO LYS LEU	PRO LYS LEU	PRO	PBO	PBO	PRO LYS	LYS ARG	PRO LYS SER	PRO LYS SER	PRO THA LEU	PRO LYS LEU					PRO LYS LEU	PRO LYS LEU	LYS				
53 LEU(3) LYS ASN SER SER <td< td=""><td></td><td>48 49</td><td>LEU(.98) ILE(.98)</td><td>ILE.</td><td>ILE</td><td>nε</td><td>TYR</td><td></td><td>TYA</td><td>ILEU ILE TYR</td><td>HE</td><td>TYA</td><td>TYR</td><td>TYR</td><td>TYR</td><td></td><td></td><td></td><td></td><td></td><td>TYR</td><td>ILE TYR</td><td></td><td></td><td></td><td></td></td<>		48 49	LEU(.98) ILE(.98)	ILE.	ILE	nε	TYR		TYA	ILEU ILE TYR	HE	TYA	TYR	TYR	TYR						TYR	ILE TYR				
St LEU(189) LEU (180) LEU (180) <thleu (180)<="" th=""> <thleu (180<="" td=""><td></td><td>50 51 52</td><td></td><td>ALA SER</td><td>ASP ALA SER</td><td>SER</td><td>ALA ALA SER</td><td>ALA ALA SER</td><td>GLY ALA SER</td><td>SER</td><td>ALA THR</td><td>ALA ALA SER</td><td>SER</td><td>SER</td><td>ALA SER</td><td></td><td></td><td></td><td></td><td>ASP</td><td>ALA ALA</td><td>ALA</td><td></td><td></td><td></td><td></td></thleu></thleu>		50 51 52		ALA SER	ASP ALA SER	SER	ALA ALA SER	ALA ALA SER	GLY ALA SER	SER	ALA THR	ALA ALA SER	SER	SER	ALA SER					ASP	ALA ALA	ALA				
9: GLY	8 2	54 55	LEU(.98)	LEU GLU		LEU GLN	LEU PRO	LEU GLN	LEU GLU		LEU GLN	1 611	LEU GLN	LEU GLN	LEU GLN					LEU GLU	LEU	LEU GLU				
600 SER S			GLY VAL(.98)	GLY	GLY VAL		GLY VAL	GLY VAL	GLY VAL	GLY VAL		GLY VAL	GLY VAL	GLY VAL	GLY VAL					GLY ASX	VAL					
63 GLV GLM GLY GLM		60 61	ARG(.95)	SER ARG	SER	SER	ARG	PRO SER ARG	PRO SER ARG	PRO SER ARG	PRO SER ARG	PRO SER ARG	ARG	SER ARG	SER ARG					SER	ARG					
669 THR ALA THR ALA THR THR THR THR THR 71 ADP THR THR THR THR THR THR THR 73 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 76 THR THR THR THR THR THR THR THR THR 76 LEU LEU LEU LEU LEU LEU LEU LEU LEU 77 GUV SER SER SER SER SER SER SER 78 GUU GUU GUU GUU GUU GUU GUU GUU 80 ALA(39) ALA ALA ALA ALA ALA ALA ALA 81 ALA ALA ALA ALA ALA ALA ALA A		63 64		SER GLY	SER GLY	SER GLY	SER	SER	SEH	SER	SER	SER	SER	SER	SER GLY					SEA GLY	тня					
669 THR ALA THR ALA THR THR THR THR THR 71 ADP THR THR THR THR THR THR THR 73 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 74 THR THR THR THR THR THR THR THR 76 THR THR THR THR THR THR THR THR THR 76 LEU LEU LEU LEU LEU LEU LEU LEU LEU 77 GUV SER SER SER SER SER SER SER 78 GUU GUU GUU GUU GUU GUU GUU GUU 80 ALA(39) ALA ALA ALA ALA ALA ALA ALA 81 ALA ALA ALA ALA ALA ALA ALA A		66 67	GLY(.95)	GLY. SER	GLY GLY SER	GLY	SEA SEA GLY	GLY SER	GLY SER GLY	PHE	SER	SER	SER	GLY SER	GLY SER					GLY SER	GLY					
72 THA		69 70		THR		THR	THR	THR	THB	тна	THR	THR	THR	ALA GLU	THR ASP					THR						
76 LLE (.95) LLE GLX 76 SER <	- 			THR PHE	THR PHE	PHE	THR LEU	THA LEU	LEU	PHE	LEU	LEU	LEU	LEU	LEU					VAL						
79 GLN		75	ILE(.95) SEA(.95)	北臣.	ILE SER	ILE SER	H.E	ILE	ILE SEA THA	ILE SER GLY	ILE ASN SER	ILE SEA SEA	ILE SEA SEA	ILE SER SER	ILE SER SER					GLX SER SER						
81 GLU		78		GLN	GLN	GLN	GLN	GLN PBO	LEU GLN PRO	GLN PRO	GLN	GLN PRO	GLN PRO	GLN PRO	GLN PRO											
B4 ALA(39) ALA ALA <t< td=""><td></td><td>82</td><td></td><td>GLU ASP</td><td>GLU ASP</td><td>ASP ILE</td><td>GLU ASP PHE</td><td>ASP PHE</td><td>GLU ASP ILE</td><td>ASP ILE</td><td>GLU ASP PHE</td><td>GLU ASP PHE</td><td>ASP</td><td>GLU ASP PHE</td><td>ASP SER</td><td></td><td>•</td><td></td><td></td><td>ILE</td><td></td><td></td><td></td><td></td><td></td><td></td></t<>		82		GLU ASP	GLU ASP	ASP ILE	GLU ASP PHE	ASP PHE	GLU ASP ILE	ASP ILE	GLU ASP PHE	GLU ASP PHE	ASP	GLU ASP PHE	ASP SER		•			ILE						
89 GLN GL		85 86		ALA THR TYR		TYR	A1 A	ALA THR TYR	GLY ASN TYR TYR					ALA THR TYR TYR	IYH					ALA PRO LYS						
94 LEU LEU LEU THR TYR VAL LEU PRO PR		88	CYS	GLN GLN	GLN GLN	GLN	GLN GLN	GLN - GLN	GLN GLN		LEU GLN	GLN GLN	GLN GLN	GLN GLN	CYS GLN			·		GLN						
94 LEU LEU LEU THR TYR VAL LEU PRO PR		91 92		ASP ASN	ASP TYR	GLN	11,6	TYR ASN SER	TYR ASP ASN	TYR ASP THR	SER SER	SER	ASN SEA	ASN TYR ASN	SER					ASP ASP						
95C		95 95A	,			PRO		PRO #	PRO		PHE	PRO		SER	LEU					PRO						
95F	7	95C 95D						···-												 						
90 PHE PHE PHE PHE PHE LEU 99 GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY		95F 96				TYR			H.E	ARG	TBP			PHE	ILE					түн				•		
102 THR(.57) THR T		98 99	GLY	PHE	PHE	PHE	PHE		PHE	PHE GLY GLN	PHE			PHE GLY GLY	PHE GLY GLN					LEU GLY GLY						
104 VAL VAL LEU VAL VAL 105 ASP GLU GLU GLU GLU 106 PHE ILE ILE MET 106A ILE ILE ILE MET 106A ILE ILE ILE ILE 107 LYS LYS LYS LYS	-	101		GLY THR	GLY THA	тня	GLY THR		THR	LYS	THR	•••		GLY THR	GLY THR ARG					GLY THR						
1054		105		VAL ASP PHE	VAL GLU ILE	GLN	GLU		VAL GLU ASN	LEU GLU ILE	VAL GLU			VAL ASP ASN	LEU GLU ILE					VAL GLU MET						
		107		LYS	LYS	 тня	LYS		LYS .	LYS		•••		LYS	LYS											


Ġ.

~															•			-							
HUM -	AN KA	APPA LIGH			27 CAR A	28 TEI	(cont'd 29 BJ 48) 30 CON	31 TRA	32 AMYLOID LEP	33 F- GUI #	34 OU (IOC)	35 DEE	38 GAL (I)	37 JOH	38 KER	39 LAY	40 BRA	41 WES	42 Vb CL	43 Vb' 'CL	44 HK102 'CL	45 EU	45 DEN	47 PAU
F R 1	0 1 2 3 4 5 6 7 8 9 101123 4 5 6 7 8 9 101123 1 4 15 6 7 1 1 1 1 2 0 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	ALLER AND A CONTRACT OF A SUBARA AND A CONTRACT OF A CONTR	PERT RAROA AURAR LYPGL ALGE TILERE SELENAR LYPGLA RELEAS VGAAV TILER	ALLAN TOSPAS SUBAS VOLANA HLA	THA ILE THA	ALLON TOSPS SUSAR ALYPOL REAL	SENT ANAGA AURAA LYPGL ALLAN TOSPS SUSAS VGAAV TILA	ALLIGN THESE SUBSES VOLSAN THER	ALLEN HARAGE RURAA LYPGAL ALLEN TOSPES SLEADE VGAAV THIT	ALLAN THURADA AURAA AUPAGL A	ALLENT RARAR RURAR LYXGL REARS	AUGM THERE AURAR LYXGL RERS	ALLAN THARAR AURAR LYPOL RERS	ALLAN TOSAS SUSAS VOAAV TILLY	ALLIE HNAOR RUAAR LYPGL ARE TO SELENAR LLEAR ALLEAR	ALLEN TANACA RURAA LYPG HEAS	ASENT RNAOR RURIR LYPGL REAV TILTYS	ALLAN TOSPES SUSTER ALYS	ALLENT ANAROR RIACAR LYPOL RUNTGSPRS SVELAR LYPOL LYPOL RUNTGSPRS SVELAR LYPOL RUNTGSPRS SV	ALLGN TOSPS SUSAS VOASAAV TILTY	ALGM TOSPS SUSAS VOARA HITTO	*P AILENT GLET THR GLEAR THR SEC PROR LEER ALAR LEER ALAR VGLSP CARA V THLER AAROL THLER AAROL THLER CARA VGLSP VTHLER CARA VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VGLSP VTHLER VTHLER VGLSP VTHLER VGLSP VTHLER VTHLER VGLSP VTHLER VTHLER VGLSP VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VTHLER VT	AILGNE ANAROA TURAR LYPGL AR LERS	ALLON THORAGE FURAR LYPGL REAS	ALLON THURDE FURAS AVANA HILTO
СОЯ	223 224 25 26 277 278 277 277 277 277 277 277 277 277	GLN GLN ASER GLN ASE GLY TYRU LEU	ALA ALA SER GLN ASX SER ILE SER	<u> </u>	<u></u> <u>CYS</u>	CYS	CYS				ALA	ARG ALA SER GLX THR ERR SERR LER SERR LEU ASX	ARG ALA GLY GLN SER VAL ASN LYSA LEU ASN	ARG ALA SERN GLI IIIIIGLY GLY ARGNP ASSNP LE GASNP LE ARSNP LE THR		GLN ALA SEA GLN IIIII ASP ILE LYS	GLN ALA SER GU: ASN ASN ALA AVR LESN LESN TRP		ARG AREN I IIII SPERSPUA	ARG ALA SER GLI I I I I I I I I I I I I I I I I I I	ARG ALAR ALAR ALAR ALAR ALAR ALAR ALAR A	ARG ALAR SELN SER ILER SER LEU ALA TRYR	ARG ALARN ALARN SEL IIIIR SILSNAPU ALARN ALARN	SER	ILE SER SER LEU ALA
	35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50	Ϋ́́́					ALA PRO LYS GLY ASP ALA					TAP TYAX GGLX SOOY GLYSA OX ASXU LEU ILE TYAAAAF ALAA SE	LYS PRO GLY LYSA PRO LYSA LYSA LEU	PAC LYS GLU LEU ILE TYF		-	TYRNN GLNS GLYS PROSULALA PROSULEU LEUULE TYP ALA SEF		TYA GLN GLN SGLN SGLN SGLN SGLN SGLN SC SC SC SC SC SC SC SC SC SC SC SC SC	TYA GLN G LYSO GLYS GLYS ALA PROS LEU LEU LEU LEU LEU	TYR GLN GLN SPRO GLYS ALA PROS LEU LEU ILE TYR ALA	GLNN SOULSA PRUSA PRUSULLER TASA	TYRNN CARACTER STATES	GUN GUN CUS GLY GLU GLU GLU CUS CUS CUS CUS CUS CUS CUS CU	GLX GLX PRO GLX ALA PRO GLX ALA PRO LEU ILE TYA GLX
CDR №	51 52 53 54 55 56						SEP SEP LEU ASX SEF	L		-		ASX LEU HIS SEF	LEL LYS SEF	LEL GLN SEF	 				SER LEU GLU ASN	GUN	LEU GLN SER	SER LEU GLU SER GLY	SEF LEL GLU SEF	J LEL J GLU J SEF GLY	, ,
F R 3	578 996123 456677123 456778 901223 4888 8456777778 901223 48567888 88878						GLV VAL SEFE GLV SEFE SEFE SEFE SEFE SEFE SEFE SEFE SEF					GVA PERTER VELEC HOMENEN FEELU DECLET UNIVEL	V PSERHEF LEPLY HITHER HERLELERALSH LTHITYY VPSERHEF LEPLY HITHER HERLELERALSH LTHITYY	DAGEN YRYRY ROERU RIRYU JOUDE ARARS	- OFFER LAND - CARLER AVAN ALEGA		V REALEST TAPHTH HEADER COROLS AND A CERCERS COSO TAPHTH HEADER COROLS ATTYING A CERCERS AND A CERCERS	- DAGEN VAVAY ADERE ALLARD NOUP - ARARS	GV PSARS GSGSG TOPTIL THSSI GPGAR ATTROGG	PRC SEF SEF SEF SEF SEF SEF SEF SEF SEF SEF	PERCENT AFREE HALLS ALLS ALLS ALLS ALLS ALLS ALLS ALL	VALORGERYRYRYRUERU RUERU KOPPE AFTER VREARHER YRYRYRUERU RUERU KOPPE AFTER VREARHER YRYRYRUERU RUERU ACHYFRY VREARHER YRYRYRU RUERU RUERU ACHYFRY VREARHER YRYRU RUERU RUERU ACHYFRY	SEI GLI PRI ASI PRI ASI PRI ASI PRI TY TY CY	DAGUE YAYAY AUERU AMARAU NOPPE AAARAS	ערשויית אלאאל ערשיים ביינערן צבטהה הענגנט
C D R 3	89 90 91 92 93 95 95 95 95 95 95 95 95 95 95 95 97 98			`			GLUSEI SEI ASI ARH PH						SEYTH HA	R SE YYO PR	R RO	TY AS A S PA PA PA TH GL	HPP UO	PO	GLL GLL HIS SEF VAL PR L L L L L L L L L L L L		A ALA N ASI A SEI PHI D PAC	A TYPH A ASH A SEF TYPO SEF 	LY ME GL	H PHA PA	
Р. Я. 4	98 99 100 101 102 103 104 105 106 106 107 108 109	<u>,</u>				-	GL GL TH LYY GL VA LY	YNYR SLUL					X PR GLYA R GUXE ME S TH	OY GLU AR TH SLU GL IL SLU GL	NYA SLUE	GL GL TH LY VA AS LE LY	O GL Y GL H TH S LY GL U VA			Y Y R L S				(S L) (L LE (U GL (L IL (L IL (S L)	R U U E 'S

н: -	JMAN H	(APPA 48 HBJ 4	49	50		52 MON			55 S- GUI	56 AMYLOID BAN	57 BJ 19	58 BEL	59 JBL	60 PAP	61 CAR	62 MEV	63 BI	64 AMYLOID ES305	65 CRA	66' DAV	67. FIN	68 Ka	69 Vd 'CL	70 LUX	71 NE	72 Va CL
F R I	01234 56789 011234 156789 011234 156789 01223	SPENT ANAROA TURAR LYPGL ALERAS VGAARAV TULERAS VGAARAV TULERAS VGAARAV TULERS	ASLENT RNROR LURAR LYPGL RERS	ASLEAT ANAROA HURAA LYPGI ASLEAT ANAROA HURAA LYPGI ALARAV HURAAV HURAAV HURAAV HURAY	ALGA TIGSPAS TURAR LYPGL REAS ILSAS VGASAL REAS VGASAL TILLSAS VGASAL TILLTCY	ANLENT ANAROR TURAR LYPG	ASULLE RURAOR FURA SULLE HLEROR FURAS VLLAR LYPO	ASLENT RNARO & RURAR LYPO ASLENT RNARO & SUBARA LYPO ASLENT RNARO & SUBARA VILLAN ASLENT RNARO & SUBARA VILLAN ASLENT RNARO & SUBARA SUBARA VILLAN	ALGA THANAOR AURAA LYXGL AEAAS	ALLG HANAGAR AUPAGA ALAS	* ALLGI HARROR RURAR LYAGL RURAS ALLGI HARROR RURAR LYAGL RURAS VGAAV HILTY	ALLAN ANHOR RURAR LYPGA REAS	ALLEN ANAROR RURA VALYPGA AN THE BY	ASLENT RORA RUA ALYPO	ALLAN HURACE LUSAS VGAAV SILLY	A VALUE ANAROR AUARA LAVARA A LEAR SALAR ALAVARA A LEAR ALAVARA ALAVA ALAVA ALAVARA ALAVARA ALAVA ALAVARA ALAVARA ALAVARA ALAVARA ALAVAR	ALLANT RANGOR OURAR LYPILANT GERAR LYPILANT READ	* ALLAND TGLENURANDER RUDRA SELAR LURANDES SELAR SLASSANDER LIR TGLENRAS VGA VIH TCY	ALLANT ANTRON AUTAAN UCAS	ALLENT ANAROR RUA ALLENT ANAROR RUA SERENT ALLEN VGARAL REAS	ALLAN TOSPAS SUS IN VOAAN TILTY	ALLGM TOSPES HEELAR LYPGL ALLAN TOSPES HEELAR VGAAV TILTY	ALLEN RARA BURAR LYPGA REAS	ALLU RNACOR OURAR LYPGLU AND LURAR LYPGLE AND LEAR VILLER	ALLANDA RUAAR LUAAR LAXAA HEAS	INTER ANT AND
CDR 1	24 25 26 27 27A 27B 27CD 27C 27F 28 29 30 31 32 33 34	ARG ALA SELN G SERN SERN SERN ASSNP LEU ALA	ARG ALA SERX ASX ASXA ASXA LEU ALA	ARG ALARG	ARLA ALA GLN IIIIII SEA SEA SEA SEA				ARG ALA	ARG ALA SER GLN SER VAL TYR VAL ALA	GLARN SGLI SPENREUN	GLX ALA SER GLX IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	ARG ALAR SERN SELER SELER THREU		ARG ALARN	ARG ALA SER GLI SER SER VAL SER VAL ASP LEU ASN	GLARN	GLN SER GLU ALA * ASN TYR LEU		ASP ALARN SGLN ASX SERP LEU LEU	ASP ALARN SELN 	GLU ALARN SERN THR LEURR LEUR ASN	ARG ALA SER GLN GLY ILER SER LEU ALA		GLX ALA SER GLX ASX ILE SER ASX LEU ASX	ARG ALA SER GLI ILEAR ALA SER LEAR ALA LEU ALA
FR2 CD	35 36 37 38 39 40 41 42 43 44 45 46 47 46 47 46 47 46 50 51 52 53	TRP TYLN GGLUS PROY LYSO GGLA ROS LEU LEE SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA SCA LYSA LYSA SCA LYSA LYSA LYSA LYSA LYSA LYSA LYSA LYS	TRP TYR GLX GLX LYS PRO	TRP TYR GLN GLY ARG PRO						THP PHE GLN LYS PRO GLY LYS ALA PRO LYS LEU LEU LEU TYR ASP ALA SER SER	TTGLA PRYSA OSUULE PRYSA PRYSUL PRYSA PRYSA PRYSA PRYSA PRYSA PRYSE PRYSA PRYSE PRYSA PRYSE PRYSA PRYS	GLY PRO LEU LEU ASP ALAR THR	THP TYR GLN LYS GLN LYS ALA PRO LYS ALA PRO LYS LEU TYR		TAP TYLN SOULSA PRUSA PRUSA VEL R LYSER SSE	TRP TYPN GLN LYPN GLN LYPN GLN LYPN LYPN LYPN LYPN LEU LELE PHE ASP THRA ASN	TRP TYRN GLN LYSO GLN LYSO CLYS ALA PROS PHE LEU TYR ASP ALA GLN			TRP TYR GLN GLN TYR PRO	TAP TYA GLN GLN TYA PRO	TRANSON SOLVER RALARD	TAPA TYRN GGLYSOY LYRN LYRN LYRN LYRN LYRN LYRN LYRN LYR		TRP TYR GLX LYS PROY GLX ALA PRO LYS VAL MET ILE TYR	TYTUN SOYSA OSUUL RIPARA
CDR2	54 56 56 57 58 58 58 58 58 58 58 58 58 58	SEU GLUG ARU VAL OSER SELY VAL SEC SELY		1						LEUNA GSELY VA ORANEH YRANA GSELSEL HAPHEU HAERRU NOUDPE ANARTH GSELSEL HAPHEU HAERRU NOUDPE ANARTH LU THLERRU NOUDPE ANARTHYRS	SEE LIGUR LIGUR VAL XR GSERGE GSERGELGUR SASXES SASXES TLYS	LUYAR GUL GOSELYAR GOSELYAR GUL AND THIS SLE LOOK AND THIS SARHER AND THIS SLE LOOK AND AND THIS SLE LOOK AND ANTYTYYY			SE LULR VARABLE VA	LEUNER UN DESARTER YYGERY HAPPERU HAPP	LEULE IS A SUBSECT AND A SUBSE	,		•		GV PSAPS GGGSG TAPTP TILSSV GPGAP ATTTC	THEURERY APPER GERYRY THE THE THE APPER GERY APPER GERY APPER GERY APPER GERY APPER			S LOS GV REARS GSGSG TARTL TISSL GRGAR ATTYC
CDR3	89 991 992 934 955 955 955 955 955 955 955 955 97 98 99 99 99 99 99									GLNANASE RO	PHE ASP LEU PRO TYR THR PHE	GLNRXANS EO			ASN THR PHE	GLN GLN SER THR ASN GLU VAL THR PHE	GLN GLN TYR ASN LEU PRO TYR TYR THR PHE					ARG THR PHE	GLN GLN LEU ASN SER TYR PRO			GLNE GLNE AS TYRO
F R 4	99 100 101 102 103 104 105 106 106 106 107 108 109									GLY GLN THR LYS JLE LYS ARG	GLY PRO GLY THR LYS VAL GLU LEU	GLY GLY GLY THR GLU VAL GLU VAL LYS		•	PRO GLY THA LYS VAL ASP ILE LYS	GLY GLY THR THR VAL ASP ILE LYS ARG	GLY GLN GLY THR LYS GLU ILE LYS ARG THR					GLY GLN GLY THR VAL ASP L				

ни	MAN K	АРРА 73 N1	UGHT 74 PW	CHAINS S	UBGRO 76 ALE	77 5HE	(cont'd 78 ADA) 79 KUE	80 GO	81 BOL	92 Ri	83 Ve CL	84 0CO	85 V13 'CL	86 V18A 'CL	87 V19A CL	88 V 198 'CL	89 V188 CL	90 HF6- 21/28	91 SAC	92- WAG	93 нвэ 1	94 AMYLOID 547	95 WEB	96 HOE
F A ۱	0 1 2 3 4 5 6 7 8 9 11 12 3 4 5 6 7 8 9 11 12 3 4 5 6 7 8 9 0 11 2 3 4 5 6 7 8 9 0 11 2 3 4 5 6 7 8 9 0 11 12 3 4 5 6 7 11 11 11 11 11 11 11 11 11 11 11 11 1	ALLE RNROR AURA, LYPGAL RUSA SLUSA, VGAAV TIELY	A LAND TRANCH RURAR LYPGL AND TYPE	ASP ILEN MET A GLNA BIB SERU SERA SERU SERA SER VGLY ARGL IIE CYS	ASLEXT RARAS SALAR LYPEGA ARA RUARS SALAR LYPEGA ARA RUARS SALAR LYPEGA ARA RUARS VIENTES	ASERAD RARON RURAR LYXGL RORNS ASERAD RURAR LYXGL RORNS AS VGARA HARYS	ALLAN THURPOR AUAN VALAPOL AND ALLAN THURPOR AUAN VALAPOL AND VALAPOL AND	ALLAN TOSPES TOPAS VELADA THITO A	ASALE HARAOR ALAR LYXAL HERSIG	SPEN HELLAR ALLYXGL REAS	PARENT ANAROR TURAR LYSING	ALLAN THERE RABRAR TYPESA REAS	ASELXT RXROR RUN OF ALXAND AND AND AND AND AND AND AND AND AND	INTERPOR SURAAR TYPE ANALAS IS ANALA	A STOR AND A START A START A STORE A START A S	alin alu AN THE SEA ALA VE GUY THE ASS SAALA VE GUN	Lailept AN Ca Rais A a VO stric AR Co Ca Svota A VO stric AR CO CA	aiginalu RRN HLU THLUR protection of the second sec	ASA val MET THAN SEAO SEEO SEEO SEEO SEEO SEEO SEEO SEEO	ALLUE HNAOR RURAR LYP	ALLAND RADAR AUGAS	ASP ALLOT RANGOUNT RURAN THURADA SULAR LUSA ALLAN VGLAN	ASE GLAT THAN SEROR SEROR SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEEDRA SEED	ASP ILENT THEROFE ERI GERAR SERI SERAR SERAR VAL	ASPE ILLN THN SPROA SEA LEERA SEA SALER VAL
CDR1	24 25 26 27 27 27 27 27 27 27 27 27 27 27 27 27	GLU ALAR SELR VAL LEUU SE LYN SE LYN THRE ALA	ALA SER GL	ILE PBO	ARG	GLX ALA SEAX GLI I SEAX ASX SEAX ASX LEAX ASX LEAX ASX	ALA SER GLX IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	ALER LER LER LASE	ALA SER	ALA SER GLX 	ALA SER GLI IIIIISER SER	ALA SERN IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	ALA SER GL: 	ILE SER SER TYR LEU ALA	SERANAL TY ERNNAL TY ERNNAL TY ERNNAL LE EN	ALARU	ALA SERU GUIIIIISE SERPU SERPU LALA	ALERNAL	SELAU LAPA SGUEE LAPA SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION SUCCESSION						
FR 2	35 36 37 38 39 40 41 42 43 44 45 46 47 48 49	THP TYR GLN UYSO LYSS LYSS LEU LEU LEU LEU TYR	TRP GLN GLN LYS	TYR		TRP TYR HIS GLX LYS PRO	TAP TYA GLX TYA	TAP TYAN GUN LYS GLU LYS ALA PRO GLU LYS LEU LEU TYR LYS)) 			TAPA TYAN GLNS CHORE CAN CAN CAN CAN CAN CAN CAN CAN CAN CAN		TAP TYR GLN LYS PRO GLY LYS ALA PRO GLU LEU LEU LEU LEU	GLY GLN PRO PRO LYS LEU LEU ILE	TYR GLN GLS LEUY GLN GLN GLN C GLN C C C C C C C C C C C C C C C C C C C	LYS PRO GLN PRO LYS LEU LEU ILE TYR	PRO LYS LEU LEU ILE TYR	TYR GLN GLN PRO						
CDAN	50 51 52 53 54 55	ASP ALA SER ASN LEU GLU THF						LYS ALA SER THA LEU GLU THF	1))			ALA SEF THF LEU SE	, , , , , ,	ALA SER THR LEU GLN SEF	LEU ALA SER	LEU ALA SEF	LYS LEU ALA SEF								
۴ Я З	58 578 90123 456678 90123 456678 90123 456678 90123 456678 88678 88678 88678 88678 88678 88678 88678 88678 88678	THEYAL OF A SELLER A SECTION OF					-	GV REARES VELVEL ELARGE TURGEL LARGEN LINY YY	A CARLEN VELLEY MALENT A NOPPE ARREN			GV PSAPS GSGSG TAPTI TISCI GSGAP ATTYC	אין סרסשיר אראשיי אישטרא אייטאא אייטאר אייטאט אראטא אראטא אייט איין אייטאט אראטא אייט איין אייט אייט אייט אייט	GV REARES GEORGE TARTER TERESCE GEORGAP ATTYCE	LA CHARTER YRYRY SABERU FERSUN AFUS A ATTYRY	UN DESAME ANY SUBAL A STATE A SOLA A ATTYCH	CLA OFFICIENT AND AND A CONSISTENT AND A	ULU CROENT ANTRY ANTRA CYLL NSOPPA ARRAS			s				
	89 90 91 92 93 94 95 95 95 95 95 95 95 95	GLL GGLYATH PA LEI A BC DEF										GLU GLU TYY STY PR		GLU GLU TYI SE PH PR	H TYI R TYI E SEL O GL GL THI AS	H ITH R ILE R TYI Y SE R SE Y GL R AS	GL Y GL A TY	N GLL NY GLL A TY A SEL SEL SEL SEL TY A SEL SEL TY		GLUSA ALLT :: : : :	XXXA AR				
1	98 99 100 101 102 102 104 105 106 106 106		EYLYA SLUR SLUR			•		PHUGGULH GGGGT LLESUL LLESUL	YNYR SUPE - S	· .			-		-										
	108	AF T)	ig IR					AR Th	เคี			-			-					11					

•																		
IMAN K	арра 97 LOD	98 HBJ 10	CHAIP 99 BEN	100 GR	101 MAA	102 MUK		104* MAR	105 AMYLOID	106 BJ1	107 HBJ 6	108 PEN	109 AMYLOID MS	210 CL*	111 GM131 'CL #	# OF SEQUENCES	# OF AMINO ACIDS	OCCURRENCES OF MOST COMMO AMINO ACID
0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16	ASP ILE GLN THR GLN SERO SER IN SER ALA VAL	ASP ILENT GMET RURAR SPRE RURAS LEERAS SELSERAS	ASPENT RNROR RURA	ASPENT RALAROR ALLAR SPACE ALLAR SPACE ALLAR SPACE ALLAR SPACE ALLAR SPACE ALLAR SVAL	ASP ILEN IEU THR GLEN FRAG SER ILER ALA	ASP ILENT HLRROG SPROR SPROR SEEUS SEEUS	ASP ILE GLEN THR GLEN SER SER LEU	ASP ILN IGLN IED THAN SEROR SER LEU	ASP ILE GET PGER PGER PGER SEA SEU	ASX MET THR GLN	ASP ILE GLN MET THR GLN	ASP ILE GLN MET	ASP Ø GLN MET THR			1 109 107 108 108 109 108 107 105 105 105 104 103 102 101 97 93 93 90	1 4 3 4 4 3 2 3 4 5 5 4 4 7 32 4 6 3	109(ASP): 99(ASP) 109(ASP): 99(ASP) 101(ILE) 97(GLN): 93(MET) 105(THR) 107(GLN): 100(GLN 103(FER) 90(FER) 91(AER) 94(SER) 94(SER) 93(ASP): 94(ASP) 92(GLY) 87(ASP): 94(ASP)
17 18 19 20 21	-									ARG VAL THR ILE THR CYS						91 91 88 88 83	4 4 7 1	88(VAL) 87(THR) 84(ILE) 75(THR) 83(CYS) 43(ARG)
22 23 24 25 26 27 27 27 27 27 27 8									-	GLN ALA SER GLN						75 75 72 72 4 4	5 4 4 4 2 2	71(ALA) 57(SER) 66(GLN) : 53(GL 3(SER) 3(VAL)
27C 27D 27E 27F								•		ASP					. ·	2 2 1 72	2 2 1 8 5	1(+) 1(+) 1(SER) 25(ASP) : 22(SE 51(II F)
28 29 30 31 32 33								•	• .	ILE ASN LYS						71 68 66 66 64 60	10 : 11 10 7 4 7 : 1	24(SEH) 33(TYR) 60(LEU) 3 24(ASN) : 22(A)
34 35 36 37 38																63 61 60 58 55 57	1 2 4 4 4 4	63(TRP) 57(TYR) 56(GLN) : 49(G 55(GLN) : 50(G 50(LYS) 54(PRO)
39 40 41 42 43 44 45 45 45 47																44 46 47 47 47 46 45 43	4352 16722	50(LYS) 54(PRO) 35(LYS) 42(ALA) 47(PRO) 41(LYS) 53(LEU) 41(LYS) 44(LEU) 44(LEU) 42(LE) 42(LE)
48 49 50 51 52 53						_										45 45 45 44 43 44 44	· 5 4 5: 7	8 15(+) 39(ALA) 41(SER) 6 18(SER) 43(LEU) 20(GLU) 27(SER)
54 55 56 57 58 59 60 61 62 63 64 65																- 42 43 43 42 42 43 43 43 43 43 43 43	7 1 2 4 1 3 3 3 7 1 4 3 3 3 3	27(SER) 43(GLY) 42(VAL) 39(PRO) 42(SER) 41(APKE) 36(SER) 36(SER) 38(SER) 38(SER) 38(SER) 38(SER)
66 67 68 69 70 71 72 73 73																41 41 41 40 40 40 40 40	3:443 5:443 43	38(GLY) 98(THR) 6 25(ASP) : 23/ 36(PHE) 37(THR) 31(LEU) 37(THR) 38(LE) 37(SEP)
74 75 76 77 80 81 82 83										•					LEU GLN PRO GLU ASF PHE	40 40 40 40 40 40	2 5 2 3 3 1 4 2	27(SER) 35(LEU) 3 39(GLN) : 35(33(PRO) 5 29(GLU) : 26(2 40(ASP) : 37(/ 28(PHE)
84 85 86 87 88										TY TY CI	'S				- THE - TYE - TYE - CYS - GU	42 41 5 42 42 43 43 43	4 22 1 3 : 3 :	
89 90 91 92 93 93 94 95 95 95 95	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5													AS IL Pf		2 46 5 46 9 46 5 45 5 5 4 4 4	10 : 8 9 10 3 2 3 1	12(LEU) 35(PRO) 2(+) 2(+) 2(+) 2(TYR) 1(ASP)
95 95 97 97 97	5F 6 7 9		,							Ť	ΎΑ			G TI P G	LY LE HA TH HE PH LY GL LN GL	E 31 Y 31 Y 31	- 12 	29(PHE) 31(GLY) 5 18(GLN) : 17 31(GLY)
100 101 102 103 104 104 104 104 104 104	0123456													L V G	LY GL HA TH YS TH AL VA LU GL LE ME 	R 31 R 31 L 30 U 30 T 31	3:	2 30(THR 23(LYS) 23(VAL) 2 20(GLU) : 19 7 15(ILE) 2 29(LYS)
10	7													A	NG AR		. 2	2 22(ARG) 20(THR)



12. 2.

47 ANTIBODY SPECIFICITIES: HUMAN KAPPA LIGHT CHAINS SUBGROUP I A WEA: ANTI-3,4-PYRUVYLATED GALACTOSE MONOCLONAL COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVIT 251 LOW: 39) LAY: ANTI-HUMAN GAMMA G1 AND G3 GLOBULINS: PO IDIOTYPE 39) HEI: COLD AGGLUTININ WITH ANTI-GD (MEMBRANE-GLYCOLIPID-DEPENDENT) ACTIVITY 66) DAV: ANTI-HUMAN GAMMA G GLOBULIN 67) FIN: ANTI-HUMAN GAMMA G GLOBULIN 92) WAG: ANTI-DINITROPHENYL 104) MAR: ANTI-LIPOPROTEIN LIPASE ALLOTYPE: HUMAN KAPPA LIGHT CHAINS SUBGROUP 1 79) KUE: INV(2) CLASS: HUMAN KAPPA LIGHT CHAINS SUBGROUP I 8) WEA: 'IGM-KAPPA 33) F-GUI: IGG3-KAPPA 55) S-GUI: IGG3-KAPPA 74) PW: IGG1-KAPP IGG1.KAPPA REFERENCE: HUMAN KAPPA LIGHT CHAINS SUBGROUP I 1) ROY: HILSCHMANN, & CRAIGLC. (1965) PROC.NAT.ACAD.SCI.USA.53.1403-1409; HILSCHMANN,N. (1967) Z.PHYSIOL.CHEM.348,1077-1080; HILSCHMANN,N. BARNIKOL,H.U.HESS.M.LANGER.B.PONSTINGLH.STEINMETZ-KAYNE,M.SUTER,L. & WATANABES. (1969) PROC. 5TH FEBS SYMP., 15,57-74, (CHECKED BY AUTHOR WHO PROVIDED ADDITIONAL RESIDUES TO THOSE PUBLISHED AND CORRECTED RESIDUES 65 AND 67 AS GIVEN IN THE TABLE) GIVEN IN THE INGLE? 2) AU: SCHIECHL,H. & HILSCHMANN,N. (1971) Z.PHYSIOL.CHEM.352.111-115: (1972) Z.PHYSIOL.CHEM.353.345-370. (CHECKED BY AUTHOR) 3) RE: PALM,W. & HILSCHMANN,N. (1973) Z.PHYSIOL.CHEM.354.1651-1654: (1975) Z.PHYSIOL.CHEM.356.167-191. (CHECKED BY AUTHOR) 4) HAU: WATANABE,S. & HILSCHMANN,N. (1970) Z.PHYSIOL.CHEM.351.1291-1295. (CHECKED BY AUTHOR) a) Add: WATANABES. B INISONWANA, (1970) 22 THOLOLONEW TAD. TAD. (1970) 1105 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (1974) 2010 (7) AG: TITANI,K.,SHNODA,T. & PUTNAM,F.W. (1985) J.BIOL.CHEM.,224.3535550. (CHECKED BY AUTHOR 02/3265)
 8) WEA: GONI,F. & FRANGIONE,B. (1983) PROC.NAT.ACAD.SCI.USA.80.4837-4843. (CHECKED BY AUTHOR 03/23/84)
 9) HX137CL: BENTLEY,D.L. & RABBITTS,T.H. (1983) CELL.32.181-189.
 10) HX134'CL: BENTLEY,D.L. & RABBITTS,T.H. (1983) CELL.32.181-169.
 11) DAUDI'CL: KLOBECK.H.G.,COMBRIATO,G. & ZACHAU.H.G. (1984) NUC.ACIDS RES. 12.18.6995-7006.
 12) WALKER'CL: KLOBECK.H.G.,COMBRIATO,G. & ZACHAU.H.G. (1984) NUC.ACIDS RES. 12.18.6995-7006.
 12) WALKER'CL: MEDBECK.H.G.,COMBRIATO,G. & ZACHAU.H.G. (1984) NUC.ACIDS RES. 12.18.6995-7006.
 12) WALKER'CL: KLOBECK.H.G. (2018) AUTHOR 02/22/85 WHO 13) HF3-16/6: ATKINSON,P.M.,LAMPMAN,G.W.,FURIE,B.C.,NAPARSTEK,Y.,SCHWARTZ,R.S.,STOLLAR,B.D. & FURIE,B. (1985) J.CUN.INVEST.,75.1138-1143. (CHECKED BY AUTHOR 08/21/85) 14) HF2-1/13B: ATKINSON.P.M. LAMPMAN.G.W., FURIE, B.C., NAPARSTEK, Y., SCHWARTZ, R.S., STOLLAR, B.D. & FURIE, B. (1985) J.CLIN. INVEST., 75, 1138-1143. (CHECKED BY AUTHOR 09/21/85) 15) HF2-10/2: ATKINSON,P.M. LAMPMAN, G.W.,FURIE, B.C., NAPARSTEK,Y., SCHWARTZ,R.S., STOLLAR, B.D. & FURIE, B. (1985) J.C.LIN.INVEST., 75, 1138-1143. (CHECKED BY AUTHOR 08/21/55) 15) HF2-1/17: ATKINSON,P.M. LAMPMAN,G.W.,FURIE,B.C.,NAPARSTEK,Y.,SCHWARTZ,R.S.,STOLLAR,B.D. & FURIE,B. (1985) J.CUN.INVEST.,75,1138-1143. (CHECKED BY AUTHOR 08/21/55) 17) BJ26: ALESCIO-ZONTAL, & BAGLIONI,C. (1970) EUR.J.BIOCHEM.,15,450-463. (CHECKED BY AUTHOR) 18) RFZ: SMITHES,O., GIBSON,D.,FANNING,E.M.,GOODFLIESH,R.M.,GILMAN,J.G. & BALLANTYNE,D.L. (1971) BIOCHEMISTRY,10,4912-4921. (CHECKED BY AUTHOR) 19) PSM: SEON.B.K. (1982) MOL.IMMUNOL., 19.83-86. (CHECKED BY AUTHOR 05/23/83) SEDNIE SEDNIK, (1982) MOLIMMUNUL. ISIOSOBI, (CHEDRED DI AUTHOR 051201401 OLIMMUNOL.,18,793-805.
 SEM IGG: KUAN.T.K.,TUNG,E.,WANG,I.Y. & WANG,A.C. (1981) IMMUNOL.,44,255-271. (CHECKED BY AUTHOR 05/26/83)
 ESM IGM: KUAN.T.K.,TUNG,E.,WANG,I.Y. & WANG,A.C. (1981) IMMUNOL.,44,255-271. (CHECKED BY AUTHOR 05/26/83) 23) WAT: STEVENS,F.J.,WESTHOLM,F.A.,PANAGIOTOPOULOS,N.,SCHIFFER,M.,POPP.R.A. & SOLOMON,A. (1981) J.MOL.BIOL.,147,185-193. (CHECKED BY AUTHOR 05/26/1983) 24) AMYLOID VII.BE GLENNER.G.G.TERRY,W.,HERADA.M.,ISERSKY,C. & PAGE.D. (1971) SCIENCE,172,1150-1151. (CHECKED BY AUTHOR 09/22/78) 25) LOW: CAPRA,J.D. KENDE,J.M.,WILLIAMS,R.G.,R.,FEIZI,T. & KUNKEL.H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR WHO CORRECTED RESIDUE 16 AS GIVEN IN TABLE) 26) DIE: CAPRA,J.D. & KUNKEL.H.G. (1970) PROC.NAT.ACAD.SCI.USA.67,87-92. (CHECKED BY AUTHOR) CAPRAJD, & KUNKELHG, (1970) FROCINATIACADISCIUSALDIAS, (URLED BT AUTHOR)
 CAPRAJD, & KUNKELHG, (1970) FROCINATIACADISCIUSALDIAS, (URLED BT AUTHOR)
 TEI: CAPRAJD, & KUNKELHG, (1970) FROCINATIACADISCIUSALDIAS, (URLED BT AUTHOR)
 BJ48: ALESCID-ZONTAL, & BAGUIONIC, (1970) FURJ.BIOCHEM., (15450-463, (CHECKED BY AUTHOR)
 BJ48: ALESCID-ZONTAL, & BAGUIONIC, (1970) FURJ.BIOCHEM., (15450-463, (CHECKED BY AUTHOR)
 OCN: NIALL, HD, & EDMAN, P. (1957) NATURE, 216, 262-263, (CHECKED BY AUTHOR 07(25/79)
 TRA: NIALL, HD, & EDMAN, P. (1957) NATURE, 216, 262-263, (CHECKED BY AUTHOR 07(25/79) 32) AMYLOID LEP: LIANJ.B.SKINNER.M.BENSON.M.D. & COHEN.A.S. (1977) BIOCHIM.BIOPHYS.ACTA.491.167.176.
 33) F.GUI: WANG.A.C.FUDENBERG.H.A. & CREYSSEL.R. (1982) ACTA HAEMAT.68.187-195. (CHECKED BY AUTHOR 05/26/83)
 34) OU(IOC): KOHLER.H.SHIMIZU.A.PAUL.C. & PUTNAM.F.W. (1970) SCIENCE.169.56-59. (KAPLAN.A.P. & METZGER.H. (1969) BIOCHEMISTRY.8.3944-3951.) (CHECKED BY AUTHOR 05/15/83) (CHECKED BY AUTHOR 08/15/63)
35) DEE: MILSTEIN.C. & DEVERSON.E.V. (1971) BIOCHEM.J.,123,945-958. (CHECKED BY AUTHOR)
36) GALIDI: LAURE.C.J.WATANABE.S. & HILSCHMANN.N. (1973) Z.PHYSIOL.CHEM.354,1503-1504. (CHECKED BY AUTHOR)
37) JOH: CAPRA.J.D. & KUNKEL.H.G. (1970) PROC.NAT.ACAD.SCI.USA.67.87-92. (CHECKED BY AUTHOR)
39) KER: MILSTEIN.C. (1966) BIOCHEM.J.,101,352-368. (CHECKED BY AUTHOR BWIDDED ADDITIONAL RESIDUES TO THOSE PUBLISHED)
39) LAY: KAPLAN.A.P. & METZGER.H. (1969) BIOCHEMISTRY.8,3944-3951. (CHECKED BY AUTHOR) KLAPPER.D.G. & CAPRA.J.D. (1976) ANN.IMMUNDL.(INST.PASTEUR), 127C.261-271. (CHECKED BY AUTHOR 09/01/79)
40) BRA: WANG.A.C.WELLSJ.V., FUDENBERG.H.H. & GERGELYJ. (1974) IMMUNDCHEM..11.341-345. (CHECKED BY AUTHOR)
41) WES: KRATZIN.M.YANG.C.Y.KRUSCHEJ.U. & HILSCHMANN.N. (1980) Z.PHYSIOL.CHEM.351,1591-1598.
42) V5'CL: PECH.M.JAENICHEN.H.-R.POHLENZ.H.-D.NEUMAIER.P.S.KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR)
43) WES: VARD.A.C.Y. RUBCHENZ.H.D.NEUMAIER.P.S.KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR)
44) WES: VARD.A.C.Y. RUBCHENZ.H.D.NEUMAIER.P.S. KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR)
45) WICL: PECH.M. JAENICHEN.H.-R.POHLENZ.H.D.NEUMAIER.P.S. KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR)
46) WES: VARD.A.C.Y. KRUSCHEJ.U. A HILSCHMANN.N. (1980) Z.PHYSIOL.CHEM.351,1591-1598.
47) WES: KARTZIN.H.YANG.C.Y.KRUSCHEJ.U. A HILSCHMANN.N. (1980) Z.PHYSIOL.CHEM.351,1591-1598.
48) VICL: PECH.M.JAENICHEN.H.-R.POHLENZ.H.D.NEUMAIER.P.S. KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR)
49) VICL: PECH.M. JAENICHEN.H.-R.POHLENZ.H.D.NEUMAIER.P.S. KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176,189-204. (CHECKED BY AUTHOR) 43) Vb"CL: PECH.M. JAENICHEN,H.R.,POHLENZ,H.D.,NEUMAIER,P.S.,KLOBECK,H.-G. & ZACHAU,H.G. (1984) J.MOL.BIOL.,176,189-204. (CHECKED BY AUTHOR 12/14/84) AUTHOR 12/14/84) 44) HK102'CL: BENTLEY,DL, & RABBITTS,T.H. (1980) NATURE,288,730-733. (CHECKED BY AUTHOR 11/30/82) 45) EU: GOTTLIEB,D.,CUNNINGHAM,B.A.,RUTISHAUSER,U. & EDELMAN,G.M. (1970) BIOCHEMISTRY,9,3155-3161. (CHECKED BY AUTHOR) 46) DEN: YANG,C.Y.,PAULY,E.,KRATZIN,H. & HILSCHMANN,N. (1981) Z.PHYSIOL.CHEM.,362.1131-1146. 47) PAU: DAYHOFF,M.O. (1972) ATLAS OF PROTEIN SEQUENCE & STRUCTURE,5,D-245. SUBMITTED BY SMITHIES,O.,GIBSON,D.M. AND FANNING,E.M. (CHECKED BY AUTHOR) 48) HBJ4: SMITH,G.P.,HOOD,L. & FITCH,W.M. (1971) ANN.REV.BIOCHEM.40,965-1012. 40) FDA. HUNKER OF DECORDENT & SCHERE BERG MALL (1971) ANN.REV.BIOCHEM.40,965-1012. 49) FRA: MEINKE.G.C. SIGRIST.P.H. & SPIEGELBERG.H.L. (1974) IMMUNOCHEM. 11.457-460. (CHECKED BY AUTHOR WHO PROVIDED ADDITIONAL RESIDUES TO THOSE PUBLISHED); MEINKE.G.C. & SPIEGELBERG.H.L. (1976) IMMUNOCHEM. 13.915-919. (CHECKED BY AUTHOR 10/17/77) 50) GR': FAIR,D.S.,SLEDGE,C.,KRUEGER,R.G.,MANN,K.G. & HOOD,L.E. (1975) BIOCHEMISTRY,14,5581-5568. 51) PAUL: SMITH,G.P.,HOOD,L. & FITCH,W.M. (1971) ANN,REV.BIOCHEM.,40,969-1012. 51) PAUL: SMITH, G.P. HOOD, L. & FITCH, W.M. (1971) ANN.REV.BIOCHEM., 40.969-1012.
52) MON: NIALL, H.D. & EDMAN, P. (1967) NATURE 216,262-263. (CHECKED BY AUTHOR 07/25/79)
53) HEI: RIESEN, W.F., MAJANIEMI, J.HUSER, H., BRAUND, G. & ROELCKED. (1976) SCAND, J.IMMUNOL., 6,145-148. (CHECKED BY AUTHOR 10/19/79)
54) POT: CAPRA, J.D. & KUNNEL, H.G. (1970) PROCNAT, ACAD, SCI, USA, 67, 9-22. (CHECKED BY AUTHOR WHO CORRECTED RESIDUE 9 AS GIVEN IN TABLE)
55) S-GUI: WANG, A.C., FUDENBERG, H.H. & CREYSSEL, R. (1982) ACTA HAEMAT, 66, 137-195. (CHECKED BY AUTHOR 05/26/83)
56) AMYLOID BAN: DWULET, F.E., O'CONNOR, T.P. & BENSON, M.D. (1986) MOL.IMMUNOL, 23, 73-78.
57) BJ19: ALESCIO-ZONTAL, & BAGLIONIC. (1970) EUR, JBIOCHEM, 15: 450-463. (CHECKED BY AUTHOR)
59) BL1: MILSTEINC, (1960) PROC. STH FEBS SYMP, 15: 43-56. (CHECKED BY AUTHOR WHO PROVIDED ADDITIONAL RESIDUES TO THOSE PUBLISHED AND CORRECTED RESIDUES 1, 38, 27, 79 AND 82 AS GIVEN IN TABLE)
50) BL1: GEONI BY, (1920) MOLIMILINGI (19 2936. (CHECKED BY AUTHOR WHO PROVIDED ADDITIONAL RESIDUES TO THOSE PUBLISHED BY AUTHOR DY ADDITIONAL RESIDUES 1, 38, 27, 79 AND 82 AS GIVEN IN TABLE) AND CORRECTED RESIDUES 1.3.8.27.19 AND 82 AS GIVEN IN TABLE! SEON.B.K. (1982) MOLIMMUNOL.19.83-86. (CHECKED BY AUTHOR 05/23/83) NIALLH.D. & EDMAN.P. (1967) NATURE,216,262-263. (CHECKED BY AUTHOR 07/25/79) MILSTEIN.C.P. & DEVERSON.E.V. (1974) EUR.JBIOCHEM.48,377-391. (CHECKED BY AUTHOR) EULITZ.M. & LINKE.R.P. (1982) Z.PHYSIOL.CHEM.363,1347-1358. (CHECKED BY AUTHOR 10/10/83) 59) JBL: 60) PAP: 61) CAR: 62) MEV:

48 REFERENCE: HUMAN KAPPA LIGHT CHAINS SUBGROUP I (contid) 63) BI: BRAUN,H.,LEIBOLD,W.,BARNIKOL,H.U. & HILSCHMANN,N. (1971) Z.PHYSIOL,CHEM.,352,647-651; (1972) Z.PHYSIOL,CHEM.,353,1284-1306. (CHECKEO BY AUTHOR WHO PROVIDED AN ADDITIONAL RESIDUE TO THOSE PUBLISHED AND CORRECTED RESIDUE 72 AS GIVEN IN TABLE) 69) Vorce PECH.M. JAENICHEN.H.-R., POHLENZ,H.-D., NEUMAIER,P.S., KLOBECK,H.-G. & ZACHAU,H.G. (1984) J.MOL.BIOL., 176, 189-204. (CHECKED BY AUTHOR 12/14/84) 12/1904) 70 J.UX: NIALL.H.D. & EDMAN.P. (1967) NATURE.216.262-263. (CHECKED BY AUTHOR 07/25/79) 71) ТИЕ: MATTHEWS.J.B. & JEFFERIS.R. (1977) IMMUNOCHEM..14.793-797. (CHECKED BY AUTHOR 08/10/79) 72) Va"CL: PECH.M.JAENICHS.H.H.-R.,POHLENZ.H.-D.,NEUMAIER.P.S.,KLOBECK.H.-G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176.189-204. (CHECKED BY AUTHOR 12/14/84) AUTHOR 12/18/4) 73) NI:- SHINODA,T. (1973) J.BIOCHEM.,73.433-448. (CHECKED BY AUTHOR) 74) FW: PICKAL,WANG,A,C,FROHLICHMAN,R. & FUDENBERG,H.H. (1982) ACTA HAEMAT..68,207-214. (CHECKED BY AUTHOR 05/26/83) 75) AMYLOID X: GLENNER,G,G,TERRY,W,HERADA,M,JSERSKY,C. & PAGE,D. (1971) SCIENCE.172.1150-1151. (CHECKED BY AUTHOR 09/22/78) 76) ALE: MILSTEIN,C,MILSTEIN,C,P,& FEINSTEIN,A. (1969) NATURE.221.151-154. (CHECKED BY AUTHOR 09/21/72) 76) ALE: MILSTEIN,C.MILSTEIN,C.P.& FEINSTEIN.A. (1969) NATURE.221.151-154. (CHECKED BY AUTHOR)
77) THE: MEINKE.G.C. & SPIEGELBERG.H.L. (1976) IMMUNOCHEM..13.915-919. (CHECKED BY AUTHOR 10/17/77)
78) ADA: MEINKE.G.C. & SPIEGELBERG.H.L. (1976) IMMUNOCHEM..13.915-919. (CHECKED BY AUTHOR 10/17/77)
79) KUE: EULITZ.M.KLEY.H.P. & ZEITLER.H.J. (1979) Z.PHYSIOL.CHEM.360.725.704. (CHECKED BY AUTHOR 07/17/79)
80) GG: WANG.A.C..FUDENBERG.H.H. & CREYSELLR. (1974) EUR.JIMMUNOL.4.446-448. (CHECKED BY AUTHOR 07/17/79)
81) BOL: WANG.A.C..WELLS.J.V.FUDENBERG.H.H. & GERGELY.J. (1974) IMMUNOCHEM..11.341-345. (CHECKED BY AUTHOR)
82) RI: PICK.A.J.WANG.A.C..FROHLICHMAN.R. & FUDENBERG.H.H. (1982) ACTA HAEMAT.68.207-214. (CHECKED BY AUTHOR)
83) Ve'CL: PECH.M.J.AENICHEN.H.R.,POHLENZ.H.-D.,NEUMAIR.P.S..KLOBECK.H.G. & ZACHAU.H.G. (1984) J.MOL.BIOL..176.189-204. (CHECKED BY AUTHOR 12/14/84) 84) OCO: WANGA.C.,WELLS.J.V.,FUDENBERG.H.H. & GERGELY.J. (1974) IMMUNOCHEM.,11.341-345. (CHECKED BY AUTHOR) 85) V13'CL: JAENICHEN,H.-R.,FECH,M.,LINDENMAIER,W.,WILDGRUBER.N. & ZACHAU,H.G. (1984) NUC.ACIOS RES.,12.5249-5263. (CHECKED BY AUTHOR 12/14/84) 12/14/84) BO V18A*CL: HEIDMANN.O. & ROUGEON.F. (1984) NATURE.311.74-76. 67) V19A*CL: HEIDMANN.O. & ROUGEON.F. (1984) NATURE.311.74-76. 80) V18B*CL: HEIDMANN.O. & ROUGEON.F. (1984) NATURE.311.74-76. 89) V18B*CL: HEIDMANN.O. & ROUGEON.F. (1984) NATURE.311.74-76. 07 1189 С. ПЕЛМИНИ, А ПООССИЛГ. (1389) НАГОЛЕЗТ (1470. 90) HF8-21/28: ATKINSON, M., LAMPMAN, G.W., FURIE, B.C., NAPARSTEK, Y., SCHWARTZ, R.S., STOLLAR, B.D. & FURIE, B. (1985) J.C.LIN.INVEST., 75.1138-1143. (СНЕСКЕФ ВУ АИТНОЙ 08/21/65) 90) HF8-21/28: ATKINSON,P.M. LAMPMAN,G.W.,FURIE,B.C.,NAPARSTEK,Y.,SCHWARTZ,R.S.,STOLLAR,B.D. & FURIE,B. (1985) J.CLIN.INVEST.,75.1138-1143. (CHECKED BY AUTHOR 092/169)
91) SAC: SMITHES,O.,GIBSON,O.M.,FANNING,E.M.,PERCY,M.E.,PARR,D.M. & CONNELL,G.E. (1971) SCIENCE,172.574-577. (CHECKED BY AUTHOR)
92) WAG: KAPLAN,A,P. & METZGER,H. (1959) BIOCHEMISTRY.8.3944-3951. (CHECKED BY AUTHOR)
93) HBJ1: HOODL.,GRAY,W.R.,SANDERS,B.G. & DREYER,W.J. (1967) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,133-145.
94) AMYLOID 547: WESTERMARK P.SLETTEN,K. & NATVIG,J.B. (1981) ACTA PATH.MICROBIOL.SCANO.C69,199-203. (CHECKED BY AUTHOR 10/17/77)
96) HOE: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH,E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.,66,842-847. (CHECKED BY AUTHOR 10/17/77)
97) LOD: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH,E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.,66,842-847. (CHECKED BY AUTHOR 10/17/77)
98) HBJ10: HOODL.,GRAY,W.R.,SANDERS,B.G. & DREYER,W.J. (1987) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,133-145.
99) BEN: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
90) GR: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
910) GR: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
910) GR: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
9110 MAA: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
9100 GR: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTHOR)
9110 MAA: CAPRA,J.D.,KEHOE,J.M.,WILLIAMS,R.C.,JR.,FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.69,40-43. (CHECKED BY AUTH 100) FEN: MUULINA, & FOUGEREAU, M. (1973) NATURE NEW BIOLOGY.258,176.176. 109) AMYLOID MS: PICKAL,SCHREIBMAN,S.,LAVIE,G. & FROHLICHMAN,R. (1973) PROTIDES BIOL.FLUIDS.20,63-72. 110) CL: SOLOMONA, MCLAUGHLIN,CL. & CAPRAJD. (1975) J.CLINICAL INVESTIGATION.55,579-586. (CHECKED BY AUTHOR) 111) GM131'CL: MORIN,J.W.,BLACK,A.,WU,M. & BEYCHOK,S. (1985) PROC.NAT.ACAD.SCI.USA.82.7025-7029. NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP I IDENTICAL SETS OF FRAMEWORK SEGMENTS: L SETS OF FRAMEWORK SEGMENTS: SET 1: ROY(1)AU(2)[REI3]HAUIA]HK101°CL[5].SCW[8].AG[7].WEA[8].HK137°CL[9].HK134°CL[10].DAUDI°CL[11].WALKER°CL[12]. SET 1: ROY(1)AU(2)[REI3]HAUIA]HK201°CL[5].HF2-1/17]16].BJ26[17].RF2(16].PSM[19].HOM[20].ESM [GG[21].ESM [GG[22].ESM [GG[21].ESM [GG[22].ESM [GG[21].ESM [GG[22].ESM [GG[22] FR1: G. C. (169), L. (2) IDENTICAL)
 ROY(1), L(2) IDENTICAL)
 ROY(1), L(2), WALKERTCL(12), Vb°CL(42), Vb°CL(43), HK102°CL(44), KA(66), Vd°CL(69), Va°CL(72), Va°CL(83), (10) IDENTICAL)
 HK107°CL(3), HK104°CL(10), (2) IDENTICAL)
 HK107°CL(3), HK104°CL(10), (2) IDENTICAL)
 HK107°CL(3), HK104°CL(10), (2) IDENTICAL)
 Y134°CL(86), IDENTICAL TO 7 MOUSE V-KAPPA-II; PC1229(NZB)(1), PC2680(NZB)(2), PC7132(NZB)(9), MOPC70(5), PC2413(NZB)(11), V134°CL(86), (10) IDENTICAL TO 7 MOUSE V-KAPPA-II; PC1229(NZB)(1), PC304(2), K22-213(0), V20°CL(36), K16-187(641), V138°CL(86), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V130°CL(136), K16-187(641), V138°CL(86), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P010°CL(86), V138°CL(86), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P130°CL(86), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P130°CL(80), V138°CL(80), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P130°CL(80), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P130°CL(80), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P130°CL(80), (2) IDENTICAL HUMAN V-KAPPA-I; ALSO 4 HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P110°CL(3), IDENTICAL HUMAN V-KAPPA-IV, V10°CL(11), VKAPPA IV GERMLINE CL(2), P110°CL(3), IDENTICAL HUMAN V-KAPPA-IV, V10°CL(3), IDENTICAL HUMAN V-KAPPA-F82: SET 5: V19BC PB1 TEF 210(NZB)[23],H36-15[28],2242[29],V-2161.5KB'CL[30],V-21C9.5KB'CL[31], [A.BY][35],10.4(A.TH)[39],H35-5(48],40.C(A.TH)[52],M0PC53[54],A8PC22[55], B16KB'CL[56],11939[62]: 1 MOUSE V:KAPPAVI: BFPC61A'CL[64], AND 15 RABBIT V-KAPPA: K49-501[45],3547[47],K4820[57],K30-267[61],311[65],4422[66],1709'CL[68], NZB)[21].PC 19],3368[20],BS-5[38],BS-1[39 ,4363[85],120[103],K-25[112],) 4192(7),3930(39),120(100),027(101),757 1: HAU(4),HK107(CL[5],HK137(CL[9],HK134(CL[10],Vb'CL[42],Vb''CL[43],Va''CL[72], (7 IDENTICAL) 2: V96'CL[63],V13''CL[65], (2 IDENTICAL) 3: V198'CL[68],V18'CL[69], (2 IDENTICAL) FR3: SEI 3: V198 CL[88],V188 CL[89]. (2 IDENTICAL)
SET 1: AU[2],GAL(1)[39],CL-(110]. (3 IDENTICAL HUMAN V-KAPPA-I: ALSO 2 HUMAN V-KAPPA-II: GM 607 'CL[5], RPM1-6410'CL[16]; 7 HUMAN V-KAPPA-II: I: AU[2],GAL(1)[39],CL-(110]. (3 IDENTICAL HUMAN V-KAPPA-I: ALSO 2 HUMAN V-KAPPA-II: GM 607 'CL[5],RPM1-6410'CL[16]; 7 HUMAN V-KAPPA-II: VLAPAIII: WOL[2]/PAY[7],PEI11],GLO[15],CDR[20],REE[57],VKAPPA3'CL[52]; AND 1 HUMAN V-KAPPA-IV: PB17IV'CL[3].
SET 2: HAU[4], (DENTICAL TO 1 HUMAN V-KAPPA-II: KOPAI-I: ALSO 2 HUMAN V-KAPPA-II: NIM3],FR[14]; 6 HUMAN V-KAPPA-III: NEU[5].
SET 3: AG[7],DEN[46],BI63], (0)[2],FR(2],11,RC(2],41-CL[28]; AND 1 HUMAN V-KAPPA-IV: LEN[4].)
SET 4: WEOTIFICATION (10)[2],FR(2],11,RC(2],12-CL[28]; AND 1 HUMAN V-KAPPA-IV: LEN[4].)
SET 4: WEOTIFICATION (10)[39], (2 IDENTICAL)
SET 5: WEALHER CL[12],OU(100)[34], (2 IDENTICAL HUMAN V-KAPPA-I; ALSO 1 HUMAN V-KAPPA-II: TEW[1].)
SET 6: WES[41],MEV[62], (2 IDENTICAL) FR4: IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS 1: AU[2].NE[7]1.SHE[7]1. (3: IDENTICAL) 2: WEA[8].GAL[0].36]. (2: IDENTICAL) 3: MK]34(2).10[.V5(L42].V5 CL[43].2] (3: IDENTICAL) 4: HF2-16[6]:3].HF2-1713B[14].HF2-18/2[15].HF2-177[16]. (4: IDENTICAL) 5: V4CL[69].V5(CL[33]. (2: IDENTICAL) COA1: SET SET 5: Vd'CL[69],Ve'CL[83]. (2 IDENTICAL) 1: HK101°CL[5],HK137°CL[9],HK134'CL[10],WALKER'CL[12],Vb'CL[42],Vb''CL[43]. (6 IDENTICAL) 2: AG[7],NI731. (2 IDENTICAL DIDENTICAL) 3: HK102°CL[44],Va''CL[72]. (2 IDENTICAL) 4: Vd'CL[69],Vo'CL[63],V13°CL[85], (3 IDENTICAL) 5: V18A'CL[66], (IDENTICAL TO 1 RABBIT V-KAPPA: 4153-1[24].) 6: V18A'CL[67], (IDENTICAL TO 1 RABBIT V-KAPPA: 4163-5[4].) 1: HK101°CL[5],HK134'CL[10], (2 IDENTICAL) 2: LAY[39], (IDENTICAL TO 1 HUMAN V-KAPPA-III: POM[48].) 3: Vb'CL[42],Vb''CL[43], (2 IDENTICAL) COR2: CDR3: IDENTICAL SETS OF J-MINIGENES: SETS OF J-MINIGENES: SET 1: AU[2]. [IDENTICAL TO 1 HUMAN V-KAPPA-II: RPM1-6410'CL[16]: 2 HUMAN V-KAPPA-III: PIE[11].VKAPPA3'CL[82]: AND 1 HUMAN V-KAPPA-III: F8[71V'CL[3]. SET 2: AG[7]. [IDENTICAL TO 1 HUMAN V-KAPPA-II: GOT[6]. SET 3: WALKEPRCL[12]. [IDENTICAL TO 1 HUMAN V-KAPPA-II: TEW[1].) SET 4: DEN[46].BI(63). (2 IDENTICAL HUMAN V-KAPPA-II: ALSO 1 HUMAN V-KAPPA-II: FR[14]: AND 3 HUMAN V-KAPPA-III: GAR'[10].FLO[12]. IARC/IBL41'CL[28].

NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP I (cont'd) GENERAL NOTES:

SEE SIGNAL PEPTIDE TABLE IF # OCCURS AT POSITION 0.

SPECIFIC NOTES:

AT POSITION

27C 27D 50 92 95A 95B

b) HK101'CL: THE SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN FOETAL LIVER DNA.
 7) AG: THE AMINO ACID RESIDUES AT POSITIONS 39 AND 41 WERE REPORTED BY THE AUTHORS AS GLY AND LYS RESPECTIVELY; HOWEVER, THE PROOF WAS NOT ABSOLUTE. THUS, THEY ARE OMITTED.

PROOF WAS NOT ABSOLUTE. THUS, THEY ARE OMITTED. 9) HK137'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN FETAL DNA. 10) HK134'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN FETAL DNA. 17) BJ26: ACID RESIDUES AT POSITIONS 39 AND 4) OF BJ26 WERE REPORTED BY THE AUTHORS AS GLY AND LYS RESPECTIVELY. SINCE THIS PROTEIN WAS SEQUENCED BEFORE THE SEQUENCES OF MANY OTHER PROTEINS WERE KNOWN AT THESE TWO POSITIONS, WE HAVE OMITTED THEM.

OMITTED THEM. 33) Figure The Sequences of Figurand Sigure FROM the same patient. 44) HK102'CL: The Sequence is obtained by translating the nucleotide sequence of a Clone of Human Foetal Liver DNA. 55) Sigure The Sequences of Figurand Sigure FROM the Same Patient. 56) APVLOID BAN: AMINO ACID RESIDUES FOUND AT POSITIONS 104 AND 105 ARE VALLEU AND GLN.GLU RESPECTIVELY. 57) BJ19: The Amino Acid Residues at Positions 39 AND 41 WERE REPORTED BY THE AUTHORS AS GLY AND LYS RESPECTIVELY. 59) JULI THES ADVISED BEFORE THE SEQUENCES OF MANY OTHER PROTEINS WERE KNOWN AT THESE TWO POSITIONS, WE HAVE SUBJECT THE AND ACID RESIDUES AT POSITIONS 104 AND THE REPORTED BY THE AUTHORS AS GLY AND LYS RESPECTIVELY. 59) JULI THE AMINO ACID RESIDUES DEFORE THE SEQUENCES OF MANY OTHER PROTEINS WERE KNOWN AT THESE TWO POSITIONS, WE HAVE OMITED THEM. 59) JBL: THE AMINO ACID RESIDUE FOUND AT POSITION 34 WAS ALA OR SER. 64) AMYLOID ES305: THE AMINO ACID RESIDUES AT POSITIONS 21 AND 29 WERE ILE OR LEU. 74) PW: THE SEQUENCE WAS FROM A PATIENT WITH TRANSITIONAL CELL CARCINOMA OF THE URINARY BLADDER. 82) AIL: THE SEQUENCE WAS FROM A PATIENT WITH TRANSITIONAL CELL CARCINOMA OF THE URINARY BLADDER. 109) AMYLOID MS: THE AMINO ACID RESIDUE AT POSITION 2 MS WAS ILE OR LEU. 111) GM131'CL: FROM AN EPSTEIN-BARR VIRUS-TRANSFORMED HUMAN LYMPHOID CELL LINE

308

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

RESIDUES

(LEU.VAL) (TRP.GLU) (ALA.ASP) YR.ASP.AS (SER.GLY)

-								.	•			:	50												
}	IUMAN	I KAPPA UGH INVARIANT RESIDUES	•	2	3	UP 11 CUM	5 GM 607 'CL	6 BAT	7 BATES	8. 808	9 SLO	10* WILS	11 GLI	12 AMYLOID TEW	13 RAI	14* FR #	15 YOS	16 RPM1- 6410 'CL	17 MAN	18 KIR	19 HYL	20 MAG	21 TVE	22 EID	23 Gal (II)
FA 1	0 1 2 3 4 5 6 7 8 9 9 1 1 2 3 4 1 5 6 7 8 9 9 1 1 1 3 3 1 1 5 6 7 8 9 9 1 1 1 2 3 4 5 6 7 8 9 9 0 1 1 1 3 1 4 5 6 7 8 9 9 0 1 1 2 3 4 5 6 7 8 9 9 0 1 1 1 2 3 4 5 6 7 8 9 9 0 1 1 1 2 3 4 5 6 7 8 9 9 0 1 1 1 2 3 4 5 6 7 8 9 9 0 1 1 1 2 3 4 5 8 7 8 9 9 0 1 1 1 2 3 1 1 1 1 3 1 5 8 7 8 9 9 0 1 1 1 2 3 1 1 1 1 2 3 1 1 1 1 1 1 1 1 1	SER PRO LEU SER LEU PRO(.96) VAL(.96) THR GLY PRO ALA SER ILE CYS	AND	ALLALT RANGEROU RUOALR OYUUOA SULERSOU SULE PALE RUOALR OYUUOA SULERSOU SULERSON SULERSON	TGSPAL RUOLA OYUOA REAS	THR GLN SER PRO	A ILALT RNROU RUOLAR OYUUOA A SILERS	ASP LALT ANA THEROU RUOLA OY UNDA RUDAL SELEAS AR	ASPE IVALT RINA OY L SEROU RUDONALA OY UNIT ROLU SEROU VALA OY UNIT ROLU PALA SEREAS ARG	ASPELELT RUSEROU RUCOLAR OYUUOA REERS G	APELLT RNR BUD RDOLR OY DOA REERS	ASPE VAL THUR SERVER SE	APELLT RNROU RUOLA OYUUOA AUERS	ALVAR TOSEL SLERAL ROYUDOR A SLERS	ASEALT RURADU RUOLA RULERS G NAME HURADU RUOLA RULERS G SEAL SLEPAT RELERS G	ASPELLT RUNAROU RUOLA RULAN TOUS RUU SLEVAL RULAN ASILAN TOUS RUU SLEVAL PALLON ARULAN AND AND AND AND AND AND AND AND AND A	ASE ALL ALL ALL ALL ALL ALL ALL ALL ALL AL	A SALT ANARAU AUDOLA DY ANALAS	ASP LEEA LERU VAL PROAL SEE SEAS CRG	ASP ILE VAL METR GLEAU SEEO VAL	ASP ILE VAL METR GLN SERO LEU PRO VAL	ASP ILE VAL METR GLERO LEU VAL	ASP ILE VAL MET THN SER PRO LE SER VAL	ASP ILAL MET THN SEAO LEU SEAO LEU VAL	ASP NLE VAL MET THR LEU SER LEU PRO VAL
CDR 1	24 25 26 27, 27, 27, 27, 27, 27, 27, 27, 27, 27,	B LEU D I	AR GRANNEL USR PYEPRU N	ARG SEER ALEU LEUX SER XSER AGLX AGLX AGLX AGLX ASP	AR GRANDER UPAR SELRAU LERR - P YTAS AS LAS LAS TYPE AS TYPE AS	A SERNAU UPRAY SEL UPRAY GASEL LASELYP YANA GASHRAU A SELYPU A SEL	A SEENRU USA SELE LISE S YANRU P	SERNAU SERNAU LEUSA SEL LEUSA SEL LEUSA SEL SER SEL SER SEL SER SEL SER SER SER SER SER SER SER SER SER SER	SER GLN LEU HIS ASX GLY ASX TYR	ALA SER GLX ARG IIIIII VAL LEU	SER SER SER LEU ARG HIS ASX	SER SER SER SER LEU LEU	SER SER GLN SER LEU	SEA SEA GLX		SEEN AS AS A SEEN A SEE		SERNALARA SELLALARA SEU VYSE SP A GASHARU A SP A CLNARA LAS							
FA2	34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49	GLY SER PRO LEU ILE TYR	TRP TYR LEU GLN LYS PRO GLY GLN	ASP TRP TLEUX GLXSOY GLXR PGLXU SER OGLUEL LEU TYR	TRP TYRE LEU GLN LYS GLN SER GLN SER GLN LEU LEU TYR	ASPRESSION SALES PROBUSE	TRA LEUN SOGLN SEAO GLUU LEE TYR	TRP TYPA LEUX LYS PRO GLY GLX PRO GLX								TAP TYA LEU GLN PAO GLN SER PRO GLN SER PRO GLU LEU LEU LEU ILE TYR		TREENN GOYNA OGOLA AROYNA OGOLA AROYNA OGOLA AROYNA AROU ILE AROU ILE TYR				,			
CORN	50 51 52 53 54 55 58	SER ARG	ARG	LEU GLY SER ASN ARG ALA SER	LEU GLY SER ASN ARG ALA SER	THR LEU SER TYR ARG ALA SER	LEU GLY SER ASN ARG ALA SER									LEU SER SER TYR ARG ASP SER		LYS VAL SER ASN ARG ASP SER							
FR3	578 901123 45878 901123 45878 90123 45878 90123 45878 90123 45878 90123 45878 90123 45878 90123 458888 8888888888888888888888888888888	SER SER GLY VAL PRO ARG PHE SER GLY SER GLY THR PHE LEU ILE ARG VAL VAL VAL CLY VAL CLY VAL CLY VAL CYS	OV PARAPS OSOSO TAPTAL LILSAN OAGAN GVITTO	GLA ROARDER YERY RAXERD SEERGL XAAPHE GERYRY RAXERY RAXERY RAXERY RAXERY RAXERY CALAXX YAAPHE	GV PAAPS GSGSA TAPTL ILSAA GPGAV GVTTC	GLAL OPPORTATION OF A CANADA C	GV PAAPS GSOSG TAPTA LISAAV GAGAV GVTTC								SER ARG	GV PAAPS ASGEST HAPTEL UNTARA LAUPL YLARAST		GUN PARPER YAYAN APERAL ULLERA ULUPL YARAPS GEGEE HAPEHAU SERAL ULUPL YARAPS GEGEE HAPEHAU SERAL ULUPL YARAS	VAL GLX ASLA GLLX GLLX GLLL TYPA CYS						
C D R 3	89 90 91 92 93 94 95 95 95 95 95 95 95 95 95 95 95 95 95	МЕТ	ALA PRO		SER PRO	MET GLN ARGUU ILEO	MET GLN ALA LEU GLN THR PRO			·						GLA ALA SEA PRO		MET GLN GLY THR HIS TAP SER			•				
FR4	96 97 98 99 100 101 102 103 104 105 106 106 107 108 109	THR PHE GLY GLY THR ILE ARG THR	PHE I GLY G GLN G GLY G THA T ARG J LEU G LU G LU G LYS I	LEU THA SLY SLY SLY SLY SLY SLU SLU SLU SLU	PHE GLY GLN GLY THR LYS GLU ILE LYS	LYS LEU GLU ILE ARG	GLN PHE GLY GLY THR UYS USL USL LYS ARG									THA PHE GLY GLY THA GLY LEU GLZ LYS ARA THA		PHE GLN GLN GLN GLN THA LYS GLU IL- LYS ARG	VAL GLU ILE LYS						

MA	N K/	арра	ыснт	CHAIL	45 SU	BGROI	JPH	cont d)				
		24* GIL	25 MEH	26 SC	27 Тн	28 SYV		808 2		# OF SEQUENCES	# OF AMINO ACIDS	OCCURRENCES OF MOST COMMON AMINO ACID	VARIABILIT
	0	ASP	ASP iLE VAL	ASP ILE VAL	ASP	ASP ILE	ASP ILE	ASP ILE VAL	ASP	31 30 30	1223	31(ASP) 29(ILE) 29(VAL)	1. 2.1 2.1 3.2
	2 3 4	MET	MET	MET	VAL MET	MET	VAL MET THR	leu	mei thr	30	3	28(MC+)	1.
	567	GLN SER PRO	THR GLN SER PRO	THR GLN SER	THR GLN SER	THR GUN	GUN			27 25	1	28(THR) 27(GLN) 25(SER) 24(PBO)	1. 1. 1.
	ลี 9	LEU	PRO		PRO					24 25 24	i	24(PRO) 25(LEU) 24(SEB)	1) 1.
	10	SER LEU ser								24 24 23 17	2	24(SER) 24(LEU) 23(PRO) 22(VAL) 17(THR)	1. 2.1 2.1
	12 13 - 14	301									21	17(THR) 16(PRO)	î. 2.1
	15 16 17-									17 17 17	1 2	17(GLY) 16(GLU) 17(PRO)	1. 2.1 3.
	17- 18 19									17	1	17(ALA)	1.
	20 21 22 23	•								17 17 17	1 2	17(SER) 17(LE) 16(SER) 17(CYS)	1. 2.1
	22 23 24									17	1	16(ARG)	<u>1.</u> 1.
•	25									14	2	13(SER) 14(SER) 14(GLN) : 12(GLN) 10(SER)	2.2 1. 1. : 2.3
	25 26 27 27A									14 12 12	1:2 3	10(SER) 12(LEU)	
										12	3	001 510	
	27C 27D 27E 27F									7 2 10	5224	5(HIS) 6(SER) 1(+) 7(ASP): 4(+)	5.7 : 10.
	28									10	9 4:5	B(GLY) 5(ASN) : 3(ASP)	3.8 7.2 : 15. 9. : 12.
	29 30 31 32									. 9	4 1	8(GLY) 5(ASN) : 3(ASP) 4(ASN) : 3(++) 9(TYR) 8(LEU)	9. : 12. 1. 1.
	33 34									8	2		2.7 : 4.
	35 36 37									8 8 8	1 2 2	8(ASN) 4(+) 8(TAP) 7(TYR) 7(LEU) 8(GLN) : 8(GLN)	2.3
	38 39									8 8	- 1 <u>-</u> 2 2	8(GLN) : 8(GLN) 7(LYS) 7(PRO)	1. : 2.3 2.3 2.3
	40 41									8	2 1 1:2	8(GLY) 8(GLY) 8(GLN): 6(GLN) 6(SER)	1 2
	42 43									8 6 7	1	6(SER) 7(PRO)	1. 1.
	44 45 46									· 7 · 7 7 7 7	321	7(PRO) 5(GLN) : 3(+) 6(LEU) 7(LEU) 7(ILE)	4.2 : 7. 2.3 1. 1.
	46 47 48									7 7 6	1	6(TYB)	1
	49 50									6 6 7 7 7	3 4	4(LEU) 3(GLY) 7(SER) 5(ASN)	4.5 8.
	51 52 53										12	7(SEH) 5(ASN)	1. 2.8 1.
	54 55			·						7 7 7	1 2 1	7(ARG) 5(ALA) 7(SER)	2.8 1.
	56 57 58									7	1	7(GLY) 7(VAL)	1.
	59 60									7 7 7 8	2.	7(PRO) 6(ASP) 7(ARG) 8(PHE)	2.3 1.
	61 62 63		•							8	1	B(SEN)	1:
	64 65									8 8 8	2	7(GLY) 8(SER) 8(GLY)	2.3 1. 1.
	66 67 68									8	1 2	7(GLY) 8(SER) 8(GLY) 8(SER) 7(GLY)	2.3
	69 70									777	1 2	7(THR)	1, 1, 2,
	69 70 71 72 73									8 8 8	1	8(LEU)	1.
	74 75 76									8	3	6(LYS) 6(ILE) 7(SER) 8(ARG)	4. 1. 2.3
										8 8 8.	2 1 1		1.
	78 79										22	6(GLU) : 4(+) 7(ALA)	2.7 : 4 2.3 1. : 2. 1. : 2.
	79 80 81 82 83									8 8 8 8	1 : 2 1 : 2 1 : 2	8(GLU) : 8(GLU) 8(ASP) : 6(ASP) 8(VAL)	1. : 2. 1. : 2. 1.
	83 84								·		1	BIGLY) BIVAL	1:
	84 85 86 87									8 8 8 8	1	8(VAL) 8(VAL) 8(VAL) 8(TYR) 8(TYR) 8(CYS)	1. 1. 1.
	88										1 2	7(ME)	1. 2.
	89 90 91 92 93									7 7 7 7 7	1:2 3 2 3	5(ALA) 5(LEU) 5(GLN) : 4(GLN)	1. : 2. 4.2 2.8 4.2 : 5
	93 94									7	5 2	2(+) 6(PAO)	18. 2.3
	94 95 95A 95B 95C									·			
	95D												
	95E 95F 96									3	6	2(TYA) 7(THA)	21. 1.
	<u>97</u> 98									. 7	1	7(PHE) 7(GLY) 6(GLN) 7(GLY) 7(GLY) 7(THR)	1.
	99 100 101 102									7 7 7 7 7	1 2 1 1	6(GLN) 7(GLY) 7(THR)	1. 2.3 1.
3	102 103											5(LYS) 4(+)	4.2 4.
	103 104 105 106									7 8 8	/1 ; 2 /1 ; 2	5(LYS) 4(+) 2 8(GLU) : 7(GLU) 8(ILE)	1. : 2
_	108A 107	۰ 									2	7(LYS)	2.3
	108									7	1	7(ARG) 4(THR)	1.

	52
-	THAT THAT THAT THAT THAT THAT THAT THAT
ANTIBOD	Y SPECIFICITIES: HUMAN KAPPA LIGHT CHAINS SUBGROUP II
P) ROB	COLD AGGLUTININ WITH ANTI-PRID ACTIVITY
	COLD AGELDINN WITH ANTI-BLOOD GROUP I ACTIVITY S: COLD AGELUTINN WITH ANTI-BLOOD GROUP I ACTIVITY ANTI-PHOSPHOCHOLINE(BINDING CONSTANT=6.4X10EXP4)
14) FR:	ANTI-PHOSPHOCHOLINE(BINDING CONSTANT) ANTI-IGG COLD ACCLUTININ WITH ANTI-PR2 ACTIVITY (ABC MEMBRANE ANTIGEN ON HUMAN, RAT AND GUINEA PIG ERYTHROCYTES INACTIVATED BY COLD ACCLUTININ WITH ANTI-PR2 ACTIVITY (ABC MEMBRANE ANTIGEN ON HUMAN, RAT AND GUINEA PIG ERYTHROCYTES INACTIVATED BY
24) GIL:	ANTI-IGO
27) 18:	COLD AGGLUTININ WITH ANTI-PRZ ACTIVITY (NOO MARIANA PROTEOLYTIC ENZYMES AND NEURAMINIDASE)
DECEBEN	ICE: HUMAN KAPPA LIGHT CHAINS SUBGROUP II
	ICE: HUMAN KAPPA LIGHT CHAINS SUBGROUP II
1) TEM	
2) NUL 2) NUM	DREYER,W.J.,GRAY,W.R. & HOODL. (1987) COLD SPAND CHECKED BY AUTHOR 10/18/77) E EULITZ.M. & KLEY,HP. (1977) IMMUNOCHEM14.289-297. (CHECKED BY AUTHOR 10/18/77)
ALCUN	A HILSCHMANN, & CRAIGLIG, THOSE CHECKED BY AUTHOR
.,	HILSCHMANN, (1959) NATURE.35.199-205. (CHECKED & STAUCTURE.309.73-76. 607-7CL: KLOBECK.H.G.SOLOMONA. & ZACHAU.H.G. (1984) NATURE.309.73-76. 107-7CL: MADOFF.M.O. (1972) ATLAS OF PROTEIN SEQUENCE & STAUCTURE.5.0-246. SUBMITTED BY SMITHIES.OGIBSON.D.M. AND FANNING.E.M. 107-7CL: MADOFF.M.O. (1972) ATLAS OF PROTEIN SEQUENCE & STAUCTURE.5.0-246. SUBMITTED BY SMITHIES.OGIBSON.D.M. AND FANNING.E.M. 107-7CL: MADOFF.M.O. (1972) ATLAS OF PROTEIN SEQUENCE & STAUCTURE.5.0-246. SUBMITTED BY SMITHIES.OGIBSON.D.M. AND FANNING.E.M. 107-7CL: MADOFF.M.O. (1972) ATLAS OF PROTEIN SEQUENCE & STAUCTURE.5.0-246. SUBMITTED BY SMITHIES.OGIBSON.D.M. AND FANNING.E.M. 107-7CL: MADOFF.M.O. (1971) ANN.REV.BIOCHEM.40.969-1012. 107-7CL: MADOFF.M.O. (1971) ANN.REV.BIOCHEM.40.969-1012.
5) GM	607-CL: KLOBECK, H.G. SUCUMUNA, G EPROTEIN SEQUENCE & STAUCTURE, S.D-246. SUBMITTED BY SMITHES. O. GIBSON, O.W. AND THE SECUENCE AS TRUCTURE, S.D-246.
6) B A1	CHECKED BY AUTHOR
7) BAT	(CHECKED BY AUTHOR) res: SMITH.G.P.HOOD.L. & FITCH.W.M. (1971) ANN.REV.BIOCHEM.40,969-1012. res: SMITH.G.P.HOOD.L. & FITCH.W.M. (1973) VOX SANG.24.432-440. (CHECKED BY AUTHOR)
8) ROI	B: GERGELY J. WANG, A.C. & FOODSTOOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOLIMMUNOT METHOD WHILE BICK ALL & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOT METHOD WHILE BICK BICK BICK BICK BICK BICK BICK BICK
9) SLC	 CAPRAJD.KEHOEJ.M. WILLIAMS.R.C.JR.FEIZIT. & KUNKEL.H.G. (1972) PROC.NATACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR) CAPRAJD.KEHOEJ.M.WILLIAMS.R.C.JR.FEIZIT. & KUNKEL.H.G. (1972) PROC.NATACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR)
	AUTHOR VOIDS IN THAT AND B C. IN FEIZIT, & KUNKELH.G. (1972) PHOL NAT. ACAD. SCI. SCI. SCI. SCI. SCI. SCI. SCI. SCI
10) Wit	LS: CAPRAJD.KEHOE.J.MWILLIAMS.R.C., JR.FEIZIT. & KUNKELH.G. (1973) FROM THE AUTHOR 10/17/77) FRANGIONE.B.FRANKLINE.C. & PRELLIF. (1976) SCAND.JIMMUNOL.5.823-627. (CHECKED BY AUTHOR 10/17/77)
11) GLI 12) AM	LS: CAPRAJD.KEHOEJM.WILLIMAS.LOUGHT.B.C. A PRELLIF. (1976) SCAND.JIMMUNOL.5.823-627. (CHECKED BY AUTHOR 10/1///) FRANGIONE.B.FRANKLIN.E.C. & PRELLIF. (1976) SCAND.JIMMUNOL.5.823-627. (CHECKED BY AUTHOR 12/12/16/12/12/16/12 FRANGIONE.B.FRANKLIN.E.C. & PRELLIF. (1976) SCAND.JIMMUNOL.5.823-627. (CHECKED BY AUTHOR 12/12/16/12/12/16/12 VILDO TEW: TERRY.W.D.PAGE.D.L.KIMURA.S.ISOBE.T.OSSERMAN.E.F. & GLENNER,G.G. (1973) J.CLIN.INVEST.52.1276-1281. (CHECKED BY AUTHOR VILDO TEW: TERRY.W.D.PAGE.D.L.KIMURA.S.ISOBE.T.OSSERMAN.E.F. & GLENNER,G.G. (1973) J.CLIN.INVEST.52.1276-1281.
	03/02/04)
13) RA	I: MILSTEIN.C.P. & MILSTEIN.C. (1971) CLOSINGHAM
14) FR	 MILSTEIN.C.P. & MILSTEIN.C. (1971) BIOCARLINS, LEUEN DE MILSTRY, 14,1052-1057; RIESEN.W.F. BRAUN.D.G. & JATON.J.C. (1976) PUBLISHED) RIESEN.W. RUDIKOFF.S. ORIOL.R. & POTTER.M. (1975) BIOCHEMISTRY, 14,1052-1057; RIESEN.W.F. BRAUN.D.G. & JATON.J.C. (1976) RIESEN.W. RUDIKOFF.S. ORIOL.R. & POTTER.M. (1975) BIOCHEMISTRY, 14,1052-1057; RIESEN.W.F. (SHECK) BY AUTHOR 1200577) RIESEN.W. RUDIKOFF.S. ORIOL.R. & POTTER.M. (1975) BIOCHEMISTRY, 14,1052-1057; RIESEN.W.F. (SHECK) BY AUTHOR 1200577) RIESEN.W. RUDIKOFF.S. ORIOL.R. & POTTER.M. (1975) BIOCHEMISTRY, 14,1052-1057; RIESEN.W.F. (SHECK) BY AUTHOR 1200577) RIESEN.W. RUDIKOFF.S. ORIOL.R. & FOTTER.M. (1976) BIOCHEMISTRY, 15,3829-3833. (CHECKED BY AUTHOR 1200577) RUBAS, C., TUNG.E., WANGJ., FUDENBERG.H.M., PICK.A.L. & FROEHLICHMAN.R. (1980) CANCER IMMUNOTIMER., 381-86. (CHECKED BY AUTHOR 03/10/01) WANGA, C., TUNG.E., WANGJ., FUDENBERG.H.M., PICK.A.L. & FROEHLICHMAN.R. (1980) CANCER IMMUNOTIMER., 381-86. (CHECKED BY AUTHOR 03/10/01) WANGA, C., TUNG, E., WANGJ., FUDENBERG.H.M., PICK.A.L. & FROEHLICHMAN.R. (1980) CANCER IMMUNOTIMER., 381-86. (CHECKED BY AUTHOR 03/10/01) WANGA, C., TUNG, E., WANGJ., FUDENBERG.H.M., PICK.A.L. & FROEHLICHMAN.R. (1980) CANCER IMMUNOTIMER., 381-86. (CHECKED BY AUTHOR 03/10/01)
	S: WANG.A.C. TUNG.E. WANGI, FUDENBERG, FUSION STATES TO BE LEDER P. (1980) CELL, 22, 197-207; KLOBECK, H.G.MEINDLA. COMBRIATO.G.
17) MA	N: MILSTEIN,C. (1969) PROC. 5TH FEBS SYMP.15.43-56. (CHECKED BY AUTHOR 12/05/77) R: SLETTEN,K.,HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77)
18) KI	R: SLETTENK, HANNESTAD,K. & HARBOE,M. (1974) SCAND.JIMMUNOL.3.215-218. (CHECKED BY AUTHOR 12/05/77) L: SLETTENK, HANNESTAD,K. & HARBOE,M. (1974) SCAND.JIMMUNOL.3.215-218. (CHECKED BY AUTHOR 12/05/77) L: SLETTENK, HANNESTAD,K. & HARBOE,M. (1974) SCAND.JIMMUNOL.3.215-218. (CHECKED BY AUTHOR 12/05/77)
19) HY	L: SLETTEN,K. HANNESTAD,K. & HARBOE.M. (1974) SCAND.JIMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77) AG: SLETTEN,K. HANNESTAD,K. & HARBOE.M. (1974) SCAND.JIMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77)
20) MA	AG: SLETTEN.K.,HANNESTAD.K. & HARBOE.M. (1974) SCAND.JIMMUNOL3,215-218. (CHECKED BY AUTHOR 12/05/77) E: SLETTEN.K.,HANNESTAD.K. & HARBOE.M. (1974) SCAND.JIMMUNOL3,215-218. (CHECKED BY AUTHOR 12/05/77)
20 EU	E: SLETTEN,K. HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77) D: SLETTEN,K. HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77) D: SLETTEN,K. HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL3.215-218. (CHECKED BY AUTHOR 12/05/77)
221 04	MUNDE MILSTEIN, C. JAHVIS, J.M. & MILO (CHARGE CHARGE) IMMUNOLOGY, 35,447-453. (CHECKED OF HOTHER
24) GI	L: ABRAHAM.G.N., BHOWN, P., JOHNSTON, SCIENCE AND LIMMUNOL 3,215-218. (CHECKED BY AUTHOR 1205/7)
25) MI	EN: SLETTEN.K. HANNESTAU.K. & HANDOUTINE THORAGE (CHECKED BY AUTHOR)
261 61	SEON B.K. YAULY, & PRESSWORKS, (1014)
071 71	A CEPCELY J WANGAU & FUDENDERIGANIN (TTT)
28) 51	AT: SLETTEN,K.,HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL.,3,215-218. (CHECKED BY AUTHOR 12/05/77) Λ: SLETTEN,K.,HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL.,3,215-218. (CHECKED BY AUTHOR 12/05/77) Λ: SLETTEN,K.,HANNESTAD,K. & HARBOE,M. (1974) SCAND.J.IMMUNOL.,3,215-218. (CHECKED BY AUTHOR 12/05/77)
29) LL	IT: SLETTEN.K.HANNESTAD.K. & PARDOLINI, (1973) NATURE NEW BIOLOGY,246,176-178. DB2: MOULIN.A. & FOUGEREAU.M. (1973) NATURE NEW BIOLOGY,246,176-178.
30) H	DB2: MOULIN.A. & FOUGEREAU.M. (1973) NATURE NEW BIOLOGY,246,176-178.
NOTES	HUMAN KAPPA LIGHT CHAINS SUBGROUP II
IDENTI	CAL SETS OF FRAMEWORK SEGMENTS:
FRI	
FR	2: SET 1: MIL[2].NIM[3].GM 607 CL[3]. (3 IDENTICAL SET 2: MIL[2].FR[14]. (2 IDENTICAL)
FR	THE AND TO THE POINT AND CLUBE IZ WENTIONE TO THE AND THE WAS SOMETHING TO THE AND THE WAS SOMETHING TO THE AND THE
FR	4: SET : GM BU/ DEDI. WEIZI DEVIJ DAVIJ DEVIJ GLOTISI CURIZOL HEISI // VARPA III. VARPA III. WEIZI DEVI GLOTISI CURIZOL HEISI // VARPA III. NEUISI.
•	SET 2: NIM3],FRI 19, (2: IVEN TIGAL PURCHATCH 24) CL (28); AND 1 HUMAN V-KAPPA-IV: LENGED; GOTGIG,GARTIOL,FLOT[2],FR4(21),JARGED 24 VICE 20(102),JUIGO(134).
	SET 3: TEWITI, (IDENTICAL TO 2 HUMAN V-KAPPA-R WALKEN OLITION TO A THE AND A
IDENT	ICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS:
	R1: R2: SET 1: MIL[2].NIM[3].GM 607 'CL[5]. (3 IDENTICAL)
	83:

UDH3: IDENTICAL SETS OF J-MINIGENES: SET 1: RPM1-6410°CL[16]. (IDENTICAL TO 1 HUMAN V-KAPPA-I: AU[2]; 2 HUMAN V-KAPPA-III: PIE[11],VKAPPA3°CL[62]; AND 1 HUMAN V-KAPPA-IV: PB17IV°CL[3]). SET 2: TEWI11. (IDENTICAL TO 1 HUMAN V-KAPPA-I: WALKER°CL[12]). SET 3: FR[14]. (IDENTICAL TO 2 HUMAN V-KAPPA-I: DEN[46],Bi[63]; AND 3 HUMAN V-KAPPA-III: GAR'[10],FLO[12],JARC/BL41°CL[26].)

SPECIFIC NOTES:

SPECIFIC NUTES: 12) AMYLOID TEW: IT HAS THE SAME SEQUENCE AS THAT OF TEW SO FAR AS THE SEQUENCED POSITIONS ARE CONCERNED. 14) FR: AN IDIOTYPIC ANTIBODY TO FR NOT INHIBITABLE BY PHOSPHORYLCHOLINE REACTED BETTER WITH THE FR HEAVY CHAIN THAN WITH THE LIGHT CHAIN. THE CROSS-REACTION WITH MOPCIE? WAS 10,000 TIMES WEAKER. (RISEN.W.F. (1979) EUR.JIMMUNOL.9.421-425.) 16) RPM1-6410°CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN ADULT DNA.

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

AT POSITION	AESIDUES
27F 28 31 34 45 79 94 104	(GLY,ASN) : (GLY,ASP) (ASP,ASN) (THR,ASP) (ASP,ASN) (GLU,GLN) (GLU,GLN) (THR,SER) (LEU,VAL)

and Street of

									•																	
АМИН	12	PPA LIGHT	CHAIN 1 11	NOL	3.	P III NG9 CL	5". NEU	607	7. PAY	8ª SON	9.' 9'	10 GAR	11' PIE	12" FLO	13- LOP	14 · SCA	15° GLO	15 SAL	17 WIL	18 · MA	19, NIC	20 · CUR	21 FR4	22" DRE	23 PER	24 . CAM
	4 5 T 6 7 5	'AL(.96) HR SER 'PO	GLU ILE VAL LEU THR GLN SER PRO GLY	GLU ILE VAL LEU THR GLN SERO GLY	GLU ILE VAL LEU THR GLN SER PRO GLY	# GLU ILE VAL LEU THR SER GLY	GLU ILE VAL LEU THA SEA PRO GLY THA	GLU ILE VAL LEU THR GLN SER PRO GLY THR	GLU ILE VAL LEU THR SERO GLY THR	GLU ILE VAL LEU THR GLN SER PRO GLY THR	GLU ILE VAL LEU THR GLN SER PRO GLY THR	GLU ILE VAL LEU THR GLN SER PRO GLY THR	GLU ILE VAL LEU THR GLN SER PRO GLY THR	GLU ILE VAL LEU THR SERO GLN THR	GLU ILE VAL LEU THR SER PRO GLY THR	GLU ILE VAL LEU THR SEA GLN SEA GLN THA	GLU VAL LEU THR GLN SER PRO GLY THR LEU	GLU ULE VAL LEU TGLN SERO GLY TLEU	GLU ILE LEU THR SERO GLY THER SERO THEUR	GLU ILE VAL LEU THR GLN SERO GLY THR LEU	GLU ILE VAL LEU THR SER PRO GLY THR LEU	LEU	VAL LEU THR GLN SER PRO GLY THR LEU	GLU ILE VAL LEU THR SERO GLY THR LEU	GLU ILE VAL THR SEROY GLY THR LEU	GLE VAL VAL TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TGLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLANA TCLA
F .	14 5 14 5 15 F 16 0 17 18 19 20	EU(.99) SER SER(.97) PRO(.98) SLY	THR LEERU SERU GLEGR GLEGR GLEGR ALL CLEGR LEU	THRURSER LEER OY GLUG ALG ALA LEU	THR LEERUR SEOR GLUG ARGA ALA THRU	THRU LEERUR SELUR GLUG ALLA SER LEER SER	LEU SER LEU SER GLV GLU ARG ALA THR LEU	LEU SER LEU SER PRO GLU ARG ALA THR LEU	LEU SER GLU GLU ARG ALA THEU SER	LEUR SER SEV SER GLU SER GLU ARG ALA THRU	LEU SER PRO GLU ARG ALA THR	LERUR OYUGA SEUR OYUGA AL HEUR	THRUSER LER PROY GLUG ALA THE SER	LEUR SEU SER GLU GLU GLU GLU ARA ALA TLER	LEER OY UGA A THER	ALA THR LEU SER	SER LEUR PROY GLUG ARA ALA THEU SEA	SER LEU SER GLU ARG ALA THR LEU SER	SER LEU SER GLU GLU GLU GLU GLU GLU GLU GLU GLU SER CYS	LEUR SEUR PRUS GLUGA ALA HEURS CYS	SER LEU SE O GLU ARA ALA LEURS CYS	SER LEER PROY GLUG ARLA THEU SERS	SER LER PGLUG ARLA THEURS CYS	SER LEU SER GLY GLX ARG ALA THR LEU SER CYS	SER SEUS PROY GLX ARG ALA THRU SER CYS	SER PRO GLY GLU ANG ALA THR LEU SER
	22 23 24	ALA(.98)	SER CYS ARG ALA SER	SER CYS ARG ALA SER	SER CYS ARG ALA SER	ARG ALA SER GLN	SER CYS ARG ALA SER GLN	SER CYS ARG ALA SER SER	ARG ALA SER GLN	ALA SER GLN	SER CYS ARG ALA SER GLN SER	SEA CYS ARG ALA SER GLN SER	CYS ARG ALA SER GLN SER	CYS ARG ALA SER GLN	CYS ARG ALA SER GLN	ARG ALA SER GLN	ALA	CYS ARG ALA GLY SER	ARG ALA SER GLN SER	ARG ALA SER GLN SER	ARG ALA SER GLN SER	ARG ALA SER GLN	ARG ALA SER GLN SER	ARG ALA SER GLX SER	ARG ALA SER GLX	ARG ALA SER GLN SER
C DR 1	27A 277B 277C 277C 277E 277E 277F 28 29 30 31 322 33 32 33 34		GLER 	VAL	VAL SER SER TYR	LEU ALA	SER ARG TYR LEU ALA	VAL AAG SEA SEA TYR LEU		VAL SEAR	VAL SEA SEA TYP	VAL SER SER LEU	VAL SEA SEA TYAL	VAL SEF SEF LEL AL/	VAL SEF	SEP SEP ASM TYP LEU	SER SER SER TYR LEU ALA	SEP VAL SEP VAL ALA TRF	SER ASN SER MET LEU ALA	, .			ALA TRP	LEU ALA		VAL SEAR SEAR TAP TAP
F # 2	35 36 37 38 39 40 41 42 43 44 45 46 47 48	GLY(.96) LEU(.96) LEU(.96)	TRIF GGLU GGLU PRI GGL AR AR LEELILI TY	TYPE GLI GLI PAC GLI PAC GLI CO GLI CO CO CO LEI LEI LEI	TYRE TYRE TYRE TYRE TYRE TYRE TYRE TYRE	TYPE GLASSO PRO GLASSO PRO GLASSO PRO GLASSO PRO GLASSO PRO GLASSO PRO GLASSO PRO LEL	TYPAN GOLY SOLVA DOUL	TYPE	TYPEN GLIN GLIN GLIN GLIN GLIN GLIN GLIN GLIN	TYPELL GLYRCLUL GLYRCLUL GLYRCLUL GLYRCLUL GLYRCLUL GLUL GLUL GLUL GLUL GLUL GLUL GLUL	TYPEN GLIYER	TYPEN GLA GLASS PRODUCED SOPRESSON ALL SOPRESSON ALL SOPRE	TYPH GLA GLY GLY GLY GLY GLY GLY GLY GLY GLY GLY	TYPE GLI YELE Y	N GLIYO PRO PRO GLIYO GLI GLIYO CALI GLIYO CALI CALI CALI CALI CALI CALI CALI CALI	N GLI N GLI S PRO Y GLI N GLI A O	N GLN N GLN LYS		i GLN			GLN GLS PRC GLN GLN ALA ARG LEL LLE TYP	GLN GLN ARC PRC GLN GLN ALA PRC			GLN
CDR2	49 50 51 52 53 54 55 56	ARG(.95)	VA AL SE SE AR AL TH	A AL R SE R SE G AR A AL	Y GL A AL R SEI R SEI G AR A AL R TH	Y GL A AL A TH A SE G AR AL A TH	Y GL A AL A SE R SE G AR A TH	A AL/ R SE R SE G AR A AL R TH	A AL R SE G AR A AL R TH Y GL	A AL R SE R SE G AR AL TH Y GL	A R G A R	GL AL SE SE AR AL TH	A AL R SE G AR A AL Y GL	A AL R SE R SE G AR A AL R TH	A R G A R		AL/ SEI AR AL TH GL					GL ALA SEF SEI ARC ALA THI GL	A AU A AU F TH Y GL	Pi Y E		
5 F	57 58 9601 623 634 655 667 68 9701 772 733 745 669 771 778 90811 833 8455 8455 8455 8455 8455 8455 8455	GLY(.96) PRO ARG GLY SER(.95) GLY SER(.95) GLY THR(.95) PHE THR(.95) PHE THR(.95) ILE(.95) GLU(.95) ASP ALA VAL(.95) TYR	PAS GISGSIG TI APPTLL T - SGLL GPGAP X VI	E OPGER YRYRRYRRYRRU RERYU UOUPE ALRR	E OPGER YRYRYRPERU RERYU UOUPE ALAR	L PARAPHE LESALEG TASHTLE TILSAL GPGAP AVIT	OPGER YRARY ASTAL TVSAL GPGAP AVTT	OPGER YRYRY RPERD RLEGU LOUDE ALYR CARPS GSGSG TAPTLE THISRL GPGAP AVTT	E OPGER YRYRY RPERU RERGU UOUPE ALAYR	OPORER YRYRY RPERU RERGU LOUPE ALAR Oporer yryr rperu rergu Loupe Alar		THUSIALL GPGAP AVTT	OPGER YRYRY RPERD RERGU LOUPE ALYR	OPGER YRYRY RPERU RERGU UOUPE ALAYR			RAAPS GSGSG TAPTL TUSAL GPGAP AVTTC	ALANT HEROTO NOUTE HERE A LEVEL A LEVEL					PGER YRYRY RPERU REAGU VOUPE ALRAS Apps Gsgsg Hapte Thistee Gipgaap Avtpic	PGER YRYRY RREAU REAGU LOU		
C D R S	95 95 95	A IB IC		SER I SER I SER I SER I	YS CONTRACT	LN GGTGA	LN GOTTON SERVICE	LN GG YR T LN GG YR T LN G YR T SLA S FRO FRO T	LN GUNNER	ILN CONTRACTOR	LN YR SER SER ROOR T	GOTOS: SF	LN GETCH	LN CONTRACTOR	LN SLA SLA SLA SEA SEA SEA SEA SEA SEA SEA SEA SEA SE		GTGS SP	LN LN YR YR E R R C C C C C C C C C C C C C C C C				GTGS SP	LN GT GG SP			
ş	95 95 96 97 91 90 100 100 100 100 100 100 100 100 1	5 5 7 8 9 9 9 0 1 1 2 2 3 4 5 6		THR GLY GLY GLY THR LYS VAL GLU LEU	ARG C THA T GLY C GLY C GLY C GLY C GLY C THA S LYS C UAL C GLU C	SUN HR SLN SLN SER SLYSL VAL VAL ULE		CYS A	ARG IHA GLY GLY GLY GLY THA LYS LYS	LEU THR GLY GLY GLY THR LYS VAL	TYR THR GLY GLY GLY THR LYS VAL GLU ILE	1	THR THR SLY SLN SLY THR LYS LEU GLU	TRP THR GLY GLN GLY THR LYS	TYR THR GLY GLY THR LYS LEU ILE			EU HR SLY SLY HR LYS ILR SLU ILE					HA P HE F SLY (SLN (SLY (SLY (NAL 1 SLU (ILE)	ALN ARO ALY ALLY ALLY ALLY ALLY ALLY ALLY ALLY		,
	10 10 10	5A 17 LYS(.9 18 ARG	5)	LYS ARG THR	LYS	LYS ARG	 	LYS	LYS ARG THR	LYS	LYS ARG THR		LYS	LYS ARG THR	LYS ARG THR			LYS ARG THR			,		YS ARG (HR	LYS		

.

.0	AN KJ	25° STE	26* GJ		28 IARC/ BL41 'CL	29 RAD	30 DIL	31 CAS	32 MCE' #	33 KEA	34 SMI	35 <u>*</u> AJ	36 BRO 166	37 NIG	38 IKE	39 TIL	40 AMYLOID KSA	41 POL	*	43- SHE #	44 JH #	45 WIN				VAND
	0 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 15 6 7 8 9 20 1 2 2 2 2 3	GLU ILALU RINAROY THEARUR OYLUGA ALUR SER GLUGA ALUR SER	GLUE ANNA SARA A	GLE TALARDY RURAUA OYXGA AU TALARDY THERE A CYNGARAU TALARDY RURAUA CYNGARAU TALARDYN	GILLU ANAGOY AURA AURA GILALU ANAGOY AURA AURA GILSPAC AURA AURA SPACES ALL SCIENCE	GLE GLE VAL HUNROY TGSERUY TLEEUR OY SEE POLY SEE OY SARA AU THEERS CARDON AND AND AND AND AND AND AND AND AND AND	GLUE ANAROY RURUR OY POON AURS	GLEAU ANA OY RURUR OY PGA RURUR OY PGA RURAL SPAL HERE OY PGA RURAL SPAL HERE	GILALU ANROY AUA SA COLLAR COLLAR AND A COLLAR AND A COLLAR AND A COLLAR AND A COLLAR A COLLA	GLELU RNROA RURUR OYUGA RURS GLELU TGSPRA TUSUS PGGAAA TUSUS ARUSUS PGGAAA TUSUS ARURS	GILELU RANAROA RURALA HURALA H	GUELU ANAGO AURUA OYUGA RURSIG GUAL HURAGA HURUA OYUGA RURSIG ALSUS PGGAAA THEEYG A	GLELU RUADY RUADY AURILS POLAL THUR SPAL THUR OSPAL THUR OGAAL HUR ARG	INSELLU ANAROY RURUR OYUGA RURUR ILSELS PGGRAA TLEE	GILALU ANARO ARURUA OYUGA AURS	GUELU HNRO I AURUR OYUGI AURUS OGUAL AURON I AURUS OGUAL AURON I AURUS OGUA TUSY OGUA	OLAN RNAOL AURIAR OYUGA AURSON TUSAN	ULELU HUROA RURUR OYXGA AURSY SPRATESEE RUXGA AURSY GGAAA HEESY AR	GLEL UNANT HERA RUALR OYUGA RUARS VE RALLE OYUGA RUARS A THEESE OYUGA RUARS	GLUEL HRNRRO # RURAL ROY USE OF THELES OF RULUGA RURAL RUBAL RURAL RURAL RURAL RURAL RURAL RUVAL RUPAL	GLELL ULAL RANA RUR OYUUGA RUR THER RANA THEESE ROYUUGA RUR THESE RAL RUR THE	SPELU RNRO RURUR OYUGA RUR THLEESLE RLLUGA RUR	IVSELLU RXRAO A THEESER RUXGA HURSE A HURSE ROLXGA HURSE A HURSE ROLXGA HURSE A HURSE	GONAL ANAROY RUADA OYU A BURS G	GLEVER RNAROY THERE ROYUGAAL HEERS GAAL	GLELLE HANA AND A CONTRACT OF A CONTRACT ON A CONTRACT OF
	24 25 26 27 27A	•	ARG ALA SER	ARG ALA SER GLX SER	ARG ALA SER GLN	ALA SER GLN VAL	ALA SEA GLN SEA		ALA SER GLN	ALA GLY SER		ALA SER GLN SER	ALA SER GLN SER		ALA SER GLN SER	ALA SER GLN	ALA SER GLX THR	ALA SER GLX	ALA SER		ALA		ALA SER GLX	SER GLN	SER GLN SER	GL
CDR 1	27B 27C 27D 27E 27F 28 29 30 31 32 33			VAL SER SER TYR LEU	SER VALA SER VALA SER VALA	SER SER SER SER SER SER SER LEU	LEU ASX SERS LYSA LEU		SER	ASX ALA ALA SER LEU ALA		VAL SER		·	VAL SERN SERN SERN LEU ALA	SER VAL SER SER TYR LEU	 LÉU	SER VAL ARG TYR LEU ALA			-		SER VAL SER LEU TYR LEU ALA	HIS VAL ILE ALA GLY TYR	ILE SER ASN SER TYR LEU ALA	AS ILI SES LE ALE TF
F R 2	34 35 36 37 38 39 40 41 42 43 44 45 46 47			ALA	ALA TRPANS GLYSGUNA GLYSGUNA GLYSGUNA CLED	ALA TRP TYR GLN GLN GLN GLN GLN ARG ARG	LYS PRO GLY GLX THR PRO ARG LEU			THP TYR GLX GLX LYS PRO			TAP TYR GLN GLN LYS PAO		TRP TYR GLN GLN ARG PRO	TRP TYR GLN	TGGL ROYNA GGL SEAUNA GGLA PRAEUU GGL PRAEUU LLE							TAP TYR GLN GLN LYS PRC	LYS PRO SER GLY SER PRO ARG LEU LEU	
	48 49 50 51 52 53				ARG ASP ALA SER SER		ILE TYR								•		GLY GLY THR SER THR								TYR GLY ALA SER THR ARG	
CDR2	54 55 56 57				ARG ALA ASN GLY	ALA THR GLY											ARG ALA THR GLY ILE								ALA THA GLY	
F A 3	5 59068123 4568 6901223 456677 7901223 456688 88888888888888888888888888888888				E OPGHER YRYRY RPHERU LELERGU JOUJPE ALMRRY Raare Gsegse taarti llesreu Joujpe Almrry Lilsaai opgaa Avytyd	TH PARGER YRURY REPERU REFOUNDED AND ANTES											PRO ALA ARG PHE SER GLY						•		PALAGEE LIEUTEL HELEFEL LELISUUTELISH JAYYYYYUUUU	NAME AND ALTER A MARCH ALTERS NAME
CDRS	89 90 91 92 93 94 95 95 95 95 95 95 95 95 95 95 95 95 95				GLNAFAFAF GULNAFAFAF SEAC											·							-		ASI ASI PRI PRI PRI PRI PRI PRI PRI TH	NN PO
FR 4	97 98 99 100 101 102 103 104 105 106				THE PHE GLI GLI GLI THE LYS LEU GLU	R TH PHI N GL N GL N GL N GL N GL N GL N GL N GL					<u>. </u>														PH GLL GLL AR VA GLL LY	

E-VIELONG

	50 AMYLOID SO124	51 DOV	52 SHM	53 GRA	54* GOE 11	LOW.	56 VER	57 REE	58 WE #	59 ноw	60 HS4	61 HBJ 5	62. TEH	63' CRA (III)	64 PLA	65" PIN	66 MCE	ь7 HAC	68 K EV15 CL	69 BER	70- BOR	DRI	WAL	73. GOL	ĠĂG
0 1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 6 7 8 9 10 11 12 3 4 5 10 11 12 3 4 5 10 11 12 3 4 5 10 11 12 3 4 5 11 12 3 4 5 10 11 12 3 11 12 3 4 5 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 11 12 3 11 11 12 3 11 11 12 3 11 11 12 3 11 11 12 3 11 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 11 2 3 11 12 3 11 12 3 11 12 3 11 12 3 11 11 2 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 12 3 11 1 12 3 11 1 12 3 11 1 12 3 11 1 12 3 11 1 1 12 3 11 1 12 3 11 1 12 3 1 1 1 1	GILELABI ANAROS GILY M TGERBAR OLUGA TGESPASTALSES PGGARS HUAR SVS PGGARS TLEP	GLU LAL AND	GLE VAL THERO AL A THERA SPRA THEER OY UNA THEER OY UNA THEER OY UNA THEERS	GLU: WALL THAN SEAD THEUR THAN SEAD THEUR SEAD SEAD SEAD SEAD SEAD SEAD SEAD SEAD	aspelu ANRA aspeluation of the second	GLE HAN ROAD AND A CONTRACT A CONTRACTACT A CONTRACTACT A CONTRACT A CONTRACTACT A CONTRACTACT A CONTRACTACT A CONTRACTACT A CONTRACTACT A CONTRACTACTACTACTACTACTACTACTACTACTACTACTACTA	GLUT LU RNRROY RURA OY UL TURALU RNRROY RURALU RURALU RURALU RURALU RULALU RULA	GLUURALU RNROA RURAL OYU HANROA RURAL SOU TUSO	aspe all gint TGESPRO SEERA SEERA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA SUBRA	GLU RUR GLU RUR GLU TGLEROY RULE HNN RGEROY RULE RUR GSPRGL RUL SPRCH RUR GLU	GLU GLU VAL TGLEROY GSEROY GLU THEUROUR SPRCY THEUROUR SPRCY THEUROUR SPRCY THEUROUR SPRCY THEUROUR SCOUPSING	GLE ALU ALURALU ASX RURALU ASX RURALU ASX RURALU ASX RURALSE AGLX	GLELU ANROY AURUR TGSPRGL TLEERER O	GLUE VALU RUNRUS GLUE TGLEROY TGLEROY TLEERUR LSECRO	GLU ILE VALUER GLER GLER SEEO GLY SEEN SEEN SEEN SEEN SEEN SEEN SEEN SEE	GLE VALU ANNROY TGLERUS THERUS SPACE THERUS SPACE SEER SPACE SEER SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE SPACE	GILE ANAROY AUL TGLERU VAL TGLEROY AULAEU	GILE VALU VALU THRNA SEROY THEU	Ghr thr L THRNA GLEAD Blaeet A Blae Blae Blae Blae Blae Blae Blae Blae	GLU GLELUALU LEUTHRALU THRALUAROS FROY GLA	GLU ILE VAL THR SERO ala THR	GLU GILE VALU THA GLNA PRO	GLU GLE VAL LEHR TGLR PRO	GLE VALU R VALU	GLU ILE VAL LEV GLN SER
223 24 25 26 27 27A 27B 27C 27E 28 29 30 31 32 33 33	SEYS ARG ALA	ARG ALA SER GLU SER GLU SER SER SER SER SER SER LEU ALÀ	ARG ALA SEA GLU		CYS	SER ALLY SER ALLY GLI IIIIISER VASLAR VASLAR VALA	CYS LYS SER LEU SER LEU SER	ARG ALA SERU SERU SERU LER ASSP VAL	ALA SER GLN THR GLY				•						CYS LYS ALAR SELN IIII ASP MET ASP MET ASP		-				
35 367 3 90 40 40 40 40 40 40 40 40 40 40 40 40 40		TRP TYR GLX GLX LYS PRO	TRP			TRP TYR GLX LYS PRO	TRP TYR GLN GLN LEU LEU ILEU ILEU ILEU	ARG LEU LEU TYR ALA		~									THP TYAN GLNN GLUS GLUG ALA ALA ILEE ILE GLUG ALA						•
50 51 52 53 54 55 56 57 58 59							VAL SER THR ARG ALA GLY ILE PRO	SEA THR ARG PRO THA ASP ILE PRO											THA THA LEU VAL PRO GLY ILE PRO						<u> </u>
59 661 663 665 666 665 666 665 667 68 665 667 772 776 776 776 88 88 88 88 88 88 88 88 88 88 88 88 88							AAPT GAAASG TAAPTL TUSAL GPGAAP ALAPPS AAPT GAAASG TAAPTL TUSAL GPGAAP ALAPPS	AARBER AARBER S GEL RLURA S GEL RLURA S GEL RLURA S GEL RLURA S GEL RLUPA S GEL RLUPA S GEL RLUPA S GLUPA S GL							-	-			DOGUER LARYARY RAPURAU HENRORU UDEURA A ARTARUSULA DARHE LEEUYA HONHAU HENRORU UDEURA A ARTARUSULA DARHS GSGTG TAARTU TILASAI GSGAAA ATTARO LGH GSGAA ATTARO LGH			•			
89 901 93 954 955 955 955 955 955 955 955 955 955			-		-				3										ASP ASN PHE PRO		•				
98 99 100 101 102 103 104 105 106 106A 107							PHE GLU GLU THF LYS LEU ASF VAI	PHE GLY GLM GLY GLY CLY CLY CLY CLY CLY CLY CLY CLY CLY C						. –											

MAN K	APPA						01	82	# OF	# OF	OCCURRENCES	VARIABILITY
	75 DOB	76 HS6	77 HÐJ 12	78 BUR (K)	79 LEG	80 86 AI	WR #	VKAPPA3	SEQUENCES	AMINO ACIDS	OF MOST COMMON AMINO ACID	
0 1 2	GLU	GLU	GLU	GLU	GLU				79 79 79 79	3:4 5 4	74(GLU) : 73(GLU) 74(ILE) 76(VAL)	3.2 : 4.3 5.3 4.2 3.6
3	ile met	LEU	VAL VAI THR	LEU	VAL LEU				79 77	3	65(LEU) 77(THR) 75(GLN) : 69(GLN) 75(SER)	2.1 : 2.2
5 6 7	THR GLN SER	THR GLN	GLN					•	77 77 75 74 69	21	74(PRO)	1. 1. 9. : 11.
8 9 10	PRO ale								69 70 68	6:7 .4 2	46(GLY) 66(THR) 67(LEU)	4.2 2.
11 12 13									67 67 68	1 5 2	67(SER) 52(LEU) 64(SER)	1. 6.4 2.1
14	-								58 62 62	2	65(PRO) 62(GLY) 56(GLU) : 50(GLU) 51(ARG)	2. 1. 3.3 : 5.
16 17 18						ARG			62 58 60	3:4 7 2	52(ALA)	8. 2.3
20						ata			59 60 60	523	53(THA) 57(LEU) 58(SER)	5.6 2.1 3.1
21 22 23	<u> </u>					LEU SER CYS ARG			50 51	4	50(CYS) 47(ARG)	4.3
24 25 26						ALA SER			52 49 47	223	51(ALA) 46(SER) 43(GLN) : 37(GLN) 29(SER)	2. 2.1 3.3 : 3.8
27 27A 27B						SER			32	4	29(SER)	
270												
27E 27F 28						LEU			47 44	7:8 6	25(VAL) 27(SER) 24(SER)	13 : 15. 9.8
29 30 31						SER GLY ASN TYR			40 39 40	7 10 8	24(SER) 24(SER) 28(TYR)	12. 16. 11.
32 33						LEU			41	4 5	36(LEU) 37(ALA)	4.6 <u>5.5</u> 1.
34 35 36						TRP TYR GLN	TRP TYR GLN		38 39 39	1 : 2 2 : 3	38(TRP) 39(TYR) 39(GLN) : 33(GLN) 36(GLN) : 30(GLN)	1. 2. 2.1 3.
.37 38 39						GLN	GLN		39 37 33	2:3 3 3	29(LYS) 32(PBO)	3.4 3.2
40						PRO GLY GLN	GLY		34 27 27 26	2 4 3	26(GLY) 24(GLN) : 23(GLN) 23(ALA)	3.4 3.2 2.1 4.5 : 4. 3.4
43						PRO	ALA PRO		7	з	25(PBO)	3.2 3.3 2.1 2.1
45 46 47						ARG LEU LEU MET	LEU LEU ILE		26 24 23 22	3223	24(ARG) 23(LEU) 22(LEU) 20(ILE)	3.3
48 49 50						TYR GLY	PHE ASP		. <u>22</u> 21 20	4 5 3	19(TYR) 16(GLY) 16(ALA) 18(SER) 16(SER)	4.6 6.6 3.8
52						VAL SER SER	SER		20 21	2	18(SER) 16(SER) 19(ARG)	3.8 2.2 2.6 2.1
54 55						ARG ALA THR	ALA		20 23 22	2 3 4	21(ALA) 19(THR) 22(GLY)	3.3 4.6 2.1
56 57 58						GLY	GLY VAL	PRO	23 23 23	23	21(ILE) 23(PBO)	3.3
59 60 61						ASP		ASP ARG PHE	23 23 23	5	17(ASP) 23(ARG) 23(PHE) 21(SER)	6.8 1. 1.
62 63						PHE SER GLY		SER GLY	23	2	21(SEA) 23(GLY) 21(SEB)	2.2 1. 2.1
64 65 66 67						SER		SER ALA SER GLY	22 22 22 22 22 22 22 22	. 2	23(GLY) 21(SER) 17(GLY) 21(SER) 22(GLY)	5.2 2.1 1,
68						SER GLY ALA		GLY THR ASP	22 22 21 21	· 2 2	21(THR) 19(ASP)	2.1 2.2 1
69 70 F 71 8 72								PHE THA LEU	21 21 21	1 1	21(PHE) 21(THR) 21(LEU)	1.
3 73								THA	21 21	2 2 3	20(THR) 20(ILE) 19(SER)	2.1 2.1 3.3
75 76 77						ARG LEU		SER ARG LEU	21 22 22	5 3	16(ARG)	6.9 3.3 2.1 : 2
79 80 81						GLX		GLU PRO	22 22 22 22	2 2 2 1 3	21(GLU) : 20(GLU) 19(PRO) 21(GLU) 21(GLU) 22(ASP) 20(PHE)	2.1 : 2 2.3 2.1
81 83 83						GLU ASP PHE		GLU ASP PHE	22	3	22(ASP) 20(PHE) 22(ALA)	
84 85	5					ALA VAL TYR		ALA VAL TYR TYR	22 22 22 22 22 22	2 1 2	22(ALA) 21(VAL) 22(TYR) 20(TYR) 22(CYS)	2.1 1. 2.2
8 8 8	3				·	CYS GLN		GLN		1	22(CYS) 21(GLN) 22(GLN)	2.1
8 9 9	1					GLN GLN TYR GLY		GLN TYF GLY	22 22 22	2 1 2 5 5	21(GLN) 21(GLN) 22(GLN) 22(TYR) 16(GLY) 12(SER)	1, 2.2 6.9 6.8
999	3					SER SER PRO		ASN SEP GLN	21	. 3	18(SER) 18(PRO)	4.7 3.5
D 9	5 5A 5B								1	1	i(PRO)	
9	5C 5D											
9 9 9	5E 5F 6 7					PHE THR			1 20	10 2 1	20(2HE)	48. 2.1 1. 1.
9	8 9					PHE		PHE GL GL	20 20 20	122	20(GLY)	2.2
10 10 F 10	2					GLN GLY SER		GL' THF LYS	{ 20	2		1. 2.2 2.2
A 10	4					LYS LEU GLU		GL	J 20		18(LYS) 11(VAL) 18(GLU) 18(ILE)	3.6 2.2 3.3
10	6 6A					LYS		۱L E L Y	· 20	2	19(1 YS)	2.1
10								AR	G 16 10		16(ARG) 10(THR)	1.

.

-		
	57	
	SPECIFICITIES: HUMAN KAPPA LIGHT CHAINS SUBGROUP III	
2) WOL:	ANTI-HUMAN GAMMA G GLOBULIN; WA IDIOTYPE	
E1 412711.	ANTI-HUMAN GAMMA G GLOBULIN; WA IDIOTYPE CRYOGLOBULIN WITH ANTI-IGG ACTIVITY: B IDIOTYPE (KUNKELH.G. WINCHESTER.R.J. JOSLIN,F.G. & CAPRA.J.D. (1974) J.EXP.MED. 139.128)	
C) COT	CONOCIONIUM WITH ANTHIGG ACTIVITY, DIPUTITE	
7) PAY:	CRYOGLOBULIN WITH ANTI-IGG ACTIVITY, B IDIOTRE	
and the state	CONCELORITIN WITH ANTI-LOW-DENSITY LIPOPROTEIN ACTIVITY, O ISOCTOR	
111 015	CHYOGLOBULIN WITH ANTHIGG ACTIVITY: B IDIOTYPE CRYOGLOBULIN WITH ANTHIGG ACTIVITY: B IDIOTYPE	
14) SCA:	CHYOGLOBULIN WITH ANTI-LOW-DENSITY LIPOPROTEIN ACTIVITY; B IDIOTYPE CRYOGLOBULIN WITH ANTI-LOW-DENSITY LIPOPROTEIN ACTIVITY; B IDIOTYPE	
	CRYOGLOBULIN WITH ANTI-LOW DENSITY LIPOPHOTEIN ACTIVITY IN ANTI-IGG ACTIVITY; B IDIOTYPE ANTI-HUMAN GAMMA G GLOBULIN; WA IDIOTYPE; CATVITY (GROUP 1) COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY (GROUP 1)	
100 110	COLD AGGIUTININ WITH ANTI-BLOOD GROUP SMALL FACTOR	
	COVOCIONIN WITH ANTI-IGG AUTVILLE O DIDUTIES	
22) DBE	COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY	
	A CONTRACT ANTERIOOD GROUP LAGIVIT	
000 010 0	COLD ACCULITININ WITH ANTI-BLOOD GROUP LACIANTI (ATTECAN)	
	COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY	
100 01 4	CONCELOBILINE WITH ANTI-IGG ACTIVITY; B DUDITES	
43) SHE':	CRYOGLOBULIN WITH ANTI-IGG ACTIVITY BUDGITHE	
48) POM:	ANTI-HUMAN GAMMA GI GLOBULIN; PO IDIOTYPE ANTI-HUMAN GAMMA GI GLOBULIN; PO IDIOTYPE ANTI-MEASLES VIRUS (WOODFOLK STRAIN): ANTI-SUBACUTE SCLEROSING PANENCEPHALITIS VIRUS (LEC STRAIN)	
54) GOEII 62) TEH:	ANTHIMASLES VIAS (MOOFICE) ANTALA	
63) CRA(II); ANTI-HUMAN GAMMA G GLOBULIN	
64) PLA:	ANTI-HUMAN GAMMA G GLOBULIN ANTI-HUMAN GAMMA G GLOBULIN	
55) PIN: 70) BOB:	COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY	
71) DBI:	ANTI-HUMAN GAMMA G GLOBULIN	
72) WAL:	ANTI-HUMAN GAMMA G GLOBULIN ANTI-HUMAN GAMMA G GLOBULIN	
74) GAG:	COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY	
	HUMAN KAPPA LIGHT CHAINS SUBGROUP III	
	IGM-KAPPA IGM-KAPPA	
	і ідм-карра	
	IGM-KAPPA	
	IGM-KAPPA	
	IGM-KAPPA	
	IGM-KAPPA IGM-KAPPA	
	· IGM-KAPPA	
	: IGM-KAPPA	
	IGM-KAPPA .	
	: IGM-KAPPA	
REFEREN		
REFEREN 1) Tł:	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTED L RADNIKOL H I WATANABES, & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY	
REFEREN 1) Tł: 2) WOL	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER.L.BARNIKOL.H.U.WATANABE.S. & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM350.27.5-278; (1972) Z.PHYSIOL.CHEM353.189-208. (CHECKED BY AUTHOR) AUTHOR)	
1) 71:	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER.L.BARNIKOL.H.U.WATANABE.S. & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM350.27.5-278; (1972) Z.PHYSIOL.CHEM353.189-208. (CHECKED BY AUTHOR) AUTHOR)	
1) Tł: 2) WOL 3) SIE;	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L.BARNIKOL,H.U.WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS,D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVIMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822.	
1) Tł: 2) WOL 3) SIE;	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS D.W. & CAPRA.J.D. (1961) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS.D.W. & CAPRA.J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA.J.D. (1975) ADVIMMUNDOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS.D.W. & CAPRA.J.D. (1991) BIOCHEMISTRY.20.5816-5822. YOL: BENTLEY.D.L. (1984) INATURE.307.77-80.	
1) Tł: 2) WOL 3) SIE;	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS D.W. & CAPRA.J.D. (1961) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS.D.W. & CAPRA.J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA.J.D. (1975) ADVIMMUNDOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS.D.W. & CAPRA.J.D. (1991) BIOCHEMISTRY.20.5816-5822. YOL: BENTLEY.D.L. (1984) INATURE.307.77-80.	
1) Tł: 2) WOL 3) SIE;	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U., WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM., 350.275-276; (1972) Z.PHYSIOL.CHEM., 353,189-208. (CHECKED BY AUTHOR) ANDREWS D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADV.IMMUNICLOGY.20.1-AD, (CHECKED BY AUTHOR); ANDREWS.D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) ENDITION ON A SERVICE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81, FRANCINE, C., SOLOMONA, & FRANGIONE, B. (1983) J.IMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/23/81; GONIF, CHENP, P., PONS-ESTEL, B., CARSOND, A. & FRANGIONE, B. (1983) J.IMMUNOL, 131, 132-1325. (CHECKED BY AUTHOR 09/23/81; PONS-ESTEL, B., CONS.MA, & FRANGIONE, B. (1983) J.IMMUNOL, 131, 132-1325. (CHECKED BY AUTHOR 09/23/81; PONS-ESTEL, B., CONS.MA, & FRANGIONE, B. (1984) J.EXP.MED., 160.893-904; GONIF, CHENP, P., PONS-ESTEL, B., CARSOND,A. & SERVICE, 1994 J.E	
1) Tł: 2) WOL 3) SIE; 4) NG9 5) NEU 6) GOT	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L.BARNIKOLH,U.WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) SANDREWS,D.W. & CAPRA,J.D. (1961) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVIMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR); ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); MOREWS.D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1954) BIOCHEMISTRY.20.5816-5822. VCL: BENTLEY.D.L. (1984) NATURE.307.77-80. S. LEDFORD,D.K. GONIF, P.ZPOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, CHEN, P.P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON, A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, P.ZPONS-ESTEL, B., COLOMON,A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,A. & FRANGIONE, B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,A. & FRANGIONE, B. (1984) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,A. & FRANGIONE, B. (1985) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,D.A. & FRANGIONE, B. (1984) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,D.A. & FRANGIONE, B. (1985) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CONSON,D.A. & FRANGIONE, B. (1986) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 09/23/240); PONS-ESTEL, B., CONSON	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U., WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM., 350.275-276; (1972) Z.PHYSIOL.CHEM., 353,189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADV.IMMUNOLOGY.20.1-AD, (CHECKED BY AUTHOR); ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81) BARDINE, SOLOMONA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR (922384); PONS-ESTEL B., CORDNA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR (932384); PONS-ESTEL B., CONDNA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR (932384); PONS-ESTEL B., CONDA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR (932384); PONS-ESTEL B., CONDA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR (9329, 1905) SENTEL B., CONDA, & FRANGIONE,B. (1983) JIMMUNOL, 131, 1322-1325. (CHECKED BY	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U.,WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353,189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1961) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS,D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVIMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR); ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81, WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81, WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81, WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81, WHO SUGGESTED THAT SECUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR, 1981) BIOCHEMISTRY.20,5816-5822. VCL: BENTLEY.DL. (1984) NATURE.307.77-80. I: LEDFORD.DK.GONIFPIZZOLATO.M.FRANKLIN.E.C. SOLOMON.A. & FRANGIONE B. (1985) J.IMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23 841; PONS-ESTEL B., CARSON.DA. & FRANGIONE B. (1985) J.IMMUNOL134,1322-1325. (CHECKED BY AUTHOR (1992) B41; PONS-ESTEL B., CONS.A, & FRANGIONE B. (1993) J.IMMUNOL134,1322-1325. (CHECKED BY AUTHOR (1923 844); GONIFPIZZOLATO.M., FRANKLIN.E.C. SOLOMONA, & FRANGIONE B. (1983) J.IMMUNOL131,1322-1325. (CHECKED BY AUTHOR (1923 844); GONIFPIZPOLATO.M., FRANKLIN.E.C. SOLOMONA, & FRANGI	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE.S. & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA.J.D. (1981) BIOCHEMISTRY 20.5816-5822. CAPRA.J.D. (1975) ADV.IMMUNICOGY 20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA.J.D. (1975) ADV.IMMUNICOGY 20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. CCHECKED BY AUTHOR 08/25/81 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR): A SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81); GONIF.CHEN.P.P.PONS-ESTEL.B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); GONIF.CHEN.P.P.PONS-ESTEL.B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONC.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONSON.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONS	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE.S. & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA.J.D. (1981) BIOCHEMISTRY 20.5816-5822. CAPRA.J.D. (1975) ADV.IMMUNICOGY 20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS.D.W. & CAPRA.J.D. (1975) ADV.IMMUNICOGY 20.1-40. (CHECKED BY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. CCHECKED BY AUTHOR 08/25/81 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR): A SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (CHECKED BY AUTHOR 08/25/81); GONIF.CHEN.P.P.PONS-ESTEL.B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); GONIF.CHEN.P.P.PONS-ESTEL.B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/81); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CARSON.D.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONC.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONSON.A. & FRANGIONE B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL B., CONS	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-206. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1961) PROC.NAT.ACAD.SCI.USA.76.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS,D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVI,MMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD.SCI.USA.76.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. VCL: BENTLEY,D.L. (1984) INATURE.307.77-60. U: LEDFORD,D.K.,GONI,F., PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.ACT, A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, CHEN,P.P., PONS-ESTEL,B.,CONF.OND,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/33/84); PONS-ESTEL, B., CARSON,D.A. & FRANGIO	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-206. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1961) PROC.NAT.ACAD.SCI.USA.76.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS,D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVI,MMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD.SCI.USA.76.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. VCL: BENTLEY,D.L. (1984) INATURE.307.77-60. U: LEDFORD,D.K.,GONI,F., PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.S.OLOMON,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/61): PONS-ESTEL,B.,CONF.ACT, A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, CHEN,P.P., PONS-ESTEL,B.,CONF.OND,A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL, B., CARSON,D.A. & FRANGIONE, B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/33/84); PONS-ESTEL, B., CARSON,D.A. & FRANGIO	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L. BARNIKOL,H.U.WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS,D.W. & CAPRA,J.D. (1993) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVI,MMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS,D.W. & CAPRA,J.D. (1994) NATURE.307.77-80. NCL: BENTLEY,D.L. (1984) NATURE.307.77-80. IEDEFORD,D.K.,GONIF, P.P.ZOLATOM, FRANKLIN, E.C., SOLOMON,A. & FRANGIONE B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/81); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1986) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); GONIF, P.,PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); PONS-ESTEL, B.,CARSON,D.A. & FRANGIONE,B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); PONS-ESTEL, B.,CARSON,A. & FRANGIONE,B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); PONS-ESTEL, B.,CARSON,A. & FRANGIONE,B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/32/844); PONS-ESTEL,	
1) TI; 2) WOL 3) SIE; 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WE! 10) GAF 11) PIE;	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L. BARNIKOL,H.U.WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL,CHEM.350.275-276; (1972) Z.PHYSIOL,CHEM.353,189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS,D.W. & CAPRA,J.D. (1993) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADV.IMMUNOLOGY.201-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS,D.W. & CAPRA,J.D. (1991) BIOCHEMISTRY.20.5816-5822. NCL: BENTLEY,D.L. (1984) NATURE.307.77-80. LEDFORD,D.K.,GONI,F.,PZZOLATO,M.,FRANKLIN,E.C.,SOLOMON,A. & FRANGIONE B. (1985) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/25/81): FNNSESTEL B.,CARSON,D.A. & FRANGIONE B. (1985) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,CHEN,P.P.,PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,CHEN,P.P.,PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,CHEN,P.P.,PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1985) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,PZZOLATO,M.,FRANKLIN,E.C. SOLOMON,A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,PZZOLATO,M.,FRANKLIN,E.C. SOLOMON,A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,PZZOLATO,M.,FRANKLIN,E.C. SOLOMON,A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); GONI,F.,PIZOLATO,M.,FRANKLIN,E.C. SOLOMON,A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL,B.,CARSON,D.A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL,B.,GONI,F.,SOLOMON,A. & FRANGIONE,B. (1983) JIMMUNOL131,1322-1325. (CHECKED BY AUTHOR 09/23/84	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L. BARNIKOL,H.U.,WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL,CHEM.350.275-276; (1972) Z.PHYSIOL,CHEM.353,189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD,SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS,D.W. & CAPRA,J.D. (1993) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVIIMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD,SCI.USA.78.3799-3803. CAFECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. YCL: BENTLEY,D.L. (1984) NATURE.307.77-80. Y: LEDFORD,D.K.,GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1985) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/81); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1985) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/81); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); GONIF, CHEN,P.P.,PONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); FONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); FONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/344); PONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/344); PONS-ESTEL,	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER,L. BARNIKOL,H.U.,WATANABE,S. & HILSCHMANN,N. (1969) Z.PHYSIOL,CHEM.350.275-276; (1972) Z.PHYSIOL,CHEM.353,189-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD,SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS,D.W. & CAPRA,J.D. (1993) BIOCHEMISTRY.20.5816-5822. CAPRA,J.D. (1975) ADVIIMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRA,J.D. (1991) PROC.NAT.ACAD,SCI.USA.78.3799-3803. CAFECKED BY AUTHOR 08/25/81) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. YCL: BENTLEY,D.L. (1984) NATURE.307.77-80. Y: LEDFORD,D.K.,GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1985) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/81); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1985) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/81); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); FONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); GONIF, CHEN,P.P.,PONS-ESTEL,B.,COARSON,D.A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/25/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1986) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIF, PIZZOLATO,M., FRANKLIN, E.C., SOLOMON,A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); FONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/84); FONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/344); PONS-ESTEL, B.,COARSON,D.A. & FRANGIONE,B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 09/23/344); PONS-ESTEL,	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE.S. & HILSCHMANN.N. (1969) Z.PHYSIOL.CHEM.350.275-276; (1972) Z.PHYSIOL.CHEM.353,189-208. (CHECKED BY AUTHOR): A DEREWSD.W. & CAPRALD. (1981) PROC.NAT.ACAD.SCI.USA.78.3795-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWSD.W. & CAPRALD. (1991) BIOCHEMISTRY 203616-5822. CARRALD. (1970) BIOCHEMISTRY 203616-5822. CARRALD. (1970) A DEVISION WHO SUGGESTED THAT THE SECOURCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); (NDREWSD.W. & CAPRALD. (1981) BIOCHEMISTRY.203616-5822. NCI. BENTLEY.DL. (1984) NATURE 207.77-80. ELEFORD.D.K., GONLF, PIZZOLATO.M. FRANKLIN.E.C. SOLOMONA. & FRANGIONE.B. (1985) JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/25/81); EONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/25/81); SOLOMON.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/25/81); FONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/25/81); SOLOMON.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIFCHEN.P.P., PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIFCHEN.P.P., PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.JIMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIFCHEN.P.P., PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.IMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); GONIFCHEN.P.P., PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.IMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.EKP MED. 150.993.904. (1) ELDFORD.D.K., GONIF., PIZZOLATO.M., FRANKLIN.E.C., SOLOMONA. & FRANGIONE.B. (1983) J.IMMUNOL. 131.1322-1325. (CHECKED BY AUTHOR 09/23/84); PONS-ESTEL.B., CARSON.D.A. & FRANGIONE.B. (1984) J.EKP MED. 150.993.904. (2) ELEDFORD.D.K., GONIF., FIZZOLATO.M., FRANKLIN.E.C., SOLOMONA. & FR	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLO	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL H. U. WATANABE, S. & HILSCHMANN, N. (1969) Z.PHYSIOL, CHEM. 350,275-276; (1972) Z.PHYSIOL, CHEM. 353,189-208. (CHECKED BY AUTHOR) (1973) BIOCHEMISTRY 20,301-40. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/61); ANDREWS, D.W. & CAPRAJD. (1981) BIOCHEMISTRY 20,301-40. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD. (1981) DEVELOPMENT 20,35816-5822. NCL: BENTLEY, D.L. (1984) NATURE 20,777-80. S. LEDFORD, D.K. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:35.4079-4079. S. LEDFORD, D.K. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:31.1322-1325. (CHECKED BY AUTHOR 079.) S. LEDFORD, M. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:31.1322-1325. (CHECKED BY AUTHOR 079.) S. LEDFORD, M. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:31.1322-1325. (CHECKED BY AUTHOR 079.) S. LEDFORD, M. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:31.1322-1325. (CHECKED BY AUTHOR 079.) S. LEDFORD, M. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:31.1322-1325. (CHECKED BY AUTHOR 079.) S. LEDFORD, M. GONIF, PIZZOLATO, M. FRANKLIN, E.C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL:131.1322-1325. (CHECKED BY AUTHOR 059394) SONIF, CHEN, P.P., PONS-ESTEL, B., CARSON, D.A. & FRANGIONE B. (1983) JIMMUNOL.:131.1322-1325. (CHECKED BY AUTHOR 059394) SONIF, CHEN, P.P., PONS-ESTEL, B., CARSON, D.A. & FRANGIONE B. (1983) JIMMUNOL.:131.1322-1325. (CHECKED BY AUTHOR 05	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WIL 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTER L. BARNIKOL, H.U., WATANABE, S. & HILSCHMANN, N. (1959) Z.PHYSIOL, CHEM. 350.275-276; (1972) Z.PHYSIOL, CHEM. 353,189-208. (CHECKED BY AUTHOR) SUTER L. BARNIKOL, H.U., WATANABE, S. & HILSCHMANN, N. (1959) Z.PHYSIOL, CHEM. 350.275-276; (1972) Z.PHYSIOL, CHEM. 353,189-208. (CHECKED BY AUTHOR) SUTER L. BARNIKOL, H.U., WATANABE, S. & HILSCHMANN, N. (1959) Z.PHYSIOL, CHEM. 350.275-276; (1972) Z.PHYSIOL, CHEM. 351,799-3803. CHARAJD, (1975) ADV. IMMUNOL OGY 20,1-40. (CHECKED BY AUTHOR): ANDREWS, D.W. & CAPRAJD, (1981) PROC.NAT.ACAD. SCI.USA, 78.3799-3803. CHERCKED BY AUTHOR 082561 WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS, D.W. & CAPRAJD, (1981) BIOCHEMISTRY 20,5016-5822. CICL BENTLEY, D.L. (1984) NATURE 307.77-80. CIE DEORD, D.K. GONIF, PIZZOLATO, M. FRANKLINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3941) FOR (1981) FID. (1981) FID. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3941) FID. (1981) FID. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3940) FID. (1985) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3940) FID. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3940) FID. (1985) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3940) FID. (1985) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 09/3940) FID. (19	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WIL 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, BARNIKOL, H.U., WATANABE, S. & HILSCHMANN, N. (1959) Z.PHYSIOL, CHEM. 350,275-276; (1972) Z.PHYSIOL, CHEM. 353,159-208. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1981) PROC, NAT, ACAD, SCI, USA, 78,3795-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1981) PROC, NAT, ACAD, SCI, USA, 78,3795-3803. (CHECKED BY AUTHOR 08/25/81); MUSCING 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1981) PROC, NAT, ACAD, SCI, USA, 78,3795-3803. (CHECKED BY AUTHOR 08/25/81); MUSCING 08/25/81]; MUSCING 0	
1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE; 12) FLO 13) LOF 14) SCF 15) GLO 16) SAL 17) WIL 18) MA 19) NIC	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, GARNIKOL, H.U., WATANABELS, & HILSCHMANN, (1969) Z.PHYSIOL, CHEM. 350,275-276; (1972) Z.PHYSIOL, CHEM. 353,189-208. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1981) PROCNAT, ACAD, SCI, USA, 78,3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1993) PROCNAT, ACAD, SCI, USA, 78,3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1993) PROCNAT, ACAD, SCI, USA, 78,3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1993) PROCNAT, ACAD, SCI, USA, 78,3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1993) PROCNAT, ACAD, SCI, USA, 78,3799-3803. (CHECKED BY AUTHOR 08/25/81); ANDREWS, D.W. & CAPRA, J.D. (1993) DIJOHENNISTUX, J.S.BIFJASZZ. COLDEN, J., CHECKE, D.Y., J.M., J. (1993) DIJOHENNISTUX, J.S.BIFJASZZ. COLDEN, J., CONIF, F., PIZZOLATO, M., FRANKLINE, C., SOLOMON, A. & FRANGIONE B. (1983) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/25/801); GONIF, CHEN, P.P., ZOLATO, M., FRANKLINE, C., SOLOMON, A. & FRANGIONE B. (1984) J.M.MUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/25/801); GONIF, CHEN, P.P., ZOLATO, M., FRANKLINE, C., SOLOMON, A. & FRANGIONE B. (1983) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/23/801); GONIF, CHEN, P.P., P.D.SCHESTEL, B., CARSON, D.A. & FRANGIONE, B. (1984) J.LEXP. MEO. 160,893-504. (GONIF, CHECKED BY AUTHOR 03/23/801); GONIF, CHEN, P.P., P.D.SCHESTEL, B., CARSON, D.A. & FRANGIONE, B. (1984) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/23/801); GONIF, CHEN, P.P., P.ZOLATO, M., FRANKLINE, C., SOLOMONA, & FRANGIONE, B. (1983) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/33/801); GONIF, CHEN, P.P., ZOLATO, M., FRANKLINE, C., SOLOMONA, & FRANGIONE, B. (1983) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/33/801); GONIF, CHEN, P.P., PONSESTEL, B., CARSON, D.A. & FRANGIONE, B. (1983) J.IMMUNOL, 131, 1322-1325. (CHECKED BY AUTHOR 03/33/801); GONIF, CHEN, P.P., PONSESTEL, B., CARSON, D.A. & F	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WIA 18) MA 19) NIC 20) CUI 	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL_BARNIKOL.HU.WATANABES. & HUSCHMANN.N. (1969) Z.PHYSIOL.CHEM.350.275276; (1972) Z.PHYSIOL.CHEM.353,189-208. (CHECKED BY AUTHOR) : ANDREWS.D.W. & CAPRAJ.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS.D.W. & CAPRAJ.D. (1991) BIOCHEMISTRY.205816-5322. CAPRAJ.D. (1975) ADVIMMUNOLOGY.301-J0 (CHECKED BY AUTHOR). ANDREWS.D.W. & CAPRAJ.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-9803. (CHECKED BY ACTION OF THE CONSTRUCTION OF THE CHECKED BY AUTHOR). ANDREWS.D.W. & CAPRAJ.D. (1991) PROC.NAT.ACAD.SCI.USA.78.3799-9803. (CHECKED BY ACTION OF THE CONSTRUCTION OF THE SECURICE DETERMINED IN 1975 WAS INCORRECT AND SHOLLD BE DELETED; (CHECKED BY ACTION OF THE CONSTRUCTION OF THE CONSTRUCTION OF THE SECURICE DETERMINED IN 1975 WAS INCORRECT AND SHOLLD BE DELETED; (CHECKED BY ALCONSTRUCTION OF THE CONSTRUCTION OF THE SECURIC DETERMINED IN 1975 WAS INCORRECT AND SHOLLD BE DELETED; (CHECKED BY, ALCONSTRUCTION OF THE CONSTRUCTION OF THE SECURIC DETERMINED IN 1975 WAS INCORRECT AND SHOLLD BY AUTHOR 1 EEDFORD DK.GONE, FAZOLATO M. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUEL P.P.P.DONS-ESTELB.CARSON.D.A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUEL P.P.ZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUZZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUZZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUZZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1980) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUZZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1983) JIMMUNOL.131.1322-1325. (CHECKED BY AUTHOR 1 EEDFORD DK.GONE, FUZZOLATOM. FRANKLIN E.C. SOLOMON A. & FRANGIONE B. (1983) JIMMUNOL.131.1322-1325. (CH	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WHA 18) MAC 19) NIC 20) CUI 21) FR4 	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUJTERL_BARNIKOL,H.U.WATANABES, & HILSCHMANN,N. (1959) Z.PHYSIOL,CHEM.350,275-276; (1972) Z.PHYSIOL,CHEM.353,189-208, (CHECKED BY AUTHOR) : ANDREWS D.W. & CAPRA.J.D. (1959) PROC.NAT.ACAD.SCI.USA.76.3799-3003. (CHECKED BY AUTHOR 08/25/81): ANDREWS D.W. & CAPRA.J.D. (1951) BIOCHEMISTRY.203616-5822. CAPRA.J.D. (1957) ADV.IMMUNDLOGY.20.1-40. (CHECKED BY AUTHOR) MOREWS D.W. & CAPRA.J.D. (1951) PROC.NAT.ACAD.SCI.USA.76.3799-3003. (CHECKED BY AUTHOR 08/25/81 WHO BOCHEMISTRY.203616-5822. CAPRA.J.D. (1975) ADV.IMMUNDLOGY.20.1-40. (CHECKED BY AUTHOR) MOREWS D.W. & CAPRA.J.D. (1951) PROC.NAT.ACAD.SCI.USA.76.3799-3003. (CHECKED BY AUTHOR DECHEMISTRY 20.3616-5822. CAPRA.J.D. (1975) ADV.IMMUNDL.GY.20.777-80. CHECKED BY AUTHOR BOOTARS FEEL B.,CARSON D.A. & FRANGIONE B. (1983) JIMMUNDL. 131.1322-1325. (CHECKED BY AUTHOR D.K., GONIF, F.PZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL. 131.1322-1325. (CHECKED BY AUTHOR D.K., GONIF, F.PZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL. 131.1322-1325. (CHECKED BY AUTHOR D.K., GONIF, F.PZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.K., GONIF, F.PZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304); GONIF, F.PZZOLATO, M.FRANKLINE C. SOLOMONA. & FRANGIONE B. (1983) JIMMUNDL.131.1322-1325. (CHECKED BY AUTHOR D.Y32304);	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI' 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WIL 18) MAI 19) NIC 20) CUI 21) FAR 22) DRI 	CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SULTERL BARNIKOLHU, WATANABES, & HUSCHMANN, (1969) Z.P.HYSIOL, CHEM, 350,275-276; (1972) Z.P.HYSIOL, CHEM, 353,189-208, (CHECKED BY AUTHOR) (1991) BIOCHMISTRY, 200,11691) PROC.NATACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 08/25/81): ANDREWS.D.W. & CAPRA.J.D. (1991) BIOCHMISTRY, 200,516-5922. (CAPRA.J.D. (1973) ADV.IMMUNOL.GY.20.140, CONCENCE DY AUTHOR): ANDREWS.D.W. & CAPRA.J.D. (1981) PROC.NATACAD.SCI.USA.78.3799-3803. (CHECKED BY AUTHOR 950; (1981) BIOCHMISTRY, 200,516-5927. (CHECKED BY AUTHOR 957,500; 200,000,4,5; FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1985) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1985) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322-1325. (CHECKED BY AUTHOR 95394); PONS-ESTEL,B., CARSON, D.A. & FRANGIONE,B. (1983) JIMMUNOL, 131,1322	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WIL 18) MA 19) NIC 20) CUI 21) FA- 22) DRI 23) PEI 24) CA 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, BARNIKOLHU, WATANABELS, & HILSCHMANN, (1989) Z.PHYSIOL, CHEM.350,275-276; (1972) Z.PHYSIOL, CHEM.353,189-208. (CHECKED BY AUTHOR) ANDREWS, D.W., & CAPRAJD, (1991) PROCINATACAD SCILUSA,78,3799-3803. (CHECKED BY AUTHOR) 09/25/81; ANDREWS, D.W. & CAPRAJD, (1991) BROCINATACAD SCILUSA,78,3799-3803. (CHECKED BY AUTHOR) 09/25/81; ANDREWS, D.W. & CAPRAJD, (1991) BROCINATACAD SCILUSA,78,3799-3803. (CHECKED BY AUTHOR) 09/25/81; ANDREWS, D.W. & CAPRAJD, (1971) MOLTAND, 2016; DS210, 2010; DS10,000, Y20,140, (DHECKED BY AUTHOR); ANDREWS, D.W. & CAPRAJD, (1991) BROCINATACAD SCILUSA,78,3793-9803. CAPRAJD, (1972) MAY, MAYAMAD, (1981) BIOCHEMISTRY, 2018; DS220; THAT THE SCOURNOE DE TERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS, D.W. & CAPRAJD, (1981) BIOCHEMISTRY, 2018; DS210; DS10, Y20, Y20, Y20, Y20, Y20, Y20, Y20, Y2	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI' 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLO 16) SAI 17) WIL 18) MA 19) NIC 20) CUI 21) FAR 22) DRI 23) PEA 24) CA 25) STI 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUITERL BARNINGL.HU., WATANABES, & HILSCHMANN,N. (1989) Z.PHYSIOL.CHEM.350.275-278; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS, D.W. & CAPRAJ,D. (1989) PROC.NAT.ACAD.SCIUSA.78.3799-3803. (CHECKED BY AUTHOR 08/25/61); ANDREWS,D.W. & CAPRAJ,D. (1991) PROC.NAT.ACAD.SCIUSA.78.3799-3803. (CAPPAJ,D. (1975) ADVIMMUNCIO, COMO SUGGESTED THAT THE SECUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED; ANNREWS,D.W. & CAPRAJ,D. (1991) PROC.NAT.ACAD.SCIUSA.78.3799-3803. (CL: BENTLEY,D.L. (1984) NATURE.307.77-60. Y.L. (1984) C.N. (1984) NATURE.307.77-60. Y.L. (1984) C.N. (1984) NATURE.307.77-60. Y.CL: BENTLEY,D.L. (1984) NATURE.307.77-60. Y.CL: BENTLEY,D.L. (1984) J.B.MALING, C.S.OLOMON,A. & FRANGIONE B. (1983) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 03/2384): GONI,F., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 03/2384): GONI,F., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 03/2384): GONI,F., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, S., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, S., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, S., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, J., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, J., CHEN,P.P., PONS-ESTEL,B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/3294); GONI, J., CHEN,P.P., PONS-ESTEL, B., CARSON,D.A. & FRANGIONE B. (1984) J.IMMUNOL131.1322-1325. (CHECKED BY AUTHOR 05/32934); GONI, J.	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WH 18) MA 19) NIC 20) CUI 21) FR4 22) PEI 23) PEI 24) CA 25) GLC 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, BARNIKOL HU, WATANABES, & HILSCHMANN, 1989) Z.PHYSIOL,CHEM.350,275-278: (1972) Z.PHYSIOL,CHEM.353,199-208. (CHECKED BY AUTHOR) ANDREWS,D.W. & CAPRAJD, (1989) PROC.NAT.ACAD.SCI.USA.76.3799-3803. (CHECKED BY AUTHOR 08/25/61): ANDREWS,D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3798-3803. CAPRAJD, (1972) ATOM REVISIOL 067.201.40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3798-3803. CAPRAJD, (1973) ATOM REVISIOL 067.201.40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3798-3803. CAPRAJD, (1974) ATOM REVISION OF 201.40. (CHECKED BY AUTHOR): ANDREWS,D.W. & CAPRAJD. (1981) PROC.NAT.ACAD.SCI.USA.78.3798-3803. CALE BENTLEY,D.L. (1984) INATURE.307.77-80. CELE BENTLEY,D.L. (1984) NATURE.307.77-80. CELE DENTLEY,D.L. (1984) INATURE 307.77-80. CELE DENTLEY,D.L. (1984) C.C. COLL, PAZZOLATO, M. FRANKLINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DESTORD, CONL, F. PIZZOLATO, M. FRANKLINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEDORD,D.K. CONLE, F. ZZOLATO, M. FRANKLINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEDORD,D.K., CONLE, F. ZZOLATO, M. FRANKINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEDORD,D.K., CONLE, F. ZZOLATO, M. FRANKINE C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEDORD,D.K., CONLE, F. ZZOLATO, M. FRANKINE, C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEORD,D.K., CONLE, F. ZZOLATO, M. FRANKINE, C. SOLOMONA, & FRANGIONE B. (1983) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEORD,D.K., CONLE, F. ZZOLATO, M. FRANKINE, C. SOLOMONA, & FRANGIONE B. (1984) JIMMUNOL131.1322-1325. (CHECKED BY AUTHOR) DEORD,D.K., SOLO	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WH 18) MAC 20) CUI 21) FRA 22) DRI 23) PEI 24) CA 25) STI 26) GJ 27) TA 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, BARNIKOL-HU, WATANABES, & HILSCHMANNN. (1989) Z.PHYSIOLCHEM.350.275-276; (1972) Z.PHYSIOLCHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS, D.W., & CAPRALED, (1981) PROCLMAT ACAO.SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 0#25/81); ANDREWS, D.W., & CAPRALD, (1975) ADVINUMUCIOOY.201-40. (CHECKED BY AUTHOR); ANDREWS, D.W., & CAPRALD, (1981) PROC.MAT ACAO.SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 0#25/81) WHO SUCCESTED THAT THE SECURNCE DETERMINED IN 1975 MAS INFCORRECT AND GHOULD BE DELETED); ANDREWS, D.W., & CAPRALID, (1981) BIOINTON, 2016 1981 CIC: BENTLEY, D.L. (1984) MATURE, C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#25/81 WHO SUCCESSTED THAT THE SECURNCE DETERMINED IN 1975 MAS INFCORRECT AND GHOULD BE DELETED); ANDREWS, D.W., & CAPRALE, B.CARSOND, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#25/81 WHO SUCCESSTED THAT THE SECURNCE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#36/81 BUNNOL, 135.4075-4079. I. EDFORD, D.K., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1986) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1983) AUTHOR 0018. CHECKED STANDAUCUSS 0019-4079. I. EDFORD, D.K., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, STANDAUCUSS 0019-4099. I. EDFORD, M., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; DONK, SOLUM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; DONK, SOLUM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 1	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WH 18) MAC 20) CUI 21) FRA 22) DRI 23) PEI 24) CA 25) STI 26) GJ 27) TA 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL, BARNIKOL-HU, WATANABES, & HILSCHMANNN. (1989) Z.PHYSIOLCHEM.350.275-276; (1972) Z.PHYSIOLCHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS, D.W., & CAPRALED, (1981) PROCLMAT ACAO.SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 0#25/81); ANDREWS, D.W., & CAPRALD, (1975) ADVINUMUCIOOY.201-40. (CHECKED BY AUTHOR); ANDREWS, D.W., & CAPRALD, (1981) PROC.MAT ACAO.SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 0#25/81) WHO SUCCESTED THAT THE SECURNCE DETERMINED IN 1975 MAS INFCORRECT AND GHOULD BE DELETED); ANDREWS, D.W., & CAPRALID, (1981) BIOINTON, 2016 1981 CIC: BENTLEY, D.L. (1984) MATURE, C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#25/81 WHO SUCCESSTED THAT THE SECURNCE DETERMINED IN 1975 MAS INFCORRECT AND GHOULD BE DELETED); ANDREWS, D.W., & CAPRALE, B.CARSOND, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#25/81 WHO SUCCESSTED THAT THE SECURNCE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1985) AUTHOR 0#36/81 BUNNOL, 135.4075-4079. I. EDFORD, D.K., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1986) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 1983) AUTHOR 0018. CHECKED STANDAUCUSS 0019-4079. I. EDFORD, D.K., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, STANDAUCUSS 0019-4099. I. EDFORD, M., GONH, F. PIZOLATOM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; PONS 55TEL B.CARONDAU, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; DONK, SOLUM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 131.1322-1325. (CHECKED BY AUTHOR 0923081; DONK, SOLUM, FRANKLINE C. SOLOMONA, & FRANCIONE B. (1985) JIMMUNOL, 1	
 1) TI: 2) WOL 3) SIE: 4) NG9 5) NEU 6) GOT 7) PAY 8) SON 9) WEI 10) GAF 11) PIE: 12) FLO 13) LOF 14) SCA 15) GLC 16) SAI 17) WH 18) MA 19) NIC 20) CUI 21) FR4 22) PEI 24) CA 25) STI 26) GJ 27) TA 28) IAI 29) RA 29) RA 	 CE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III SUTERL BARNIKOL-HULWATANAGES. & HILSCHMANNN. (1989) Z.PHYSIOL.CHEM.350.275-278; (1972) Z.PHYSIOL.CHEM.353.189-208. (CHECKED BY AUTHOR) ANDREWS D.W., & CAPPALD. (1981) PROCLMAT ACAD.SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 08/25/61); ANDREWS.D.W. & CAPPA.J.D. (1987) BD/ (1975) AD/ IMMUNICIOGY 201-40. (CHECKED BY AUTHOR); ANDREWS.D.W. & CAPPA.J.D. (1981) PROCLMAT ACAD SCLUSA.78.3799-3803. (CHECKED BY AUTHOR 08/25/61) WHO SUGGESTED THAT THE SEQUENCE DETERMINED IN 1975 WAS INCORRECT AND SHOULD BE DELETED); ANDREWS.D.W. & CAPPA.J.D. (1981) BD/CHEMISTRY.2015/05/15/201 VILI: BENTLEY.D.L. (1984) NATURE 30.77762. VILI: BENTLEY.D.L. (1984) NATURE 30.77762. VILI: DENTLEY.D.L. (1985) NATURE 30.77762.	

10.0

REFERENCE: HUMAN KAPPA LIGHT CHAINS SUBGROUP III (cont'd) 32) MCE: MIDDAUGH,C.R.,KEHOEJ,M.,PRYSTOWSKY,M.B.,GERBER-JENSON,B.,JENSON,J.C. & LITMAN,G.W. (1978) IMMUNOCHEM., 15, 171-187. (CHECKED BY AUTHOR 10/22/80) AUTHOR 10/22/80) 3) KEA: WANG.A.C. & FUDENBERG.H.H. (1975) IMMUNOL.COMMUN.4.433.497. (CHECKED BY AUTHOR 09/23/7): WANG.A.C. TUNG.E..WANG.I.,FUDENBERG. H.H.,PICK.AI. & FROEHLICHMAN.R. (1975) IMMUNOL.IMMUNOLTHER.381.456. (CHECKED BY AUTHOR 03/18/81) 3) SMI: NIALL.H.O. & EDMAN.P. (1987) NATURE.216.282.263. (CHECKED BY AUTHOR 07/25/79) 3) AJ: CAPRA.J.D.,KEHOE.J.M.,WILLIAMS.R.C.,JR.,FEIZI.T. & KUNKEL.H.G. (1972) PROC.NAT.ACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR) 3) AJ: CAPRA.J.D.,KEHOE.J.M.,WILLIAMS.R.C.,JR.,FEIZI.T. & KUNKEL.H.G. (1972) PROC.NAT.ACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR) 3) AJ: CAPRA.J.D.,KEHOE.J.M.,WILLIAMS.R.C.,JR.,FEIZI.T. & KUNKEL.H.G. (1972) PROC.NAT.ACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR) 3) AJ: CAPRA.J.D.,KEHOE.J.M.,WILLIAMS.R.C.,JR.,FEIZI.T. & KUNKEL.H.G. (1972) PROC.NAT.ACAD.SCI.USA.69.40-43. (CHECKED BY AUTHOR) 3) BRO'IGG: HOPPERJ.E.,NOYES.C.,MEINRIKSON.R. & KESSEL.J.W. (1976) JIMMUNOL.16.733.746. (CHECKED BY AUTHOR 01/25/83) 3) KE: CAPRA.J.D. (1975) ADV.IMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR 07/25/79) 3) IKE: CAPRA.J.D. (1975) ADV.IMMUNOLOGY.20.1-40. (CHECKED BY AUTHOR) 3) TIL: PINK.J.R.L.,WANG.A.C. & FUDENBERG.H.H. (1971) ANN.REV.MED..22.145-170. (CHECKED BY AUTHOR) 3) TIL: PINK.J.R.L.,WANG.A.C. & FUDENBERG.H.H. (1971) ANN.REV.MED..22.145-170. (CHECKED BY AUTHOR) 4) AMYLOID, KSAI: SLETTENK, WESTERMARK,P.,PITKANEN,P.,THYRESSON.N. & OLSTAD.O.K. (1983) SCAND.J.IMMUNOL.,18,557-560. (CHECKED BY AUTHOR) 40] AMYLOID KSA: SLETTEN,K.,WESTERMARK,P.,PITKANEN,P.,THYRESSON,N. & OLSTAD.O.K. (1983) SCAND.J.IMMUNOL., 18,557-560. (CHECKED BY AUTHOR 04/26/84) WANG.A.C., WELLS.J.V., FUDENBERG.H.H. & GERGELY.J. (1974) IMMUNOCHEM..11.341-345. (CHECKED BY AUTHOR) LEDFORD.D.K..GONI, F., PIZZOLATO, M., FRANKLIN, E.C., SOLOMON, A. & FRANGIONE, B. (1983) J.IMMUNOL..131.1322-1325. (CHECKED BY AUTHOR 03/23/84) 41) POL: LEDFORD.D.K., GONI, F., PIZZOLATO, M., FRANKLIN, E.C., SOLOMON, A. & FRANGIONE, B. (1983) J.IMMUNOL., 131, 1322-1325. (CHECKED BY AUTHOR 03/23/84) 42) CLA: 43) SHE': IEMMERSUN.H. KAPLAN.B. DENTON.M.D. ANDERAS.P. ANDERSON.B. & MARGOLIASH.E. (1978) BIOCHEMISTRY.18.4676-4683. NIALL.H.D. & EOMAN.P. (1967) NATURE.216.262-263. (CHECKED BY AUTHOR 07/25/79) WANG.A.C. WELLS.J.V. FUDENBERG.H.H. & GERGELY.J. (1974) IMMUNOCHEM..11.341-345. (CHECKED BY AUTHOR) WANG.A.C. TUNG.E. WANG.I. FUDENBERG.H.H., PICK.A.I. & FROEHLICHMAN.R. (1980) CANCER IMMUNOLIMMUNOTHER..9.81-86. (CHECKED BY AUTHOR 03/10/81) JEMMERSON,R.,KAPLAN,B.,DENTON,M.D.,ANDERAS,P.,ANDERSON,B. & MARGOLIASH.E. (1979) BIOCHEMISTRY, 18,4676-4683. 44) JH: 45) WIN: 46) LEA: AUTHOR 03/18/81) 48) POM: KLAPPER.D.G. & CAPRA.J.O. (1976) ANN.IMMUNOL.(INST.PASTEUR),127C.261-271. (CHECKED BY AUTHOR 08/01/79) 49) VAND: SEON.B.K.GALANI,S.,HENDERSON,E.S. & PRESSMAN,D. (1977) IMMUNOCHEM.,14.567-572. (CHECKED BY AUTHOR 08/28/78) 50) AMYLOID S0124; SLETTEN,K.,WESTERMARK,P.,PITKANEN,P.,THYRESSON,N. & OLSTAD.O.K. (1989) SCAND,JIMMUNOL.,18.557-560. (CHECKED BY AUTHOR 04/25/84) WANG, A.C., TUNG, E., WANG, J., FUDENBERG, H.H., PICK, A.L. & FROEHLICHMAN, R. (1960) CANCER IMMUNOL IMMUNOTHER. 9.81-86. (CHECKED BY AUTHOR 03/18/81) WANG, A.C., TUNG, E., WANG, I., FUDENBERG, H.H., PICK, A.J. & FROEHLICHMAN, R. (1980) CANCER IMMUNOLIMMUNOTHER., 9,81-86. (CHECKED BY AUTHOR 03/18/81) 51) DOV: AUTHOR 03/18/81) NIALLH.D. & EDMAN.P. (1967) NATURE.216.262-263. (CHECKED BY AUTHOR 07/25/79) STROSBERG.A.D.KARCHER.D. & LOWENTHAL.A. (1975) J.IMMUNOL..115.157-160. (CHECKED BY AUTHOR) WANG.A.C., TUNG.E., WANG.I., FUDENBERG.H.H., PICK.A.I. & FROEHLICHMAN.R. (1980) CANCER IMMUNOL.IMMUNOTHER..9.81-86. (CHECKED BY AUTHOR 03/18/81) 52) SHM: 53) GRA: 54) GOEII: 5) LOW: WANG.A.C., TUNG.E., WANG.L.FUDENBEHG.H.H., PICK.AI. & FHUEHLICHMAN.R. (1980) CARCER IMMUNOUTHER.36466. (CHECKEN AUTHOR 03/18/3)
56) VER: CHERSIA. & NATALLP.G. (1978) IMMUNOCHEMISTRY.15.585-589. (CHECKED BY AUTHOR 09/13/79)
57) REE: PRELLI, TUMMOLO, D., SOLOMON,A. & FRANGIONE,B. (1980) SCAND.JIMMUNOL., IN PRESS.
58) WE: DWORSKY,E., SLETTEN,K.: HARBOE,M. & WETTELAND,P. (1980) SCAND.JIMMUNOL. 12.281-287. (CHECKED BY AUTHOR 02/28/1984)
59) HOW: KAPLAN,A.P. & METZGER,H. (1989) BIOCHEMISTRY.8.3944-3951. (CHECKED BY AUTHOR)
50) H58: HOODL.,GRAY,W.R.,SANDERS,B.G. & DREYER,W.J. (1967) COLD SPRING HARBOR SYMP. OUANTITATIVE BIOL.32.133-145.
51) HBJ5: HOODL.,GRAY,W.R.,SANDERS,B.G. & DREYER,W.J. (1967) COLD SPRING HARBOR SYMP. OUANTITATIVE BIOL.32.133-145.
52) TEH: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
54) PLA: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) PLA: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) PLA: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) PLA: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) PLA: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) MCE: CAPRA,J.O.,KENGEJ.M., WELCH.E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66.842-847. (CHECKED BY AUTHOR 10/17/77)
56) MCCE: CAPRA,J.O.,KENGEJ.M., WILLIAMS,R.C.J.R.,FEIZIT. & KUNKEL,H.G. (1972) PROC.NATACAD.SCI.USA.89.40-43. (CHECKED BY AUTHOR 10/17/77) 94 PLA: JOHNSTON,S.L., ABRAHAM,G.N. & WELCH,E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.86.842-847. (CHECKED BY AUTHOR 10/17/77)
65) PIN: JOHNSTON,S.L., ABRAHAM,G.N. & WELCH,E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.86.842-847. (CHECKED BY AUTHOR 10/17/77)
66) MCE: CAPRA,J.O., KEHOE,J.M., WILLIAMS,R.C., JR., FEIZI,T. & KUNKEL,H.G. (1972) PROC.NAT.ACAD.SCI.USA.89.40-43. (CHECKED BY AUTHOR)
67) HAC: HOOD,L. & TALMAGE,D.W. (1970) SCIENCE, 169,325-334.
68) K. EVIS'CL: STAVNEZEH,J., KEKISH.O. BATTER,D., GRENIEH,J., BALAZSI, HENDERSON,E. & ZEGERS,B.J.M. (1986) NUC.ACIDS RES.,13,3495-3514.
69) BER: WANG,A.C., WELLS,J.V., FUDENBERG,H.H. & GERGELY,J. (1974) IMMUNOCHEM.,11,341-345. (CHECKED BY AUTHOR)
70) BOR: GERGELY,J., WANG,A.C., & FUDENBERG,H.H. & GERGELY,J. (1974) IMMUNOCHEM., 11,341-345. (CHECKED BY AUTHOR)
71) DRI: CAPRA,J.D. (1975) ADV.IMMUNOLOGY.20,1-40. (CHECKED BY AUTHOR)
72) WAL: CAPRA,J.D. (1975) ADV.IMMUNOLOGY.20,1-40. (CHECKED BY AUTHOR)
73) GOL: CAPRA,J.D. (1975) ADV.IMMUNOLOGY.20,1-40. (CHECKED BY AUTHOR)
74) GAG: CAPRA,J.D. (1975) ADV.IMMUNOLOGY.20,1-40. (CHECKED BY AUTHOR)
75) DOB: HOOD,L., & TALMAGE,D.W. (1970) SCIENCE, 1863,225-334.
76) H38: HOOD,L., & TALMAGE,D.W. (1970) SCIENCE, 1863,225-334.
76) H38: HOOD,L., & TALMAGE,D.W. (1970) SCIENCE, 1863,225-334.
76) H39: HOOD,L., & TALMAGE,D.W. (1970) SCIENCE, 1863,225-334.
76) H39: HOOD,L., GRAY,W.R., SANDERS,B.G. & DREYER,W.J. (1987) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,133-145.
77 HBJ12: HOOD,L., GRAY,W.R., SANDERS,B.G. & DREYER,W.J. (1987) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,133-145.
78) BUR(K): MOULINA. & FOUGEREAU.M. (1973) NATURE NEW BIOLOGY.246,176-178. (CHECKED BY AUTHOR)
79) LEG: MOULINA. & FOUGEREAU.M. (1973) NATURE NEW BIOLOGY.246,176-178. (CHECKED BY AUTHOR)
79) BUR(K): MOULINA. & FOUGEREAU.M. (1973) NATURE NEW BIOLOGY.246,176-178. (CHECKED BY AUTHOR) 82) VKAPPAS'CL: BENTLEY, D.L. & RABBITTS, T.H. (1981) CELL.24,613-623. (CHECKED BY AUTHOR 12/07/81) NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP III DENTICAL SETS OF FRAMEWORK SEGMENTS: SETS OF FRAMEWORK SEGMENTS: SET 1: TII1,WOL(2),SIE(3),NG9°CL(4),NEU(5),GOT(6),PAY(7),SON(8),WEI'(9),GAR'(10),PIE(11),FLO(12),LOP(13),SCA(14),GLO(15),SAL(16), WIL(17),MA1(8),NC(19),CUR(20),FR4(21),DRE(22),PERI23),CAM(24), (24 IDENTICAL) SET 2: GJ(26),TAK(27), (2 IDENTICAL) SET 3: RAD(22),DI(20),CAS(31), (3 IDENTICAL) SET 4: KEA(33),SM(134), (2 IDENTICAL) SET 4: KEA(33),SM(134), (2 IDENTICAL) SET 6: CLA(42),SHE(43), (2 IDENTICAL) SET 6: CLA(42),SHE(43), (2 IDENTICAL) FR1: SET 1: TITI, WOLZI, SIEJA, NGO'CL(4), NEU[5], GOTIE, SON[8], GAR'110], PIETTI, FLOT2, GLOT15, CUR(20). (12 IDENTICAL HUMAN SET 1: TITI, WOLZI, SIEJA, NGO'CL(4), NEU[5], GOTIE, SON[8], GAR'110], PIETTICAL HUMAN VKAPPA-W: Vg'CL(122).) VKAPPA-W: Vg'CL(122).) FR2: SET 1: TI[1], WOL[2]. (2 IDENTICAL) SET 2: GOTIGL, PAYT7, GAR (10), PIE(11), FLO(12), GLO(15), CUR(20). (7 IDENTICAL) FR3: SET 2: GO1(6),PAY[7],GAR [1]0],PIE[1]1,FLO]12],GLO[13],ULA[20], [7: IDENTICAL HUMAN V-KAPPA-III: ALSO 3 HUMAN V-KAPPA-II: SET 1: WOL[2],PAY[7],PIE[11],GLO[15],CUR]20],BEE[57],WAPPA3CU[32], [7: IDENTICAL HUMAN V-KAPPA-III: ALSO 3 HUMAN V-KAPPA-IV: PB17IV'CL[3], AU[2],64(1)(366),CL'1101; 2 HUMAN V-KAPPA1: CM 607 'CL[5],RPM1-6410'CL[16]; AND 1 HUMAN V-KAPPA-IV: PB17IV'CL[3], SET 2: POM[48], (IDENTICAL TO TUL],FR1(31],IARC/9L41'CL[28], (8: IDENTICAL HUMAN V-KAPPA-IV; LEN[4], DEN146[DB[68]; 2 HUMAN V-KAPPA-II; HAU[4], AND 1 HUMAN V-KAPPA-IV; LEN[4], SET 3: NOUIS(GARTI 10FL 10FL),FR1(31],IARC/9L41'CL[28], (8: IDENTICAL HUMAN V-KAPPA-IV; LEN[4], DEN146[DB[68]; 2 HUMAN V-KAPPA-IV; VJI'CL[11],
 SET 4: SON(8), (IDENTICAL TO 1 HUMAN V-KAPPA-IV; VJI'CL[1],) FR4; IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS: SETS OF COMPLEMENTARITY DETERMINING REGIONS. SET : SIEJAIKEI3BI, (2 IDENTICAL) SET 2: NG9'CLI4I,PAY(7),SON(B),WEI'19,GAR'[10],PIE[11],FLO[12],GLO[15],CUR[20],DRE[22],CAM[24]. (11 IDENTICAL) SET 3: TIL[39]. (IDENTICAL TO 1 MOUSE V:KAPPA-V: Vg'CLI[22].) SET 1: WOL[2],SIE[3],NEUI5],GOT(6],PAY[7],SON(B],GAR[10],PIE[11],FLO[12],GLO[15],CUR[20]. (11 IDENTICAL) SET 2: POMI4BI, (IDENTICAL TO 1 MOUSE V:KAPPA-V: Vn'CLI[2].) SET 2: POMI4BI, (IDENTICAL TO 1 MOUSE V:KAPPA-V: Vn'CLI[2].) SET 2: GOT(B],CUR[20]. (2 IDENTICAL) SET 3: POMI4BI, (IDENTICAL TO 1 HUMAN V:KAPPA-I: LAY[39].) SET 3: PAY[7],GLO[15], (2 IDENTICAL) SET 4: GAR'[10],FLO[12]. (2 IDENTICAL) CDR1: CDR2: SEIS OF J-MINIGENES: SET 1: PIE[1]; VKAPPA3CL[82]; (2 IDENTICAL HUMAN V-KAPPA-III; ALSO 1 HUMAN V-KAPPA-I; AU[2]; 1 HUMAN V-KAPPA-II: RPM1-6410CL[16]; AND 1 HUMAN V-KAPPA-IV: PB17IV CL[3].) SET 2: GOT[8]; (IDENTICAL TO 1 HUMAN V-KAPPA-I; AG[7].) SET 3: GAR[10]; FLO[12]; LARCIBL41*CL[28]; (3 IDENTICAL HUMAN V-KAPPA-III; ALSO 2 HUMAN V-KAPPA-I; DEN[46]; BI[63]; AND 1 HUMAN V-KAPPA-II; FR]14].) SET 4: WOL[2]; CUR[20]; (2 IDENTICAL) SET 5: PAY[7]; GLO[15]. (2 IDENTICAL) IDENTICAL SETS OF J-MINIGENES: SPECIFIC NOTES: 4) NG9'CL: THE AMINO ACID SEQUENCE IS TRANSLATED FROM THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CONA a) NGYCL: THE AMINO ACID SEQUENCE IS THANSLATED FROM THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CUNA.
 32) MCE': IT IS A CRYOIMMUNOGLOBULIN. THE AUTHORS ORIGINALLY DESIGNATED IT AS MCE. BUT IN ORDER TO DIFFERENTIATE IT FROM ANOTHER MCE SEQUENCED BY CAPRA ET AL., IT IS DENOTED AS MCE'.
 42) CLA: THE AMINO ACID RESIDUES FOUND AT POSITION 9 WERE GLY AND ALA.
 43) SHE': THE AMINO ACID RESIDUES FOUND AT POSITION 9 WERE GLY AND ALA.




ł

NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP III (coni'd)

NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP III (conid)
44) JH: THE NAME WAS GIVEN TO US BY THE AUTHORS. IT IS NOT INCLUDED IN THE PAPER.
58) WE: AT POSITIONS 20:20 AND 33 OF AMINO ACID SEQUENCE WERE FOUND BOTH LEU AND ILE. IN THE SAME SEQUENCE TWO RESIDUES WERE FOUND IN FOUND IN TOSITIONS 13:49:10:15:17:19:20:21:22 AND 20. THE SECOND RESIDUES WERE GLUVALLEU.GLY THR FRO GLUVAL THRLEU.SER AND VAL. RESPECTIVELY. A DETERMINATION WAS NOT MADE IN THE ARTICLE AS TO WHETHER THE SEQUENCE BELONGED TO SUBGROUP IOR TO SUBGROUP III.
81) AMYLOID WR: AMINO ACID RESIDUES FOUND AT POSITION 54 ARE LEU AND ALA.
82) VKAPPA3'CL: THE AMINO ACID REQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF CDNA FROM A MOUSE-HUMAN HYBRID CELL LINE.

WT PQ 1	N KAPPA LIGHT	1	2 VKAPPA IV GERMLINE	9 PB17IV CL	4 LEN	5 я.к.	6" L. TH.	77 TUR	в Ан	9 DA	10 DA-Н	11 DA-N	12 JAH	13 SCH	14 JUV	AMYLOID GAB	# OF SEQUENCES	AMINC
	0 1 ASP 2 VAL 4 5 6 GLN 7 SER 8 PRO 9	ASP ILE VAL MET THR GLER PRO ASP	CL ASP ILE VAL MET THR GLN SER PRO ASP	ASP ILE VAL MET THR GLN SER PRO ASP	ASP ILE VAL MET THR GLN SER ASP	ASP ILE VAL MET THR SERO ASX	ASP ILE VAL IEU THR SERO ASX	ASP ILE VAL MET THR GLN SER PRO SET	ASP ILE VAL MET THR GLN SEP 91x	ASP ILE VAL MET THR GLN SEA PRO ASP	ASP ILE VAL MET THR GLN SER PRO ASP	ASP ILE VAL IEU GLEAD SEROP ASS	ASP ILE VAL THR GLN SER ASP	ASP ILE VAL MET THR GLN SER PRO ate	ASP leu VAL MET THR GLN SEA PRO A90	ASP ILE VAL MET THR GLN PRO ASX	15 15 15 15 15 15 14 15 15 15	12 12 11 11 5 2
	10 LEU 12 ALA 13 VAL 14 SER 15 16 GLY 17 18 18 ALA	SER LEU ALA SE UY GLG ALA THR	SER LEUA VAL SER GLU GLU ARG ARG ALH R	SERUALAL VAR LELYUGARA ALA SEUYUGAALA ALA THE	SEA LALA VAL SER UGLUY GLUY GLUY ALA THE	ALA	GLY GLX ARG ALA THR	· THR	SEALALA VALA SEO GARGA ALA THE	SER LEA VAL SER GLY asp GLY asp ALA THR val	SELALA VALA SEO Gasin ALA THR vat	SERUALAL ALAL SEUY BIOUA ALA THOU	thr LEU ALA VAL	LEU ALA VAL	LEU ALA VAL	ALA THR ILE a30	14 14 11 11 11 11 12 12 12 12 10	1 1 2 2 1 1 2 3 1 1 3 3
	21 22 23 CYS 24 25 28 27 27A 27B	ILEN ASYS LYS SEEN SELN SELN LEA LEA LEA LEA LEA LEA	LE ASYS LYS SER SER SER SER SER LER LEUR SER LEUR SER	LENS LYS ARNRE SELEURARA SELEURARA SELEURARA SAS	LE ASYS		ASX	SET CYS ARG SER GLN SER VAL LEU	ASX CYS ARG ARG ALA GLX ARG VAL	GLN ALA	SUT CYS GLN ALA	GLN GLN ALA SER GLN VAL LEU TYR				asp CYS LYS	10 10 9 7 6 5 5 5 5 5 5 5 5	1 22 1 1 21 2
	27C LEU 27D TYA 27F SER 28 30 LYS 31 ASN 32 TYR 33 LEU	SER SER ASN LYS ASN LYS ASN LEU ALA	ASN ASN LYS ASN LYS ASN LEU ALA	SEA ASP ASN LYS ASN TYA LEU ALA	SEF ASP LYS ASP LEU AL							ASP LYS					4 5 4 4 4 4	211111111111111111111111111111111111111
	34 ALA 35 THP 36 TYR 37 GLN 38 GLN 39 LYS 40 PRO 41 GLY 42 GLN 43	ALA TRP GLN GLN GLN GLN GLN GLN GLN GLN GLN	TAP TYR GLN GLN LYS PRO GLY	TRP TYR GLN GLN PRO GLN GLN PRO	TRI TYI GU GU PRI GL GL PRI	0 T 7 7 MOY 20				PRI GLI ALI PRI	Y N A			÷			444 45555 5	
	44 PRO 45 LYS 48 LEU 47 LEU 48 ILE 49 TYR 50 TRP	PRO LYS LEU LEU ILE TYR TBP	PRO LYS LEU ILE TYR	PRO LYS LEU ILE TYR TRP ALA	PR LEE LEE TP ALE	S U U E R			<u></u>		S J I B P Y						9 5 5 5 5 5 5 5 4	
2	51 52 THR 53 THR 54 ARG 55 GLU 56 SER 57 GLY 58 VAL	ALA SER THR ARG GLL SEF GLY VAL	ARG GLU SER GLY VAL	SER THR ARG GLU SER GLY VAL	AF GL SE GI												4 4 4 4 4 4 4	
	59 PRO 80 ASP 81 ARG 62 PHE 63 GLY 64 GLY 65 SELY 66 SELY 67 SER 67 SER	PAC ASF ARC PHI SEF GLU SEL GLU THI	PRO ASP ARG PHE PHE SER SER GLY GLY GLY GLY	PRC ASP ARG PHE SEF GL\ SEF GL\ THF						SE SE SE	Ř						2445 55555 44	
FR3	69 THR 70 ASP 71 PHE 72 THR 73 LEU 74 THR 75 ILE 76 SER 77 SER 78 LEU	ASI PH TH LE TH ILE SEE SEE LE GL	ASP PHER THRU LEU THR LEU THR SER SER U	ASF PHI THE LEI THE SEE LEI GL								·				·	4 4 4 4 4 4 4 4 4 4 4 4 4	
	79 GLN 80 ALA 81 GLU 82 ASP 83 VAL 84 ALA 85 VAL 86 TYP 87 TYP 87 TYP	GLLGS AGS A ALA YYYYY	A ALA U GLU P ASP IL VAL A ALA IL VAL IR TYR	GLI GLI ASA VA AL YA TYY TYY			,	•									4 4 4 4 4 4 4 4 4	
1 000	BB CYS 89 GLN 90 GLN 91 TYR 93 93 94 95 95A	GL GL TY AS TF IL PF	N GLN GLN GLN FYR FYR FYR FYR FYR FYR FYR FYR FYR FYR	GL GL TY AS		ILN YR YR HR HR				ŗ							4 4 4 4 4	
Ā 3	95B 95C 95D 95E 95F 96	-				TYR		·								Q	3	
- F- 4	103 LYS 104 105 GLU 106 ILE	PIGGGT VG	HE			PHE GLN GLN THR LYS LEU GLU ILE											3 3 3 3 3 3 3 3 3 4 4 4 4 4	
	106A 107 108 ARG 109	L	YS	L A	YS	LYS ARG THR					ARG					·····	. 4 3 1	

HUMAN KAPPA LIGHT CHAINS SUBGROUP IV (cont'd) OCCURRENCES VARIABILITY OF MOST COMMON AMINO ACID



· · · · · · · · · · · · · · · · · · ·
ANTIBODY SPECIFICITIES. HUMAN KAPPA LIGHT CHAINS SUBGROUP IV
3) PB171V'CL: ANTI-STREPTOCOCCUS GROUP A CARBOHYDRATE WITH SPECIFICITY FOR NACETYL GLUCOSAMINE
5) R.K.: COLD AGGLUTININ WITH ANTI-PRIH ACTIVITY (BBC MEMBRANE ANTIGEN ON HUMAN EATTHACCTICS INACTIVITED OF THE LECTION DISTANCE
AND NEURAMINIDASE) 6) L.TH.: COLD AGGLUTININ WITH ANTI-PR2 ACTIVITY (RBC MEMBRANE ANTIGEN ON HUMAN, RAT AND GUINEA PIG ERYTHROCYTES INACTIVATED BY PROTEOLYTIC ENZYMES AND NEURAMINIDASE)
7) TUR: COLD AGGLUTININ WITH ANTI-PR ACTIVITY
REFERENCE: HUMAN KAPPA LIGHT CHAINS SUBGROUP IV
1) VJI'CL: KLOBECK.H.G. BORNKAMMM.G.W. COMBRIATO.G. MOCIKAT.R. POHLENZ.H.D. & ZACHAU.H.G. (1985) NUC.ACIDS RES., 13,6515-6529. (CHECKED BY AUTHOR 02/25/86)
BY AUTHOR 02/25/86) 2) VKAPPA IV GERMLINE'CL: KLOBECK,H.G.,BORNKAMMM,G.W.,COMBRIATO,G.,MOCIKAT,R.,POHLENZ,H.O. & ZACHAU,H.G. (1985) NUC.ACIDS RES.,13, 6515-6529.
6515-6529. 3) PB17/VCL: MARSH.P.,MILLS,F. & GOULD.H. (1985) NUC.ACIDS RES., 13,6531-6544. (CHECKED BY AUTHOR 03/19/86 WHO CORRECTED A MISPRINT IN THE ORIGINAL PAPER FOR RESIDUE 50)
A A A A A A A A A A A A A A A A A A A
THE ALARY DE VENIOR LA MELLANC D.C. ID FEIZLT & KUNKEL H.G. (19/2) PHOUNAL ACAD, SOLUGA, S
9) DA: WANG A.C. ZHANG H.S. BONEWALD L. TUNG E. BOUVET J.P. & LIACOPOULOS.P. (1985) MIANIN WHICH STWART SO COLI (1995)
10) DA-H: BOUVETUP, LIACOPOULOSP, PILLOTU BANDAR, TUNGE, & WANGA,C. (1980) JIMMUNOL, 129, 219-220. (CHECKED BI AUTHON OBCHON,
11) DA-N: BOUVETJP, LIACOPOULOS P. PILLOT J. BANDA, R. TUNG, E. & WANG, A.C. (1980) J.IMMUNOL. (22) 159-1524.
12) JAH: SLETTENK, HANNESTAD,K, & HARBOEM, (1974) SCANDJIMMUNOL,3,219-222. (CHECKED BY AUTHOR 12/05/77)
IN AGUE OF CATTOR A HADROF M (1974) SCAND J.IMMUNUL 3.219-222. (CHECKED BY AGUNON (DOGU))
14) JUV: SLETTEN, HANNESTAU, A HAROUEIN, (1974) C. (1980) IN AMYLOID AND AMYLOIDOSIS,G.G.GLENNER,P.P.E. COSTA & F.DE FREITAS EDS., 15) AMYLOID GAB: PRAS., FRANGIORE, & FRANKLIN,E.C. (1980) IN AMYLOID AND AMYLOIDOSIS,G.G.GLENNER,P.P.E. COSTA & F.DE FREITAS EDS., EXCERPTA MEDICA AMSTERDAM OXFORD-PRINCETON.249-252. (CHECKED BY AUTHOR 11/18/81)
NOTES: HUMAN KAPPA LIGHT CHAINS SUBGROUP IV
IDENTICAL SETS OF FRAMEWORK SEGMENTS:
FR1: SET 1: VJ/CL[11]VKAPPA IV GERMLINE/CL[2].PB17IV/CL[3].R.K.[5]. (4 IDENTICAL) SET 2: LENIA]R.K.[5]. (2 IDENTICAL)
FR2: SET 1: VJ/CL[1].VKAPPA IV GERMLINE'CL[2].P817IV'CL[3].LENIA! (a IDENTICAL HUMAN V:KAPPA.IV: ALSO 2 HOMMAN V:KAPPA.IV: ADSO 2 HOMMAN V:KAPPA.IV:
PC923(NZB)(56) PC4050(NZB)(57), V-21B)5KB°CL(58), 11949(62); 1 MOUSE V-KAPPA-VI: BPPCB1A CL(54); AND 15 HABBIT V-KAPPA- K9-335-(119) 3368(20), BS-5(38), BS-1(39), K49-501(45), 3547(47), K4820(57), K30-267(61), 311(65), 4422(66), 17D9°CL(66), 4192(71, 4363(85), 120(103), K-25(112),)
FR3: SET 1: VUICLI1, VKAPPA IV GERMLINE'CL(2), PB17IV'CL(3), LEN(4). (4 IDENTICAL)
EDA: SET 1 PB17V/CU3L (IDENTICAL TO 3 HUMAN V-KAPPA-): AU[2] GAL(1)[38] CU1(1)(): 2 HUMAN V-KAPPA-3(CU3))
SET 2: LEMIA: (DENTICAL TO 1 HUMAN V-KAPPA-III: VOL21/FAY17)/BICTI1/GLUISICUAL20./REESTIGAT, AND 8 HUMAN V-KAPPA-III: SET 2: LEMIA: (DENTICAL TO 3 HUMAN V-KAPPA-III: AGI71/DENI65/BIBB3); NEU[S],GOT[6],GAR[10],FA[21],FA[21],IAR/BL4TCU23]) SET 3: VUICL[1]. (DENTICAL TO 1 HUMAN V-KAPPA-III: SONIBJ.)
IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS:
CDR1: SET 1: VJI'CL[1],VKAPPA IV GERMLINE'CL[2]. (2 IDENTICAL)
CDR1: SET 1: VJICL[1],VKAPPA IV GERMLINE CL[2], [E IDENTIONE) CDR2: SET 1: VJICL[1],VKAPPA IV GERMLINE CL[2],PB17IV/CL[3],LEN[4]. (4 IDENTICAL HUMAN V-KAPPA-IV; ALSO 1 MOUSE V-KAPPA-VI: KPNI6 'CL[70].

CDR3:

IDENTICAL SETS OF J-MINIGENES; IDENTICAL SETS OF J-MINIGENES; SET 1; FB171V'CL[3], (IDENTICAL TO 1 HUMAN V-KAPPA-I; AU[2]; 1 HUMAN V-KAPPA-II; RPM1-6410'CL[16]; AND 2 HUMAN V-KAPPA-III; PIE[11],VKAPPA3'CL[82].)


+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

AT POSITION RESIDUES 22 (SER.ASP.ASN) 96 (TRP.TYR) 104 (LEU.VAL)

.

711F?

н	UMA	I LAMBDA LIC		INS SU	BGRO	UP I																			
-	C	INVARIANT RESIDUES	NEWN	4 HA	3 LR	4 NIG -64	NEW	/ BL2 CL	wah	8 NIG -77	VOR	10 RHE	LÖC	OKA	13 AMYLOID EPS	14 HBJ 7	cox	16* КОН	17 HS 92	18 HS 78	19 NIG -51	20 HS 94	21 HBJ 11	22 BJ 98	23 MZ
	56789 10	PCA(.95) SER VAL(.95) LEU THR GLN(.95) PRO PRO SER	PCA SER VAL LEU THR GLN PRO PRO SER	PCA SER VAL LEU THR GLN PRO SER	PCA SEA VAL LEU THA GLNO PRO SEA	PCA SEA VAL LEU THR GLN PRO SER	SER VAL LEU THR	GLN SER VAL LEU THR GLN PRO PRO SER	THR GLN PRO PRO SER	PCA SER VAL LEU THR GLN PRO SER	LEU	PCA SER VAL LEU THR GLN PRO SER	PCA SER VAL LEU THR PRO SER	PCA SEA VAL LEU THR GLN PRO SER	VAL LEU THR GLN PRO SER	PCA SER VAL LEU THR GLNO PRO SER	PCA SER VAL LEU THR GLAO PRO SER	PCA SER VAL LEU THR GLX PRO SER	PCA SEA VAL LEU THR PARO SEA	PCA SER ala LEU THR GLN PRO SER	PCA SER VAL LEU THR GLN PRO SER	PCA SER VAL LEU THR GLN PRO SER	PCA SER VAL LEU THR GLN PRO SER	тня	
F R 1	11 12 13 14 15 16 17 18	SER PRO(.95) GLY VAL(.95)	VAL SEA GLY ALA PAO GLY GLN ARG VAL	VAL SEA GLY Ibr GLY GLN ARG VAL	VAL SER GLY ALA PRO GLY glu ARG VAL	VAL SER ala ALA PRO GLY GLN GLN VAL	GLY GLN Iys VAL	VAL SEA ala ALA PRO GLY GLN Iys VAL	AIB SER GLY Ihr GLY GLN ARG VAL	BIB SER GLY Thr PROY GLN ARG VAL	BIA SER GLY Ibr GLY GLN ARG VAL	BIB SEA GLY Ihr GLY GLN AAG VAL	ala SER GLY thr PRO GLY GLN ARG VAL	ala SER GLY thr PRO GLY GLN ARG VAL	leu SER ALA PRO GLY GLY ARG VAL	ala SER GLY PRO GLN GLN VAL	ala SER GLY Ihr GLY GLN ARG VAL	ala SER GLY thr PRO GLY GLX ser VAL	VAL SER GLY ALA PRO GLY GLN ARG	VAL SER GLY ALA PRO GLY GLN Ihr VAL	ala SEA GLY Val PRO GLY GLN Ser VAL	VAL SER ala ALA PRO GLY GLN ARG	ala SER GLY thr PRO GLY GLN ARG	SER als ALA VAL	ala
	20 21 22 23 24	ILE(.95) SER(.95) CYS	THR ILE SER CYS THR	THR ILE SER CYS SER	DIU ILE SEA CYS SEA	THR ILE SER CYS SER	THR ILE SER CYS SER	THR ILE SER CYS SER	THR ILE SER CYS PHE	THR ILE SER CYS SER	THR ILE SER CYS SER	THR ILE SER CYS THR	THR ILE SER CYS SER	THA ILE SEA CYS SEA	SER CYS SER	THR ILE SER CYS SER	THR ILE SER CYS SER	THR ILE SER		тня	ile ILE SER CYS SER			THR ser ite CYS	ile ILE SER CYS SER
	25 26 27 27 27 27 27	GLY A B C	GLY SER SER	GLY GLY SER	GLY	GLY SEA SER	GLY GLY SER	GLY SEA SEA	GLY SER SER	GLY SER THR	GLY GLY ASN	GLY SEA ALA	GLY SEA SEA	GLY SER GLY	GLY SER SER	GLY SEA	GLY SEA SEA				GLY SEA SEA				GLY SER SER
CDR 1	270 275 275 28 29 30 31 32 33		SER ASN ILE GLY ALA GLY ASN HIS VAL	SER GLY THR GLY ASN ASN TYR VAL		SER AS. ILE GLY ASP ASN PHE VAL	THR ASN ILE GLY ASN TYR VAL	SER ASN ILE GLY ASN TYR VAL	SER ASN ILE GLY ARG TYR TYR VAL	SER ASN ILE GLY SER ASN THR VAL	PHE ASP ILE GLY ARG ASN SER VAL	THR ASP ILE GLY SER SER SER VAL	SER ASN ILE GLY GLU THR ASN SER VAL	SER ASN ILE GLY SER HIS HIS THR VAL	SER ASN ILE GLY LYS ASN TYR VAL		SER LEUYR SESNN LEUYR SESNN VAL				SER ASN ILE GLY ARG ASN THR VAL				SER ASN MET
	34 36 37 38	TRP	LYS TRP TYR GLN GLN	TYR TYR TYR GLN GLN		SER TYP GLN GLN	SEA HIS GLN HIS	SEA TAP TYA GLN GLN	TYR TRP TYR GLN GLN	THR TRP TYR GLN HIS	ASN TRP TYP GLN VAL	ILE TRP TYR GLN GLN	SER TRP TYR GLN HIS	ASN TRP TYR HIS GLN	ASP TRP TYR GLN GLN		ASN TRP TYR ARG HIS				ASN TRP TYR GLN GLN				
F #2	39 40 41 42 43 44	PRO GLY PRO	LEU PRO GLY THR ALA PRO	LEU PRO GLY THR ALA PRO		LEU PRO GLY THR ALA PRO	LEU PRO GLY THR ALA PRO	VAL PRO GLY THA ALA PRO	LEU PRO GLY THA THA THA PRO	LEU PRO GLY THR ALA PRO	HIS PRO GLY THR ALA PRO	VAL PRO GLY LYS ALA	LEU PRO GLY THR ALA	PHE PRO GLY THR ALA	LEU PRO GLY THR ALA		LEU PRO GLY THR ALA				VAL PRO GLY ALA ALA				
	45 46 47 48 49	LEU	LYS LEU LEU ILE PHE			LYS LEU LEU ILE TYR	LYS LEU LEU ILE TYR	LYS LEU LEU ILE TYR	LYS LEU LEU ILE TYR	LYS LEU LEU ILE TYR	ARG LEU LEU ILE TYR	PRO LYS LEU LEU ILE TYR	PRO LYS LEU LEU ILE TYR	PRO LYS LEU LEU ILE TYR	PRO LYS LEU LEU RLE PHE		PRO LYS LEU VAL ILE TYR				PRO LYS LEU LEU VAL				
CDR2	50 51 52 53 54 55 56	SER	HIS ASN ASN ALA ARG	ARG ASP LYS ARG PRO SER		ASP ASN ASN LYS ARG PRO SER	GLU ASP ASN LYS ARG PRO SER	ASP ASN ASN LYS	LYS ASP ASN GLN ARG	SER ASN ASP GLN	SER SER ASP GLN ABG	TYR ASN ASP LEU	GLU ASP ASN SER	ARG ASN ASP GLN ARG PRO SER	ASN ASN ASN LYS ARG		SEA ASP SER GLN ARG PRO SEA				SER ASN ASN GLN TAP PRO				a tm
	57 58 59 60 61 62	GLY		GLY VAL PRO ASP ARG PHE		GLY ILE PRO ASP ARG	GLY ILE PRO ASP	GLY ILE PRO ASP	GLY VAL PRO ASP	GLY VAL PRO HIS	GLY VAL PRO	GLY	GLY VAL SER ASP	GLY VAL ASP ARG PHE	ABG		GLY VAL PRO ASP ARG				SER GLY VAL PRO ASP ABG				
	63 64 65 66 67	SER SER LYS SER GLY	SER VAL SER LYS SER	SER GLY SER LYS SER		PHE SEA GLY SEA LYS SEA GLY	SER	SER	SER GLY SER LYS	GIV	SER	ALA SER	SER ALA SER	SER GLY SER LYS	ARG PHE SER GLY SER LYS SER		ILE SER ALA SER LYS SER				PHE SER GLY SER LYS				
FR3	68 70 71 72 73	GLY SER ALA LEU	GLY SER ALA THR LEU	GLY THR SER ALA SER LEU		THR SER ALA THR	GLY THR SER ALA THR LEU	THR SER ALA THR	THR	ALA SER ALA	THR	THR SER ALA	THR SER	SÉR GLY THR SER ALA SER LEU	SER GLY THR SER ALA THR LEU		GLY THR SER ALA SER			;	SER GLY THR SER ALA SER				
3	74 75 76 77 78	ILE GLY LEU	ALA ILE THR GLY LEU	ALA ILE SER GLY LEU		GLY ILE THR GLY	ALA ILE THR GLY	GLY ILE THR GLY	ALA ILE SER GLY		ALA ILE SER GLY	ALA ILE SER GLY		LEU ALA ILE SER GLY LEU	GLY ILE THR GLY LEU	í	LEU ALA ILE SER GLY LEU				LEU ALA ILE SER GLY LEU				
	79 80 81 82 83	GLU	GLN ALA GLU ASP GLU	ARG SER GLU ASP GLU		THR GLY ASP GLU	ARG THR GLY ASP GLU	GLN THR GLY ASP GLU	ARG SEA GLU ASP 3LU	GLN SER GLU ASP GLU	GLN SER GLU ASN GLU	GLU SER GLU ASP GLU	GLN PRO BLU ASP GLU	GLN SER GLU ASP GLU	GLN THR GLY ASP GLU	0.50					HIS SER GLU GLU				
	84 85 86 87 88 89	TYR CYS	ALA ASP TYR TYR CYS	ALA HIS TYR HIS CYS		CYS	CYS	CYS (ALA ASP TYR TYR CYS ALA	JYS -	ALA ASP TYR PHE CYS	ASP TYR TYR CYS	CYS I	ALA ASP TYR TYR CYR	ALA ILE TYR TYR CYS		SER SP TYR TYR TYR SYS			, j F	ALA ASP IYR PHE CYS				
C	90 91 92 93 94 95	`	GLN SEA TYR ASP ARG SER	ALA ALA TRP ASP TYR ABG	Ś	ASP SER : SER :	ASP SER /	THR TRP ASN A	ALA T TRP T ASP J ASP J SER S	ASP ASP	ALA THR TRP ASP ASP SER	ASN ASP	ASP /	ALA IAP ASP ASP SER	GLY THR TYR ASP ASN ARG	S A	SP SP				ALA THR TRP ASP ASP SER				
D R 3	95A 95Đ 95C 95D 95E		LEU 	LEU SER ALA			ALA (SER EU ASN GLY			LEU I ASP / /AL C		ARG	Ğ	EN EU SP			40	SLY				
	95F 96 97 98 99 99 00	PHE	ARG VAL PHE GLY	VAL VAL PHE GLY	. F	SLY MET PHE P	VAL VAL	TRP T	AP P AL V HE F	AL PHE SLY	PAO P VAL (PHE P SLY (RO A	AL F	HE ILY	SER VAL PHE GLY GLY	P V P G	RO AL HE			ድ	RO AL HE				
F 7 R 1 4 1	01 1 02 03 04 05	GLY THR THR	GLY THR LYS LEU THR	GLY GLY THR GLN LEU THR	U T A	ARG 1 ARG 1 AL N	THR T INR T VAL L INR T	INA T	SLY O HR T HR L	ALY CONTRACT	3LY (3LY (1HR 1 .YS 1	SLY C HR T YS L EU V	HR T	HE ILY ILY HR YS EU HR	GLY GLY THR ASN VAL THR	G G T L	LY HA YS AL HA			. GGT 1.	LY LY HR YS EU				
1 1 1	06 06A 07	GLN PRO	LEU ARG	VAL LEU ARG GLN PRO	Ľ	VAL I EU L BLY (LEU L	VAL N .EU L GLY S	EU G	AL I ILN L	JAL N EU L SLY G	AL N EU L SLY G	AL N EU L	HR AL EU LY	THR VAL VAL GLY GLN PRO	v L G	HR EU LY LN RO			Ť V L G	HÀ AL EU LY LN RO				
								<i>.</i>	1 S			•	- /			. "				· P	n0				



Ù

ANTIBODY SPECIFICITIES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP I

1) NEWM: ANTI-3-(3-HYDROXY-3',7',11',15' TETRAMETHYL HEXADECYL) 2-METHYL 1.4 NAPHTHOOUINONE(VIT.K10H) 16) KOH: ANTI-HUMAN GAMMA G GLOBULIN

REFERENCE: HUMAN LAMBDA LIGHT CHAINS SUBGROUP I

1) NEWM: CHEN.B.L. & POLJAK.R.J. (1974) BIOCHEMISTRY 13,1295-1302. (CHECKED BY AUTHOR 01/24/78)

NEWM: CHRENALL'S FOLGARINJ, (13/4) BIOLCHEMISTRT, 13/129 BIOLCHEM, 245,4475-4487. (CHECKED BY AUTHOR 06/15/83)
 LA: SHINODAT, TITANIK, & PUTNAM, W. (1970) BIOLCHEM, 245,4475-4487. (CHECKED BY AUTHOR 06/15/83)
 LA: CAULIN-GLASER, T., PRELLIF, & FRANKLIN, E.C. (1982) JLAB.CLIN, MED, 99,845-851. (CHECKED BY AUTHOR 12/10/82)
 NGG48: TONOIKE, H., KAMETANI, F., HOSHIA, SHINODA, T. & ISOBE, T. (1983) BIOCHEM, BIOPHYS, RES.COMMUN., 126,1228-1234.
 NEW: LANGER, B., STEINMETZ, KAYNEM, & HILSCHMANNN, (1968) Z. PHYSIOL.CHEM, 349,945-951.
 BL2 'CL: TSUJIMOTO, Y. & CROCE, C. M. (1984) NUC.ACIDS RES., 12,8407-8414.

7) WAH: TAKAHASHI,Y.,TAKAHASHI,N.,TETAERT,D. & PUTNAM,F.W. (1983) PROC.NAT.ACAD.SCI.USA.80,3686-3690. (CHECKED BY AUTHOR 06/15/83) 8) NIG-77: TONOIKE,H.,KAMETANI,F.,HOSHI,A.,SHINODA,T. & ISOBE,T. (1985) BIOCHEM.BIOPHYS.RES.COMMUN.,126,1228-1234.

- 9) VOR: ENGELHARD.M. HESS.M. & HILSCHMANN.N. (1974) Z.PHYSIOL.CHEM. 355.85-88; ENGELHARD.M. & HILSCHMANN.N. (1975) Z.PHYSIOL.CHEM. 356, 1413-1444.

- 17) HS92: HOODLL & EIN.D. (1968) NATURE.220.764-767; (1968) SCIENCE.1662.679-681. 18) HS78: HOODLL & EIN.D. (1968) NATURE.220.764-767; (1968) SCIENCE.1662.679-681.
- 19) NIG-51: TAKAHASHI.N., TAKAYASU.T., SHINODA, T., ITO, S., OKUYAMA, T. & SHIMIZU, A. (1980) BIOMED.RES., 1, 321-333. (CHECKED BY AUTHOR 01/28/81) 20) H\$94: HOOD, L. & EIN, D. (1968) NATURE, 220, 764-767; (1968) SCIENCE, 1662, 679-681.

- HB311: HOODLL.GRAY.W.R.SANDERS.G. & DREYER.W.J. (1986) SCIENCE, 1002.619581.
 HB311: HOODLL.GRAY.W.R.SANDERS.G. & DREYER.W.J. (1987) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,133-145.
 BJ98: BAGLIONIC. (1967) BIOCHEM.BIOPHYS.RES.COMMUN.26.82-89.
 MZ: MILSTEIN.C. FRANGIONE.B. & PINK.J.R.L. (1967) COLD SPRING HARBOR SYMP. QUANTITATIVE BIOL.32,31-35. (CHECKED BY AUTHOR 10/17/77)
 MZ: MILSTEIN.C. JR. & HOODL. (1970) PROC.NAT.ACAD.SCI.USA.66.975-982.

NOTES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP I

IDENTICAL SETS OF FRAMEWORK SEGMENTS:

- EB1: SET 1: WAH[7],NIG-77[8],VOR[9],RHE[10],LOC(11),OKA[12], (6 IDENTICAL)
- SET 1: NEWMI1].AMYLOID EPS[13]. (2 IDENTICAL) SET 2: HA[2].NIG-64[4]. (2 IDENTICAL) SET 3: NIG-77[8].LOC[11]. (2 IDENTICAL) FR2:

- FR3:
- FR4:
- SET
 3: NIG-77(8),LOCI11, (2: IDENTICAL)

 SET
 1: NIG-78(8),LOCI11, (2: IDENTICAL)

 SET
 1: NEWMI1, [DENTICAL TO 1 HUMAN V-LAMBDA-II; WHI3; AND 1 HUMAN V-LAMBDA-V; BO[1],)

 SET
 2: NEWISI,UGRISI,COX152, (3: IDENTICAL)

 SET
 2: NEWISI,UGRISI,COX152, (3: IDENTICAL)

 SET
 3: NIG-77(8),LOC111, (DENTICAL TO 1 HUMAN V-LAMBDA-II; WHI3; AND 1 HUMAN V-LAMBDA-V; BO[1],)

 SET
 3: NIG-77(8),LOC111, (DENTICAL TO 1 HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; MES[2],ES492(8),TRO[14],

 SET
 3: BLZ (DISTICAL TO 1 HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; MES[2],ES492(8),TRO[14],

 SET
 3: BLZ (DISTICAL)
 10: NITCAL HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; MES[2],ES492(8),TRO[14],

 SET
 3: BLZ (DISTICAL)
 2: NEWISI, CAMBIDA-II; ALSO 5 HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; MES[2],ES492(8),TRO[14],

 SET
 3: BLZ (DISTICAL)
 2: NITCAL HUMAN V-LAMBDA-II; ALSO 5 HUMAN V-LAMBDA-II; ALSO 5

IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS:

CDR1:

CDR2:

- SET 1: NIG-64[4],BL2 'CL[6], (2 IDENTICAL) CDB3: SET 1: VOR[9].NIG-51[19]. (2 IDENTICAL)
- IDENTICAL SETS OF J-MINIGENES;

 - SET 3: VPIJAINIGENES; SET 1: NEWISI, (IDENTICAL TO 1 HUMAN V-LAMBDA-VI: AMYLOID-API[1],) SET 2: BL2 CLI6J, (IDENTICAL TO 2 HUMAN V-LAMBDA-VI: SUT[2],THO[4], AND 24 MOUSE V-LAMBDA: MOPC104E[1],J558[2],X5104[3],HOPC1[4], J988[5],H2051[6],W3159[7],Y543[1],Y548[0],Y5659[1],MOPC51[L]](2],ST913[1],Y544]14[Y5669[15],S176[16], H2020[17],RPC20[18],IIS_303[JAMBDA-CL[19],S43*CL[21],S2H5*CL[39],S2E9*CL[39],S1F12*CL[40],IG 25LAMBDA-CL[41],) SET 3: VOR[9](COX],15], (2 IDENTICAL)

SPECIFIC NOTES:

24) FUL: SOX AND HOOD HAVE REPORTED FOUR HUMAN V KAPPA AND ONE V LAMBDA CHAINS WITH ASN-SER/THR TO CONTAIN CARBOHYHDRATE.

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

AT POSITION RESIDUES 34 104 (SER.ASN) (LEU.VAL)

HUM		AMBDA LIGH	1 NIG -84	MES	з	NEI	к ⁵ я	8 81M	SLA	8 ES492	WEIA	10 тос	11 SM #	12 HS 68	13 HS 77	14 TRO	15 80H	18 NIG -58	17 VIL	18 HBJ 15	19 19 8	20 HS 70	21 WIN	80A	PRE	24 HS 86
	01234 58789	SER(.96) LEU(.96) GLN(.96) SER(.96)	PCA SER ALA LEU THR GLN ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SEA LEU FGLNO ALE SER SER	PCA SER ALA LEU THR GLN ALA SER	PCA SER ALA LEU THR GLN ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SEA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLN ALA SER	PCA SER ALA LEU THR GLN ALA SER	PCA SER ALA LEU THR GLN PRO Brg SER	PCA SER ALA LEU THR GLN PRO arg SER	PCA SER ALA LEU THR GLN PRO arg SER	HIS SER ALA LEU THR GLN PRO ALA SER	PCA SER ALA LEU THR GLO ALA SER	PCA SER ALA LEU BIB GLN PRO ALA SER	PCA SER ALA LEU SOF GLN PRO ALA SER	PCA SER ALA LEU THR GLN PRO pro arg	PCA SER ALA LEU THR GLX PRO arg SER	PCA SER ALA LEU THR GLN 9er pro SER	PCA SER pro LEU ala GLN PRO ALA SER
F A 1	10 11 12 13 14 15 16 17 18 19	SER(.96) SER(.95) PRO(.96) GLY	VAL SEA GLY SEA GLY GLN SEA ILE	VAL SER GLY SEA PROY GLN SER ILE	VAL SER GLY SER PRO GLN SER ILE	VAL SER SER SER GLN SER SER ILE	VAL SER GLY SER GLY SER GLN SER ILE	VAL SER SER GLR GLR GLN SER ILE	VAL SERY SELY SELY SER GLN SER ILE	VAL SER GLA SER GLA SER GLN SER ILE	VAL SERY SER PRLY SER GLN SER ILE	VAL SER BIA SER PROY GLN SER ILE	VAL SER SER OGLN SER GLER	VAL SER GLY SER PROY GLN SER ILE	VAL SER SER PROY GLN SER ILE	VAL SER GLA SER GLA SER GLA SER Val THR	VAL SERY SER GLN SER GLN SER SER THR	VAL SER GLY SER OGLY GLN SEA Iou THR	VAL SER GLY SER IOU GLY SER ILE ILE THR	VAL SER GLY SER OLN GLN ILE THR	VAL SER GLY SER PRO GLY GLN SER ILE THR	VAL SER GLY SER PROY GLN SEE ILE THR	VAL SER GLY SER GLN SER Val THR	VAL SEA GLY SEA PAO GLY his SEA Val THR	ala SER GLY SER PRO GLY GLN SER Vai THR	VAL SEA SEA SEA PRO GLY SEA ILE THA
	20 21 22 23 24 25 26 27	THR CYS	THR ILE SER CYS THR GLY THR THR	THA ILE SEA CYS THA GLY THA SEA	THR ILE SER CYS THR GLY THR SER	THR ILE SEA CYS THR GLY THR THR	THR ILE SER CYS	THR ILE SER CYS	THR ILE SEA CYS	ALA GLY THA HIS	THR val SER CYS ALA GLY HIS THR	THR ILE SER CYS THR GLY THR THR	THA ILE SER CYS THR GLY ASX SER	тня 	THR	ILE SER CYS THR GLY THR SER	ALA GLY THA SER	SER CYS SER GLY ALA PRO	ILE SER CYS THR GLY THR SER				THR GLY SER SER TYR	ILE SER CYS ILE GLY THA SER	ILE SER CYS	
C D R 1	27A 27B 27C 27D 27E 27F 28 29 30 31 323 31		SER ASP VAL GLY GLY ASP PHE VAL SER	SESP VAL GLY GLY ASN TYAL SEA	SEP ASSL VALY SEP ASSL STYN ASHE VAL SEP	SEP VAL GLY SEP VAL SEP VAL SEP VAL				SER ASPL ASPL ASPL ASPL ASPL ASA ALA	SASALA PRNRE SASALA PRNRE SASALA SESSILE R	ASP ILEY GLY STYR STYR VAL SER	SER VAL GLY			SER VAL GLY ALA TYR SER VAL SER	SEPLY ASS VALY GLV ASS PHEL SER	CYS ASP VAL ASP GLYS GLUS SER VAL SER	SER ASP VAL GLY TYR ASN TYR VAL SER				SER ASN VAL THR GLYR ASN HIS VAL SER	SER ASN VAL GLY ASP TYR LYS TYR VAL SER		
F R 2	34 35 36 37 38 39 40 41 42 43 44 45 46	TRP PRO PRO	TRP TYR GLN GLN HIS PRO GLY LYS ALA PRO LYS	TAPENN SOLYSA PRESULT	TRPR TYLN GLN HISO GLYS ALA PROY LYS LEU	TAPA TYPA GLN GLN ASO GLN CAS CAS CAS CAS CAS CAS CAS CAS CAS CAS				TRP TYR GLN LEU HISO GLY ILE ALA PRS LEU	TRP PHE GLN GLN HIS PRO LYS ALA PRO LYS	TAP TYR GLN GLN TYR GLY LYS ALA PROS LYSL				TAP TYR GLN GLN HIS GLY LYS ALA PRO LYS LEU	THP TYPA GLN SOL BRLYSA CLYSA CLYSA CLYSA LEU	TAPAN GUN SOY BOLYSA OSU LYSA PASU LEE	TRP PHEN GLN SOLA GLN SOLA GLN SOLA GLN PROYALA OSU LEC				TRP GLN GL SPOY LYSL PRSUL V ROSUL V ROSUL MET	TAP TYR GLX GLX PROY LYS A PROY LYS LYS LYS LYS LYS LYS LYS LYS LYS LY		
CD#2	47 48 49 50 51 52 53 54 55	ILE	LEU ILE TYR ASP VAL ASN SER ARG PRO	ILE PHE ASP VAL SER GLU ARG PRO	ILE ILE TYR ASP VAL THR TYR ARO SER	MET ILE TYR GLU GLY ASN LYS ARO SER				MET ILE PHE VAL SER ASN ARG SER	LEU ILE TYR ALA VAL THR PHE ARO SER	LEU ILE PHE ASP VAL ASN SER ARG PRO SER				MET ILE PHE VAL THR LYS ARO SER	ILE ILE TYA GLY VAL ASN LYS ARO SER	GLY PHE SER ASN	ILE SER GLU ARGN ARGN ARO SER				ILE TYR ASP VAL ASP LYS ARG PRO SER	GLU VAL SEA SEA ARG SER		
	56 57 58 59 60 62 63 65 65 65 65 65 67	ARG SER GLY SER	SER GLY ILE SEA ASNO AREE SER SER SER SER SELY SGLY	SER GLY SEP SASGER SASGER SES SES SES SES SES	GLY ILE SER	GLYL RNGER YRSGLYSGLYSGLY		-		GLY VAL SENG ARGEA SEA SEA SEA SEA SEA SEA SEA SEA SEA S	GLY GLE PAU ARG PHER SER SER SER SER SER SER	GLY VAL				GLY VAL PROP ASPG LER GLER SERS GLY GLY	GLY VAL PRO TYR ARGE FHER SER GLY SER GLY GLY	GLY VAL PRO LEU ARG PHE SEA	GLY SARAPS GELERA		PHE SER GLY SER LYS		GLY VAL PASPGER SESSEA SESSEA	GLY VAL PRO ASP ASP SER SER GLY SER SER SER SER		
FR 3	68 69 70 71 72 73 74 75 76 77 78 79	ALA LEU THR SER GLY	GLY ASN THR ALA SER LE R ILE SELY LE U GLN	ASN THR ALA SER LEU THR ILE SER LEU	ASN THR ALA SER LEU THR SER GLY LEU	LYS THR SER LEU THR ILE SER LEU				ASN ALA SER LEU THR ILE SELY LEU GLN	ASN THRA SEEU THE SEEU THE SEEU LEU LEU	ASN THR ALA SER LEU THR ILE SERY LEU GLN				ASP THR SER LEU THR SER LEU LEU ARG	ASN THA SER LE R ILE SER LE C LE C	ALA ALA SEU THE SELU SELU GLU OLN	ASN THR SERU THR SERU LER LEU GLN				ASN ALA SEU LE R SGLU SGLU GLN	LEU		
	80 81 83 84 85 86 85 86 87 88	GLU ALA TYR CYS	ALA GLU ASP GLU ALA ASP TYR TYR CYS	ALA GLU ASP GLU ALA ASP TYR TYR CYS	ALA GLU ASP GLU ALA ASP TYR TYR CYS	VAL GLU ASP GLU ALA ASP TYR TYR CYS				ALA GLU ASP GLU ALA TYR TYR CYS	PRO ASP GLU ALA ASP TYR PHE CYS MET	ALA GLU ASP GLU ALS FYRE CYS SER				ALA ASP GLU ALA ASP GLU ALA TYR CYS	ALA HIS TYR TYR CYS	ALA ASP TYR TYR	ALA ASP TYR TYR CYS				ALA ASN GLA ASNU ASPA TYPA SER SER SER	ALA ASX TYR TYR CYS		
CDR3	89 90 91 92 93 94 95 95 95 95 95 95 95 0		SEA SEA PHE THA THA ASN SEA ARG	GLY SER ASN THR	SER ASX SER THR	ALA GLY ASX SER THR				SERE SERE THR ASP THR GLN LEU	SEA TYR LEU SER ALA SEA	SEP GLY THA				CYS SER TYR ALA GLY ARG TYR SER	ARG PHE THR	SER SER	SETTAR AND				GLY GLY THA TYA SEA	SEF		
 F	95E 95F 96 97 98 99 100 101 102 103	PHE GLY GLY THR	ALA VAL PHE GLY GLY THR LYS	тня	ARG LEU PHE GLY GLY THR LYS		VAL PHE GLY THR GLY THR GLN	VAL GLY THA GLY THR AAG	GLY THA GLY THR LYS	VAL VAL PHE GLY GLY THR LYS	PHE VAL PHE GLY SER GLY	ILE ILE PHE GLY GLY THP				VAL ILE GLY GLY GLY THR	GLY GLY GLY THR	VAL ILE PHE GLY ALA GLY THR	VAL VAL PHE GLY GLY GLY THR				LEU ILE PHE GLY GLY THA LYS	PHE GLY THP GLY THP	GLY THF GLY THF	1
	103 104 105 106 106A 107 108 109	VAL LEU PRO	LTS LEU SER VAL LEU GLY GLN PRO	LEU THR VAL LEU GLY	ARG	LEU				LEU THR VAL LEU GLY GLN PAO	LYS VAL THA VAL LEU ARG GLN PRO	LEU	;		,	LEU THR VAL LEU GLY	LEU THA VAL LEU		LEU THR VAL LEU GLY GLN				LYS J.EU THR VAL LEU GLY GLN	LEU GLY		

			•				-
	_	25 WAL	25 4A CL	* 05	ES AMI	F OCCUBBENCES	VARIABILIT
	2 3 4	LEU	gin Ihr val val	26 26 26 26	3090 1000	24(PCA) 25(SER) 23(ALA) 25(LEU)	3.3 2.1 3.4 2.1
	8 9	SER	glu pro SER	26 26 26 26 26	1 3 2	23(THR) 2 26(GLN) : 25(GLN) 24(PRO) 18(ALA) 25(SER)	1, 3.4 3.3 4.3 2.1
R 1 1 1: 1 1:	1 2 3	ala SER GLY thr	leu Ihr val SER	26 26 26 26	3 2 4 2	23(VAL) 25(SER) 23(GLY) 25(SER)	3.4 2.1 4.5 2.1
16 17 18	5	GLY GLN arg	gly thr val	26 26 26 25	2 1 4 3 3	25(PRO) 26(GLY) 23(GLN) 23(SER)	2.1 1. 4.5 3.4 4.2
23			leu thr CYS	25 19 18 18	1321	25(THR) 17(ILE) 17(SER) 18(CYS)	1. 3.4 2.1 1.
25 26 27 27	A		SER SER THR	15 15 15	4 255	9(18H) 14(GLY) 10(THR) 7(SER)	6.7 2.1 7.5 11.
c <u>27</u>	ç		ALA VAL	15 15 15	4	12(SER) 11(ASP) 14(VAL)	
29 30 31 32			ècn	14 14 14		6(GLY)	7.5 12. 9.3 12. 14.
34				13 13 14	21	11(VAL) 12(SER)	14. 3.5 <u>2.2</u> 1.
38 38 39 40		(()	GLN GLN .YS	14 14 14		10(TYR) 14(GLN) : 13(GLN) 13(GLN) : 12(GLN) 10(HIS) 10(HIS)	2.8 1. : 2.2 2.2 : 3.5 7.
44		007 P	SLY SLN NLA 'RO	14 14 14		13(GLY) 11(LYS) 13(ALA)	7. 1. 2.2 5.1 2.2
46 47 48		Ļ	LA LEU LE	14 14 14		14(16)	1. 2.2 3.5 8.4 1.
50 51 52 53		S	ER	14 14 14 14 14		7(ASP) 11(VAL) 5(SER)	4.7 10. 5.1 14. 21.
56		s	ER	14 14 14	2 2 1	13(ARG) 13(PRO) 14(SER)	2.2 2.2 1. 2.2
59 60		T. P	HR BO	14 14 14 14	2	10(VAL) 7(_) 5(ASP) 14(ABG)	4.2
64 65		6	1 Y	15 15	1	14(PHE) 15(SER) 15(GLY) 15(SER)	1. 2.1 1. 1. 1.
68		G	EU LY	14		13(LYS) 13(SER) 12(GLY) 10(ASN) - 9(ASN)	3.5 2.2 2.3 5.6 ; 6.2
71 72 73			Û	14 14 14	3121	12(THR) 14(ALA) 13(SER) 14(LEU)	3.5 1. 2.2 1.
76 77		- LE SE GL	U A Y	14 14 14	2	13(ILE) 14(SER) 14(GLY)	1. 2.2 1. 1. 2.2
79 80 81 82		GL PP	N	14 14 14	3	12(GLN) 10(ALA) 11(GLU) 13(ASP)	3.5 4.2 3.8 2.2 1.
84 85 86 87		AL GL TY TY	AUAA	14 14 14 14	3 4		$3.8^{1}_{:}5.6^{1}_{2,3}$
89 90 91 92				14 14 14 14	4227	8(SER) 13(SER) 12(TYR) 5(ALA)	1. 7. 2.2 2.3 20.
94 95 95A 958		GL AL	Y	14 13 11 2	5:6 7 3 2	5(SER) 5(+) 5(+) 5(+) 1(+)	8. 14. : 17. 30.
95D 95E 95F 96 97	VAL			13 16	8 3	5(VAL) 10(VAL)	21.
98 99 100 101	PHE GLY SEF			16 18 18 18	1 1 4 1	16(PHE) 18(GLY) 10(GLY) 18(GLY)	4.8 1. 1. 7.2 1.
103 104 105 106	1 15			18 15 15 13	5 2 3 1	13(LYS) 9(LEU) 13(THR) 13(VAL)	1. 6.9 3.3 3.5 1,
106A 107 108 109				13 13 10 10	1 3 1:2	13(LEU) B(GLY)	.4.9 1. : 2.2
	FR 1 1111111111111111111111111111111111	0 1 2 3 4 5 6 7 8 9 10 12 11 12 12 3 4 5 6 7 8 9 11 12 12 23 223 223 223 225 227 227 223 22 223 22 223 22 223 22 223 22 224 22 226 227 227 CDD 228 29 331 33 34 44 43 44 43 44 49 51 558 56 577 76 777 74 756 776 99	25 WAL 0	25 26 WAL 26 CL 0 1 PCA ginr 3 vell val 4 LEU val 5 THR THR 6 GOD ginr 7 SER Inr 7 B Proo Bit 6 GOLV Reg R 10 Inr Inr 113 GELY Val R 114 thr SER SER 115 PRO PRO GLY 122 Ithr SER SER 221 Titre GLY ALA 222 ALA SER SER 221 CYS SER SER 222 TYA THR SER 221 CYS SER SER 222 ALA SER SER 227 THR <t< td=""><td>25 26 # OF 1 PCA gin 26 2 SER Ihr 26 3 val val 26 4 LEU val 26 5 THR THR 26 6 OCRO GLN 26 9 SER SER 26 10 26 11 Jag leu 26 11 Jag R 11 12 SER htr 26 14 thr SER 26 15 GLV GLV 21 16 GLV GLV 26 22 thr 18 27 21 THR 14 25 227 THR 15 277 15 277 15 278 SER 14 30</td><td>$\begin{array}{c c c c c c c c c c c c c c c c c c c$</td><td>WAL As SEGUENCES ANNO O CMOST COMMON 1 PCA III 26 32 24(PCA) 2 SER HIN 26 2 25(SER) 3 YER NINE 28 22 25(LEA) 4 THA THA 78 12 26(CLN) 25(CLN) 5 THA THA 78 12 25(CLN) 12 25(CLN) 6 GLN GLN 26 3 22(VAE) 12 7 113 GER 120 26 2 22(VAE) 13 GER 120 26 2 22(VAE) 14 thr 266 3 22(VAE) 23 22(VAE) 14 15 1 26(CLN) 12 22(SER) 12 15 GLN thr 25 3 22(SER) 12 14 15 1</td></t<>	25 26 # OF 1 PCA gin 26 2 SER Ihr 26 3 val val 26 4 LEU val 26 5 THR THR 26 6 OCRO GLN 26 9 SER SER 26 10 26 11 Jag leu 26 11 Jag R 11 12 SER htr 26 14 thr SER 26 15 GLV GLV 21 16 GLV GLV 26 22 thr 18 27 21 THR 14 25 227 THR 15 277 15 277 15 278 SER 14 30	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	WAL As SEGUENCES ANNO O CMOST COMMON 1 PCA III 26 32 24(PCA) 2 SER HIN 26 2 25(SER) 3 YER NINE 28 22 25(LEA) 4 THA THA 78 12 26(CLN) 25(CLN) 5 THA THA 78 12 25(CLN) 12 25(CLN) 6 GLN GLN 26 3 22(VAE) 12 7 113 GER 120 26 2 22(VAE) 13 GER 120 26 2 22(VAE) 14 thr 266 3 22(VAE) 23 22(VAE) 14 15 1 26(CLN) 12 22(SER) 12 15 GLN thr 25 3 22(SER) 12 14 15 1



REFERENCE: HUMAN LAMBDA LIGHT CHAINS SUBGROUP II

1) NIG-84: TONOIKE,H.,KAMETANI,F.,HOSHI,A.,SHINODA,T. & ISOBE,T. (1985) FEBS LETTERS,185.139-141.

 MES: ZHU,D., KIM,H.S. & DEUTSCH.H.F. (1983) MOL.IMMUNOL. 20.1107-1116.
 MES: ZHU,D., KIM,H.S. & DEUTSCH.H.F. (1983) MOL.IMMUNOL. 20.1107-1116.
 WH: KIEFER,C.R., PATTON,H.M., JR., MCQUIRE,B.S., JR. & GARVER,F.A. (1980) JIMMUNOL. 124,301-306. (CHECKED BY AUTHOR 02/11/80) 3) WH: KIEFER,C.R.PATTON,H.M.JR.MCQUIRE,B.S.JR. & GARVER,F.A. (1980) J.IMMUNOL.,124,301-306. (CHECKED BY AUTHOR 02/1
4) NEI: GARVER,F.A. & HLSCHMANN,N. (1971) FEBS LETTERS,16,128-132; (1972) EURJ.BIOCHEM.,26,10-32; (CHECKED BY AUTHOR)
5) KAR: FETTJ,W. & DEUTSCH,H.F. (1976) IMMUNOCHEM.,13,149-155. (CHECKED BY AUTHOR)
6) RIM: FETTJ,W. & DEUTSCH,H.F. (1976) IMMUNOCHEM.,13,149-155. (CHECKED BY AUTHOR)
7) SLA: FETTJ,W. & DEUTSCH,H.F. (1976) IMMUNOCHEM., 13,149-155. (CHECKED BY AUTHOR)
7) SLA: FETTJ,W. & DEUTSCH,H.F. (1976) IMMUNOCHEM., 13,149-155. (CHECKED BY AUTHOR)
8) SLA: FETTJ,W. & DEUTSCH,H.F. (1976) IMMUNOCHEM., 13,149-155. (CHECKED BY AUTHOR)

1/ DEX: FEITURY & DEUTSCH.H.F. (1976) IMMUNUCHEM.,13,149-155. (CHECKED BT AUTHUH) 8) ES492: TVETERAAS.T.SLETTEN.K. & WESTERMARK,P. (1985) BIOCHEM.J.,232.183.190. 9) WEIR: FETTURY & DEUTSCH.H.F. (1976) IMMUNOCHEM.,13,149-155. (CHECKED BY AUTHOR); JABUSCH.J.R. & DEUTSCH.H.F. (1982) MOL.IMMUNOL.. 9) WEIR: FETTURY & DEUTSCH.H.F. (1976) IMMUNOCHEM.,13,149-155. (CHECKED BY AUTHOR); JABUSCH.J.R. & DEUTSCH.H.F. (1982) MOL.IMMUNOL..

9) WEIR: FETTJW, & DEUTSCH.H.F. (1976) IMMUNOCHEM..13.149-155. (CHECKED BY AUTHOR): JABUSCH.J.R. & DEUTSCH.H.F. (1982) MOL.IMMUNOL... 19:901-906.
10) TOG: NABESHIMA.Y. & IKENAKA.T. (1979) MOL.IMMUNOL..16.439-444. (CHECKED BY AUTHOR 10/10/79)
11) SM: GARVER.F.A., CHANG.L...MENDICINO.J..ISOBE.T. & OSSERMAN.E.F. (1975) PROC.NAT.ACAD.SCI.USA.72.4559-4563. (CHECKED BY AUTHOR 05/31/83)
12) HS68: HOOD.L. & EIN.D. (1968) NATURE.220.764-767.
13) MS77: HOOD.L. & EIN.D. (1968) NATURE.220.764-767.
14) TRO: SCHOLZ.R. & HILSCHMANN.N. (1975) Z.PHYSIOL.CHEM..356.1333-1335.
15) BOH: KOHLER.H..RUDOFSKY.S. & KLUSKENSL. (1975) J.IMMUNOL.OGY.114.415-421. (CHECKED BY AUTHOR)
16) NIG-58: TAKAYASU.T..TAKAHASHIN.SHINODAT..OKUYAMAT. & TOMIOKA.H. (1990) J.BIOCHEM..69.421-436. (CHECKED BY AUTHOR 05/24/84)
17) VIL: PONSTINGL.H. & HILSCHMANN.N. (1959) Z.PHYSIOL.CHEM..350,1140-1152; (1971) Z.PHYSIOL.CHEM..352.859-877. (CHECKED BY AUTHOR)
18) HOJS: STANTON.T.SLEDGE.C. CAPRA.J.D., WOODS.R. CLEM.W. & HOOD.L. (1974) J.IMMUNOL...112.633-640.
19) HBJ8: HOOD.L. & EIN.D. (1968) NATURE.220.764-787.
20) HS70: HOOOL.& EIN.D. (1969) NATURE.220.764-787.
21) HS70: CHEN.B.L.CHU.Y.H..JUMPINEY.R.L. & POLJAK.R.J. (1978) BIOCHIM.BIOPHYS.ACTA.537.9-21. (CHECKED BY AUTHOR 07/16/79)
21) WIN: CHEN.B.L.CHU.Y.Y.H.JUMPINEY.R.L. & POLJAK.R.J. (1978) BIOCHIM.BIOPHYS.ACTA.537.9-21. (CHECKED BY AUTHOR 07/16/79)
23) PRE: FETT.J.W. & DEUTSCH.H.F. (1976) IMMUNOCHEM..13.149-155. (CHECKED BY AUTHOR)

(CHECKED BY AUTHOR 07/28/79) 23) PRE: FETT.J.W. & DEUTSCH.H.F. (1976) IMMUNOCHEM..13.149-155. (CHECKED BY AUTHOR) 24) HS86: HOOD,L. & EIN.D. (1968) NATURE.220.764-767. 25) WAL: FETT.J.W. & DEUTSCH.H.F. (1978) IMMUNOCHEM..13.149-155. (CHECKED BY AUTHOR) 26) 4A'CL: ANDERSON,M.L.M.,SZAJNERT,M.F.,KAPLAN.J.C.,MCCOLL,L. & YOUNG.B.D. (1984) NUC.ACIOS RES..12.6647-6661. (CHECKED BY AUTHOR) 05/16/65)

NOTES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP II

IDENTICAL SETS OF FRAMEWORK SEGMENTS:

SET 1: NIG-84[1],MES[2],WH[3],NE][4],KAR[5],RIM[6],SLA[7], (7 IDENTICAL) SET 2: TRO[14],BOH(15], (2 IDENTICAL) FR1:

SET 1: WH(3).BOH(15).NIG-58(16).BUR(22). (4 IDENTICAL) FB2:

FRO

 SET 1: WHI3I. (IDENTICAL TO 1 HUMAN V-LAMBDA-I: NEWMI1]: AND 1 HUMAN V-LAMBDA-V: BO[1].)
 SET 2: MESI2I.ES492[8].TRO]14[.VIL[17].WIN[21]. [5 IDENTICAL HUMAN V-LAMBDA-II: ALSO 4 HUMAN V-LAMBDA-II: BL2 'CL[8].RHE[10].
 OKA112].NIG-51[13]: 4 HUMAN V-LAMBDA-III: HILI1.CAP[4].BAU112].DEL[14]: 1 HUMAN V-LAMBDA-II: SH11; 3 HUMAN OKA112].NIG-51[13]: 4 HUMAN V-LAMBDA-III: HILI1.CAP[4].BAU12].DEL[14]: 1 HUMAN V-LAMBDA-II: SH11; 3 HUMAN OKA112].NIG-51[13]: 4 HUMAN V-LAMBDA-III: HILI1.CAP[4].BAU12].DEL[14]: 1 HUMAN V-LAMBDA-II: SH11; 3 HUMAN OKA112].NIG-51[13]: 0 Y-LAMBDA-VI: SUT[2].TH0[4].LBV/CL[5]: AND 24 MOUSE V-LAMBDA-III: NPC10[4].S9[2].XS104[3].HOPC114[.J598][5].H2061[6].
 W3159[7].Y5431[8].Y5485[9].Y5830[10].Y5669[11].MOPC511[L]][2].S178[13].Y5441[14].Y5608[15].S1761[0].H202[17].
 RPC2018[.G. 302LAMBDA'CL]91543CU[2].S2H5CU[39].S1F12CL[149].IG 25LAMBDA'CL[41].)
 SET 3: NIG-84[1]. (IDENTICAL TO 1 HUMAN V-LAMBDA-III: GRI7].) FR4:

IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS: SET 1: MES(2), VIL 171. (2 IDENTICAL HUMAN V-LAMBDA-II; ALSO 1 HUMAN V-LAMBDA-V: MCG(3).) CDR1:

CDR2: 1: NIG-84[1], TOG[10]. (2 IDENTICAL) SET

CDB3:

IDENTICAL SETS OF J-MINIGENES:

SET 1: MES(2).TRO(114), (2 IDENTICAL HUMAN V-LAMBDA-II; ALSO 1 HUMAN V-LAMBDA-II: BAU(12).) SET 2: ES492(8).VIL(17), (2 IDENTICAL HUMAN V-LAMBDA-II; ALSO 1 HUMAN V-LAMBDA-III; DEL(14).)

SPECIFIC NOTES:


A

11) SM: IT HAS O LINKED CARBOHYDRATE ATTACHED TO SER AT POSITION 22 AND N-LINKED CARBOHYDRATE ATTACHED TO ASX AT POSITION 25.

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

T POSITION	RESIDUES
47	(ILE,MET)
53	(LYS,ASN)
59	(PRO SER)
95	(SER.ASN)
95A	(THR.SER)
95B	(LEU,ARG)

,	IUMA	N LAMBDA LI INVARIAN RESIDUES	т т	AINS 5	30868 3 PS	CAP	5		7.ª GAR	. в сн	9 X (PET	10 KERN	11 TA	12 8AU	13 AMYLOID	14 DEL	15 LYN	16 NIG	17 AMYLOID	18 MOT	19 WIG	20 WHI	21 DU	22 LON
-		0 1 2 TYR(.96) 3 4 LEU(.96) 5 6	SER TYR GLU LEU THR GLN	SEA TYA GLU LEU THR GLN	SER TYA GLU LEU THA GLN	SER TYR GLU LEU THR GLN PRO	SER TYR GLU LEU THR GLN PRO	LEU	SEA TYA GLU LEU	LEU	TYR BSP LEU	TYR ala LEU	SER TYR ala LEU THR	TYR oly LEU THR	758 TYR ASP LEU THR	TYR Val LEU Ser GLN	TYR GLU LEU THR GLN	-68 TYR asp LEU THR GLN	808 TYR 85p LEU THR GLN	Phe TYR GLU LEU THR	SER phe gly val ser	TYR val LEU THR GLX	TYR GLX LEU THR	TYR Ser LEU THR
	1	7 8 PRO 9 SER	PRO PRO SER	PRO PRO SER	PRO PRO SER	PRO PRO SER	SER	PRO	Iys GLN PRO PRO SER	THR GLN PRO SER	SER	THR GLN PRO PRO SER	THR GLN PRO PRO SER	THR GLN PRO PRO SER	GLN PRO PRO SER	GLN PRO PRO SER	GLN PRO PRO SER	GLN ala PRO SER	GLN PRO SER	GLN PRO PRO SER	Ser GLN PRO PRO SER	GLX ala PRO SER	GLX PRO PRO SER	GLN PRO PRO SER
F F 1		1 SER 2 SER 3 VAL(.96) 4 PRO(.95)	VAL SEA VAL SER PRO	VAL SEA VAL SER PRO	VAL SER VAL SER PRO	VAL SER VAL SER PRO	VAL SER VAL SER PRO GLY	VAL SER VAL SER	VAL SER VAL SER PRO	VAL SER VAL SER PRO	PRO	PBO	VAL SER VAL SER PRO	leu SER VAL SER PRO	Met SER VAL SER PRO	VAL SER VAL ela PRO	VAL SER VAL phe PRO	Jeu SER VAL SEA PRO GLY	VAL SER VAL SER PRO GLY	VAL SER leu ala ala	VAL SER VAL SER PRO	leu SER VAL ala	VAL SEA VAL SER PRO	VAL SER VAL SER PRO
	18	7 3 9 ALA(.95)	GLY GLN THR ALA	GLY GLN THR ALA	GLÝ GLN THR ALA	GLY GLN THR ALA	GLY GLN THR ALA	GLY GLN THR ALA	GLY GLN THR ALA	GLY GLN THR ALA	GLY GLN THR ALA	GLY GLN THA ALA	GLÝ GLN THR ALA	GLY GLN THR ALA	GLY GLX THR ALA	GLY GLN THA ALA	GLY GLN pro gly	GLY GLN THR ALA	GLY THR ALA	ala GLY GLN THR ALA	GLY GLN THR ALA	PRO GLY GLX THR ALA	PRO GLY GLX	
	2222		ARG ILE THR CYS	ARG ILE THR CYS	ARG ILE THR CYS	ARG ILE THR CYS	ILE THR CYS	ser ILE THR CYS	ARG ILE THR CYS	ARG	Ser ILE THR CYS	Val ILE THA CYS		LE THR CYS	Ser ILE THR	ARG ILE THR CYS	ILE THR CYS	tyr ILE THR CYS	ILE THR	met ILE THR CYS	Ser ILE THR CYS	ARG ILE THA CYS		
	24 25 26 27 27 27		SER ALA ASN ALA	SER GLY ASP ALA	SER GLY ASP ALA	SER GLY ASP ALA	SEA GLY ASP	SER GLY ASX	SER GLY ASP VAL		SER GLY ASP LYS	SEA GLY ASP ASN		SER GLY ASP LYS	SEA GLY GLX ASX	GLY GLY ASP GLY	SER GLY ASP ALA	SER GLY ASP ASN		GLU GLY ASN ASP		GLX ASX ASX		
CDR	27 27 27	C D													' '					 -::				
î	27 28 29 30		LEU PRO ASN GLN	LEU	LEU	LEU	LEU	LEU	LEU		LEU GLY ASP	LEU GLU LYS THR	,	LEU GLY GLU	LEU	ILE GLY GLY	LEU	LEU GLY		ILE GLY GLU		ILE GLX GLX		
	31 32 33 34		GLN TYR ALA TYR	PRO ASP LEU TYR VAL	THR ASN LYS TYR ALA	ALA GLU TYR ALA	GLX GLX	GLX GLX	LYS LYS TYR ALA		LYS ASP VAL	VAL		GLN TYR VAL		SER VAL	ASP LYS TYR VAL	ASN GLU PHE VAL		ARG SER VAL		GLX	TYR VAL	
_	35 36 37 38	TAP	TRP TYR GLN GLN	TRP TYR GLN GLN	TYR TRP	TYR TRP TYR GLN GLN			TYR TRP TYR GLN GLU		CYS TRP TYR GLN	SEA TRP PHE GLN		CYS TRP TYR GLN	·	HIS TRP TYR GLN	TRP TYR GLX HIS	SER TAP TYR GLN GLN		HIS TAP TYR GLN		•	CYS TRP	
f	39 40 41 42		LYS PAO GLY	LYS		LYS PRO GLY			ARG SER GLY		GLN ARG PRO GLY	GLN ARG PRO GLY GLN		GLN LYS PRO GLY		GLN LYS PRO GLY	1.15	GLN PRO GLY GLX SER		GLN LYS PRO GLY				
2	43 44 45	PRO	ARG ALA PRO VAL			GLN ALA PRO VAL			GLN ALA PRO VAL		GLN SER PRO VAL LEU	PRO		GLN SER PRO VAL		GLN ALA PRO	PRO LEU	PRO		GLN ALA PRO				
	46 47 48 49	VAL	MET VAL ILE TYR			MET VAL ILE TYR			LEU VAL VAL TYR		LEU VAL ILE TYR	LEU VAL ILE TYR		LEU VAL ILE TYR		VAL LEU VAL VAL HIS	VAL	ALA LEU VAL ILE TYR		PRO VAL ILE TYR				
CDR	50 51 52 53		LYS ASP THR GLN			GLU THR ASN LYS			GLU ASP SER GLY	ASP THR GLY	GLN ASP ASN GLN	HIS THR SER GLU		HIS ASP SER LYS		GLU ASP ASN	GLX	ASX THR SER LYS		ASP ASP ALA ASP				
Fi 2 	54 55 56 57	ARG	ARG PRO SER GLY			ARG PRO SER GLY			ARG PRO SER GLU	ARG PRO SER THR	ARG SER SER GLY	ARG PRO SER GLU		ARG PRO SER GLY		ARG PRO ALA	ARG PRO	ARG PRO SER		ARG PRO SER	•			
	58 59 60 61	PRO	ILE PRO GLN ARG			ILE PRO GLU ARG			ILE PRO GLU	ILE PRO GLU	ILE PRO GLU ARG	ILE PRO GLU ARG		ILE PRO GLU ARG		ILE PRO GLU		GLY ILE PRO GLU ARG		GLY VAL PRO ALA				
	62 63 64 65	SER	PHE SER SER			PHE SER GLY SER				AAG PHE SEA GLY SER	GLY SER	GLY SER SER SER		PHE SER GLY SER		PHE SEA GLY SEA	PHE	SER GLY		ARG PHE SER GLY				
	66 67 68 69	SER GLY	SER THR SER GLY			THR SER GLY THR			SER SER GLY	THR SER GLY	ASN SER GLY	SER GLY		ASN SER GLY		ASN SER GLY		LYS SER GLY		TYR ASN SER GLY				
F R 3	70 71 72 73	LEU	THR THR VAL THR LEU			THR VAL THA LEU			LYS	THR THR ALA THR LEU	ASN THR ALA THR LEU	ALA THR ALA THR LEU		THR THR ALA THR LEU		ASN THA ALA ALA LEU		ASN THR ALA THR LEU		ASN SEA ALA				
5	74 75 76 77	THR ILE	THR ILE SER GLY VAL		•	THR ILE SER GLY			THR ILE SER		THR ILE SER GLY	THR ILE SER GLY		THR ILE SER		THA JLE SER	-	IHR ILE SER		LEU THR ILE ASN				
	78 79 80 81		VĂĹ GLN ALA GLU			GLN GLN GLU			ALA GLN	VAL	THA GLN ALA MET	ALA GLN SER	(GLY THR GLN ALA MET		ARG VAL GLU ALA GLY	ç	GLY THR GLU GEA		ARG VAL GLU ALA				
	82 83 84 85	ALA	ASP GLU ALA ASP TYR		i	ASP GLU		•	ASP GLU	GLX	GLU	VAL ASP GLU ALA	ć	ASP GLU		ASP GLU	ć	NET ASX SLU ALA	ć	GLY ASP GLU ALA				
	86 87 88 89	TYR CYS	TYR TYR CYS GLN		(ALA ASP TYR TYR CYS SER			CYS	TYR CYS	ALA ASP TYR TYR CYS	ALA ASP TYR PHE CYS		ALA ASP TYR TYR CYS		ALA ASP TYR TYR CYS	(ALA ASP TYR PHE CYS				
	90 91 92 93		ALA TRP ASP ASN		5	SER ALA SER		,	SEA THR ASP ILE ASN	SER ALA	GLN ALA TRP ASP SER	GLN THR TAP ASP THR	ń	SLN RP SP SER		GLU VAL TAP ASP	ہر بر	LA HP SX		SER FRP ASP				
CDR3	94 95 95A 95B		SER ALA		500	SER SLN SLY				ARG	MET SER	ILE THR	ר ר	YA		ASP ARG 7HR ALA	ן מ מ	1LX LE AG SP	(5 1	SLY SEA TYA	•			
3	95C 95D 95E	•														HIS				SLU 				
	95F 96 97 98 99	PHE GLY	SER ILE PHE GLY		}	AET AL		L F	280 .EU 24E		VAL VAL PHE	ALA ILE PHE	P				۷ ۷ ۹	AL AL HE	\\ F	AL AL V	HE			
FR	100 101 102 103	GLY THR	GLY GLY THA LYS		C T			· 1	3LY (3LY ("HR 1	IHR I	GLY GLY GLY THR	GLY GLY GLY THR	0 G T		(GLY GLY GLY THR	G T	LY LY LY	G T G	ILY G HR A ILY G HR TI	LY LA LY HR			
4	103 104 105 106 106A	VAL LEU			L T V	YS EU HR AL EU		Lor Vor	Λ. ١	AL N	ARG LEU THR VAL LEU	LYS LEU THR VAL LEU	۲ ۷	YS EU HA AL EU	L	LYS EU THR VAL	L T V	YS EU HR AL	· ¥	'AL HR 'AL	ня			
****	107 108 109	GLN PRO	GLY GLN PRO		G	LY LN RO			LY ILN RO		SER	SER GLN PRO	G	LY LN RO		.EU GLY	L.	EU	G	EU LY LN RO				_



•

ANTIBODY SPECIFICITIES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP III 7) GAR: ANTI-RIBOFLAVIN

- REFERENCE: HUMAN LAMBDA LIGHT CHAINS SUBGROUP III

1) HIL: LOPEZ DE CASTRO, J.A., CHIU, Y.Y.H. & POLJAK, R.J. (1978) BIOCHEMISTRY, 17.1718-1723. (CHECKED BY AUTHOR 07/16/79)

2) YO: TISCHENDORF,F.W.,TISCHENDORF,M.M. & WITTMANN-LIEBOLD,B. (1976) Z.NATURFORSCH.31C,759-760. 3) PS: KOCHWA,S.,TERRY,W.D.,CAPRA,J.D. & YANG,N.Y. (1971) ANN.N.Y.ACAD.SCI.,190.49-70. (CHECKED BY AUTHOR)

(a) PS: ROCHWAS, TEHHY, W.D., CAPRAJ, D. & TARGN, T. (1971) ANKINT, ACAD, SOL, 1904, 10. (CRECKED BT AUTHOR)
(b) CAP: ZHUD, XIM, R.S. & DEUTSCH, F.F. (1983) MOLIMMUNOL, 20, 1107-1116.
(c) LOY A: WOLFENSTEIN-TODEL, C., FRANKLIN, E.C. & RUDDERS, R.A. (1974) JIMMUNOL, 112,871-876. (CHECKED BY AUTHOR)
(c) LOY G: WOLFENSTEIN-TODEL, C., FRANKLIN, E.C. & RUDDERS, R.A. (1974) JIMMUNOL, 112,871-876. (CHECKED BY AUTHOR)
(c) LOY G: WOLFENSTEIN-TODEL, C., FRANKLIN, E.C. & RUDDERS, R.A. (1974) JIMMUNOL, 112,871-876. (CHECKED BY AUTHOR)
(c) GAR, M.G., GUIRE, B.G., JR, OSSERMAN, E.F. & GARVER, F.A. (1983) JIMMUNOL, 113, 1871-1875. (CHECKED BY AUTHOR)
(c) CHECKED BY AUTHOR 02/20/84)
(c) CHECKED, J., MARK, WATANABE, S., HARA, H. & KITAGAWA, M. (1972) JIMMUNOCHEM, 9, 207-210.

 CH: OKADA,T.,NO2O,Y.,IITANIK,WATARABES.,BARAH. & KITAGAVAR. (1972) IMMONOCIENT.,3207-800.
 X(PET): MILSTEINC,CLEGGJ,B. & JARVISJ,M. (1968) BIOCHEM.J.(110.631-652) (CHECKED BY AUTHOR)
 KERN: PONSTINGL,H.,HESS,M. & HILSCHMANN,N. (1968) Z.PHYSIOL,CHEM.,349,867-871; (1971) Z.PHYSIOL,CHEM.,352,247-266. (CHECKED BY AUTHOR)
 KERN: PONSTINGL,H.,HESS,M. & HILSCHMANN,N. (1968) Z.PHYSIOL,CHEM.,349,867-871; (1971) Z.PHYSIOL,CHEM.,352,247-266. (CHECKED BY AUTHOR)
 KERN: PONSTINGL,H.,HESS,M. & HILSCHMANN,N. (1970) Z.PHYSIOL,CHEM.,351,763-767; BACZKO,K.,BRAUN,D.G. & HILSCHMANN,N. (1974) Z.PHYSIOL,CHEM.,351,763-767; BACZKO,K.,BRAUN,D.G. & HILSCHMANN,N. (1974)
 AMYLOID 758: NATVIGJ,B.,WESTERMARK,P.,SLEYTEN,K.,HUSBY,G. & MICHAELSON,T. (1981) SCAND,J.IMMUNOL.,14,89-94. (CHECKED BY AUTHOR) 14) DEL: EULITZ,M. (1974) EUR.J.BIOCHEM. 50,49-69. (CHECKED BY AUTHOR 10/18/77)

I. LULL, I. (L.), I. (L.),

17) AMYLOID 808: WESTERMARK,P.NATVIG,J.B.,ANDERS,R.F.,SLETTEN,K. & HUSBY,G. (1976) SCAND.J.IMMUNOL.,5,31-36. (CHECKED BY AUTHOR)
18) MOT: KOJIMA,M.,ODANI,S. & IKENAKA,T. (1980) MOL.IMMUNOL.,17,1407-1414.
19) WIG: FETT,J.W. & DEUTSCH,H.F. (1975) IMMUNOCHEM.,13,149-155. (CHECKED BY AUTHOR)
20) WHI: WANG,A.C.,WELLS,J.V.,FUDENBERG,H.H. & GERGELY,J. (1974) IMMUNOCHEM.,11,341-345. (CHECKED BY AUTHOR)
21) DU: BUCHWALD,B.M. (1971) CANJ,BIOCHEM.,49,500-902. (CHECKED BY AUTHOR)
22) LON: JOHNSTON,S.L.,ABRAHAM,G.N. & WELCH,E.H. (1975) BIOCHEM.BIOPHYS.RES.COMMUN.66,842-847. (CHECKED BY AUTHOR)
23) SG: TISCHENDORF,F.W.,TISCHENDORF,M.M. & WITTMANN-LIEBOLDB. (1976) ZNATURFORSCH.31C.758-760.
24) GIM: HESS,M.,HILSCHMANN,A,RIVET,L.,RIVET,C. & ROPARTZ.C. (1971) NATURE NEW BIOLOGY,234.58-61. (CHECKED BY AUTHOR)
25) 111: LANGER,B.,STEINMETZ-KAYNEM. & HILSCHMANN,N. (1968) Z.PHYSIOL.CHEM.,349.945-951. (CHECKED BY AUTHOR)
26) T19: HESS,M.,HILSCHMANN,N.R,RIVET,L.,RIVET,C. & ROPARTZ.C. (1971) NATURE NEW BIOLOGY,234.58-61. (CHECKED BY AUTHOR)
27) VIN: PINKJ,R.L. & MILSTEN,C. (1967) PROC. 571 HEBS N.PHISOL.CHEM. 2010,254.58-61. (CHECKED BY AUTHOR)
28) T19: LANGER,B.,STEINMETZ-KAYNEM. & HILSCHMANN,N. (1968) Z.PHYSIOL.CHEM. 2010,254.58-61. (CHECKED BY AUTHOR)
29) T19: LINS,CHMANN,N.RIVET,L.,RIVET,C. & ROPARTZ.C. (1971) NATURE NEW BIOLOGY,234.58-61. (CHECKED BY AUTHOR)
20) T19: NIKJ,R.L. & MILSTEN,C. (1967) PROC. 571 HEBS SCHARMAN, (1968) Z.PHYSIOL.CHEM. 2010,254.58-61. (CHECKED BY AUTHOR)
20) T19: LANGER,B., TEINMETZ-KAYNEM, & HILSCHMANN, (1968) Z.PHYSIOL.CHEM. 2010,254.58-61. (CHECKED BY AUTHOR)
20) T19: NIKJ,R.L. & MILSTEN,C. (1967) PROC. 571 HEBS SCHARMAN, (1968) Z.PHYSIOL.CHEM. 2010,254.58-61. (CHECKED BY AUTHOR)

27) VIN: PINK, J.R.L. & MILSTEIN, C. (1989) PROC. 5TH FEBS SYMP., 15, 177-182. (CHECKED BY AUTHOR) 28) MIL: LANGER, B., STEINMETZ-KAYNE, M. & HILSCHMANN, N. (1968) Z.PHYSIOL.CHEM., 349, 945-951. (CHECKED BY AUTHOR)

HUMAN LAMBDA LIGHT CHAINS SUBGROUP III NOTES:

IDENTICAL SETS OF FRAMEWORK SEGMENTS:

SET 1: HIL[1], YO[2], PS[3], CAP[4], (4 IDENTICAL) SET 2: LOY A[5], LOY G[6]. (2 IDENTICAL) FR1:

FR2:

FR3: FB4:

SH[1]: 3 HUMAN PC1[4],J698[5],H2061[6], 20[17],

SET 1: HIL[1].CAP[4].BAU[12].DEL[14]. [4 IDENTICAL HUMAN V-LAMBDA-III: ALSO 4 HUMAN V-LAMBDA-I: BL2 'CL[6].RHE[10].OKA[12]. NIG-51[19]: 5 HUMAN V-LAMBDA-II: MES[2].E5492[8].TRO[14].VL17].WIN2[1: 1 HUMAN V-LAMBDA-IV: SH[1: 3 HUMAN V-LAMBDA-VI: SU[2].THO[4].LBV CL[6]: AND 24 MOUSE V-LAMBDA'IV.CAP[1].2569[3].756[6].HZOC1[4].DS9[5].H2O51[1 V-LAMBDA-VI: SU[2].THO[4].LBV CL[6]: AND 24 MOUSE V-LAMBDA'IV.CAP[1].2569[3].756[6].HZOC1[4].DS9[5].H2O51[1 ND20].ND20[3].H2O31[3].HAMBDA'LE10].S05CL[21].S2H5'CL38].S2E9CL[39].S1F12'CL[40].IG 25LAMBDA'CL[41]. SET 2: GR17L IDENTICAL TO 1 HUMAN V-LAMPDA'IV. NGA40[1].

SET 2: GARI7]. (IDENTICAL TO SET 3: KERNI10]. (IDENTICAL TO L TO 1 HUMAN V-LAMBDA-II: NIG-84[1].) L TO 1 HUMAN V-LAMBDA-VI: NIG-48[10].)


IDENTICAL SETS OF J-MINIGENES:

SET 1: BAU(12), (IDENTICAL TO 2 HUMAN V-LAMBDA-II: MES(2), TRO(14).) SET 2: DEL[14], (IDENTICAL TO 2 HUMAN V-LAMBDA-II: ES492(8), VIL(17).)

SPECIFIC NOTES:

18) MOT: THERE ARE TWO RESIDUES IN FRONT OF POSITION 1; THEY ARE VAL AND THR.

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:



	SER SER SER SER SER SER SER SER	THAR ASRO ALL VALRA LUGUEN VALRA LA LUGUEN VALRA LA LUGUEN VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA VALRA V	THE SER USE AND A CONTRACT AND A CON	005 555 5555 565 4 4 4 4	1 21 2 2 2 2 2 2 2 2 1 2 2 2 1 2 2 2 2	4(SER) 4(GLU) 5(LEU) 5(GLN) 3(PRO) 3(PRO) 3(SER) 4(VAL) 3(ALA) 3(ALA) 3(GLN) 3(GLN) 3(GLN) 3(GLN) 3(GLN) 3(LEG) 2(SER) 4(CYS) 2(SER) 4(CYS) 2(SER) 4(CLY) 3(ASF)	1, 2,5 1, 3,3 2,5 3,3 1, 5, 3,3 1, 2,5 3,3 1, 2,5 7,5 10, 1, 2,5 2,5 10, 1, 2,5 2,5 10, 2,5 1,5 2,5 3,3 1, 2,5 2,5 3,3 1, 2,5 7,5 1, 2,5 2,5 3,3 1, 2,5 2,5 2,5 3,3 1, 2,5 2,5 2,5 3,3 1, 2,5 2,5 2,5 3,3 1, 2,5 2,5 3,3 1, 2,5 2,5 2,5 2,5 2,5 3,3 1, 2,5 2,5 2,5 2,5 2,5 2,5 2,5 2,5 2,5 2,5
	0 1 VAL 2 3 4 5 5 6 6 6 6 6 6 7 7 8 8 9 9 9 1 1 1 1 1 1 1 1 1 1 1 1 1	VAL VAL SER SERL ALA SERL ALA ALA GLUX THR GUN THA ALEU GGLY THR ARG ARE THR GUN ASPR 	VAL	5555555555 74 74 75 75 75 75 75 75 75 75 75 75 75 75 75	2 2 2 1 : 2 3 4 1 2 3 3	3(PAO) 5(GLY) 5(GLN) : 3(GLN) 4(THR) 2(+) 2(ARG) 5(LE) 4(THR) 4(THR) 4(CYS) 2(SER)	2:5 2:5 3.3 1. 1. : 0.3 2:5 7.5 10. 1. 2:5 1. 2.5 1. 0.
COR1 FR2	10 ILE 12 CYS 13 CYS 14 CYS 15 TAP 26 TYR 37 GLN 38 GLN	ARG ARG ILE ILE CYS GLN GLY ASP SER I I LEU ARG GLY ASP SER I I LEU ARG GLY ASP	ser vol all LEE LLEE LLE THAR THAR ATHEN SER SER LL GLY GLY GL ASP ASP LL LY I I I I I I I I I I I I I I I I I I I	4 Y 4 A 4 A 4	1 2 1 3	2(SER)	1. 2.5 1. 8.
	26 277A 277B 277C 277C 277C 277C 277C 277C 277C	ASP SET LEU ARY GLYR ASP		-	ž	2(+)	4.
FR2	28 29 30 31 32 33 34 35 TRP 35 TRP 35 TYR 37 GLN 38 GLN	LEU ARG GLY TYR ASP	LEU LEU IL			1(ASN) : 1(ASP)	
F R 2	35 TRP 36 TYR 37 GLN 38 GLN	ALA	GLY GLY GI ASP GLN AL ASN ALA TYA AS ALA T SER II	E 4 -Y 4 -A 4 -R 4 -X 3 -X 3 -X 3 -X 3	1 2 4 3 2 : 3 2 3	2(ALA) 1(+)	2.7 2.7 16. 6. 3. : 9. 3. 9. 1.
CDR2	39 LYS 40 42 42 43 43 44 45 46 47 48	ALA TAP TYR GLN LYS PRO GLY GLN ALA PRO LEU LEU VAL	JRP TRP TYR TYR GLN GLN LYS	3 3 2 2 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	(TPP) ((LN) 2(GLN) 2(GLN) 2(LYS) 1(PRO) 1(GLN) 1(ALA) 1(ALA) 1(LEU) 1(LEU) 1(LEU) 1(LEJ) 1(VAL) 1(VAL)	1. 1. 1. 1.
	49 50 51 52 53 54 55 56	TYR GLY ARG ASN ASN ARG PRO SER		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1	1(GLY) 1(ARG) 1(ASN) 1(ASN) 1(ASN) 1(ARG) 1(PRO) 1(SER) 1(GLY)	
	57 58 59 60 61 62 63 63 64 65 68	GLY ILE PASP ARBE SEA GLY SEA SEA		, ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1(LE) 1(PRO) 1(ASP) 1(APRO) 1(APHE) 1(SER) 1(SER) 1(SER) 1(SER) 1(SER) 1(SER)	
FR3	67 68 69 70 71 72 73	SER GLY HIS THR ALA SER LEU THR		1	1	1(HIS) 1(THA) 1(ALA) 1(SER) 1(LEU)	
	74 75 76 79 80 81 82 83 84 83 84 85 86 87	ILE TGLA GLA GLA GLA GLA GLA ASP TYR TYR				(THR) (THR) (THR) (THR) (THR) (TGLY) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (GLV) (TASP) (GLV) (TYR) (CYS)	
CD R 3	88 89 90 91 92 93 93 94 95	CYS ASN SER AROP SER SER SELY LYS HIS		1 1 3 3 1 1 1 1 1 1 1 1		1 ((45N) 1 ((5ER) 1 (ASP) 1 (ASP) 1 (ASP) 1 (SER) 1 (SER) 1 (SER) 1 (SER) 1 (SER) 1 (SER) 1 (HIS)	
F F 4	95F 96 97 98 90 100 101 101 101 101 F 102 Fi 103	VAL LEU PHE GLY GLY GLY THA LYS LEU THA VAL LEU GLY	•		 	1 (YAL) 1 (LEU) 1 (JELY) 1 (GLY) 1 (GLY) 1 (JELY) 1 (JELY)	

REFERENCE: HUMAN LAMBDA LIGHT CHAINS SUBGROUP IV

- NEY: NUMAR LAMBUA LIGHT CHARTS SUBJECT IN
 SH: TITANIK, WIKLER, SHINODA, T. & PUTNAM, F.W. (1970) J.BIOL.CHEM..245.2171-2176. (CHECKED BY AUTHOR 06/15/83)
 NEY: WANG, A.C., WELLS, J.V., FUDENBERG, H.H. & GERGELY, J. (1974) IMMUNOCHEM..11.341-345. (CHECKED BY AUTHOR)
 USH: TISCHENDORF, F.W., TISCHENDORF, M.M. & WITTMANN-LIEBOLD, B. (1976) Z.NATURFORSCH.31C.758-760.
 PFA: TISCHENDORF, F.W., TISCHENDORF, M.M. & WITTMANN-LIEBOLD, B. (1976) Z.NATURFORSCH.31C.758-760.
 FRA: WANG, A.C. & FUDENBERG, H.H. (1974) J.IMMUNOGENETICS.1.303-313. (CHECKED BY AUTHOR)

NOTES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP IV

IDENTICAL SETS OF FRAMEWORK SEGMENTS: FR1: SET 1: SH[1].NEV[2]. (2 IDENTICAL)

FR1: FR2:

- FR3: FR4:
- SET 1: SHI1]. [IDENTICAL TO 4 HUMAN V-LAMBDA-II: BL2 'CLI6I.RHE[10].OKA[12],NIG-51[19]: 5 HUMAN V-LAMBDA-II: MES[2].ES492[8]. TRO[14],VIL]17],VVIN[21]: 4 HUMAN V-LAMBDA-III: HIL]11,CAP[4].BAU[12],DEL]14]: 3 HUMAN V-LAMBDA-VI: SUT[2].THO[4], LBV CLI5]: AND 24 MOUSE V-LAMBDA: MOPCT04E[1].JS508[2].XS104[3].HOPC114],J6588[5].H206116[,W3159]7].Y5431[8],Y5485[9]. Y5830[10];Y5669[11],MOPC511[12]].S7184].H.Y5606[15].S71616],H2020[17],RPC20[18],IG 303LAMBDA CL[19]. S43[CL[21],S2H5'CL[38],S2E9'CL[39],S1F12'CL[40],IG 25LAMBDA 'CL[41].)

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

AT POSITION	RESIDUES
19 27	(VAL,ALA) (LYS.SER)
30 32	(ALA.GLY ASP.GLN) (TYR.ASP.ASN) (ILE.ALA.SER)
34	(ILE ALA.SER)

• •	N LAMBDA LIGI INVARIANT RESIDUES	HT CHAINS SU		# OF EQUENCES	# OF AMINO ACIDS	OCCURRENCES OF MOST COMMON AMINO ACID	VARIABILITY
	0 1 PCA 2 SEB	PCA PCA	PCA SER ALA LEU	3 3 3	1	3(PCA) 3(SER) 3(ALA)	1. 1. 1.
	3 ALA 4 LEU 5 THR	ALA ALA LEU LEU THR THR	LEU THR GLN	3	1	3(LEU) 3(THR) 3(GLN) 3(PBQ)	1. 1. 1. 1.
	5 THR 6 GLN 7 PRO 8 PRO 9 SER	THR THR GLN GLN PRO PRO PRO PRO SER SER	PRO PRO SER	3333	1	3(PRO) 3(PRO) 3(SER)	1. 1. 1.
	10 11 ALA 12 SER 13 GLY	ALA ALA SER SER GLY GLY	ALA SER GLY	3333		3(ALA) 3(SER) 3(GLY) 3(SER)	1.
	14 SER	SER SER	SER IOU GLY		1 2 1	3(SER) 2(PRO) 3(GLY) 3(GLN) 3(SER)	1. 3. 1.
	15 16 GLÝ 17 GLN 18 SER	PRO PRO GLY GLY GLN GLN SER SER VAL VAL		33333	1 1 1	3(VAL)	1. 1. 1.
	19 VAL 20 THR 21 ILE 22 SER 23 CYS	THR THR	THR	3 3 3	1 1	3(THR) 3(ILE) 3(SER) 3(CYS)	1. 1. 1.
	24 THR	THR THR	SEA CYS THR GLY	3	1 1	3(THR) 3(GLY) 3(THR)	1.
	25 GLY 26 THA 27 SER 27A	GLY GLY THA THA SER	SER	3 3 2	1 1	3(THA) 2(SER)	1. 1.
ç	27A 27B 27C 27C SER 27C SER 27E ASP 27F VAL	SER	SER	2	1	2(SER) 2(ASP) 2(VAL) 2(GLY)	
C D R 1	28 GLY	SER ASP VAL GLY	GLY	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	1	3(+)	1. 4.
	29 30 31	ASP ASN LYS TYR	TYA ASN TYA VAL	22220	2221	1(+) 1(+) 2(TYR) 2(VAL)	4. 4. 1. 1.
	32 TYR 33 VAL 34 SER	VAL SER TRP	SER	22	1	2(SEB)	<u>.1.</u> 1. 1.
	35 TAP 36 TYR 37 GLN 38 GLN	TYA GLN GLN	TYR GLN GLN	2222	1	2(TAP) 2(TYR) 2(GLN) 2(GLN) 2(GLN)	1.
- ·	39 HIS 40 41 GLY	HIS PRO	HIS ALA GLY LYS	****	1212	2(HIS) 1(+) 2(GLY) 1(+)	4. 1. 4.
F R 2	42 43 ALA	GLY ARG ALA PRO LYS	ALA PRO LYS		1	2(ALA) 2(PRO) 2(LYS)	1. 1. 1.
	44 PRO 45 LYS 46 47	LYS LEU VAL ILE	LYS VAL ILE ILE	****	221	1(+) 1(+) 2(ILE)	4. 4. 1.
	48 ILE 49 50 GLU	PHE	TYB	2	2	1(+) 2(GLU) 2(VAL) 1(+) 1(+)	4. 1. 1. 4.
CDR2	51 VAL 52 53	VAL SEA GLY	GLU VAL ASN LYS ARG	2222	1 2 2. 1	1(+) 1(+) 2(ARG) 2(PRO)	4. 1.
2	54 ARG 55 PRO 56 SFR	ARG PRO SER GLY	PRO SER GLY	222	1	2(PRO) 2(SER) 2(GLY) 2(VAL)	1. 1. 1.
	57 GLY 58 VAL 59 PRO	VAL PRO ASP	PRO	2 2 2	.]	2(PRO) 2(ASP) 2(ASP) 2(ARG) 2(PHE) 2(SER)	1, 1, 1,
	6C ASP 61 ARG 62 PHE 63 SER	ARG PHE SER	ASP ARG PHE SER	2222	1	2(PHE) 2(SER)	1. 1. 1.
	64 GLY 65 SER	GLY SEA LYS	GLY SEA LYS SER GLY	~~~~~		2(GLY) 2(SER) 2(LYS) 2(SER)	1.
	67 SER 68 69 ASN	SER ASP ASN THP	GLY ASN THR		2	1(+) 2(ASN) 2(THR)	. 4. 1. 1,
F	71 ALA 72 SER	ALA SER LEU	ALA SER LEU	NNNNN	1	2(ALA) 2(SER) 2(LEU)	1. 1. 1.
3	74 THR 75 VAL	THR VAL SER GLY	THR VAL SER GLY	NNNNN	1	2(THR) 2(VAL) 2(SER) 2(GLY)	· 1. 1. 1.
	78 LEU	GLY LEU ARG		22	1	. 2(LEU)	1. 4. 1.
	79 80 ALA 81 GLU 82 ASP	ALA GLU ASP GLU	ALA GLU ASP GLU	22	1	1(+) 2(ALA) 2(GLU) 2(ASP) 2(GLU)	1. 1. 1.
	83 GLU	GLU ALA ASP TYR TYR TYR	ALA ASP TYP TYP	22 22 22 22 22 22 22 22 22 22 22 22 22	1	2(ALA) 2(ASP) 2(TYR) 2(TYR) 2(TYR) 2(CYS)	1. 1. 1.
_	86 TYH 87 TYR 88 CYS		CYS	22	1	2(TYR) 2(CYS) 2(SER)	1.
	89 SER 90 SER 91 TYR 92	SEA SEA TYR VAL ASP	SEF SEF TYP GLU GL	2000	1 1 2 2 2	2(SER) 2(SER) 2(SER) 2(TYR) 1(+) 1(+)	4.
ç	93	ASN	GL SEF ASF	22222	22	1(+) 1(+) 2(ASN)	4. 4.
CDR 3	94 95 95A ASN 95B 95C	ASN	ASP 			· ·	•
	950 95E 95F 96 PHE		PHI	2 2			1.
· -	97 VAL 98 PHE 99 GLY	. VAL	VA PH GL TH GL	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		2(PHE) 2(GLY) 2(GLY) 22 1(+) 22 2(GLY) 22 2(GLY) 2(THR)	
F	100 101 GLY 102 THR	GLY GLY GLY THR	114	1 .2 C		2(GLY) 2(THR) 2(LYS)	1, 1, 1, 1, 1, 1, 4, 1, 1,
F F1 4	103 LYS	LYS LEU THR VAL	LY VA TH VA	L 2 R 2	:	1 2(LYS) 2 1(+) 1 2(THP) 1 2(VAL) 1 2(LEU)	4. 1. 1.
	106 VAL 106A LEU 107	ABG		Y 2	:	2 1(+)	4. 1. 1.
	108 GLN 109 PRO	GLN PRO	GL PR	0 2		1 2(GLN) 1 2(PRO)	1.

ANTIBODY SPECIFICITIES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP V

3) MCG: ANTI-EPSILON-DNP-LYS, EPSILON-DNP-AMINOCAPROATE, DNP-LEU, TRIACETIN, SODIUM MERTHIOLATE, METHADONE, 1,10-PHENANTHROLINE, CAFFEINE, THEOPHYLLINE, DI-DNP-LYS, DNP-TRP, DNP-PHE, DI-DNP-TYR, COLCHICINE, P-NITROANILINE, P-NITROPHENYLPHOSPHORYL CHOLINE, 5-ACETYLLINEACIL, MENADIONE, MEPERIDINE, TRIBUTYRIN, OMEGA-BROMOHEPTANOATE, O-CHLOROMERCURIPHENOL, P-CHLOROMERCURIPHENOL, PHENYLMERCURIC COMPOUNDS, METHYL-MERCURIC CHLORIDE.

75

REFERENCE: HUMAN LAMEDA LIGHT CHAINS SUBGROUP V

1) BO: WIKLER.M. & PUTNAM.F.W. (1970) J.BIOL.CHEM.245.4488-4507. (CHECKED BY AUTHOR 06/15/83) 2) HBJ2: HOOD.L..GRAY.W.R.SANDERS.B.G. & DREYER.W.J. (1967) COLD SPRING HARBOR SYMP. OUANTITATIVE BIOL.32.133-145. 3) MCG: FETT.J.W. & DEUTSCH.H.F. (1974) BIOCHEMISTRY.13.4102-4114. (CHECKED BY AUTHOR)

NOTES: HUMAN LAMBDA LIGHT CHAINS SUBGROUP V

IDENTICAL SETS OF FRAMEWORK SEGMENTS: . FR1: SET 1: BO[1],HBJ2[2]. (2 IDENTICAL)

FR2: FR3:

FR4: SET 1: BO[1], (IDENTICAL TO 1 HUMAN V-LAMBDA-I: NEWM[1]; AND 1 HUMAN V-LAMBDA-II: WH[3],) SET 2: MCG[3], (IDENTICAL TO 1 HUMAN V-LAMBDA-I: LOC(11].)

IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS:

COR1: SET 1: MCG(3). (IDENTICAL TO 2 HUMAN V-LAMBDA-II: MES(2).VIL(17).)

COR2:

CDR3:

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:

AT POSITION RESIDUES

29	(GLY.ASP)
30	(TYR,ASN)
31	(LYS.ASN)
40	(PRO,ALA)
42	(LYS.ARG)
46	(LEU.VAL)
47	(ILE.VAL)
49	(TYR, PHE)
52	SER.ASNI
53	(LYS.GLY)
68	(GLY,ASP)
79	(ARG.GLN)
92	(VAL.GLU)
93	(GLY,ASP)
94	(SER.ASN)
95	(ASP.ASN)
100	
	(THR.GLY)
104	(LEU.VAL)
107	(ARG.GLY)

MA	INVARIANT	1	2	3	4	5.	6 GIO	7	8 WAN	9 WIN	10 NIG	JAM ·	12 MOR	13 KIN	# OF SEQUENCES	# OF AMINO ACIDS	OF MOS	RRENCES	VARIABILI
	RESIDUES	AMYLOID	SŨT #	AMYLOID +RS		LEV CL					-48				13			(ASN) (PHE)	3.3
	0 1 2	ASP	ASP	ASP PHE MET	asn PHE MET	asn PHE MET	ASN PHE MET	ASP PHE MET	asn PHE ile	asn PHE MET	asn Ieu MET	ASP PHE MET	asn Ieu MET LEU	ASN PHE MET LEU	13 13 13 13	2 2 1	12	(PHE) (MET) (LEU)	2.4 2.2 1.
	3 4 LEU 5	MET LEU THB	LEU THR	THR	THR	LEU THR	LEU THR GLN	LEU THR	LEU THR GLN	LEU THR GLN	LEU ile GLN PRO	LEU THR glu PRO	THR	10U GLN	13 13 13	321	12	(THA) 2(GLN) 2(PRO)	3.5 2.2 1.
	6 7 PRO	GLN PRO HIS	GLN PRO HIS	GLN PRO HIS	HIS	GLN PRO HIS SER	PRO HIS SER	GLN PRO HIS SER	SER	PÃO SER	PRO pro SER	PRO HIS SER	PRO HIS SER	PRO HIS SER	11 13	2	10	(HIS) S(SER)	2.2 1.
	9 SEA	SEA VAL	SER VAL	VAL	SER VAL	VAL		VAL	VAL	VAL SER GLU	VAL	VAL SER	leu SEA	VAL	13 13 12	2	1	2(VAL) 3(SER) 1(GLU)	2.2 1. 2.2
	11 12 SER 13 14 SER	SEA GLU SEA	SEA GLU SEA	VAL SER GLU SER	VAL SER GLU SER	SER GLU SER	VAL SER GLU SER	GLU SER	SER		GLU SER	GLU SER PRO	SER PRO		12 11 12	1	1	1(SER) 2(PRO) 2(GLY)	1. 1. 1.
	15 PAO 16 GLY	PRO	PRO GLY LYS	PRO	PRO GLY LYS	PRO GLY LYS	PRO GLY LYS THR	PRO GLY gly	PRO GLY LYS THR	PRO GLY LYS THR	GLY LYS THR	GLY LYS	GLY LYS THR	、	12 12 11 11	1000	1	0(LYS) 0(THR) 0(VAL)	3.6 2.2 2.2
	17 18 19 -	GLY LYS THR VAL	THR VAL	gin THR VAL THR	THR VAL THR	THR VAL THR	VAL	gly iys VAL ser	VAL	VAL	THR		ile ile ile		11	2 3 3		8(THR) 8(ILE) 0(SER)	4.1 3.8 1.
	20 21 22 SER	THR PHE SER CYS	ile ile SER CYS	SER	ile SER CYS	ite SER CYS	ile SER CYS	ile SER CYS	ile SER CYS		SEA CYS		SEA CYS THR		10 10 9	12	1	0(CYS) 8(THR)	1.
÷	23 CYS 24	THR · GLY	THR	THR GLY	THR ARG SER	THA	THR	SER			THR ARG THR		ALA ASN		7 7 7 7	333		3(+) 4(SER) 3(+)	7. 5.3 7.
	25 26 27 27A	SER GLY GLY	SEP ASP GLY	GLY ASP	GLY	GLY ASN SER GLY SEP		ALA			SER		GLY GLY ASN		7	33		3() 5(GLY) 4(SER)	
	27B 27C	SER	THF	SER	SER 		•				ASP				1	1		1(ASP) 1(SEA)	
	27D 27E 27F	 ILE	ILE	ILE	ILE	ILE					SEP ILE		ILE GLY		7	1	•	7(ILE) 6(ALA)	1. 2.3 4.5
	28 ILE 29 30 31	ALA ASP	ALA GLI	ALA	ALA SER TYR	ALA SEF ASM	1				ALA SEF ASN TYF	4	SER HIS PRO		6 7 7	3 4 3		4(SER) 2(-) 5(TYR)	14. 4.2 1.
	31 32 33 VAL	SER PHE VAL	TYF TYF VAI	A TYR VAL	VAL	VAL	-				VAL	-	VAL		7 7 5	1 1	,	7(VAL) 5(GLN) 6(TRP)	· 1. 1.
	34 GLN 35 TRP 35 TYR	GLN TRP TYR	GLI TRI TYI	9 R	GLN TRP TYR	TRA	2				TRE TYP ARC	3	THP	1	6 6 5	132		6(TYR) 4(GLN) 4(GLN)	1.
	37 38	GLN GLN	GLI	N N	GLN LEU ARG	GLI ARI	G				GLI ARC PRI		PRO	0	5 6	1 2 3		5(ARG) 5(PRO) 4(GLY)	1. 2.4 4.5
	39 ARG 40 41	ARG PRO GLY SER	PH GL AR	Ŷ	PRC GLY SER	VA SE	L				GL GL AL	Y Y	ASP	२	6 6 5	3		4(SER) 5(ALA) 5(PRO)	4,1 1. 1.
	42 43 ALA 44 PRO 45 THR	ALA PRO THR	AL PRI TH	A O	ALA PRO THF		р Н				PR TH TH	о я			. 5 5 5	1		5(THA) 4(THA) 4(VAL)	1. 2. 2.
	46 47	THR VAL ILE		8						•	16 10	Ĕ			5 5 5	· 2 1 2		5(ILE) 4(TYR)	1. 2.
	48 ILE 49 50	ASP	PH	E U	GLL GLL	JGL	U.				AS TH	P R			5 5 5	222		3(GLU) 4(ASP) 4(ASN)	3. 2. 2. 1.
	51 52 53 GLN	ASP ASN GLN	AS TH GL	.N	ASP	N GL	N				AS GL AF PR	N			555	1		5(GLN) 5(ARG) 5(PRO)	1
	54 ARG 55 PRO 56	ARG PRO SER	AF PF SE	A	AR	R LE	10 U				TY GL	<u>R</u>				3		3(SER) 5(GLY) 5(VAL)	5. 1 1
	57 GLY 58 VAL	GLY VAL PRO	GI V/ PF	AL.	GL' VA	L V/ 0 PF	4L 40				· VA PF	10			55	1		5(PRO) 4(ASP) 5(ARG)	. 2
	59 PRO 60 61 ARG 62 PHE	ASP ARG PHE	AS AF	SP	AS AR PH	G AF	1G				AF PH SE	iG ∔E			5 5 5	1		5(PHE) 5(SER)	1
	63 SER 64 GLY	SER GLY SER	S	ER ER	SE GL SE	Y G					GI				5 5	1		5(GLY) 5(SER) 5(SER)	1
	65 SER 66 67 SER	SER ALA	si	ÊR	# SE	R SI	ËR ER				\$1	R			5 5 5	12	:	4(SER) 5(ASN)	2
	68 69 ASN 70 SEA 71 ALA	ASN SEP	A: S	SN ER	AS SE AL	N A	SN ER				S				555	1		5(SER) 5(ALA) 5(SER)	
	71 ALA 72 SER 73 LEU	ALA SEA	S L	LA ER EU	SE LE TH	:U L	LA ER EU HR				ĩ	EU HR	•		5 ´5	1		5(LEU) 5(THR) 4(ILE)	
	74 THR 75 76 SER	THR ILE SEF GLI	1	HA LE ER	VA SE GL	AL I	EA EA				S	LE ER LY			555		1	5(SEA) 4(GLY) 5(LEU)	1
	77 78 LEU	LEL) L		L6 L)	10 L (S 1	EU YS				-	EU HA .SN			5		322	3(LYS) 4(THR) 4(GLU)	
	79 80 81 82 ASP	LYS THF GLU ASF		ILN HA ILU ISP ILU	GI	HR T LU G SP A LU G	HA LU SP SLU				A A T	SP SP HR			5 5 5		1	4(GLU) 5(ASP) 4(GLU)	
	83 84 ALA 85	GL	J 6		AI A	LA A SP A YR 1					A N	LA MET YR			555		1 2 1	5(ALA) 4(ASP) 5(TYR) 4(TYR)	
	86 TYH 87	ASI TYI TYI CYI		ILA ISP IYR IYR IYS	C C	YRI T YS C	YR CYS				F	HE YS			5 5 5		2 1 1	5(CYS)	
~	89 GLN 90 SER	GLI SEI TY		SLN EA	S	EA S YA A	SLN SER PHE ASP				5	SER IYR ASP			5 5 5 5		2 .	5(GLN) 5(SER) 4(TYR) 4(ASP) 3(SER)	
	91 92 93	SE	R A	ASP ARG	ŝ	ER A	ASN				5	SEA			5 5 3		3 4 2 3	2(ASN) 3(ASN)	1
	94 95 95A	AS HI HI	5	HIS			THA ASN BLN				í	LEU			š		3	1(+)	
F 3	958 95C 95D											<u></u>							
	95E 95F 96	VA	L		т	RP AL	GLY VAL								5 5 5		3 1 1	3(TRP) 5(VAL) 5(PHE)	
-	SA PHE	PH	E	PHE	P	LE	PHE GLY GLY GLY					PHE GLY GLY			5 5 5 5 5		1 1 1	5(PHE) 5(GLY) 5(GLY) 5(GLY)	
	100 GLY 101 GLY	GL GL TH	Y Y IA	PHE GLY GLY GLY THR	7	nhR –	GLY THR LYS					GLY THR LYS LEU			5 5 6		1 2	5(LY3) 5(LEU)	
	R 103 LYS 4 104 105 THR	LY V/ TH V/ LE	s	LYS LEU THA	1	EU HB	LEU THA					VAL			6 6 5 6		1	6(THR) 6(VAL) 6(LEU)	
	106 VAL 106A LEU	V/ LE	ιŭ IŪ	VAL LEU GLY	1 (GLY	VAL LEU GLY					LEU SER		LEU SER GLN	6		2	4(GLY) 6(GLN) 6(PRO)	
	107 108 GLN 109 PRO	G		GLN PRO		GLN PRO	GLN PRO					GLN		PRO	ē		۱	6(PRO)	

							(160				·									
AĂN HI	EAVY CHAINS	3 SUBC 1 EU	SHOUP	3	wol	б СА	6 ND CL	мот	BRO 11GG	тно тно	10* STE	11 BEN (i)	12 ZUC	13 Di	14 BOT	0.00M CL	16* MAR	17 FI	18 VU	19 WAR	20 VIL	21 DUN	22 ADA	23 NOR	SAW
01234 56789 0	LEU(.96) SER	PCA VALN GLU VALN SGLN SGLN GLU	PCAL VGLN LEU VGLN SERY ALA GLU	VALNU VALNU VALNU VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA VALNA	PCAL GLN LEU MOLN SER GLN SER ALA GLU	PCALNU GLEU LUR SGLA GLLA VAL	OIN GLN LEU VALN SER GLA GLU	PCA VAL GLN LEU VAL SELN GLN SELY ALA GLU	ULL VALN GLE U GLE VALN GLE ALN GLE ALN GLE ALN	GLU VAL GLU VAL GLU GLU GLU GLU GLU GLU	PCA VAL his LEU VAL SER Ser ALA GLU VAL	GLU GLU GLU VALR GLA GLA GLU VAL	PCA VAL VAL VAL VAL SER ALA BEU	PCA VAL GLN LEU GLY 9 ¹ y	asp ser pro LEU GLAY GLAY GLA SELY GLL	DALS U SULAY	PCA VAL GLN LEU	PCA VAL GLN LEU	PCA VAL GLN LEU	PCA VAL GLN LEU	PCA VAL GLN LEU	PCA VAL GLN LEU	PCA VAL GLN	PCA VAL GLN	PCA VAL glu LEU
10 112 112 115 117 119 212 222		VLYSO YRALS LASS VLYR GSEALS LASS VLY AESYS	GLU VAL LYS PRO GER SEAL BYAL LYS VAL SEAL BYAL LYS VAL LYS VAL LYS SEAL VAL LYS SEAL SEAL SEAL SEAL SEAL SEAL SEAL SEA	VAL LYS PRO GLIB SEAL SEAL VYS VAL SCYS	VALS LYSO YRRL GSEAL & VALS SEAL & VALS SE	VAL arg LYSO GLI ala SEL VAL BR SC SC LYS LYS SC LYS SC LYS SC LYS SC LYS SC LYS SC LYS SC LYS SC SC SC SC SC SC SC SC SC S	VAL arg LYS PRO GLA BRA SEAL arg VAL PRO LYS VAL CYS	VAYSO YRA BI BESYS	VAL LYS PR UYS GLUP SEUS IN SE LYS	GLU VAL LYS PRO GLU SER BIG SER SER LYS	LYS PAO GLY Blo SER LYS VAL SER CYS Bro ALA	SOF PRO GLY	UYS PRO GLY 919 	LYS LYS	gly ile u glyr u	pro pro 111111111									
23 24 25 26 27 28 29		LYS ALA SEA GLY GLY THA PHE SER	UYS Ibr SER GLY OLY THR PHE SER	SER GLY THR PHE ASD	SER GLY GLY THR PHE Val	SER GLY tyr THR PHE SER	ALA SER GLY Iyr THR PHE II0	SEA GLY #5P BSP PHE 85D	giy SER GLY Iyr	giy phe GLY tyr	ALA				lys glu glu ala arg				5. 		<u>.</u>				
30 31 32 33 34 35 35A 35B		ARG SER ALA ILE ILE	GLY TYA THA ILE SER	SEA TYA TYA MET HIS	ASP TYR LYS GLY LEU	HIS TYR ALA MET	ASP SEA TYA ILE HIS	THA TYA ASP ILE HIS TAP						SER	LEU SER GLY ARG ASP	=									
36 37 39 40 41 42 43 44 45 46 47	GLY LEU GLU TAP	TAP VAL ARG GLN ALA PROY GLY GLY LEU TAP	GLY ARG GLY LEU	GLY GLY GLY LEU	LYS GLY LEU		ILE ARG GLN ALA PRO GLY HIS GLY LEU GLU TRP	VAL ARG GLN ALA PRO GLY ARG GLY LEU GLU TRF	-					ILE ARG GLN PRO GLY S GLY GLU GLU GLU TRP VAL	VAL SEA GLNO										
48 49 50 51 52 52A 52B		MET GLY ILE VAL PRO	SER PRC ALA	MET GLY ILE ILE ASN	GLY GLY ILE PRO		VAL GLY TRP ILE ASN PRO							GLU GLU ILE ASF											
52C 53 54 55 56 57 58 59 60		MET PHE GLY PRO ASN TYP ALA GLN LYS	ASF PRO PRO PHE GLN	GLY GLY SEF THF SEF TYF	ASN GLU GLU VAL	1	ASN SEP GLY THP ASN TYP ALA	ASI ARI THI THI TYI GL						TYPE SEF GLI THE ASE TYPE											
61 62 63 64 65 66 67 68 69	ARG	PHE GLN GLN ARC VAL THF	GLUS ARC VAL	J GL	VAI VAI		ARC PHE GLN GLN ARC VAL THE ME	ALI ALI ALI ALI ALI ALI ALI ALI ALI ALI						LYS SEF ARG ME	1										
70 71 72 73 74 75 76 77	SER	ILE THE ASE GLU SEE THE ASE THE ASE THE ASE THE	A LEI P LYI P PRI A SEI A SEI A SEI A SEI A SEI A SEI A SEI	A THI AND ASI ASI ASI ASI ASI ASI ASI ASI ASI ASI	A SEI SP LEY A PAR A SEI A SEI		THE ARC AU SEI SEI SEI THI AU	A TH ARS AS AS SE SE TH TH TH	AGPAR AAAAL					LEI AS THI SE VA AS LEI PH SE	JPAR LNUER		•								
78 79 80 81 82 824 824 826 83 84	в	ME GLI SEI SEI AR SE	T ME GLU LE VA A AS UG AS	T ME U GL U LE L SE	T ME U GL U LE SE		MESE ASE ASE ASE ASE ASE ASE ASE	T ME P GL U LE G TH A	T U U R A					SE E SE VA TH	R		-								
820 83 84 85 86 87 88 89 80 81 92 93 94	ALA	GL AS TH PH PH CY GL	A A A A A A A A A A A A A A A A A A A	Y ALAN YAY YALAN YALANYAN YALAN YALAN YALAN YALAN YALAN YALAN YALAN YALAN YALANY	AP AS A TH A AL A AL A TY A TY	R A L R	SE AL VA PTY CY -AL		A E A A A G		•			TH AL VA TY C)			-								
95 98 99 99 100 100 100		GL TY GL TY SE	Y GI R Y GI GI R V GI R V AS		GU GU GU GU GU GU GU GU GU GU GU GU GU G		SE AS PF TF SE AS TY	A HTSAT	ly sharpa spaya		×		-												
100 100 100 100 100 100 100 100 100	00 00 00 00 00 00 00 00 00 00 00 00 00	PF					Ŷ	A SIA LI AR LI AR					-		ag 2					•		×			
10: 10: 10: 10: 10: 10: 10: 10: 10: 10:	3 4 5 6 GLY 7 8	GTA GGUV	LU T GG GV LLYN GV LLYU L ALA	RP LY LN LN EU AL HR		RPY LN LN EAU AHN AER	TOG GTT VI	HP T LY G LN G HAR L HAR L	RP LLN LHR EU LE LE LE LE LE LE		•			- 66 - 1 - 1	LY ER LY LY EU AL		······································								

-															
HUMAN H	25. KOH	26 AIC	27	28 VAU	29 LE®	SAC	31 DEE	32 LEA	33 HAR	34 HUS	# OF SEQUENCE	# OF) OF	OCCURRENCES MOST COMMON AMINO ACID	VARIABILITY
01 12 34 56 89 90 101 123 14 167 19 201 221 223 224 225 223 224 225 229 300		PCA VAL Ieu	PCA medi GLN	Post val			VAL arg ilə		 pca LEU		30 30 225 14 14 15 15 14 14 14 14 14 14 14 12 11 12 12 11 10 100 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	5668 42184 38528 34853 438255 38468 57		21(PCA) 25(VAL) 22(GLN) 24(LEU) 11(VAL) 11(VAL) 14(SER) 14(GLY) 12(ALA) 12(ALA) 12(ALA) 12(ALA) 12(CLV) 12(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(LYS) 13(7.1 7.2 7.9 2.1 5.1 2.8 1. 2.5 3.5 2.3 3.5 2.3 2.2 2.2 3.5 2.2 2.2 3.5 2.2 2.2 3.5 2.2 2.2 3.5 2.2 2.2 3.5 2.3 3.7 3.7 14. 3.2 8.5 2.3 1. 1. 2.6 3.7 2.2 3.5 2.3 3.7 2.2 3.5 2.3 3.7 1. 2.8 3.5 2.3 3.7 2.2 3.5 2.2 3.5 2.3 3.7 2.2 3.5 2.2 3.5 2.3 3.7 2.2 3.5 2.3 3.7 2.2 3.5 2.3 3.7 2.2 3.5 2.3 3.7 2.2 3.5 2.2 2.3 3.7 2.2 2.2 3.5 2.2 2.3 3.5 2.2 2.3 3.7 2.2 2.2 3.5 2.2 2.3 3.7 2.2 2.3 3.7 3.7 3.7 3.7 3.7 3.7 3.7 3.7 3.7 3
31 C 33 D 34 F 35			=		11111						8 8 8 8	2 6 4 5		4(ILE) 3(HIS)	24. 8. 13.
1 354 355 36 36 37 38 39 40 41 F 42 2 43 44 45 45 46 46 46 47 48	3										8 8 8 8 8 8 8 8 8 8 7 7 7 7 7 7 7 7 7 7		22	7(TRP) 5(VAL) 7(AGG) 7(GLN) 6(ALA) 7(PRO) 7(GLV) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU) 7(GEU)	2.3 4.8 2.3 4. 2.3 2.3 14. 1. 1. 1. 1. 1. 1. 2.3 2.3 4.9. 4.2 2.3 2.3 2.3 2.3 2.3 2.3 2.3 2.3 2.3 2
50 52 52 52 52 52 52 52 52 52 52 52 52 52							• • • • • • • • • • • • • • • • • • •				e 	7	38 3 65 35463 43434 5	4(PRO) 2(SER) 2(+) 4(GLY) 4(THR) 2(ASN) 5(TYR) 3(ALA) 3(ALA) 3(PHE) 3(PHE) 4(GLN) 3(GLY)	21. 18. 5.3 18. 7. 21. 4.2 8. 4.2 6. 12. 5. 7.
FR3	67890123456778930122A8 823456778930122A8 823888888888888888888888991923							TYR TYR Cys Thy Gly			MET ASER UGLX ALARGL GLXX THRA VALR TYPR CALA	76667 77777 77777 777888 888888 88899 9999	1000 00051 00000 00050 044 01000 1000	7(ARG) 5(YAL) 5(THR) 3(+) 4(THR) 3(+) 7(5ER) 3(+) 4(ALA) 5(CLU) 7(5ER) 4(ALA) 5(CLU) 7(5ER) 4(ALA) 5(CLU) 7(5ER) 6(5ER) 6(5ER) 6(5ER) 7(LEU) 4(ARG) 6(VAR) 6(VAR) 6(VAR) 6(VAR) 6(VAR) 6(VAR) 6(VAR) 6(VAR) 6(ALA) 8(CYS) 8(B(ALA) 6(ALA) 8(CYS) 8(B(ALA) 6(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(B(ALA) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(CYS) 8(C	4, 1, 4, 2,3 2,3 1, 2,3 4,5 18,
00	93 94 95 95 97 98 98 97 98 98 98 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 98 99 90 90 90 90 90 90 90 90 90 90 90 90							GLY MET			ASX AAR SX AAR S	7777666555442222113788889	56 655554 43422 22110	6(TRP) 6(GLY)	21. 21. 15. 15. 15. 15. 15. 15. P) 9. 3. 2.7: 4.

ANTIBODY SPECIFICITIES: HUMAN HEAVY CHAINS SUBGROUP I

2) SIE: ANTI-HUMAN GAMMA & GLOBULIN; WA IDIOTYPE ANTI-HUMAN GAMMA & GLOBULIN; WA IDIOTY

WOL: 10) STE: COLD AGGLUTININ WITH ANTI-BLOOD GROUP I ACTIVITY

16) MAR: ANTI-LIPOPROTEIN LIPASE 25) KOH: ANTI-HUMAN GAMMA G GLOBULIN

CLASS: HUMAN HEAVY CHAINS SUBGROUP I

1) EU: IGG1-KAPPA 2) SIE: IGM-KAPPA 4) WOL: IGM-KAPPA 5) CA: IGG1-6) ND'CL: IGE-7) HOT: IGG-8) BRO'IGG: IGG-KAPPA 10) STE: IGG1-11) BEN(I): IGG3-12) ZUC: IGG3-13) DI: IGM-14) BOT: IGM 15) OMM'CL: 1003-16) MAR: IGM 19) WAR: IGG1-IGG3-LAMBDA 20) VIL: 21) DUN: IGG4-22) ADA: IGA-23) NOR: IGA-24) SAW: 25) KOH: 1662 IGM-LAMBDA 26) RIC: 27) WIS: IGG3-IGG3-28) VAU: IGG1-29) LEB: 30) SAC: 1001

IGG1-KAPPA 34) HUS: 1663-

REFERENCE: HUMAN HEAVY CHAINS SUBGROUP I

1) EU: CUNNINGHAM, B.A., RUTISHAUSER, U., GALL, W.E., GOTTLIEB, P.D., WAXDAL, M.J. & EDELMAN, G.M. (1970) BIOCHEMISTRY, 9.3161-3170. (CHECKED BY AUTHOR)

AUTHOR)
AUTHOR)
SIE: ANDREWS,O.W. & CAPRA,J.D. (1981) PROC.NAT.ACAD.SCI.USA.76.3799-3803; ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822. (CHECKED BY AUTHOR 11/15/82); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.582-5830. (CHECKED BY AUTHOR 11/15/82); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.582-5830.
HOYCL: RECHAVIG.RAM.O.GLAZERL.ZAKUT,R. & GIVOL,D. (1983) PROC.NAT.ACAD.SCI.USA.80.855-859. (CHECKED BY AUTHOR 01/04/83)
WOL: ANDREWS,D.W. & CAPRA,J.D. (1961) PROC.NAT.ACAD.SCI.USA.70.3030; ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5816-5822.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830. (CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED BY AUTHOR 11/15/92); ANDREWS,D.W. & CAPRA,J.D. (1981) BIOCHEMISTRY.20.5822-5830.
(CHECKED NUCH, A YON BAHR-LINDSTROM,H. (1974) PROGRESS IN IMMUNOL CONCEPTS AND DEVELOPMENTS. BACH.M., ED., PP.1-36. MARCEL BAHR-LINDSTROM,H. (1978) IN IMMEDIATE HYPERSENSITIVITY MODERN CONCEPTS AND DEVELOPMENTS. BACH.M., ED., PP.1-36. MARCEL DEKKER,NEW YORK; KENTEN,J.H.,MOLGAARD,H.Y.,HOUGHTON,M.DERBYSHRE,R.B., VINEY,J, BELL,L.O. & GOULD.H.J. (1982) PROC.NAT.ACAD.SCIUSA.73,6661-6665.
(MOT.I. (1982) MOL.IMMUNOL...19.1095-1103; KOJIMA,M.KOIDE,T.ODANI,S. & ONO.T. (1986) MOL.IMMUNOL..23,169-174.

KOJIMA, M. ODANI, S. & ONO, T. (1982) MOL.IMMUNOL., 19, 1095-1103; KOJIMA, M. KOIDE, T., ODANI, S. & ONO, T. (1986) MOL.IMMUNOL., 23, 169-174. (CHECKED BY AUTHOR 08/08/86) 7) MOT:

BI BROILD BI ROTHON WOULD' B) BROILD BI ROPPERJE.NOVES.C.HEINRIKSON,R. & KESSELJ.W. (1976) J.IMMUNOL..116.743-746; HOPPERJE. & BRAHN,E. (1977) J.IMMUNOL..119. 847-849. (CHECKED BY AUTHOR 09/25/78 WHO POINTED OUT THAT BRO' IS SAME AS BRIGG AND SUGGESTED THAT IT SHOULD BE RENAMED AS BROIGG) BRO'IGG: HOPPERJ.E., NOYES.C. HEINRIKSON,R. & KESSELJ.W. (1976) JIMMUNOL.115,742-745; HOPPERJ.E. & BRAHNE, (1977) JIMMUNOL_119, 847-849; (CHECKED BY AUTHOR 09/29/78 WHO POINTED OUT THAT BRO'IS SAME AS BRIGG AND SUGGESTED THAT IT SHOULD BE RENAMED AS BROIGS.
PTHO: HOPPERJ.E. & BRAHNE, (1977) JIMMUNOL, 119,847-849; (CHECKED BY AUTHOR)
STE: FISHER,C.E.PALM.W.H. & PRESS.E.M. (1989) FEBS LETTERS,520-22 (CHECKED BY AUTHOR)
STE: FISHER,C.E. FRANK,W.H. & PRESS.E.M. (1989) NATURE224,897-999; (CHECKED BY AUTHOR)
BEN(I): KAPLANA,P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
DIC HEMISTRY, 1324224389. (CHECKED BY AUTHOR 09/2783)
BOT: BARNIKOL-WATANABE,S., MIHAESCO.C., BARNIKOL,H.U. & HLISCHMANN,N. (1984) Z.PHYSIOL.CHEM.365,105-118.
BOM: KAPLANA,P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
STE: SANKOL-WATANABE,S., MIHAESCO.C., BARNIKOL,H.U. & HLISCHMANN,N. (1984) Z.PHYSIOL.CHEM.365,105-118.
BOM: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: MONTGOMERY,P.C., BELLO,A.C. & ROCKEY,J.H. (1970) BIOCHIM.BIOPHYS.ACTA.200,258-286. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-811. (CHECKED BY AUTHOR)
WI: KAPLAN,A.P., HOOD, L.TERRY,W.D. & METZGER,H. (1971) IMMUNOCHEMISTRY,8,801-8

WANG.A.C. & FUDENBERG.H.H. (1975) ARCH.BIOCHEM.BIOPHYS., 168.657.654. (CHECKED BY AUTHOR 09/23/77)

NOTES: HUMAN HEAVY CHAINS SUBGROUP I

IDENTICAL SETS OF FRAMEWORK SEGMENTS

1: VAU[28].LEB[29]. (2 IDENTICAL) SET

EU[1].HG3'CL[3]. (2 IDENTICAL) WOL[4]. (IDENTICAL TO 2 HUMAN V-H-III: TIL[4].TE([10].) FR2: SET 1: SET 2:

FA3:

SET 2: WOLL4). (IDENTICAL TO 2 HUMAN V-H-III: (ILL4). ETITUS) SET 1: NOCL41. (IDENTICAL TO 1 HUMAN V-H-III: CEGECUT061.) SET 1: WOLL41. (IDENTICAL TO 2 HUMAN V-H-III: MCE'[4],NZU[15]; 4 HUMAN V-H-III: TIL[4],DOB[31],WEA[33],NIE[34]; AND 1 MOUSE V-H-III: MOPC/421.(80-7) SET 2: NOCL43. (IDENTICAL TO 2 HUMAN V-H-III: HIG1'CL(10]; 1 HUMAN V-H-III: U266'CL(105]; AND 1 MOUSE V-H-IIA: HDEX12[15].) FR4:

IDENTICAL SETS OF COMPLEMENTARITY DETERMINING REGIONS:

CDB1:

34) HUS:

COR2 CDR3:

SET 1: HG3 CU3). (IDENTICAL TO 1 HUMAN V-H-III: LAMBDA-VH26 CU2I: 1 MOUSE V-H-IB: PJ14 CU22); AND 5 MOUSE V-H-IB: 166-2 CU3). 185-1 CU51,102 CU151,23 CU181,3 CU261.) SET 2: NO CU51, (IDENTICAL TO 1 HUMAN V-H-III: U266 CU1061.)

IDENTICAL SETS OF J-MINIGENES: SET 1: NO'CLI6J. (IDENTICAL TO 1 HUMAN V-H-I): HIG1'CL[10]; AND 1 HUMAN V-H-III: U266'CL[106].)


NOTES: HUMAN HEAVY CHAINS SUBGROUP I (cont'd) SPECIFIC NOTES:

3) HG3'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN FETAL LIVER GENOMIC DNA.


163

3) HOUCL: THE ANIMO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF MOUSE CDNA. IT
6) ND'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF MOUSE CDNA. IT
6) ND'CL: THE AMINO ACID SEQUENCE DETERMINED EARLIER EXCEPT THAT THE AMINO ACID SEQUENCE DETERMINATION GAVE PCA AT POSITION 1, VAL AT 2, VAL AT 34, GLY AT 35, ILE AT 49 AND HIS AT 49.
7) MOT: PAPAIN CLEAVES BETWEEN ARG 56 AND THR 57, AND BETWEEN ARG 62 AND SER 63.
12) ZUIC: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.
14) BOT: IT WAS FROM A CASE OF IGM HEAVY CHAIN DISEASE.
15) OMM'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CELL LINE CDNA. IT
15) OMM'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CELL LINE CDNA. IT
16) OMM'CL: THE AMINO ACASE OF HEAVY CHAIN DISEASE.
17) MOT: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.
18) OMM'CL: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CELL LINE CDNA. IT
19) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
10) MOT: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.
11) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
12) DISTICUT: THE AMINO ACID SEQUENCE IS OBTAINED BY TRANSLATING THE NUCLEOTIDE SEQUENCE OF A CLONE OF HUMAN CELL LINE CDNA. IT
10) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
11) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
12) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
14) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
15) MANCH AND ACASE OF HEAVY CHAIN DISEASE.
16) WAS FROM A CASE OF HEAVY CHAIN DISEASE.
17) WIS IT WAS FROM A CASE OF HEAVY CHAIN DISEASE. WAS FROM A CASE OF HEAVY CHAIN DISEASE.
27) WIS: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE. ITS RESIDUES AT POSITIONS 108 AND 109 ARE ASN AND CYS RESPECTIVELY, WHICH DO NOT CORRESPOND TO THE USUAL RESIDUES FOUND AT THESE POSITIONS IN HUMAN HEAVY CHAIN SUBGROUP I.
28) VAU: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.
29) LEB: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.
30) SAC: IT WAS FROM A CASE OF HEAVY CHAIN DISEASE.

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:



HUMAN	HEAVY CHAIN	SUBGR	OUP II 2 3 AW OL	мсе. *	CE-1 CL	6 НЕ 1	7 SUP-T1 VH-JA CL	B* NEWM	9 WAH	10 HIG1 CL	11 CAR	12 SA	13 10	14 SPA #	15 NZU #	18 EAI	N OF SEQUENCES	# OF AMINO ACIDS	OCCURRENCES OF MOST COMMON AMINO ACID
0 1 2 3 4 5 8 7 7 8 9 10 1 12 3 14 4 5 8 7 7 8 9 10 1 12 3 14 4 5 8 7 7 8 9 10 1 12 1 13 14 1 16 6 1 1 17 18 9 20 2 2 2 2 4 2 2 2 6 6 2 7 2 8 9 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 2 9 2 7 2 8 2 9 2 9 2 9 2 7 2 8 2 9 2 9 2 9 2 9 2 9 2 9 2 9 2 9 2 9	GLY LEU VAL LEU LEU THR CYS GLY	VT J AGSSGR ALV "R TOTTT ITOTT SOISS ALRU GURYO ALUSO RINRUH URVSTR VTL AGSSGR ALVYR TGTTLT UTOTA SOF SGPHER	CALRU GUERY O ALUAYA MULAUR URSHE RYERU GUERY O ALUAYA MULAUR URSHE RYERU SULERY O ALUAYA MULAUR URSHE RYERU SULERY O ALUAYA MULAUR SURSHE RYERU SULERY SURSHE RYERU SULERY SURSHE RYERU SULERY SURSHE RYERU SULERY SULERY S	INAU SUBYO AULSO ANDUR URSRE RYE	LE RGURY AGURY CAULAN ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAYS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS ALEVAS	PCALRU SUNTO TLEAUR TOTLET LITCTHO SELEN	A	PCAL DUL GLEXARY GSELG GELGO YUL GO ULANGO YARU ULAGO GSELGO YUL GO GSELGO YUL GO GSEL	areiginu ruhayo yuusoo	ginu non py a structure of the structure	LEU THA CYS THA Val SEA GLY	THR CYSR CYSR Val SER SER SER		PCA 9lu 9lu val ARGU SER 			12 12 12 12 12 12 12 12 12 12 12 12 12 1	3442 5 5 4 4 4 4 4 3 12 31122 33212 11133 21534 4	8(PCA) 9(YAL) 6(THR) 11(LEU) 9(SER) 9(SER) 10(GLU) 9(GLU) 9(SER) 10(VLL) 10(VLL) 10(VLL) 9(PRO) 4(+1) 10(VLL) 9(PRO) 5(GLU) 9(PRO) 5(GLU) 9(PRO) 5(GLU) 9(PRO) 5(GLU) 9(THR) 10(VLL) 11(LEU) 11(LEU) 11(LEU) 11(SER) 12(SER) 5(LEU) 2(SER) 5(LEU) 2(SER) 5(LEU) 2(SER) 5(LEU) 2(SER) 5(LEU) 2(SER)
30 31 32 0 33 0 34 8 34 1 35 35 35	A	SER S SER C THR C GLY T MET M CYS C VAL	SER SE	R THR SER G GLY T VAL G GLY L VAL R GLY	SER VAL SER	thr THR ASP GLY VAL ALA VAL GLY	SER GLY TYR TYR TRP GLY	SER ASN ASP TYR TYR TYR THR	ARG THA GLY TYR TYR TYR GLY	SEA GLY TYA TAP SEA TAP							10 10 10 10 10 8 7 10	5 7 4 8 3 3	4(THR) 2() 5(GLY) 4(MET) 2() 6(VAL) 4(GLY) 10(TRP)
36 37 38 39 40 41 F 42 2 43 44 45 46 45 46 47 48 49	TRP ARG PRO GLY LEU GLU TRP	TRP ILE ARG GLN PROOI GLY LGLY GLY LGLU GLU TRP LGLA ARG	ALA A	E ILE ARG ARG GLN ARG ARG ARG ARG PRO L S ALLEU L U C L U C L U C L U C L U C L U C L U C L U C C L U C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C C L N C L N C C L N C L N C C L N C L N C C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N C L N D L N D L N D L N D L N D L N D L N D L N D L N D L D D L D D D D	TRP ILE AGLN PRO GLY ALA UGUP LEU ALE ALA ALA ALA ILE	TAP ILE ARG GLY PAO GLY ARG ALG LEU TRP LEU ALA TRP LEU	TAP ILE AAG GUN PRO PRO PRO CLS USU GLY USU GLY USU TAP ILE CLY SILE	TRP VAL ARG GLN PROO PROO PROO PROO PROO PROO PROO PRO	TAP ILE ARG GLY PRO PRO GLY LES GLY ILE GLY VAL	ILE ARG GLN PRO GLY ARG GLY GLU GLU GLU GLU							10 10 10 10 10 10 10 10 10 10	2 12311 3211 22 746	9(LE) 10(ARG) 9(GLN) 8(PRO) 10(PRO) 10(GLY) 6(LYS) 5(+) 10(LEU) 10(GLU) 10(TRP) 6(LEU) 6(ALA) 3(ARG) 6(ILE) 3(ASP) : 3(ASP)
5012 522555 5455 55455 55455 55455 55455 66666	26 26 35 35 37 39 30 12 20 21 22 23 24 24 24 24 24 24 24 24 24 24 24 24 24	ASP TRP ASP ASP LYSP TYP	ILE AAAAUPTT STSL	LEN TARF FRANK	ASP TRPP ASP ASP ASP TYPR TYPR THAR SEU	LEUR TRA ASSIGN ROAUS	T HI - R YARAA NOAUS	PHE TYRS GLYA SSPA ASPA ASPA THROU LARG	TYPE TYPE THE GLIVE SEE TYPE ASSO SEE ASSO	HISA SELEPT							9 9 10 10 10 10 10 10 10 10 10	2 55 24 64 4 :32 5 21 3 3	1(∸) 4(TRP) 5(ASP) : 4(ASP) 6(ASP) 5(ASP) 4(LYS) 6(TYR) 7(TYR)
8 8 8 8 8 8 8 8 8 8 9 9 9 9 9 9 9	5 7 ARG 9 9 0 1 1 2 3 4 SER 5 6 6 7 ASN 6 9 9 0 1 LEU 1 2 2 7 2 9 0 0 1 LEU 1 2 2 7 3 4 4 5 6 8 9 9 0 1 1 2 2 7 8 9 9 0 0 1 1 2 2 7 9 9 0 0 1 1 2 2 7 9 9 0 0 1 1 2 2 7 9 9 0 0 1 1 2 2 7 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 1 2 3 4 9 9 0 0 1 2 3 4 9 9 0 0 1 2 3 4 9 9 9 0 0 1 1 2 2 7 9 9 0 0 1 2 3 4 9 9 9 0 0 1 2 2 7 9 9 9 0 0 1 2 2 7 9 9 9 0 0 1 2 2 7 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	THR	TALST SLAAS LAGVY LINTA VGPGA TATTT CAARLAV SLASTS LAGVY LINTA LAPPAR SALAL USTAR LYPERA ALTY CAARLAV	HR GEUR BYSNER SSUAL ULT ILTAS MERGAVAS EGUTYYN YN	HAGUA ASSAR SAN L JALAN TROLP RYRRE SAG	SEROURAL REGPERER SNNLL UHETERN TEPOOLP ALTER SNNLL UHETERN TAASPRE SNNLL UHETERN TAASPAAL ALTEVE YSLL	SEROLA RALARA SANDARA LARAAP RALARA YAYA	SAVTN LVAFER SZLER LRELSS VTAAA TAVTT SAA	GLACINE RESULTE ESERE ESALS TALETY YAAR	SAVTIL SLOPPE SUSPER UNUGR TRAAP RATER SAG					MEPONA ASPAL APROL ASGLYRA ARG ARG ARG		10 10 10 10 10 10 10 10 10 10	1034 34001 01000 15453 05001 00010 1000	10(ARG) 6(LEU) 8(THR) 7(LSER) 5(LE) 5(LSER) 9(THR) 10(ZSER) 7(LYS) 10(ASN) 9(GLN) 8(VAL) 9(GLN) 9(GLN) 10(LEU) 3(+) 10(LEU) 3(+) 11(LSER) 7(LYS) 10(ASR) 9(GLN) 10(LEU) 3(+) 11(LSER) 10(ASR) 9(GLN) 11(ASR) 9(GLN) 11(ASR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(THR) 9(TH
CDRR310 CDRR310 FR11	14 15 16 16 17 18 18 19 19 10 10 10 10 10 10 10 10 10 10	ARDE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE. ALE.	SERS VIEN	AR P P P P P P P P P P P P P P P P P P P	GOOPEERYNUYY ALABER YNUYY ALABER YNUYYY ALABER YNUYYYY ALABER YNUYYYY ALABER YNUYYYY ALABER YNUYYYY ALABER YNUYYYY ALABER YNUYYYYY ALABER YNUYYYYY ALABER YNUYYYYA ALABER YNUYYYYYYYYYYYYYYYYYYYYYYYYYYYYYYYYYYY	ARGE LEUS	ALG GGO VAR ARA	ASA LEDUCYE	NU CAYSE AUGHE SASSIL SVIFEGE GTTV.H	OOORA PEYRY RPPY & PLAY N 188	1977 1979 1979 1979 1979 1979 1979 1979				PROOP PTRAFER AFU VALUE PT SASE UGUYA PT SASE UGUYA PSECOFUL VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE VALUE	ASI VA GLI AR TH VA SE	11 11 11 11 11 10 10 10 77 66 54 34 8 12 12 12 12 12 12 12 12 12 12	977788 66554 243334 34 2224 243334 34 2224 2233	2(+) 4(PRO) 4(PRO) 3(ARG) 2(+) 2(+) 2(+) 2(+) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-) 2(-)



ANTIBODY SPECIFICITIES: HUMAN HEAVY CHA IBGROUP II B) NEWM: ANTI-3-(3-HYDROXY-3,7,11,15".TETRAMETHYL HEXADECYL) 2-METHYL 1.4 NAPHTHOOUINONE(VIT.K10H)

CLASS: HUMAN HEAVY CHAINS SUBGROUP II

- 1) COR: IGG1-2) DAW: IGG1-LAMBDA
- 3) OU: IGM-KAPPA
- 4) MCE': IGM-KAPPA 6) HE: IGG1-
- 6) NEWM: IGG1-LAMBDA 9) WAH: IGD-LAMBDA
- 12) SA: IGG2-LAMBDA
- 15) NZU: IGM-
- 16) ERI: IGD-

REFERENCE: HUMAN HEAVY CHAINS SUBGROUP II

- 1) COR: PRESS.E.M. & HOGG.N.M. (1970) BIOCHEM.J.,117,641-650. (CHECKED BY AUTHOR) 2) DAW: PRESS.E.M. & HOGG.N.M. (1970) BIOCHEM.J.,117,641-660. (CHECKED BY AUTHOR) 3) OU: PUTNAM.F.W.,SHIMIZU,A.,PAUL.C.,SHINODA,T. & KOHLER,H. (1971) ANN.N.Y.ACAD.SCI.,190,83-103. (CHECKED BY AUTHOR 06/15/83) 3) OU: PUTNAM.F.W.,SHIMIZU,A.,RAUL.C.,SHINODA,T. & KOHLER,H. (1971) ANN.N.Y.ACAD.SCI.,190,83-103. (CHECKED BY AUTHOR 06/15/83) 4) MCC: GERBER-JUENSON,B.,KAZIN,A.,KEHOE,J.M.,SCHEFFEL,C.,ERICKSON,B.W. & LITMAN,G.W. (1981) J.IMMUNOL.,126,1212-1216. (CHECKED BY AUTHOR 12/15/80)

- 12/15/80) 5) TE-1 'CL: TAXAHASHI,N.,NOMA,T. & HONJO,T. (1984) PROC.NAT.ACAD,SCI.USA.81,5194-5198. 6) HE: CUNNINGHAM,B.A. GOTTLIEB,P.D.,FFLUMM,M.N. & EDELMAN,G.M. (1971) PROGRESS IN IMMUNOLOGY (B.AMOS.ED.),ACADEMIC PRESS,N.Y.,PP.3-24. 6) HE: CUNNINGHAM,B.A. GOTTLIEB,P.D.,FFLUMM,M.N. & EDELMAN,G.M. (1971) PROGRESS IN IMMUNOLOGY (B.AMOS.ED.),ACADEMIC PRESS,N.Y.,PP.3-24. 6) HE: CUNNINGHAM,B.A. GOTTLIEB,P.D.,FFLUMM,M.N. & EDELMAN,G.M. (1971) PROGRESS IN IMMUNOLOGY (B.AMOS.ED.),ACADEMIC PRESS,N.Y.,PP.3-24. 7) SUP.T1 VH-JACL: DENNY,C.T.,YOSHIKAI,Y.,MAK,T.W.,SMITH,S.D.,HOLLIS,G.F. & KIRSCH,I.R. (1986) NATURE 320,549-551. 7) SUP.T1 VH-JACL: DENNY,C.T.,YOSHIKAI,Y.,MAK,T.W.,SMITH,S.D.,HOLLIS,G.F. & KIRSCH,I.R. (1986) NATURE 320,549-551. NUMARCH, DENNING, C. L., VOSHIKALY, MAK, T.W., SMITH, S.D., HOLLIS, G.F. & KIRSCH, I.R. (1966) NATURE, 320,549-551.
 SUP-TI VH-JA*CL: DENNY, C.T., YOSHIKALY, MAK, T.W., SMITH, S.D., HOLLIS, G.F. & KIRSCH, I.R. (1976) PROC. NAT. AGAD. SCI.USA, 71, 340,0344. (CHECKED BY AUTHOR WHO CORRECTED RESIDUES 6,9, 15, 16, 24, 26, 27, 29 THROUGH 328, 056 0, 48 GIVEN IN TABLE). CONFECTED RESIDUES 6,9, 15, 16, 24, 26, 27, 29 THROUGH 328, 056 0, 48 GIVEN IN TABLE). COTTOR FILLS BOOK, AND HAS CORRECTED RESIDUES 6,9, 15, 16, 24, 26, 27, 29 THROUGH 328, 056 0, 48, 054,056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48, 056 0, 48,

NOTES: HUMAN HEAVY CHAINS SUBGROUP I

IDENTICAL SETS OF FRAMEWORK SEGMENTS:

- FR1: SET 1: SUP-T1 VH-JA'CL[7].WAH[9]. (2 IDENTICAL)
- FR2:
- SET 1: MCE'[4],NZU[15], (2 IDENTICAL HUMAN V-H-II; ALSO 1 HUMAN V-H-I: WOU[4]; 4 HUMAN V-H-III: TIL[4],DOB[31],WEA[33],NIE[34]; AND 1 MOUSE V-H-IIIA; MOPC47A[48].) SET 2: HIGT'CL[10], (IDENTICAL TO 1 HUMAN V-H-I: ND'CL[6]; 1 HUMAN V-H-III: U266'CL[106]; AND 1 MOUSE V-H-IIA; HDEX12[15].) FR4:

IDENTICAL SETS OF J-MINIGENES: SET 1: HIGT'CL(10), (IDENTICAL TO 1 HUMAN V-H-I: ND'CL(6); AND 1 HUMAN V-H-III: U266'CL(106).)

SPECIFIC NOTES:

4) MCE': IT IS A CRYOIMMUNOGLOBULIN AND IS DESIGNATED BY THE AUTHORS AS MCE. IN ORDER TO DIFFERENTIATE IT FROM ANOTHER MCE SEQUENCED BY CAPRA ET AL., IT IS DENOTED AS MCE.

5) CE-1 'CL: CELL LINE CESS
7) SUP.TI VH.JA'CL: IT IS FROM A PATIENT SUFFERING FROM CHILDHOOD T.CELL LYMPHOMA WITH INV(14)(1).2:032.2). THE INVERSION ON CHILDHOOG T.CELL LYMPHOMA WITH INV(14)(1).2:032.2). THE INVERSION ON CHILDHOGO THE CHILDHOGO TO CHIL TU CHILDHOGO TO CHILDHOGO TO C

+ THE FOLLOWING WERE EQUALLY AND MOST FREQUENTLY OCCURRING:



ΗU	MAN I	INVARIANT RESIDUES	15 SUB 1* TUR	2 LAMBDA -VH26 'CL		4 TIL	was	6 HF2- 1/13B	7 HF2- 1/17	8 HF2- 18/2	9 H11 'CL	10 TEI	11 BRO IGM	12 GR'	13 WAT	14* LAY	15 GRA'	16 FR #	17 MU	18 VIN	19 HF3- 16/6	20 BEN (III)	21 ZAP	22 JON	23 KEA
	0 1 2 3 4 5 6 7 8 9	LEU(.97) SER(.97) GLY(.99) GLY(.98)	GLU VAL GLN LEU GLU GLU GLU GLY	GLU GLU VAL GLN LEU GLU SER GLY GLY	GLU VAL GLN LEU GLU GLU GLY	GLU VAL GLN LEU GLU SER GLY GLY	GLU VAL GLN LEU GLU SER GLY GLY	GLU VAL GLN LEU GLU GLU GLU GLY	GLU VAL GLN LEU LEU GLU SER GLY GLY	GLU VAL GLN LEU gln SER GLY	GLU VAL GLN LEU Val GLU SER GLY GLY	GLU VAL GLN LEU VBI GLU SER GLY	GLU VAL GLN LEU Val GLU SER GLY	GLU VAL GLN LEU Val GLU SER GLY GLY	GLU VAL GLN LEU Val GLU SER GLY GLY	ala VAL GLN LEU GLU SER GLY GLY	GLU VAL GLN LEU Val GLU SERY GLY	GLU VAL GLN LEU VBI ASP SER GLY GLY	GLU GLU GLN LEU VAL GLN VAL GLN VAL GLN VAL GLN VAL GLN VAL	GLU VAL GLN LEU Val GLU SER GLV GLY	GLU VAL GLN LEU GLN LEU GLY GLY	GLU VAL GLN LEU Val GLU SER GLY GLY	GLU VAL GLN LEU Val GLU SER GLY GLY 8la	ASP VAL GLN LEU Val GLU SER GLY GLY	GLU VAL GLN LEU Val GLX GLY GLY GLY
FR 1	10 11 12 12- 14 15 16 17	PRO(.95) GLY	GLY LEU VAL GLN PRO GLY GLY SER LEU	GLY LEU VAL GLN PRO GLY SER LEU	GLY LEU VAL GLN PRO GLY GLY SER	GLY LEU GLN PRO GLY SEU ARG	GLY LEU VAL GLN PRO GLY SER LEU	GLY VAL GLN PRO GLY SER LEU	GLY LEU VAL GLN PRO GLY SER LEU	GLY LEU VAL GLN PRO GLY SER LEU	GLY LEU VAL GLN PRO GLY SER LEU ARG	GLY LEU VAL GLN PRO GLY SEU SEU ARG	GLY LEU VAL PRO GLY SEU	GLY LEU VALN PRO GLY GLY SEU SEU	GLY LEU GLN P GLY GLN GLY GLY SEU SEU	GLY LEU GLN PRO GLY SER LEU ARG	GLY LEU VAL GLN PRO GLY SER LEU ARG	GLY LEU VAL GLN PRO GLY SER LEU ARG	GLY LEU VAL Iys PAO GLY SEU ARG	GLY LEU GLN PRO GLY SER LEU ARG	GLY LEU VAL GLN PRO GLY SER LEU ARG	LEU ala GLN PRO GLY SER LEU ARG	LEU VAL GLN PRO GLY SER GLY SER GLY AHG	GLY LEU VAL Iys PRO GLY SEU ARG	ULU VAL IVS PRO GLY GLY SEA LEU ARG
	-18 19 20 21 22 23 24 25 26	LEU(.97) CYS SER(.98) GLY(97)	ARG LEU SER CYS ALA ALA SER GLY	ARG LEU SER CYS ALA ALA SER GLY	LEU ARG LEU SER CYS ALA ALA SER GLY	LEU SER CYS ALA ALA SER GLY	ARG LEU SER CYS ALA ALA SER GLY PHE	ARG LEU SER CYS ALA ALA SER GLY	ARG LEU SER CYS ALA ALA SER GLY	ARG LEU SER CYS ALA ALA SER GLY	LEU SER CYS ALA ALA SER GLY	ARG LEU SER CYS ALA ALA SER GLY PHE	ARG LEURS CYS ALA SERS ALA SERY PHE	ARG LEU SER CYS ALA ALA SEP GLY PHE	ARG LEURS CLA ALA SERY ALA SELY PHE	LEU SER CYS ALA SER GLY PHE	LEU SER CYS ALA ALA SEP GLY PHE	LEU SEA CYS ALA ALA SER GLY PHE	LEU SER CYS ALA ALA SER GLY PHE	LEU SER CYS ALA ALA SER GLY PHE	LEU OIN CYS ALA ALA SER GLY PHE	LEU ala CYS ALA ALA SER GLY PHE	LEU SER CYS ALA ALA SER GLY PHE	LEU SER CYS ALA ALA SER GLY PHE	LEU SER CYS ALA ALA SER GLY PHE
CDRI	26 27 29 30 31 32 33 34	GLY(97) PHE(98)	PHE THR PHE SER VAL LEU SER	PHE THR PHE SER SER TYR ALA MET	PHE THR PHE SER SER ALA MET	PHE THR PHE SER THR TYR VAL MET	PHE ser PHE SER THR ASP ALA MET TYR	PHE Jys PHE SER SER TYR ALA MET SER	PHE Iys PHE SER TYR ALA MET SER	PHE THR PHE SER SER TYR ALA MET SER	PHE THR PHE SER TYR TRPT MET HIS	THA PHE SER THA SER ALA VAL TYR	THR PHE SER TYR ASN MET ASN	THR PHE SER ALA ASX TYR MET	THR PHE ASX THR THR THR MET VAL	THR PHE SER ALA SER ALA MET SER	THR IIO SEA LYS THR VAL TYA GLU	THR PHE SER ASX PHE TYR MET ASP	THR PHE Ihr ARG GLY GLY LEU GLU	THR val SER THR ASN TYR MET	SER PHE SER PRO SER ALA MET SER	THR PHE SER THR THR PHE MET ARG	THR PHE SER THR SER ARG PHE	THR PHE SER THR ALA TRP MET LYS	IIe PHE pro TYR
	35 35A 35B 36 37 38 39 40 41	TRP VAL(.95) ARG(.97) GLN(.97)	SER TRP VAL ARG GLN ALA PRO	SER TRP VAL ARG GLN ALA PRO	SER TRP VAL ARG GLN GLY	SEA TRP VAL ARG GLN ALA PRO GLY	TRP VAL ARG GLN ALA PRO GLY	THP VAL ARG GLN ALA	TRP VAL ARG GLN ALA	TRP VAL ARG GLN ALA	TRP VAL ARG GLN ALA PRO GLY	TRP VAL ARG GLN ALA PRO GLY	THP VAL ARG GLN VAL THR GLY	TRP VAL PRO GLY ALA PRO GLY		TAP VAL ARG GLN ALA PRO GLY	TRP VAL ARG GLN ALA PRO GLY	TAP VAL ARG GLN ALA PRO GLY	TRP VAL ARG GLN ALA PRO GLY		TRP VAL ARG GLN ALA	TRP VAL ARG	TRP VAL ARG GLN ALA PRO GLY	TRP VAL ARG GLN ALA PRO GLY	
F R 2	42 43 44 45 46 47 48 49 50	GLY . LYS(.97) LEU(.97) TRP	GLY LYS GLY GLU GLU TRP VAL SER GLY	GLY LYS GLY LEU GLU TRP VAL SER ALA	LYS GLY LEU GLU TRP VAL ALA	LYS GLY LEU GLX TRP VAL GLY ALA	LYS GLY LEU GLÙ TRP VAL ALA				LYS GLY LEU VAL TRP VAL SER ARG	LYS GLY GLU TAP VAL GLY	LYS GLY LEU GLU TRP VAL SER ALA	ARG GLY LEU		LYS GLY LEU GLU TRP VAL ALA TRP LYS	LYS GLY LEU GLU TRP VAL THR TYR VAL	LYS GLY LEU GLU TRP VAL ALA ARG	LYS ALA LEU GLX TAP VAL LEU VAL PHE				LYS GLY LEU GLU TAP VAL GLU PHE ARG	LYS GLY LEU GLU TRP VAL VAL TAP ARG	
ç	50 51 52 52A 52B 52C 53 54 55	LYS	ARG LEU ASN ALA SER SER	ILE SER GLY SER GLY GLY	LYS TYR GLU ASN GLY		TRP LYS TYR GLN GLU ALA SER ASN				ILE ASN SER GLY SER	ARG TYA GLU GLY SER LER LER	ILE GLY THR ALA GLY ASP			GLU ASN GLY ASN ASP	GLN GLN VAL SER LYS PHE		VAL				GLN GLY SER ALA ILE	VAL GLU GLN VAL VAL GLU	
CDR 2	56 57 58 59 60 61 62 63 64		ASNU LEISE ALL SELA SELA GLN	SER THR TYR GLP ASER VAL LYS	ASP LYSS HTY ALPA SEAL ASP ASP ASP ASP	VAL SER SER TYR ALA SER VAL SER LYS	SER HIS PHE ASPR THR VAL ASN				SER THR TYR ALP SER VAL VAL	HIS TYR ALA VAL SER VAL GLN	GLN TYR ALA SEA VAL LYS GLY			LYS HIS TYR ALA ASP SER VAL ASN GLY	SER TYR ALA VAL SER VAL GLN GLY		LYS PHE TYA GLU SEQ ASN		ŕ.		SER HIS TYR ALA SER VAL GLN ALA	GLU LYS ALA PHE ALA SER VAL ASN GLY	
	65 67 68 69 70 71 72 73 74	GLY(.97) ARG PHE(.97) ILE(.97) SER(.97) ARG	GLY ARG PHE THR ILE SERG ARSN ASP	GLY ARG PHE THR ILE SER ARG ASP	GLY ARG PHE THR ILE SERG ASP	GLY ARG PHER THR ILE SERG ASP ASP	GLY ARG PHE THR ILE SER ARG ASN ASP				GLY ARG PHE THR ILE SERG ASP ASN	GLY ARGE PHER ILE SERGN ASP	ARG PHE THR ILE SER ARG ASN ASP			ARG PHE THR ILE SERG ASN ASP	ARG PHR THE SERGN ASSPR					,	ARG PHE THR ILE SEG ASP SER	ARGE THE SEG ASP ASP SER	
FRG	75 76 77 78 79 80 81		ASP SER LYS ASN THR LEU TYR LEU GLN	ASN SER LYS ASN THR LEN LEN MET	ASP SER LYS ASN THEU TYR LEU	ASP SER LYSN THR IIII	SER LYS ASN THR LEU TYR LEU GLN				ALA LYS ASNA LEU TYA LEU MET	SER LYS ASHRUA TYEUN LEUN MEU LEUN	SER LYS ASR LEU TYR LEU ASN MET			SER LYS ASN THR LEU TYA LEU GLN MET	LYS ASN THR LEU TYR LEU GLU MET LYS	MET					LYS ASN THR LEU TYR LEU GLN MET ASN	LYS ASN THR LEU TYR LEU GLN MET	MET
3	62 82A 82B 82C 83 84 85 86 85 86	MET(.95)	LEU SER LEU GLN ALA GLU ASP THB	ASN SER ARG ALA GLU ASP THB	MET ASN SER LEU GLN ALA GLD ASP THR	ASN SER LEU ARG ALA GLU ASP THR	MET ASN ARG LEU GLU ALA GLU ASP THR				ASN SER LEU ARG ALA GLU ASP THR	LEU GLU PRO GLU ASP	ASN SER LEU ARG ALA GLU ASP THB	,		ASN GLY LEU GLN ALA GLU VAL SER ALA	GLY GLU PRO GLU ASP	ASN SER LEU ARG ALA GLX ASX THR		ALA GLU ASP THR			THR GLY GLU ALA GLU ASP THR	SER VAL THR PRO GLU ASP THR	ASN ASN LEU ARG VAL GLX ASX THR ALA
	88 89 90 91 92 93 94 95	TYR(.98) Tyr(.95) Cys	ALA LEU TYR TYR CYS ALA ARG LEU	ALA VAL TYR TYR CYS ALA LYS	ALA LEU TYR TYR CYS ALA ARG ASP ALA	ALA VAL TYR TYR CYS ALA LYS GLY LYS	ALA VAL TYR TYR CYS ALA ARG PHE ARG				ALA VAL TYR TYR CYS ALA ARG	ALA VAL TYR TYR CYS ALA ARG VAL THR	ALA VAL TYR TYR CYS ALA ARG SER PRO			TYR TYR CYS ALA ARG ASP ALA	ALA VAL TYR TYR CYS ALA ARG HIS ILE	ALA VAL TYR TYR CYS ALA ARG		ALA VAL TYR TYR CYS ALA ARG			ALA VAL TYA TYA CYS ALA ARG THA ARG	ALA VAL TYR TYR CYS ALA ARG VAL VAL	VAL TYA TYA CYS ALA ARG ASX ARG
CDR	96 97 98 99 100 100A 100B 100C 100D		SER VAL THR ALA VAL ALA PHE		GLY PRO TYR VAL SER PRO THR PHE	VAL SER ALA TYR TYR PHE	GLN PRO PHE VAL GLN					PRO ALA ALA SER LEU THR	VAL SER LEU VAL ASP GLY TRP LEU TYR			GLY PRO TYR VAL SER PRO THR PHE	TYR VAL' THR LEU TYR TYR TYR						PRO GLY GLY TYR PHE SER	YAL SEA THA SEA	LEU GLY PRO THR ALA CYS SER VAL
3	100E 100F 100G 100H 100J 100J 100K 101		 		I I I I I HE ALA	ASX TYR	PHE PHE ASP VAL					PHE SER ALA VAL	TYA TYA GLY CLY SER			PHE ALA HIS	PHE MET ALA VAL						ASP VAL	MET ASP VAL	
F 17 4	102 103 104 105 106 107 108 109 110	GLY(.97) GLY(.97) VAL(.97)	TRP GLY GLN GLY THR LYS VAL SER		TYA GLY GLN GLN THA LEU VAL THA	TRP GLY GLX GLY THR LEU VAL THR	PHE GLY GLN GLY THR LEU VAL THR					TRP GLY GLN GLY THR LEU VAL THR	TRP GLY GLN GLY THR LEU			TAP GLY GLN GLY THR LEU VAL THR	TRP GLY GLN GLY THR LEU					• .	GLY GLY GLN GLY THR LEU VAL SEA	TAP GLY GLN GLN THR PRO VAL THR	
	111 112 113	SER				VAL SEA SER					 	343													-

GP 1806



ĩ,

PATENT DOCKET NO. 709

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Paul J. Carter et al.

Serial No. 07/715,272

Filed: 14 June 1991

Immunoglobulin Variants For:

Group Art Unit: 1806

Examiner: L. Feisee

RECEIVED NOV 0 5 1993 GROUP 1800

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-1896

NOTICE OF APPEAL

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

w020.u

Applicant hereby appeals to the Board of Appeals and Interferences from the decision dated May 19, 1993, of the Primary Examiner finally rejecting claims 1-11 and 17-21.

The Commissioner is hereby authorized to charge Deposit Account No. 07-0630 in the amount of \$270 to cover the fees for this appeal and to charge the deposit account for any further fees in regard to this patent application. A duplicate copy of this Notice is enclosed for this purpose.

> Respectfully submitted, GENENTECH, INC.

Janet E. Hasok

119

Janet E. Hasak Reg. No. 28,616

Dated: October 15, 1993

CERTIFICATE OF MAILING (37 CFR 1.8a) I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231. ms Date: October 15, 1993 ILA 0 Louise Strasbaugh RP14167 11/04/93 07715272 07-0630 140 270.00CH

1806

PATENT DOCKET 709



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

}

)

)

In re Application of Paul J. Carter et al. Serial No. 07/715,272 Filed: 14 June 1991 For: Immunoglobulin Variants Group Art Unit: 1806

Examiner: L. Feisee 🖁

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-1896

PETITION AND FEE FOR EXTENSION OF TIME (37 CFR 1.136(a))

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Applicant petitions the Commissioner of Patents and Trademarks to extend the time for response to the Office action dated May 19, 1993 for an additional month, from September 19, 1993 to October 19, 1993. The extended time for response does not exceed the statutory period.

Please charge Deposit Account Number 07-0630 in the amount of \$250 to cover the cost of the second month extension fee less the first month extension fee paid in relation to the request for a one month extension of time filed on September 20, 1993. Any deficiency or overpayment should be charged or credited to this deposit account. <u>A duplicate of this sheet is enclosed.</u>

Respectfully submitted,

GENENTECH, INC.

Janet E. Hasak Reg. No. 28,616

Date: October 15, 1993

RP14166 11/04/93

07-0630 140 116

250.00CH

CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and

345

Trademarks, Washington, D.C. 20231 tras LS. Strasbaugh Louise

Date: October 15, 1993

07715272

w021.u

	· · · · · · · · · · · · · · · · · · ·
,	
	•
、 、	
· ·	
	UNITED STATES DEPARTMENT OF COMMERCE Patent and Tredomark Office Address: COMMISSIONER OF PATENTS AND TRADEMARKS
· · ·	Washington, D.C. 20231
	SERIAL NUMBER HILING DATE HIRST WANKED AFTER DATE
	<u></u>
	12M2/1021 GENENTECH, INC. ATTN: COROLYN R. ADLER
	ALTINE CLINE IN INCLUSION 460 POINT SAN BRAND BLVD. SOUTH SAN FRANCISCO, CO 94050 DATE MALEO:
	Below is a communication from the EXAMINER in charge of this application
- ,	Below is a communication from the EXAMPLEA in Given of the EXAMPLEA IN GIVEN OF PATENTS AND TRADEMARKS
	ADVISORY ACTION
	THE PERIOD FOR RESPONSE:
	b) explore three months from the date of the final rejection or as of the mailing date of this Advisory Action, whichever is tater. In no b) explore three months from the date of the final rejection or as of the mailing date of the linal rejection.
	Any extension of time must be obtained by filing a polition under 37 CFR 1.135(a), the proposed response and the appropriate fae. The date on which the response, the pebilon, and the fee have been filed is the date of the response and also the date for the purposes of detarmining the period of extension and the corresponding amount of the fee. Any extension fee pursuent to 37 CFR 1.17 will be calculated from the date of the originally set shortened statutory period for response or as set forth in b) above.
	Appellant's Brief is due in accordance with 37 CFR 1, $192(6)$. Applicant's response to the final rejection, filed 2725 has been considered with the following effect, but it is not deemed to place the application in condition for allowance:
	The emported emperiments to the claim and /or specification will not be entered and are initial rejuction with the claim and /or specification will not be entered and are initial rejuction.
·	a. There is no convinding showing under 37 CFR 1.116(b) why the proposed amendment is necessary and was not center presented.
	 A They raise new issues that would require hurder consideration and/or search. (See Note). A They raise the issue of new matter. (See Note).
	d. A Thoy are not deemed to place the application to better form for appeal by materially reducing or simplifying the tesues for noosal.
	e. They present additional claims without cancelling a corresponding number of finally rejected claims.
	NOTE: John Ras not been defined with respect to the Datrable domain of a conservent and body
	Newly proposed or amended claims would be allowed II submitted in a separately filed amendment cancelling the non-allowable claims.
	3. If Upon the filling an appeal, the proposed amendment in will be entered and the status of the claims will found the status of the status of the claims will found the status of the status of the claims will found the status of the status o
	Claima allowed: 12 and 12 SUPERVISORY PATENT EXAMINER: Claima objected to: 10717-21 GROUP 180 19/11/92
:	claims rejected: / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / / /
	A the set of the set o
: <u>-</u> - به المراجع	introduction of the carbonal alling on Main it is not clear
	 The affidavit or exhibit will not be considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and content of a considered because applicant has not anothing out and construct anothing out and construct anothing out anothing out and construct anothing out anothing
	- The proposed drawing correction his has not been approved by the examinar.
346	that the consensus antibody to the application wonder be denticed to the antibody of the Queen at 2 the declars be denticed to an add with other of that the talked



For:

Sir:

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

RECEIVED

DEC 2 9 1993

In re Application of

Paul J. Carter et al.

Serial No. 07/715272

Filed: June 14, 1991

Immunoglobulin Variants GROUP 1863

Group Art Unit: 1806

Examiner: L. FEISEE

PATENT DOCKET 709

460 Point San Bruno Boulevard South San Francisco, CA 94080 (415) 225-1896

AMENDMENT PURSUANT TO 37 CFR § 1.116(a)

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Pursuant to 37 CFR § 1.116(a), please cancel claims 1-11, 17, and 19-21 of the above application. Claim 18 was canceled in the Amendment filed September 20, 1993. Applicants trust that the above-mentioned application with allowed claims 12 and 13 will be in condition for allowance following the entry of this amendment and look forward to receiving the Notice to this effect.

> Respectfully submitted, GENENTECH, INC.

ner E. Hasah

Dated: December 13, 1993

Janet E. Hasak Reg. No. 28,616

CERTIFICATE OF MAILING

I hereby cortify that this correspondence is being deposited with the United States Postal Service in first class envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, on the date shown below.

Dated: 13 DEC 1993

Has buise Strasbaugh



UNITED STATES DEPARTMENT OF COMMERCE Patent and Trademark Office

Address: COMMISSIONER OF PATENTS AND TRADEMARKS Washington, D.C. 20231

	FIRST NAME	APPLICANT	TAT	TORNEY DOCKET NO.
SERIAL NUMBER FILING DATE				
		n [EX	AMINER
		Г	ART UNIT	PAPER NUMBER
		L		25
	· .			
			ATE MAILED	ι,
E	XAMINER INTERVIEW SUM	IMARY RECOR	D	
Il participants (applicant, applicant's representat	tive, PTO personnel):		,	
1. Jan du. Lap				
1 Materse	(4)			
1/1/98	. قى		· ,	
ate of interview				
ype: Pelephonic D Personal (copy. is.gi	iven to 🔲 applicant 🗌 applicar	t's representative).		
xhibit shown or demonstration conducted:	No. If yes, brief descri	ption:	•	
xhibit shown or demonstration conducted.	у на со со со со со со с а со			•
		١		
		was not reache	ed.	
Agreement 🛛 was reached with respect to som	e or all of the claims in question.			
Claims discussed:			· · · · · · · · · · · · · · · · · · ·	
dentification of prior art discussed:	•			
		H or any other com	ments:	
Description of the general nature of what was ag	reed to if an agreement was reached		than	aux.
Finality	1 would K	R WI	<i>FV</i> [00 c.	
A.I.A.	11. pomine			
ATCADA DI	maing			
•	9			
(A fuller description, if necessary, and a copy attached. Also, where no copy of the amendme				
Unless the paragraphs below have been checke NOT WAIVED AND MUST INCLUDE THE S last Office action has already been filed, then a	d to indicate to the contrary, A F	ORMAL WRITTEN	N RESPONSE TO T	HE LAST OFFICE ACTION
It is not necessary for applicant to provi	de a separate record of the substanc	e of the interview.		
Since the examiner's interview summary	v above (including any attachment	s) reflects a comple	ete response to each	of the objections, rejections and form is considered to fulfill t

Since the examiner's interview summary above (including any attachments) reflects a complete response to each or the objections, rejections and requirements that may be present in the last Office action, and since the claims are now allowable, this completed form is considered to fulfill th response requirements of the last Office action.

PTOL-413 (REV. 1-94)

Examiner's Signature

Patent and Tridemark Office Address: COMINSESTVER OF PATENTS AND TRADEN BERIAL NUMBER FILING DATE FIRST NAMED INVENTOR Address: 07/715.272 06/14/91 CARTER P 709 EXAMINER FEISEE, L	• •				ADEDADT	
07/715.272 06/14/91 CARTER P 709 EXAMPLE 18M2/0203 ATUNIT PARENT GENENTECH. INC. ATUNIT PARENT ACO POINT SAN BRUNO BLUD. 26 300 1806 SOUTH SAN FRANCISCO, CA 94080 1806 02/03/94 This ba a communication from the asaminer in charge of your application. 02/03/94 02/03/94 COMMISSIONER OF PATENTS AND TRADEMARKS 1 02/03/94 02/03/94 A shortered statutory particle for response to fall action is set to explice	~~~~~			Potent and 1 Address: COMM	Tedemark DI	FICE
ISM2/0203 EXAMINER GENENTECH. INC. ATTUNIT ATTN: CAROLYN R. ADLER 26 460 POINT GAN BRUND BLVD. COMMISSIONER OF PARCISCO, CA 94080 1806 SOUTH GAN FRANCISCO, CA 94080 1806 This is a communication from the avaniner in charge of your application. COMMISSIONER OF PATENTS AND TRADEMARKS 02/03/94 This a positization has been examined Responsive to communication filed on 12/17/13 This action in A thorae of Roferences Clied by Examiner, PTO-982. 1 days from the date of the information on how to Effect Drawing Obarges, PTO-1474. action and patients information and how to Effect Drawing Obarges, PTO-1474. biolise of Informal Patent Application, PTO-149. Part INE FOLLOWING ATTACHMENT(S) ARE PART OF THIS ACTION: i action of References Clied by Examiner, PTO-982. 2 biolise of Informal Patent Application, PTO-149. Importantion on How to Effect Drawing Obarges, PTO-1474. E biolise of Informal Patent Application, PTO-149. action patients in action of moving in the date of the above, claima Claims 1 12 Other above, claima are withdrawn from concerned Standard OF Action 12 12 Action and patient Application, PTO-149. are regioted. Claims 12<	BERIAL NUMBER	FILING DATE	FIRST NAMED		·····	ATTORNEY DOCKE
16M2/0203 ARTUNIT PAPER NL GENENTECH. INC. ATTN: CAROLYN R. ADLER 26 460 POINT BAN BRUND BLVD. SOUTH BAN FRANCISCO, CA 94080 1806 SOUTH BAN FRANCISCO, CA 94080 1806 02/03/94 This is a communication from the examinor in charge of your application. .02/03/94 .02/03/94 COMMISSIONER OF PATENTS AND TRADEMARKS 1 .02/03/94 A is not need statutory partical for memory and the state of the application. .02/03/94 .02/03/94 A is not need statutory partical for response will cause the application. .02/03/94 .02/03/94 A is not need statutory partical for response will cause the application. .02/03/94 .02/03/94 A is not need statutory partical for response will cause the application. .02/03/94 .02/03/94 A is not need statutory partical for response will cause the application. .02/03/94 .02/03/94 A is not need statutory partical for response will cause the application. .03/04/04/04 .03/04/04/04 .03/04/04/04 .03/04/04/04 .02/04/04/04 A is not need statutory partical for response will cause the application. .0 .0 .0 .0 .0 .0 .0 .0 .0 .0 <td>07/715.272</td> <td>06/14/91</td> <td>CARTER</td> <td></td> <td>FEISEE,</td> <td>EXAMINER</td>	07/715.272	06/14/91	CARTER		FEISEE,	EXAMINER
SOUTH SAN FRANCISCO, CA 94050 1806 DATE MALLED: 02/03/94 COMMISSIONER OF PATENTS AND TRADEMARKS 0 Image: Communication has been examined Image: Communication float on the examined of the sequence of the s			18M2/0203			
This is a communication from the examined in charge of your application. COMMISSIONER OF PATENTS AND TRADEMARKS Image: Commission of the patent of the action is a communication filed on						
A shortened statutory period for response to this action is set to expire	This is a communication COMMISSIONER OF P.	I from the examiner in charg ATENTS AND TRADEMAR	ge of your application. IKS I		'n	02/03/94
A shortened statutory period for response to this action is set to expire			1	ution Moder	12/17/93	
Part I THE FOLLOWING ATTACHMENT(5) ARE PART OF THIS AGTION: 1. Motice of References Cited by Examiner, PTO-892. 2. Motice of Draftsman's Patent Drawing Review 3. Notice of Art Cited by Applicant, PTO-1449. 2. Motice of Informal on the to Effect Drawing Changes, PTO-1474. 2. Motice of Informal Patent Application, PTO-1 Part II SUMMARY OF ACTION 1. Claims 1. 1. Claims 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	A shortened statutory pe	ziod for response to this act	tion is set to expire	3_month(s),	days	from the date of this lotte
1. Motice of References Ciled by Examiner, PTO-892. 2. Holice of Draftsman's Petent Drawing Review 3. Information on How to Effect Drawing Changes, PTO-1474. 4. Notice of Informal Patent Application, PTO-1 8. Information on How to Effect Drawing Changes, PTO-1474. 6. Notice of Informal Patent Application, PTO-1 9. Information on How to Effect Drawing Changes, PTO-1474. 6. Notice of Informal Patent Application, PTO-1 9. Claims 1. D. DM. 13 are pending in the claims 1. Claims 1. I.		· ·			1001 00 0.0.0. 10.	3
1. D DMD 13 are pending in the DI the above, claims 01 the above, claims are withdrawn from corrections, in the above, claims 2. D Claims 1 - 11, 14 - 21 1. D Claims are withdrawn from corrections, in the above, claims 2. D Claims 1 - 11, 14 - 21 1. D Claims are rejected. 3. D Claims 1 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -	3. Notice of Art	Cited by Applicant, PTO-14	49.			
Of the above, claims are withdrawn from corrections 2. [Claims 1 - 11, 14 - 21 have been cancell have been cancell 3. [Claims are rejected. 6. [Claims /2 - 3/21/3 are objected to. are objected to. 6. [Claims are objected to. 7. [Claims are objected to. 8. [Claims are objected to. 9. [Claims application has been filed with informal drawings under 37 C.F.R. 1.85 which are acceptable for examination purposes. 8. [] Formal 'drawings have been received on	Part II SUMMARY OF	ACTION	1' A			
2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	1. Claims	12	2 ond 13	,		
2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	Of the abc	ove, ciaime		•		re withdrawn from consid
9. Claims	2. Di Ciaima	1-11,10	1. 1.1.1			
Claims	,					
8. Claims		12 200	113			
G. Claimsare subject to restriction or election require G. Claimsare subject to restriction or election require G. Claimsare subject to restriction or election require G. This application has been filed with informal drawings under 37 C.F.R. 1.85 which are acceptable for examination purposes. G. Formal drawings are required in response to this Office action. G. The corrected or substitute drawings have been received on Under 37 C.F.R. 1.84 these drawings are is acceptable; Cont acceptable (eee explanation or Notice of Drattsman's Patent Drawing Review, PTO-948). The proposed additional or substitute sheel(s) of drawings, filed on has (have) been Capproved by the examiner; Caseptable; Correction, filed, has been Capproved; Caseptaved by the examiner; Caseptaved by the examiner; where explanation). The proposed drawing correction, filed, has been Capproved; Caseptaved (eee explanation). Acknowledgement is made of the claim for priority under 35 U.S.C. 119. The certified copy has Deen received Care Deen Deen Direction, earch to the market or consolution as to the market to closed accordance with the practice under Ex pane Casyle, 1935 C.D. 11; 453 O.G. 213.						
 7. This application has been filed with informal drawings under 37 C.F.R. 1.85 which are acceptable for examination purposes. 8. Formal drawings are required in response to this Office acton. 9. The corrected or substitute drawings have been received on Under 37 C.F.R. 1.84 these drawing have been received on Under 37 C.F.R. 1.84 these drawing have been received on Under 37 C.F.R. 1.84 these drawing have been received on Under 37 C.F.R. 1.84 these drawing have been received on Under 37 C.F.R. 1.84 these drawing have been received on Under 37 C.F.R. 1.84 these drawing field on In a corrected or substitute sheel(s) of drawings, filed on has (have) been has proved by the examiner; desproved by the examiner (see explanation). 11. The proposed drawing correction, filed, has been has been approved; desproved (see explanation). 12. Acknowledgement is made of the claim for priority under 35 U.S.C. 110. The certified copy has been received, field on 13. Since this application appears to be in condition for allowance except for formal matters, prosocution as to the merits is closed accordance with the practice under Ex parte Quayle, 1935 C.D. 11; 453 O.G. 213. 						
8. Formal drawings are required in response to this Office action. 9. The corrected or autostitute drawings have been received on Under 37 C.F.R. 1.84 these drawing are acceptable; not acceptable (see explanation or Notice of Draftsman's Patent Drawing Review, PTD-948). 10. The proposed additional or substitute sheel(s) of drawings, filed on has (have) been approved by the examiner; disapproved by the examiner; (see explanation). 11. The proposed drawing correction, filed, has been approved; disapproved (see explanation). 12. Acknowledgement is made of the claim for priority under 35 U.S.C. 119. The certified copy has been received not been been filed in parent application, serial no; filed on; filed on 13. Since this application appears to be in condition for allowance except for formal matters, presocution as to the ments to closed accordance with the practice under Ex parte Quayle, 1935 C.D. 11; 453 O.G. 213.						
 9. The corrected or substitute drawings have been received on Under 37 C.F.R. 1.64 these dramatic are acceptable; D not acceptable (see explanation or Notice of Drattsman's Patent Drawing Review, PTO-948). 10. The proposed additional or substitute sheel(s) of drawings, filed on has (have) been Deproved by the examiner; Deserving correction, filed has been Deproved; Disperved (see explanation). 11. The proposed drawing correction, filed has been Deproved; Disperved (see explanation). 12. Acknowledgement is made of the claim to priority under 35 U.S.C. 119. The certified coop has D been received D not bee D been filed in parent application, serial no; filed on 13. Since this application appears to be in condition for allowance except for formal matters, prosecution as to the merits to closed accordance with the practice under Ex parte Quayle, 1935 C.D. 11; 453 O.G. 213. 	,					
are acceptable; and acceptable (eee explanation or Notice of Drattsman's Patent Drawing Review, PTD-948), 10. The proposed additional or substitute sheel(s) of drawings, filed on, has (have) been approved by the examiner; deseptorved by the examiner; deseptorved by the examiner; deseptorved by the examiner; deseptorved (eee explanation). 11. The proposed drawing correction, filed, has been approved; disapproved (eee explanation). 12. Acknowledgement is made of the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim tor priority under 35 U.S.C. 119. The certified copy has a been received and the daim tor priority under 35 U.S.C. 119. The certified copy has a been received and the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim tor priority under 35 U.S.C. 119. The certified copy has a been received and the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim for priority under 35 U.S.C. 119. The certified copy has a been received and the daim for allowance except for formal matters, presecution as to the merits to closed accordance with the practice under Ex parte Queyle, 1935 C.D. 11; 453 O.G. 213.	· · ·				. Under 37	C.F.R. 1.84 these drawlr
examiner; desapproved by the examiner (see explanation). 11. The proposed drawing correction, filed, has beenapproved;disapproved (see explanation). 12. Acknowledgement is made of the daim for priority under 35 U.S.C. 119. The certified copy has been received not bee; fied on; field on	are 🔲 acceptab	ile; 🗋 not acceptable (see r	explanation or Notice of I	Drațtsman's Paten	t Drawing Review,	PTO-948),
 12. ☐ Acknowledgement is made of the claim for priority under 35 U.S.C. 119. The certified copy has ☐ been received ☐ not been been filed in parent application, serial no				۰.	, has (have) been	approved by the
 Deen filed in parent application, serial no; filed on; 13. Since this application appears to be in condition for allowance except for formal matters, prosecution as to the marks is closed accordance with the practice under Ex parte Quayle, 1935 C.D. 11; 453 O.G. 213. 					red; 🖸 disepprove	d (ass explanation).
accordance with the practice under Ex parte Quayle, 1935 C.D. 11; 453 O.G. 213.	· 🖸 been filed in p	azent application, serial no.	· <u></u> ;	filed on	copy has Dibeen	received D not been re
14. 💭 Other					ra, prosecution as	to the ments to closed in
					•	
		, .				

PTOL-328 (Pev. 293)

.)

10 B G

Serial No. 715272

Art Unit 1806

The finality of the previous Office action is withdrawn in view of the following new grounds of rejection.

Claims 12 and 13 are pending in this application, and these claims are humanized light and heavy chain variable regions of a 5 previously referenced antibody 4D5.

Claims 12 and 13 are rejected under 35 U.S.C. § 103 as being unpatentable over Hudziak et. al. or Fendly et. al. in view of Queen et. al.

Hudziak et. al. and Fendly et. al. both teach the production and characterization of the 4D5 antibody (see Hudziak et. al. 1166-1167 and Fendly et. al. pages 1553-1554). Hudziak et. al. suggests the possible therapeutic role of the 4D5 antibody in human neoplasias which overexpress p185-HER2 (pages 1171, last paragraph) while Fendly et. al. disclose the possible use of anti-p185 HER2 antibodies for <u>in vivo</u> radioimaging for detection of relevant primary tumors. They do not describe the production of these antibodies in the humanized form.

Queen et. al. teach the production of antibodies against 1L-2 receptor in the humanized form, using computer modeling in order to modification of certain framework regions in determine the 20 conjunction with CDR grafting. The antibodies produced are than to be used for in vivo administration to human patients, either for diagnosis or therapy. It is known in the art that murine and even chimeric antibodies have characteristics which may severely limit 25 their use in human therapy. As foreign proteins, murine and chimeric antibodies may elicit immune reactions that reduce or destroy their therapeutic efficacy and/or evoke allergic. or The probable need for hypersensitivity reactions in patients. readministration of such therapeutic modalities in neoplastic

disorders increases these risks. The result would be tissue injury by virtue of antigen-antibody deposition.

facie obvious to one of ordinary It would have been prima skill in the art at the time the invention was made to make humanized antibodies having the sequences of the 4D5 antibody. The methods of Queen et. al. were clear and self explanatory, and resulted in a high affinity antibody. One of ordinary skill in the art would have been motivated to humanize the 4D5 antibody in light diagnostic applicability. therapeutic and its potential of specific amino acid claims are drawn to Although the sequences, it is maintained that the differences in amino acid sequence which would have been obtained using the method of Queen al. would not have been patentably distinct from the claimed et. amino acid sequences. Absent sufficient factual evidence to the 15 contrary the claims are obvious over the cited prior art.

35 U.S.C. § 101 reads as follows:

Whoever invents or discovers any new and useful process, machine, manufacture, or composition matter or any new and useful improvement of thereof, may obtain a patent therefore, subject conditions and requirements of this to the title.

Claims 12 and 13 are rejected under 35 U.S.C. § 101 because the claimed invention lacks patentable utility. These claims are drawn to a light chain variable region polypeptide and a heavy 25 chain variable region polypeptide which in and of themselves have no patentable utility. The specification does not disclose any

з 351

10

practical utility for the individual polypeptides and does not present evidence that these polypeptides are capable of binding in any particular manner when not in association with each other.

Claims 12 and 13 are directed to an invention not patentably 5 distinct from claims 1, 3-9, and 40 of commonly assigned 07/977,453.

Specifically, the claims of the instant invention are drawn to the humanized version of the 4D5 antibody which is disclosed in copending application.

Commonly assigned 07/977,453, discussed above, would form the 10 basis for a rejection of the noted claims under 35 U.S.C. § 103 if the commonly assigned case qualifies as prior art under 35 U.S.C. § 102(f) or (g) and the conflicting inventions were not commonly owned at the time the invention in this application was made. In 15 order for the examiner to resolve this issue, the assignee is required under 37 C.F.R. 1.78(c) and 35 U.S.C. 5 132 to either show that the conflicting inventions were commonly owned at the time the invention in this application was made or to name the prior inventor of the conflicting subject matter. Failure to comply with this requirement will result in a holding of abandonment of the 20 A showing that the inventions were commonly owned at application. the time the invention in this application was made will preclude a rejection under 35 U.S.C. § 103 based upon the commonly assigned case as a reference under 35 U.S.C. § 102(f) or (g).

⁴ 352

Claims 12 and 13 are provisionally rejected under the judicially created doctrine of obviousness-type double patenting as being unpatentable over claims 1,3-9, and 40 of copending application Serial No. 07/977,453 in view of Queen et. al.. The instant claims are drawn to the heavy chain and light chain 5 the 4D5 antibody. Copending application variable regions of 07/977,453 claims an antibody with the same characteristics as 4D5, and also states within the claims that 4D5 antibody was useful for diagnosis and therapy of tumors expressing the p185 HER2 antigen on their surface. The induction of HAMA responses upon repeated 10 administration of rodent antibodies has led to the desirability of producing antibodies which are even more "near human" than chimeric antibodies. Queen et. al. describes the production of antibodies which contain essentially the CDR of rodents and are grafted into These antibodies are also mutated in human framework regions. 15 certain framework residues in order to produce functional and high affinity molecules. The procedure in Queen et. al. clearly teaches the particular framework residues that need to be changed in order to yield high affinity antibodies, and they teach how to determine the appropriate residues using computer modeling programs. This 20 protocol is adaptable to any number of antibodies. Therefore, not production of non-immunogenic 4D5 antibodies was the only desirable, but the procedure for producing the antibodies was also well known and practiced. It would have been prima facie obvious

15

to one of ordinary skill in the art at the time the invention was made to use the claims of the copending application in combination with the reference of Queen et. al. in order to obtain high affinity functional humanized antibodies.

patenting rejection is The obviousness-type double 5 judicially established doctrine based upon public policy and is primarily intended to prevent prolongation of the patent term by prohibiting claims in a second patent not patentably distinct from claims in a first patent. In re Vogel, 164 U.S.P.Q. 619 (CCPA A timely filed terminal disclaimer in compliance with 37 1970). 10 C.F.R. 1.321(b) would overcome an actual or provisional rejection on this ground provided the conflicting application or patent is shown to be commonly owned with this application. See 37 C.F.R. 1.78(d).

Claims 12 and 13 are provisionally rejected under 35 U.S.C. § 103 as being obvious over copending application Serial No. 07/977,453 in view of Queen et. al.

The instant claims are drawn to the heavy chain and light chain variable regions of the 4D5 antibody. Copending application 20 07/977,453 discloses an antibody with the same characteristics as 4D5, and also discloses that 4D5 antibody is useful for diagnosis and therapy of tumors expressing the p185 HER2 antigen on their surface. The induction of HAMA responses upon repeated administration of rodent antibodies has led to the desirability of

⁶ 354

5

producing antibodies which are even more "near human" than chimeric Queen et. al. describes the production of antibodies antibodies. which contain essentially the CDR of rodents and are grafted into These antibodies are also mutated in human framework regions. certain framework residues in order to produce functional and high affinity molecules. The procedure in Queen et. al. clearly teaches the particular framework residues that need to be changed in order to yield high affinity antibodies, and they teach how to determine the appropriate residues using computer modeling programs. This protocol is adaptable to any number of antibodies. Therefore, not 10 4D5 antibodies the production of non-immunogenic only was desinable, but the procedure for producing the antibodies was also well known and practiced. It would have been prima facie obvious to one of ordinary skill in the art at the time the invention was made to use the claims of the copending application in combination 15 with the reference of Queen et. al. in order to obtain high affinity functional humanized antibodies.

Copending application Serial No. 07/977,453 has a common assignee with the instant application. Based upon the earlier effective U.S. filing date of the copending application, it would 20 constitute prior art under 35 U.S.C. § 102(e) if patented. This provisional rejection under 35 U.S.C. § 103 is based upon a presumption of future patenting of the conflicting application.

This provisional rejection might be overcome either by a showing under 37 C.F.R. 1.132 that any unclaimed invention disclosed in the copending application was derived from the inventor of this application and is thus not the invention "by 5 another", or by a showing of a date of invention prior to the effective U.S. filling date of the copending application.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Lila Feisee whose telephone number is (703) 308-2731.

Any inquiry of a general nature or relating to the status of this application should be directed to the Group receptionist whose telephone number is (703) 308-0196.

PRIMARY EXAMINER **GROUP 1800**

Feisee/lf January 11, 1994

10

15

TO SEPARATE. D TOP AND BOTTOM EDGES, SNAP-APART AN DISCARD CARBON ROUP ART UNIT ATTACHMENT TO PAPER NUMBER U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE FORM PTO-892 (REV. 2-92) SERIAL N H 01 15 NOTICE OF REFERENCES CITED al et **U.S. PATENT DOCUMENTS** FILING DATE IF SUB-CLASS CLASS DOCUMENT NO. DATE NAME С D F G н FOREIGN PATENT DOCUMENTS PERTINENT SHTS. PP. DWG SPEC. SUÐ-CLASS NAME CLASS DOCUMENT NO. DATE COUNTRY L м N 0 P a OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.) 86: PAJAS 10029 2 Dille n 10-138 ancir an th this office action. on 707.05 (a).) 7. . · A come of this sheard ăă. 357

		UNITED STATES DEPARTMEN Patent end Trademark Office Address: COMMISSIONER OF I Washington, D.C. 202	PATENTS AND THADEMARKS	
SERIAL NUMBER FILI	ING DATE FIRS	T NAMED APPLICANT	ATTORNEY DOCKET-NO.	
07/715,27	2 06/14/91			709
		18M2/0906 ART UN	FEISEE,	L
460 POINT South San	OLYN R. ADLER SAN BRUNO BLVD FRANCISCO, CA	DATE MAILEC	27 1806	
1. The Notice	e of Appeal filedis:			~~~09/06/94
A. 🗋	Not acceptable for reason(s) that	at:		· · ·
	(1) The Appeal fee required not submitted with the	t by 35 U.S.C. 41 (a)(6) and 37 CF Notice of Appeal.	R 1.17(e) was	· ·
	(2) The submitted fee of \$	is insufficient. The ap	peal fee required by	
	(3) The Notice of Appeal w	as not timely filed.		
	(4) The Appeal fee received	d on was not timely	tiled.	
-	(5) The Appeal is not in a have not been finally	ompliance with 37 CFR 1.191 in or twice rejected.	that the claims	
	(6) A Notice of Allowability	was mailed by the Office on	· ·	
в. 🗖	ONE MONTH from the date of PERIOD OF THE LAST OFFICE NO EXTENSION OF THIS ONE or (b) BUT THE PERIOD FOR R	ted as indicated. Applicant is giv this letter OR the TIME REMAIN ACTION, whichever is longer, to MONTH PERIOD MAY BE GRAN ESPONSE SET IN THE LAST ACT not timely completed, the applic	ING IN THE RESPONSE complete the appeal. TED UNDER 37 CFR 1.136(a) FION MAY POSSIBLY	
	(1) The Notice of Appeal	is not signed.		
	(2) dentification of the app 37 CFR 1.191 (b).	ealed claim or claims is required u	ndər	
2. The Brief fi	led is NOT :	acceptable for the reason(s) indica	ted below.	·
	I in this application will be dismisse . Extensions of time may be obtain	ed unless the applicant makes the E ined under 37 CFR 1.136(a).	Brief	
A. 🛄	The Brief and/or Brief fee is untin	nely. See 37 CFR 1.192.		,
в. 🗖	The requisite fee which must acc See 37 CFR 1.17(!).	ompany the Brief has been omitted		
c.□	The submitted Brief fee of fee required by 37 CFR 1.17(1) is	is not the proper amount.	The Brief	
3 The Appeal	I in this application is DISMISSED	because		
A. 🗌		ired under 37 CFR 1.17(I) was not I the period for obtaining an extens FR 1.136 has expired.		hille
B	The Brief was not filed, or was no of time to file the brief under 37 C	t timely filed and the period for obta FR 1.136 has expired.	• • • • • • • • • • • • • • • • • • • •	L. LACEY
4 As the resul	It of the dismissal in "3" above, this		SUPERVISURY	PATENT EXAMINER
- A. A.	is abandoned since there are no a	llowed claims.	on.	
В. 🗌	is being returned to the examiner to the examiner to the merits. Prosecution on the merits	for disposition since it contains allow is CLOSED.	wed	1199

(PTOL-461, Rev. 4/89)

20. . . .

	REQUEST FOR ACLESS OF ABANDONED APPLIC ON UNDER 37 CFR 1.14
	Create at the All the All
	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
с ¹	Group Art Unt Examiner
: *	Paper No. #28
	Assistant Communicationer for Platents
	Washington, DC 20231
	I have by request access under 37 CFR 1.14(a)(3)(Iv) to the application file record of the above- identified ABANDONED application, which is: (CHECX CNE;
	(A) referred to in United States Patent Number 5821337 column
• -	
	(B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No filed filed on page of
	paper number
	(C) an application that daims the benefit of the filing date of an application that is open to public inspection, i.e., Application No.
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public.
	(D) an application in which the applicant has filed an authonization to lay open the complete
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public.
	 (D) an application in which the applicant has filed an authomization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address:
	 (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: 2001 Jeffenson DHVIS Hwy Anl- Va. 22202
	 (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: 2001 Jeffenson DHVIS Hwy Anl- Va. 22202
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: <u>2001 Jeffenson DHVis Hwy Anl-Va-22202</u> Suite # 806 Brian William 10/19/99
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondences concerning this request to the following address: <u>2001</u> Jefferson DHVis Huy Anl- Va. 22202 <u>50:1e # 806</u>
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: <u>2001</u> Jefferson <u>Physics Huy</u> Anl- Ua. 22202 <u>5uite # 806</u> <u>Brian Willingham</u> <u>Brian Willingham</u> <u>FOR PTO USE ONLY</u>
	(D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: <u>2001</u> Jefferson <u>Physics Huy</u> Apl- Uo. 22202 <u>5uite # 806</u> <u>Brien Willight</u> <u>10 /19 / 99</u> Signature Bright Williphe help

In re Application of Fied Application Number RECEIVED 6-14-91 NOV 3 0 1999 Brammer Group Art Linit File Information Unit Paper No. 1229 Assistant Commissionar for Patents Washington, DC 20231 I hereby request access under 37 CFR 1.14(a)(3)(Iv) to the appointion file record of the above-Identified ABANDONED application, which is: (CHECX ONE) \mathcal{V} (A) referred to in United States Patent Number <u>5821337</u> column (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No. ______ filed ______ on page _____ of paper number (C) an application that claims the benefit of the filing date of an application that is open to public Inspection, I.e., Application No. ______ filed _____ or (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: - 30-99 FOR PTO USE ONLY Typed or primed name N Approved by: (Inmiais) Unit: Burgen Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the entime case. Any comments on the employed to the you are required to complete this form should be sent to the Chief Information Officer. Pr and Tratemate Office. Wearington. DC 20231 DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDREES. SEND TC: Assistant Commensationer for Patence, Wearington. DC 20221

PTO/S8/68 ed for use through 10/31/99. Office: U.S. DEPARTMENT OMB 0651-003 Patent and Trademark Office; to respond to a colecton of information unit A LOSOLAVE Under the Paperwork Reduction Act of 1995, no persons are reduc REQUEST FOR ACCESS OF ABANDONED APPLICATION UNDER 37 CFR 1.14(a) In re Application of Filed Application Number Group Art Unit Examiner 20 Paper No. Assistant Commissioner for Patents Washington, DC 20231 thereby request access under 37 CFR 1.14(a)(3)(iv) to the application file record of the above-Mentified ABANDONED application, which is: (CHECK ONE) (A) referred to in United States Patent Number 6054297 ., column . . (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., _____ filed __ Application No. _ paper number ____ (C) an application that claims the benefit of the filing case of an application that is open to public Inspection, i.e., Application No. (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: Date Signature INE OR FOR PTO USE ONLY Typed or printed name Approved by: _ (initials) Unit: Burden Hour Statement: This form is estimated to taxe 0.2 hours to condicte. The will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer. Patent and Trademark Office. Washington, DC 20231. JO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents. Washington, DC 20231.

ì

PTC/SB/68 (11-66) Approved for use through 10/31/99, OMB 0651-0031 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Under the Pederwork Reduction Act of 1995, no persons are required to respond to a collection of information unities & discisive a weld CMB control number REQUEST FOR ACCESS OF ABANDONED APPLICATION UNDER 37 CFR 1.14(a) In re Application of RECED Application Number Filed SEP 2 8 2001 Group Art Unit Examine Filo Infermation Unit Paper No. Assistant Commissioner for Patents Washington, DC 20231 I hereby request access under 37 CFR 1.14(a)(3)(iv) to the application file record of the aboveidentified ABANDONED application, which is: (CHECK ONE) (A) referred to in United States Patent Number column (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No. ____, filed _____ on page ___ ____ of paper number __ (C) an application that claims the benefit of the filing date of an application that is open to public inspection, i.e., Application No. filed . . 01 (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: Date Signature FOR PTO USE ONLY Typed or printed name Approved by: (initial - inite Burgan Hour Statement: This form is estimated to taxe 0.2 hours to complete. The will vary capencing upon the needs of the individual case. Any comments on the emount of time you are required to complete inits form should be asont to the Chief Information Officer. Patent and Trademark Office, Washington, OC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Weshington, OC 20231.

PTO/SB/68 (04-01) Approved for use through 10/31/2002. OMB 0651-0031 U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number REQUEST FOR ACCESS TO AN APPLICATION UNDER 37 CFR 1.14(e) e Application of hlice RECEIVED Examiner Art Unif MAR 1 1 2002 File Information Unit Paper No Assistant Commissioner for Patents Washington, DC 20231 1. I hereby request access under 37 CFR 1.14(e)(2) to the application file record of the above-identified ABANDONED Application, which is not within the file jacket of a pending Continued Prosecution Application (CPA) (37 CFR 1.53(d)) and is: (CHECK ONE) United States Patent Application Publication No. 5821337, page ____, line____, (A) referred to in: ___, column _____, line _____, or United States Patent Number____ an International Application which was filed on or after November 29, 2000 and which designates the United States, WIPO Pub. No. _____, page _____, line_____ (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11(b) or 1.14(e)(2)(i), i.e., Application No._____, paper No. _____, page _____, line _____ 2. I hereby request access under 37 CFR 1.14(e)(1) to an application in which the applicant has filed an authorization to lay open the complete application to the public. Date FOR PTO USE O hame Approved ned or prin ed (ials) Unit: Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patient and Trademark Office. Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

1

	ved for use through 10/31/2002. ON 3 (U.S. Patent and Transk Officer U.S. Patent OF CO	MMERCE
	are required to respond to a collection of information unless it disclays a valid OME contr TO AN APPLICATION UNDER 37 CFR 1.14(e)	<u>ni number.</u>
	In re Application of	
RECEIVED	Application Number 07/715.272 Jun.14	199
JUN 2 4 2002	Art Unit Examiner	•
File Information Unit	L.	
	Paper No	
Assistant Commissioner for Patents Washington, DC 20231		
ABANDONED Application, which is not Application (CPA) (37 CFR 1.53(d)) an (A) referred to in: United States Patent Application Pu United States Patent Number	R 1.14(e)(2) to the application file record of the above-identified of within the file jacket of a pending Continued Prosecution and is: (CHECK ONE)	¢.
designates the United States, W	VIPO Pub. No, page, line, open to public inspection as set forth in 37 CFR 1.11(b) or	
	, paper No, page, line	
	R 1.14(e)(1) to an application in which the applicant the complete application to the public.	
Chris Luby Signature	<u>6(24102</u> Date	
Chors Riley	FOR PTOUSE ONLY	
Typed or printed name	Approved by: A. Bryan (Mittels)	<i>t</i> .

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS, SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

.

a. 5 20	Jurden Hour Statement: This form is estimated to take 0.2 hours to complete a mount of time you are required to complete this form should be sent 10231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRE	ete. Time will vary depending upon the needs of the individual case. Any comments on to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231.
	Typed or printed name	Approved by:
• •-		FOR PTO USE ONLY
	Bain Willigh	8/21/02
	 I hereby request access under 37 CFR 1.1 has filed an authorization to lay open the c 	4(e)(1) to an application in which the applicant omplete application to the public.
	1.14(e)(2)(i), i.e., Application No	, paper No, page, line
	· · · · · · · · · · · · · · · · · · ·	to public inspection as set forth in 37_CFR 1.11(b) or
		iled on or after November 29, 2000 and which Pub. No, page, line
		ation No, page, line, 1 2 1 3, column, line, or
	(A) referred to in: United States Patent Application Publica	ation No, page, line,
	ABANDONED Application, which is not with Application (CPA) (37 CFR 1.53(d)) and is:	(e)(2) to the application file record of the above-identified in the file jacket of a pending Continued Prosecution (CHECK ONE)
	Assistant Commissioner for Patents Washington, DC 20231	
	File Information Unit	Paper No. #34
	AUG 2 1 2002	Art Uhit Examiner
	RECEIVED	Application Number 511ed 5-14-91
	REQUEST FOR ACCESS TO A	AN APPLICATION UNDER 37 CFR 1.14(e)

.

PTO/S3/66 (04 Approved for use through 10/31/2002, GMS 0651-0 U.S. Patent and Tradamark Office; U.S. DEPARTMENT OF COMMER Under the Reduction Act of 1995, no persons are required to respond to a collection of information unless it disclave a valid OMB control number I
REQUEST FOR ACCESS TO AN APPLICATION UNDER 37 CFR 1.14(e)
Application of RECEIVED Application Number 07/2/5 272 Filed Application of
OCT 1 8 2002 Art Unit de Exeminer feisee File Information Unit
Assistant Commissioner for Patents Washington, DC 20231
I hereby request access under 37 CFR 1.14(e)(2) to the application file record of the above-identified ABANDONED Application, which is not within the file jacket of a pending Continued Prosecution Application (CPA) (37 CFR 1.53(d)) and is: (CHECK ONE)
(A) referred to in:
United States Patent Application Publication No. <u>6407213</u> , page, line,
United States Patent Number, column, line, or
an International Application which was filed on or after November 29, 2000 and which

designates the United States, WIPO Pub. No. _____, page ____, line ____, (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11(b) or

1.14(e)(2)(i), i.e., Application No._____, paper No. _____, page _____, line _

2. I hereby request access under 37 CFR 1.14(e)(1) to an application in which the applicant has filed an authorization to lay open the complete application to the public.

Typed or printed name

Date

FOR PT	O USE ONLY
Approved by:	
Unit:	(Iniüels)

Surden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Tredemark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

PTO/SB/68 (04-01) Approved for use through 10/31/2002. OMB 0651-0031 U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respon collection of information unless it displays a valid OMB control number REQUEST FOR ACCESS TO AN APPLICATION UNDER 37 CFR 1.14(e) In re Application of RECEIVED Application Number Filed NOV 01 2002 Art Unit Éxaminer File Information Unit Paper No. Assistant Commissioner for Patents Washington, DC 20231 I hereby request access under 37 CFR 1.14(e)(2) to the application file record of the above-identified ABANDONED Application, which is not within the file jacket of a pending Continued Prosecution Application (CPA) (37 CFR 1.53(d)) and is: (CHECK ONE) (U/A) referred to in: United States Patent Application Publication No. _____, page _____, line____ United States Patent Number _____, column _____, line _____, or an International Application which was filed on or after November 29, 2000 and which designates the United States, WIPO Pub. No. _____, page _____, line _____ (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11(b) or 1.14(e)(2)(i), i.e., Application No._____, paper No. _____, page _____, line ____ 2. I hereby request access under 37 CFR 1.14(e)(1) to an application in which the applicant has filed an authorization to lay open the complete application to the public. Stonature Date FOR PTO USE ONL Typed or printed name Approved by Unit: Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer. U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

PTO/SB/68 (04-01) Approved for use Ihrough 10/31/2002. OMB 0651-0031 U.S. Palent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

	SS TO AN APPLICATION UNDER 37 CFR 1.14(e)
1.	In re Application of
RECEIVED	Application Number Filed
DEC 1 0 2002	07-715372 06-14-81
DEC 1 8 2002	Art Unit Examiner
File Information Unit	
	Paper No37
Assistant Commissioner for Pate Washington, DC 20231	ents
ABANDONED Application, which Application (CPA) (37 CFR 1.53)	7 CFR 1.14(e)(2) to the application file record of the above-identified h is not within the file jacket of a pending Continued Prosecution (d)) and is: (CHECK ONE)
(A) referred to in:	ling
United States Patent Applica	tion Publication No, page, line
United States Patent Numbe	c_{10} c
an International Application	which was filed on or after November 29, 2000 and which
	Dade line
designates the United Sta	ates, WIPO Pub. No, page, line
(B) referred to in an application	that is open to public inspection as set forth in 37 CFR 1.11(b) or
(B) referred to in an application	that is open to public inspection as set forth in 37 CFR 1.11(b) or
(B) referred to in an application	
(B) referred to in an application	that is open to public inspection as set forth in 37 CFR 1.11(b) or
(B) referred to in an application 1.14(e)(2)(i), i.e., Application	that is open to public inspection as set forth in 37 CFR 1.11(b) or
(B) referred to in an application 1.14(e)(2)(i), i.e., Application	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant
(B) referred to in an application 1.14(e)(2)(i), i.e., Application	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant
 (B) referred to in an application 1.14(e)(2)(i), i.e., Application I hereby request access under has filed an authorization to lay 	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant y open the complete application to the public.
(B) referred to in an application 1.14(e)(2)(i), i.e., Application	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant
 (B) referred to in an application 1.14(e)(2)(i), i.e., Application I hereby request access under has filed an authorization to lay <i>Monit Rhays</i> Signature 	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant y open the complete application to the public. $D_2 - (R \cdot D_2)$ Date D_3
 (B) referred to in an application 1.14(e)(2)(i), i.e., Application I hereby request access under has filed an authorization to lay 	that is open to public inspection as set forth in 37 CFR 1.11(b) or on No, paper No, page, line 37 CFR 1.14(e)(1) to an application in which the applicant y open the complete application to the public.

P

'.r

、 ·



(12) United States Patent

Carter et al.

US 6,407,213 B1 (10) Patent No.:

(45) Date of Patent:

Jun. 18, 2002

(54)	METHOD ANTIBOL	FOR MAKING HUMANIZED	EP EP EP
(75)	Inventors:	Paul J. Carter; Leonard G. Presta, both of San Francisco, CA (US)	EP EP GB
(73)	Assignee:	Genentech, Inc., South San Francisco, CA (US)	W(W(W(W(
(*)	Notice:	Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.	WC WC WC WC
(21)	Appl. No.	08/146,206	w
(22)	PCT Filed	Jun. 15, 1992	W
(86)	PCT No.:	PCT/US92/05126	W W
	§ 371 (c)((2), (4) D		w w w
	Re	lated U.S. Application Data	w W
(63)	Continuatio Jun. 14, 19	on-in-part of application No. 07/715,272, filed on 191, now abandoned.	-, W W W

- Int. Cl.⁷ C07K 16/00 (51)
- U.S. Cl. 530/387.3; 435/69.6; 435/69.7; (52)
- 435/70.21; 435/91; 536/23.53; 424/133.1 Field of Search 435/69.6, 69.7, (58)
- 435/70.21, 91, 172.2, 240.1, 240.27, 252.3, 320.1, 328; 536/23.53; 424/133.1; 530/387.3

References Cited (56)

U.S. PATENT DOCUMENTS

1016567 4	3/1089	Cabilly et al.
4,816,567 A	711000	Urdal et al 530/388.22
4,845,198 A		Ciudi di d
5,132,405 A		
5,225,539 A	7/1993	Winter 530/389.3
5,225,559		Queen et al 530/387.3
5,530,101 A		
5,558,864 A	9/1996	Denaig of an
5,585,089 A	12/1996	Queen et al 424/133.1
	10/1007	Hudziak et al
5,677,171 A	10/1007	Queen et al 530/387.2
5,693,762 A	• 12/1997	
5,714,350 A	2/1998	Co et al 435/69.6
6 777 007 A	6/1998	Hudziak et al 424/130.1
5,772,997 A		Carter et al 530/387.3
5,821,337 A	10/1998	
5,834,598 A	11/1998	Lowman ct al 530/399
5.859.205 A	1/1999	Adair et al 530/387.3
J. 839.200 m	11 1 1 1 1 1 1	• • • • • • • • • • • • • • • • • • • •

FOREIGN PATENT DOCUMENTS

AU	85058/91	3/1992	C07K/15/12
EP	120694	10/1984	
EP	125023 A1	11/1984	
ÉP	0 239 400 *	9/1987	C12N/15/00
EP	323806 A1	7/1989	
EP	328404 A1	8/1989	A61K/39/395
EP	338745 A1	10/1989	
EP	365209 A2	4/1990	
EP	365997 A2	5/1990	
EP	368684	5/1990	
EP	403156 A1	12/1990	
EP	438310 A2	7/1991	
EP	438312 A2	7/1991	
EP	440351 A2		
EP	0 460 167 B1	12/1991	
EP	0 519 596 A1	12/1992	1

EP	0 592 106 A1	4/1994	
EP	0 620 276	10/1994	
EP	682040 Al	11/1995	0.00000000
EP	451216 BI	1/1996	C12P/21/08
EP	432249 B1	9/1996	
GB	2 188941	10/1987	
WO	WO 87/02671	5/1987	
WO	WO 88/09344	12/1988	
wo	WO 89/01783	3/1989	
wo	WO 89/06692	7/1989	
wo	WO 89/09622	10/1989	
wo	WO 90/07861	7/1990	
wo	90/07861	7/1990	C12P/21/00
WO	WO 91/07492	5/1991	
wo	WO 91/07500	5/1991	
wo	WO 91/09966	7/1991	C12P/21/08
wo	WO 91/09968	7/1991	C12P/21/08
wo	WO 91/09967	11/1991	
wo	WO 92/01047	1/1992	
WO	WO 92/04380	3/1992	
WO	WO 92/04381	3/1992	
wo	WO 92/05274	4/1992	
wo	WO 92/11383	7/1992	
WO	WO 92/11018	9/1992	A61K/35/14
wo	WO 92/15683	9/1992	
wo	WO 92/16562	10/1992	
wo	WO 92/22653	12/1992	•
wo	WO 93/02191	2/1993	
wo	94/11509	5/1994	
wo	WO 94/12214	6/1994	

OTHER PUBLICATIONS

dit .

Riechmann et al. [Nature 332:323-327 (1988)]* Queen et al. [Proc. Natl. Acad. Sci. 86:10029-10033 (1989)].*

Roitt [Immunology, published 1985, by Gower Medical

Publishing Ltd. (London, England) p. 5.5].* Tramontano et al. [J. Mol. Biol. 215:175-182 (1990)].*

"Biosym Technologies" in New Products, Chemical Design Automation 3 (Dec. 1988).

"Polygen Corporation" in New Products, Chemical Design Automation 3 (Nov. 1988).

Adair et al., "Humanization of the murine anti-human CD3 monoclonal antibody OKT3" Hum. Antibod. Hybridomas 5:41-47 (1994).

Chothia et al., "Principles of protein-protein recognition" Nature 256:705-708 (1975).

Chothia et al., "Transmission of conformational change in insulin" Nature 302:500-505 (1983).

Corti et al., "Idiotope Determining Regions of a Mouse Monoclonal Antibody and Its Humanized Versions" J. Mol. Biol. 235:53-60 (1994).

(List continued on next page.)

Primary Examiner-Anthony C. Caputa

Assistant Examiner-Minh-Tam Davis (74) Attorney, Agent, or Firm-Wondy M. Lee

ABSTRACT (57)

Variant immunoglobulins, particularly humanized antibody polypeptides are provided, along with methods for their preparation and use. Consensus immunoglobulin sequences and structural models are also provided.

82 Claims, 9 Drawing Sheets

Under the Paperwork Reduction Act of 1995, no persons are required to re	Approved for use through 10/31/2002, OM U.S. Patent and Trademark Office; U.S. DEPARTMENT OF (esoond to a collection of information unless it disolays a valid OMB cor
REQUEST FOR ACCESS TO AN A	PPLICATION UNDER 37 CFR 1.14(e)
	In re Application of
	("De et
	Application Number Filed
	57/715272 June 1
	Art Unit Examiner
	-
	28
	Paper No. 38
Assistant Commissioner for Patents Washington, DC 20231	
1, defining (61), DO 2023 (
1. hereby request access upder 37 CEP 1 14/oV2V	· · ·
	to the application file record of the above-identified
ABANDONED Application, which is not within the f Application (CPA) (37 CFR 1.53(d)) and is: (CHEC	In a cket of a pending Continued Prosecution
(A) referred to in:	· ·
United States Patent Application Publication No	. 6407213. , page, line
United States Patent Number	, line, line, or
an International Application which was find an	, column, line, or
an International Application which was filed on the designators the United Status, WIDO D. 1. 1.	or after November 29, 2000 and which
	o, page, line
(B) referred to in an application that is open to publ	ic inspection as set forth in 37 CFR 1.11(b) or
1.14(e)(2)(i), i.e., Application No	, paper No, page, line
,	
. I hereby request access under 37 CFR 1.14(e)(1) t	0 an application in which the applicant
	o an application in which the applicant application to the public.
 I hereby request access under 37 CFR 1.14(e)(1) t has filed an authorization to lay open the complete 	o an application in which the applicant application to the public.
	o an application in which the applicant application to the public.
	o an application in which the applicant application to the public.
	to an application in which the applicant application to the public.
Unality request decess under of CFR 1.14(9)(1) has filed an authorization to lay open the complete	to an application in which the applicant application to the public. $5/20/03$
	to an application in which the applicant application to the public. $\frac{5\sqrt{20}/03}{Date}$
Unality request decess under of CFR 1.14(9)(1) has filed an authorization to lay open the complete	application to the public. $\frac{52003}{}$
Usletta Poinderte Usletta Poinderte	application to the public. $\frac{52003}{}$
Unality request decess under of CFR 1.14(9)(1) has filed an authorization to lay open the complete	application to the public. <u>5/20/03</u> Date FOR PTO USE ONLY
Usletta Poinderte Usletta Poinderte	application to the public. 5/20/03 Date

: 1 1

	• •
	PTO/S5/63 (07-03) Approved for use inrough 7/31/2002. OM6 0651-0031
Under the Paperwork Reduction Act of 1995, no persons are required to rescon	U.S. Patani and Trademark Officé: U.S. DÉPARTMENT OF COMMERCE d to a collection of information unless it disclays a valid OME control number.
REQUEST FOR ACCESS TO AN ABANDONED	APPLICATION UNDER 37 CFR 1.14
	In re Application of
Bring completed form to: File Information Unit Crystal Plaza Three, Room 1D01 2021 South Clark Place Arlington, VA Telephone: (703) 308-2733 File Information Unit	Application Number 071715272 6-14-91 Paper No. <u>#39</u>
I hereby request access under 37 CFR 1.14(a)(1)(iv) to the appl application, which is identified in, or to which a benefit is claim attachment):	ication file record of the above-identified ABANDCNED red, in the following document (as shown in the
United States Patent Application Publication No.	, page,line
United States Patent Number $(0.039055$ column	n, ilne, or
WIPO Pub. No, page, li	
 purchased from the Office of Public Records upon payment of <u>For published applications that are still pending</u>, a member of the file contents; the pending application as originally filed; or any document in the file of the pending application. <u>For unpublished applications that are still pending</u>: (1) If the <u>benefit of the pending application is claimed</u> under application that has: (a) issued as a U.S. patent, or (b) p patent application publication, or an international patent Article 21(2), a member of the public may obtain a copy the file contents; the pending application as originally filed; or any document in the file of the pending application (2) If the application is <u>incorporated by reference or otherwis</u> registration, a U.S. patent application publication, or an in accordance with PCT Article 21(2), a member of the pub the pending application as originally filed. 	the public may obtain a copy of: 35 U.S.C. 119(e), 120, 121, or 365 in another ublished as a statutory invention registration, a U.S. application publication in accordance with PCT of: <u>the identified</u> in a U.S. patent, a statutory invention international patent application publication in
' Signature	Date
Typed or printed name	FOR PTO USE ONLY
Registration Number, if applicable	Approved by:
$\sum \frac{1}{1} $	Unit: FFT U.
Telephone Number	
This collection of information is required by 37 CFR 1.14. The information is required to	obtain or relain a benefit by the public which is to file (and by the USPTO

This collection of information is required by 37 CFR 1.14. The information is required to obtain or relatin a benefit by the public which is to file (and by the USP to b process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete. Including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Palent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA.

If you need assistance in completing the for 174 all 1-800-PTO-9199 and select option 2.

		<u>ericano de a influentan di Informatian Linese in declave e velta CAIS converte.</u> ED APPLICATION UNDER 37 OFR 1.14
	F	CAREY
Ring completed form to:	REED	Acciloation Murroer : Fires
File Information Unit	- b -	07/715272 00/14/91
Nystal Plaza Three, Room 100 021 South Clark Place	1 NOV 1 9 2003	Office of the
dington, VA	1 . K	440
elephone: (703) 308-2733	File Information Unit	Facer No. 4170
	i	
hereby request access under	-37 CFR 1.14(a)(1)(iv) to the ap	poliostion file record of the above-identified ABANDC) almed. In the following document (as shown in the
pplication, which is identified tachmenti:	I IE, CT 10 Which a Denemble D.e	
		, page,line
United States Patent N	lumber 605429 , col	ilimo, line,or
	,	
WIPO Pub. No	, page	_,
	hout house to Pending	Applications (37 CFP. 1.14):
Related intomation a	analications is not svelleble to	i the public but copies may be available and may be
numbered from the Office of	of Public Records upon payment	t of the appropriate fee (37 CFR 1.19(5)), as tokews:
For published applications	that are still cending, a member	r of the public may obtain a copy of:
the file contents;		. ~
the pending application	n as originally filed; or	
eny document in the fi For unpublished application	le of the pending application.	
(1) lette benefit of the pe	inding application is claimed und	der 36 U.S.C. 119(e), 120, 121, or 365 in another
sanlication that has (a) issued as a U.S. patent, or (b)	Noublished as a statutory invention registration, a U.S.
patent application put	lication, or an international pate	ant application publication in accordance with PCT
Article 21(2), a memb the file contents	er of the public may obtain a cog	5y 0i.
	», plication as originally filed; or	· · · · · · · · · · · · · · · · · · ·
eav document i	in the file of the pending soplication	tion.
(2) If the application is inc	corporated by reference or other	rvise identified in a U.S. patent, a statutory invention
registration, a U.S. pa	Article 21(2), a member of the p	n International patent application publication in public may obtain a conviot
the pending an	plication as originally filed.	
		/
11 Aid	() (d)	11/10/10
	Demanda	
11 Willa F		/ Apata
SKIEDS	A 1	FOR PTO USE ONLY
Signational Signational	nonlal la	
Dislotter B	inclexter	
Signation Signation Julio Han Han Typed or printed n	ame	Ant -
	·	Approved by:
	r. if applicable	Approved by:
	·	Approved by:
	·	Approved by:
Registration Number	r. if applicable	Approved by:

e dan dalar

ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, R. 272 If you need astistance in completing the form call 1-200-PT(7-9199 and select option 2.

U.S. Pa	Approved for use through 7/31/2003, OMS 0881-00 tent and Trademark Office: U.S. DEPARTMENT OF COMMER
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collect	
EQUEST FOR ACCESS TO AN ABANDONED APPLI	
In re A	Application of
	ation Number Filed
notal Plaza Three Boom 1001	7/715272 6-14-91
D21 South Clark Place DFC, 0 4 2003	Paper No. #4.41
alenhone: (703) 308-2733	Paper No. 77-77
File Information Unit	
- $ -$	file record of the above-identified ABANDONE
oplication, which is identified in, or to which a benefit is claimed, in a	ne following document (as shown in the
tachment):	ling
United States Patent Application Publication No.	_, page, ine
United States Patent Number 6467213, column	
WIPO Pub. No, page, line	· · · ·
 For unpublished applications that are still pending: (1) If the benefit of the pending application is claimed under 35 U.S application that has: (a) issued as a U.S. patent, or (b) published patent application publication, or an international patent application Article 21(2), a member of the public may obtain a copy of: the file contents; the pending application as originally filed; or any document in the file of the pending application. (2) If the application is incorporated by reference or otherwise identification, a U.S. patent application, or an international patent application, the pending application publication publication, or an international patent application with PCT Article 21(2), a member of the public may the pending application as originally filed. 	tified in a U.S. patent, a statutory invention in
$M \sim n I I$	12-4-03
Miller Inter	
Milligue from	Date
Signature	·
WAYNE CROTEAN	Date FOR PTO USE ONLY
Signature WAYNE CROTERN Typed or printed name	FOR PTO USE ONLY
Typed or printed name	·
WAYNE CROTEAN	FOR PTO USE ONLY
Typed or printed name	FOR PTO USE ONLY Approved by:

•.

amount of time you require to complete this form and/or suggestions on redecing VA 22313-1450. DO NOT SEND FEES OR COMPLETED FOR Trademark Office, U.S. Department of Commerce, P.O. 6ox 1450. Alexandriz, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FOR ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA. If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

	•			PTC/SE/SE/ST- ugh 7/S1/2003, OME (881-3) EPARTMENT OF COMMER
Under the Pagerwork Reduction Act of 1	996. no cersons are required to	respond to a collection of information	n uniess R discli	evs a valid CMB ponici nomi
REQUEST FOR ACCESS	TO AN ABANDON	ED APPLICATION L	JNDER 3	7 CFR 1.14
		In re Application of		
·	RECEIVED			Filed
	EUEIVED	Application Number		
le Information Unit rystal Plaza Three, Room 1DP1	MAR 0 9 2004	07/715,	LTC	
021 South Clark Place			Paper	No. FA2
elephone: (703) 308-2733 File	e Information Unit			····
nereby request access under 37 oplication, which is identified in, tachmeni):	OF 10 WINCH & Denem 10			
United States Patent Appli	ication Publication No	, paga,	line	
United States Patent Num	ber 6,054,297	, column, line,		or
WIPO Pub. No	page	, line		
WI-O P db. No	· ·	and a second		
Related Information abo Direct access to pending app purchased from the Office of P For published applications tha the file contents; the pending application a any document in the file of	blications is not available ublic Records upon payn at are still bending, a mer as originally filed; or of the pending application	nent of the appropriate fee (3 nber of the public may obtain n.	37 CFR 1.19 n a copy of:	(b)), as follows:
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of <u>For unpublished applications</u> (1) If the <u>benefit of the pending</u> application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the (2) If the application is incon registration, a U.S. paten	blications is not available ublic Records upon payn at are still bending, a mer of the pending application that are still pending: ling application is claimed issued as a U.S. patent, ation, or an international of the public may obtain cation as originally filed; of the file of the pending approved by rejerence or proceed by rejerence or	nent of the appropriate fee (nber of the public may obtain <u>1</u> under 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publicatio a copy of:	20, 121, or 3 ry invention n in accorda . patent, a st pplication pu	(b)), as follows: 55 in another registration, a U.S. note with PCT
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of For unpublished applications. (1) If the benefit of the pend application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the contents is incon registration, a U.S. paten	blications is not available ublic Records upon payn at are still bending, a mer as originally filed; or of the pending application that are still bending: issued as a U.S. patent, ation, or an international of the public may obtain cation as originally filed; of the file of the pending approved the file of the pending approved to the polication publication rit application publication rit application publication	nent of the appropriate fee (nber of the public may obtain <u>1</u> under 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: or otherwise identified in a U.S. or an international patent a	20, 121, or 3 ry invention n in accorda . patent, a st pplication pu	(b)), as follows: 55 in another registration, a U.S. note with PCT
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of For unpublished applications. (1) If the benefit of the pend application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the contents is incon registration, a U.S. paten	blications is not available ublic Records upon payn at are still bending, a mer as originally filed; or of the pending application that are still bending: issued as a U.S. patent, ation, or an international of the public may obtain cation as originally filed; of the file of the pending approved the file of the pending approved to the polication publication rit application publication rit application publication	nent of the appropriate fee (nher of the public may obtain n. <u>1</u> under 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: or otherwise identified in a U.S. , or an international patent a the public may obtain a cop	$\frac{1}{9}$	(b)), as follows: 55 in another registration, a U.S. note with PCT
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of For unpublished applications. (1) If the benefit of the pend application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the contents is incon registration, a U.S. paten	blications is not available ublic Records upon payn at are still bending, a mer as originally filed; or of the pending application that are still bending: issued as a U.S. patent, ation, or an international of the public may obtain cation as originally filed; of the file of the pending approved the file of the pending approved to the polication publication rit application publication rit application publication	nent of the appropriate fee (nher of the public may obtain n. <u>1</u> under 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: or otherwise identified in a U.S. , or an international patent a the public may obtain a cop	37 CFR 1.19 n = copy of: 20, 121, or 3i ry invention in in accordation provided in the interval of the interva	(b)), as follows: 65 in another registration, a U.S. nce with PCT atutory invention iblication in
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of For unpublished applications (1) If the benefit of the pending application that has: (a)) patent application public Article 21(2), a member the file contents; the pending applic any document in the (2) If the application is incorregistration, a U.S. paten accordance with PCT Article application and the pending applic the pending application a	SCCMMAN bilications is not available ublic Records upon payne at are still bending, a mer as originally filed; or of the pending application that are still pending: ling application is claimed issued as a U.S. patent, ation, or an international of the public may obtain cation as originally filed; of the file of the pending approvated by reference or nt application publication ricei 21(2), a member of location as originally filed.	nent of the appropriate fee (nher of the public may obtain n. <u>1</u> under 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: or otherwise identified in a U.S. , or an international patent a the public may obtain a cop	37 CFR 1.19 n = copy of: 20, 121, or 3i ry invention in in accordation provided in the interval of the interva	(b)), as follows: 55 in another registration, a U.S. note with PCT
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application at any document in the file of <u>For unpublished applications</u> (1) If the <u>benefit of the pending</u> application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the (2) If the application is incon registration, a U.S. paten	SERMAN	Lunder 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: of otherwise identified in a U.S. or an international patent a the public may obtain a cop	r = 100 GeV	(b)), as follows: 65 in another registration, a U.S. nce with PCT atutory invention iblication in
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application and any document in the file of For unpublished applications. (1) If the benefit of the pending application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the file contents; the pending applic any document in the secondance with PCT Article 2D Article 2D registration, a U.S. paten accordance with PCT Article 2D the pending applic Gay Way Signature CARY WAS	SCAMAN if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable	Approximate the public out of the appropriate fee (independent of the public may obtain the public may obtain the public as a statutor publication a copy of: Different application publication a copy of: Different application publication a U.S. (or an international patent a copy of: Different application publication a the public may obtain a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a Copy of: Different application a U.S. (or an international patent a copy of: Different application a Copy of: Different applicati	37 CFR 1.19 n = copy of: 20, 121, or 3i ry invention in in accordation provided in the interval of the interva	(b)), as follows: 65 in another registration, a U.S. nce with PCT atutory invention iblication in
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application a any document in the file of For unpublished applications (1) If the benefit of the pending application that has: (a)) patent application public Article 21(2), a member the file contents; the pending applic any document in the (2) If the application is incorregistration, a U.S. paten accordance with PCT Article accordance with PCT Article application the pending application applicati	SCAMAN if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable if applicable	Lunder 35 U.S.C. 119(e), 12 or (b) published as a statutor patent application publication a copy of: of otherwise identified in a U.S. or an international patent a the public may obtain a cop	r = 100 GeV	(b)), as follows: 65 in another registration, a U.S. nce with PCT atutory invention iblication in
Direct access to pending app purchased from the Office of P For published applications that the file contents; the pending application and any document in the file of For unpublished applications. (1) If the benefit of the pending application that has: (a) patent application public Article 21(2), a member the file contents; the pending applic any document in the file contents; the pending applic any document in the secondance with PCT Au the pending applic Comp Warm Signature CARY WARM	SCAMAN if applicable SCAMAN if applicable SCADA if applicable SCADA Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Interpretation Int	Approximate the public out of the appropriate fee (independent of the public may obtain the public may obtain the public as a statutor publication a copy of: Different application publication a copy of: Different application publication a U.S. (or an international patent a copy of: Different application publication a the public may obtain a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a U.S. (or an international patent a copy of: Different application a Copy of: Different application a U.S. (or an international patent a copy of: Different application a Copy of: Different applicati	r = 100 GeV	(b)), as follows: 65 in another registration, a U.S. nce with PCT atutory invention iblication in

REQUEST FOR ACCESS T	TO AN ABANDONED	APPLICA	ION UNDE	ER 37 CFR 1.14
		In ta Applie	ellon ci	00
		LG	ALTERI	ØL –
ring completed farm to:		Application	Numbe:	Filed
ile Information Unit Irystal Plaza Three, Room 1001		107/21	5272	1. huslar
021 South Clerk Place		01/21	JATE	
rlington, VA				It U:
elephone: (703) 308-2733			P :	spsr No. 91-7-
nerzpy request access under 37 Cf oplication, which is identified in, or tachment):	FR 1.14(a)(1)(iv) to the appl r to which a benafit is claim	i cation file rac and, in the follo	ord of the abo wing docume	və identified ABAN nt (as shawn in the
United States Patent Applica	Ilon Publication No.		, li	ne,
United States Patent Number	6639055 , colum	in, I	inz,	10
WIPO Pub. No.				
				,
For published applications that err the file contents; the pending application as ori any document in the file of the For unpublished applications that (1) If the <u>benefit of the pending a</u> application that has: (a) issue	ginally fied; or a pending application. <u>ate:still panding:</u> <u>oplication is cisimed</u> under 3 d as a U.S. patent. or (b) pu	he public may (IS U.S.C. 119(6 blished as a sta), 120, 121, or Julary avenior	165 in enginer registration a V S
 For published applications that end the file contents; The pending application as oriany document in the file of the file of the file of the published apolications that; (1) If the benefit of the pending a application that has: (a) issue patent application publication, Article 21(2), a member of the file contents; The pending application is the file (2) If the application is incorporation registration, a U.S. patent application, a U.S. patent application, a U.S. patent application. 	e still pending, a member of I ginally filed; or a pending application. <u>are still pending:</u> collection is claimed under 2 d as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. Ed by reforence or otherwise lication publication for an int	he public may i 15 U.S.C. 119(s blished as a sta oplication public f: <u>Identified</u> in a l	blain a copy a), 120, 121, or lulary העפחונסר ation in accord J.S. patent, a s	165 In enailher A registrallon, a U.S Janze with PCT
For published applications that end the file contents; the pending application as ori- any document in the file of the For unpublished apolications that: (1) If the benefit of the pending a application that has: (a) issue patent application publication, Article 21(2), a member of the the file contents; the pending application any document in the file	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may i 15 U.S.C. 119(s blished as a sta oplication public f: <u>Identified</u> in a l	blain a copy a), 120, 121, or lulary העפחונסר ation in accord J.S. patent, a s	165 In enailher A registrallon, a U.S Janze with PCT
 For published applications that end the file contents; The pending application as oriany document in the file of the file of the file of the published apolications that; (1) If the benefit of the pending a application that has: (a) issue patent application publication, Article 21(2), a member of the file contents; The pending application is incorporation registration, a U.S. patent application with PCT Article 2 	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may i 15 U.S.C. 119(s blished as a sta oplication public f: <u>Identified</u> in a l	blain a copy a), 120, 121, or lulary העפחונסר ation in accord J.S. patent, a s	165 In enailher A registrallon, a U.S Janze with PCT
 For published applications that end the file contents; The pending application as oriany document in the file of the file of the file of the published apolications that; (1) If the benefit of the pending a application that has: (a) issue patent application publication, Article 21(2), a member of the file contents; The pending application is incorporation registration, a U.S. patent application with PCT Article 2 	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may i 15 U.S.C. 119(s blished as a sta oplication public f: <u>Identified</u> in a l	blain a copy a), 120, 121, or lulary העפחונסר ation in accord J.S. patent, a s	165 In enailher A registrallon, a U.S Janze with PCT
 For published applications that end the file contents; The pending application as oriany document in the file of the file of the file of the published apolications that; (1) If the benefit of the pending a application that has: (a) issue patent application publication, Article 21(2), a member of the file contents; The pending application is incorporation registration, a U.S. patent application with PCT Article 2 	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may (15 U.S.C. 119(s olished as a sta oplication public f: <u>Identified</u> in a t anational pater may obtain a c	blain a copy a), 120, 121, or lulary העפחונסר ation in accord J.S. patent, a s	165 In enailher A registrallon, a U.S Janze with PCT
For published applications that error the file contents; The pending application as ori- any document in the file of the For unpublished applications that is (1) If the benefit of the pending a application that has: (a) issue patent application publication. Article 21(2), a member of the the file contents; the pending application any document in the file (2) If the application is incorporate registration, a U.S. patent app accordance with PCT Article 2 the pending application	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may (15 U.S.C. 119(s olished as a sta oplication public f: <u>Identified</u> in a t anational pater may obtain a c	blain a copy o), 120, 121, or lutary invention ation in accord J.S. patent, a s it application pr copy of: $\frac{2}{04}$ Data	165 in enaiher nregistration, a U.S fance with PCT talutory invention ublication in
For published applications that error the file contents; The pending application as ori- any document in the file of the For unpublished applications that is (1) If the benefit of the pending a application that has: (a) issue patent application publication. Article 21(2), a member of the the file contents; the pending application any document in the file (2) If the application is incorporate registration, a U.S. patent app accordance with PCT Article 2 the pending application	e still pending, a member of I ginally filed; or a pending application. <u>are still pending</u> : <u>oplication is claimed</u> under 2 id as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or as originally filed; or as of the pending application. <u>ed by reforence or otherwise</u> lication publication, or an int 1(2), a member of the public	he public may (15 U.S.C. 119(s olished as a sta oplication public f: <u>Identified</u> in a t anational pater may obtain a c), 120, 121, or lutary invention ation in accord J.S. patent, a s of application pr apy of: 2/04	165 in enaiher nregistration, a U.S fance with PCT talutory invention ublication in
For published applications that error the file contents: The pending application as oriany document in the file of the For unpublished apolications that (1). If the benefit of the pending a application bublication, and that the file contents; The pending application publication, Article 21(2), a member of the file contents; The pending application any document in the file (2). If the application is incorporative registration, a U.S. patent application any document in the file (2). If the application, a U.S. patent application, a U.S. patent application. Signature Signature Signature Typed or printed name	e still pending, a member of a ginally filed; or a pending application. <u>are still panding: collection is claimed</u> under 3 d as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or a of the pending application. ed by reference or githerwise liceton publication, or an int (1(2), o member of the public as originally filed.	IS U.S.C. 119(8 olished as a sta polication public f: <u>Identified</u> in a t amational pater may obtain a c	blain a copy o), 120, 121, or lutary invention ation in accord J.S. patent, a s it application pr copy of: $\frac{2}{04}$ Data	165 in enaiher nregistration, a U.S fance with PCT talutory invention ublication in
For published applications that error the file contents; the pending application as oriany document in the file of the For unpublished applications that (1) If the benefit of the pending a application that has: (a) issue patant application publication, Article 21(2), a member of the the file contents; the pending application is incorporative registration, a U.S. patant application of the file content of the pending application. Signature Signature Made Jonanni (2) If the application is incorporative registration, a U.S. patant application.	e still pending, a member of a ginally filed; or a pending application. <u>are still panding: collection is claimed</u> under 3 d as a U.S. patent, or (b) pu , or an international patent a e public may obtain a copy o as originally filed; or a of the pending application. ed by reference or githerwise liceton publication, or an int (1(2), o member of the public as originally filed.	IS U.S.C. 119(8 olished as a sta polication public f: <u>Identified</u> in a t amational pater may obtain a c	blain a copy o), 120, 121, or lutary invention ation in accord J.S. patent, a s it application pr copy of: 2/04 Data FOR PTOL	165 in enaiher nregistration, a U.S fance with PCT talutory invention ublication in
For published applications that error the file contents: The pending application as oriany document in the file of the For unpublished apolications that (1). If the benefit of the pending a application bublication, and that the file contents; The pending application publication, Article 21(2), a member of the file contents; The pending application any document in the file (2). If the application is incorporative registration, a U.S. patent application any document in the file (2). If the application, a U.S. patent application, a U.S. patent application. Signature Signature Signature Typed or printed name	e still pending, a member of I ginally fied; or a pending application. <u>are still pending:</u> oplication is claimed under 3 d as a U.S. patent, or (b) pu , or an international patent a public may obtain a copy of as originally filed; or a of the pending application. ed by reforence or ginerwise lication publication, or an int (1(2), o member of the public as originally filed.	IS U.S.C. 119(8 olished as a sta polication public f: <u>Identified</u> in a t amational pater may obtain a c	blain a copy o), 120, 121, or lutary invention ation in accord J.S. patent, a s it application pr copy of: 2/04 Data FOR PTOL	165 in enaiher nregistration, a U.S fance with PCT talutory invention ublication in

Tranman Drice U.S. Department of Commerce, P.O. Eos 1430, Alexandra, VA 22113-1410 DO NGT SEND FEES OF COMPLETED FORMIS TO THIS SODDIESS. BRING TO: File Intermation Unit. Crystal Plaza Three, Room 1001, 2021 South Clark Placa, Arlington, VA.

ly you need assistance in completing the form will I too and a sec

Unda, the Panenwork Jaduction Abi d	TO AN AMALIOMIEN	APPLICATION UN	DER 37 CFR 1.1º
Under the Paterners Reductive Ant a	TO AN ABANDONED	In to Application di	
	_	Application Number	File
Sring completed form to:	REOF	Application 100000	6/14/91
ile Information Unit Inisial Plaza Three, Room 1001	''ECEIVER	AT/715272	
ort South Clark Place	MANA		PEDSPNO. +44
ulinning VA	MAY 3 - 2004		PEDSP No. 777
elephone: (703) 308-2733		1	
nerapy request access under 37 C	LI'lle Informationabili	ration lile racial of the	above-Identified ABLNDO
herapy raquest access under 37 C pplication, which is Identified In, i	or to which a benafit is claim	In the following doci	iment (as shown in the
nachment);	,		
A the time Ontent Andle	allon Publication No.	, p202,	linz
United States Haight Applic		1t	
United States Patent Numb	or <u>6407213</u> , column	n, IIA 2,	
	page, lin	· · · · · · · · · · · · · · · · · · ·	
WIPO PUB. No.			
purchased from the Office of Pue For published applications that i the file contents; inte pending application as i any document in the file of For unpublished applications that (1) If the benefit of the pending application that has: (a) iss output application publication	ariginally Alad; or the pending application, at are still pending; a <u>application is claimed</u> under 3 ued ex a U.S. patent, or (b) put on, or an International patent a	he public may obtain a co 15 U.S.C. 119(a), 120, 12 blianed es a statutory num pplication publication in a	1. ur 365 in endiner antion redistration, a U S
purchased from the Drice of Pue For published applications that in the file contents; the pending application as in any document in the file of For unpublished applications that (1) If the benefit of the pending application that has: (a) iss patch application publication amole 21(2), a member of the file contents; the pending application any document in the	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at ane still pending</u> : <u>a application is claimed</u> under 3 upon or an international patent at the public may obtain a copy a ion as originally filed; or file of the pending application.	he public may obtain a co 15 U.S.C. 119(a), 120, 12 blighted as a statutory hou polication publication in a f; hereinized in a U.S. pater	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any occument in the file of For unpublished application as is application that has: (a) iss patch application publication Amole 21(2), a member of the file contents; the pending application any document in the (2) If the application is incorpor registration, a U.S. patent a accordance with PET Articl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blianed as a statutory nou pplication publication in a f: <u>Identified</u> in a U.S. pater gmational patent applicat	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Difice of PUE For published applications that is the pending application as is any occument in the file of For unpublished application as is (1) If the benefit of the pending application that has: (a) is patch application publication Amplication publication the file contents; the pending application any document in the (2) If the application is incompo- conviction at 15 material	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blianed as a statutory nou pplication publication in a f: <u>Identified</u> in a U.S. pater gmational patent applicat	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any occument in the file of For unpublished application as is application that has: (a) iss patch application publication Amole 21(2), a member of the file contents; the pending application any document in the (2) If the application is incorpor registration, a U.S. patent a accordance with PET Articl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blianed as a statutory nou pplication publication in a f: <u>Identified</u> in a U.S. pater gmational patent applicat	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any occument in the file of For unpublished application as is application that has: (a) iss patch application publication Amole 21(2), a member of the file contents; the pending application any document in the (2) If the application is incorpor registration, a U.S. patent a accordance with PET Articl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blianed as a statutory nou pplication publication in a f: <u>Identified</u> in a U.S. pater gmational patent applicat	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any occument in the file of For unpublished application as is application that has: (a) iss patch application publication Amole 21(2), a member of the file contents; the pending application any document in the (2) If the application is incorpor registration, a U.S. patent a accordance with PET Articl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blianed as a statutory nou pplication publication in a f: <u>Identified</u> in a U.S. pater gmational patent applicat	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any document in the file of For unpublished applications that (1) If the benefit of the pending application that has: (a) iss patent application publication and file contents; the pending application any document in the (2) If the application is income registration, a U.S. patent a accordance with PCT Amicl Inc pending application ince pending applications ince pending applications accordance with PCT Amicl Ince pending applications accordance with PCT Amicl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blished as a statutory nou pplication publication in at f; identified in a U.S. pater smallonal patent applicat c may obtain a copy of; S/4/	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any document in the file of For unpublished applications that (1) If the benefit of the pending application that has: (a) iss patent application publication and file contents; the pending application any document in the (2) If the application is income registration, a U.S. patent a accordance with PCT Amicl Inc pending application ince pending applications ince pending applications accordance with PCT Amicl Ince pending applications accordance with PCT Amicl	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a co is U.S.C. 119(a), 120, 12 blished as a statutory nou pplication publication in at f; identified in a U.S. pater smallonal patent applicat c may obtain a copy of; S/4/	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any document in the file of For unpublished application as is application that has: (a) iss patch application publication and is contents; the pending application any document in the (2) If the application, a U.S. patent a accordance with PCT Article Ince pending application application, a U.S. patent a accordance with PCT Article Ince pending application Symptone Symptone Article Jurgen	and Records upon payment of o <u>are still pending</u> , a member at t the pending application, <u>at are still pending</u> : <u>a application is claimed</u> under 3 sued es a U.S. patent, or (b) puil on, or an international patent at the public may obtain a copy a lion as originally filed; or file of the pending application, <u>rated by reference or atherwise</u> application, or an int a 21(2), a member of the public	he public may obtain a construction of the public may obtain a construction of the publication of the public	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Difice of PUE For published applications that is the pending application as is any occument in the file of For unpublished applications that (1) If the benefit of the pending application that has: (a) iss patch application publication Amole 21(2), a member of the file contents; the pending application any document in the (2) If the application is incorpor registration, a U.S. patent a accordance with PCT Article Inc pending sobjication Symptone Symptone Rock Jonna Typed or printed name	and Records upon payment of o are still pending, a member at t criginally Alad; or the pending application, at are still pending: a application is claimed under 3 sued as a U.S. patent, or (b) put on, or an international patent at the public may obtain a copy a lon as originally filed; or file of the pending application, raised by reference or stherwise application publication, or an int a 21(2), a member of the public ion as originally filed.	he public may obtain a co is U.S.C. 119(a), 120, 12 blished as a statutory nou pplication publication in at f; identified in a U.S. pater smallonal patent applicat c may obtain a copy of; S/4/	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention
purchased from the Drice of Pue For published applications that is the pending application as is any document in the file of For unpublished application as is application that has: (a) iss patch application publication and is contents; the pending application any document in the (2) If the application, a U.S. patent a accordance with PCT Article Ince pending application application, a U.S. patent a accordance with PCT Article Ince pending application Symptone Symptone Article Jurgen	plicable	he public may obtain a construction of the public may obtain a construction of the publication of the public	apy of: 1. or J65 in enoined antion registration, a U S ceordance with PC T at, a statutory invention

Linnan Drie, U.S. Desament ni Commare, P.O. Ist 1130, Alesandes, VA 23313-1418 OG NOT SIND FEES DA COMPLETEU FOAN 2000/FSS - BRING 70 - File Intermetion Unit, Crystel Plaza Three, Room 1001, 2021 South Clark Plaza, Arlington, V.a.

If you need assistance in completing the torus only to one -----

Ŋ

.

. •

6-4513 PTO/SE/68 (07-03) Approved for use through 7/31/2003. OMB 0651-0031 U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OME control number. REQUEST FOR ACCESS TO AN ABANDONED APPLICATION UNDER 37 CFR 1.14 In re Application of RECEIVED aligetion Number Bring completed form to: Crystal Plaza Three, Room D01 JUN 0 7 2004 2021 South Clark Place ." File Information Unit Paper No._ Arlington, VA Telephone: (703) 308-2733 I hereby request access under 37 CFR 1.14(a)(1)(iv) to the application file record of the above-identified ABANDONED application, which is identified in, or to which a benefit is claimed, in the following document (as shown in the attachment): United States Patent Application Publication No._____, page, _____ line ____ United States Patent Number (407213, column _____, line, _____ or , line ____, page ____ WIPO Pub. No.____ Related Information about Access to Pending Applications (37 CFR 1.14): Direct access to pending applications is not available to the public but copies may be available and may be purchased from the Office of Public Records upon payment of the appropriate fee (37 CFR 1.19(b)), as follows: For published applications that are still pending, a member of the public may obtain a copy of: the file contents: the pending application as originally filed; or any document in the file of the pending application. For unpublished applications that are still pending: (1) If the benefit of the pending application is claimed under 35 U.S.C. 119(e), 120, 121, or 365 in another application that has: (a) issued as a U.S. patent, or (b) published as a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of: the file contents; the pending application as originally filed; or any document in the file of the pending application. (2) If the application is incorporated by reference or otherwise identified in a U.S. patent, a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of: the pending application as originally filed. attire Typed or printed hame Registration Number, if applicabl File Information Unit Unit Telephone Number This collection of information is required by 37 CFR 1.14. The information is required to obtain or ratain a benefit by the public which is to file (and by the USPTO

This collection of information is required by 37 CFR 1.14. The information is required to obtain or retain a benefit by the public which is to file (and by the USF) to process) an application. Confidentially is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case, Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burdan, should be sent to the Chiaf Information Officer, U.S. Patent and amount of time you require to complete this form and/or suggestions for reducing this burdan. VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS Trademant: Office, U.S. Department of Commerces, P.O. 80x 1450, Alexandria, VA 22313-1450, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA.

PTO/S8/68 (07-03-Approved for use through 7/51/2003, OME 0551-0031 U.S. Fätent and Tradsmark Office: U.S. DEPARTMENT OF COMMERCE Under the Paparwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control rum REQUEST FOR ACCESS TO AN ABANDONED APPLICATION UNDER 37 CFR 1.14 In re Application of Filed Application Number ENED Bring completed form to: 76-File Information Unit Crystal Plaza Three, Room 1001 JUN 0.9.2004 2021 South Clark Place Arlington, VA Telephone: (703) 308-2733 File Information / thit I hereby request access under 37 CFR 1.14(a)(1)(iv) to the application file record of the above-identified ABANDONED application, which is identified in, or to which a benefit is claimed, in the following document (as shown in the attachment): United States Patent Application Publication No. _____, page, _____ line _____ United States Patent Number 15 61 407, 213 B1, column _____, line, _____ or WIPO Pub. No._____, page _____, line _____. Related Information about Access to Pending Applications (37 CFR 1.14): Direct access to pending applications is not available to the public but copies may be available and may be purchased from the Office of Public Records upon payment of the appropriate fee (37 CFR 1.19(b)), as follows: For published applications that are still pending, a member of the public may obtain a copy of: the file contents; the pending application as originally filed; or any document in the file of the pending application. For unpublished applications that are still cending: (1) If the benefit of the pending application is claimed under 35 U.S.C. 119(a), 120, 121, or 365 in another application that has: (a) issued as a U.S. patent, or (b) published as a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of: the file contents; the pending application as originally filed; or any document in the file of the pending application. (2) If the application is incorporated by reference or otherwise identified in a U.S. patent, a statutory invention registration, a U.S. patent application publication, or an international patent application publication in accordance with PCT Article 21(2), a member of the public may obtain a copy of: the pending application as originally filed. Date Signature THE FRANKFRANKAN-Typed or printed name Approved and Registration Number, if applicable Unit Telephone Number

This collection of information is required by 37 CFR 1.14. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will very depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA.

	tene to a collection of information unless is disclays a valid GMS constitution;
EQUEST FOR ACCESS TO AN ABANDONE!	
RECEIVED RECEIVED AUG 1 6 2004 AUG 1 6 2004 File Information Unit File Information Unit	In re Application of CARTEY Application Number Filed ON 1500 01491 Faper No. #17
ereby request access under 37 CFR 1.14(a)(1)(iv) to the ap plication, which is identified in, or to which a benefit is cla achment): United States Patent Application Publication No.	
United States Patent Number 61012R_, col	
WIPO Pub. No, page	
 patent application publication, or an international pater Article 21(2), a member of the public may obtain a contents; the pending application as originally filed; or any document in the file of the pending application (2) If the application is incorporated by reference or other registration, a U.S. patent application publication, or a accordance with PCT Article 21(2), a member of the pending application is pater or the pending application. 	bublished as a statutory invention registration, a con- nt application publication in accordance with PCT py of: don. wise identified in a U.S. patent, a statutory invention in international patent application publication in
the pending application as originally filed.	BILO Date FOR PRODECTIVED Approved by: AUC/16/2004 File Information

This collection of information is required of all other first and 37 U.S.C. 122 and 37 OFR 1.14. This collection is estimated to take 12 minutes to complete, induction to process) an application. Condentially is governed by 35 U.S.C. 122 and 37 OFR 1.14. This collection is estimated to take 12 minutes to complete, induction gainarity, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and mount of time you require to complete this form and/or suggestions for reducing this burden. should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 32315-1450, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADORESS. BRING TO: File Information Unit, Crystal Plaze Three, Room 1001, 2021 South Clark Place, Arillagton, VA, ADORESS. BRING TO: File Information Unit, Crystal Plaze Three, Room 1001, 2021 South Clark Place, Arillagton, VA,

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2. 379

Under the Peperwork Reduction Act of 199	· · · · · · · · · · · · · · · · · · ·		oprovéd for use tinn temark Office: U.S. nation unless it disc		
Under the Pederwork Reduction Act of 199 EQUEST FOR ACCESS T	O AN ARANDONED	APPLICATIO	N UNDER 3	7 CFR 1.14	1
EQUESTFORACCESCT		In re Application	i cî		
ng completed form to: e Information Unit ystal Plaza Three, Room 1001 21 South Clark Place ington, VA lephone: (703) 308-2733	RECEIVED NOV 0 1 2004 File Information Unit		JJZ Paper	1	
ereby request access under 37 C plication, which is identified in, o achment):	FR 1.14(a)(1)(iv) to the appl r to which a benefit is clain ation Publication No. <u>680</u>				18
United States Patent Applica	etion Publication No. <u>Cour</u>	no line	ż,	or	
United States Patent Number	37, Will	ine .			
WIPO Pub. No	, page, I	111-			
patent application publicat Article 21(2), a member of the file contents; the pending applica any document in the (2) If the application is incoro registration, a U.S. patent	are still <u>bending</u> , a member of originally filed; or the pending application.	if the public may ob if 35 U.S.C. 119(e). published as a stat t application publica y of: on. <u>vise identified</u> in a l	ntain a copy or: , 120, 121, or 3 utory invention ation in accorda J.S. patent, a si nt apolication pu	65 in another registration, a U ance with PCT	J.S.
Typed or printed name	drisuez	A	NON=BO	Kaut	
Registration Number, if (73) (18-27) Telephone Num his collection of information is required by 37 his collection of information is required by 37	27				w the USPT
tis callsction of information is required by 37 and application. Canādenizelity is go process) an application. Canādenizelity is application and submitting the cample			Contract of the local division of the local		

	STO/SE/86 (07-02)
	Approved for use through 7/31/2003. ONE 0851-003
Under the Pepervork Reduction Act of 1998. no persons are required to rescond	to a collection of information unless it disclays a valic office and
Under the Pederwork Reduction Act of 1984. An entropy and interest and interest of the second	APPLICATION UNDER STOLIC TIC
	in re Application of Physics R
	Application Number Filed
Bring completed form to: File Information Unit	071715202 7114191
Crystal Plaza Three, Room 1001	
2021 South Clark Flace	Paper No. ##
Telephone: (703) 308-2733	
I hereby request access under 37 CFR 1.14(a)(1)(iv) to the apoli application, which is Identified in, or to which a benefit is claim attachment):	cation file record of the above-identified ABANDONED ed, in the following document (as shown in the
	, page,line
United States Patent Number (0401213, colum	
WIPO Pub. No, page, lir	ne NUV 2 9 2004 `
Related Information about Access to Pending A Direct access to pending applications is not available to the purchased from the Office of Public Records upon payment of <u>For oublished applications that are still pending</u> , a member of the file contents; the pending application as originally filed; or any document in the file of the pending application. <u>For unpublished applications that are still pending</u> . (1) If the <u>benefit of the pending application is claimed</u> under application that has: (a) issued as a U.S. patent, or (b) p patent application publication, or an international patent Article 21(2), a member of the public may obtain a copy the file contents; the pending application as originally filed; or any document in the file of the pending application (2) If the application, a U.S. patent application, or an accordance with PCT Article 21(2), a member of the public the pending application as originally filed.	the appropriate fea (37 CFR 1.19(b)), as follows: the public may obtain a copy of: 35 U.S.C. 119(e), 120, 121, or 365 in another ublished as a statutory invention registration, a U.S. application publication in accordance with PCT of: n. <u>ise identified</u> in a U.S. patent, a statutory invention international patent application publication in blic may obtain a copy of:
A CHART	FOR PTO USE ONLY
Typed or printed name	Approved by RECEIVED
Registration Number, if applicable	Unit NOV 2 9 2004
Telephone Number	File Information Unit
Telephone Number This collection of information is required by 37 CFR 1.14. The information is required to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR getharing, preparing, and submitting the completed application form to the USPTO. getharing or require to complete this form and/or suggestions for reducing this amount of time you require to complete this form and/or suggestions for reducing this amount of time, you require to complete this form and/or suggestions for reducing this amount of time, you require to complete this form and/or suggestions for reducing this amount of time. U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA.22 DDRESS. BRING TO: File Information Unit, Crystal Plaza Three, R.38	burden, should be sent to the Chief Inclineson Chief STER SCENS TO THIS

•

PTO/SB/68 (11-04) 1gh 7/31/2006. OMB 0651-0031

Approvad for use through the fillede.	0.0.0
Land Tendemork Office: U.S. DEPARTMENT	
L L	

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF CONVINCENCE to a collection of information unless it displays a valid OMB control number.

Under the Paperwork Reduction Act of 1995, no pers	ABANDONED APPLICATION UNDER 37 CFR 1.14
	In re Application of
ring completed form to:	
ile Information Unit, Room 2E04 900 Crystal Drive	Application Number Filed
rlington, VA 22202-3514	07/113272 June 14,1991
elephone: (703) 308-2733	Paper No. \$60
hereby request access under 37 CFR 1.14 pplication, which is not within the file jacl nd which is identified in, or to which a be	4(a)(1)(iv) to the application file record of the above-identified ABANDONE ket of a pending Continued Prosecution Application (CPA) (37 CFR 1.53(c enefit is claimed, in the following document (as shown in the attachment):
United States Patent Application Pu	ublication No, page, line,
United States Patent Number	0 5429,7column, line, or
	, page, line
 available through the Public Patent App web site (www.uspto.gov). Terminals it The member of the public may also be the appropriate fee. Such copies must appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending application. For unpublished applications that are (1) If the benefit of the pending application, application that has: (a) issued are patient application publication, publication, publication, publication publication, or Article 21(2), a member of the puoriginally filed; or any document (2) If the application is incorporated 	a power to inspect, cannot of the application file, then the file is made public is entitled to a copy of the application file, then the file is made olication Information Retrieval system (Public PAIR) on the USPTO internet that allow access to Public PAIR are available in the Public Search Room, entitled to obtain a copy of all or part of the application file upon payment of the purchased through the Office of Public Records upon payment of the <u>Il pending</u> , a member of the public may obtain a copy of: lication as originally filed; or any document in the file of the pending <u>still pending</u> : <u>ication is claimed</u> under 35 U.S.C. 119(e), 120, 121, or 365 in another is a U.S. patent, or (b) published as a statutory invention registration, a U.S. an international patent application publication in accordance with PCT ublic may obtain a copy of: the file contents; the pending application as in the file of the pending application. by reference or otherwise identified in a U.S. patent, a statutory invention ration publication, or an international patent application publication publication as (2), a member of the public may obtain a copy of the pending application as
<u> </u>	$\frac{\Delta a v. 18. 05}{Date. 18. 05}$
Signatu	FOR PTO USE ONLY
Typed or prin	
Typed or print	
	Approved by: (initials); }
Registration Numb	
71418.0	1330
Telephone	
USPTO to process) an application. Confidentiality is got complete, including gathering, preparing, and submitting comments on the amount of time you require to comple	1 and 1.14. The information is required to obtain or retain a benefit by the public which is to file (and verned by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 12 minutes to g the completed application form to the USPTO. Time will vary depending upon the individual case. A ste this form and/or suggestions for reducing this burden, should be sent to the Chief Information Offic Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED ormation Unit, Room 2E04, 2900 Crystal Drive, Arlington, Virginia.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

	PTO/SB/68 (07-03 Approved for use through 7/31/2003. OMB 0651-003
Under the Paperwork Reduction Act of 1995, no persons are required to respo	and to a collection of information unless it displays a valid of the control number
EQUEST FOR ACCESS TO AN ABANDONED	APPLICATION UNDER 37 CFR 1.14
	Lin to Application of
R	Carter et al
ing completed form to:	Application Number Filed 07/715,272 6/14/91
le Information Unit ystal Plaza Three, Room 1D01 JAN 3 1 2005	07(115,272 614(91
	H E I
File Information view	Paper No. 77-51
hereby request access under 37 CFR 1.14(a)(1)(iv) to the a	pplication file record of the above-identified ABANDONE
pplication, which is identified in, or to which a benefit is ch	aimed, in the following document (as shown in the
ttachment):	
United States Patent Application Publication No.	, page, line,
United States Patent Number <u>4,800,738</u> , co	olumn, line, or
WIPO Pub. No, page	line
WIPO Pub. No, page	
 patent application publication, or an international publication, or an international publication 21(2), a member of the public may obtain a the file contents; the pending application as originally filed; or any document in the file of the pending appli (2) If the application is incorporated by reference or ot registration, a U.S. patent application publication, or any document in the file 21(2), a member of the pending application application for a second and with PCT Article 21(2), a member of the pending application application publication, a the pending application publication application application publication, a the pending application publication, a the pending application publication application publication, a the pending application publication, a the pending application publication application publication, a the pending application publication, a the pending application publication application publication, a the pending application publication, a the pending application application publication, a the pending application publication, a the pending application application publication, a the pending application appli	(b) positionates are publication in accordance with PCT copy of: ication. <u>herwise identified</u> in a U.S. patent, a statutory invention or an international patent application publication in
the pending application as originally filed.	ne public may obtain a copy of:
the pending application as originally filed.	
Rayhine K. Petitt	Lan. 31, 2005
Kayfine K. Petitt Signature	Jan. 31, 2005
the pending application as originally med. Kayfine K. Petitt Signature	Jan. 31, 2005
Rayfine K. Petitt	Jan. 31, 2005
the pending application as originally med. Kayfine K. Petitt Signature Rayline K. Petitt Typed or printed name n/a	Jan. 31, 2005
the pending application as originally med. Kayfine K. Petitt Signature Rayline K. Petitt Typed or printed name	JAN 3 1 2005
the pending application as originally med. Kayfine K. Petutt Signature Rayline K. Petitt Typed or printed name <u>n/a</u> Registration Number, if applicable	JAN 31, 2005
the pending application as originally med. Kayfine K. Petitt Signature Rayline K. Petitt Typed or printed name n/a	Jan. 31, 2005 Date FOR'PTO'USE ONLY JAN 3 1 2005 Approved by: (initials) File Information 1/0.2

This collection of information is required by 37 CFR 1.14. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTC to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTC. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450, DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA.

 $\cdot l$

PTO/S8/62 (07-03) Approved for use through 7/31/2003, OMB 0651-0031 U.S. Patent and Tradecond Office: U.S. DEPARTMENT OF COMMERCE to a collection of inform unders it diseases a valid CMB control number.

RECEIV APR 1 5 20 APR 1 5 20 APR 1 5 20 APR 1 5 20 APR 1 5 20 File Information File Information File Information File Information CFR 1.14(a)(1)(iv plication, which is identified in, or to which a beat iachment): United States Patent Application Publication United States Patent Number <u>SSA/3</u> WIPO Pub. No, page Related Information about Access to Direct access to pending applications is not purchased from the Office of Public Records upor	Application Number D I D I D I D I D I D I D I D I D I D	line
e Information Unit ystal Plaza Three, Room 1D01 21 South Clark Place ington, VA lephone: (703) 308-2733 ereby request access under 37 CFR 1.14(a)(1)(iv plication, which is identified in, or to which a ber tachment): United States Patent Application Publication United States Patent Number, page WIPO Pub. No, page	Unit 07/7/54 Unit 07/7/54 to the application file record of the efit is claimed, in the following docu No, page, 37, column, line,	Paper No. <u>#52</u> above-identified ABANDO ument (as shown in the
21 South Clark Place ington, VA lephone: (703) 308-2733 ereby request access under 37 CFR 1.14(a)(1)(iv plication, which is identified in, or to which a be- tachment): United States Patent Application Publication United States Patent Number, page WIPO Pub. No, page	to the application file record of the efit is claimed, in the following doct No, page, 37, column, line,	above-identified ABANDO ument (as shown in the line
ereby request access under 37 CFR 1.14(a)(1)(iv plication, which is identified in, or to which a ber tachment): United States Patent Application Publication United States Patent Number, page	No, page, 37, column, line, , line	line
United States Patent Application Publication United States Patent Number, page	No, page, 37, column, line, , line	line
United States Patent Application Publication United States Patent Number <u>589/3</u> WIPO Pub. No, page	37. column, line,	line or
United States Patent Number, page, p	37. column, line,	or
WIPO Pub. No, page	, into	
WIPO Pub. No, page	, into	
Related Information about Access to	Num Applications (37 CFF	· · · · · · · · · · · · · · · · · · ·
Related Information about Access to		2 1.14):
 billect access from the Office of Public Records upor For published applications that are still pending the file contents; the pending application as originally filed; any document in the file of the pending applications that are still pending (1) If the benefit of the pending application that has: (a) issued as a U.S. patent application publication, or an interrest Article 21(2), a member of the public may the file contents; the pending application as originall any document in the file of the pending application as originall (2) If the application is incorporated by reference registration, a U.S. patent application as prigrated by reference the pending application applica	lication. <u>Ig:</u> <u>laimed</u> under 35 U.S.C. 119(e), 120, atent, or (b) published as a statutory ir bitional patent application publication ir bitain a copy of: filed; or ing application. <u>ice or otherwise identified</u> in a U.S. pa ication, or an international patent appli ber of the public may obtain a copy o	121, or 365 in another ovention registration, a U.S. accordance with PCT
	4-15	5-05
fillupy pit in	Date	and an and a second
Signature)	F	OR PTO USE ONLY
Typed or printed name		A
	Approv	d bRECHINIals
Registration Number, if applicable	1 mile	APR 1 5 2005
415-1077	Unit: _	HIN & J GWVJ
Telephone Number This collection of information is required by 37 CFR 1.14. The init to process) an application. Confidentiality is governed by 35 U.S.0		File Information Unit

anount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information United. Some amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information United. Some amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information United. Trademark Office, U.S. Department of Commerce, P.O. Box 1450. Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORM Trademark Office, U.S. Department of Commerce, P.O. Box 1450. Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORM ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1D01, 2021 South Clark Place, Arlington, VA.

Best Available Copy	racad la s collection of information unless is clasicys a valid whis contra number.
REQUEST FOR ACCESS TO AN ABANDONE	DAPPLICATIC UNDER 37 OFR 1.14
	Inte Application of
Bal Complete fam lo: Figlaformation Unit Official Plaza Three, Room 1001 2021 South Clark Place Arlington, VA	Application Number 07/7/5072 6/4/9/ Paper No. 3
Talephone: (703) 308-2733 File Information Unit	
I hereby request access under 37 OFR 1.14(a)(1)(iv) to the ap application, which is identified in, or to which a benafit is dia attachment):	
United States Patent Application Publication No.	, <u>P2</u> 25,Nic
United States Patent Number 6407-2/3_ col	
WIPO Pub. No dege	, line
 Related Information about Access to Pending Direct access to pending explications is not available to the output of the seconds upon payment of the file contents; the gending application as originally filed; or any document in the file of the pending application is claimed under application that has: (a) issued as a U.S. patent, or (b) patent application publication as originally filed; an Article 21(2), a member of the pending application as originally filed; or any document in the file of the pending application is claimed under application publication, or an international patent Article 21(2), a member of the pending application as originally filed; or any document in the file of the pending application as originally filed; or (b) patent application publication as originally filed; or any document in the file of the pending application as originally filed; or (b) patent application publication as originally filed; or any document in the file of the pending application as originally filed; or any document in the file of the pending application as originally filed; or any document in the file of the pending application as originally filed; or any document in the file of the pending application as originally filed; or any document in the file of the pending application or an accordance with PCT Article 21(2), a member of the publication, or an accordance with PCT Article 21(2), a member of the publication, or any accordance with PCT Article 21(2), a member of the publication. 	of the appropriate fee (37 CFR 1.19(b)), as follows: of the public may obtain a copy of: er 35 U.S.C. 119(e), 120, 121, or 365 in another oublished as a statutory invention registration, a U.S. It application publication in accordance with FCT by of: on.
Picky Jen	9/12/05
RICKY GARCIA	ECR.FIQUEE CHLY
Typed or printed name	REC.
	Approved by <u>Rectored 2005</u>
Registration Number, if applicable	Unit: File information L' it
the information is required by 37 CFR 1.14. The information is required	to obtain or ratein a banafi by the public which is to file (and by the USP and to the USP and to the standard to attract to have no minutes to boostates, inducing
The collection of information is required by GT CFR 1.14. The information is required to process) an application, Confidentially is governed by 15 U.S.C. 122 and 17 CFR quitaring, preparing, and submitting the completed application form to the USPTO. If shown of time you require to complete this form and/or suggestions investiging this shown of time you require to complete this form and/or suggestions investigating the predemark Office, U.S. Department of Commerce, P.J. Box 1460, Alexandriz, VA.	inter ville contraction of the incident case . Any contraction is of the second s
epitering, preparing, and submitting the complexed application form to the option of the status of time you require to complexe this form and/or suggestions for relating this indone of time you require to complexe this form and/or suggestions for relating this indone of time you require to complexe this form and/or suggestions for relating this indone of time you require to complexe this form and/or suggestions for relating this indone of time you require to complexe this form and/or suggestions for relating the indone of time you require to complexe the submitted of the indone of time you require to complexe the submitted of the indone of the submitted of the indone of the submitted of the indone of the submitted of the indone of indone	a a line se not stab test og tokal ETta SVPM
ADDRESS, BRING TO: File Information Ghit, Unyacat Free 385	

Ň

Undarth	te Pepervork Ret	luction Am of 195	E. no persons.	are required	b rescord	c a collectio	េនាធ ការប្រ ត នាំ តែតែកា	รปอส นภ!	ess it disc	cugh 7/31/2 DEPARTNIS liays a veilid	ONE 100	irat aum:
and the second		CCESS TO										
					1		lication		7	<u></u>		
					·			C	44	81L	5K	
Rrina comolei	ted form to war		N KANA N	1	Į	Applicati	on Mund		0.00	Filed	1.	
File Informal	tion Unit	RECE				NT	[]]]	5	12) []	///	5/7
Crystel Plezi	a Three, Roc Clady Plade	m 1D01	7 2006		L	- El for			/-		/	
		MAR 2	. 2000			,			Paper	No	450	1
felephone: ((703) 308-27:	3	ation Un	it l					rehei	110. 24		۷
•		ile Informa						م حمات ال		dontified	ARAN	DONE
hereby req	juest access	under 37 CF	R 1.14(a)(1 to which a	i benefit i	s claimed	tion file 1, in the f	ollowing	i qocn 1 qocn	ment (is shown	In the	20112.
ttachment)					11	nna	12	-		**		
1.1-	the of Charles D	atent Applicati	ion Dublica		NH	/////	$\langle \mathcal{L} \rangle$		line			
											•	
Un	lited States P	etent Number	<u></u>		, column		, line, _	,,,	••••••	or		
WE	PO Pub. No.		, pa	ge	, line		•					
•••	• • • • • • • • • • •							`				
Direct a purchas For pub the	access to pe sed from the C <u>alished apolic</u> e file contents	tion about nding applica Office of Public ations that are ; plication as off	ations is no c Records u e still oendi	ot availabi ipon payn <u>no</u> , a men	e to the p ant of the		iate fee l	37 CF	R 1.19(ay be lows:	
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>bilished application</u> e file contents e pending apply by document i <u>bublished apply</u> the <u>benefit of</u> pplication that atent applicat rticle 21(2), a the file c like pend any docu the application, a coordance wit	nding applica office of Public ations that are control to a sori- n the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing applicatior ument in the fil n is incorvora J.S. patent application b PCT Article	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio in a U.S oatent a	20, 121 ry inve n in ac	, or 36 , or 36 ntion re- cordan	5 in anoth gistration ce with P	er I, a U.S CT ention	-
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>bilished application</u> e file contents e pending apply by document i <u>bublished apply</u> the <u>benefit of</u> pplication that atent applicat rticle 21(2), a the file c like pend any docu the application, a coordance wit	nding applica office of Public ations that are in the file of the offication as ori in the file of the offications that. the vending a has: (a) issue on publication member of the ontents; ing applicatior ing incorrora	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio in a U.S oatent a	20, 121 ry inve n in ac	, or 36 , or 36 ntion re- cordan	5 in anoth gistration ce with P	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>bilished application</u> e file contents e pending apply by document i <u>bublished apply</u> the <u>benefit of</u> pplication that atent applicat rticle 21(2), a the file c like pend any docu the application, a coordance wit	nding applica office of Public ations that are control to a sori- n the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing applicatior ument in the fil n is incorvora J.S. patent application b PCT Article	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio in a U.S oatent a	20, 121 ry inve n in ac	, or 36 , or 36 ntion re- cordan	5 in anoth gistration ce with P	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>alished applicate</u> e file contents e pending app ny document is <u>abublished applicate</u> the <u>benefit of</u> pplication that atent application the file c the pend any docu the application gistration, a t coordance witt the pend	nding applica office of Public ations that are blication as ori in the file of the <u>blications that</u> the <u>bending</u> a has: (a) issue on publication member of the ontents; ing applicatior imment in the fil n is <u>incoroora</u> J.S. patent application	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio in a U.S oatent a	20, 121 n a co 20, 121 ny inve in in ac	, or 36 , or 36 ntion re- cordan	5 in anoth gistration ce with P	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>alished applicate</u> e file contents e pending app ny document is <u>abublished applicate</u> the <u>benefit of</u> pplication that atent application the file c the pend any docu the application gistration, a t coordance witt the pend	nding applica office of Public ations that are control to a sori- n the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing applicatior ument in the fil n is incorvora J.S. patent application b PCT Article	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto publicatio in a U.S patent a in a cop	20, 121 n a co 20, 121 ny inve in in ac	, or 36 ntion re- cordan	5 in anoth gistration ce with P	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>bilished acolic</u> e file contents e pending ap ny document i <u>bublished acolic</u> the <u>benefit</u> of pplication that atent application the file c the application, a l ccordance with the pend Sig	nding applica office of Public ations that are in the file of Public blication as ori in the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing application ing application J.S. patent application J.S. patent application application for a since of the patents; ing application ing application in	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto publicatio in a U.S patent a in a cop	20, 121 ry inve in a co 20, 121 ry inve in in ac	, or 36 ntion re- cordan	5 in anoth gistration ce with P utory inve lication in	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If ap pa Ar	access to pe sed from the C <u>alished applicate</u> e file contents e pending app ny document is <u>abublished applicate</u> the <u>benefit of</u> pplication that atent application the file c the pend any docu the application gistration, a t coordance witt the pend	nding applica office of Public ations that are in the file of Public blication as ori in the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing application ing application J.S. patent application J.S. patent application application for a since of the patents; ing application ing application in	titions is no c Records L e still cendin ginally filect e pending a <u>are still cen</u> and still cen e cending a construction e of the pent ted by refar plication pu 21(2), a me	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio	20, 121 ry inve n in aco policati y of:	, or 36 ntion re- cordan	5 in anoth gistration ce with P utory inve lication in	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If aF pa Ar (2) If re ac	access to pe access to pe and from the C <u>bilished apolic</u> e file contents e pending app ny document i <u>ublished apolic</u> the <u>benefit</u> data the <u>benefit</u> data atent application any doct the application signification coordance with the pend Signification Signification Signification	nding applica office of Public ations that are plication as ori- n the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing applicatior imment in the fil n is <u>incorvora</u> J.S. patent application ing a	titions is no c Records L e still cendir ginally filec e pending a <u>are still cendir</u> and as a U.S h, or an inte e public ma h as origina e of the per ted by refar plication pu 21(2), a men n as origina	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto publicatio in a U.S patent a in a cop	20, 121 ry inve n in aco policati y of:	, or 36 ntion re- cordan	5 in anoth gistration ce with P utory inve lication in	er I, a U.S CT ention	
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If aF pa Ar (2) If re ac	access to pe access to pe and from the C <u>bilished apolic</u> e file contents e pending app ny document i <u>ublished apolic</u> the <u>benefit</u> data the <u>benefit</u> data atent application any doct the application signification coordance with the pend Signification Signification Signification	nding applica office of Public ations that are in the file of Public blication as ori in the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing application ing application J.S. patent application J.S. patent application application for a since of the patents; ing application ing application in	titions is no c Records L e still cendir ginally filec e pending a <u>are still cendir</u> and as a U.S h, or an inte e public ma h as origina e of the per ted by refar plication pu 21(2), a men n as origina	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio	20, 121 ry inve n in aco policati y of:	, or 36 ntion re- cordan	5 in anoth gistration ce with P utory inve lication in	er I, a U.S CT ention	•
Direct a purchas <u>For pub</u> the the an <u>For unp</u> (1) If aF pa Ar (2) If re ac	access to pe access to pe and from the C <u>bilished apolic</u> e file contents e pending app ny document i <u>ublished apolic</u> the <u>benefit</u> data the <u>benefit</u> data atent application any doct the application signification coordance with the pend Signification Signification Signification	nding applica office of Public ations that are plication as ori- n the file of the blications that the vending a has: (a) issue on publication member of the ontents; ing applicatior imment in the fil n is <u>incorvora</u> J.S. patent application ing a	titions is no c Records L e still cendir ginally filec e pending a <u>are still cendir</u> and as a U.S h, or an inte e public ma h as origina e of the per ted by refar plication pu 21(2), a men n as origina	ot availabl ipon paym ng, a men di; or application is claimed b. patent, c mational ; ay obtain a illy filed; o nding app rence or o blication, ember of t	e to the p nent of the aber of the under 35 or (b) pub patent ap a copy of: r lication. thetwise	U.S.C. 1 ished as plication ;	19(e), 12 a statuto bublicatio	20, 121 ry inve n in aco policati y of:	, or 36 ntion re- cordan	5 in anoth gistration ce with P utory inve lication in	er , a U.S CT	

getnenng, grepenng, end sublinking the completed application form of the data for reducing this burden, should be sent to the Chief Information Officer, U. amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U. Tredemark Office, U.S. Department of Commance, P.O. Sox 1450. Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORM ADDRESS. BRING TO: File Information Unit, Crystal Plaza Three, Room 1001, 2021 South Clark Place, Arlington, VA

۰,

Under the Paperwork (Cooperation		tion of information unless it displays a valid OMB control number.
Under the Paperwork Reduction AC OF 1933, in Page REQUEST FOR ACCESS TO AN	ABANDONED APPLI	
Bring completed form to: File Information Unit, Room 2E04 2900 Crystal Drive Arlington, VA 22202-3514 Felephone: (703) 308-2733	In re Application of Application Number EC 07/7/52 SEP ((a)(1)(iv) to the Epideiga (2)	FIVED Filed Paper No. 753 Paper No
and which is identified in, or to which a be	ifent is clained, in the feat	
United States Patent Number	321337, column	_, page, line, , line, or
WIPO Pub. No Related Information About Ac		
A member of the public, acting without a through the FIU. If the member of the p available through the Public Patent App web site (www.uspto.gov). Terminals the through the Public Patent App web site (www.uspto.gov).	lication Information Retrieval s	ystem (Public PAIR) on the USPTO internet
The member of the public may also be of the appropriate fee. Such copies must l appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending appli- application. For unpublished applications that are s (1) If the benefit of the pending appli- application that has: (a) issued as patent application publication, or Article 21(2), a member of the pu- originally filed; or any document in (2) If the application is incorporated by	an anow access of a copy of all of be purchased through the Offi <u>pending</u> , a member of the put ication as originally filed; or an <u>still pending</u> : <u>cation is claimed</u> under 35 U.S is a U.S. patent, or (b) publishe an international patent applica blic may obtain a copy of: the n the file of the pending applic by reference or otherwise iden	or part of the application file upon payment of ce of Public Records upon payment of the olic may obtain a copy of: y document in the file of the pending c.C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as
The member of the public may also be of the appropriate fee. Such copies must l appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending appli application. For unpublished applications that are s (1) If the benefit of the pending appli application that has: (a) issued as patent application publication, or Article 21(2), a member of the pu- originally filed; or any document in registration, a U.S. patent applica- accordance with PCT Article 21(2)	an anow access of a copy of all of be purchased through the Offi <u>pending</u> , a member of the put ication as originally filed; or an <u>still pending</u> : <u>cation is claimed</u> under 35 U.S is a U.S. patent, or (b) publishe an international patent applica blic may obtain a copy of: the n the file of the pending applic by reference or otherwise iden	or part of the application file upon payment of ce of Public Records upon payment of the olic may obtain a copy of: y document in the file of the pending c.C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. <u>tiffied</u> in a U.S. patent, a statutory invention incel extent application publication in
The member of the public may also be end the appropriate fee. Such copies must appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending application. For unpublished applications that are still application. For unpublished applications that are still application that has: (a) issued as patent application publication, or a Article 21(2), a member of the publication is incorporated to registration, a U.S. patent application accordance with PCT Article 21(2) originally filed.	at allow access of a copy of all of be purchased through the Offi <u>pending</u> , a member of the put ication as originally filed; or an <u>still pending</u> : <u>cation is claimed</u> under 35 U.S is a U.S. patent, or (b) publishe an intermational patent applica blic may obtain a copy of: the in the file of the pending applic by reference or otherwise iden tion publication, or an intermat c), a member of the public may	or part of the application file upon payment of ce of Public Records upon payment of the olic may obtain a copy of: y document in the file of the pending c.C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. <u>tiffied</u> in a U.S. patent, a statutory invention incel extent application publication in
The member of the public may also be e the appropriate fee. Such copies must l appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending appli- application. For unpublished applications that are s (1) If the benefit of the pending appli- application that has: (a) issued as patent application publication, or Article 21(2), a member of the pu- originally filed; or any document in registration, a U.S. patent applica accordance with PCT Article 21(2) originally filed.	The file of the pending application of the purchased through the Office pending, a member of the publication as originally filed; or an estill pending: cation is claimed under 35 U.S is a U.S. patent, or (b) publishes an international patent application are obtain a copy of: the normal publication, or an internation publication, or an internation, a member of the public may estimate the public may obtain a copy of the publication, or an internation publication, or an internation publication, or an internation of the public may estimate the public may est	C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. tified in a U.S. patent, a statutory invention ional patent application publication in obtain a copy of the pending application as tion. tified in a U.S. patent, a statutory invention ional patent application publication in obtain a copy of the pending application as the pending application as
The member of the public may also be end the appropriate fee. Such copies must appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending application. For unpublished applications that are so (1) If the benefit of the pending application that has: (a) issued as patent application publication, or a Article 21(2), a member of the publication, or is incorporated to registration, a U.S. patent application accordance with PCT Article 21(2) originally filed. Signature Set The	an anow access of a copy of all of the purchased through the Offi pending, a member of the publication as originally filed; or an setill pending: cation as originally filed; or an setill pending: cation is claimed under 35 U.S is a U.S. patent, or (b) publishes an intermational patent applica blic may obtain a copy of: the n the file of the pending application publication, or an intermately, a member of the public may re $N \ll S$	C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. Discussion accordance with PCT file contents; the pending application as ation. Discussion a copy of the pending application as atom.
The member of the public may also be e the appropriate fee. Such copies must l appropriate fee (37 CFR 1.19(b)). For published applications that are still the file contents; the pending applic application. For unpublished applications that are s (1) If the benefit of the pending applic application that has: (a) issued as patent application publication, or Article 21(2), a member of the pui originally filed; or any document in registration, a U.S. patent applica accordance with PCT Article 21(2) originally filed.	at allow access of a copy of all of be purchased through the Offi <u>pending</u> , a member of the pub- ication as originally filed; or an <u>still pending</u> : <u>cation is claimed</u> under 35 U.S is a U.S. patent, or (b) publishe an international patent applica blic may obtain a copy of: the in the file of the pending applica by reference or otherwise iden ition publication, or an internat t), a member of the public may	Provide the application file upon payment of the application file upon payment of the oblic records upon payment of the oblic may obtain a copy of: y document in the file of the pending S.C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. Iffied in a U.S. patent, a statutory invention in a obtain a copy of the pending application as attended as a copy of the pending application as a statutory invention in a copy of the pending application as a copy of the pending application a

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Under the Paperwork Reduction Act of 1995, no REQUEST FOR ACCESS TO	AN ABANDONED APPLIC	nt and Trademark Office; U.S. DEPAR INJERT OF COMMERCE ion of information unless it displays a valid OMB control number.
REQUEST FOR ACCESS TO	In re Application of	
Bring completed form to:		
File Information Unit, Room 2E04 2900 Crystal Drive	Application Number	777 Filed
Arlington, VA 22202-3514	07/7152	12 Configure
Telephone: (703) 308-2733	/	Paper No/
application, which is not writing the inc	a benefit is claimed, in the follow	le record of the above-identified ABANDONED prosecution Application (CPA) (37 CFR 1.53(d)) ing document (as shown in the attachment):
i i marina karatan karata	- Bublication No	, page, line,
United States Patent Number	64072K3column	, line, or
	, page, line	
web site (www.uspto.gov). Termin The member of the public may also the appropriate fee. Such copies r appropriate fee (37 CFR 1.19(b)). For published applications that ar the file contents; the pending application. For unpublished applications that (1) If the benefit of the pending application that has: (a) issue patent application publication Article 21(2), a member of th originally filed; or any docum (2) If the application is incorpor	and that allow documents a copy of all of nust be purchased through the Offi <u>e still pending</u> , a member of the pul- application as originally filed; or an <u>application is claimed</u> under 35 U.S. red as a U.S. patent, or (b) publishe n, or an international patent applica- he public may obtain a copy of: the nent in the file of the pending applica- ated by reference or otherwise ider	C. 119(e), 120, 121, or 365 in another d as a statutory invention registration, a U.S. tion publication in accordance with PCT file contents; the pending application as ation. <u>tified</u> in a U.S. patent, a statutory invention ional patent application publication in y obtain a copy of the pending application as
11 any	alla	9-1-06
- Street Street	gnature	Datě
WALKE	Clotom	- BECENFD
Typed or	r printed name	Approved by
Registration	Number, if applicable	SEP O(IIIIall)6
.115-	1777	Unit: File Information Unit
Telep	hone Number	
This collection of information is required by 37 CF USPTO to process) an application. Confidentiality complete, including gathering, preparing, and sub	R 1.11 and 1.14. The information is required is governed by 35 U.S.C. 122 and 37 CFR 1 mitting the completed application form to the complete this form and/or suggestions for redu	to obtain or retain a benefit by the public which is to file (and by t 11 and 1.14. This collection is estimated to take 12 minutes to USPTO. Time will vary depending upon the individual case. Any cing this burden, should be sent to the Chief Information Officer, VA 22313-1450. DO NOT SEND FEES OR COMPLETED Crystal Drive, Arlington, Virginia.



(12) United States Patent Carter et al.

(10) Patent No.: (45) Date of Patent:

US 6,407,213 B1 Jun. 18, 2002

- METHOD FOR MAKING HUMANIZED (54) ANTIBODIES
- Inventors: Paul J. Carter; Leonard G. Presta, (75) both of San Francisco, CA (US)
- Genentech, Inc., South San Francisco, Assignee: (73)CA (US)
- Subject to any disclaimer, the term of this Notice: (*) patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- 08/146,206 (21) Appl. No.:

(56)

- Jun. 15, 1992 (22) PCT Filed:
- PCT/US92/05126 (86) PCT No.: § 371 (c)(1), Nov. 17, 1993 (2), (4) Date:

Related U.S. Application Data

- Continuation-in-part of application No. 07/715,272, filed on Jun. 14, 1991, now abandoned. (63)
- (51) Int. Cl.⁷ C07K 16/00
- (52) U.S. Cl. 530/387.3; 435/69.6; 435/69.7; 435/70.21; 435/91; 536/23.53; 424/133.1
- (58) 320.1, 328; 536/23.53; 424/133.1; 530/387.3

References Cited

U.S. PATENT DOCUMENTS

4,816,567 A	3/1989	Cabilly et al.
4,845,198 A	7/1989	Urdal et al 530/388.22
4,043,190 1		Huston et al 530/387.3
5,132,405 A		Winter 530/389.3
5,225,539 A		Queen et al 530/387.3
5,530,101 A		Bendig et al 424/133.1
5,558,864 A		ACBOIL OF THE
5,585,089 A	12/1996	
5,677,171 A	10/1997	Hudziak et al
5,693,762 A	12/1997	Queen et al 530/387.2
5,714,350 A	2/1998	Co et al 435/69.6
5,772,997 A	6/1998	Hudziak et al 424/130.1
	10/1998	Carter et al 530/387.3
5,821,337 A	11/1008	Lowman et al 530/399
5,834,598 A	1/1000	Adair et al
5 850 205 A	1/1999	August of the information of the

FOREIGN PATENT DOCUMENTS

AU	85058/91	3/1992	C07K/15/12
EP	120694	10/1984	
EP	125023 AI	11/1984	
EP	0 239 400 *	9/1987	C12N/15/00
EP	323806 A1	7/1989	
EP	328404 A1	8/1989	A61K/39/395
EP	338745 A1	10/1989	
EP	365209 A2	4/1990	
EP	365997 A2	5/1990	•
EP	368684	5/1990	
EP	403156 A1	12/1990	
EP	438310 A2	7/1991	
EP	438312 A2	7/1991	
EP	440351 A2	8/1991	
EP	0 460 167 B1	12/1991	
EP	0 519 596 Al	12/1992	

EP	0 592 106 A1	4/1994	
EP	0 620 276	10/1994	
EP	682040 A1	11/1995	
EP	451216 B1	1/1996	C12P/21/08
EP	432249 B1	9/1996	
GB	2 188941	10/1987	
wo	WO 87/02671	5/1987	
wo	WO 88/09344	12/1988	
wo	WO 89/01783	3/1989	
wo	WO 89/06692	7/1989	
wo	WO 89/09622	10/1989	
WO	WO 90/07861	7/1990	
wo	90/07861	 7/1990 	C12P/21/00
wo	WO 91/07492	5/1991	
wo	WO 91/07500	5/1991	
wo	WO 91/09966	7/1991	C12P/21/08
wo	WO 91/09968	7/1991	C12P/21/08
wo	WO 91/09967	11/1991	
wo	WO 92/01047	1/1992	· ·
wo	WO 92/04380	3/1992	
wo	WO 92/04381	3/1992	
wo	WO 92/05274	4/1992	
wo	WO 92/11383	7/1992	
wo	WO 92/11018	9/1992	A61K/35/14
WO	WO 92/15683	9/1992	· · · · ·
wo	WO 92/16562	10/1992	
wo	WO 92/22653	12/1992	
ŵO	WO 93/02191	2/1993	
wo	94/11509	5/1994	
wo	WO 94/12214	6/1994	

OTHER PUBLICATIONS

Riechmann et al. [Nature 332:323-327 (1988)].*

Queen et al. [Proc. Natl. Acad. Sci. 86:10029-10033 (1989)].*

Roitt [Immunology, published 1985, by Gower Medical Publishing Ltd. (London, England) p. 5.5].*

Tramontano et al. [J. Mol. Biol. 215:175-182 (1990)].*

"Biosym Technologies" in New Products, Chemical Design Automation 3 (Dec. 1988).

"Polygen Corporation" in New Products, Chemical Design Automation 3 (Nov. 1988).

Adair et al., "Humanization of the murine anti-human CD3 monoclonal antibody OKT3" Hum. Antibod. Hybridomas

Sci41-47 (1994). Chothia et al., "Principles of protein-protein recognition" Nature 256:705-708 (1975).

Chothia et al., "Transmission of conformational change in insulin" Nature 302:500-505 (1983).

Corti et al., "Idiotope Determining Regions of a Mouse Monoclonal Antibody and Its Humanized Versions" J. Mol. Biol. 235:53-60 (1994).

(List continued on next page.)

Primary Examiner-Anthony C. Caputa Assistant Examiner-Minh-Tam Davis

(74) Attorney, Agent, or Firm-Wendy M. Lee

ABSTRACT (57)

Variant immunoglobulins, particularly humanized antibody polypeptides are provided, along with methods for their preparation and use. Consensus immunoglobulin sequences and structural models are also provided.

82 Claims, 9 Drawing Sheets