

PDR®
59
EDITION
2005

PHYSICIANS' DESK REFERENCE®

Executive Vice President, PDR: David Duplay

Vice President, Sales and Marketing: Dikran N. Barsamian
Senior Director, Pharmaceutical Sales: Anthony Sorce
National Account Manager: Marion Reid, RPh
Senior Account Managers: Frank Karkowsky, Suzanne E. Yarrow, RN
Account Managers: Marjorie A. Jaxel, Kevin McGlynn, Elaine Musco, Lois Smith, Eileen Sullivan, Richard Zwickel
Senior Director, Brand and Product Management: Valerie E. Berger
Director, Brand and Product Management: Carmen Mazzatta
Associate Product Managers: Michael Casale, Andrea Colavecchio
Senior Director, Publishing Sales and Marketing: Michael Bennett
Director of Trade Sales: Bill Gaffney
Associate Director of Marketing: Jennifer M. Fronzaglia
Senior Marketing Manager: Kim Marich
Direct Mail Manager: Lorraine M. Loening
Manager of Marketing Analysis: Dina A. Maeder
Promotion Manager: Linda Levine
Vice President, Regulatory Affairs: Mukesh Mehta, RPh
Vice President, PDR Services: Brian Holland
Director of PDR Operations: Jeffrey D. Schaefer
Director of Operations: Robert Klein
Clinical Content Operations Manager: Thomas Fleming, PharmD

Manager, Editorial Services: Bette LaGow
Drug Information Specialists: Min Ko, PharmD; Greg Tallis, RPh
Project Editors: Neil Chesanow, Harris Fleming
Senior Editor: Lori Murray
Production Editor: Gwynned L. Kelly
Manager, Production Purchasing: Thomas Westburgh
Production Manager: Gayle Graizzaro
Production Specialist: Christina Klingler
Senior Production Coordinator: Gianna Caradonna
Production Coordinator: Yasmin Hernández
Senior Index Editors: Noel Deloughery, Shannon Reilly
Format Editor: Michelle S. Guzman
Traffic Assistant: Kim Condon
PDR Sales Coordinators: Nick W. Clark, Gary Lew
Production Design Supervisor: Adeline Rich
Senior Electronic Publishing Designer: Livio Udina
Electronic Publishing Designers: Bryan C. Dix, Rosalia Sberna
Production Associate: Joan K. Akerlind
Digital Imaging Manager: Christopher Husted
Digital Imaging Coordinator: Michael Labryere
Finance Director: Mark S. Ritchin
Director of Client Services: Stephanie Struble

THOMSON

PDR

Copyright © 2005 and published by Thomson PDR at Montvale, NJ 07645-1742. All rights reserved. None of the content of this publication may be reproduced, stored in a retrieval system, resold, redistributed, or transmitted in any form or by any means (electronic, mechanical, photocopying, recording, or otherwise) without the prior written permission of the publisher. Physicians' Desk Reference®, PDR®, Pocket PDR®, PDR Family Guide to Prescription Drugs®, PDR Family Guide to Women's Health and Prescription Drugs®, and PDR Family Guide to Nutrition and Health® are registered trademarks used herein under license. PDR® for Ophthalmic Medicines, PDR® for Nonprescription Drugs and Dietary Supplements, PDR® Companion Guide, PDR® Pharmacopoeia, PDR® for Herbal Medicines, PDR® for Nutritional Supplements, PDR® Medical Dictionary, PDR® Nurse's Drug Handbook, PDR® Nurse's Dictionary, PDR® Family Guide Encyclopedia of Medical Care, PDR® Family Guide to Natural Medicines and Healing Therapies, PDR® Family Guide to Common Ailments, PDR® Family Guide to Over-the-Counter Drugs, PDR® Family Guide to Nutritional Supplements, and PDR® Electronic Library are trademarks used herein under license.

Officers of Thomson Healthcare, Inc.: *President and Chief Executive Officer:* Robert Cullen; *Chief Financial Officer:* Paul Hilger; *Chief Technology Officer:* Fred Lauber; *Executive Vice President, Medical Education:* Jeff MacDonald; *Executive Vice President, Micromedex:* Jeff Reihl; *Executive Vice President, PDR:* David Duplay; *Senior Vice President, Business Development:* Robert Christopher; *Senior Vice President, Marketing:* Timothy Murray; *Vice President, Human Resources:* Pamela M. Bilash

ISBN: 1-56363-497-X

**DOCKET
ALARM**

Find authenticated court documents without watermarks at docketalarm.com.

CONTENTS

Manufacturers' Index (White Pages) **1**

Section 1

Lists all pharmaceutical manufacturers participating in *PHYSICIANS' DESK REFERENCE*. Includes addresses, phone numbers, and emergency contacts. Shows each manufacturer's products and the page number of those described in *PDR*.

Brand and Generic Name Index (Pink Pages) **101**

Section 2

Gives the page number of each product by brand and generic name.

Product Category Index (Blue Pages) **201**

Section 3

Lists all fully described products by prescribing category. An overview of the headings appears on pages 201 and 202.

Key to Controlled Substances Categories217

Gives the definition of each category and the prescribing limitations that apply.

Key to FDA Use-in-Pregnancy Ratings217

Provides at-a-glance description of each risk/benefit rating.

U.S. Food and Drug Administration Telephone Directory.....218

Gives numbers of key reporting programs and information services.

Poison Control Centers219

A national directory arranged alphabetically by state and city.

Vaccine Adverse Event Reporting Form225

Includes master copy and instructions for completion.

Product Identification Guide (Gray Pages) **301**

Section 4

Presents full-color, actual-size photos of tablets and capsules, plus pictures of a variety of other dosage forms and packages. Arranged alphabetically by manufacturer.

Product Information (White Pages) **401**

Section 5

The main section of the book. Includes entries for some 3,000 pharmaceuticals. Listings are arranged alphabetically by manufacturer.

Diagnostic Product Information **3437**

Section 6

Gives usage guidelines for a variety of diagnostic agents. Arranged alphabetically by manufacturer.

FDA MedWatch Form.....3441

Includes master copy and instructions for completion.

Treatment—Standard symptomatic treatment may be undertaken if overdose occurs. If the patient develops a dramatic increase in blood pressure, 5 to 10 mg of phenolamine mesylate has been shown to be effective in lowering blood pressure for the short time that control would be needed. It is unknown whether GlucaGen® is dialyzable, but such a procedure is unlikely to provide any benefit given the short half-life and nature of the symptoms of overdose.

DOSE AND ADMINISTRATION

GlucaGen® should be reconstituted with 1 ml of Sterile Water for Reconstitution (if supplied) or with 1 mL Sterile Water for Injection, USP.

Using the syringe, withdraw all of the Sterile Water for Reconstitution (if supplied) or 1 mL Sterile Water for Injection, USP and inject into the GlucaGen® vial. Roll the vial gently until powder is completely dissolved and no particles remain in the fluid. The reconstituted fluid should be clear and of water-like consistency. The reconstituted GlucaGen® gives a concentration of approximately 1 mg/ml GlucaGen®. The reconstituted GlucaGen® should be used immediately after reconstitution. Discard any unused portion.

For the treatment of hypoglycemia: For adults and for pediatric patients weighing 55 lb (25 kg) or more, administer 1 mg by subcutaneous, intramuscular, or intravenous injection.^{1,2} According to the literature, ½ adult dose (0.5 mg) is recommended for pediatric patients weighing less than 55 lb (25 kg) or younger than 6–8 years old.^{2,3,4,5,6} Emergency assistance should be sought if the patient fails to respond within 15 minutes after subcutaneous or intramuscular injection of glucagon. The glucagon injection may be repeated while waiting for emergency assistance.¹ Intravenous glucose MUST be administered if the patient fails to respond to glucagon. When the patient has responded to the treatment, give oral carbohydrate to restore the liver glycogen and prevent recurrence of hypoglycemia.

Directions for Use as a Diagnostic Aid: Reconstitute as indicated above. Discard any unused portion. When the diagnostic procedure is over, give oral carbohydrate to restore the liver glycogen and prevent occurrence of secondary hypoglycemia.

Time of maximal glucose concentration
 Intravenous: 5 to 20 minutes
 Intramuscular: 30 minutes
 Subcutaneous: 30 to 45 minutes

Time for GI smooth muscle relaxation¹
 Intravenous: 0.25 to 2 mg (IU)—45 seconds.
 Intramuscular:
 1 mg (IU)—8 to 10 minutes
 1 mg (IU)—4 to 7 minutes

Duration of action—
 Hyperglycemic action—60 to 90 minutes
 Smooth muscle relaxation—

Intravenous:
 0.25 to 0.5 mg (IU)—9 to 17 minutes
 1 mg (IU)—22 to 25 minutes
Intramuscular:
 1 mg (IU)—12 to 27 minutes
 2 mg (IU)—21 to 32 minutes

Stability and storage
Before Reconstitution: The GlucaGen® package may be stored up to 24 months at controlled room temperature 20° to 25°C (68° to 77°F) prior to reconstitution. Avoid freezing and protect from light. GlucaGen® should not be used after the expiry date on the vials.

After Reconstitution: Reconstituted GlucaGen® should be used immediately. Discard any unused portion. If the solution shows any sign of gel formation or particles, it should be discarded.

HOW SUPPLIED

The GlucaGen® Diagnostic Kit includes:
 1 vial containing 1 mg (1 IU) GlucaGen® [glucagon (rDNA origin) for injection]
 1 vial containing 1 ml Sterile Water for Reconstitution
 NDC 55390-004-01

The GlucaGen® 10-pack includes:
 10x1 vial containing 1 mg (1 IU) GlucaGen® [glucagon (rDNA origin) for injection]
 NDC 55390-004-10
 Edition March 2001

REFERENCES

1. Drug Information for the Health Care Professional. 17th ed. Rockville, Maryland: The United States Pharmacopoeial Convention, Inc; 1997; Vol. 1: 1A:1516-1518.
2. Gibbs et al: Use of Glucagon to terminate insulin reactions in diabetic children. *Nbr Med J* 1958;43:56-57.
3. Carson MJ, Koch R. Clinical studies with glucagon in children. *J Pediatr* 1955;47:161-170.
4. Shipp JC, et al: Treatment of insulin hypoglycemia in diabetic campers. *Diabetes* 1964; 13:645-648.
5. Aman J, Wranne I. Hypoglycemia in childhood diabetes II: Effect of subcutaneous or intramuscular injection of different doses of glucagon. *Acta Paediatr Scand* 1988;77: 545-553.
6. Aynsley-Green AS, Eyre JA, and Soitesz G. Hypoglycemia

Berlex, Inc.
 6 WEST BELT
 WAYNE, NJ 07470
 www.Berlex.com

Direct Inquiries to:
 1-(888) BERLEX-4

BETASERON®
 [bay-ta-seer-on]
 Interferon beta-1b

DESCRIPTION

Betaseron® (Interferon beta-1b) is a purified, sterile, lyophilized protein product produced by recombinant DNA techniques. Interferon beta-1b is manufactured by bacterial fermentation of a strain of *Escherichia coli* that bears a genetically engineered plasmid containing the gene for human interferon beta_{ser17}. The native gene was obtained from human fibroblasts and altered in a way that substitutes serine for the cystine residue found at position 17. Interferon beta-1b has 165 amino acids and an approximate molecular weight of 18,500 daltons. It does not include the carbohydrate side chains found in the natural material.

The specific activity of Betaseron is approximately 32 million international units (IU)/mg Interferon beta-1b. Each vial contains 0.3 mg of Interferon beta-1b. The unit measurement is derived by comparing the antiviral activity of the product to the World Health Organization (WHO) reference standard of recombinant human interferon beta. Mannitol, USP and Albumin (Human), USP (15 mg each/vial) are added as stabilizers.

Lyophilized Betaseron is a sterile, white to off-white powder, for subcutaneous injection after reconstitution with the diluent supplied (Sodium Chloride, 0.54% Solution).

CLINICAL PHARMACOLOGY

General

Interferons (IFNs) are a family of naturally occurring proteins, produced by eukaryotic cells in response to viral infection and other biologic agents. Three major groups of interferons have been distinguished: alpha, beta, and gamma. Interferons alpha and beta comprise the Type I interferons and interferon gamma is a Type II interferon. Type I interferons have considerably overlapping but also distinct biologic activities. The bioactivities of IFNs are mediated by their interactions with specific receptors found on the surfaces of human cells. Differences in bioactivities induced by IFNs likely reflect divergences in the signal transduction process induced by IFN-receptor binding.

Biologic Activities

The mechanism of action of Interferon beta-1b in patients with multiple sclerosis is unknown. Interferon beta-1b receptor binding induces the expression of proteins that are responsible for the pleiotropic bioactivities of Interferon beta-1b. A number of these proteins (including neopterin, β₂-microglobulin, MxA protein, and IL-10) have been measured in blood fractions from Betaseron-treated patients and Betaseron-treated healthy volunteers. Immunomodulatory effects of Interferon beta-1b include the enhancement of suppressor T cell activity, reduction of pro-inflammatory cytokine production, down-regulation of antigen presentation, and inhibition of lymphocyte trafficking into the central nervous system. It is not known if these effects play an important role in the observed clinical activity of Betaseron in multiple sclerosis (MS).

Pharmacokinetics

Because serum concentrations of Interferon beta-1b are low or not detectable following subcutaneous administration of 0.25 mg or less of Betaseron, pharmacokinetic information in patients with MS receiving the recommended dose of Betaseron is not available. Following single and multiple daily subcutaneous administrations of 0.5 mg Betaseron to healthy volunteers (N=12), serum Interferon beta-1b concentrations were generally below 100 IU/mL. Peak serum Interferon beta-1b concentrations occurred between one to eight hours, with a mean peak serum interferon concentration of 40 IU/mL. Bioavailability, based on a total dose of 0.5 mg Betaseron given as two subcutaneous injections at different sites, was approximately 50%.

After intravenous administration of Betaseron (0.006 mg to 2.0 mg), similar pharmacokinetic profiles were obtained from healthy volunteers (N=12) and from patients with diseases other than MS (N=142). In patients receiving single intravenous doses up to 2.0 mg, increases in serum concentrations were dose proportional. Mean serum clearance values ranged from 9.4 mL/min•kg⁻¹ to 26.9 mL/min•kg⁻¹ and were independent of dose. Mean terminal elimination half-life values ranged from 8.0 minutes to 4.3 hours and mean steady-state volume of distribution values ranged from 0.25 L/kg to 2.88 L/kg. Three-times-a-week intravenous dosing for two weeks resulted in no accumulation of Interferon beta-1b in sera of patients. Pharmacokinetic parameters after single and multiple intravenous doses of Betaseron were comparable.

Betaseron dose. Biologic response marker levels peaked between 40 and 124 hours and remained elevated above baseline throughout the seven-day (168-hour) study. The relationship between serum Interferon beta-1b levels or induced biologic response marker levels and the clinical effects of Interferon beta-1b in multiple sclerosis is unknown.

CLINICAL STUDIES

The safety and efficacy of Betaseron have been assessed in three multicenter trials. Study 1 evaluated Betaseron in relapsing-remitting MS (RRMS) patients and Studies 2 and 3 assessed Betaseron in secondary progressive MS (SPMS) patients.

The effectiveness of Betaseron in relapsing-remitting MS (Study 1) was evaluated in a double blind, multicentric, randomized, parallel, placebo controlled clinical investigation of two years duration. The study enrolled MS patients, aged 18 to 50, who were ambulatory (EDSS of ≤ 5.5), exhibited a relapsing-remitting clinical course, met Poser's criteria⁷ for clinically definite and/or laboratory supported definite MS and had experienced at least two exacerbations over two years preceding the trial without exacerbation in the preceding month. Patients who had received prior immunosuppressant therapy were excluded.

An exacerbation was defined as the appearance of a new clinical sign/symptom or the clinical worsening of a previous sign/symptom (one that had been stable for at least 30 days) that persisted for a minimum of 24 hours.

Patients selected for study were randomized to treatment with either placebo (N=123), 0.05 mg of Betaseron (N=125), or 0.25 mg of Betaseron (N=124) self-administered subcutaneously every other day. Outcome based on the 372 randomized patients was evaluated after two years.

Patients who required more than three 28-day courses of corticosteroids were removed from the study. Minor analgesics (acetaminophen, codeine), antidepressants, and oral baclofen were allowed ad libitum, but chronic nonsteroidal anti-inflammatory drug (NSAID) use was not allowed.


The primary protocol-defined outcome measures were 1) frequency of exacerbations per patient and 2) proportion of exacerbation free patients. A number of secondary clinical and magnetic resonance imaging (MRI) measures were also employed. All patients underwent annual T2 MRI imaging and a subset of 52 patients at one site had MRIs performed every six weeks for assessment of new or expanding lesions. The study results are shown in Table 1.

[See table 1 at top of next page]
 Of the 372 RRMS patients randomized, 72 (19%) failed to complete two full years in their assigned treatments.

Over the two-year period, there were 25 MS-related hospitalizations in the 0.25 mg Betaseron-treated group compared to 48 hospitalizations in the placebo group. In comparison, non-MS hospitalizations were evenly distributed among the groups, with 16 in the 0.25 mg Betaseron group and 15 in the placebo group. The average number of days of MS-related steroid use was 41 days in the 0.25 mg Betaseron group and 55 days in the placebo group (p=0.004).

MRI data were also analyzed for patients in this study. A frequency distribution of the observed percent changes in MRI area at the end of two years was obtained by grouping the percentages in successive intervals of equal width. Figure 1 displays a histogram of the proportions of patients, which fell into each of these intervals. The median percent change in MRI area for the 0.25 mg group was -1.1%, which was significantly smaller than the 16.5% observed for the placebo group (p=0.0001).

Distribution of Change in MRI Area



In an evaluation of frequent MRI scans (every six weeks) on 52 patients at one site, the percent of scans with new or expanding lesions was 29% in the placebo group and 6% in the 0.25 mg treatment group (p=0.006).

The exact relationship between MRI findings and clinical status of patients is unknown. Changes in lesion area often do not correlate with changes in disability progression. The prognostic significance of the MRI findings in this study has not been evaluated.

Continued on next page

Information on Berlex products (appearing here) is based on the most current information available at the time

Betaseron—Cont.

Studies 2 and 3 were multicenter, randomized, double-blind, placebo controlled trials conducted to assess the effect of Betaseron in patients with SPMS. Study 2 was conducted in Europe and Study 3 was conducted in North America. Both studies enrolled patients with clinically definite or laboratory-supported MS in the secondary progressive phase, and who had evidence of disability progression (both Study 2 and 3) or two relapses (Study 2 only) within the previous two years. Baseline Kurtzke expanded disability status scale (EDSS) scores ranged from 3.0 to 6.5.² Patients in Study 2 were randomized to receive Betaseron 0.25 mg (n=360) or placebo (n=358). Patients in Study 3 were randomized to Betaseron 0.25 mg (n=317). Betaseron 0.16 mg/m² of body surface area (n=314, mean assigned dose 0.30 mg), or placebo (n=308). Test agents were administered subcutaneously, every other day for three years.

The primary outcome measure was progression of disability, defined as a 1.0 point increase in the EDSS score, or a 0.5 point increase for patients with baseline EDSS \geq 6.0. In Study 2, time to progression in EDSS was longer in the Betaseron treatment group (p=0.005), with estimated annualized rates of progression of 16% and 19% in the Betaseron and placebo groups, respectively. In Study 3, the rates of progression did not differ significantly between treatment groups, with estimated annualized rates of progression of 12%, 14%, and 12% in the Betaseron fixed dose, surface area-adjusted dose, and placebo groups, respectively. Multiple analyses, including covariate and subset analyses based on sex, age, disease duration, clinical disease activity prior to study enrollment, MRI measures at baseline and early changes in MRI following treatment were evaluated in order to interpret the discordant study results. No demographic or disease-related factors enabled identification of a patient subset where Betaseron treatment was predictably associated with delayed progression of disability. In Studies 2 and 3, like Study 1, a statistically significant decrease in the incidence of relapses associated with Betaseron treatment was demonstrated. In Study 2, the mean annual relapse rates were 0.42 and 0.63 in the Betaseron and placebo groups, respectively (p<0.001). In Study 3, the mean annual relapse rates were 0.16, 0.20, and 0.28, for the fixed dose, surface area-adjusted dose, and placebo groups, respectively (p<0.02).

MRI endpoints in both Study 2 and Study 3 showed lesser increases in T2 MRI lesion area and decreased number of active MRI lesions in patients in the Betaseron groups. The exact relationship between MRI findings and the clinical status of patients is unknown. Changes in MRI findings often do not correlate with changes in disability progression. The prognostic significance of the MRI findings in these studies is not known.

Safety and efficacy of treatment with Betaseron beyond three years are not known.

INDICATIONS AND USAGE

Betaseron (Interferon beta-1b) is indicated for the treatment of relapsing forms of multiple sclerosis to reduce the frequency of clinical exacerbations.

CONTRAINDICATIONS

Betaseron is contraindicated in patients with a history of hypersensitivity to natural or recombinant interferon beta, Albumin (Human), USP, or any other component of the formulation.

WARNINGS**Depression and Suicide***

Betaseron (Interferon beta-1b) should be used with caution in patients with depression, a condition that is common in people with multiple sclerosis. Depression and suicide have been reported to occur with increased frequency in patients receiving interferon compounds, including Betaseron. Patients treated with Betaseron should be advised to report immediately any symptoms of depression and/or suicidal ideation to their prescribing physicians. If a patient develops depression, cessation of Betaseron therapy should be considered.

In the three randomized controlled studies there were three suicides and eight suicide attempts among the 1240 patients in the Betaseron treated groups compared to one suicide and four suicide attempts among the 789 patients in the placebo groups.

Injection Site Necrosis

Injection site necrosis (ISN) has been reported in 5% of patients in controlled clinical trials (see **ADVERSE REACTIONS**). Typically, injection site necrosis occurs within the first four months of therapy, although post-marketing reports have been received of ISN occurring over one year after initiation of therapy. Necrosis may occur at a single or multiple injection sites. The necrotic lesions are typically three cm or less in diameter, but larger areas have been reported. Generally the necrosis has extended only to subcutaneous fat. However, there are also reports of necrosis extending to and including fascia overlying muscle. In some lesions where biopsy results are available, vasculitis has been reported. For some lesions debridement and, infrequently, skin grafting have been required.

TABLE 1
Two Year RRMS Study Results
Primary and Secondary Clinical Outcomes

Efficacy Parameters	Treatment Groups			Statistical Comparisons p-value			
	Placebo (N=123)	0.05 mg (N=125)	0.25 mg (N=124)	Placebo vs 0.05 mg	0.05 mg vs 0.25 mg	Placebo vs 0.25 mg	
Primary End Points							
Annual exacerbation rate	1.31	1.14	0.90	0.005	0.113	0.0001	
Proportion of exacerbation-free patients†	16%	18%	25%	0.609	0.288	0.034	
Exacerbation frequency per patient	0† 1 2 3 4 ≥5	20 32 20 15 15 21	22 31 28 15 7 16	29 39 17 14 9 8	0.151	0.077	0.001
Secondary Endpoints††							
Median number of months to first on-study exacerbation	5	6	9	0.299	0.097	0.010	
Rate of moderate or severe exacerbations per year	0.47	0.29	0.23	0.020	0.257	0.001	
Mean number of moderate or severe exacerbation days per patient	44.1	33.2	19.5	0.229	0.064	0.001	
Mean change in EDSS score‡ at endpoint	0.21	0.21	-0.07	0.995	0.108	0.144	
Mean change in Scripps score‡‡ at endpoint	-0.53	-0.50	0.66	0.641	0.051	0.128	
Median duration in days per exacerbation	36	33	35.5	ND	ND	ND	
% change in mean MRI lesion area at endpoint	21.4%	9.8%	-0.9%	0.015	0.019	0.0001	

ND: Not done

† 14 exacerbation free patients (0 from placebo, six from 0.05 mg, and eight from 0.25 mg) dropped out of the study before completing six months of therapy. These patients are excluded from this analysis.

†† Sequelae and Functional Neurologic Status, both required by protocol, were not analyzed individually but are included as a function of the EDSS.

‡ EDSS scores range from 1-10, with higher scores reflecting greater disability.

‡‡ Scripps neurologic rating scores range from 0-100, with smaller scores reflecting greater disability.

Some patients have experienced healing of necrotic skin lesions while Betaseron therapy continued; others have not. Whether to discontinue therapy following a single site of necrosis is dependent on the extent of necrosis. For patients who continue therapy with Betaseron after injection site necrosis has occurred, Betaseron should not be administered into the affected area until it is fully healed. If multiple lesions occur, therapy should be discontinued until healing occurs.

Patient understanding and use of aseptic self-injection techniques and procedures should be periodically reevaluated, particularly if injection site necrosis has occurred.

Anaphylaxis

Anaphylaxis has been reported as a rare complication of Betaseron use. Other allergic reactions have included dyspnea, bronchospasm, tongue edema, skin rash and urticaria (see **ADVERSE REACTIONS**).

Albumin (Human), USP

This product contains albumin, a derivative of human blood. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of Creutzfeldt-Jakob disease (CJD) also is considered extremely remote. No cases of transmission of viral diseases or CJD have ever been identified for albumin.

PRECAUTIONS**Information for Patients**

All patients should be instructed to carefully read the supplied Betaseron Medication Guide. Patients should be cautioned not to change the dose or schedule of administration without medical consultation.

Patients should be made aware that serious adverse reactions during the use of Betaseron have been reported, including depression and suicidal ideation, injection site necrosis, and anaphylaxis (see **WARNINGS**). Patients should be advised of the symptoms of depression or suicidal ideation and be told to report them immediately to their physician. Patients should also be advised of the symptoms of allergic reactions and anaphylaxis. Patients should be advised to promptly report any break in the skin, which may be associated with blue-black discoloration, swelling, or drainage of fluid from the injection site, prior to continuing their Betaseron therapy.

Patients should be informed that flu-like symptoms are common following initiation of therapy with Betaseron. In

Female patients should be cautioned about the abortifacient potential of Betaseron (see **PRECAUTIONS, Pregnancy, Teratogenic Effects**).

Instruction on Self-injection Technique and Procedures

Patients should be instructed in the use of aseptic technique when administering Betaseron. Appropriate instruction for reconstitution of Betaseron and self-injection should be provided, including careful review of the Betaseron Medication Guide. The first injection should be performed under the supervision of an appropriately qualified health care professional.

Patients should be cautioned against the re-use of needles or syringes and instructed in safe disposal procedures. A puncture resistant container for disposal of used needles and syringes should be supplied to the patient along with instructions for safe disposal of full containers.

Patients should be advised of the importance of rotating areas of injection with each dose, to minimize the likelihood of severe injection site reactions, including necrosis or localized infection; (see **Picking an Injection Site** section of the Medication Guide).

Laboratory Tests

In addition to those laboratory tests normally required for monitoring patients with multiple sclerosis, complete blood and differential white blood cell counts, platelet counts and blood chemistries, including liver function tests, are recommended at regular intervals (one, three, and six months) following introduction of Betaseron therapy, and then periodically thereafter in the absence of clinical symptoms. Thyroid function tests are recommended every six months in patients with a history of thyroid dysfunction or as clinically indicated. Patients with myelosuppression may require more intensive monitoring of complete blood cell counts, with differential and platelet counts.

Drug Interactions

No formal drug interaction studies have been conducted with Betaseron. In the placebo controlled studies in MS, corticosteroids or ACTH were administered for treatment of relapses for periods of up to 28 days in patients (N=664) receiving Betaseron.

Carcinogenesis, Mutagenesis, and Impairment of Fertility
Carcinogenesis: Interferon beta-1b has not been tested for its carcinogenic potential in animals.

Mutagenesis: Betaseron was not mutagenic when assayed

...sult in increased transformation frequency in an *in vitro* model of tumor transformation.
Impairment of fertility: Studies in normally cycling, female rhesus monkeys at doses up to 0.33 mg/kg/day (32 times the recommended human dose based on body surface area, body surface dose based on 70 kg female) had no apparent adverse effects on either menstrual cycle duration or associated hormonal profiles (progesterone and estradiol) when administered over three consecutive menstrual cycles. The validity of extrapolating doses used in animal studies to human doses is not known. Effects of Betaseron on normally cycling human females are not known.

Pregnancy - Teratogenic Effects
Pregnancy Category C: Betaseron was not teratogenic at doses up to 0.42 mg/kg/day when given to pregnant female rhesus monkeys on gestation days 20 to 70. However, a dose related abortifacient activity was observed in these monkeys when Interferon beta-1b was administered at doses ranging from 0.028 mg/kg/day to 0.42 mg/kg/day (2.8 to 40 times the recommended human dose based on body surface area comparison). The validity of extrapolating doses used in animal studies to human doses is not known. Lower doses were not studied in monkeys. Spontaneous abortions while on treatment were reported in patients (n=4) who participated in the Betaseron RRMS clinical trial. Betaseron given to rhesus monkeys on gestation days 20 to 70 did not cause teratogenic effects; however, it is not known if teratogenic effects exist in humans. There are no adequate and well-controlled studies in pregnant women. If the patient becomes pregnant or plans to become pregnant while taking Betaseron, the patient should be apprised of the potential hazard to the fetus and it should be recommended that the patient discontinue therapy.

Nursing Mothers
 It is not known whether Betaseron is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from Betaseron, a decision should be made to either discontinue nursing or discontinue the drug, taking into account the importance of drug to the mother.

Pediatric Use
 Safety and efficacy in pediatric patients have not been established.

Geriatric Use
 (Clinical studies of Betaseron did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently than younger patients.)

ADVERSE REACTIONS

In all studies, the most serious adverse reactions with Betaseron were depression, suicidal ideation and injection site necrosis (see **WARNINGS**). The incidence of depression of any severity was approximately 34% in both Betaseron-treated patients and placebo-treated patients. Anaphylaxis and other allergic reactions have been reported in patients using Betaseron (see **WARNINGS**). The most commonly reported adverse reactions were lymphopenia (lymphocytes <1500/mm³), injection site reaction, asthenia, flu-like symptom complex, headache, and pain. The most frequently reported adverse reactions resulting in clinical intervention (e.g., discontinuation of Betaseron, adjustment in dosage, or the need for concomitant medication to treat an adverse reaction symptom) were depression, flu-like symptom complex, injection site reactions, leukopenia, increased liver enzymes, asthenia, hypertonia, and myasthenia.

Because clinical trials are conducted under widely varying conditions and over varying lengths of time, adverse reaction rates observed in the clinical trials of Betaseron cannot be directly compared to rates in clinical trials of other drugs, and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates.

The data described below reflect exposure to Betaseron in the three placebo controlled trials of 1115 patients with MS treated with 0.25 mg or 0.16 mg/m², including 1041 exposed for greater than one year. The population encompassed an age range from 18-65 years. Sixty-five percent (65%) of the patients were female. The percentages of Caucasian, Black, Asian, and Hispanic patients were 94.0%, 4.3%, 0.2%, and 0.8%, respectively.

The safety profiles for Betaseron-treated patients with SPMS and RRMS were similar. Clinical experience with Betaseron in other populations (patients with cancer, HIV positive patients, etc.) provides additional data regarding adverse reactions; however, experience in non-MS populations may not be fully applicable to the MS population.

Injection Site Reactions
 In three controlled clinical trials, injection site reactions occurred in 86% of patients receiving Betaseron with injection site necrosis in 5%. Inflammation (53%), pain (18%), hypersensitivity (3%), necrosis (5%), mass (2%), edema (3%), and non-specific reactions were significantly associated with Betaseron treatment (see **WARNINGS** and **PRECAUTIONS**). The incidence of injection site reactions tended to decrease over time, with approximately 76% of patients experiencing the event during the first three months of treatment.

like symptom complex at the end of the studies. For patients who experienced a flu-like symptom complex in Study 1, the median duration was 7.6 days.

Laboratory Abnormalities

In the three clinical trials, leukopenia was reported in 18% and 5% of patients in Betaseron- and placebo-treated groups, respectively. No patients were withdrawn or dose reduced for neutropenia in Study 1. Three percent (3%) of patients in Studies 2 and 3 experienced leukopenia and were dose-reduced. Other laboratory abnormalities included SGPT greater than five times baseline value (10%), and SGOT greater than five times baseline value (3%). In Study 1, two patients were dose reduced for increased liver enzymes; one continued on treatment and one was ultimately withdrawn. In Studies 2 and 3, 1.5% of Betaseron patients were dose-reduced or interrupted treatment for increased liver enzymes. Three (0.3%) patients were withdrawn from treatment with Betaseron for any laboratory abnormality including two (0.2%) patients following dose reduction (see **PRECAUTIONS, Laboratory Tests**).

Menstrual Irregularities

In the three clinical trials, 82 (14%) of the 577 pre-menopausal females treated with Betaseron and 74 (18%) of the 405 pre-menopausal females treated with placebo reported menstrual disorders. One event was reported as severe, all other reports were mild to moderate severity. No patients withdrew from the studies due to menstrual irregularities. Table 2 enumerates adverse events and laboratory abnormalities that occurred among all patients treated with 0.25 mg or 0.16 mg/m² Betaseron every other day for periods of up to three years in the controlled trials at an incidence that was at least 2% more than that observed in the placebo patients.

TABLE 2
Adverse Reactions and Laboratory Abnormalities

Adverse Reaction	Placebo (n=789)	Betaseron (n=1115)
Body as a Whole		
Injection site reaction	29%	85%
Asthenia	54%	61%
Flu-like symptom complex	41%	60%
Headache	48%	57%
Pain	42%	51%
Fever	22%	36%
Chills	11%	25%
Abdominal pain	13%	19%
Chest pain	7%	11%
Malaise	4%	8%
Injection site necrosis	0%	5%
Cardiovascular System		
Peripheral edema	12%	15%
Vasodilation	6%	8%
Hypertension	4%	7%
Peripheral vascular disorder	4%	6%
Palpitation	2%	4%
Tachycardia	2%	4%
Digestive System		
Nausea	25%	27%
Constipation	18%	20%
Diarrhea	16%	19%
Dyspepsia	12%	14%
Hemic and Lymphatic System		
Lymphocytes < 1500/mm ³	70%	88%
ANC < 1500/mm ³	5%	14%
WBC < 3000/mm ³	4%	14%
Lymphadenopathy	4%	8%

Weight gain	5%	7%
Musculoskeletal System		
Myasthenia	43%	46%
Arthralgia	29%	31%
Myalgia	16%	27%
Leg cramps	2%	4%
Nervous System		
Hypertonia	40%	50%
Dizziness	21%	24%
Insomnia	19%	24%
Incoordination	18%	21%
Anxiety	8%	10%
Nervousness	5%	7%
Respiratory System		
Dyspnea	4%	7%
Skin and Appendages		
Rash	18%	24%
Skin disorder	10%	12%
Sweating	6%	8%
Alopecia	2%	4%
Urogenital System		
Urinary urgency	10%	13%
Metrorrhagia*	8%	11%
Menorrhagia*	6%	8%
Impotence**	7%	9%
Urinary frequency	5%	7%
Dysmenorrhea*	5%	7%
Prostatic disorder**	1%	3%

* pre-menopausal women
 ** male patients

The following adverse events have been observed during postmarketing experience with Betaseron and are classified within body system categories:

Body General: *fatal capillary leak syndrome; **Cardiovascular:** cardiomyopathy, deep vein thrombosis, pulmonary embolism; **Digestive:** hepatitis, pancreatitis, vomiting; **Endocrine:** hypothyroidism, hyperthyroidism, thyroid dysfunction; **Hemic and Lymphatic System:** anemia, thrombocytopenia; **Metabolic and Nutritional:** Gamma GT increase, hypocalcemia, hyperuricemia, triglyceride increase; **Nervous:** ataxia, confusion, convulsion, depersonalization, emotional lability, paresthesia; **Respiratory:** bronchospasm, pneumonia; **Skin and Appendages:** pruritus, skin discoloration, urticaria; **Urogenital:** urinary tract infection, urosepsis.

*The administration of cytokines to patients with a pre-existing monoclonal gammopathy has been associated with the development of this syndrome.

Immunogenicity
 As with all therapeutic proteins, there is a potential for immunogenicity. Serum samples were monitored for the development of antibodies to Betaseron during the RRMS study. In patients receiving 0.25 mg every other day 56/124 (45%) were found to have serum neutralizing activity at one or more of the time points tested. The relationship between antibody formation and clinical safety or efficacy is not known.

These data reflect the percentage of patients whose test results were considered positive for antibodies to Betaseron using a biological neutralization assay that measures the ability of immune sera to inhibit the production of the interferon-inducible protein, MxA. Neutralization assays are highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of neutralizing activity in an assay may be influenced by several factors including sample handling, timing of sample collection,

Continued on next page

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.