

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 July 2001 (19.07.2001)


PCT

(10) International Publication Number  
WO 01/51049 A1

- (51) International Patent Classification<sup>7</sup>: A61K 31/44, 31/445 (US). SHAW, Leslie, M. [US/US]; 705 Sunnyside Avenue, Audubon, PA 19403 (US).
- (21) International Application Number: PCT/US01/01537 (74) Agents: COLBY, Gary, D. et al.; Akin, Gump, Strauss, Hauer & Feld, L.L.P., One Commerce Square, 22nd floor, 2005 Market Street, Philadelphia, PA 19103-7986 (US).
- (22) International Filing Date: 12 January 2001 (12.01.2001)
- (25) Filing Language: English (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (26) Publication Language: English (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (30) Priority Data: 60/176,086 14 January 2000 (14.01.2000) US
- (71) Applicant (*for all designated States except US*): THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): WASIK, Mariusz, A. [US/US]; 210 Cedarbrook Road, Ardmore, PA 19003

[Continued on next page]

(54) Title: O-METHYLATED RAPAMYCIN DERIVATIVES FOR ALLEVIATION AND INHIBITION OF LYMPHOPROLIFERATIVE DISORDERS



(57) Abstract: The present invention relates to methods of alleviating and inhibiting a lymphoproliferative disorder in a mammal, the method comprising administering one or more rapamycin derivatives (including rapamycin) to the mammal (see figure 8). Further, the invention provides a method for identifying agents which are useful for alleviating and inhibiting lymphoproliferative disorders, as well as a method for identifying agents which are capable of inhibiting metastasis of lymphatic tumors in a mammal.

WO 01/51049 A1



**Published:**

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## TITLE

5 O-Methylated Rapamycin Derivatives for  
Alleviation and Inhibition of Lymphoproliferative Disorders

## BACKGROUND OF THE INVENTION

Post-transplant lymphoproliferative disorders (PTLDs) which usually involve expansion of B lymphocytes infected with the Epstein-Barr virus (EBV), are a life-threatening complication of the immunosuppressive therapy  
10 necessary to inhibit graft rejection (Morrison et al., 1994, Am. J. Med. 97:14-24; Warnke et al., 1995, AFIP Fascicle 14:531-535). PTLDs comprise a whole spectrum of lymphoproliferative disorders ranging from a polyclonal atypical lymphoid hyperplasia to a monoclonal, overtly malignant B-cell lymphoma  
15 (Morrison et al., 1994, Am. J. Med. 97:14-24; Warnke et al., 1995, AFIP Fascicle 14:531-535; Curtis et al., 1999, Blood 94:2208-2216; Harris et al., 1997, Semin. Diagn. Path. 14:8-14). Less advanced forms of PTLDs respond to a less aggressive course of immunosuppressive therapy (Morrison et al., 1994, Am. J. Med. 97:14-24; Sigal et al., 1992, Ann. Rev. Immunol. 10:519-60). However, lowering the dose  
20 of standard immunosuppressive drugs, which nullify the body's ability to reject and destroy foreign tissue, can jeopardize the survival of a graft. Moreover, this modification in treatment with conventional agents is not effective against malignant, lymphoma-type PTLDs which are usually fatal for the graft recipient.

Lymphoma causes significant morbidity and mortality, accounting  
25 for more than 50,000 new diagnoses annually in the United States alone. Many lymphomas are either Hodgkin's or non-Hodgkin's lymphomas, which can be derived from peripheral, mature B, T, or NK lymphomas. Based on their natural course, non-Hodgkin's lymphomas are classified into low, intermediate, and high grades. Low grade lymphomas are usually slowly progressive, but are essentially  
30 non-curable. The current 5-year disease-free, post-therapy survival rate for the intermediate and high grade lymphomas is approximately 60%. These aggressive types of lymphoma result in a rapid demise of the patients who do not respond to therapy. Prognosis of lymphomas occurring in patients who are

immunocompromised such as AIDS and post-transplant patients, is particularly poor. Therefore, new treatment modalities are needed to improve cure rate of lymphoma.

SDZ RAD (40-O-{2-hydroxyethyl}-rapamycin) is one of a class of rapamycin derivatives which exhibit immunosuppressive activities (PCT application WO 94/09010; Schuurman et al., 1997, Transplantation 64:32-5; Schuler et al. 1997, Transplantation 64:36-42; Sedrani, et al., 1998, Transplant. Proc. 30:2192-2194; Schuurman et al., 1998, Transplant Proc. 30:2198-2199; Hausen et al., 1999, J. Heart Lung Transplant 18:150-159). Compounds of this class, including rapamycin, have several points of action in normal T lymphocytes. They inhibit primarily down-stream signaling events mediated by the IL-2 receptor (Seghal, 1998, Clin. Biochem. 31:335-340) and other cytokine receptors (Sakata et al., 1999, Immunology Letters 68:301-309), but also affect cell-cycle progression at the early G<sub>1</sub> phase (Terada et al., 1993, J. Cell Physiol. 154:7-15; Flanagan et al., 1993, Ann. N.Y. Acad. Sci. 696:31-37). The multi-faceted immunosuppressive activities exhibited by SDZ RAD and other O-alkylated rapamycin derivatives compounds make these compounds versatile immunosuppressive agents.

There is a significant need for more effective therapeutic and prophylactic methods for limiting the severity and frequency of lymphoproliferative disorders such as lymphomas and PTLDs. The present invention satisfies this need.

#### BRIEF SUMMARY OF THE INVENTION

The invention includes a method of alleviating a lymphoproliferative disorder in a human patient. The method comprises administering to the patient, in an amount sufficient to alleviating the disorder, a rapamycin derivative having the chemical structure shown in Formula I in Figure 8. In a preferred embodiment, the rapamycin derivative is 40-O-(2-hydroxy)ethyl-rapamycin. Numerous other useful rapamycin derivatives (including rapamycin itself) are described in this disclosure. Lymphoproliferative disorders that can be alleviated using this method include, for example, PTLDs and lymphatic cancers such as lymphomas. The method can also be used to alleviate lymphoproliferative disorders caused or associated with

W3 01/31/15 PCT/US01/0137  
treatment of the patient by immunosuppressive therapy (e.g., immunosuppressive therapy associated with tissue transplantation).

In another method included in the invention, a lymphoproliferative disorder is inhibited or prevented in a patient at risk for developing such a disorder (e.g., an immunocompromised patient or a patient undergoing immunosuppressive therapy).

In these methods, the rapamycin derivative can be co-administered (in a single composition or in discretely-administered compositions) with a second pharmacologically active agent, such as an immunosuppressive agent.

Immunosuppressive agents are known for use in methods of inhibiting graft rejection, and those known methods can be improved by administering both the immunosuppressive agent and a rapamycin derivative disclosed herein to a patient who has received a graft.

The invention also includes a method of inhibiting metastasis of a lymphatic tumor in a human patient afflicted with a lymphatic cancer. This method comprising administering to the patient, in an amount sufficient to inhibit lymphocyte proliferation, a rapamycin derivative having the chemical structure shown in Formula I.

In another aspect, the invention includes a method of assessing whether an agent is useful for alleviating or inhibiting a lymphoproliferative disorder in a human patient. This method comprising transforming a B lymphocyte with an Epstein-Barr virus, injecting the lymphocyte into a mouse having a severe combined immunodeficiency, administering the agent to the mouse, and monitoring tumor growth in the mouse for at least about 21 days. If one or more of tumor regression, tumor eradication, and absence of a second tumor is observed in the mouse, then this is an indication that the agent is useful for alleviating or inhibiting a post-transplant lymphoproliferative disorder in a mammal.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph which illustrates SDZ RAD-mediated inhibition of in vitro proliferation of PTLD-like EBV+ B cells. BC-1 is an EBV+/HSV8+ B-

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.