

**ECCO 11**

the European  
Cancer  
Conference

*Lisbon*

21-25 October 2001

Vol 37 Supplement 6

October 2001

ISSN 0959-8049

**ABSTRACT BOOK**

Univ. of Minn.  
Bio-Medical  
Library

11 14 01

# EJC

*The Official Journal of*

EORTC

European Organization for  
Research and Treatment of Cancer

ESO

European School of Oncology

EACR

European Association for  
Cancer Research


FECS

Federation of European  
Cancer Societies

EUSOMA

European Society of Mastology

EUROPEAN JOURNAL OF CANCER



FEDERATION OF  
EUROPEAN  
CANCER  
SOCIETIES



PERGAMON

**DOCKET  
ALARM**

Find authenticated court documents without watermarks at [docketalarm.com](http://docketalarm.com).

**Aim of the Study:** We prospectively studied circulating T, N-demethyl T, N-dedimethyl T and 4-hydroxy T (D-T, DD-T and 4-OH-T) that to be matched with lipids, coagulation, blood counts, steroid hormone binding globulin (SHBG), various hormones, vaginal karyopyknotic index (KPI) (our previous studies BMJ ii: 1351-2, 1977; Cytopatol 9: 263-270, 1998) and transvaginal ecosonography endometrial morphology.

**Patients** were 85, consecutively enrolled after radical surgery for M<sub>0</sub>PBC, and given adjuvant T 20 mg/day for 5 years.

**Results:** T induces most of its agonist estrogen-like effects within 2 w.s therapy. A steady state of the biological events is reached in about 4 w.s and persists up to 60 mo.s. Cholesterol, LDL-C, HDL-C, LDL-C/HDL-C ratio, apolipoprotein B/apolipoprotein A ratio, fibrinogen and antithrombin III activity dropped from 2 w.s to 60 mo.s. HDL-C was unchanged, SHBG and KPI increased early, while FSH, LH and prolactin decreased. A late rise of Hb, PCV occurred from the 18<sup>th</sup> mo. Platelets were lower from the 4<sup>th</sup> w (all quoted results:  $p \leq 0.005$ ). Plasma concentration of T and its metabolites shows interpersonal variations; the steady state was reached at the 4<sup>th</sup> w. Positive relations ( $p \leq 0.005$ ) existed between 4-OH-T and fibrinogen and KPI, inverse relation between 4-OH-T and antithrombin III, D-T and fibrinogen, DD-T and fibrinogen. Agonistic properties are mainly exerted by the three metabolites. Further analyses is ongoing.

*In praise of GOCNE (Gruppo Oncologico Cooperativo Nord-Est), CRO-Aviano (PN), Italy. This investigation is part of the study programs of and partially supported by AOI-Associazione Oncologica Italiana, Aviano (PN), Italy.*

240

POSTER

#### Pharmacokinetics (PK) of iriflufen using two different intermittent dosing schedules as a 30 minute (MIN) infusion in advanced solid tumors (AST): Preliminary data

F. Lokiec<sup>1</sup>, E. Raymond<sup>2</sup>, J. Alexandre<sup>3</sup>, E. Brain<sup>1</sup>, M. Ould Kaci<sup>4</sup>, S. Lagree<sup>4</sup>, S. Smith<sup>5</sup>, K. Vanderbilt<sup>5</sup>, E. Cvitkovic<sup>4</sup>, J.L. Misset<sup>3</sup>, <sup>1</sup>Center René Huguenin, Saint-Cloud; <sup>2</sup>Institut Gustave-Roussy, Villejuif; <sup>3</sup>Hôpital Paul Brousse, Villejuif; <sup>4</sup>CAC, Kremlin-Bicêtre, France; <sup>5</sup>MGI Pharma, Minnesota, USA

**Purpose:** Iriflufen, an acylfulvene analog of illudin S, has shown promising anti-tumor activity during preclinical/clinical development, with delayed thrombocytopenia, severe asthenia and nausea/vomiting as treatment-limiting toxicities (tox). PK analysis of the 5 min daily  $\times$  5 or intermittent weekly dosing q3 or 4w schedules (sch) has shown a short mean plasma half-life ( $t_{1/2}$ ) range: 4-6 min; with a large interpatient (pt) variability. A 30 min administration corresponds to approximately 5  $t_{1/2}$ . Therefore, by the end of a 30 min infusion, a steady state will be reached. Within the ongoing Phase I study, we decided to use the 30 min infusion to improve the PK and pharmacodynamic (PD) analyses and to investigate a possible correlation between observed tox and C<sub>max</sub>.

**Methods:** Pts with AST were treated with the same schedules (sch) previously explored with the 5 min infusion duration (B: D1, 8, q3w and C: D1, 15 q4w) using the following dosing levels (DL in [mg/m<sup>2</sup>/d]). Sch B: DL2 [18], DL3 [21]; Sch C: DL2 [24], DL3 [28]. During each pt's first 3 infusions, 10 plasma samples were collected up to 5 hours post-infusion. As of 04/2001, 24 pts were included in DL2 and DL3 of both sch.

**Results:** PK analysis has been performed for day 1 in the first 17 pts. Total body clearance appeared stable up to 28 mg/m<sup>2</sup>.

Dose mg/m <sup>2</sup>	Sch (DL)	N eval. pts	C <sub>max</sub> ng/ml	AUC ng/mlxh	Clearance l/h/m <sup>2</sup>	T <sub>1/2</sub> beta min
18	B (DL2)	7	112 ± 58	38.8 ± 18.1	568 ± 279	7.1 ± 3.6
21	B (DL3)	2	115 ± 4	35.9 ± 1.4	586 ± 23	1.9 ± 0.1
24	C (DL2)	7	202 ± 181	68.7 ± 61.6	532 ± 293	8.5 ± 6.8
28	C (DL3)	1	214	70.3	398	4.7

**Conclusions:** Preliminary results show that AUCs and total body clearances are similar when iriflufen is administered as a 5 or a 30 minute infusion, despite its short terminal half-life. Fully updated PK analysis and the PK/PD relationships will be presented.

241

POSTER

#### Pharmacokinetic Interactions between gemcitabine (GEM) and vinorelbine (VNR) in patients with advanced stage solid tumors

M. Airolidi<sup>1</sup>, S. Novello<sup>2</sup>, E. Buffa<sup>2</sup>, M.C. Verleno<sup>2</sup>, S. Marchionatti<sup>1</sup>, L. Cattel<sup>2</sup>, <sup>1</sup>S. Giovanni A.S. Hospital, Medical Oncology, Turin, Italy; <sup>2</sup>University of Turin, Postgraduate School in Clinical Pharmacy, Turin, Italy

**Purpose:** GEM and VNR are both active as single agents in patients with advanced stage of solid tumors. Because of their different mechanism of action, good tolerability and feasible administration on an outpatient basis they should be an interesting combination for palliative chemotherapy. The aim of this study is to determine possible pharmacokinetic interactions between GEM and VNR.

**Methods:** A total of 11 patients with advanced non small cell lung cancer or metastatic breast cancer were treated with GEM (1h i.v. infusion, 1000 mg/m<sup>2</sup>) followed by VNR (10 min i.v. slow bolus, 25 mg/m<sup>2</sup>) on days 1, 8, every 3 weeks; 5 patients received single-agent GEM (1h i.v. infusion, 1000 mg/m<sup>2</sup>) as a control group. GEM and VNR were measured in blood samples taken at several time points after the starting of treatment and immediately added with enzymatic inhibitor of deaminase THU. Plasma levels were quantified by liquid extraction followed by HPLC analysis and pharmacokinetic data were processed by KineticTM 1.1 computer program

**Results:** In the schedule GEM+VNR average C<sub>max</sub> of GEM was 26278±4671 ng/ml, AUC=20305±3111 ng/ml/h, C<sub>tot</sub>=75,48±14,56 l/h, t<sub>1/2</sub>alpha=5,1±5,3 min; t<sub>1/2</sub>beta=18,2±3,4 min. The steady state was reached about 30 min after the beginning of administration; C<sub>max</sub> of VNR was 813±387 ng/ml, AUC=224±75 ng/ml/h, C<sub>tot</sub>=192,74±72,51 l/h; C<sub>max</sub> of single-agent GEM was 29295±5681 ng/ml, AUC=24327±7346 ng/ml/h, C<sub>tot</sub>=75,20±17,78 l/h, t<sub>1/2</sub>alpha=3,0±5,1 min, t<sub>1/2</sub>beta=20,0±5,2 min.

**Conclusions:** C<sub>max</sub>, AUC and Cl values of GEM followed by VNR agreed with the values shown in monotherapy; C<sub>p/t</sub> curve for GEM in combination was best described by a biphasical model with a rapid distribution and elimination. C<sub>p/t</sub> curve for VNR showed rapid distribution; moreover, interpatient variability was important for all parameters in accordance to the literature data; these preliminary data have shown that treatment with GEM+VNR do not alter the pharmacokinetic behaviour of both drugs compared with single agent therapy. Further investigation is needed to relate pharmacokinetic data to toxicity of the treatment.

242

POSTER

#### Pharmacokinetics of escalating doses of CCI-779 in combination with 5-fluorouracil and leucovorin in patients with advanced solid tumors

J. Boni<sup>1</sup>, J. Korth-Bradley<sup>1</sup>, C.J.A. Punt<sup>2</sup>, A. Hanauske<sup>3</sup>, K. Weigang-Köhler<sup>4</sup>, C. Thielert<sup>5</sup>, J. Frisch<sup>5</sup>, <sup>1</sup>Wyeth-Ayerst Research, Clinical Pharmacokinetics, Radnor, PA, USA; <sup>2</sup>University Medical Center, Nijmegen, The Netherlands; <sup>3</sup>Onkologische Tagesklinik, Munich, Germany; <sup>4</sup>Klinikum Nuernberg, Nuernberg, Germany; <sup>5</sup>Wyeth Oncology, Munich, Germany

**Purpose:** To evaluate the pharmacokinetics of 5-fluorouracil (5-FU) and CCI-779, administered in combination to patients with advanced solid tumors.

**Methods:** Weekly treatment spanned 6 weeks of a 7-week cycle. During day 1 of each treatment cycle, a 1 hour intravenous (IV) infusion of 200 mg/m<sup>2</sup> leucovorin (LV) was followed by a 24 hour continuous infusion of 2600 mg/m<sup>2</sup> 5-FU. Starting on cycle 1/day 8, 15 to 75 mg/m<sup>2</sup> of IV CCI-779 preceded LV/5-FU treatment. Concentrations of 5-FU were measured throughout the 24 hour infusion period in cycle 1/weeks 1, 2 and 4; CCI-779 and its primary metabolite sirolimus (rapamycin) were followed for 8 days during cycle 1/weeks 2 and 4. Parameters were derived using noncompartmental methods.

**Results:** For 15 patients, mean±SD steady-state 5-FU plasma concentration was 667±202 ng/mL (Week 1 without CCI-779) and 666±248 ng/mL (Week 2 with CCI-779); mean clearance (CL) was 325±113 L/h (Week 2). For 13 patients, mean CCI-779 CL increased with increasing dose (13 L/h [low dose] to 41 L/h [highest dose]) (CV ~ 30%). Mean half-life was 17±4 hours. The mean ratio of sirolimus-to-CCI-779 AUC was 2.4 to 3.8. No period effects were observed.

**Conclusion:** Inclusion of CCI-779 in the standard 5-FU/LV regimen did not affect 5-FU pharmacokinetic disposition, nor did 5-FU change CCI-779 pharmacokinetic parameters.