

PTO/SB/05 (08-08)

Approved for use through 06/30/2010. OMB 0651-0032
U.S. Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

15949
101508
U.S. PTO

**UTILITY
PATENT APPLICATION
TRANSMITTAL**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.	056291-5004-02
First Inventor	John R. Evans
Title	FORMULATION
Express Mail Label No.	

APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents.</i>	ADDRESS TO: Commissioner for Patents P.O. Box 1450 Alexandria VA 22313-1450
--	---

1. Fee Transmittal Form (e.g., PTO/SB/17)
2. Applicant claims small entity status.
See 37 CFR 1.27.
3. Specification [Total Pages 23]
Both the claims and abstract must start on a new page
(For information on the preferred arrangement, see MPEP 608.01(e))
4. Drawing(s) (35 U.S.C. 113) [Total Sheets 1]
5. Oath or Declaration [Total Sheets 1]
a. Newly executed (original or copy)
b. A copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 18 completed)
i. DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s)
name in the prior application, see 37 CFR
1.63(d)(2) and 1.33(b).
6. Application Data Sheet. See 37 CFR 1.76
7. CD-ROM or CD-R in duplicate, large table or
Computer Program (Appendix)
 Landscape Table on CD
8. Nucleotide and/or Amino Acid Sequence Submission
(if applicable, items a. - c. are required)
a. Computer Readable Form (CRF)
b. Specification Sequence Listing on:
i. CD-ROM or CD-R (2 copies); or
ii. Paper
c. Statements verifying identity of above copies

- ACCOMPANYING APPLICATION PARTS**
9. Assignment Papers (cover sheet & document(s))
Name of Assignee _____
 10. 37 CFR 3.73(b) Statement Power of
(when there is an assignee) Attorney
 11. English Translation Document (if applicable)
 12. Information Disclosure Statement (PTO/SB/08 or PTO-1449)
 Copies of citations attached
 13. Preliminary Amendment
 14. Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
 15. Certified Copy of Priority Document(s)
(if foreign priority is claimed)
 16. Nonpublication Request under 35 U.S.C. 122(b)(2)(B)(i).
Applicant must attach form PTO/SB/35 or equivalent.
 17. Other: _____

18. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in the first sentence of the specification following the title, or in an Application Data Sheet under 37 CFR 1.76:


Continuation Divisional Continuation-in-part (CIP) of prior application No.: 10/872,784

Prior application information: Examiner San Ming R. Hui Art Unit: 1617

19. CORRESPONDENCE ADDRESS

The address associated with Customer Number: 09629 OR Correspondence address below

Name		City		State		Zip Code	
Address		Country		Telephone		Email	

Signature		Date	October 15, 2008
Name (Print/Type)	Donald J. Bird	Registration No. (Attorney/Agent)	25,323

This collection of information is required by 37 CFR 1.53(b). The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.
If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

FORMULATION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound

5 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-
aqueous ester solvent which is miscible in the ricinoleate vehicle.

10 Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with
15 antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

20 The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such
25 compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989).
30 One of these, 7α -[9-(4,4,5,5,5-pentafluoropentyl sulphinyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available

antioestrogens. *In vitro* findings and early clinical experience with 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

5 7α -[9-(4,4,5,5,5-Pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely
10 blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

Because fulvestrant has none of the oestrogen-like stimulatory activity that is
15 characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose
20 which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross
25 the blood-brain barrier.

European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35
30 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a

pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml⁻¹ (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations:

In the formulations within Table 1 a number of different oils are used to solubilise the compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

20

25

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

<u>PRODUCT NAME</u>	<u>STEROID</u>	<u>DOSE</u>	<u>TYPE</u>	<u>COMP'</u>	<u>SOURCE</u>	<u>OIL</u>	<u>BzBz</u>	<u>BzOH</u>	<u>EtOH</u>	<u>DOSE</u>	<u>DOSING</u>
SUSTANON 100	Testosterone propionate	30mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
	Testosterone phenylpropionate	60mg									
	Testosterone isocaproate	60mg									
	Testosterone decanoate	100mg									
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%		1 or 2ml	1 week	
TOCOGESTAN	Hydroxy progesterone enantate	200mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%			2ml	< 1week
	Progesterone	50mg									
	α -Tocopherol	250mg									
TROPHOBOLENE	Estrapronicate	1.3mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%			1ml	15 to 30 days
	Nandrolone undecanoate	50mg									
	Hydroxyprogesterone heptanoate	80mg									
NORISTERAT	Norethisterone oenanthoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES		1ml	8 weeks	
BENZO- GYNOESTRYL PROGESTERONE -RETARD	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis				1ml	1 week
	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES			1 or 2ml	1 week
GRAVIBINAN	Estradiol 17- β -valerate Hydroxyprogesterone caproate	5mgml ⁻¹ 250mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES			1 or 2ml	1-2 weeks

Z70635

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis	75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%	
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestrogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%	

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
 5 % are w/v and * approximate as measured directly from a single sample

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

10

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically

significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

5 Currently guidelines recommend that no more than 5mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

10 The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of $>50\text{mgml}^{-1}$ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of $<5\text{ml}$ - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into
15 a concentration of at least 50mgml^{-1} - see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

20 Therefore, we present as a feature of the invention a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

25 Another feature of the invention is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

30 Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of

formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for
 5 intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of
 10 fulvestrant.

For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of
 15 the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Preferred pharmaceutical formulations of the invention are as described above wherein:

- 20 1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml^{-1} .
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
3. The total amount of fulvestrant in the formulation is 250mg and the total volume of
 25 the formulation is 5-5.25ml.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably up to 0.15ml will also be present in the formulation. Typically the formulation will be
5 presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and,
10 preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-
15 28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the
20 formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral
25 administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

30 It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain

some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

5 Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v, at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or
 10 less, 50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-
 15 25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

20 Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.

17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.
2. Our findings from studies using ¹⁴C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

5 We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still achieved by the formulation of the invention.

10 By use of the term "therapeutically significant levels" we mean that blood plasma concentrations of at least 2.5 ngml⁻¹, ideally at least 3 ngml⁻¹, at least 8.5 ngml⁻¹, and up to 12 ngml⁻¹ of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 15 ngml⁻¹.

By use of the term "extended release" we mean at least two weeks, at least three 15 weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

It will be understood that the attendant physician may wish to administer the 20 intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally 25 containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

Table 3Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

5 **Solubility comparisons of fulvestrant in oil based formulations with and without alcohols**

		Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
10	Formulation ^(a)	Complete vehicle	Vehicle minus alcohols
	Castor oil based	81.2	12.6
15	Miglyol 812-N based	86.8	1.7
	Sesame seed/Castor oil (1:1) based	70.1	4.4
	Sesame seed oil based	45.7	0.7
20	Arachis oil-based	40.2	< 0.2

25 ^(a) Complete Vehicle Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

		Days						
30	Formulation ^a	2	3	4	7	10	30	51
35	Formulation F1 castor oil based	0	0	0	0	0	0	0
	Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
40	Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

45 ^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant-plasma profile over five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

Preferably 5ml of the intramuscular injection is administered.

A further feature of the invention is use of fulvestrant in the preparation of a pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent
5 indications such as breast cancer and gynaecological conditions, such as endometriosis.

In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical
10 structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a
15 pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

20 Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

25 Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay
30 as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary

pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram below

5

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol

10 10% weight per volume of ethanol

15% weight per volume of benzyl benzoate

250mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil

FLOW DIAGRAM OF MANUFACTURING

Ingredients/Components

Fulvestrant
Alcohol
Benzyl Alcohol

Benzyl Benzoate

Castor Oil

Process

STAGE 1: DISSOLUTION OF
ACTIVE AGENT

STAGE 2: MIX

STAGE 3: MAKE TO
WEIGHT

STAGE 4: STERILE FILTRATION
(0.2µm)
INTO BULK RECEIVING VESSEL

STAGE 5: STERILE (0.2µm)
IN-LINE FILTRATION

STAGE 6: ASEPTIC FILLING,
AND STOPPERING

STAGE 7: VISUAL
INSPECTION

References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids* 1989; 54:71-99.
- 5 2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- 10 4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- 15 5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112: R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

Claims

1. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration of at least 2.5ngml^{-1} for at least 2 weeks.
2. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for at least 4 weeks.
3. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for 2 to 5 weeks.
4. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of fulvestrant.
5. A pharmaceutical formulation as claimed in claim 1 to 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.
6. A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of a pharmaceutically-acceptable alcohol.
7. A pharmaceutical formulation as claimed in any claim from 1 to 6 which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
8. A pharmaceutical formulation as claimed in claim 7 which contains 50%w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent .

9. A pharmaceutical formulation as claimed in claim 7 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 5 10. A pharmaceutical formulation as claimed in claim 7 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
11. A pharmaceutical formulation as claimed in claim 7 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 10 12. A pharmaceutical formulation as claimed in claim 7 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
13. A pharmaceutical formulation as claimed in claim 7 which contains 25% w/v or less
15 of a pharmaceutically-acceptable non-aqueous ester solvent.
14. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- 20 15. A pharmaceutical formulation as claimed in any claim from 1 to 14 wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
16. A pharmaceutical formulation as claimed in any claim from 1 to 15 wherein the
25 pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
17. A pharmaceutical formulation as claimed in any claim from 1 to 16 wherein the total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.

18. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 5 19. A pharmaceutical formulation as claimed in claim 18 wherein the total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5 to 5.25ml.
20. A pharmaceutical formulation as claimed in any of claims 1-19 wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of
10 formulation, 10% weight of benzyl alcohol per volume of formulation and 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.
21. A method of treating a benign or malignant diseases of the breast or reproductive tract by administration to a human in need of such treatment by intramuscular a pharmaceutical
15 formulation as claimed in claims 1 to 19.
22. A method as claimed in claim 21 for treating breast cancer.
23. A syringe or vial containing a pharmaceutical formulation as defined in claim 20.

ABSTRACT**TITLE: Formulation**

The invention relates to a novel sustained release pharmaceutical formulation adapted
5 for administration by injection containing the compound
 7α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more
particularly to a formulation adapted for administration by injection containing the compound
 7α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in
10 solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-
aqueous ester solvent which is miscible in the ricinoleate vehicle.




Figure 1

FOR UTILITY/DESIGN
CIP/PCT NATIONAL/PLANT
ORIGINAL/SUBSTITUTE/SUPPLEMENTAL
DECLARATIONS

RULE 63 (37 C.F.R. 1.61)
DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the INVENTION ENTITLED

FORMULATION

the specification of which (CHECK applicable BOX(ES))
X A. is attached hereto.
BOX(ES) → B. was filed on _____ as U.S. Application No. _____ /
→ C. was filed as PCT International Application No. PCT/ _____ / _____ on _____
and (if applicable to U.S. or PCT application) was amended on _____

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56. Except as noted below, I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International Application which designated at least one other country than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate, or PCT International Application, filed by me or my assignee disclosing the subject matter claimed in this application and having a filing date (1) before that of the application on which priority is claimed, or (2) if no priority claimed, before the filing date of this application:

PRIOR FOREIGN APPLICATION(S)

Number	Country	Day/MONTH/Year Filed	Date first Laid-open or Published	Date Patented or Granted	Priority NOT Claimed
0000313.7	GB	10 January 2000			
0008837.7	GB	12 April 2000			

If more prior foreign applications, X box at bottom and continue on attached page.

Except as noted below, I hereby claim domestic priority benefit under 35 U.S.C. 119(a) or 120 and/or 365(c) of the indicated United States applications listed below and PCT international applications listed above or below and, if this is a continuation-in-part (CIP) application, insofar as the subject matter disclosed and claimed in this application is in addition to that disclosed in such prior applications, I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56 which became available between the filing date of each such prior application and the national or PCT international filing date of this application:

PRIOR U.S. PROVISIONAL, NONPROVISIONAL AND/OR PCT APPLICATION(S)

Application No. (series code/serial no.)	Day/MONTH/Year Filed	Status	Priority NOT Claimed
		pending, abandoned, patented	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

And I hereby appoint Pillsbury Madison & Sutro LLP, Intellectual Property Group, 1100 New York Avenue, N.W., Ninth Floor, East Tower, Washington, D.C. 20005-3918, telephone number (202) 861-3000 (to whom all communications are to be directed), and the below-named persons (of the same address) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent, and I hereby authorize them to delete names/numbers below of persons no longer with their firm and to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct the above firm and/or a below attorney in writing to the contrary.

Paul N. Kokufis	16773	Dale S. Lazar	28872	Mark G. Paulson	30793	Michael R. Dzwonczyk	36787
Raymond F. Lippitt	17519	Paul E. White, Jr.	32011	Stephen C. Glazier	31361	W. Patrick Bengtsson	32456
G. Lloyd Knight	17698	Glenn J. Perry	28458	Paul F. McQuade	31542	Jack S. Barufka	37087
Carl G. Love	18781	Kendrew H. Colton	30368	Ruth N. Morduch	31044	Adam R. Hess	41835
Kevin E. Joyce	20508	G. Paul Edgell	24238	Richard H. Zaitlen	27248		
George M. Sirilla	18221	Lynn E. Eccleston	35861	Roger R. Wise	31204		
Donald J. Bird	25323	Timothy J. Klima	34852	Jay M. Finkelstein	21082		
Peter W. Gowdey	25872	David A. Jakopin	32995	Anita M. Kirkpatrick	32617		

(1) INVENTOR'S SIGNATURE:

John R Evans

Date: *25th January 2001*

John	R	EVANS
First	Middle Initial	Family Name
Residence	Macclesfield, Cheshire	United Kingdom
	City	State/Foreign Country
Post Office Address	Alderley Park, Macclesfield, Cheshire, SK10 4TG, United Kingdom	
(include Zip Code)	SK10 4TG	
		Country of Citizenship

(2) INVENTOR'S SIGNATURE:

Rosalind A Grundy

Date: *25th January 2001*

ROSALIND	U	GRUNDY
First	Middle Initial	Family Name
Residence	Macclesfield, Cheshire	United Kingdom
	City	State/Foreign Country
Post Office Address	Charter Way, Macclesfield, Cheshire, SK10 2NA, United Kingdom	
(include Zip Code)	SK10 2NA	
		Country of Citizenship

FOR ADDITIONAL INVENTORS, "X" box and proceed on the attached page to list each additional inventor.

See additional foreign priorities on attached page (incorporated herein by reference).

Atty. Dkt. No. PM

(M#)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re <u>Continuation</u> Application of:)	Group Art Unit: Not Assigned
)	
John R. Evans et. al)	Examiner: Not Assigned
)	
Continuation of Application No. 10/872,784)	
)	
Application No. Not Assigned)	
)	
Filed: October 15, 2008)	
)	
For: FORUMLATION)	<u>Date: October 15, 2008</u>

PRELIMINARY AMENDMENT

Prior to examination on the merits, please amend the above-referenced application as follows:

Amendments to the Specification begin on page 2 of this paper.
Remarks begin on page 3 of this paper.

Amendments to the Specification:

On page 1, please after the title, please insert the following paragraph:


This application is a Continuation Application of copending U.S. Patent Application No. 10/872,784, filed June 22, 2004, which claims benefit of U.S. Patent Application No. 09/756,291, filed January 9, 2001 which claims the benefit of Great Britain Application No. 0008837.7 filed April 12, 2000 and Great Britain Application No. 0000313.7, filed January 10, 2000, all of which are incorporated herein by reference in their entireties.

REMARKS

The specification has been amended to update the priority data. Applicants submit that the amendments to the specification do not introduce prohibited new matter.

Dated: **October 15, 2008**
Morgan, Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel: 202-739-3000
Fax: 202-739-3001

Respectfully submitted
Morgan, Lewis & Bockius LLP



Donald J. Bird
Registration No. 25,323

Filing Date: 10/15/08

Approved for use through 7/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875	Application or Docket Number 12/285,887
---	---

APPLICATION AS FILED – PART I			SMALL ENTITY		OR		OTHER THAN SMALL ENTITY	
(Column 1) (Column 2) (Column 3)			RATE (\$)	FEE (\$)		RATE (\$)	FEE (\$)	
FOR	NUMBER FILED	NUMBER EXTRA	N/A			N/A	330	
BASIC FEE (37 CFR 1.16(a), (b), or (c))	N/A	N/A	N/A			N/A	540	
SEARCH FEE (37 CFR 1.16(k), (l), or (m))	N/A	N/A	N/A			N/A	220	
EXAMINATION FEE (37 CFR 1.16(o), (p), or (q))	N/A	N/A	N/A			N/A	156	
TOTAL CLAIMS (37 CFR 1.16(i))	23	minus 20 =	3	x\$26		x\$52	156	
INDEPENDENT CLAIMS (37 CFR 1.16(h))	2	minus 3 =	*	x\$110		x\$220		
APPLICATION SIZE FEE (37 CFR 1.16(s))	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$270 (\$135 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR							
MULTIPLE DEPENDENT CLAIM PRESENT (37 CFR 1.16(j))				195		390		
				TOTAL		TOTAL	1246	

* If the difference in column 1 is less than zero, enter "0" in column 2.

APPLICATION AS AMENDED – PART II					SMALL ENTITY		OR		OTHER THAN SMALL ENTITY		
(Column 1) (Column 2) (Column 3)					RATE (\$)	ADDITIONAL FEE (\$)		RATE (\$)	ADDITIONAL FEE (\$)		
AMENDMENT A	CLAIMS REMAINING AFTER AMENDMENT	MINUS	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA							
	Total (37 CFR 1.16(i))	*	Minus **	=	X =		X =				
	Independent (37 CFR 1.16(h))	*	Minus ***	=	X =		X =				
	Application Size Fee (37 CFR 1.16(s))										
	FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))					N/A		N/A			
					TOTAL ADD'T FEE		TOTAL ADD'T FEE				

APPLICATION AS AMENDED – PART II					SMALL ENTITY		OR		OTHER THAN SMALL ENTITY		
(Column 1) (Column 2) (Column 3)					RATE (\$)	ADDITIONAL FEE (\$)		RATE (\$)	ADDITIONAL FEE (\$)		
AMENDMENT B	CLAIMS REMAINING AFTER AMENDMENT	MINUS	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA							
	Total (37 CFR 1.16(i))	*	Minus **	=	X =		X =				
	Independent (37 CFR 1.16(h))	*	Minus ***	=	X =		X =				
	Application Size Fee (37 CFR 1.16(s))										
	FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))					N/A		N/A			
					TOTAL ADD'T FEE		TOTAL ADD'T FEE				

* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.

** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".

*** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".

The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

MULTIPLE DEPENDENT CLAIM FEE CALCULATION SHEET Substitute for Form PTO-1360 (For use with Form PTO/SB/06)	Application Number 12/285,887	Filing Date 10/15/2008
Applicant(s)		

* May be used for additional claims or amendments

CLAIMS	AS FILED		AFTER FIRST AMENDMENT		AFTER SECOND AMENDMENT			AS FILED		AFTER FIRST AMENDMENT		AFTER SECOND AMENDMENT	
	Indep	Depend	Indep	Depend	Indep	Depend		Indep	Depend	Indep	Depend	Indep	Depend
1	1												
2		1											
3		1											
4	1												
5		2											
6		2											
7		2											
8		2											
9		2											
10		2											
11		2											
12		2											
13		2											
14		2											
15		2											
16		2											
17		2											
18		2											
19		2											
20		2											
21		2											
22		2											
23		2											
24													
25													
26													
27													
28													
29													
30													
31													
32													
33													
34													
35													
36													
37													
38													
39													
40													
41													
42													
43													
44													
45													
46													
47													
48													
49													
50													
Total													
Indep	2												
Total													
Depend	40												
Total													
Claims	42												

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the PTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 7 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY. DOCKET NO, TOT CLAIMS, IND CLAIMS. Row 1: 12/285,887, 10/15/2008, 1617, 0.00, 056291-5004-02, 23, 2

CONFIRMATION NO. 1199

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004

FILING RECEIPT



Date Mailed: 11/04/2008

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Applicant(s)

John R. Evans, Macclesfield, UNITED KINGDOM;
Rosalind U. Grundy, Macclesfield, UNITED KINGDOM;

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 10/872,784 06/22/2004

Foreign Applications

UNITED KINGDOM 0008837.7 04/12/2000
UNITED KINGDOM 0000313.7 01/10/2000

If Required, Foreign Filing License Granted: 11/03/2008

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is US 12/285,887

Projected Publication Date: To Be Determined - pending completion of Missing Parts

Non-Publication Request: No

Early Publication Request: No

Title

Formulation

Preliminary Class

514

PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4158).

LICENSE FOR FOREIGN FILING UNDER

Title 35, United States Code, Section 184

Title 37, Code of Federal Regulations, 5.11 & 5.15

GRANTED

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as

set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

NOT GRANTED

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (12/285,887), FILING OR 371(C) DATE (10/15/2008), FIRST NAMED APPLICANT (John R. Evans), ATTY. DOCKET NO./TITLE (056291-5004-02)

CONFIRMATION NO. 1199

FORMALITIES LETTER

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 11/04/2008

NOTICE TO FILE MISSING PARTS OF NONPROVISIONAL APPLICATION

FILED UNDER 37 CFR 1.53(b)

Filing Date Granted

Items Required To Avoid Abandonment:

An application number and filing date have been accorded to this application. The item(s) indicated below, however, are missing. Applicant is given TWO MONTHS from the date of this Notice within which to file all required items and pay any fees required below to avoid abandonment.

- The statutory basic filing fee is missing. Applicant must submit \$330 to complete the basic filing fee for a non-small entity. If appropriate, applicant may make a written assertion of entitlement to small entity status and pay the small entity filing fee (37 CFR 1.27).

The application is informal since it does not comply with the regulations for the reason(s) indicated below.

The required item(s) identified below must be timely submitted to avoid abandonment:

- A substitute specification in compliance with 37 CFR 1.52, 1.121(b)(3), and 1.125, is required. The substitute specification must be submitted with markings and be accompanied by a clean version (without markings) as set forth in 37 CFR 1.125(c) and a statement that the substitute specification contains no new matter (see 37 CFR 1.125(b)). The specification, claims, and/or abstract page(s) submitted is not acceptable and cannot be scanned or properly stored because:
- The application contains drawings, but the specification does not contain a brief description of the several views of the drawings as required by 37 CFR 1.74 and 37 CFR 1.77(b)(7).

Applicant is cautioned that correction of the above items may cause the specification and drawings page count to exceed 100 pages. If the specification and drawings exceed 100 pages, applicant will need to submit the required application size fee.

The applicant needs to satisfy supplemental fees problems indicated below.

The required item(s) identified below must be timely submitted to avoid abandonment:

- Additional claim fees of \$156 as a non-small entity, including any required multiple dependent claim fee, are required. Applicant must submit the additional claim fees or cancel the additional claims for which fees are due.

- To avoid abandonment, a surcharge (for late submission of filing fee, search fee, examination fee or oath or declaration) as set forth in 37 CFR 1.16(f) of **\$130** for a non-small entity, must be submitted with the missing items identified in this notice.

SUMMARY OF FEES DUE:

Total additional fee(s) required for this application is **\$1376** for a non-small entity

- **\$330** Statutory basic filing fee.
- **\$130** Surcharge.
- The application search fee has not been paid. Applicant must submit **\$540** to complete the search fee.
- The application examination fee has not been paid. Applicant must submit **\$220** to complete the examination fee for a non-small entity.
- Total additional claim fee(s) for this application is **\$156**
 - **\$156** for **3** total claims over 20.

Replies should be mailed to:

Mail Stop Missing Parts
Commissioner for Patents
P.O. Box 1450
Alexandria VA 22313-1450

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web.
<https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at **1-866-217-9197** or visit our website at <http://www.uspto.gov/ebc>.

If you are not using EFS-Web to submit your reply, you must include a copy of this notice.

/tnguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NUMBER	FILING OR 371(C) DATE	FIRST NAMED APPLICANT	ATTY. DOCKET NO./TITLE
12/285,887	10/15/2008	John R. Evans	056291-5004-02

CONFIRMATION NO. 1199

IMPROPER CPOA LETTER

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 11/04/2008

NOTICE REGARDING POWER OF ATTORNEY

This is in response to the Power of Attorney filed 10/15/2008. The Power of Attorney in this application is not accepted for the reason(s) listed below:

- The Power of Attorney you provided did not comply with the new Power of Attorney rules that became effective on June 25, 2004. See 37 CFR 1.32.

/tnguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Prior Examiner: San-Ming R. Hui
)	
FOR: FORMULATION)	Date: June 4, 2009

SECOND PRELIMINARY AMENDMENT

Prior to calculation of the excess claims fee and examination on the merits, please amend the claims of the above-referenced application as follows:

Amendments to the Claims begin on page 2 of this paper.

Remarks begin on page 6 of this paper.

IN THE CLAIMS:

This listing of claims will replace all prior versions and listing of claims in this application.

Listing of the claims:

Claim 1 (**original**): A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration of at least 2.5ngml^{-1} for at least 2 weeks.

Claim 2 (**original**): A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for at least 4 weeks.

Claim 3 (**original**): A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for 2 to 5 weeks.

Claim 4 (**original**): A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of fulvestrant.

Claim 5 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 to 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.

Claim 6 (**original**): A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of a pharmaceutically-acceptable alcohol.

Claim 7 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 6~~ which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 8 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 50%w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent .

Claim 9 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 10 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 11 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 12 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 13 (**original**): A pharmaceutical formulation as claimed in claim 7 which contains 25% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.

Claim 14 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 13~~ wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.

Claim 15 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 14~~ wherein the pharmaceutically-acceptable non-aqueous ester solvent

is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

Claim 16 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 15~~ wherein the pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.

Claim 17 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 16~~ wherein the total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.

Claim 18 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any claim from 1 to 13~~ wherein the total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.

Claim 19 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~claim 18~~ wherein the total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5 to 5.25ml.

Claim 20 (**currently amended**): A pharmaceutical formulation as claimed in claim 1 or claim 4 ~~any of claims 1-19~~ wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of formulation, 10% weight of benzyl alcohol per volume of formulation and 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.

Claim 21 (**currently amended**): A method of treating a benign or malignant diseases of the breast or reproductive tract by administration to a human in need of such treatment by intramuscular a pharmaceutical formulation as claimed in claim 1 or claim 4 ~~claims 1 to 19~~.

Claim 22 (**original**): A method as claimed in claim 21 for treating breast cancer.

Claim 23 (**original**): A syringe or vial containing a pharmaceutical formulation as defined in claim 20.

REMARKS

Claim Amendments

The claims have been amended to remove improper multiple dependencies.

These amendments have been made without waiver or prejudice to Applicants' right to prosecute any subject matter deleted thereby in one or more continuing applications. Following entry of these amendments, claims 1-23 remain pending in this application.

EXCEPT for issue fees payable under 37 C.F.R. § 1.18, the Director is hereby authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 C.F.R. §§ 1.16 and 1.17 which may be required, including any required extension of time fees, or credit any overpayment to Deposit Account 50-0310. This paragraph is intended to be a **CONSTRUCTIVE PETITION FOR EXTENSION OF TIME** in accordance with 37 C.F.R. § 1.136(a)(3).

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: June 4, 2009
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000
DJB:

By: /Donald Bird/
Donald J. Bird
Registration No. 25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Prior Examiner: San-Ming R. Hui
)	
FOR: FORMULATION)	Date: June 4, 2009

STATEMENT ACCOMPANYING SUBSTITUTE SPECIFICATION

In response to the Notice to File Missing Requirements, attached are a clean copy and a marked up copy of the substitute specification in which headings have been inserted. Both copies contain the amendments that are set forth in the Preliminary Amendment filed on October 15, 2008. The attached copies of the substitute specification do not include prohibited new matter. In particular, referring to the “marked up” copy:

- At page 1, the text under CROSS-REFERENCE TO RELATED APPLICATIONS was previously inserted by the Preliminary Amendment of October 15, 2008.
- At page 1, the deleted text under “Field of the Invention” has been copied to page 6 under SUMMARY OF THE INVENTION.
- At page 6, the text under SUMMARY OF THE INVENTION is copied from the original specification at page 1, lines 3-9.
- At page 6, the text under BRIEF DESCRIPTION OF THE DRAWING is copied from the original specification at page 15, lines 3-5.
- The claims in this substitute specification are as originally presented. However, the Examiner’s attention is drawn to the accompanying Second Preliminary Amendment wherein the claims are amended to remove improper multiple dependencies.

If there are any additional fees due in connection with the filing of this response, please charge the fees to our Deposit Account No. 50-0310. If a fee is required for an extension of time

under 37 C.F.R. §1.136 not accounted for above, such an extension is requested and the fee should also be charged to our Deposit Account.

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: **June 4, 2009**
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000

By: /Donald Bird/
Donald J. Bird
Registration No.25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a Continuation Application of copending U.S. Patent Application
5 No. 10/872,784, filed June 22, 2004, which claims benefit of U.S. Patent Application No.
09/756,291, filed January 9, 2001 which claims the benefit of Great Britain Application No.
0008837.7 filed April 12, 2000 and Great Britain Application No. 0000313.7, filed January
10, 2000, all of which are incorporated herein by reference in their entireties.

10 BACKGROUND OF THE INVENTION

Field of the Invention

The invention relates to a novel sustained release pharmaceutical formulation adapted
for administration by injection containing the compound

7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol.

15

Description of the Related Art

Oestrogen deprivation is fundamental to the treatment of many benign and malignant
diseases of the breast and reproductive tract. In premenopausal women, this is achieved by
the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in
20 postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with
antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present
in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such
as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the
25 partial agonism they display, which results in an incomplete blockade of oestrogen-mediated
activity (Furr and Jordan 1984, May and Westley 1987).

The potential for nonsteroidal antioestrogens to display agonistic properties prompted
the search for novel compounds that would bind ER with high affinity without activating any
of the normal transcriptional hormone responses and consequent manifestations of oestrogens.
30 Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like
ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such
compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for

the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7 α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989).

5 One of these, 7 α -[9-(4,4,5,5,5-pentafluoropentyl sulphanyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have
10 promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts
15 thereof and any possible solvates of either thereof.

Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the
20 uterotrophic activity of tamoxifen.

Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a
25 reduction of tumour invasiveness.

In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does
30 not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-
5 1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

10 Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml⁻¹ (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

15 Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

In the formulations within Table 1 a number of different oils are used to solubilise the
20 compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

25

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

<u>PRODUCT NAME</u>	<u>STEROID</u>	<u>DOSE</u>	<u>TYPE</u>	<u>COMP.</u>	<u>SOURCE</u>	<u>OIL</u>	<u>BzBz</u>	<u>BzOH</u>	<u>EtOH</u>	<u>DOSE</u>	<u>DOSING</u>
SUSTANON 100	Testosterone propionate Testosterone phenylpropionate Testosterone isocaproate Testosterone decanoate	30mg 60mg 60mg 100mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%			1 or 2ml	1 week
TOCOGESTAN	Hydroxy progesterone enantate Progesterone α -Tocopherol	200mg 50mg 250mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%			2ml	< 1week
TROPHOBOLENE	Estrapronicate Nandrolone undecanoate Hydroxyprogesterone heptanoate	1.3mg 50mg 80mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%			1ml	15 to 30 days
NORISTERAT	Norethisterone oentanhoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES			1ml	8 weeks
BENZO- GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis				1ml	1 week
PROGESTERONE -RETARD	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES			1 or 2ml	1 week
GRAVIBINAN	Estradiol 17- β -valerate Hydroxyprogesterone caproate	5mgml ⁻¹ 250mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES			1 or 2ml	1 - 2 weeks

Z70635

- 5 -

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis		75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%		
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestrogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%		

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
5 % are w/v and * approximate as measured directly from a single sample

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

DETAILED DESCRIPTION OF THE INVENTION

Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06

Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

Currently guidelines recommend that no more than 5mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of $>50\text{mgml}^{-1}$ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of $<5\text{ml}$ - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50mgml^{-1} - see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly

lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

Therefore, we present as a feature of the invention a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 5 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Another feature of the invention is a pharmaceutical formulation comprising 10 fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically- 15 acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

20 Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient 25 amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 30 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Preferred pharmaceutical formulations of the invention are as described above

5 wherein:

1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 10 3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably
 15 up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v, at
 20 least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and
 25 preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the
5 formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral
10 administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

15 It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when
20 measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v,
25 at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected
30 from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-

25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as
5 described in the US, British, European and Japanese pharmacopoeias).

Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

10 By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after
15 intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles.
- 5 The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.
2. Our findings from studies using ^{14}C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of
10 administration.

It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra
15 Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended
20 release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

By use of the term “therapeutically significant levels” we mean that blood plasma concentrations of at least 2.5 ngml^{-1} , ideally at least 3 ngml^{-1} , at least 8.5 ngml^{-1} , and up to 12 ngml^{-1} of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less
25 than 15 ngml^{-1} .

By use of the term “extended release” we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks)
30 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a
5 good release profile or lack of precipitation of drug after injection at the injection site.

Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor
10 oil.

Table 3**Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C**

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

5 **Solubility comparisons of fulvestrant in oil based formulations with and without alcohols**

		Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
10	Formulation ^(a)	Complete vehicle	Vehicle minus alcohols
	Castor oil based	81.2	12.6
15	Miglyol 812-N based	86.8	1.7
	Sesame seed/Castor oil (1:1) based	70.1	4.4
	Sesame seed oil based	45.7	0.7
20	Arachis oil based	40.2	< 0.2

25 ^(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

		Days						
30	Formulation ^a	2	3	4	7	10	30	51
35	Formulation F1 castor oil based	0	0	0	0	0	0	0
	Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
40	Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

45 ^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile over
5 five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

10 Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per
15 volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

20 A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-
25 acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

Preferably 5ml of the intramuscular injection is administered.

A further feature of the invention is use of fulvestrant in the preparation of a
30 pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent
5 indications such as breast cancer and gynaecological conditions, such as endometriosis.

In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical
10 structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a
15 pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

20 Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

25 Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay
30 as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary

pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram below

5

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol


10 10% weight per volume of ethanol

15% weight per volume of benzyl benzoate

250mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil

FLOW DIAGRAM OF MANUFACTURING



References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids* 1989; 54:71-99.
- 5 2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- 10 4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112:
15 R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

Claims

1. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of
5 formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration of at least 2.5ngml^{-1} for at least 2 weeks.
- 10 2. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for at least 4 weeks.
3. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for 2 to 5 weeks.
- 15 4. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of fulvestrant.
20
5. A pharmaceutical formulation as claimed in claim 1 to 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.
6. A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of
25 a pharmaceutically-acceptable alcohol.
7. A pharmaceutical formulation as claimed in any claim from 1 to 6 which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 30 8. A pharmaceutical formulation as claimed in claim 7 which contains 50%w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent .

9. A pharmaceutical formulation as claimed in claim 7 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 5 10. A pharmaceutical formulation as claimed in claim 7 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
11. A pharmaceutical formulation as claimed in claim 7 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 10 12. A pharmaceutical formulation as claimed in claim 7 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
13. A pharmaceutical formulation as claimed in claim 7 which contains 25% w/v or less
15 of a pharmaceutically-acceptable non-aqueous ester solvent.
14. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- 20 15. A pharmaceutical formulation as claimed in any claim from 1 to 14 wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
16. A pharmaceutical formulation as claimed in any claim from 1 to 15 wherein the
25 pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
17. A pharmaceutical formulation as claimed in any claim from 1 to 16 wherein the total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.

30

18. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 5 19. A pharmaceutical formulation as claimed in claim 18 wherein the total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5 to 5.25ml.
20. A pharmaceutical formulation as claimed in any of claims 1-19 wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of
10 formulation, 10% weight of benzyl alcohol per volume of formulation and 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.
21. A method of treating a benign or malignant diseases of the breast or reproductive tract by administration to a human in need of such treatment by intramuscular a pharmaceutical
15 formulation as claimed in claims 1 to 19.
22. A method as claimed in claim 21 for treating breast cancer.
23. A syringe or vial containing a pharmaceutical formulation as defined in claim 20.

20

ABSTRACT OF THE DISCLOSURE

The invention relates to a novel sustained release pharmaceutical formulation adapted
5 for administration by injection containing the compound
 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more
particularly to a formulation adapted for administration by injection containing the compound
 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in
solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-
10 aqueous ester solvent which is miscible in the ricinoleate vehicle.




Figure 1

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Donald J. Bird
Attorney Docket Number:	056291-5004-02

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Utility application filing	1011	1	330	330
Utility Search Fee	1111	1	540	540
Utility Examination Fee	1311	1	220	220
Pages:				
Claims:				
Claims in excess of 20	1202	22	52	1144
Independent claims in excess of 3	1201	2	220	440
Multiple dependent claims	1203	1	390	390

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Miscellaneous-Filing:				
Late filing fee for oath or declaration	1051	1	130	130
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				
Extension - 5 months with \$0 paid	1255	1	2350	2350
Miscellaneous:				
Total in USD (\$)				5544

Electronic Acknowledgement Receipt

EFS ID:	5447464
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	09629
Filer:	Donald J. Bird
Filer Authorized By:	
Attorney Docket Number:	056291-5004-02
Receipt Date:	04-JUN-2009
Filing Date:	15-OCT-2008
Time Stamp:	14:01:06
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Deposit Account
Payment was successfully received in RAM	\$5544
RAM confirmation Number	215
Deposit Account	500310
Authorized User	

The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:

Charge any Additional Fees required under 37 C.F.R. Section 1.16 (National application filing, search, and examination fees)

Charge any Additional Fees required under 37 C.F.R. Section 1.17 (Patent application and reexamination processing fees)

File Listing:					
Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1	Transmittal Letter	056291-5004-02-IDS-First.pdf	81302 eade4c534e276bda9fa6f9e5b4e8332d9142e0aa	no	2
Warnings:					
Information:					
2	Information Disclosure Statement (IDS) Filed (SB/08)	056291-5004-02-1449-First.pdf	116751 e68383a4eb651aaf482cbbaf6d53e829038a4d20	no	3
Warnings:					
Information:					
This is not an USPTO supplied IDS fillable form					
3	Foreign Reference	WO99027906.pdf	1201771 17eb0e2105e6de09d3b59735d456b173d15d799f	no	30
Warnings:					
Information:					
4	NPL Documents	PharmaceuticalDosageForms.pdf	321460 39ff9c71d0af795f5437c1a25f3481319563e66f	no	5
Warnings:					
Information:					
5	Transmittal Letter	056291-5004-02-IDS-Second.pdf	81796 fd36687d33e20f7fd5ff580e7099a596bc13416e	no	2
Warnings:					
Information:					
6	Information Disclosure Statement (IDS) Filed (SB/08)	IDSonControlledConfidentialNonCommercialTesting.pdf	792753 cb3373ff7c2ce4bc041ffd0dad5f368f09c2c590	no	19
Warnings:					
Information:					
This is not an USPTO supplied IDS fillable form					
7	Transmittal Letter	056291-5004-02-IDS-Third.pdf	82365 d9be9d39c492d49ab6b6d49daa15f9c47d949b80	no	2
Warnings:					
Information:					
8	Information Disclosure Statement (IDS) Filed (SB/08)	056291-5004-02-1449-Third.pdf	113011 9e24bc0dd8cf2f5039110bd1967ef93c8b7c1038	no	1
Warnings:					

Information:					
This is not an USPTO supplied IDS fillable form					
9	Foreign Reference	Third-IDS-Attachment_1_EP1250138B1.pdf	1033234 81ddb4f194370660cd8dc7eddbe333789f cfe55	no	22
Warnings:					
Information:					
10	Foreign Reference	Third-IDS-Attachment_2_EP1250138B1- Opposition.pdf	13564969 40565840d23255482e222a3baf07ab64d13 d70a3	no	323
Warnings:					
Information:					
11	Foreign Reference	Third-IDS-Attachment_3_EPSearchReport.pdf	302425 dca31854f20a13359152eac5a9511c65b2d 48cd5	no	10
Warnings:					
Information:					
12		056291-5004-02- PreliminaryAmendment.pdf	93701 ca09efc3357d1e9ee8a65e5db4da159c494 00150	yes	6
	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Preliminary Amendment		1	1	
	Claims		2	5	
	Applicant Arguments/Remarks Made in an Amendment		6	6	
Warnings:					
Information:					
13	Applicant Response to Pre-Exam Formalities Notice	056291-5004-02- StatementAccompanying- SubstituteSpecification.pdf	92679 22f9f3beba00152b5b15cbb47600c4cea46 332b0	no	2
Warnings:					
Information:					
14		056291-5004-02- SubstituteSpecification- CleanCopy.pdf	109710 5e1d509a92fd52412d6fa8ccd3b6124be3d af91f	yes	25
	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Specification		1	20	

	Claims	21	23
	Abstract	24	24
	Drawings-only black and white line drawings	25	25

Warnings:

Information:

15	Specification	056291-5004-02-SubstituteSpecification-MarkedCopy.pdf	111551 <small>eaca016da35e65b59d55b4252f29a81707b065b5</small>	no	25
----	---------------	---	---	----	----

Warnings:

Information:

16	Fee Worksheet (PTO-875)	fee-info.pdf	42584 <small>cd28591b73ad539e0e4d6b6b16512f68460b0231</small>	no	2
----	-------------------------	--------------	--	----	---

Warnings:

Information:

Total Files Size (in bytes):			18142062
-------------------------------------	--	--	----------

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Examiner: Unassigned
)	
FOR: FORMULATION)	Date: June 4, 2009

FIRST INFORMATION DISCLOSURE STATEMENT

UNDER 37 C.F.R. § 1.97(b)

Pursuant to 37 C.F.R. §§ 1.56 and 1.97(b), Applicants request that the Examiner consider this Information Disclosure Statement and the documents listed on the attached Form PTO-1449. To the best of the undersigned's knowledge, this Information Disclosure Statement is being filed before the mailing date of a first Office Action on the merits in the above-referenced application. Accordingly, Applicants do not believe that a fee is due for filing this Information Disclosure Statement.

With the exception of documents 38 and 65, copies of the listed documents were previously submitted or cited by the Examiner in parent Application No. 10/872,784. A copy of each of documents 38 and 65 are attached. Applicants respectfully request that the Examiner initial and return the Form PTO-1449, indicating that the information has been considered and made of record herein.

This submission does not represent that a search has been made or that no better art exists and does not constitute an admission that each or all of the listed documents are material or constitute "prior art." Applicants reserve the right to take appropriate action to establish the patentability of the disclosed invention over the listed documents, should one or more of the documents be applied against the claims of the present application.

Except for issue fees payable under 37 C.F.R. §1.18, the Commissioner is hereby authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 C.F.R. §§1.16 and 1.17 which may be required,

including any required extension of time fees, or credit any overpayment to Deposit Account No. 50-0310. This paragraph is intended to be a **CONSTRUCTIVE PETITION FOR EXTENSION OF TIME** in accordance with 37 C.F.R. §1.136(a)(3).

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: **June 4, 2009**
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000

By: /Donald Bird/
Donald J. Bird
Registration No. 25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

RECEIVED

PATENT
SEP 17 2002
TECH CENTER 1600/2900

ATTORNEY DOCKET NO.: 09756291-5004

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE



re PATENT APPLICATION of:

EVANS et al.

Appln. No.: 09/756,291

Filed: January 9, 2001

FOR: FORMULATION

)
) Group Art Unit: 1617
)
) Examiner: Hui, San-ming
)
)
)
)
)

September 13, 2002

Commissioner of Patents
Washington, D.C. 20231

Sir:

SECOND INFORMATION DISCLOSURE STATEMENT

Applicant wishes to make of record the following circumstances regarding the controlled, confidential and non-commercial testing of compositions meeting the definition of "pharmaceutical formulation", as used in the present method of treatment claims, which was carried out in the United States more than one year before the filing date of the present application in preparation for and during the testing (IND) phase of the regulatory review of such formulation by the FDA.

1. The elected invention as presently claimed is directed toward a method for treating a benign or malignant disease of the breast or reproductive tract of a human by intramuscular injection of a particular pharmaceutical formulation comprising the active drug fulvestrant in a vehicle comprising ricinoleate, a pharmaceutically-acceptable alcohol, and a pharmaceutically-acceptable non-aqueous ester solvent miscible in ricinoleate, as detailed in the claims.

09/16/2002 BABRAHA1 00000047 500310 09756291

05 FD:125 180.00 CH

2. Fulvestrant is the international non-proprietary (generic) name for the compound 7-alpha-[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]estra-1,3,5(10)-triene-3,17-beta-diol, which compound is encompassed by claims of U.S. Patent No. 4,659,516 issued to Bowler *et al.* in 1987 (hereinafter the "Bowler '516 patent").
3. The present specification acknowledges that fulvestrant is included among the steroid derivatives disclosed in European Patent Application No. 0 138 504 (corresponding to the Bowler '516 patent) as being effective antioestrogenic agents. The Bowler '516 patent notes at the bottom of column 7 that compositions of the disclosed steroid derivatives may be in a form suitable for oral or parenteral administration, and that compounds having antioestrogenic effect may have value in the treatment of, *e.g.*, anovulatory infertility, breast tumors and menstrual disorders.
4. However, certain characteristics of fulvestrant make it very difficult to formulate a pharmaceutically acceptable and effective composition for administration to humans. In particular, fulvestrant is an extremely lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low, placing severe limitations on the manner and mechanism by which it can be administered.
5. Subsequent to grant of the Bowler '516 patent, applicants developed an injectable extended release formulation of fulvestrant by which it became feasible to effectively utilize the known pharmacological properties of fulvestrant in the treatment of benign or malignant diseases of the breast or reproductive tract in humans, as presently claimed. This injectable extended release formulation of fulvestrant was subjected to extensive *in*

vitro and *in vivo* testing in animals, and eventually in clinical trials as detailed below, leading up to the first FDA approval of this formulation in April 2002.

6. In brief chronology, fulvestrant was initially put into development by Imperial Chemical Industries PLC (hereinafter "ICI"), under the product designation ICI 182,780. Development of fulvestrant was continued by Zeneca Limited (formed from ICI in 1993) under the product designation ZD9238. By December 6, 1996, preliminary testing of an injectable formulation containing fulvestrant as active ingredient had progressed to the point that an IND (Investigational New Drug) application was filed with the FDA for FASLODEX[®] (fulvestrant) Injection. As of the January 5, 1997 effective date of the IND application, clinical testing could, for the first time, commence in human subjects in the United States.
7. Clinical testing under the IND continued on behalf of AstraZeneca (formed by merger in 1999) until it was believed that sufficient evidence of safety and efficacy of the formulation had been obtained, and on March 28, 2001 an NDA (New Drug Application) was submitted to the FDA. Meanwhile, the subject application for patent, Application No. 09/756,291, was filed in the United States on January 9, 2001, claiming priority from GB Application 0000313.7, filed January 10, 2000, and GB Application 0008837.7, filed April 12, 2000. Thereafter, on April 25, 2002, the NDA for the Faslodex (injectable fulvestrant formulation) was approved by the FDA, whereupon Faslodex was approved for commercial marketing for the treatment of certain breast cancers.
8. The injectable fulvestrant formulation constituting Faslodex comes within the definition of "pharmaceutical formulation" as used in the method of treatment claims presently

pending in this application. The April 25, 2002 approval date constitutes the earliest possible date for commercial marketing in the United States of a formulation for use in accordance with the present claims.

9. This FDA approval came after the present application was filed, and was the culmination of many years of testing and gathering of data on the injectable formulation of fulvestrant (ICI 182,780 or ZD9238), both in the United States and abroad, in animals and eventually in human clinical trials. As will be evident below, all such testing in the United States more than one year before this application was filed was carried out under agreements which imposed obligations of confidentiality on the involved institutions and/or investigators, gave AstraZeneca strict control over the permitted use and disposition of the test samples of formulation, and provided that AstraZeneca was entitled to all information or data derived from the testing.¹ Moreover, all persons enrolled in the clinical trials were advised of the experimental nature of the formulation, and acknowledged this in signed informed consent forms as a precondition to their enrollment. AstraZeneca received no payment for the samples, and was not otherwise compensated for the use of these samples in the clinical trials. Under these conditions and the applicable case law discussed later below, these tests of the fulvestrant formulation in the United States did not constitute a “public use” under 35 U.S.C. § 102(b) of the present invention because the tests were carried out under strict obligations of confidentiality, and the tests and the use and disposition of the formulation, remained

¹ Reference to AstraZeneca hereinafter should be understood to refer to AstraZeneca and/or its predecessors in interest, ICI and Zeneca, unless the context indicates otherwise.

under the control of AstraZeneca throughout the entire period. These tests did not place the formulation in the public domain or cause the public to believe that the formulation of the invention was freely available, and certainly did not constitute a commercial exploitation of the invention more than one year before this application was filed.

10. Prior to the January 5, 1997 effective date of the IND application, all testing of fulvestrant formulation in the United States was necessarily carried out *in vitro* or in animals, and therefore cannot come within the scope of the present method of treatment claims. Nevertheless, it should be noted that all such testing was carried out under strict conditions of confidentiality and limitations of use imposed by a Statement of Proposed Investigation (SOPI) form that each investigator was required to sign as a condition to receiving samples of fulvestrant formulation.

11. The SOPI forms used by ICI in the early 1990s required a statement of proposed use of the material (necessarily not including any use in humans) that had to be approved by ICI, and stated just above the required signature of the investigator:

“If samples are supplied, I undertake:-

1. to make available all results to ICI;
2. that the results will not be submitted for publication or disclosed in any other way prior to disclosure to ICI;
3. to use the samples only for the purposes described above and not to pass the samples or any portion thereof to any investigators for any other purpose;
4. not to use the samples for any commercial purpose or for any study requested by a commercial organization.”

12. The SOPI forms used by Zeneca and AstraZeneca (even after the effective date of the IND, for any samples provided to investigators outside of formal protocols for clinical or compassionate use trials discussed below) similarly required a statement of proposed use of the material that had to be approved by Zeneca or AstraZeneca, and explicitly stated, "Laboratory studies/tests on animals only. (Not for human use)." Again, just above the signature of the investigator, the following undertaking was printed on each form:

- “1. All results acquired as a direct result of the use of the sample(s) will be promptly furnished to AstraZeneca.
2. The results will not be submitted for publication or disclosed in any other way without prior consent from AstraZeneca, which will not be unreasonably withheld.
3. The sample(s) will only be used for the purpose described above and shall not be passed to a third party. Any unused material will be returned to AstraZeneca.
4. The sample(s) will not be used to support the development of any commercial product containing the compound(s) supplied by AstraZeneca.
5. AstraZeneca shall be granted first option of a license to all rights in any discoveries or inventions made as a direct result of the investigations described above (whether patentable or not). In particular, the option will include an option for a license under any patents and patent applications relating to the use of the sample(s).
6. AstraZeneca requires assurance from all external investigators that all studies carried out on behalf of AstraZeneca and/or involving AstraZeneca compounds are carried out in compliance with all animal welfare legislation, regulations and policies applicable in that country/state. Please let us have your confirmation in writing that this is the case. We would also like to receive any additional

information on your in-house animal welfare arrangements which you are able to provide.”

13. It is understood that no investigator receiving fulvestrant formulation pursuant to a SOPI, at least in the United States and prior to the filing of the present application for patent, was informed of the components and/or proportions thereof constituting the injectable vehicle in which the fulvestrant was carried, and that no investigator publication of results approved by AstraZeneca included such a disclosure.
14. Two clinical studies involving Faslodex were carried out at least in part in the United States prior to the filing date of the present application for patent.
 - Clinical Study 9238IL/0021 began, in the United States, in April 1997 and extended to June 2000; was carried out in 69 centers involving 414 patients; and had the objective of comparing the effect, in terms of time to progression, of two doses of Faslodex (125 and 250 mg) with one dose of Arimidex (1 mg) in postmenopausal women with advanced breast cancer.
 - Clinical Study 9238IL/0025 began, in the United States, in November 1998 and extended to July 2001; was carried out in 32 centers involving 51 patients; and had the objective of comparing the effect, in terms of time to progression, of Faslodex (250 mg) with Nolvadex (20 mg) as first-line therapy in postmenopausal women with advanced breast cancer.
15. Each clinical study was carried out under a Clinical Study Agreement entered into by each Institution and Investigator taking part in the study.

16. A representative Clinical Study Agreement for Clinical Study 9238IL/0021 provided in relevant part:

“The clinical Study to be performed pursuant to this Agreement shall be that set forth in the Protocol entitled “A Double-blind, Randomized, Multicenter Trial comparing the Efficacy and Tolerability of 125 and 250 mg of FASLODEX™ (Long-acting ICI 182,780) With 1 mg ARIMIDEX™ (Anastrozole) in Postmenopausal Women With Advanced Breast Cancer” (hereinafter referred to as “Protocol”). Institution shall use its best efforts to ensure that the work required under the Protocol is properly performed in accordance therewith.”

* * * * *

“ZENECA reserves the right to terminate this Agreement and Study at any time in its sole discretion upon thirty (30) days prior written notice. However, ZENECA may terminate this agreement upon five (5) days written notice for safety, regulatory or ethical reasons. In the event of termination, all unused Study materials shall be returned to ZENECA and ZENECA shall reimburse Institution for all actual costs reasonably incurred up until the effective termination date.”

* * * * *

“All rights to all data, inventions or discoveries Institution may make or conceive in the course of their work for ZENECA in their performance under this Agreement and using product in accordance with the detailed protocol provided by ZENECA will be the property of ZENECA and will be assigned to ZENECA, and Institution will assist ZENECA, at ZENECA’s expense, by executing rightful papers for obtaining proper patent protection in such inventions or discoveries in any country which ZENECA at ZENECA’s option, desires to obtain patent protection. All control of and decisions regarding such patent filings and prosecution, whether U.S. or foreign, and all costs and fees associated therewith, shall be exercised and/or borne by ZENECA.”

* * * * *

“It may be necessary for Zeneca to disclose to Institution certain information considered proprietary or confidential (hereinafter ‘Confidential Information’) to aid Institution in effecting or completing their performance under this Agreement. Institution agrees to maintain in confidence all Confidential Information Institution obtains from ZENECA relating to this Agreement and not to disclose any of said Confidential Information to a third party for a period of three (3) years after the termination of this Agreement without the prior written consent of ZENECA. Notwithstanding the foregoing, it is understood that Confidential Information shall not include the following: (i) information that is now publicly available, (ii) information that later becomes publicly available, after it has become publicly available, (iii) information which Institution obtain from some third party not under any obligation to ZENECA with respect to such information, or (iv) information which Institution already have in their possession, prior to any disclosure by ZENECA, as evidenced by written records, (v) is independently developed by Institution or (vi) is required by law or regulation to be disclosed, provided, however, that Institution notifies and consults with Zeneca prior to such disclosure.

“Subject to the provisions of confidentiality set forth in Section 6(d) above, ZENECA agrees to grant Institution the right to publish its findings in the scientific literature, provided that ZENECA shall have the right to review, at least 30 days prior to submission for publication, copies of any and all final draft manuscripts which are authored or co-authored by Institution or by anyone in their research group and which are based in whole or in part on research conducted under this Agreement. Upon request by ZENECA, in order to protect intellectual property rights, Institution agrees to delay submission of such final draft manuscripts for publication for a period not exceeding six (6) months from the date on which ZENECA receives such final draft manuscripts. Institution agrees to implement any reasonable suggestions made to preserve ZENECA’s

right in its Confidential Information before any disclosure for publication or presentation; Investigator and Institution agrees to take appropriate cognizance of any other suggestions by ZENECA before any disclosure for publication or presentation.”

17. A representative Clinical Study Agreement for Clinical Study 9238IL/0025 similarly provided in relevant part:

“The clinical Study to be performed pursuant to this Agreement shall be that set forth in the Protocol which is attached hereto as Exhibit A and incorporated herein by reference. Institution and Investigator shall use their best efforts to ensure that the work required under the Protocol is properly performed in accordance therewith.”

* * * * *

“Zeneca reserves the right to terminate this Agreement and Study at any time in its sole discretion upon five (5) days prior written notice. In the event of termination, all unused Study materials shall be returned to Zeneca and Zeneca shall reimburse Institution and Investigator for all actual costs reasonably incurred up until the effective termination date.”

* * * * *

“All rights to all data, inventions or discoveries Institution and Investigator may make or conceive in the course of their work for Zeneca in their performance under this Agreement will be the property of Zeneca and will be assigned to Zeneca, and Institution and Investigator will assist Zeneca, at Zeneca’s expense, by executing rightful papers for obtaining proper patent protection in such inventions or discoveries.”

* * * * *

“It may be necessary for Zeneca to disclose to Investigator and Institution certain information considered proprietary or confidential (hereinafter “Confidential

Information”) to aid Investigator and Institution in effecting or completing their performance under this Agreement. Confidential Information shall also include Study data; however, Investigator’s and Institution’s right to publish pursuant to Section (d) below* shall not be affected by this provision. Investigator and Institution agree to maintain in confidence all Confidential Information Investigator and Institution obtain from Zeneca relating to this Agreement and not to disclose any of said Confidential Information to a third party without the prior written consent of Zeneca. Notwithstanding the foregoing, it is understood that Confidential Information shall not include the following: (i) information that is now publicly available, (ii) information that later becomes publicly available, after it has become publicly available, (iii) information which Investigator and Institution obtain from some third party not under any obligation to Zeneca with respect to such information, or (iv) information which Investigator and Institution already have in their possession, prior to any disclosure by Zeneca, as evidenced by written records.

“Nothing herein shall prevent Investigator and Institution from complying with the legal obligation to disclose Confidential Information so long as Investigator and Institution (i) provide Zeneca prompt notice of its intent to disclose (or to resist disclosure) (ii) take reasonable steps to require the recipient to preserve the confidential nature of the information once disclosed and (iii) afford Zeneca the opportunity to attempt to prevent the disclosure (whether or not Investigator and Institution have sought to resist disclosure) or obtain protection for the information disclosed.”

* * * * *

*[(d)] “Subject to the provisions of confidentiality set forth in Section 6(c) above, Zeneca agrees to grant Investigator and Institution the right to publish their findings in the scientific literature, provided that Zeneca shall have the right to review, at least 30 days prior to submission for publication, copies of any and all final draft manuscripts which are authored or co-authored by Investigator and

Institution or by anyone in their research group and which are based in whole or in part on research conducted under this Agreement. In the event it is necessary for Zeneca to prepare a patent application(s) and other documentation, and upon request by Zeneca, Investigator and Institution agree to delay submission of such final draft manuscripts for publication for a period not exceeding six (6) months from the date on which Zeneca receives such final draft manuscripts. Investigator and Institution agree to implement any reasonable suggestions made to preserve Zeneca's right in its Confidential Information before any disclosure for publication or presentation; Investigator and Institution agree to take appropriate cognizance of any other suggestions by Zeneca before any disclosure for publication or presentation."

* * * * *

"Zeneca shall be entitled to make copies, at Zeneca's expense, of any and all documents and data generated from the Study. In addition, Institution and Investigator agree to allow Zeneca to audit the Study records (including administrative files and source documents such as hospital charts, office records and written results of laboratory and diagnostic tests) of Institution and Investigator at mutually convenient times.

18. An additional clinical study involving Faslodex was commenced in the United States more than one year prior to the filing date of the present application for patent, being Clinical Study 9238IL/0037, a compassionate-use trial under a protocol initially entitled "An Open-label, Treatment-use Protocol of 250 mg of FASLODEX™ (Long-acting ICI 182,780) in Postmenopausal Women With Advanced Breast Cancer." It is understood that as of one year prior to the filing date of this application, seven subjects had been enrolled in Clinical Study 9238IL/0037.

19. A “Confidentiality and Proprietary Rights Agreement” was entered into by each Investigator prior to his involvement in Clinical Study 9238IL/0037, in which the Investigator acknowledged that “he will have access to and obtain knowledge of certain proprietary and confidential Information of Zeneca and that as a condition of receiving such information” the parties agreed, in part as here relevant:

“1. ‘Confidential Information’ shall mean all information (a) disclosed by Zeneca to Investigator, either orally or in writing or (b) obtained by the Investigator from a third party or any other source, regarding the protocol entitled ‘An Open-label, Treatment-use Protocol of 250 mg of FASLODEX™ (Long-acting ICI 182,780) in Postmenopausal Women With Advanced Breast Cancer, Study No. 9238IL/0037’ (‘Study’)

“Confidential Information shall not include information that: (i) was already in the possession of Investigator before disclosure thereof by Zeneca to Investigator as evidenced by Investigator’s written records (ii) is independently developed by Investigator as evidenced by Investigator’s written records, (iii) is or becomes publicly available through no fault of Investigator, or (iv) is obtained by Investigator from a third party under no obligation not to disclose same.

“Nothing herein shall prevent Investigator from complying with a legal obligation to disclose Confidential Information so long as Investigator (i) provides Zeneca prompt notice of its intent to disclose (or to resist disclosure) (ii) takes reasonable steps to require the recipient to preserve the confidential nature of the information once disclosed and (iii) affords Zeneca the opportunity to attempt to prevent the disclosure (whether or not Investigator has sought to resist disclosure) or obtain protection for the Information disclosed.

“2. The purpose of the disclosure of Confidential Information is to allow Investigator to participate in the Treatment-use Protocol.

“3. Investigator agrees to maintain in strictest confidence and to take all reasonable steps to maintain the confidentiality of the Confidential Information. Investigator also agrees not to disclose Confidential Information to any third party, and to use Confidential Information only for the purposes stated in paragraph 2 of this Agreement.

“4. Investigator recognizes that all documents and records received by Investigator from Zeneca and all copies of such records and documents shall be Zeneca’s property exclusively. The Investigator shall at all times keep all such documents, records and copies of documents and records in Investigator’s custody and subject to Investigator’s control and shall surrender the same upon request by Zeneca.

”5. Investigator shall not disclose any Confidential Information to any of its employees, except employees of Investigator who have a need to know the Confidential Information for the purposes stated in paragraph 2 of this Agreement and who have assumed an obligation to maintain Zeneca’s Confidential Information in confidence at least to the extent that Investigator is bound hereunder. Investigator shall advise each such employee of the confidential nature of the Confidential Information received from Zeneca and the existence and importance of the confidentiality provisions of this Agreement and shall be responsible for ensuring that such employees maintain the Confidential Information in confidence in accordance with the terms of this Agreement.

“6. Because of the unique nature of the Confidential Information, Investigator understands and agrees that Zeneca will suffer irreparable harm in the event that Investigator fails to comply with any of its obligations contained hereinabove and that monetary damages will be inadequate to compensate Zeneca for such breach. Accordingly, Investigator agrees that Zeneca shall have the right to seek immediate injunctive relief to enforce the confidentiality obligations contained herein.

“7. All rights to all data, inventions or discoveries Investigator may make or conceive in the course of Investigator participation in the Study will be the property of Zeneca and will be assigned to Zeneca, and Investigator will assist Zeneca, at Zeneca’s expense, by executing rightful papers for obtaining proper patent protection in such inventions or discoveries. Investigator agrees to make no claim which will restrict the rights of Zeneca to use and disclose to others any information, knowledge, and ideas which are disclosed to Zeneca by Investigator in the course of performance of the Study.

“8. Subject to the provisions of confidentiality set forth herein, Zeneca agrees to grant Investigator the right to publish his findings in the scientific literature, provided that Zeneca shall have the right to review, at least 30 days prior to submission for publication, copies of any and all final draft manuscripts which are authored by Investigator or by anyone in his research group and which are based in whole or in part on research conducted pursuant to this Study. In the event it is necessary for Zeneca to prepare a patent application(s) and other documentation, and upon request by Zeneca, Investigator agrees to delay submission of such final draft manuscripts for publication for a period not exceeding six (6) months from the date on which Zeneca receives such final draft manuscripts. Investigator agrees to implement any reasonable suggestions made to preserve Zeneca’s right in its Confidential Information before any disclosure for publication or presentation; Investigator agrees to take appropriate cognizance of any other suggestions by Zeneca before any disclosure for publication or presentation.”

20. The Protocols referenced with respect to the above-noted Studies No. 9238IL/0021, No. 9238IL/0025 and No. 9238IL/0037 provided details of, *inter alia*, the:

- criteria for the selection and screening for eligibility of subjects for entry into the trial, as well as exclusion criteria;

- route, dose and regimen for administration of the respective drugs to individual subjects;
- procedures for drug accountability, including maintenance of accurate records on receipt and disposition of investigational materials, and return or destruction of any unused drug;
- frequency and procedures for clinical and laboratory evaluations;
- regular recordation of data on case report forms, record retention and submission of records to AstraZeneca; and
- trial monitoring and data verification by representatives of AstraZeneca.

21. These Protocols furthermore required that each subject be given appropriate information on the treatment prior to its commencement, including the experimental aspects of the treatment and the risks involved, and sign an informed consent form approved by AstraZeneca, and conforming to the requirements of 21 C.F.R. 50.20 *et seq.* , which requires as a basic element of informed consent, that each subject be provided with, *inter alia*, a “statement that the study involves research, an explanation of the purposes of the research and the expected duration of the subject's participation, a description of the procedures to be followed, and identification of any procedures which are experimental.” 21 C.F.R. 50.25(a)(1).

In evaluating the above circumstances in context of 35 U.S.C. § 102(b), the Examiner’s attention is called to MPEP ¶ 2133.03 “Rejections Based on ‘Public Use’ or ‘On Sale’”, and particularly MPEP ¶ 2133.03(a) “Public Use”, section *B.* headed “*Use by Third*

Parties Deriving the Invention from Applicant.” It is respectfully submitted that the above circumstances *do not* constitute a “public use” of the presently claimed invention under the criteria set forth in the MPEP, and as established by decisions of the Federal Circuit, because of the strict confidentiality and control imposed and maintained by AstraZeneca throughout the relevant trial periods. MPEP ¶ 2133.03(a)B. provides:

An Invention Is in Public Use If the Inventor Allows Another To Use the Invention Without Restriction or Obligation of Secrecy

"Public use" of a claimed invention under 35 U.S.C. 102(b) occurs when the inventor allows another person to use the invention without limitation, restriction or obligation of secrecy to the inventor." *In re Smith*, 714 F.2d 1127, 1134, 218 USPQ 976, 983 (Fed. Cir. 1983). The presence or absence of a confidentiality agreement is not itself determinative of the public use issue, but is one factor to be considered along with the time, place, and circumstances of the use which show the amount of control the inventor retained over the invention. *Moleculon Research Corp. v. CBS, Inc.*, 793 F.2d 1261, 1265, 229 USPQ 805, 809 (Fed. Cir. 1986). See *Ex parte C*, 27 USPQ2d 1492, 1499 (Bd. Pat. App. & Inter. 1992) (Inventor sold inventive soybean seeds to growers who contracted and were paid to plant the seeds to increase stock for later sale. The commercial nature of the use of the seed coupled with the "on-sale" aspects of the contract and apparent lack of confidentiality requirements rose to the level of a "public use" bar.); *Egbert v. Lippmann*, 104 U.S. 333, 336 (1881) (Public use found where inventor allowed another to use inventive corset insert, though hidden from view during use, because he did not impose an obligation of secrecy or restrictions on its use.).

The samples of fulvestrant formulation provided under the SOPI forms was not for human use, and therefore outside of the scope of the present method of use claims.

Nevertheless, the tests conducted on these samples by the third party Investigators did not constitute a “public use”. Through the SOPI forms, AstraZeneca maintained strict confidentiality over the samples and tests conducted therewith, maintained control over the use and disposition of the samples, and was entitled to all data developed in the course of the

tests. (¶¶ 10-13, *supra*). Moreover, AstraZeneca received no payment or other commercial benefit from providing these samples


Similarly, the three clinical trials or studies conducted in human subjects did not constitute a “public use” under the definition thereof set out in the MPEP as developed by the courts. Prior to the release of any materials or formulations on which to carry out these studies, the institutions and/or investigators involved were required to sign an agreement whereunder strict confidentiality was required, and all information provided to or developed by the institution/investigator during the course of such studies remained or became the property of AstraZeneca. (¶¶ 16, 17 and 19, *supra*). Through the Clinical Study Agreements, and the Protocols under which all three studies were conducted, AstraZeneca maintained full control over the use and disposition of the study materials or formulation that it provided to the institutions/investigators throughout the course of these studies, and the right to receive the data and records that were produced. (¶¶ 16, 17 and 20, *supra*). Moreover, each subject of these studies was fully informed of the experimental nature of the formulation and its use, as acknowledged in signed informed consent forms, and clearly did not have any basis to believe that the formulation or its use in the treatments was in the public domain or otherwise freely available. (¶ 21, *supra*). Again, AstraZeneca received no payment for the formulation used in these studies, and these studies did not constitute a commercial exploitation of the formulation.

Therefore, under the case law as developed by the courts, and its application by the Patent and Trademark Office as set out in the above-quoted paragraph from the MPEP, it is

respectfully submitted that the foregoing circumstances do not constitute a "public use"

under 35 U.S.C. § 102(b).

Respectfully Submitted,
Morgan Lewis & Bockius LLP



September 13, 2002
Morgan Lewis & Bockius LLP
Customer No. **009629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000
DJB:

By:

Donald J. Bird
Registration No. 25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Examiner: Unassigned
)	
FOR: FORMULATION)	Date: June 4, 2009

THIRD INFORMATION DISCLOSURE STATEMENT

UNDER 37 C.F.R. § 1.97(b)

Pursuant to 37 C.F.R. §§ 1.56 and 1.97(b), Applicants request that the Examiner consider this Information Disclosure Statement and the documents listed on the attached Form PTO-1449. To the best of the undersigned's knowledge, this Information Disclosure Statement is being filed before the mailing date of a first Office Action on the merits in the above-referenced application. Accordingly, Applicants do not believe that a fee is due for filing this Information Disclosure Statement.

Copies of the listed documents were previously submitted or cited by the Examiner in parent Application No. 10/872,784. Accordingly, no copies of the listed documents are provided herewith. Applicants respectfully request that the Examiner initial and return the Form PTO-1449, indicating that the information has been considered and made of record herein.

Also submitted with this Information Disclosure Statement as Attachments I to III are the following documents: Attachment I: A copy of EP 1250138B1, which is the European Patent that granted on the European counterpart of the subject Application; Attachment II: A copy of documents from the EPO file relating to the European Opposition pending against EP 1250138B1; and Attachment III: A copy the Supplementary European Search Report received by Applicant in European Application 05016921.8, which is a divisional of the application from which EP 1250138B1 granted.

This submission does not represent that a search has been made or that no better art exists and does not constitute an admission that each or all of the listed documents are material or

constitute "prior art." Applicants reserve the right to take appropriate action to establish the patentability of the disclosed invention over the listed documents, should one or more of the documents be applied against the claims of the present application.

Except for issue fees payable under 37 C.F.R. §1.18, the Commissioner is hereby authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 C.F.R. §§1.16 and 1.17 which may be required, including any required extension of time fees, or credit any overpayment to Deposit Account No. 50-0310. This paragraph is intended to be a **CONSTRUCTIVE PETITION FOR EXTENSION OF TIME** in accordance with 37 C.F.R. §1.136(a)(3).

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: **June 4, 2009**
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000

By: /Donald Bird/
Donald J. Bird
Registration No. 25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date
	1. US 3,164,520	January 5, 1965	Huber			
	2. US 4,212,863	July 15, 1980	Cornelius			
	3. US 4,388,307	June 14, 1983	Cavanak			

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
	4. EP 0310542A1	April 5, 1989	EPO			Yes

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

	5.	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
	6.	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
	7.	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
	8.	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
	9.	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 [pages 238-255 provided] (1998)
	10.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
	11.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
	12.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
	13.	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

Examiner	Date Considered
----------	-----------------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Third IDS

Attachment I



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 250 138 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
19.10.2005 Bulletin 2005/42

(51) Int Cl.⁷: **A61K 31/565**, A61P 35/00,
A61K 47/14, A61K 47/44

(21) Application number: **01900186.6**

(86) International application number:
PCT/GB2001/000049

(22) Date of filing: **08.01.2001**

(87) International publication number:
WO 2001/051056 (19.07.2001 Gazette 2001/29)

(54) **FULVESTRANT FORMULATION**
FULVESTRANT FORMULIERUNG
PREPARATION DE FULVESTRANT

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

• **GRUNDY, Rosalind, Ursula**
Macclesfield, Cheshire SK10 4TG (GB)

(30) Priority: **10.01.2000 GB 0000313**
12.04.2000 GB 0008837

(74) Representative: **Giles, Allen Frank et al**
AstraZeneca AB,
Global Intellectual Property
151 85 Sodertälje (SE)

(43) Date of publication of application:
23.10.2002 Bulletin 2002/43

(56) References cited:
EP-A- 0 346 014 WO-A-96/19997
WO-A-97/21440

(60) Divisional application:
05016921.8

• **JOHN C. WATERTON; ET AL.: "A Case of
Adenomyosis in a Pigtailed Monkey Diagnosed
by Magnetic Resonance Imaging and treated
with the Novel Pure Antiestrogen, ICI 182,780"**
LABORATORY ANIMAL SCIENCE, vol. 43, no. 3,
1993, pages 247-251, XP000998289

(73) Proprietor: **AstraZeneca AB**
151 85 Södertälje (SE)

(72) Inventors:
• **EVANS, John, Raymond**
Macclesfield, Cheshire SK10 4TG (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 250 138 B1

Description

[0001] The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

[0002] Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

[0003] An alternative approach to oestrogen withdrawal is to antagonise oestrogens with antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen (ER), compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

[0004] The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

[0005] Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989). One of these, 7α -[9-(4,4,5,5,5-pentafluoropentyl sulphinyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

[0006] 7α -[9-(4,4,5,5,5-Pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

[0007] Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

[0008] Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

[0009] In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

[0010] European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

[0011] Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml^{-1} (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

[0012] Currently there are a number of sustained release injectable steroidal formulations which have been com-

EP 1 250 138 B1

mercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

[0013] In the formulations within Table 1 a number of different oils are used to solubilise the compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

5

10

15

20

25

30

35

40

45

50

55

55 50 45 40 35 30 25 20 15 10 5

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

PRODUCT NAME	STEROID	DOSE	TYPE	COMP.	SOURCE	OIL	BzBz	BzOH	EtOH	DOSE	DOSING
SUSTANON 100	Testosterone propionate	30mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
	Testosterone phenylpropionate	60mg									
	Testosterone isocaproate	60mg									
	Testosterone decanoate	100mg									
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%		1 or 2ml	1 week	
TOCOGESTAN	Hydroxy progesterone enantate	200mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%		2ml	< 1week	
	Progesterone	50mg									
TROPHOBOLENE	α-Tocopherol	250mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%		1ml	15 to 30 days	
	Estrapronicate	1.3mg									
	Nandrolone undecanoate	50mg									
	Hydroxyprogesterone heptanoate	80mg									
NORISTERAT	Norethisterone oenanthoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES		1ml	8 weeks	
BENZO-GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis			1ml	1 week	
PROGESTERONE -RETARD	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES		1 or 2ml	1 week	
GRAVIBINAN	Estradiol 17-β-valerate	5mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES		1 or 2ml	1 - 2 weeks	
	Hydroxyprogesterone caproate	250mgml ⁻¹									

4

EP 1 250 138 B1

55

50

45

40

35

30

25

20

15

10

5

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis		75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%		
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%		

5

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
% are w/v and * approximate as measured directly from a single sample

EP 1 250 138 B1

[0014] In US 5,183,814 Example 3 an oil based injection formulation of fulvestrant is described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

[0015] Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 -

SOLUBILITY OF FULVESTRANT	
SOLVENT	SOLUBILITY
	(mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

[0016] As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

[0017] However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

[0018] Currently guidelines recommend that no more than 5mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

[0019] The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of >50mgml⁻¹ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of <5ml - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50 mgml⁻¹ - see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

[0020] Therefore, we present as a feature of the invention a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0021] Another feature of the invention is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0022] Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection

comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0023] Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

[0024] For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

[0025] Preferred pharmaceutical formulations of the invention are as described above wherein:

1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

[0026] It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

[0027] Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

[0028] The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

[0029] The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

[0030] The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

[0031] It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopoeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopoeia contains not less than 99.5% ethanol by volume

when measured at 15.56°C.

[0032] Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v, at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

[0033] It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

[0034] Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzy! alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

[0035] By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

[0036] We have surprisingly found that the above formulations of the invention provide, after intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

[0037] This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.

2. Our findings from studies using ¹⁴C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

[0038] It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

[0039] It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

[0040] We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period

can still achieved by the formulation of the invention.

[0041] By use of the term "therapeutically significant levels" we mean that blood plasma concentrations of at least 2.5 ngml⁻¹, ideally at least 3 ngml⁻¹, at least 8.5 ngml⁻¹, and up to 12 ngml⁻¹ of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 15 ngml⁻¹.

5 **[0042]** By use of the term "extended release" we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

10 **[0043]** It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

[0044] Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

15 **[0045]** Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

20

25

30

35

40

45

50

55

55
50
45
40
35
30
25
20
15
10
5

Table 3

Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C

	% w/v								
Ethanol (96%)	5	5	10	10	10	10	15	15	
Benzyl Alcohol	5	5	5	5	10	10	15	15	
Benzyl Benzoate		15		15		15		15	
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102	

10

EP 1 250 138 B1

[0046] The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

Solubility comparisons of fulvestrant in oil based formulations with and without alcohols		
Fulvestrant Solubility mg ml ⁻¹ @ 25°C		
Formulation (a)	Complete vehicle	Vehicle minus alcohols
Castor oil based	81.2	12.6
Miglyol 812-N based	86.8	1.7
Sesame seed/Castor oil (1:1) based	70.1	4.4
Sesame seed oil based	45.7	0.7
Arachis oil based	40.2	< 0.2

(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

Formulation ^a	Days						
	2	3	4	7	10	30	51
Formulation F1 castor oil based	0	0	0	0	0	0	0
Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

[0047] Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

[0048] Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

[0049] Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

[0050] A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

[0051] A further feature of the invention is the treatment of a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

[0052] Preferably 5ml of the intramuscular injection is administered.

[0053] A further feature of the invention is use of fulvestrant in the preparation of a pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably

treating breast cancer.

[0054] Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

[0055] As described above fulvestrant is useful in the treatment of oestrogen-dependent indications such as breast cancer and gynaecological conditions, such as endometriosis.

[0056] In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro-7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

[0057] A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro-7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro-7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

[0058] Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example


[0059] Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram below

[0060] Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol
 10% weight per volume of ethanol
 15% weight per volume of benzyl benzoate
 250mg of fulvestrant for each 5ml of finished formulation
 and the remaining amount as castor oil

FLOW DIAGRAM OF MANUFACTURING



References

[0061]

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. Steroids 989; 5471-99.

2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. American New York Academy Science 1990a; 595: 348-56.

3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. Regulatory mechanisms in breast cancer. Boston: Kluwer Academic, 1990b: 239-57.

4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. Journal Steroid Biochemistry 1990c; 37: 771-5.

5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. Journal Endocrinology 1987; 112: R7-10.

6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. Journal Steroid Biochemistry 1988;

3: 141-7.

Claims

- 5
1. A pharmaceutical formulation comprising fulvestrant in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.
 - 10
 2. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.
 - 15
 3. A pharmaceutical formulation as claimed in claim 1 or 2 wherein the blood plasma fulvestrant concentration attained is at least 2.5ngml^{-1} for at least 2 weeks..
 - 20
 4. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of fulvestrant.
 - 25
 5. A pharmaceutical formulation as claimed in claim 1 to 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.
 6. A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of a pharmaceutically-acceptable alcohol.
 - 30
 7. A pharmaceutical formulation as claimed in claim 5 which contains 15-25% w/v of a pharmaceutically acceptable alcohol.
 8. A pharmaceutical formulation as claimed in claim 5 which contains 17-23% w/v of a pharmaceutically acceptable alcohol.
 - 35
 9. A pharmaceutical formulation as claimed in any claim from 1 to 8 which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 - 40
 10. A pharmaceutical formulation as claimed in claim 9 which contains 50%w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 11. A pharmaceutical formulation as claimed in claim 9 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 - 45
 12. A pharmaceutical formulation as claimed in claim 9 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 13. A pharmaceutical formulation as claimed in claim 9 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 - 50
 14. A pharmaceutical formulation as claimed in claim 9 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 15. A pharmaceutical formulation as claimed in claim 9 which contains 25% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
 - 55
 16. A pharmaceutical formulation as claimed in claim 9 which contains 10-25% w/v of a pharmaceutically acceptable

non-aqueous ester solvent.

- 5
17. A pharmaceutical formulation as claimed in claim 9 which contains 12-18% w/v of a pharmaceutically acceptable non-aqueous ester solvent.
- 10
18. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 15-25% weight of a pharmaceutically-acceptable alcohol per volume of formulation, 10-25 % weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.
- 15
19. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 17-23% weight of a pharmaceutically-acceptable alcohol per volume of formulation, 12-18 % weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.
- 20
20. A pharmaceutical formulation as claimed in any claim from 1 to 19 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- 25
21. A pharmaceutical formulation as claimed in any claim from 1 to 20 wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
- 30
22. A pharmaceutical formulation as claimed in any claim from 1 to 21 wherein the pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
- 35
23. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 15-25% weight of a pharmaceutically-acceptable alcohol per volume of formulation, 10-25 % weight of benzyl benzoate in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.
- 40
24. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 17-23% weight of a pharmaceutically-acceptable alcohol per volume of formulation, 12-18 % weight of benzylbenzoate in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.
- 45
25. A pharmaceutical formulation according to claim 23 or 24 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- 50
26. A pharmaceutical formulation according to claim 25 wherein the ethanol and benzyl alcohol are present at about equal % weight per volume of formulation.
- 55
27. A pharmaceutical formulation as claimed in any claim from 1 to 26 wherein the total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
28. A pharmaceutical formulation as claimed in any claim from 1 to 27 wherein the total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
29. A pharmaceutical formulation as claimed in claim 28 wherein the total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5 to 5.25ml.
30. A pharmaceutical formulation as claimed in any of claims 1-29 wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of formulation, 10% weight of benzyl alcohol per volume of formulation, and the formulation contains 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.
31. A pharmaceutical formulation adapted for intramuscular injection, as defined in any claim from 1 to 30, for use in medical therapy.

32. Use of fulvestrant in the preparation of a pharmaceutical formulation, as defined in any claim from 1 to 30, for the treatment of a benign or malignant disease of the breast or reproductive tract.

33. A syringe or vial containing a pharmaceutical formulation as defined in claim 30.

5

Patentansprüche

1. Pharmazeutische Formulierung, enthaltend Fulvestrant in einer Ricinoleat-Trägersubstanz, ein pharmazeutisch annehmbares nichtwässriges Esterlösungsmittel und einen pharmazeutisch annehmbaren Alkohol, wobei die Formulierung zur intramuskulären Anwendung und Erzielung einer mindestens 2 Wochen anhaltenden, therapeutisch signifikanten Fulvestrantkonzentration im Blutplasma geeignet ist.
2. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 30 Gew.-% oder weniger eines pharmazeutisch annehmbaren Alkohols und mindestens 1 Gew.-% eines in einer Ricinoleat-Trägersubstanz mischbaren, pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels, sowie eine zur Herstellung einer nach Injektion zur Erzielung einer mindestens 2 Wochen anhaltenden, therapeutisch signifikanten Fulvestrantkonzentration im Blutplasma geeigneten Formulierung ausreichende Menge einer Ricinoleat-Trägersubstanz.
3. Pharmazeutische Formulierung nach Anspruch 1 oder 2, bei der die im Blutplasma erzielte Fulvestrantkonzentration mindestens 2 Wochen lang mindestens 2,5 ngml⁻¹ beträgt.
4. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 30 Gew.-% oder weniger eines pharmazeutisch annehmbaren Alkohols und mindestens 1 Gew.-% eines in einer Ricinoleat-Trägersubstanz mischbaren, pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels, sowie eine zur Herstellung einer Formulierung mit mindestens 45 mgml⁻¹ Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz.
5. Pharmazeutische Formulierung nach Anspruch 1 bis 4, die 25% w/v oder weniger eines pharmazeutisch annehmbaren Alkohols enthält.
6. Pharmazeutische Formulierung nach Anspruch 5, die 20% w/v oder weniger eines pharmazeutisch annehmbaren Alkohols enthält.
7. Pharmazeutische Formulierung nach Anspruch 5, die 15-25% w/v eines pharmazeutisch annehmbaren Alkohols enthält.
8. Pharmazeutische Formulierung nach Anspruch 5, die 17-23% w/v eines pharmazeutisch annehmbaren Alkohols enthält.
9. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 8, die 60% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
10. Pharmazeutische Formulierung nach Anspruch 9, die 50% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
11. Pharmazeutische Formulierung nach Anspruch 9, die 45% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
12. Pharmazeutische Formulierung nach Anspruch 9, die 40% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
13. Pharmazeutische Formulierung nach Anspruch 9, die 35% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
14. Pharmazeutische Formulierung nach Anspruch 9, die 30% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.

EP 1 250 138 B1

15. Pharmazeutische Formulierung nach Anspruch 9, die 25% w/v oder weniger eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
- 5 16. Pharmazeutische Formulierung nach Anspruch 9, die 10-25% w/v eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
- 10 17. Pharmazeutische Formulierung nach Anspruch 9, die 12-18% w/v eines pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels enthält.
- 15 18. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 15-25 Gew.-% eines pharmazeutisch annehmbaren Alkohols und 10-25 Gew.-% eines in einer Ricinoleat-Trägersubstanz mischbaren, pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels, sowie eine zur Herstellung einer Formulierung mit mindestens 45 mgml^{-1} Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz.
- 20 19. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 17-23 Gew.-% eines pharmazeutisch annehmbaren Alkohols und 12-18 Gew.-% eines in einer Ricinoleat-Trägersubstanz mischbaren, pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittels, sowie eine zur Herstellung einer Formulierung mit mindestens 45 mgml^{-1} Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz.
- 25 20. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 19, bei der es sich bei dem pharmazeutisch annehmbaren Alkohol um ein Gemisch aus Ethanol und Benzylalkohol handelt.
- 30 21. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 20, bei der das pharmazeutisch annehmbare nichtwässrige Esterlösungsmittel unter Benzylbenzoat, Ethyloleat, Isopropylmyristat, Isopropylpalmitat oder einem beliebigen Gemisch davon ausgewählt ist.
- 35 22. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 21, bei der es sich bei dem pharmazeutisch annehmbaren nichtwässrigen Esterlösungsmittel um Benzylbenzoat handelt.
- 40 23. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 15-25 Gew.-% eines pharmazeutisch annehmbaren Alkohols und 10-25 Gew.-% Benzylbenzoat in einer Ricinoleat-Trägersubstanz, sowie eine zur Herstellung einer Formulierung mit mindestens 45 mgml^{-1} Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz.
- 45 24. Pharmazeutische Formulierung zur intramuskulären Injektion, enthaltend Fulvestrant, jeweils bezogen auf das Volumen der Formulierung 17-23 Gew.-% eines pharmazeutisch annehmbaren Alkohols und 12-18 Gew.-% Benzylbenzoat in einer Ricinoleat-Trägersubstanz, sowie eine zur Herstellung einer Formulierung mit mindestens 45 mgml^{-1} Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz.
- 50 25. Pharmazeutische Formulierung nach Anspruch 23 oder 24, bei der es sich bei dem pharmazeutisch annehmbaren Alkohol um ein Gemisch aus Ethanol und Benzylalkohol handelt.
- 55 26. Pharmazeutische Formulierung nach Anspruch 25, bei der der gew.-%ige Anteil an Ethanol und Benzylalkohol pro Volumen Formulierung jeweils etwa gleich ist.
27. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 26, bei der das Gesamtvolumen der Formulierung 6 ml oder weniger und die Fulvestrantkonzentration mindestens 45 mgml^{-1} ausmachen.
28. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 27, bei der die Gesamtmenge an Fulvestrant in der Formulierung 250 mg oder mehr und das Gesamtvolumen der Formulierung 6 ml oder weniger ausmachen.
29. Pharmazeutische Formulierung nach Anspruch 28, bei der die Gesamtmenge an Fulvestrant in der Formulierung 250 mg und das Gesamtvolumen der Formulierung 5 bis 5,25 ml ausmachen.
30. Pharmazeutische Formulierung nach einem der Ansprüche 1 bis 29, bei der es sich bei dem pharmazeutisch annehmbaren Alkohol um ein Gemisch von, jeweils bezogen auf das Volumen der Formulierung, 10 Gew.-% Ethanol

nol und 10 Gew.-% Benzylalkohol handelt, und die Formulierung pro Volumen 15 Gew.-% Benzylbenzoat enthält, und es sich bei der Ricinoleat-Trägersubstanz um Rizinusöl handelt.

5 31. Pharmazeutische Formulierung zur intramuskulären Injektion gemäß Definition eines der Ansprüche 1 bis 30 zur Verwendung in der medizinischen Therapie.

10 32. Verwendung von Fulvestrant bei der Herstellung einer wie in einem der Ansprüche 1 bis 30 definierten pharmazeutischen Formulierung zur Behandlung gutartiger oder bösartiger Erkrankungen der Brust oder des Reproduktionstrakts.

33. Spritze oder Fläschchen, enthaltend eine wie in Anspruch 30 definierte pharmazeutische Formulierung.

Revendications

15 1. Préparation pharmaceutique comprenant du fulvestrant dans un véhicule de ricinoléate, un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique et un alcool acceptable d'un point de vue pharmaceutique, dans laquelle la préparation est adaptée à une administration intramusculaire et atteint une concentration en fulvestrant dans le plasma sanguin significative d'un point de vue thérapeutique pendant au moins 2 semaines.

20 2. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 30% ou moins en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, au moins 1% en poids d'un solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique et miscible dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation qui soit capable, après injection, d'atteindre une concentration en fulvestrant dans le plasma sanguin significative d'un point de vue thérapeutique pendant au moins 2 semaines.

25 3. Préparation pharmaceutique selon la revendication 1 ou 2, dans laquelle la concentration en fulvestrant dans le plasma sanguin atteinte est de 2,5 ng.ml⁻¹ au moins pendant 2 semaines au moins.

30 4. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 30% ou moins en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, au moins 1% en poids d'un solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique et miscible dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation à 45 mg.ml⁻¹ au moins de fulvestrant.

35 5. Préparation pharmaceutique, selon les revendications 1 à 4, qui contient 25% p/v ou moins d'un alcool acceptable d'un point de vue pharmaceutique.

40 6. Préparation pharmaceutique, selon la revendication 5, qui contient 20% p/v ou moins d'un alcool acceptable d'un point de vue pharmaceutique.

45 7. Préparation pharmaceutique, selon la revendication 5, qui contient de 15 à 25% p/v d'un alcool acceptable d'un point de vue pharmaceutique.

8. Préparation pharmaceutique, selon la revendication 5, qui contient de 17 à 23% p/v d'un alcool acceptable d'un point de vue pharmaceutique.

50 9. Préparation pharmaceutique, selon l'une quelconque des revendications 1 à 8, qui contient 60% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.

10. Préparation pharmaceutique, selon la revendication 9, qui contient 50% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.

55 11. Préparation pharmaceutique, selon la revendication 9, qui contient 45% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.

12. Préparation pharmaceutique, selon la revendication 9, qui contient 40% p/v ou moins d'un solvant d'ester non

EP 1 250 138 B1

aqueux acceptable d'un point de vue pharmaceutique.

- 5
13. Préparation pharmaceutique, selon la revendication 9, qui contient 35% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.
14. Préparation pharmaceutique, selon la revendication 9, qui contient 30% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.
- 10
15. Préparation pharmaceutique, selon la revendication 9, qui contient 25% p/v ou moins d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.
16. Préparation pharmaceutique, selon la revendication 9, qui contient de 10 à 25% p/v d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.
- 15
17. Préparation pharmaceutique, selon la revendication 9, qui contient de 12 à 18% p/v d'un solvant d'ester non aqueux acceptable d'un point de vue pharmaceutique.
18. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 15 à 25% en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, 10 à 25% en poids d'un solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique et miscible dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation à 45 mg.ml⁻¹ au moins de fulvestrant.
- 20
19. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 17 à 23% en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, 12 à 18% en poids d'un solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique et miscible dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation à 45 mg.ml⁻¹ au moins de fulvestrant.
- 25
20. Préparation pharmaceutique selon l'une quelconque des revendications 1 à 19, dans laquelle l'alcool acceptable d'un point de vue pharmaceutique est un mélange d'éthanol et d'alcool benzylique.
- 30
21. Préparation pharmaceutique selon l'une quelconque des revendications 1 à 20, dans laquelle le solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique, est choisi parmi : le benzoate de benzyle ; l'oléate d'éthyle ; le myristate d'isopropyle ; le palmitate d'isopropyle ; ou un mélange de n'importe lesquels d'entre eux.
- 35
22. Préparation pharmaceutique selon l'une quelconque des revendications 1 à 21, dans laquelle le solvant d'ester non aqueux, acceptable d'un point de vue pharmaceutique, est le benzoate de benzyle.
- 40
23. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 15 à 25% en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, 10 à 25% en poids de benzoate de benzyle, dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation à 45 mg.ml⁻¹ au moins de fulvestrant.
- 45
24. Préparation pharmaceutique adaptée à une injection intramusculaire comprenant du fulvestrant, 17 à 23% en poids d'un alcool, acceptable d'un point de vue pharmaceutique, par volume de préparation, 12 à 18% en poids de benzoate de benzyle, dans un véhicule de ricinoléate, par volume de préparation et une quantité suffisante d'un véhicule de ricinoléate, de sorte à élaborer une préparation à 45 mg.ml⁻¹ au moins de fulvestrant.
- 50
25. Préparation pharmaceutique selon la revendication 23 ou 24, dans laquelle l'alcool, acceptable d'un point de vue pharmaceutique, est un mélange d'éthanol et d'alcool benzylique.
26. Préparation pharmaceutique selon la revendication 25, dans laquelle l'éthanol et l'alcool benzylique sont présents avec des % en poids environ égaux par volume de préparation.
- 55
27. Préparation pharmaceutique selon l'une quelconque des revendications 1 à 26, dans laquelle le volume total de la préparation est de 6 ml ou moins et la concentration en fulvestrant est de 45 mg.ml⁻¹ au moins.

EP 1 250 138 B1

28. Préparation pharmaceutique selon l'une quelconque des revendications 1 à 27, dans laquelle la quantité totale de fulvestrant dans la préparation est de 250 mg ou plus et le volume total de la préparation est de 6 ml ou moins.

5 **29.** Préparation pharmaceutique selon la revendication 28, dans laquelle la quantité totale de fulvestrant dans la préparation est de 250 mg et le volume total de la préparation est de 5 à 5,25 ml.

10 **30.** Préparation pharmaceutique selon l'une quelconque des revendications 1 à 29, dans laquelle l'alcool, acceptable d'un point de vue pharmaceutique, est un mélange de 10% en poids d'éthanol par volume de préparation, de 10% en poids d'alcool benzylique par volume de préparation, la préparation contient 15% en poids de benzoate de benzyle par volume de préparation et le véhicule de ricinoléate est de l'huile de castor.

31. Préparation pharmaceutique adaptée à une injection intramusculaire, selon l'une quelconque des revendications 1 à 30, à utiliser dans une thérapie médicale.

15 **32.** Utilisation de fulvestrant dans l'élaboration d'une préparation pharmaceutique, selon l'une quelconque des revendications 1 à 30, destinée au traitement d'une maladie bénigne ou maligne du sein ou de l'appareil reproducteur.

33. Seringue ou flacon contenant une préparation pharmaceutique, selon la revendication 30.

20

25

30


35

40

45

50

55



INFORMATION DISCLOSURE CITATION (Use several sheets if necessary)	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	PTO Form 1449 June 4, 2009	Filing Date: October 15, 2008

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date
	1.	2,822,316	February 4, 1958	Richter et al.		
	2.	2,983,649	May 9, 1961	Ercoli et al.		
	3.	3,541,209	November 17, 1970	Neumann et al.		
	4.	RE 28,690	January 20, 1976	Lehmann et al.		
	5.	4,048,309	September 13, 1977	Chen et al.		
	6.	4,048,310	September 13, 1977	Chen et al.		
	7.	4,659,516	April 21, 1987	Bowler et al.		
	8.	4,888,331	December 19, 1989	Elger et al.		
	9.	5,095,129	March 10, 1992	Ottow et al.		
	10.	5,183,814	February 2, 1993	Dukes		
	11.	5,484,801	January 16, 1996	Al-Razzak et al.		
	12.	5,733,902	March 31, 1998	Schneider		
	13.	5,929,030	July 27, 1999	Hamied et al.		
	14.	20010006963	July 5, 2001	Lachnit-Fixson et al.		

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
15.	EP 0138504	Apr., 1985	EP			
16.	EP 0346014	Dec., 1989	EP			
17.	EP 0819431	Mar., 1999	EP			
18.	EP 0905143	Mar., 1999	EP			
19.	FR 6241	Sep., 1968	France			Abstract
20.	GB 817241	Jul., 1959	GB			
21.	GB 1126892	Sep., 1968	GB			
22.	GB 1207571	Oct., 1970	GB			
23.	GB 1569286	Jun., 1980	GB			
24.	JP 43-27327	Nov., 1992	Japan			
25.	JP 09-208496	Dec., 1997	Japan			Abstract
26.	JP 10-203982	Apr., 1998	Japan			
27.	JP 10-152438	Jun., 1998	Japan			Abstract
28.	JP 11-501649	Feb., 1999	Japan			
29.	JP 11-158200	Jun., 1999	Japan			
30.	SU 549118	Mar., 1977	Soviet Union			Abstract
31.	SU 676284	Jul., 1979	Soviet Union			Abstract
32.	WO 95/12383	May., 1995	WIPO			Abstract

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

Examiner	Date Considered
----------	-----------------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009				Attorney Docket No. 056291-5004-02		Application No. 12/285,887			
				Applicants: John R. EVANS et al.					
				Filing Date: October 15, 2008			Group Art Unit: 1617		
U.S. PATENT DOCUMENTS									
Initial		Document No.	Date	Name	Class	Sub-Class	Filing Date		
FOREIGN PATENT DOCUMENTS									
		Document No.	Date	Country	Class	Sub-Class	Translation		
	33.	WO 96/19997	Jul., 1996	WIPO			Abstract		
	34.	WO 97/21440	Jun., 1997	WIPO					
	35.	WO 97/37653	Oct., 1997	WIPO			Abstract		
	36.	WO 97/40823	Nov., 1997	WIPO					
	37.	WO 98/11902	Mar., 1998	WIPO			Abstract		
	38.	WO 99/27906	Jun., 1999	WIPO					
	39.	ZA 681014	Feb., 1968	South Africa					
	40.	ZA 682530	Apr., 1968	South Africa					
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)									
	41.	Anschel, "Lösungsmittel und Lösungsvermittler in Injektionen", Pharm, Ind., 1965, Vol. 27 (11a), pp. 781-787							
	42.	Davis et al., "17-Alpha-Hydroxyprogesterone-Caproate:...with Chemically Pure Progesterone", J. Clin. Endocrinol. And Metabolism, 1955, Vol. 15, pp. 923-930							
	43.	Dukes et al., "Antiuterotrophic effects of pure antioestrogen. ICI 182,780, ...the uterus in ovariectomized monkeys", J. Endocrinology, 1992, Vol. 135, pp. 239-247							
	44.	Dukes et al., "Antiuterotrophic effects of the pure antioestrogen ICI 182, 780 ...quantitative magnetic resonance imaging"; J. Endocrinology, 1992, Vol. 138, pp. 203-209							
	45.	Howell et al., "Pharmacokinetics, pharmacological and anti-tumour effects of the specific anti-oestrogen ICI 182780 in women with advanced breast cancer", British Journal of Cancer, 1996, Vol. 74, pp. 300-308							
	46.	Howell et al., "Response to a specific antioestrogen (ICI 182780) in tamoxifen-resistant breast cancer", The Lancet, Jan. 7, 1995, pp. 29-30							
	47.	Mackey et al, "Tolerability of intramuscular injections of testosterone ester in oil vehicle", Human Reproduction, vol. 10, no. 4, pp. 869-865, 1995							
	48.	Martindale, 32nd Ed., "Alcohol", Pharmaceutical Press, 1999, pp. 1099-1101							
	49.	Martindale, 32nd Ed., "Benzoates" and "Benzyl Alcohol"; Pharmaceutical Press, 1999, pp. 1102-1104							
	50.	Martindale, 32nd Ed., "Caster Oil"; 32nd Ed., Pharmaceutical Press, 1999, p. 1560							
	51.	Migally, "Effect of Castor Oil and Benzyl Benzoate Used as a Vehicle for Antiandrogens on the Adrenal Cortex", Archives of Andrology 2, 1979 pp. 365-369							
	52.	Osborne et al., "Comparison of the Effects of a Pure Steroidal Antiestrogen With Those of Tamoxifen in a Model of Human Breast Cancer", Journal of the National Cancer, May 1995, Vol. 87, No. 10, pp. 746-750							
	53.	Pellegrino, "Use of 17 α Hydroxyprogesterone Caproate in Threatened Abortion", Current Therapeutic Research, Vol. 4, No. 6, June, 1962, pp. 301-305							
	54.	Piver et al., "Medroxyprogesterone Acetate (Depo-Provera) vs. . . . Women with Metastatic Endometrial Adenocarcinoma", Cancer, Vol. 45, American Cancer Society, 1980, pp. 268-272							
	55.	Remington's Pharmaceutical Sciences, 18th ed., 1990, p. 219							
Examiner				Date Considered					
Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.									

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS							
Initial		Document No.	Date	Name	Class	Sub-Class	Filing Date

FOREIGN PATENT DOCUMENTS							
		Document No.	Date	Country	Class	Sub-Class	Translation

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)	
56.	Riffkin et al., "Castor Oil as a Vehicle for Parenteral Administration of Steroid Hormones", Journal of Pharmaceutical Sciences, Vol. 53, No. 8, August 1964, pp. 891-895
57.	Robertson et al., "A partially-blind, randomised, multicentre study comparing the anti-tumor effects of single doses (50, 125 and 250mg) of long-acting (LA) 'faslodex' (ICI 182,780 with tamoxifen in postmenopausal women with primary breast cancer prior to surgery"; Abstract 28, 22nd Annual San Antonio Breast Cancer Symposium: Dec. 8-11, 1999, San Antonio, Breast Cancer Research and Treatment 1999; 57 (1; special issue); p. 31
58.	Sawada et al., "Estrogen Receptor Antagonist ICI182,780 Exacerbates Ischemic Injury in Female Mouse", Journal of Cerebral Blood Flow and Metabolism, Vol. 20. No. 1, 2000, pp. 112-118.
59.	Vidal, Le Dictionnaire, "Benzo-Gynoestryl Retard", 1998 pg. 201
60.	Vidal, Le Dictionnaire, "Gravibinan", 1995, pp 660-661
61.	Vidal, Le Dictionnaire, "Parabolan", 1997, pg. 1245
62.	Vidal, Le Dictionnaire, "Trophobolene", 1997, pp. 1706-1707
63.	Wakeling et al., "A Potent Specific Pure Antiestrogen with Clinical Potential", Cancer Research, 1991, Vol. 51, pp. 3867-3873
64.	Waterton et al., "A Case of Adenomyosis in a Pigtailed Monkey... Treated with the Novel Pure Antiestrogen, ICI 182,780"; Laboratory Animal Science, 1993, Vol. 43, No. 3, 1993, pp. 247-251
65.	"Pharmaceutical dosage forms: Parenteral medications, Volume 1", 2nd edition, edited by Avis, Lieberman and Lachman, 1992
Examiner	Date Considered

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 9/08, 47/14, 47/44, 31/70, 31/365</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/27906 (43) International Publication Date: 10 June 1999 (10.06.99)</p>
<p>(21) International Application Number: PCT/US98/19016 (22) International Filing Date: 14 September 1998 (14.09.98) (30) Priority Data: 60/067,374 3 December 1997 (03.12.97) US 9809792.6 7 May 1998 (07.05.98) GB (71) Applicants (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). Merial LLC [US/US]; 2100 Ronson Road, ISD - 200F, Iselin, NJ 08830-3077 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WILLIAMS, James, B. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CHERN, Rey, T. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>	<p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: LONG ACTING INJECTABLE FORMULATIONS CONTAINING HYDROGENATED CASTOR OIL</p>		
<p>(57) Abstract</p>		
<p>This invention relates to novel, long-acting injectable formulations. These formulations comprise: (a) a therapeutic agent selected from the group consisting of, e.g., insecticides, acaricides, parasiticides, growth enhancers and oil-soluble NASIDS; (b) hydrogenated castor oil and (c) a hydrophobic carrier comprising: (i) triacetin, benzyl benzoate or ethyl oleate or a combination thereof; and (ii) acylated monoglycerides, propyl dicaprylates/dicaprates or caprylic/capric acid triglycerides or a combination thereof. Also provided herein is a method for the treatment or prevention of various disease states by the parental administration of the invention formulations.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

TITLE OF THE INVENTION
LONG ACTING INJECTABLE FORMULATIONS CONTAINING
HYDROGENATED CASTOR OIL

5 CROSS REFERENCE TO RELATED APPLICATIONS

Reference is made to provisional U.S. application Serial No. 60/067,374, filed on December 3, 1997. That application, as well as all documents cited herein and all documents cited in documents cited herein, are hereby incorporated by reference.

10

SUMMARY OF THE INVENTION

This invention is concerned with the unexpectedly long duration of activity which is observed when injectable formulations containing certain therapeutic agents are prepared using hydrogenated
15 castor oil and a combination of hydrophobic or water immiscible carriers. Thus, it is an object of this invention to provide such a prolonged therapeutic effect. An additional object is to describe the therapeutic agents which may be employed in the long acting formulations. A still further object is to provide additional components
20 which may be employed in the formulations. Additional objects will become apparent from a reading of the following description.

BACKGROUND OF THE INVENTION

The therapeutic agents which are used in the inventive
25 formulations are well known to the practitioner to which this invention pertains. Classes of therapeutic agents contemplated by the inventive formulations include insecticides, acaricides, parasiticides, growth enhancers, and oil-soluble, nonsteroidal anti-inflammatory drugs (NSAIDS). Specific classes of compounds which fall within these
30 classes include, for example, avermectins, milbemycins, nodulisporic acid and its derivatives, estrogens, progestins, androgens, substituted pyridylmethyl derivatives, phenylpyrazoles, and COX-2 inhibitors.

The avermectin and milbemycin series of compounds are potent anthelmintic and antiparasitic agents against a wide range of internal and external parasites. The compounds which belong to this series are either natural products or are semi-synthetic derivatives thereof. The structure of these two series of compounds are closely related and they both share a complex 16-membered macrocyclic lactone ring; however, the milbemycins do not contain the disaccharide substituent in the 13-position of the lactone ring. The natural product avermectins are disclosed in U.S. Patent 4,310,519 to Albers-Schonberg, *et al.*, and the 22, 23-dihydro avermectin compounds are disclosed in Chabala, *et al.*, U.S. Patent 4,199,569. For a general discussion of avermectins, which include a discussion of their uses in humans and animals, see "Ivermectin and Abamectin," W.C. Campbell, ed., Springer-Verlag, New York (1989). Naturally occurring milbemycins are described in Aoki *et al.*, U.S. Patent 3,950,360 as well as in the various references cited in "The Merck Index" 12th ed., S. Budavari, Ed., Merck & Co., Inc. Whitehouse Station, New Jersey (1996). Semisynthetic derivatives of these classes of compounds are well known in the art and are described, for example, in U.S. Patent 5,077,308, U.S. Patent 4,859,657, U.S. Patent 4,963,582, U.S. Patent 4,855,317, U.S. Patent 4,871,719, U.S. Patent 4,874,749, U.S. Patent 4,427,663, U.S. Patent 4,310,519, U.S. Patent 4,199,569, U.S. Patent 5,055,596, U.S. Patent 4,973,711, U.S. Patent 4,978,677, and U.S. Patent 4,920,148.

European Patent Application 413,538 relates to an injectable formulation containing an avermectin compound and triacetin. European Patent Application 535,734 relates to an injectable formulation containing an avermectin compound and hydrogenated castor oil in a hydrophobic carrier such as triacetin. The formulations in both European Patent Applications are said to provide efficacy against external and internal parasites in animals only for up to 42 days. Neither of these applications suggests or teaches how to manipulate the composition of the formulation in order to achieve efficacy beyond 42 days.

Nodulisporic acid and its derivatives are a class of acaricidal, antiparasitic, insecticidal and anthelmintic agents known to a practitioner of the art. These compounds are used to treat or prevent infections in humans and animals. These compounds are described, for example, in U.S. Patent 5,399,582 and WO 96/29073. Additionally, the compounds can be administered in combination with other insecticides, parasiticides, and acaricides. Such combinations include anthelmintic agents, such as those discussed above which include ivermectin, avermectin, and emamectin, as well as other agents such as thiabendazole, febantel or morantel; phenylpyrazoles such as fipronil; and insect growth regulators such as lufenuron. Such combinations are also contemplated in the present invention.

Generally, all classes of insecticides are provided for in this invention. One example of this class include substituted pyridylmethyl derivatives such as imidacloprid. Agents of this class are described, for example, in U.S. Patent 4,742,060 or in EP 892,060. It would be well within the skill level of the practitioner to decide which individual compound can be used in the inventive formulation to treat a particular infection of an insect.

Phenylpyrazoles are another class of insecticides which possess excellent insecticidal activity against all insect pests including blood-sucking pests such as ticks, fleas etc., which are parasites on animals. This class of agents kills insects by acting on the gamma-butyric acid receptor of invertebrates. Such agents are described, for example, in U.S. Patent No. 5,567,429, U.S. Patent No. 5,122,530, and EP 295,117. It would be well within the skill level of the practitioner to decide which individual compounds can be used in the inventive formulations.

Insect growth regulators are another class of insecticides or acaricides, which are also provided for in the inventive formulations. Compounds belonging to this group are well known to the practitioner and represent a wide range of different chemical classes. These compounds all act by interfering with the development or growth of the

insect pests. Insect growth regulators are described, for example, in U.S. Patent 3,748,356; U.S. patent 3,818,047; U.S. Patent 4,225,598; U.S. Patent 4,798,837; and U.S. Patent 4,751,225, as well as in EP 179,022 or U.K. 2,140,010. Again, it would be well within the skill level of the
5 practitioner to decide which individual compounds can be used in the inventive formulation.

Estrogens, progestins, and androgens refers to classes of chemical compounds which are also well known to a practitioner in this art. In fact, estrogens and progestins are among the most widely
10 prescribed drugs and are used, for example, alone or in combination for contraception or hormone replacement therapy in post menopausal women. Estrogens and progestins occur naturally or are prepared synthetically. This class of compounds also includes estrogens or progesterone receptor antagonists. Antiestrogens, such as tamoxifen
15 and clomiphene, are used to treat breast cancer and infertility. Antiprogestives are used as contraceptives and anticancer drugs, as well as to induce labor or terminate a pregnancy.

The androgens and antiandrogens structurally related to the estrogens and progestins as they are also biosynthesized from
20 cholesterol. These compounds are based on testosterone. Androgens are used for hypogonadism and promote muscle development. Antiandrogens are used, for example, in the management of hyperplasia and carcinoma of the prostate, acne, and male pattern baldness as well as in the inhibition of the sex drive in men who are sex
25 offenders. Estrogen, progestins, and androgens are described, for example, in "Goodman & Gilman's The Pharmacological Basis of Therapeutics," 9th ed., J.G. Handman and L. Elimbird, eds., Ch. 57 to 60, pp. 1411-1485, McGraw Hill, New York (1996) or in "Principles of Medicinal Chemistry," 2nd ed., W.O. Foye, ed., Ch. 21, pp. 495-559, Lea &
30 Febiger, Philadelphia (1981).

Estrogens, progestins and androgens are also used in animal husbandry as growth promoters for food animals. It is known in the art that compounds of these classes act as growth-promoting steroids

in animals such as cattle, sheep, pigs, fowl, rabbits, etc. Delivery systems to promote the growth of animals are described, for example, in U.S. Patent 5,401,507, U.S. Patent 5,288,469, U.S. Patent 4,758,435, U.S. Patent 4,686,092, U.S. Patent 5,072,716 and U.S. Patent 5,419,910.

5 NSAIDS are well known in the art. The classes of compounds which belong to this group include salicylic acid derivatives, para-aminophenol derivatives, indole and indene acetic acids, heteroaryl acetic acids, arylpropionic acids, anthranilic acids (fenamates), enolic acids, and alkanones. NSAIDS exert their activity by
10 interfering with prostaglandin biosynthesis by irreversibly or reversibly inhibiting cyclooxygenase. Also included are COX-2 inhibitors which act by inhibiting the COX-2 receptor. Compounds of this group possess analgesic, antipyretic and nonsteroidal anti-inflammatory properties. Compounds belonging to these classes are described, for example, in
15 Chapter 27 of Goodman and Gilman on pages 617 to 658 or in Ch. 22 of Foye on pages 561 to 590 as well as in U.S. Patents 3,896,145; U.S. Patent 3,337,570; U.S. Patent 3,904,682; U.S. Patent 4,009,197; U.S. Patent 4,223,299; and U.S. Patent 2,562,830, as well as the specific agents listed in The Merck Index. This invention contemplates those compounds that
20 are oil-soluble.

These and other embodiments are disclosed or are obvious from and encompassed by the following Detailed Description of the Invention.

25 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a long acting injectable formulation for use in human and veterinary medicine, depending on the selected specific therapeutic agent and the indication being treated. The inventive formulation comprises:

- 30 (a) a therapeutic agent selected from the group consisting of , e.g., insecticides, acaricides, parasiticides, growth enhancers, and oil-soluble NSAIDS,
- (b) hydrogenated castor oil, and

- (c) a hydrophobic carrier comprising:
- (i) triacetin, benzyl benzoate, or ethyl oleate or a combination thereof,
 - (ii) acylated monoglycerides, propyl dicaprylates/dicaprates and caprylic acid/capric triglycerides.
- 5 More preferred, are long-acting injectable formulations wherein
- (a) a therapeutic agent selected from the group consisting of, e.g., avermectins, milbemycins, nodulisporic acid and its derivatives, estrogens, progestins, androgens, phenylpyrazoles and COX-2 inhibitors,
 - (b) hydrogenated castor oil, and
 - (c) a hydrophobic carrier comprising:
 - (i) triacetin, benzyl benzoate, or ethyl oleate or a combination thereof,
 - (ii) acetylated monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric acid triglycerides or a combination thereof.
- 10
15

The formulations of the present invention considerably prolong the duration of activity. By the term "acyl" Applicants mean an organic acid group in which the OH of the carboxyl group is replaced by some other substituent; i.e., RCO wherein R is, for example a C₁-C₁₀-alkyl group or a carbocyclic aromatic or a heteroaromatic group. Examples of such groups include acetyl, propionyl, butyryl, isobutyryl, and benzoyl. The term "prolonged duration of activity" means that the activity of the therapeutic agent is extended beyond the time period normally achieved when the therapeutic agent is injected into a host using a conventional, prior art carrier. As conventional injectable formulations are well known in the art, a skilled practitioner could readily understand the meaning of this term. Generally, depending upon the agent, host, and disease state, activity can be prolonged for a period from up to 120 days to up to 180 days. Preferable time periods in which the duration of the agent is prolonged includes from 14 days to 180

20
25
30

days, 30 days to 150 days, 42 days to 120 days, and 60 days to 90 days. While not wishing to be bound by theory, it is believed this increase in activity is achieved because the inventive formulations significantly increase the plasma concentration in tissue for an extended period of
5 time by up to about 2 weeks to about 24 weeks, with time periods of up to about 6, 8, 10, 12, 16 and 20 weeks being observed. With respect to avermectins and milbemycins, the present formulations have been found to have a considerably prolonged duration against internal and external parasites over prior injectable formulation of avermectins or
10 milbemycins. In addition, the present formulations for avermectin and milbemycin provide significantly higher plasma levels at day 42 than prior long-acting formulations thereby producing efficacy for all relevant parasitic species.

Preferred long-acting injectable formulations comprise:
15 (a) about 1.0 to about 10.0% w/v of a therapeutic agent,
(b) about 1 to about 3% w/v of hydrogenated castor oil,
and
(c) a hydrophobic carrier comprising:
(i) about 30 to about 45% v/v of triacetin; benzyl
20 benzoate or ethyl oleate; and
(ii) about 55 to 70% of v/v of acetylated
monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric
triglycerides.

Even more preferred are the above formulations wherein
25 about 1.0 to about 5.0% w/v of a therapeutic agent is present. Especially preferred are the inventive formulations wherein about 2.5 to about 5.0% w/v of a therapeutic agent is present.

Especially preferred long-acting formulation of the present invention comprises:
30 (a) an avermectin or milbemycin compound,
(b) hydrogenated castor oil, and
(c) triacetin and acetylated monoglycerides.

In an especially preferred embodiment, the long-acting formulation comprises:

(a) about 1.0 to about 5.0% w/v of an avermectin or milbemyacin compound,

5 (b) about 0.5 to about 3.5% w/v of hydrogenated castor oil, and

(c) about 30 to about 45% v/v of triacetin and about 55 to about 70% v/v of acetylated monoglycerides.

10 In a most preferred embodiment, the long-acting formulation comprises:

(a) 3.15% w/v of ivermectin,

(b) 1% w/v of hydrogenated castor oil, and

(c) 40% v/v of triacetin and up to 60% v/v of acetylated monoglycerides.


15 Another aspect of the invention is to provide a method for the prevention or treatment of parasite or insect infestations in a host in need thereof for an extended period of time by administering a single dose of a long-acting injectable formulation comprising the appropriate therapeutic agent. That duration typically, for example, lasts from up to
20 about 1 month to about six months, depending upon the agent, host and indication being treated. Extensions of activity lasts from up to about 2 months to about 5 months and especially from up to 3 months to up to 4 months are observed. A further aspect of this invention is to promote the growth in animals by administering a single long-acting formulation
25 according to the present invention wherein the therapeutic agent is an estrogen, progestin, or androgen. Another aspect of the present invention is a method to treat inflammation, pain or fever for an extended period of time in a host in need thereof by administering a single-dose of a formulation according to the present invention wherein
30 the therapeutic agents are oil-soluble NSAIDS. An especially preferred aspect of the invention is to provide a method for the prevention or treatment of parasitic infestation in cattle for a minimum of 42 days which comprises administering to said cattle a single dose of a long-

acting injectable formulation according to the present invention where the therapeutic agent is an avermectin or milbemycin.

Therapeutic agents used in the invention formulations include all known avermectins, milbemycins, nodulisporic acid and its derivatives, estrogens, progestins, androgens, oil-soluble NSAIDS, phenylpyrazoles, substituted pyridylmethyl compounds, and agents which act as insect growth regulators, which are compatible in the inventive formulations for their intended use. The ester and amide derivatives of these compounds, where applicable, as well as their salt forms are also contemplated. Specific compounds which belong to these classes of therapeutic agents are well known to the practitioner of this art. Likewise, the specific disease state as well as the particular dose would be well known to the practitioner.

Avermectins and milbemycins share the same common 16-membered macrocyclic lactone ring; however milbemycins do not possess the disaccharide substituent on the 13-position of the lactone ring.

While many avermectin compounds are known in the art, a representative structure of the class of compounds is as follows:



20


where the broken line indicates a single or a double bond at the 22,23-positions;

R_1 is hydrogen or hydroxy provided that R_1 is present only when the broken line indicates a single bond;

R_2 is alkyl of from 1 to 6 carbon atoms or alkenyl of from 3 to 6 carbon atoms or cycloalkyl of from 3 to 8 carbon atoms;


R_3 is hydroxy, methoxy or = NOR_5 where R_5 is hydrogen or lower alkyl; and

5 R_4 is hydrogen, hydroxy or



where R_6 is hydroxy, amino, mono- or di-lower alkylamino or lower alkanoylamino.


The preferred compounds are avermectin Bla/Blb
 10 (abamectin), 22,23-dihydro avermectin Bla/Blb (ivermectin) and the 4"-acetylamino-5-ketoximino derivative of avermectin Bla/Blb. Both abamectin and ivermectin are approved as broad spectrum antiparasitic agents. The structures of abamectin and ivermectin are as follows:



15 wherein for abamectin the broken line represents a double bond and R_1 is not present and for ivermectin the double bond represents a single bond and R_1 is hydrogen; and

R_2 is isopropyl or sec-butyl.


The 4'-acetylamino-5-ketoximino derivatives of avermectin Bla/B1b has the following structural formula:




5 where R_2 is isopropyl or sec-butyl.

The avermectin products are generally prepared as a mixture of at least 80% of the compound where R_2 is sec-butyl and no more than 20% of the compound where R_2 is isopropyl.


10 Other preferred avermectins, include ememectin, epinomectin and doramectin. Doramectin is disclosed in U.S. Patent 5,089,490 and EP 214738. This compound has the following structure:



In the present formulations, ivermectin is especially preferred.
A representative structure for a milbemycin is that for milbemycin α_1 :



5 An especially preferred milbemycin is moxidectin, whose structure is as follows:



The compound is disclosed in U.S. Patent No. 5,089,490.

Insecticides contemplated by this invention are also well known in the art and such compounds include substituted pyridylmethyl derivatives and phenylpyrazoles. An especially preferred substituted pyridylmethyl derivative is imidacloprid. An especially preferred phenylpyrazole is fipronil, whose chemical name is 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylpyrazole. Fipronil is well known in the art as a flea and tick agent. Additional insecticides included by the invention include insect growth regulators. Especially preferred insect growth regulators include diflubenzuron, lufenuron, methoprene, phoxycarb, pyriproxyfen, and cyromazine.

Specific estrogen, progestin and androgen compounds are well known to the practitioner. Especially preferred compounds belonging to this class include progesterone, estradiol benzoate and trenbolone acetate.

Oil-soluble NSAIDS are also well known to the practitioner. Classes of NSAIDS which are preferred are indole and indene acetic acids and heteroaryl acetic acids. Especially preferred compounds include indomethacin, ketorolac, caprofen, flunixin, ketoprofen, meloxicam, naproxen, and phenylbutazone.

Hydrogenated castor oil is refined, hydrogenated, and deodorized castor oil, consisting mainly of the triglyceride of

hydroxystearic acid. The hydrogenated castor oil is readily prepared using normal techniques known to those skilled in the art of preparing hydrogenated castor oils and one suitable form of hydrogenated castor oil is available commercially under the trade name "Thixcin R" from NL
5 Industries. While not wishing to be bound by theory, it appears that hydrogenated castor oil, being a waxy hydrophobic solid, is left at the injection site entrapping the therapeutic agent after the hydrophobic carrier has diffused from the injection site; it is this hydrophobic hydrogenated castor oil/therapeutic agent matrix that forms a "depot" of
10 the active material which slowly diffuses from the injection site over a prolonged period of time. The hydrogenated castor oil constitutes approximately 1% w/v of the present formulation.

The hydrophobic carrier of the present formulation comprises a mixture of

- 15 (i) triacetin, benzylbenzoate, ethyl oleate or a combination thereof; and
(ii) acylated monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric triglycerides or a combination thereof.

20 These compounds as well as their sources are well known in the art. For example, triacetin (glyceryl triacetate or glycerol triacetate) and acetylated monoglycerides (available under the tradename "Myvacet 9-45" from Quest International). The ratio of component (i) to component (ii) used in the present formulation is
25 generally from 45:55 to 30:70; preferably the ratio is approximately 40:60. In addition to the hydrogenated castor oil, the therapeutic agent and the hydrophobic carrier, the formulation can contain other inert ingredients such as antioxidants or preservatives. Antioxidant such as a propyl
30 gallate, BHA (butylated hydroxy anisole), BHT (butylated hydroxy toluene) monothioglycerol and the like may be added to the present formulation. The antioxidants are generally added to the formulation in amounts of from about 0.01 to about 2.0% (w/v). Preservatives such as

the parabens (methylparaben and/or propylparaben) are suitably used in the formulation in amounts ranging from about 0.01 to about 2.0 w/v.

The long-acting injectable formulation of the present invention may be prepared by adding a dispersion of hydrogenated castor oil in acetylated monoglycerides, propyl dicaprylates/dicaprates or caprylic/capric triglycerides to a solution comprising the therapeutic agent, and any other inert ingredients, in triacetin benzyl benzoate or ethyl oleate, and mixing the liquids until uniform. Since the long acting formulation is intended for injection, it is necessary that it be sterilized. Heat sterilization is generally to be avoided in the situation where avermectin or milbemycin compounds are used since these compounds are unstable at autoclave temperatures. Rather, membrane sterilization is preferred in those situations with dissolved solids and gamma sterilization for the hydrogenated castor oil. The sterile hydrogenated castor oil is dispersed in the product aseptically and then aseptically packaged.

The instant formulation is equally applicable to other compounds used for injection as long as such compounds are soluble in the mixture of the hydrogenated castor oil and hydrophobic carrier. Additional compounds that can be used in this formulation are other antiparasitic agents and antibiotics, therapeutic vitamin and mineral supplements, and other agents that are assisted in their therapeutic effect by having their effects extended over a prolonged period of time. Again, such compounds would be well known to the practitioner.

The instant long-acting formulations are administered to a warm-blooded animals such as humans, cattle, sheep, pigs, cats, dogs, horses, and the like by intramuscular or subcutaneous injection. The amount of therapeutic agent depends on the individual therapeutic agent, the animal being treated, the disease state, and the severity of the disease state. The determination of those factors is well within the skill level of the practitioner. Generally, such preparation normally contain about 0.0005 to about 50% w/v of therapeutic agent. Preferred formulations are those containing about 0.01 to 10% w/v of therapeutic

agent and especially preferred formulations are those containing about 2.5 to about 5% w/v of therapeutic agent. For the avermectins and milbemycins, the formulations will generally be prepared to administer from about 0.1 to about 2 mg/kg, preferably from about 0.4 to about 0.85 mg/kg and most preferably from about 0.6 to about 0.7 mg/kg of the active ingredient. At a preferred dose volume of about 1 ml to treat 50 kg of animal body weight the formulation contains from about 5 to about 50 mg of the active agent per ml of solution or about 0.5 to about 10%, w/v preferably about 2.5 to about 5% w/v. However, depending upon the activity of the compound and the animal being treated, doses as low as about 0.3% w/v of the active ingredient are usable. For nodulisporic acid and its derivatives, a formulation containing about 0.0005 to about 5% w/v of the active compound is preferred.

The present formulation provides for an extended period of treatment. For avermectins and milbemycins a minimum of 42 days of activity against endo- and ectoparasites is obtained without causing tissue irritation. The extended period of time for the other therapeutic agents is readily determined by one skilled in the art and is determined by such factors as the therapeutic agent, disease state, host and severity of the infection. While the previously reported avermectin formulation containing hydrogenated castor oil in triacetin did produce prolonged plasma level compared to a formulation without hydrogenated castor oil, it did not achieve a plasma level efficacious against all relevant parasitic species at the 42 day target. In contrast the present formulation using avermectins or milbemycins surprisingly provides a significantly higher plasma at day 42 and beyond. The present formulation is also efficacious against ticks and *Dermatobia hominis* for up to 75 and 140 days, respectively.

The following example is provided in order that the invention might be more fully understood. It is not to be construed as a limitation of the invention.

EXAMPLE 1

<u>Material</u>	<u>%</u>	<u>Amount</u>
Ivermectin	3.15% w/w	17.6 gm
n-propyl gallate	0.02% w/w	0.10 gm
Thixcin R	1.0% w/w	5.0 gm
triacetin	40.0% w/w	200.0 gm
Myvacet 9-45	qs 100% w/w	qs to 500.0 gm

5 Triacetin was added to n-propyl gallate and ivermectin in an Erlenmyer flask and mixed until all of the n-propyl gallate dissolved. Myvacet 9-45 was placed in a non-glass beaker in a 50°C water bath, and mixed at a low speed with a dispersator mixer until the temperature of the content reached 50°C. Thixcin R was then added slowly to the vortex of the mixing Myvacet 9-45. When all the Thixcin R was added, the speed of the mixer was slowly increased to 60 rpm and mixing continued for 20 minutes. The beaker was removed from the water bath and allowed to cool to 30°C, while mixing continued at about 25 rpm. The triacetin solution was added to the Thixcin R/Myvacet 9-45 mixture and the liquids were mixed until uniform.

15

EXAMPLE 2

<u>Material</u>	<u>%</u>	<u>Amount/2000 L.</u>
Ivermectin	3.15% w/w	63.0 kg
triacetin	40.0% v/v	800.0 L
hydrogenated castor oil	1.0% w/w	20.0 kg
BHT	0.02 w/v	0.4 kg
methylparaben	0.18% w/v	3.6 kg
propylparaben	0.02% w/v	0.4 kg
Myvacet 9-45	qs 100% v/v	qs to 1200.0 L

20 Ivermectin, BHT, methyl and propyl paraben were dissolved in 800 L of triacetin, and the solution was sterile filtered into a 2000 L tank equipped with an agitator. Myvacet 9-45 was sterile filtered into a 150 L tank capable of maintaining a batch temperature of 60°C and

equipped with an agitator and with an aseptic addition of sterile powder capability. The gamma sterilized hydrogenated castor oil was dispersed in the Myvacet 9-45, and the dispersion was heated to 50°C, then transferred to the triacetin solution through a microfluidizer. The liquids were mixed until uniform and then aseptically packaged in low density polyethylene containers.

EXAMPLE 3

The plasma levels of ivermectin administered once subcutaneously at a dose of 630 mcg/kg bodyweight were determined in cattle for two formulations: formulation I contains ivermectin 3.15%, n-propyl gallate 0.02%, Thixcin R 1.5% and triacetin qs to 100%; formulation II has the composition given in Example 2. Ten animals were used for formulation I and six were used for formulation II. Mean plasma levels (ng/ml) are shown in the following Table:

<u>Formulation</u>	<u>Days post dosing</u>					
	<u>3</u>	<u>14</u>	<u>21</u>	<u>28</u>	<u>35</u>	<u>42</u>
I	80	18	10	6	4	2
II	21	25	22	16	13	9

The mean plasma level for formulation II was greater than 3 ng/ml on day 70.

The 42-day plasma level of formulation I (2 ng/ml) is not sufficient to produce efficacy against *Cooperia onocophora* and *Nematodirus* which require an ivermectin plasma level of 3 to 4 ng/ml.

EXAMPLE 4

To facilitate the manufacture of large scale batches the following process was developed which results in a product that meets

the same release specifications as the product manufactured in Example 2. The formula is also the same as used in Example 2. Ivermectin, BHT, methyl and propyl paraben are dissolved a mixture of the triacetin and Myvacet 9-45. The solution is sterile filtered. The
5 gamma sterilized hydrogenated castor is aseptically dispersed in sterile solution using an in-line educator/homogenizer system. Such in-line system can be a Flashblend system. The product is heated and recirculated through the system until the product temperature is from 42 to 50°C. Then the product is aseptically packaged.

10

* * *

15

Having thus described in detail preferred embodiments of the present invention, it is to be understood that the invention defined by the appended claims is not to be limited by particular details set forth in the above description as many apparent variations thereof are possible without departing from the spirit or scope thereof.

WHAT IS CLAIMED IS:

1. A long-acting injectable formulation comprising:
 - (a) a therapeutic agent selected from the group
5 consisting of insecticides, acaricides, parasiticides, growth enhancers
and oil-soluble NASIDS,
 - (b) hydrogenated castor oil, and
 - (c) a hydrophobic carrier comprising:
 - (i) triacetin, benzyl benzoate or ethyl oleate or a
10 combination thereof; and
 - (ii) acylated monoglycerides, propyl
dicaprylates/dicaprates, caprylic/capric acid triglycerides, or a
combination thereof.
- 15 2. A long-acting injectable formulation comprising:
 - (a) a therapeutic agent selected from the group
consisting of avermectins, milbemycons, nodulisporic acid and its
derivatives, estrogens, progestins, androgens, substituted pyridylmethyl
derivatives, phenylpyrazoles, and COX-2 inhibitors,
20 (b) hydrogenated castor oil, and
 - (c) a hydrophobic carrier comprising:
 - (i) triacetin, benzyl benzoate or ethyl oleate or a
combination thereof; and
 - (ii) acetylated monoglycerides, propyl
25 dicaprylates/dicaprates, caprylic/capric triglycerides, or a combination
thereof.
- 30 3. The long-acting injectable formulation according to
claim 1 comprising
 - (a) about 1.0 to about 10.0 w/v of a therapeutic agent;
 - (b) about 0.3 to about 5% w/v of hydrogenated castor oil;
 - (c) a hydrophobic carrier comprising:

- (i) about 30 to about 45% v/v of triacetin;
benzylbenzoate or ethyloleate; and
- (ii) about 55 to 70% of v/v of acetylated
monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric
5 triglycerides.
4. The long-acting injectable formulation according to
claim 2 comprising
- (a) about 1.0 to about 10.0 w/v of a therapeutic agent;
- 10 (b) about 0.3 to about 5% w/v of hydrogenated castor oil;
- (c) a hydrophobic carrier comprising:
- (i) about 30 to about 45% v/v of triacetin; benzyl
benzoate or ethyl oleate; and
- (ii) about 55 to 70% of v/v of acetylated
15 monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric
triglycerides.
5. The long-acting injectable formulation according to
claim 2 wherein about 2.5 to about 5.0% w/v of a therapeutic agent is
20 present.
6. The long-acting injectable formulation according to
claim 2 wherein the therapeutic agent is an avermectin or a
milbemycin.
25
7. The long-acting injectable formulation according to
claim 6, wherein the avermectin is ivermectin, abamectin, ememectin,
eprinomectin, or doramectin and the milbemycin is moxidectin.
- 30 8. The long-acting injectable formulation according to
claim 2, wherein the therapeutic agent is an estrogen, progestin or
androgen.

9. The long-acting injectable formulation according to claim 8, where the estrogen, progestin or androgen is estradiol benzoate, progesterone, or trenbolone acetate.

5 10. The long-acting injectable formulation according to claim 2, wherein the therapeutic agent is nodulisporic acid or its derivatives.

10 11. The long-acting injectable formulation according to claim 2, wherein the therapeutic agent is a substituted pyridylmethyl derivative or a phenylpyrazole.

15 12. The long-acting injectable formulation according to claim 11, wherein the therapeutic agent is imidacloprid or fipronil.

13. The long-acting injectable formulation according to claim 2, wherein the therapeutic agent is a COX-2 inhibitor.

20 14. The long-acting injectable formulation according to claim 1, wherein the therapeutic agent is an oil-soluble, nonsteroidal anti-inflammatory drug.

25 15. The long-acting injectable formulation according to claim 14, wherein the therapeutic agent is carprofen, flunixin, ketoprofen, meloxicam, naproxen or phenylbutazone.

16. The long-acting injectable formulation according to claim 1, wherein the therapeutic agent is an insect growth regulator.

30 17. The long-acting injectable formulation according to claim 16, wherein the therapeutic agent is diflubenzuron, lufenuron, methoprene, phenoxy carb, pyriproxyfen, and cyromazine.

18. The long-acting injectable formulation according to claim 1, which further comprises an antioxidant or a preservative.

5 19. The long-acting injectable formulation according to claim 2 where about 1 to about 3.0 % w/v of hydrogenated castor oil is present and hydrophobic carrier comprises about 40% v/v of triacetin, benzylbenzoate or ethyloleate and about 60% v/v of acetylated monoglycerides, propyl dicaprylates/dicaprates, or caprylic/capric triglycerides.

10

20. The long-acting injectable formulation of claim 2 which comprises:

- 15 (a) about 1.0 to about 5.0% w/v of an avermectin compound,
(b) about 1 to about 3% w/v of hydrogenated castor oil, and
(c) about 30 to about 45% v/v of triacetin and 55 to 70% v/v of acetylated monoglycerides.

20 21. The long-acting injectable formulation of claim 2 which comprises:

- 25 (a) about 3.15% w/v of ivermectin,
(b) about 1% w/v of hydrogenated castor oil, and
(c) about 40% of triacetin and up to about 60% v/v of acetylated monoglycerides.

22. The long-acting injectable formulation of claim 2 which further comprises an antioxidant.

30 23. The long acting injectable formulation of claim 2 which further comprises a preservative.

24. The long acting injectable formulation of claim 22 wherein said antioxidant is selected from n-propyl gallate, BHA, BHT and monothioglycerol.

5 25. The long-acting injectable formulation of claim 23 wherein said preservative is selected from the parabens.

26. The long-acting injectable formulation of claim 21 which further comprises an antioxidant selected from n-propyl gallate,
10 BHA, BHT and monothioglycerol.

27. The long-acting injectable formulation of claim 26 which further comprises a preservatives selected from the parabens.

15 28. The long acting injectable formulation of claim 21 which further comprises BHT and one or more preservatives from the parabens.

29. A method for the prevention or treatment of parasitic
20 infestation in a host in need thereof, which comprises parentally administering a single dose of a long-acting injectable formulation of claim 6 to said host.

30. A method for the prevention or treatment of parasitic
25 infestation in a host in need thereof for a minimum of 42 days which comprises administering to said host a single dose of a long-acting injectable formulation of claim 1.

31. A method for the prevention or treatment of parasitic
30 infestation in cattle for a minimum of 42 days which comprises administering to said cattle a single dose of a long-acting injectable formulation of claim 2.

32. A method for the prevention or treatment of parasitic infestation in cattle for a minimum of 42 days which comprises administering to said cattle a single dose of a long-acting injectable formulation of claim 21.

5

33. A method for treating or preventing insect infestation for an extended period of time in a host in need thereof which comprises parentally administering a single-dose of a long-acting injectable formulation according to claim 10 to said host.

10

34. The method according to claim 33, wherein the insects are fleas.

35. A method for treating or preventing insect infestation for an extended period of time in a host in need thereof which comprises parenterally administering a single-dose of a long-acting injectable formulation according to claim 11 to said host.

15

36. The method according to claim 35 wherein the injectable formulation as the therapeutic agent is imidacloprid or fipronil and the insects are fleas.

20

37. A method for promoting growth in animals which comprises administering a single dose of a long-acting injectable formulation according to claim 8 to said animal.

25

38. A method for treating inflammation, pain, or fever for an extended period of time in a host in need thereof which comprises administering a single-dose of a long-acting injectable formulation according to claim 14 to a host in need thereof.

30

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/19016

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/08 A61K47/14 A61K47/44 A61K31/70 A61K31/365

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 11709 A (ASHMONT HOLDINGS LTD ; HARVEY COLIN MANSON (NZ)) 3 April 1997 see page 4 - page 5; examples 1-6	1-38
A	EP 0 413 538 A (MERCK & CO INC) 20 February 1991 cited in the application see page 4 - page 5; example 1	1-38
A	GB 1 060 632 A (OLIN MATHISON CHEMICAL CORP.) 8 March 1967 see page 3; example 2	1-38
A	EP 0 535 734 A (MERCK & CO INC) 7 April 1993 cited in the application see page 4 - page 5; example 1	1-38

-/--

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

3 February 1999

Date of mailing of the international search report

11/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Boulois, D

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/19016

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 29-38 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the international Application that do not comply with the prescribed requirements to such an extent that no meaningful international Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/19016

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 196 13 972 A (BAYER AG) 16 October 1997 see page 6 - page 10; examples 1-10 -----	1-38
A	US 4 330 538 A (ITIL TURAN M ET AL) 18 May 1982 see column 3 - column 4; examples 2,4A -----	1-38
A	DE 25 48 413 A (SCHERING AG) 28 April 1977 see claim 3 -----	1-38

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Examiner: Unassigned
)	
FOR: FORMULATION)	Date: June 4, 2009

SECOND INFORMATION DISCLOSURE STATEMENT

UNDER 37 C.F.R. § 1.97(b)

Applicants wish to specifically call to the Examiner's attention in this continuing application the same circumstances as detailed in the Second Information Disclosure Statements filed September 13, 2002 and October 18, 2004 in parent applications, a copy of this is attached hereto for convenience. Specifically, the attached Second Information Disclosure Statement details circumstances regarding the controlled, confidential and non-commercial testing of compositions falling within the scope of the definition of "pharmaceutical formulation", as used in the present method of treatment claims, which was carried out in the United States more than one year before the filing date of the parent application in preparation for and during the testing (IND) phase of the regulatory review of such formulation by the FDA.

The present Information Disclosure Statement is being filed before the mailing date of a first Office Action, and therefore no certification under 37 CFR §1.97(e) or fee under 37 CFR §1.17(p) is required.

This Information Disclosure Statement is intended to be in full compliance with the rules, but should the Examiner find any part of its required content to have been omitted, prompt notice to that effect is earnestly solicited, along with additional time under Rule 97(f), to enable Applicant to fully comply.

Consideration by the Examiner of the circumstances detailed in the attached document is respectfully requested when taking up this continuing application for a first Action on the merits.

Except for issue fees payable under 37 C.F.R. §1.18, the Commissioner is hereby

authorized by this paper to charge any additional fees during the entire pendency of this application including fees due under 37 C.F.R. §§1.16 and 1.17 which may be required, including any required extension of time fees, or credit any overpayment to Deposit Account No. 50-0310. This paragraph is intended to be a **CONSTRUCTIVE PETITION FOR EXTENSION OF TIME** in accordance with 37 C.F.R. §1.136(a)(3).

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: **June 4, 2009**
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000

By: /Donald Bird/
Donald J. Bird
Registration No. 25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001

FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a Continuation Application of copending U.S. Patent Application No. 10/872,784, filed June 22, 2004, which claims benefit of U.S. Patent Application No. 09/756,291, filed January 9, 2001 which claims the benefit of Great Britain Application No. 0008837.7 filed April 12, 2000 and Great Britain Application No. 0000313.7, filed January 10, 2000, all of which are incorporated herein by reference in their entireties.

BACKGROUND OF THE INVENTION

Field of the Invention

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound

~~7 α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.~~

Description of the Related Art

Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any

of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be “pure” antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for
5 the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7 α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989). One of these, 7 α -[9-(4,4,5,5,5-pentafluoropentyl sulphanyl)nonyl]oestra-1,3,5-(10)triene-
10 3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-
15 dependent indications such as breast cancer and certain benign gynaecological conditions.

7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphanyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

20 Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

25 Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

30 In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone

density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

5 European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-
1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also
10 disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

Fulvestrant shows, along with other steroidal based compounds, certain physical
15 properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml⁻¹ (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

Currently there are a number of sustained release injectable steroidal formulations
20 which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

In the formulations within Table 1 a number of different oils are used to solubilise the compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have
25 been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

<u>PRODUCT NAME</u>	<u>STEROID</u>	<u>DOSE</u>	<u>TYPE</u>	<u>COMP'</u>	<u>SOURCE</u>	<u>OIL</u>	<u>BzBz</u>	<u>BzOH</u>	<u>EtOH</u>	<u>DOSE</u>	<u>DOSING</u>
SUSTANON 100	Testosterone propionate Testosterone phenylpropionate Testosterone isocaproate	30mg 60mg 60mg 100mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%			1 or 2ml	1 week
TOCOGESTAN	Hydroxy progesterone enantate Progesterone α -Tocopherol	200mg 50mg 250mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%			2ml	< 1week
TROPHOBOLINE	Estrapronicate Nandrolone undecanoate Hydroxyprogesterone heptanoate	1.3mg 50mg 80mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%			1ml	15 to 30 days
NORISTERAT	Norethisterone oentanhoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES			1ml	8 weeks
BENZO- GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis				1ml	1 week
PROGESTERONE -RETARD	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES			1 or 2ml	1 week
GRAVIBINAN	Estradiol 17- β -valerate Hydroxyprogesterone caproate	5mgml ⁻¹ 250mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES			1 or 2ml	1 - 2 weeks

Z70635

- 5 -

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis	75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%	
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%	

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
5 % are w/v and * approximate as measured directly from a single sample

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7 α -[9-(4,4,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7 α -[9-(4,4,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

DETAILED DESCRIPTION OF THE INVENTION

Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06

Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

Currently guidelines recommend that no more than 5mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of $>50\text{mgml}^{-1}$ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of $<5\text{ml}$ - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50mgml^{-1} - see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly

lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

Therefore, we present as a feature of the invention a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 5 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Another feature of the invention is a pharmaceutical formulation comprising 10 fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically- 15 acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

20 Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient 25 amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 30 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Preferred pharmaceutical formulations of the invention are as described above

5 wherein:

1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 10 3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably
 15 up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v, at
 20 least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and
 25 preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v, at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50% w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 5-60% w/v, 7-55% w/v, 8-50% w/v, 10-50% w/v, 10-45% w/v, 10-40% w/v, 10-35% w/v, 10-30% w/v, 10-

25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

10 By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after
15 intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles.
5 The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.
2. Our findings from studies using ^{14}C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of
10 administration.

It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra
15 Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended
20 release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

By use of the term "therapeutically significant levels" we mean that blood plasma concentrations of at least 2.5 ngml^{-1} , ideally at least 3 ngml^{-1} , at least 8.5 ngml^{-1} , and up to 12 ngml^{-1} of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less
25 than 15 ngml^{-1} .

By use of the term "extended release" we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks)
30 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a
5 good release profile or lack of precipitation of drug after injection at the injection site.

Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor
10 oil.

Table 3**Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C**

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100.	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

5 **Solubility comparisons of fulvestrant in oil based formulations with and without alcohols**

		Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
10	Formulation ^(a)	Complete vehicle	Vehicle minus alcohols
	Castor oil based	81.2	12.6
15	Miglyol 812-N based	86.8	1.7
	Sesame seed/Castor oil (1:1) based	70.1	4.4
	Sesame seed oil based	45.7	0.7
20	Arachis oil based	40.2	< 0.2

25 ^(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

		Days						
30	Formulation ^a	2	3	4	7	10	30	51
35	Formulation F1 castor oil based	0	0	0	0	0	0	0
	Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
40	Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

45 ^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile over
5 five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

10 Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per
15 volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

20 A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-
25 acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

Preferably 5ml of the intramuscular injection is administered.

A further feature of the invention is use of fulvestrant in the preparation of a
30 pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent indications such as breast cancer and gynaecological conditions, such as endometriosis.

In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary

pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram below

5

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol


10 10% weight per volume of ethanol

15% weight per volume of benzyl benzoate

250mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil

FLOW DIAGRAM OF MANUFACTURING



References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids* 1989; 54:71-99.
- 5 2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- 10 4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- 5 5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112: 15 R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

Claims

1. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of
5 formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration of at least 2.5ngml^{-1} for at least 2 weeks.
- 10 2. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for at least 4 weeks.
3. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for 2 to 5 weeks.
- 15 4. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml^{-1} of fulvestrant.
20
5. A pharmaceutical formulation as claimed in claim 1 to 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.
6. A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of
25 a pharmaceutically-acceptable alcohol.
7. A pharmaceutical formulation as claimed in any claim from 1 to 6 which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 30 8. A pharmaceutical formulation as claimed in claim 7 which contains 50%w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent .

9. A pharmaceutical formulation as claimed in claim 7 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 5 10. A pharmaceutical formulation as claimed in claim 7 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
11. A pharmaceutical formulation as claimed in claim 7 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
- 10 12. A pharmaceutical formulation as claimed in claim 7 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
13. A pharmaceutical formulation as claimed in claim 7 which contains 25% w/v or less
15 of a pharmaceutically-acceptable non-aqueous ester solvent.
14. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- 20 15. A pharmaceutical formulation as claimed in any claim from 1 to 14 wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
16. A pharmaceutical formulation as claimed in any claim from 1 to 15 wherein the
25 pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
17. A pharmaceutical formulation as claimed in any claim from 1 to 16 wherein the total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.

30

18. A pharmaceutical formulation as claimed in any claim from 1 to 13 wherein the total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 5 19. A pharmaceutical formulation as claimed in claim 18 wherein the total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5 to 5.25ml.
20. A pharmaceutical formulation as claimed in any of claims 1-19 wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of
10 formulation, 10% weight of benzyl alcohol per volume of formulation and 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.
21. A method of treating a benign or malignant diseases of the breast or reproductive tract by administration to a human in need of such treatment by intramuscular a pharmaceutical
15 formulation as claimed in claims 1 to 19.
22. A method as claimed in claim 21 for treating breast cancer.
23. A syringe or vial containing a pharmaceutical formulation as defined in claim 20.

20

ABSTRACT**TITLE: Formulation****ABSTRACT OF THE DISCLOSURE**

5 The invention relates to a novel sustained release pharmaceutical formulation adapted
for administration by injection containing the compound
7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more
particularly to a formulation adapted for administration by injection containing the compound
7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in
10 solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-
aqueous ester solvent which is miscible in the ricinoleate vehicle.




Figure 1

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875	Application or Docket Number 12/285,887	Filing Date 10/15/2008	<input type="checkbox"/> To be Mailed
---	---	----------------------------------	---------------------------------------

APPLICATION AS FILED – PART I			OTHER THAN SMALL ENTITY				
	(Column 1)	(Column 2)	SMALL ENTITY <input type="checkbox"/>	OR			
FOR	NUMBER FILED	NUMBER EXTRA	RATE (\$)	FEE (\$)	OR	RATE (\$)	FEE (\$)
<input type="checkbox"/> BASIC FEE <small>(37 CFR 1.16(a), (b), or (c))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> SEARCH FEE <small>(37 CFR 1.16(k), (l), or (m))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> EXAMINATION FEE <small>(37 CFR 1.16(o), (p), or (q))</small>	N/A	N/A	N/A			N/A	
TOTAL CLAIMS <small>(37 CFR 1.16(i))</small>	minus 20 =	*	X \$ =		OR	X \$ =	
INDEPENDENT CLAIMS <small>(37 CFR 1.16(h))</small>	minus 3 =	*	X \$ =			X \$ =	
<input type="checkbox"/> APPLICATION SIZE FEE <small>(37 CFR 1.16(s))</small>	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).						
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENT <small>(37 CFR 1.16(j))</small>							
* If the difference in column 1 is less than zero, enter "0" in column 2.			TOTAL			TOTAL	

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY				
	(Column 1)	(Column 2)	(Column 3)		SMALL ENTITY	OR			
AMENDMENT	10/15/2008	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	OR	RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	* 23	Minus	** 23	= 0		OR	X \$52=	0
	Independent <small>(37 CFR 1.16(h))</small>	* 2	Minus	***3	= 0		OR	X \$220=	0
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>						OR		
					TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	0

	(Column 1)	(Column 2)	(Column 3)		SMALL ENTITY	OR			
AMENDMENT	06/04/2009	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	OR	RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	* 23	Minus	** 23	= 0		OR	X \$52 =	0
	Independent <small>(37 CFR 1.16(h))</small>	* 2	Minus	*** 3	= 0		OR	X \$220 =	0
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>						OR		
					TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	0

* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.
 ** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".
 *** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".
 The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.

Legal Instrument Examiner:
 /HELENA PAYTON/

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**
 If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Electronic Acknowledgement Receipt

Adjustment date: 06/10/2009 MTEKLEMI
 06/05/2009 INTEFSW 00000215 500310 12285887
 05 FC:1101 440.00 CR

EFS ID:	5447464
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	09629
Filer:	Donald J. Bird
Filer Authorized By:	
Attorney Docket Number:	056291-5004-02
Receipt Date:	04-JUN-2009
Filing Date:	15-OCT-2008
Time Stamp:	14:01:06
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Deposit Account
Payment was successfully received in RAM	\$5544
RAM confirmation Number	215
Deposit Account	500310
Authorized User	

The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:

Charge any Additional Fees required under 37 C.F.R. Section 1.16 (National application filing, search, and examination fees)

Charge any Additional Fees required under 37 C.F.R. Section 1.17 (Patent application and reexamination processing fees)



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 7 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY. DOCKET NO, TOT CLAIMS, IND CLAIMS. Row 1: 12/285,887, 10/15/2008, 1617, 2754, 056291-5004-02, 23, 2

CONFIRMATION NO. 1199

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004

UPDATED FILING RECEIPT



Date Mailed: 06/17/2009

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Applicant(s)

John R. Evans, Macclesfield, UNITED KINGDOM;
Rosalind U. Grundy, Macclesfield, UNITED KINGDOM;

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 10/872,784 06/22/2004 PAT 7,456,160

Foreign Applications

UNITED KINGDOM 0008837.7 04/12/2000
UNITED KINGDOM 0000313.7 01/10/2000

If Required, Foreign Filing License Granted: 11/03/2008

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is US 12/285,887

Projected Publication Date: 09/24/2009

Non-Publication Request: No

Early Publication Request: No

Title

Formulation

Preliminary Class

514

PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4158).

LICENSE FOR FOREIGN FILING UNDER**Title 35, United States Code, Section 184****Title 37, Code of Federal Regulations, 5.11 & 5.15****GRANTED**

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as

set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

NOT GRANTED

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NUMBER	FILING OR 371(C) DATE	FIRST NAMED APPLICANT	ATTY. DOCKET NO./TITLE
12/285,887	10/15/2008	John R. Evans	056291-5004-02

CONFIRMATION NO. 1199

WITHDRAWAL NOTICE

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 08/03/2009

Letter Regarding a New Notice and/or the Status of the Application

If a new notice or Filing Receipt is enclosed, applicant may disregard the previous notice mailed on 11/04/2008. The time period for reply runs from the mail date of the new notice. Within the time period for reply, applicant is required to file a reply in compliance with the requirements set forth in the new notice to avoid abandonment of the application.

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web. <https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at **1-866-217-9197** or visit our website at <http://www.uspto.gov/ebc>.

If the reply is not filed electronically via EFS-Web, the reply must be accompanied by a copy of the new notice.

If the Office previously granted a petition to withdraw the holding of abandonment or a petition to revive under 37 CFR 1.137, the status of the application has been returned to pending status.

/tnguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NUMBER	FILING OR 371(C) DATE	FIRST NAMED APPLICANT	ATTY. DOCKET NO./TITLE
12/285,887	10/15/2008	John R. Evans	056291-5004-02

CONFIRMATION NO. 1199

WITHDRAWAL NOTICE

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 08/03/2009

Letter Regarding a New Notice and/or the Status of the Application

If a new notice or Filing Receipt is enclosed, applicant may disregard the previous notice mailed on 06/17/2009. The time period for reply runs from the mail date of the new notice. Within the time period for reply, applicant is required to file a reply in compliance with the requirements set forth in the new notice to avoid abandonment of the application.

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web. <https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at **1-866-217-9197** or visit our website at <http://www.uspto.gov/ebc>.

If the reply is not filed electronically via EFS-Web, the reply must be accompanied by a copy of the new notice.

If the Office previously granted a petition to withdraw the holding of abandonment or a petition to revive under 37 CFR 1.137, the status of the application has been returned to pending status.

/tnguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 7 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY. DOCKET NO, TOT CLAIMS, IND CLAIMS. Row 1: 12/285,887, 10/15/2008, 1617, 2754, 056291-5004-02, 23, 2

CONFIRMATION NO. 1199

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004

FILING RECEIPT



Date Mailed: 08/03/2009

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Applicant(s)

John R. Evans, Macclesfield, UNITED KINGDOM;
Rosalind U. Grundy, Macclesfield, UNITED KINGDOM;

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 10/872,784 06/22/2004 PAT 7,456,160

Foreign Applications

UNITED KINGDOM 0008837.7 04/12/2000
UNITED KINGDOM 0000313.7 01/10/2000

If Required, Foreign Filing License Granted: 11/03/2008

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is US 12/285,887

Projected Publication Date: None.

Non-Publication Request: No

Early Publication Request: No

Title

Formulation

Preliminary Class

514

PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4158).

LICENSE FOR FOREIGN FILING UNDER

Title 35, United States Code, Section 184

Title 37, Code of Federal Regulations, 5.11 & 5.15

GRANTED

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as

set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

NOT GRANTED

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (12/285,887), FILING OR 371(C) DATE (10/15/2008), FIRST NAMED APPLICANT (John R. Evans), ATTY. DOCKET NO./TITLE (056291-5004-02)

CONFIRMATION NO. 1199

FORMALITIES LETTER

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 08/03/2009

NOTICE TO FILE CORRECTED APPLICATION PAPERS

Filing Date Granted

An application number and filing date have been accorded to this application. The application is informal since it does not comply with the regulations for the reason(s) indicated below. Applicant is given TWO MONTHS from the date of this Notice within which to correct the informalities indicated below. Extensions of time may be obtained by filing a petition accompanied by the extension fee under the provisions of 37 CFR 1.136(a).

The required item(s) identified below must be timely submitted to avoid abandonment:

- A substitute specification in compliance with 37 CFR 1.52, 1.121(b)(3), and 1.125, is required. The substitute specification must be submitted with markings and be accompanied by a clean version (without markings) as set forth in 37 CFR 1.125(c) and a statement that the substitute specification contains no new matter (see 37 CFR 1.125(b)). The specification, claims, and/or abstract page(s) submitted is not acceptable and cannot be scanned or properly stored because:
- The specification contains drawings or flow diagrams (37 CFR 1.58(a)) on page(s) 19. Drawings or flow diagrams cannot be embedded in the specification and should be submitted separately in accordance with 37 CFR 1.84. (Both a substitute specification in compliance with 37 CFR 1.125 and new drawings in compliance with 37 CFR 1.84 and 1.121(d) are required).

Applicant is cautioned that correction of the above items may cause the specification and drawings page count to exceed 100 pages. If the specification and drawings exceed 100 pages, applicant will need to submit the required application size fee.

Replies should be mailed to:

Mail Stop Missing Parts
Commissioner for Patents
P.O. Box 1450
Alexandria VA 22313-1450

Registered users of EFS-Web may alternatively submit their reply to this notice via EFS-Web.
<https://sportal.uspto.gov/authenticate/AuthenticateUserLocalEPF.html>

For more information about EFS-Web please call the USPTO Electronic Business Center at **1-866-217-9197** or visit our website at <http://www.uspto.gov/ebc>.

If you are not using EFS-Web to submit your reply, you must include a copy of this notice.

/tnguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 1199
)	
John R. EVANS <i>et al.</i>)	
)	
Application No.: 12/285,887)	Group Art Unit: 1617
)	
Filed: October 15, 2008)	Prior Examiner: Unassigned
)	
FOR: FORMULATION)	Date: March 3, 2010

STATEMENT ACCOMPANYING SUBSTITUTE SPECIFICATION

In response to the Notice to File Corrected Application Papers, attached are a clean copy and a marked up copy of the second substitute specification in which the flow diagram on page 19 has been removed and resubmitted as Figure 2. The attached copies of the substitute specification do not include prohibited new matter. In particular, referring to the "marked up" copy:

- At page 19, the flow diagram was deleted to create a new formal Figure 2.
- At page 6, "Brief Description" of the new Figure 2 was inserted.
- At page 18, the "below" reference was changed to refer to "Figure 2."
- Figure 1 was modified to designate "1/2" at the top

If there are any additional fees due in connection with the filing of this response, please charge the fees to our Deposit Account No. 50-0310. If a fee is required for an extension of time under 37 C.F.R. §1.136 not accounted for above, such an extension is requested and the fee should also be charged to our Deposit Account.

Respectfully Submitted,
Morgan Lewis & Bockius LLP

Date: **March 3, 2010**
Morgan Lewis & Bockius LLP
Customer No. **09629**
1111 Pennsylvania Avenue, N.W.
Washington, D.C. 20004
Tel. No.: 202-739-3000

By: /Donald Bird/
Donald J. Bird
Registration No.25,323
Tel. No.: (202) 739-5320
Fax No.: (202) 739-3001




Figure 1

FLOW DIAGRAM OF MANUFACTURING




Figure 2

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Donald J. Bird
Attorney Docket Number:	056291-5004-02

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Pages:				
Claims:				
Miscellaneous-Filing:				
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				
Extension - 5 months with \$0 paid	1255	1	2350	2350

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Miscellaneous:				
Total in USD (\$)				2350

Electronic Acknowledgement Receipt

EFS ID:	7135227
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	09629
Filer:	Donald J. Bird
Filer Authorized By:	
Attorney Docket Number:	056291-5004-02
Receipt Date:	03-MAR-2010
Filing Date:	15-OCT-2008
Time Stamp:	17:40:20
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Deposit Account
Payment was successfully received in RAM	\$2350
RAM confirmation Number	4396
Deposit Account	500310
Authorized User	

The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows:

Charge any Additional Fees required under 37 C.F.R. Section 1.16 (National application filing, search, and examination fees)

Charge any Additional Fees required under 37 C.F.R. Section 1.17 (Patent application and reexamination processing fees)

File Listing:					
Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1		056291-5004-SubstituteSpecification-Clean.pdf	217584 59458866282b262bd94b557102e80ff69a76e4aa	yes	21
Multipart Description/PDF files in .zip description					
Document Description			Start	End	
Specification			1	20	
Abstract			21	21	
Warnings:					
Information:					
2	Specification	056291-5004-SubstituteSpecification-Marked.pdf	184205 44acc667cffffa425dc7c9d0f3e2455b24d192da	no	22
Warnings:					
Information:					
3	Miscellaneous Incoming Letter	056291-5004-StatementAccomSubstituteSpecification.pdf	91194 a21304a1693605cb782d50e3929611c7bd54f64d	no	1
Warnings:					
Information:					
4	Drawings-only black and white line drawings	Figure1.pdf	50623 8f7fd8fb2fb09a76e666027ded6c1d066e656e6	no	1
Warnings:					
Information:					
5	Drawings-only black and white line drawings	Figure2.pdf	56407 3050fd5f0cdb31d94da5f8cb7e1225c57d59bbbc	no	1
Warnings:					
Information:					
6	Fee Worksheet (PTO-875)	fee-info.pdf	29814 237e4bf3c5593664445d0c482604308e86a54b9	no	2
Warnings:					
Information:					
Total Files Size (in bytes):			629827		

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

Application No. 12/285,887

Second Substitute Specification

Clean Version

FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a Continuation Application of copending U.S. Patent Application
5 No. 10/872,784, filed June 22, 2004, which claims benefit of U.S. Patent Application No.
09/756,291, filed January 9, 2001 which claims the benefit of Great Britain Application No.
0008837.7 filed April 12, 2000 and Great Britain Application No. 0000313.7, filed January
10, 2000, all of which are incorporated herein by reference in their entireties.

10 BACKGROUND OF THE INVENTION

Field of the Invention

The invention relates to a novel sustained release pharmaceutical formulation adapted
for administration by injection containing the compound
7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol.

15

Description of the Related Art

Oestrogen deprivation is fundamental to the treatment of many benign and malignant
diseases of the breast and reproductive tract. In premenopausal women, this is achieved by
the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in
20 postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with
antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER)
present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal
antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness
25 is often limited by the partial agonism they display, which results in an incomplete blockade
of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

The potential for nonsteroidal antioestrogens to display agonistic properties prompted
the search for novel compounds that would bind ER with high affinity without activating any
of the normal transcriptional hormone responses and consequent manifestations of oestrogens.
30 Such molecules would be “pure” antioestrogens, clearly distinguished from tamoxifen-like
ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such
compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for

the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphonyl side chain in the 7 α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989).

5 One of these, 7 α -[9-(4,4,5,5,5-pentafluoropentyl sulphanyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have
10 promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphonyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable
15 salts thereof and any possible solvates of either thereof.

Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the
20 uterotrophic activity of tamoxifen.

Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a
25 reduction of tumour invasiveness.

In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does
30 not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-
5 1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

10 Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml^{-1} (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

15 Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

In the formulations within Table 1 a number of different oils are used to solubilise the
20 compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

25

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

<u>PRODUCT NAME</u>	<u>STEROID</u>	<u>DOSE</u>	<u>TYPE</u>	<u>COMP'.</u>	<u>SOURCE</u>	<u>OIL</u>	<u>BzBz</u>	<u>BzOH</u>	<u>EtO H</u>	<u>DOSE</u>	<u>DOSING</u>
SUSTANON 100	Testosterone propionate	30mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
	Testosterone	60mg									
	phenylpropionate										
	Testosterone isocaproate	60mg									
PROLUTON DEPOT	Testosterone decanoate	100mg	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%		1 or 2ml	1 week	
	Hydroxy progesterone hexanoate	250mgml ⁻¹									
TOCOGESTAN	Hydroxy progesterone enantate	200mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%		2ml	< 1week	
	Progesterone	50mg									
	α -Tocopherol	250mg									
TROPHOBOLINE	Estrapronicate	1.3mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%		1ml	15 to 30 days	
	Nandrolone undecanoate	50mg									
	Hydroxyprogesterone heptanoate	80mg									
NORISTERAT	Norethisterone oentanhoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES		1ml	8 weeks	
BENZO- GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis			1ml	1 week	
PROGESTERONE -RETARD	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES		1 or 2ml	1 week	
GRAVIBINAN	Estradiol 17- β -valerate	5mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES		1 or	1 - 2	
	Hydroxyprogesterone caproate	250mgml ⁻¹							2ml	weeks	

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis	75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%	
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestrogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%	

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
5 % are w/v and * approximate as measured directly from a single sample

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

15

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

Figure 2 shows a process flow diagram associated with the Formulation Example.

DETAILED DESCRIPTION OF THE INVENTION

Table 2 shows the solubility of fulvestrant in a number of different solvents.

25

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20

Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major
5 constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a
10 therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

Currently guidelines recommend that no more than 5mls of liquid is injected
15 intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high
20 concentrations of an alcohol concentrations of $>50\text{mgml}^{-1}$ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of $<5\text{ml}$ - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50mgml^{-1} - see Table 3 below. The finding is surprising since

the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

Therefore, we present as a feature of the invention a pharmaceutical formulation
5 comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

10 Another feature of the invention is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for
15 intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma
20 fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for
intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous
25 ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

For the avoidance of any doubt when using the term % weight per volume of
formulation for the constituents of the formulation we mean that within a unit volume of the
30 formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Preferred pharmaceutical formulations of the invention are as described above

5 wherein:

1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml^{-1} .
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 10 3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably
 15 up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v,
 20 at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and
 25 preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the
5 formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral
10 administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

15 It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of
20 ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v,
25 at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected
30 from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-

25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as
5 described in the US, British, European and Japanese pharmacopoeias).

Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

10 By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after
15 intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the

injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release
5 rate was not sufficiently high to be clinically significant.

2. Our findings from studies using ^{14}C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

It would be expected that ethanol will dissipate at least as quickly, if not more rapidly,
10 from the injection site.

It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

15 We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

20 By use of the term “therapeutically significant levels” we mean that blood plasma concentrations of at least 2.5 ngml^{-1} , ideally at least 3 ngml^{-1} , at least 8.5 ngml^{-1} , and up to 12 ngml^{-1} of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 15 ngml^{-1} .

By use of the term “extended release” we mean at least two weeks, at least three
25 weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

It will be understood that the attendant physician may wish to administer the
30 intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results
5 clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

Table 3Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25⁰C

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

5 **Solubility comparisons of fulvestrant in oil based formulations with and without alcohols**

		Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
10	Formulation ^(a)	Complete vehicle	Vehicle minus alcohols
	Castor oil based	81.2	12.6
15	Miglyol 812-N based	86.8	1.7
	Sesame seed/Castor oil (1:1) based	70.1	4.4
	Sesame seed oil based	45.7	0.7
20	Arachis oil based	40.2	< 0.2

25 ^(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

		Days						
30	Formulation ^a	2	3	4	7	10	30	51
35	Formulation F1 castor oil based	0	0	0	0	0	0	0
	Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
40	Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

45 ^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile
5 over five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

10 Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per
15 volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

20 A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-
25 acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

Preferably 5ml of the intramuscular injection is administered.

A further feature of the invention is use of fulvestrant in the preparation of a
30 pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent
5 indications such as breast cancer and gynaecological conditions, such as endometriosis.

In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical
10 structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a
15 pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

20 Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

25 Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay
30 as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary

pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram of Figure 2.

5

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol

10 10% weight per volume of ethanol

15% weight per volume of benzyl benzoate

250mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil

15

References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids* 1989; 54:71-99.
- 5 2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- 10 4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- 15 5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112: R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

ABSTRACT OF THE DISCLOSURE

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound

- 5 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

Application No. 12/285,887

Second Substitute Specification

Marked Version

FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a Continuation Application of copending U.S. Patent Application
5 No. 10/872,784, filed June 22, 2004, which claims benefit of U.S. Patent Application No.
09/756,291, filed January 9, 2001 which claims the benefit of Great Britain Application No.
0008837.7 filed April 12, 2000 and Great Britain Application No. 0000313.7, filed January
10, 2000, all of which are incorporated herein by reference in their entireties.

10 BACKGROUND OF THE INVENTION

Field of the Invention

The invention relates to a novel sustained release pharmaceutical formulation adapted
for administration by injection containing the compound
7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol.

15

Description of the Related Art

Oestrogen deprivation is fundamental to the treatment of many benign and malignant
diseases of the breast and reproductive tract. In premenopausal women, this is achieved by
the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in
20 postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with
antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER)
present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal
antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness
25 is often limited by the partial agonism they display, which results in an incomplete blockade
of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

The potential for nonsteroidal antioestrogens to display agonistic properties prompted
the search for novel compounds that would bind ER with high affinity without activating any
of the normal transcriptional hormone responses and consequent manifestations of oestrogens.
30 Such molecules would be “pure” antioestrogens, clearly distinguished from tamoxifen-like
ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such
compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for

the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphonyl side chain in the 7 α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989).

5 One of these, 7 α -[9-(4,4,5,5,5-pentafluoropentyl sulphanyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have
10 promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphonyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable
15 salts thereof and any possible solvates of either thereof.

Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the
20 uterotrophic activity of tamoxifen.

Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a
25 reduction of tumour invasiveness.

In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does
30 not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-
5 1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

10 Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml^{-1} (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

15 Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

In the formulations within Table 1 a number of different oils are used to solubilise the
20 compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

25

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

<u>PRODUCT NAME</u>	<u>STEROID</u>	<u>DOSE</u>	<u>TYPE</u>	<u>COMP'.</u>	<u>SOURCE</u>	<u>OIL</u>	<u>BzBz</u>	<u>BzOH</u>	<u>EtO</u> <u>H</u>	<u>DOSE</u>	<u>DOSING</u>
SUSTANON 100	Testosterone propionate	30mg	Androgen	Organon	ABPI Data Sheet Comp.1999	Arachis		0.1ml		1ml	3 weeks
	Testosterone phenylpropionate	60mg									
	Testosterone isocaproate	60mg									
	Testosterone decanoate	100mg									
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp.1999	Castor	up to 46%		1 or 2ml	1 week	
TOCOGESTAN	Hydroxy progesterone enantate	200mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%			2ml	< 1week
	Progesterone	50mg									
	α -Tocopherol	250mg									
TROPHOBOLINE	Estrapronicate	1.3mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%			1ml	15 to 30 days
	Nandrolone undecanoate	50mg									
	Hydroxyprogesterone heptanoate	80mg									
NORISTERAT	Norethisterone oentanhoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp.1999	Castor	YES		1ml	8 weeks	
BENZO- GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis				1ml	1 week
PROGESTERONE -RETARD	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES			1 or 2ml	1 week
GRAVIBINAN	Estradiol 17- β -valerate	5mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES			1 or	1 - 2
	Hydroxyprogesterone caproate	250mgml ⁻¹								2ml	weeks

PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis	75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹ 40mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964) 53(8) 891	Castor	78% 58%	20% 40%	2% 2%	
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progestrogen	DMS	J.Pharm. Sci.(1964) 53(8) 891	Castor	YES	YES	up to 2%	

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal
5 % are w/v and * approximate as measured directly from a single sample

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

Figure 2 shows a process flow diagram associated with the Formulation Example.

DETAILED DESCRIPTION OF THE INVENTION

Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20

Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major
5 constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a
10 therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

Currently guidelines recommend that no more than 5mls of liquid is injected
15 intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high
20 concentrations of an alcohol concentrations of $>50\text{mgml}^{-1}$ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of $<5\text{ml}$ - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50mgml^{-1} - see Table 3 below. The finding is surprising since

the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

Therefore, we present as a feature of the invention a pharmaceutical formulation
5 comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

10 Another feature of the invention is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for
15 intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma
20 fulvestrant concentration for at least 2 weeks.

Further features of the invention include a pharmaceutical formulation adapted for
intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or
less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1%
(preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous
25 ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

For the avoidance of any doubt when using the term % weight per volume of
formulation for the constituents of the formulation we mean that within a unit volume of the
30 formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Preferred pharmaceutical formulations of the invention are as described above

5 wherein:

1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
- 10 3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably
 15 up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v,
 20 at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and
 25 preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the
5 formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral
10 administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

15 It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of
20 ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v,
25 at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected
30 from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-

25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as
5 described in the US, British, European and Japanese pharmacopoeias).

Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

10 By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after
15 intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the

injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.

2. Our findings from studies using ^{14}C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

By use of the term “therapeutically significant levels” we mean that blood plasma concentrations of at least 2.5 ngml^{-1} , ideally at least 3 ngml^{-1} , at least 8.5 ngml^{-1} , and up to 12 ngml^{-1} of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 15 ngml^{-1} .

By use of the term “extended release” we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results
5 clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

Table 3Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25⁰C

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

5 **Solubility comparisons of fulvestrant in oil based formulations with and without alcohols**

		Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
10	Formulation ^(a)	Complete vehicle	Vehicle minus alcohols
	Castor oil based	81.2	12.6
15	Miglyol 812-N based	86.8	1.7
	Sesame seed/Castor oil (1:1) based	70.1	4.4
	Sesame seed oil based	45.7	0.7
20	Arachis oil based	40.2	< 0.2

25 ^(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of formulation on precipitation of fulvestrant at the injection site

		Days						
30	Formulation ^a	2	3	4	7	10	30	51
35	Formulation F1 castor oil based	0	0	0	0	0	0	0
	Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
40	Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

45 ^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile
5 over five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

10 Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per
15 volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

20 A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-
25 acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

Preferably 5ml of the intramuscular injection is administered.

A further feature of the invention is use of fulvestrant in the preparation of a
30 pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent
5 indications such as breast cancer and gynaecological conditions, such as endometriosis.

In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical
10 structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol; 35% or less weight of a
15 pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol.

20 Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

25 Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μ m porosity. The sterile filtrate is kept under a nitrogen overlay
30 as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary

pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

| *See also process flow diagram ~~below~~ of Figure 2.*

5

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

10% weight per volume of benzyl alcohol

10 10% weight per volume of ethanol

15% weight per volume of benzyl benzoate

250mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil

FLOW DIAGRAM OF MANUFACTURING

Ingredients/Components

Fulvestrant
Alcohol
Benzyl Alcohol

Benzyl Benzoate

Castor Oil

Process

STAGE 1: DISSOLUTION OF
ACTIVE AGENT

STAGE 2: MIX

STAGE 3: MAKE TO
WEIGHT

STAGE 4: STERILE FILTRATION
(0.2µm)
INTO BULK RECEIVING VESSEL

STAGE 5: STERILE (0.2µm)
IN-LINE FILTRATION

STAGE 6: ASEPTIC FILLING,
AND STOPPERING

STAGE 7: VISUAL
INSPECTION

References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids* 1989; 54:71-99.
- 5 2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- 10 4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- 15 5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112: R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

ABSTRACT OF THE DISCLOSURE

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound

- 5 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 7 columns: APPLICATION NUMBER, FILING or 371(c) DATE, GRP ART UNIT, FIL FEE REC'D, ATTY. DOCKET NO, TOT CLAIMS, IND CLAIMS. Row 1: 12/285,887, 10/15/2008, 1617, 2754, 056291-5004-02, 23, 2

CONFIRMATION NO. 1199

UPDATED FILING RECEIPT

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Date Mailed: 03/11/2010

Receipt is acknowledged of this non-provisional patent application. The application will be taken up for examination in due course. Applicant will be notified as to the results of the examination. Any correspondence concerning the application must include the following identification information: the U.S. APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please submit a written request for a Filing Receipt Correction. Please provide a copy of this Filing Receipt with the changes noted thereon. If you received a "Notice to File Missing Parts" for this application, please submit any corrections to this Filing Receipt with your reply to the Notice. When the USPTO processes the reply to the Notice, the USPTO will generate another Filing Receipt incorporating the requested corrections

Applicant(s)

John R. Evans, Macclesfield, UNITED KINGDOM;
Rosalind U. Grundy, Macclesfield, UNITED KINGDOM;

Power of Attorney: None

Domestic Priority data as claimed by applicant

This application is a CON of 10/872,784 06/22/2004 PAT 7,456,160

Foreign Applications

UNITED KINGDOM 0008837.7 04/12/2000
UNITED KINGDOM 0000313.7 01/10/2000

If Required, Foreign Filing License Granted: 11/03/2008

The country code and number of your priority application, to be used for filing abroad under the Paris Convention, is US 12/285,887

Projected Publication Date: 06/17/2010

Non-Publication Request: No

Early Publication Request: No

Title

Formulation

Preliminary Class

514

PROTECTING YOUR INVENTION OUTSIDE THE UNITED STATES

Since the rights granted by a U.S. patent extend only throughout the territory of the United States and have no effect in a foreign country, an inventor who wishes patent protection in another country must apply for a patent in a specific country or in regional patent offices. Applicants may wish to consider the filing of an international application under the Patent Cooperation Treaty (PCT). An international (PCT) application generally has the same effect as a regular national patent application in each PCT-member country. The PCT process **simplifies** the filing of patent applications on the same invention in member countries, but **does not result** in a grant of "an international patent" and does not eliminate the need of applicants to file additional documents and fees in countries where patent protection is desired.

Almost every country has its own patent law, and a person desiring a patent in a particular country must make an application for patent in that country in accordance with its particular laws. Since the laws of many countries differ in various respects from the patent law of the United States, applicants are advised to seek guidance from specific foreign countries to ensure that patent rights are not lost prematurely.

Applicants also are advised that in the case of inventions made in the United States, the Director of the USPTO must issue a license before applicants can apply for a patent in a foreign country. The filing of a U.S. patent application serves as a request for a foreign filing license. The application's filing receipt contains further information and guidance as to the status of applicant's license for foreign filing.

Applicants may wish to consult the USPTO booklet, "General Information Concerning Patents" (specifically, the section entitled "Treaties and Foreign Patents") for more information on timeframes and deadlines for filing foreign patent applications. The guide is available either by contacting the USPTO Contact Center at 800-786-9199, or it can be viewed on the USPTO website at <http://www.uspto.gov/web/offices/pac/doc/general/index.html>.

For information on preventing theft of your intellectual property (patents, trademarks and copyrights), you may wish to consult the U.S. Government website, <http://www.stopfakes.gov>. Part of a Department of Commerce initiative, this website includes self-help "toolkits" giving innovators guidance on how to protect intellectual property in specific countries such as China, Korea and Mexico. For questions regarding patent enforcement issues, applicants may call the U.S. Government hotline at 1-866-999-HALT (1-866-999-4158).

LICENSE FOR FOREIGN FILING UNDER

Title 35, United States Code, Section 184

Title 37, Code of Federal Regulations, 5.11 & 5.15

GRANTED

The applicant has been granted a license under 35 U.S.C. 184, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" followed by a date appears on this form. Such licenses are issued in all applications where the conditions for issuance of a license have been met, regardless of whether or not a license may be required as

set forth in 37 CFR 5.15. The scope and limitations of this license are set forth in 37 CFR 5.15(a) unless an earlier license has been issued under 37 CFR 5.15(b). The license is subject to revocation upon written notification. The date indicated is the effective date of the license, unless an earlier license of similar scope has been granted under 37 CFR 5.13 or 5.14.

This license is to be retained by the licensee and may be used at any time on or after the effective date thereof unless it is revoked. This license is automatically transferred to any related applications(s) filed under 37 CFR 1.53(d). This license is not retroactive.

The grant of a license does not in any way lessen the responsibility of a licensee for the security of the subject matter as imposed by any Government contract or the provisions of existing laws relating to espionage and the national security or the export of technical data. Licensees should apprise themselves of current regulations especially with respect to certain countries, of other agencies, particularly the Office of Defense Trade Controls, Department of State (with respect to Arms, Munitions and Implements of War (22 CFR 121-128)); the Bureau of Industry and Security, Department of Commerce (15 CFR parts 730-774); the Office of Foreign Assets Control, Department of Treasury (31 CFR Parts 500+) and the Department of Energy.

NOT GRANTED

No license under 35 U.S.C. 184 has been granted at this time, if the phrase "IF REQUIRED, FOREIGN FILING LICENSE GRANTED" DOES NOT appear on this form. Applicant may still petition for a license under 37 CFR 5.12, if a license is desired before the expiration of 6 months from the filing date of the application. If 6 months has lapsed from the filing date of this application and the licensee has not received any indication of a secrecy order under 35 U.S.C. 181, the licensee may foreign file the application pursuant to 37 CFR 5.15(b).



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 4 columns: APPLICATION NUMBER (12/285,887), FILING OR 371(C) DATE (10/15/2008), FIRST NAMED APPLICANT (John R. Evans), ATTY. DOCKET NO./TITLE (056291-5004-02)

CONFIRMATION NO. 1199

PUBLICATION NOTICE

9629
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004



Title:Formulation

Publication No.US-2010-0152149-A1

Publication Date:06/17/2010

NOTICE OF PUBLICATION OF APPLICATION

The above-identified application will be electronically published as a patent application publication pursuant to 37 CFR 1.211, et seq. The patent application publication number and publication date are set forth above.

The publication may be accessed through the USPTO's publically available Searchable Databases via the Internet at www.uspto.gov. The direct link to access the publication is currently http://www.uspto.gov/patft/.

The publication process established by the Office does not provide for mailing a copy of the publication to applicant. A copy of the publication may be obtained from the Office upon payment of the appropriate fee set forth in 37 CFR 1.19(a)(1). Orders for copies of patent application publications are handled by the USPTO's Office of Public Records. The Office of Public Records can be reached by telephone at (703) 308-9726 or (800) 972-6382, by facsimile at (703) 305-8759, by mail addressed to the United States Patent and Trademark Office, Office of Public Records, Alexandria, VA 22313-1450 or via the Internet.

In addition, information on the status of the application, including the mailing date of Office actions and the dates of receipt of correspondence filed in the Office, may also be accessed via the Internet through the Patent Electronic Business Center at www.uspto.gov using the public side of the Patent Application Information and Retrieval (PAIR) system. The direct link to access this status information is currently http://pair.uspto.gov/. Prior to publication, such status information is confidential and may only be obtained by applicant using the private side of PAIR.

Further assistance in electronically accessing the publication, or about PAIR, is available by calling the Patent Electronic Business Center at 1-866-217-9197.

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 5 columns: APPLICATION NO., FILING DATE, FIRST NAMED INVENTOR, ATTORNEY DOCKET NO., CONFIRMATION NO.
12/285,887 10/15/2008 John R. Evans 056291-5004-02 1199

9629 7590 12/21/2010
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004

EXAMINER

HUI, SAN MING R

ART UNIT PAPER NUMBER

1628

MAIL DATE DELIVERY MODE

12/21/2010

PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Office Action Summary	Application No. 12/285,887	Applicant(s) EVANS ET AL.	
	Examiner San-ming Hui	Art Unit 1628	

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) OR THIRTY (30) DAYS, WHICHEVER IS LONGER, FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) Responsive to communication(s) filed on ____.
- 2a) This action is **FINAL**. 2b) This action is non-final.
- 3) Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

Disposition of Claims

- 4) Claim(s) 1-23 is/are pending in the application.
4a) Of the above claim(s) ____ is/are withdrawn from consideration.
- 5) Claim(s) ____ is/are allowed.
- 6) Claim(s) 1-23 is/are rejected.
- 7) Claim(s) ____ is/are objected to.
- 8) Claim(s) ____ are subject to restriction and/or election requirement.

Application Papers

- 9) The specification is objected to by the Examiner.
- 10) The drawing(s) filed on ____ is/are: a) accepted or b) objected to by the Examiner.
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
- 11) The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

Priority under 35 U.S.C. § 119

- 12) Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
a) All b) Some * c) None of:
1. Certified copies of the priority documents have been received.
2. Certified copies of the priority documents have been received in Application No. 10/872784.
3. Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

* See the attached detailed Office action for a list of the certified copies not received.

Attachment(s)

- | | |
|---|---|
| 1) <input type="checkbox"/> Notice of References Cited (PTO-892) | 4) <input type="checkbox"/> Interview Summary (PTO-413)
Paper No(s)/Mail Date. ____. |
| 2) <input type="checkbox"/> Notice of Draftsperson's Patent Drawing Review (PTO-948) | 5) <input type="checkbox"/> Notice of Informal Patent Application |
| 3) <input checked="" type="checkbox"/> Information Disclosure Statement(s) (PTO/SB/08)
Paper No(s)/Mail Date <u>9/4/09</u> . | 6) <input type="checkbox"/> Other: ____. |

DETAILED ACTION

This is a continuation of US 10/872,784, filed 6/22/2004, now patent 7,456,160, which is a continuation of US 09/756,291, filed 1/9/2001, now patent 6,774,122. The instant application also claims the benefit of UNITED KINGDOM 0000313.7, filed 01/10/2000 and UNITED KINGDOM 0008837.7, filed 04/12/2000.

Claims 1-23 are pending.

Claim Rejections - 35 USC § 103

The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

The factual inquiries set forth in *Graham v. John Deere Co.*, 383 U.S. 1, 148 USPQ 459 (1966), that are applied for establishing a background for determining obviousness under 35 U.S.C. 103(a) are summarized as follows:

1. Determining the scope and contents of the prior art.
2. Ascertaining the differences between the prior art and the claims at issue.
3. Resolving the level of ordinary skill in the pertinent art.
4. Considering objective evidence present in the application indicating obviousness or nonobviousness.

Claims 1-23 are rejected under 35 U.S.C. 103(a) as being unpatentable over Dukes (EP 0 346 014) in view of Lehmann et al. (US Patent Re. 28,690), GB 1 569 286 (herein after referred as '286), Osborne et al., Journal of National Cancer Institute,

Art Unit: 1628

1995;87(10):746-750, and Remington (Remington's Pharmaceutical Sciences, 18th ed., 1990, page 219), all of the references are of record in the parent application.

Dukes teaches antiestrogen agents, including fulvestrant, are useful in treating postmenopausal symptoms such as urogenital atrophy affecting the vagina (See page 3, lines 56-page 4, line 1; also page 7, line 28-29). Dukes teaches that antiestrogen agent, including fulvestrant, may be used in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol (See page 7, line 20-24).

Dukes does not expressly teach the dosage of fulvestrant to be 45mg. Dukes does not expressly teach the employment of benzyl benzoate, in the percent amount of 60% w/v or less, or 50% w/v or less, or 45% w/v or less, 40% w/v or less, or 35% w/v or less, or 30% w/v or less, 25% w/v or less, or 10-25% w/v, or 12-18% w/v, as part of the vehicle herein. Dukes does not expressly teach the total amount of the fulvestrant-containing composition administered. Dukes does not expressly teach weight amount of castor oil and benzyl alcohol. Dukes does not expressly teach the employment of ethanol as part of the vehicle herein. Dukes does not expressly teach the dosage of fulvestrant to be 250mg. Dukes does not expressly teach the plasma concentration of fulvestrant herein.

Lehmann et al. teaches that benzyl benzoate and castor oil are well-known solvent useful as conventional carriers for steroids (See col. 1, line 21-26).

'286 teaches an intramuscular injection of testosterone derivative containing castor oil/benzoate in a ratio of 6:4 (See page 1, line 17).

Osborne et al. teaches fulvestrant as useful in treating human breast cancer (See pages 747- 748, Result Section).

Remington teaches that ethanol is one of the most commonly used solvents in pharmaceutical industry (See page 219).

It would have been obvious to one of ordinary skill in the art at the time the invention was made to employ benzyl benzoate, ethanol, castor oil, and benzyl alcohol, in the herein claimed weight percent, with fulvestrant in the dosage herein, in a method of treating postmenopausal symptoms such as urogenital atrophy in the vagina.

One of ordinary skill in the art would have been motivated to employ benzyl benzoate, ethanol, castor oil, and benzyl alcohol, in the herein claimed weight percent, with fulvestrant, in the dosage herein, in a method of treating postmenopausal symptoms such as urogenital atrophy or treating breast cancer because fulvestrant is known to be useful in treating urogenital atrophy, a benign disease of the female reproductive tract in the vagina and breast cancer. Castor oil and benzyl alcohol are known to be effective as vehicle for fulvestrant. Ethanol is a commonly used pharmaceutical solvent. Benzyl benzoate is known to be effective as solvent for steroidal compounds. Since fulvestrant is an estrogen derivative, benzyl benzoate would be reasonably expected to be useful as a solvent for fulvestrant. Therefore, combining one or more agents, which are known to be useful as commonly used solvents, such as benzyl benzoate, ethanol, castor oil, and benzyl alcohol, together and incorporated such combination with an estrogen derivatives, fulvestrant, would be reasonably expected to be useful in formulating a pharmaceutical composition. Furthermore, employing such

Art Unit: 1628

fulvestrant-containing composition to treat urogenital atrophy in vagina would be reasonably expected to be effective. Moreover, the optimization of result effect parameters (e.g., amount of excipients, dosage range, and dosing regimens) is obvious as being within the skill of the artisan, absent evidence to the contrary.

One of ordinary skill in the art would have been motivated to maintain the plasma concentration of fulvestrant herein because maintaining the therapeutic plasma level of the active compounds would be considered obvious as being within the purview of the skilled artisan, absent evidence to the contrary. Furthermore, the fulvestrant composition is known to be administered through injection, therefore, putting the composition into a syringe for delivering the fulvestrant composition would be considered obvious.

Double Patenting

The nonstatutory double patenting rejection is based on a judicially created doctrine grounded in public policy (a policy reflected in the statute) so as to prevent the unjustified or improper timewise extension of the “right to exclude” granted by a patent and to prevent possible harassment by multiple assignees. A nonstatutory obviousness-type double patenting rejection is appropriate where the conflicting claims are not identical, but at least one examined application claim is not patentably distinct from the reference claim(s) because the examined application claim is either anticipated by, or would have been obvious over, the reference claim(s). See, e.g., *In re Berg*, 140 F.3d 1428, 46 USPQ2d 1226 (Fed. Cir. 1998); *In re Goodman*, 11 F.3d 1046, 29

Art Unit: 1628

USPQ2d 2010 (Fed. Cir. 1993); *In re Longi*, 759 F.2d 887, 225 USPQ 645 (Fed. Cir. 1985); *In re Van Ornum*, 686 F.2d 937, 214 USPQ 761 (CCPA 1982); *In re Vogel*, 422 F.2d 438, 164 USPQ 619 (CCPA 1970); and *In re Thorington*, 418 F.2d 528, 163 USPQ 644 (CCPA 1969).

A timely filed terminal disclaimer in compliance with 37 CFR 1.321(c) or 1.321(d) may be used to overcome an actual or provisional rejection based on a nonstatutory double patenting ground provided the conflicting application or patent either is shown to be commonly owned with this application, or claims an invention made as a result of activities undertaken within the scope of a joint research agreement.

Effective January 1, 1994, a registered attorney or agent of record may sign a terminal disclaimer. A terminal disclaimer signed by the assignee must fully comply with 37 CFR 3.73(b).

Claims 21-22 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-9 of U.S. Patent No. 6,774,122 ('122). Although the conflicting claims are not identical, they are not patentably distinct from each other because '122 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition.

Claims 21-22 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-12 of U.S. Patent No. 7,456,160 ('160). Although the conflicting claims are not identical, they are not patentably distinct

Art Unit: 1628

from each other because '160 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to San-ming Hui whose telephone number is (571) 272-0626. The examiner can normally be reached on Mon - Fri from 9:00 to 5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Brandon Fetterolf can be reached on (571) 272-2919. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

San-ming Hui
Primary Examiner
Art Unit 1628

Application/Control Number: 12/285,887
Art Unit: 1628

Page 8

/San-ming Hui/
Primary Examiner, Art Unit 1628

Index of Claims *1228588 7*	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed

-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010							
	1	✓							
	2	✓							
	3	✓							
	4	✓							
	5	✓							
	6	✓							
	7	✓							
	8	✓							
	9	✓							
	10	✓							
	11	✓							
	12	✓							
	13	✓							
	14	✓							
	15	✓							
	16	✓							
	17	✓							
	18	✓							
	19	✓							
	20	✓							
	21	✓							
	22	✓							
	23	✓							

Search Notes *1228588 7*	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

SEARCHED			
Class	Subclass	Date	Examiner
514	177, 178	12/19/10	SH

SEARCH NOTES		
Search Notes	Date	Examiner
EAST and inventor search in PALM	12/19/10	SH

INTERFERENCE SEARCH			
Class	Subclass	Date	Examiner

--	--



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
 United States Patent and Trademark Office
 Address: COMMISSIONER FOR PATENTS
 P.O. Box 1450
 Alexandria, Virginia 22313-1450
 www.uspto.gov

BIB DATA SHEET

CONFIRMATION NO. 1199

SERIAL NUMBER 12/285,887	FILING or 371(c) DATE 10/15/2008 RULE	CLASS 514	GROUP ART UNIT 1628	ATTORNEY DOCKET NO. 056291-5004-02	
APPLICANTS John R. Evans, Macclesfield, UNITED KINGDOM; Rosalind U. Grundy, Macclesfield, UNITED KINGDOM; ** CONTINUING DATA ***** This application is a CON of 10/872,784 06/22/2004 PAT 7,456,160 ** FOREIGN APPLICATIONS ***** UNITED KINGDOM 0008837.7 04/12/2000 UNITED KINGDOM 0000313.7 01/10/2000 ** IF REQUIRED, FOREIGN FILING LICENSE GRANTED ** 11/03/2008					
Foreign Priority claimed <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No 35 USC 119(a-d) conditions met <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Verified and Acknowledged <u>/SAN-MING R HUI/</u> Examiner's Signature	<input type="checkbox"/> Met after Allowance Initials	STATE OR COUNTRY UNITED KINGDOM	SHEETS DRAWINGS 1	TOTAL CLAIMS 23	INDEPENDENT CLAIMS 2
ADDRESS MORGAN LEWIS & BOCKIUS LLP 1111 PENNSYLVANIA AVENUE NW WASHINGTON, DC 20004 UNITED STATES					
TITLE Formulation					
FILING FEE RECEIVED 2754	FEES: Authority has been given in Paper No. _____ to charge/credit DEPOSIT ACCOUNT No. _____ for following:			<input type="checkbox"/> All Fees <input type="checkbox"/> 1.16 Fees (Filing) <input type="checkbox"/> 1.17 Fees (Processing Ext. of time) <input type="checkbox"/> 1.18 Fees (Issue) <input type="checkbox"/> Other _____ <input type="checkbox"/> Credit	

EAST Search History

EAST Search History (Prior Art)

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	76861	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L2	310	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L3	2043	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L4	2	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L5	6	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L6	13851	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L7	1744228	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L8	7012	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L9	4	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L10	7	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L11	13	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L12	1562	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L13	2	"6774122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L14	910	514/177.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L15	1322	514/178.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L16	1979621	castor oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L17	76861	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L18	310	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L19	2043	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L20	13851	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L21	1744228	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L22	7012	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L23	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L24	13	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L25	1562	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L26	76861	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L27	4282	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L28	2482	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L29	1306	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L30	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L31	2692	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L32	2692	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L33	76861	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L34	310	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L35	2043	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L36	2	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L37	6	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L38	13851	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L39	1744228	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L40	7012	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L41	4	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L42	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L43	13	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L44	1562	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L45	76861	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L46	4282	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L47	2482	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L48	1306	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L49	3	(((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L50	2692	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L51	81494	breast adj cancer	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L52	1783	breast adj cancer and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L53	281	breast adj cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33

L54	1131	cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:33
L55	2	"7456160".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:38
L56	2	"6,774,122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2010/12/19 22:38

EAST Search History (Interference)

< This search history is empty >

12/ 19/ 10 10:49:53 PM

C:\ Documents and Settings\ SHui\ My Documents\ EAST\ Workspaces\ 12-285887.wsp

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date
	1. US 3,164,520	January 5, 1965	Huber			
	2. US 4,212,863	July 15, 1980	Cornelius			
	3. US 4,388,307	June 14, 1983	Cavanak			

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
	4. EP 0310542A1	April 5, 1989	EPO			Yes

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

	5.	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
	6.	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
	7.	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
	8.	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
	9.	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 [pages 238-255 provided] (1998)
	10.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
	11.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
	12.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
	13.	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

Examiner	/San Ming Hui/	Date Considered	12/19/2010
----------	----------------	-----------------	------------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date
	1.	2,822,316	February 4, 1958	Richter et al.		
	2.	2,983,649	May 9, 1961	Ercoli et al.		
	3.	3,541,209	November 17, 1970	Neumann et al.		
	4.	RE 28,690	January 20, 1976	Lehmann et al.		
	5.	4,048,309	September 13, 1977	Chen et al.		
	6.	4,048,310	September 13, 1977	Chen et al.		
	7.	4,659,516	April 21, 1987	Bowler et al.		
	8.	4,888,331	December 19, 1989	Elger et al.		
	9.	5,095,129	March 10, 1992	Ottow et al.		
	10.	5,183,814	February 2, 1993	Dukes		
	11.	5,484,801	January 16, 1996	Al-Razzak et al.		
	12.	5,733,902	March 31, 1998	Schneider		
	13.	5,929,030	July 27, 1999	Hamied et al.		
	14.	20010006963	July 5, 2001	Lachnit-Fixson et al.		

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
15.	EP 0138504	Apr., 1985	EP			
16.	EP 0346014	Dec., 1989	EP			
17.	EP 0819431	Mar., 1999	EP			
18.	EP 0905143	Mar., 1999	EP			
19.	FR 6241	Sep., 1968	France			Abstract
20.	GB 817241	Jul., 1959	GB			
21.	GB 1126892	Sep., 1968	GB			
22.	GB 1207571	Oct., 1970	GB			
23.	GB 1569286	Jun., 1980	GB			
24.	JP 43-27327	Nov., 1992	Japan			
25.	JP 09-208496	Dec., 1997	Japan			Abstract
26.	JP 10-203982	Apr., 1998	Japan			
27.	JP 10-152438	Jun., 1998	Japan			Abstract
28.	JP 11-501649	Feb., 1999	Japan			
29.	JP 11-158200	Jun., 1999	Japan			
30.	SU 549118	Mar., 1977	Soviet Union			Abstract
31.	SU 676284	Jul., 1979	Soviet Union			Abstract
32.	WO 95/12383	May., 1995	WIPO			Abstract

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

Examiner	Date Considered
----------	-----------------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
33.	WO 96/19997	Jul., 1996	WIPO			Abstract
34.	WO 97/21440	Jun., 1997	WIPO			
35.	WO 97/37653	Oct., 1997	WIPO			Abstract
36.	WO 97/40823	Nov., 1997	WIPO			
37.	WO 98/11902	Mar., 1998	WIPO			Abstract
38.	WO 99/27906	Jun., 1999	WIPO			
39.	ZA 681014	Feb., 1968	South Africa			
40.	ZA 682530	Apr., 1968	South Africa			

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

41.	Anschel, "Lösungsmittel und Lösungsvermittler in Injektionen", Pharm, Ind., 1965, Vol. 27 (11a), pp. 781-787
42.	Davis et al., "17-Alpha-Hydroxyprogesterone-Caproate:...with Chemically Pure Progesterone", J. Clin. Endocrinol. And Metabolism, 1955, Vol. 15, pp. 923-930
43.	Dukes et al., "Antiuterotrophic effects of pure antioestrogen. ICI 182,780, ...the uterus in ovariectomized monkeys", J. Endocrinology, 1992, Vol. 135, pp. 239-247
44.	Dukes et al., "Antiuterotrophic effects of the pure antioestrogen ICI 182, 780 ...quantitative magnetic resonance imaging"; J. Endocrinology, 1992, Vol. 138, pp. 203-209
45.	Howell et al., "Pharmacokinetics, pharmacological and anti-tumour effects of the specific anti-oestrogen ICI 182780 in women with advanced breast cancer", British Journal of Cancer, 1996, Vol. 74, pp. 300-308
46.	Howell et al., "Response to a specific antioestrogen (ICI 182780) in tamoxifen-resistant breast cancer", The Lancet, Jan. 7, 1995, pp. 29-30
47.	Mackey et al, "Tolerability of intramuscular injections of testosterone ester in oil vehicle", Human Reproduction, vol. 10, no. 4, pp. 869-865, 1995
48.	Martindale, 32nd Ed., "Alcohol", Pharmaceutical Press, 1999, pp. 1099-1101
49.	Martindale, 32nd Ed., "Benzoates" and "Benzyl Alcohol"; Pharmaceutical Press, 1999, pp. 1102-1104
50.	Martindale, 32nd Ed., "Caster Oil"; 32nd Ed., Pharmaceutical Press, 1999, p. 1560
51.	Migally, "Effect of Castor Oil and Benzyl Benzoate Used as a Vehicle for Antiandrogens on the Adrenal Cortex", Archives of Andrology 2, 1979 pp. 365-369
52.	Osborne et al., "Comparison of the Effects of a Pure Steroidal Antiestrogen With Those of Tamoxifen in a Model of Human Breast Cancer", Journal of the National Cancer, May 1995, Vol. 87, No. 10, pp. 746-750
53.	Pellegrino, "Use of 17 α Hydroxyprogesterone Caproate in Threatened Abortion", Current Therapeutic Research, Vol. 4, No. 6, June, 1962, pp. 301-305
54.	Piver et al., "Medroxyprogesterone Acetate (Depo-Provera) vs. . . . Women with Metastatic Endometrial Adenocarcinoma", Cancer, Vol. 45, American Cancer Society, 1980, pp. 268-272
55.	Remington's Pharmaceutical Sciences, 18th ed., 1990, p. 219

Examiner	Date Considered
----------	-----------------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
John R. Evans et al.) Group Art Unit: 1628
Application No.: 12/285,887) Examiner: HUI, San Ming R.
Filed: October 15, 2008)
For: FORMULATION) Confirmation No.: 1199
)

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

**REVOCAION OF POWER OF ATTORNEY
STATEMENT UNDER 37 C.F.R. § 3.73(b)
AND GRANT OF NEW POWER OF ATTORNEY**

The undersigned, a representative authorized to sign on behalf of the assignee owning all of the interest in this patent application, hereby revokes all previous powers of attorney or authorization of agent granted in this application before the date of execution hereof.

As required by 37 C.F.R. § 3.73(b), the undersigned verifies that AstraZeneca AB is the assignee of the entire right, title, and interest in the patent application identified above by virtue of an assignment from the inventors recorded in parent Application No. 10/872,784 in the U.S. Patent and Trademark Office at Reel 015906, Frame 0402.

The undersigned representative of the Assignee hereby grants its power of attorney to the patent practitioners associated with **FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER, L.L.P., Customer Number 22,852**, to prosecute

Application No.: 12/285,887
Attorney Docket No.: 11285.0056-00000

this application and to transact all business in the Patent and Trademark Office connected therewith, and to receive the Letters Patent.

Please send all future correspondence concerning this application to Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., Customer No. 22,852.

Dated: 27 May 2011

By: _____



Name: DR ALLEN F GILES

Title: AUTHORISED REPRESENTATIVE

AstraZeneca AB

Electronic Acknowledgement Receipt

EFS ID:	10182478
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	09629
Filer:	Carlos M. Tellez
Filer Authorized By:	
Attorney Docket Number:	056291-5004-02
Receipt Date:	27-MAY-2011
Filing Date:	15-OCT-2008
Time Stamp:	13:09:13
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	no
------------------------	----

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1	Power of Attorney	Executed_power_11285-0056.pdf	36929 <small>1df66963cd97026c75078f395b2ba7dac73a9f6a</small>	no	2

Warnings:

Information:

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NUMBER	FILING OR 371(C) DATE	FIRST NAMED APPLICANT	ATTY. DOCKET NO./TITLE
12/285,887	10/15/2008	John R. Evans	11285.0056-00000

CONFIRMATION NO. 1199

POA ACCEPTANCE LETTER

22852
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413



Date Mailed: 06/07/2011

NOTICE OF ACCEPTANCE OF POWER OF ATTORNEY

This is in response to the Power of Attorney filed 05/27/2011.

The Power of Attorney in this application is accepted. Correspondence in this application will be mailed to the above address as provided by 37 CFR 1.33.

/snguyen/

Office of Data Management, Application Assistance Unit (571) 272-4000, or (571) 272-4200, or 1-888-786-0101

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
)
John R. Evans et al.) Group Art Unit: 1628
)
Application No.: 12/285,887) Examiner: HUI, San Ming R.
)
Filed: October 15, 2008) Confirmation No.: 1199
)
For: FORMULATION) **VIA EFS-WEB**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

**RESPONSE AND AMENDMENT UNDER 37 C.F.R. § 1.111 AND
PETITION FOR EXTENSION OF TIME**

In reply to the non-final Office Action mailed December 21, 2010 (“Office Action”), and pursuant to 37 C.F.R. § 1.111, Applicants hereby respectfully request reconsideration of this application in view of the following amendments and remarks. Applicants hereby petition for a three-month extension of time to respond to the Office Action. The requisite fee is being paid concurrently with this Response.

Amendments to the Claims are reflected in the listing of claims, which starts on page 2 of this paper. **Remarks** follow the amendment sections of this paper and start on page 9.

AMENDMENTS TO THE CLAIMS

Please add new claims 24-53. Please also cancel claims 1-23 without prejudice or disclaimer. This listing of claims will replace all prior versions and listings of claims in the application.

Claims 1-23 (Cancelled)

24. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation comprising:
- at least 45 mgml⁻¹ of fulvestrant;
 - a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
 - 12 - 18% w/v of benzyl benzoate; and
 - a sufficient amount of castor oil vehicle;
- wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least two weeks.
25. (New) The method of claim 24, wherein the ethanol and benzyl alcohol are present in the same weight/volume amounts.
26. (New) The method of claim 24, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.
27. (New) The method of claim 24, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.

28. (New) The method of claim 24, wherein the therapeutically significant blood plasma fulvestrant concentration is attained for at least 4 weeks.
29. (New) The method of claim 24, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
30. (New) The method of claim 24, wherein the method further comprises once monthly administration of the formulation.
31. (New) The method of claim 24, wherein the formulation comprises:
about 50 mgml⁻¹ of fulvestrant;
about 10% w/v of ethanol;
about 10% w/v of benzyl alcohol; and
about 15% w/v of benzyl benzoate;
wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.
32. (New) The method of claim 31, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
33. (New) The method of claim 32, wherein the therapeutically significant blood plasma fulvestrant concentration is attained for at least 4 weeks.
34. (New) The method of claim 33, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.

35. (New) The method of claim 34, wherein the method further comprises once monthly administration of the formulation.
36. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation consisting essentially of:
- at least 45 mgml⁻¹ of fulvestrant;
 - a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
 - 12 - 18% w/v of benzyl benzoate; and
 - a sufficient amount of castor oil vehicle;
- wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least two weeks.
37. (New) The method of claim 36, wherein the ethanol and benzyl alcohol are present in the same weight/volume amounts.
38. (New) The method of claim 36, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.
39. (New) The method of claim 36, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
40. (New) The method of claim 36, wherein the therapeutically significant blood plasma fulvestrant concentration is attained for at least 4 weeks.

41. (New) The method of claim 36, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
42. (New) The method of claim 36, wherein the method further comprises once monthly administration of the formulation.
43. (New) The method of claim 36, wherein the formulation consists essentially of:
 - about 50 mgml⁻¹ of fulvestrant;
 - about 10% w/v of ethanol;
 - about 10% w/v of benzyl alcohol; and
 - about 15% w/v of benzyl benzoate;wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.
44. (New) The method of claim 43, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
45. (New) The method of claim 44, wherein the therapeutically significant blood plasma fulvestrant concentration is attained for at least 4 weeks.
46. (New) The method of claim 45, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
47. (New) The method of claim 46, wherein the method further comprises once monthly administration of the formulation.

48. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment 5 – 5.25 mL of a formulation comprising:
- 4 - 6% w/v of fulvestrant;
 - a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
 - 12 - 18% w/v of benzyl benzoate; and
 - a sufficient amount of castor oil vehicle;
- wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least two weeks.
49. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising monthly intramuscular administration to a human in need of such treatment of a formulation comprising:
- 4 - 6% w/v of fulvestrant;
 - a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
 - 12 - 18% w/v of benzyl benzoate; and
 - a sufficient amount of castor oil vehicle;
- wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least two weeks.
50. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising monthly intramuscular administration to a human in need of such treatment of a formulation comprising:
- 4 - 6% w/v of fulvestrant;

a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
12 - 18% w/v of benzyl benzoate; and
a sufficient amount of castor oil vehicle;

wherein the formulation is administered in a divided dose; and

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml^{-1} for at least two weeks.

51. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment 5 – 5.25 mL of a formulation consisting essentially of:

4 - 6% w/v of fulvestrant;
a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;
12 - 18% w/v of benzyl benzoate; and
a sufficient amount of castor oil vehicle;

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml^{-1} for at least two weeks.

52. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising monthly intramuscular administration to a human in need of such treatment of a formulation consisting essentially of:

4 - 6% w/v of fulvestrant;
a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;

12 - 18% w/v of benzyl benzoate; and

a sufficient amount of castor oil vehicle;

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml^{-1} for at least two weeks.

53. (New) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising monthly intramuscular administration to a human in need of such treatment of a formulation consisting essentially of:

4 - 6% w/v of fulvestrant;

a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;

12 - 18% w/v of benzyl benzoate; and

a sufficient amount of castor oil vehicle;

wherein the formulation is administered in a divided dose; and

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml^{-1} for at least two weeks.

REMARKS

I. Status of the claims and amendments

Upon entry of the instant amendments, claims 24-53 will be pending in this application. Claims 1-23 are being cancelled in this Response without prejudice or disclaimer. New claims 24-53 are being added in this Response. Claims 24-35 and 48-50 are directed to methods for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation *comprising* various components. Claims 36-47 and 51-53 are identical to claims 24-35 and 48-50 except that the phrase “formulation *consisting essentially of*” replaces the phrase “formulation *comprising*” the various components.

New **claim 24** finds support, for example, in original claim 21. The recitation in claim 24 regarding “at least 45 mg ml⁻¹ of fulvestrant” finds support, for example, in the specification at ¶ [0028].¹ The recitation regarding “a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol” finds support, for example, in the specification at ¶¶ [0035], [0036], and [0042]. The recitation regarding “12 - 18% w/v of benzyl benzoate” finds support, for example, in the specification at ¶¶ [0040] and [0042]. The recitation regarding “a sufficient amount of castor oil vehicle” finds support, for example, in the specification at ¶ [0027]. The recitation regarding “a therapeutically significant blood

¹ Unless otherwise specified, all citations to the instant specification refer to the pagination from the published application US 2010/0152149.

plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least two weeks” finds support, for example, in the specification at ¶¶ [0027] and [0051].

New **claim 25** finds support, for example, in the specification at ¶ [0036]. New **claim 26** finds support, for example, in the specification at ¶ [0051]. New **claim 27** finds support, for example, in the specification at ¶ [0064]. New **claim 28** finds support, for example, in the specification at ¶ [0052]. New **claim 29** finds support, for example, in the specification at ¶ [0063]. New **claim 30** finds support, for example, in the specification at ¶ [0023]. New **claim 31** finds support, for example, in the specification at ¶ [0051] and the “Formulation Example” (e.g., ¶¶ [0072] to [0076]). New **claim 32** finds support, for example, in the specification at ¶ [0064]. New **claim 33** finds support, for example, in the specification at ¶ [0052]. New **claim 34** finds support, for example, in the specification at ¶ [0063]. New **claim 35** finds support, for example, in the specification at ¶ [0023].

Claims 48-50 share various limitations with claim 24, for which support has been identified above. The recitation in **claim 48** regarding administration of “5 – 5.25 mL” of formulation finds support, for example, in the specification at ¶ [0034]. The recitation in **claim 48** regarding “4 - 6% w/v of fulvestrant” finds support, for example, in the specification at ¶ [0025]. The recitation in **claim 49** regarding “monthly intramuscular administration” finds support, for example, in the specification at ¶ [0023]. The recitation in **claim 50** regarding “the formulation [being] administered in a divided dose” finds support, for example, in the specification at ¶ [0053].

Because **claims 36-47 and 51-53** are identical to claims 24-35 and 48-50 except for the transitional phrase (“consisting essentially of” instead of “comprising”), the same

disclosure from the instant specification cited above for claims 24-35 and 48-50 provides support for claims 36-47 and 51-53. The instant amendments do not add new matter.

II. Rejection under 35 U.S.C. § 103

The Office rejected claims 1-23 under 35 U.S.C. § 103(a) as being unpatentable over European Patent Specification No. EP 0 346 014 ("*Dukes*") in view of US Reissue Patent No. 28,690 ("*Lehmann*"), Great Britain Patent Specification No. GB 1 569 286 ("GB 1 569 286"), Osborne et al., Journal of National Cancer Institute, 87(10):746-750 (1995) ("*Osborne*"), and Remington's Pharmaceutical Sciences, 18th ed., p. 219 (1990) ("*Remington*").

Applicants cancelled claims 1-23 in the instant Response. Thus, this rejection is now moot. However, in an effort to advance prosecution, and to the extent that the Office is considering applying the arguments from the outstanding obviousness rejection to the newly added claims, Applicants will address the rejections in the Office Action below.

According to the Office, *Dukes* teaches that "antiestrogen agents, including fulvestrant, are useful in treating postmenopausal symptoms such as urogenital atrophy affecting the vagina." Office Action at 3. The Office further argues that *Dukes* "teaches that antiestrogen agent, including fulvestrant, may be used in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol." *Id.* The Office acknowledges, however, that among other deficiencies, *Dukes* "does not expressly teach the employment of benzyl benzoate . . . as part of the vehicle herein." *Id.*

In an attempt to cure the shortcomings in *Dukes*, the Office cites *Lehmann* as teaching “that benzyl benzoate and castor oil are well-known solvent useful as conventional carriers for steroids.” Office Action at 3. In the Office's view GB 1 569 286 “teaches an intramuscular injection of testosterone derivative containing castor oil/benzoate in a ratio of 6:4” (*id.*); *Osborne* “teaches fulvestrant as useful in treating human breast cancer” (*id.* at 4); and *Remington* teaches “that ethanol is one of the most commonly used solvents in pharmaceutical industry” (*id.*).

The Office concludes that:

One of ordinary skill in the art would have been motivated to employ benzyl benzoate, ethanol, castor oil, and benzyl alcohol, in the herein claimed weight percent, with fulvestrant, in the dosage herein, in a method of treating postmenopausal symptoms such as urogenital atrophy or treating breast cancer because fulvestrant is known to be useful in treating urogenital atrophy, a benign disease of the female reproductive tract in the vagina and breast cancer.

Office Action at 4. According to the Office:

[C]ombining one or more agents, which are known to be useful as commonly used solvents, such as benzyl benzoate, ethanol, castor oil, and benzyl alcohol, together and incorporated such combination with an estrogen derivatives, fulvestrant, would be reasonably expected to be useful in formulating a pharmaceutical composition.

Office Action at 4. The Office further argues that “the optimization of result effect[ive] parameters (e.g., amount of excipients, dosage range, and dosing regimens) is obvious as being within the skill of the artisan, absent evidence to the contrary.” *Id.* at 5. Applicants respectfully traverse this rejection.

II.A One of ordinary skill in the art would not have combined the cited references in the manner proposed in the rejection

The Office relies on *Dukes* to argue that formulations comprising fulvestrant “in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol” were known in the art. Office Action at 3. The Office then states that one of ordinary skill in the art would have added ethanol and benzyl benzoate to *Dukes* formulation to arrive at the formulation recited in the claims. *Id.* The Office, however, provides no explanation for why one of ordinary skill in the art would have modified *Dukes* formulation in the proposed manner. As will be explained below, one of ordinary skill in the art would not have modified *Dukes* formulation at least because the addition of benzyl benzoate would have been expected to reduce the solubility of fulvestrant in the formulation. Because fulvestrant is difficult to formulate, one of ordinary skill in the art would not have prepared a formulation in which fulvestrant was expected to have a lower solubility than that in the initial formulation. *See, e.g.*, specification at ¶ [0014].

The passage from *Dukes* cited by the Office as disclosing fulvestrant formulations indicates that when administering a pure antiestrogen by periodic intramuscular injection, an oily solution of the pure antiestrogen “containing arachis or castor oil [and] an alcohol such as benzyl alcohol” is preferred. *Dukes* at 7, ll. 19-23. *Dukes* also discloses two different formulations of fulvestrant in its working examples. Example 1 from *Dukes* discloses fulvestrant “in a mixture of propylene glycol: ethanol: water: poloxamer 407.” *Dukes* at 8, ll. 35-37. Example 2 discloses a formulation “contain[ing] 50 mg of [fulvestrant], 400 mg of benzyl alcohol and sufficient castor oil to

bringing the solution to a volume of 1 ml" (*Dukes* castor oil formulation"). *Dukes* at 9, II. 21-23.

No explanation for why one of ordinary skill in the art would have modified the *Dukes* castor oil formulation in any way is set forth in the Office Action. The Office seems to imply that one of ordinary skill in the art would have added ethanol and benzyl benzoate to *Dukes* castor oil formulation simply because ethanol and benzyl benzoate were "known to be useful as commonly used solvents." Office Action at 3. The focus in an obviousness rejection, however, is not on what one of ordinary skill in the art *could* have done, but rather "on what a person of ordinary skill in the pertinent art would have known at the time of the invention, and on what such a person *would have reasonably expected to have been able to do* in view of that knowledge." M.P.E.P. § 2141.11 (emphasis added).

In this regard, one of ordinary skill in the art attempting to improve any of the *Dukes* formulations would have determined the solubility of fulvestrant in any test solvent before adding the solvent to the formulation. See, for example, Table 2 of the instant application, which reports solubility of fulvestrant in castor oil, benzyl alcohol, ethanol, and benzyl benzoate. According to Table 2, the solubility of fulvestrant in benzyl benzoate is 6.15 mgml^{-1} , whereas the corresponding solubility in benzyl alcohol is $>200 \text{ mgml}^{-1}$. Thus, based on solubility data, one of ordinary skill in the art would have realized that fulvestrant is more than one order of magnitude more soluble in benzyl alcohol than in benzyl benzoate. Therefore, incorporating benzyl benzoate into *Dukes* castor oil formulation at the expense of reducing the concentration of benzyl alcohol, as would be required to arrive at the formulation recited in the instant claims

starting from *Dukes* disclosure, would have been expected to *decrease* the solubility of fulvestrant in the resulting formulation. None of the references cited by the Office suggests otherwise. As mentioned before, decreasing the solubility of fulvestrant in a given formulation would exacerbate the problem of finding a suitable formulation for fulvestrant.

Thus, none of the cited references would have suggested to one of ordinary skill in the art the modification of any of *Dukes* formulations by the addition of benzyl benzoate. For at least this reason, the Office has not made a *prima facie* case of obviousness and Applicants respectfully request that this rejection be withdrawn.

II.B The Office has not made the necessary factual findings to support a conclusion of obviousness

By arguing that one of ordinary skill in the art would have added ethanol and benzyl benzoate to *Dukes* castor oil formulation simply because the additional solvents were known in the art (Office Action at 3), the Office seems to be basing the rejection in a “combination of prior art elements” rationale. *See, e.g.*, M.P.E.P. § 2143.A. However, an obviousness rejection under this rationale requires, among other requisites: (1) “a finding that . . . each element [in the combination] merely performs the same function as it does separately” and (2) “a finding that one of ordinary skill in the art would have recognized that the results of the combination were predictable.” *Id.* The Office has not met either of these requirements.

First, the Office has not shown that the solvents proposed to be added to *Dukes* castor oil formulation, ethanol and benzyl benzoate, would have performed the same function in the resulting formulation as they performed separately. As explained in the

previous section, the addition of benzyl benzoate to a given fulvestrant formulation would have been expected to reduce the solubility of fulvestrant in the original formulation. Table 3 in the instant specification compares the solubility of fulvestrant in various formulations comprising ethanol, benzyl alcohol, and castor oil in the presence and absence of benzyl benzoate. In each case, the solubility of fulvestrant in the solution containing benzyl benzoate *increased* compared to the solubility of fulvestrant in the corresponding formulation without benzyl benzoate. *Id.* The trend shown in Table 3 demonstrates that benzyl benzoate is *not* “perform[ing] the same function as it does separately.”

Second, because the addition of benzyl benzoate does not decrease the solubility of fulvestrant in the resulting formulation, as would have been expected, the Office cannot find that “one of ordinary skill in the art would have recognized that the results of the combination were predictable.”

Even under an “obvious to try” rationale, the Office needs to show that “one of ordinary skill in the art could have pursued the known potential solutions *with a reasonable expectation of success.*” M.P.E.P. § 2143.E. In the instant case, as shown in Section II.A above, one of ordinary skill in the art could not have expected that adding benzyl benzoate to *Dukes* castor oil formulation, while decreasing the amount of benzyl alcohol as would be required to arrive at the formulation recited in the instant claims, would have resulted in a suitable fulvestrant formulation.

For the foregoing reasons, the Office has not met its burden of proving a *prima facie* case of obviousness. Accordingly, Applicants respectfully request that this rejection be withdrawn.

III. Double Patenting Rejection

The Office rejected claims 21-22 under the nonstatutory obviousness-type double patenting doctrine as being unpatentable over: (a) claims 1-9 of U.S. Patent No. 6,774,122 and (b) claims 1-12 of U.S. Patent No. 7,456,160.

Because Applicants cancelled claims 21 and 22 in this Response, this rejection is now moot. Accordingly, Applicants respectfully request that this rejection be withdrawn.

IV. Conclusion


In view of the foregoing amendments and remarks, Applicants respectfully request reconsideration of this application and the timely allowance of the pending claims.

Please grant any extensions of time required to enter this response and charge any required fees not included with this Response to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated: June 20, 2011

By: 

Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
)
John R. Evans et al.) Group Art Unit: 1628
)
Application No.: 12/285,887) Examiner: HUI, San Ming R.
)
Filed: October 15, 2008) Confirmation No.: 1199
)
For: FORMULATION) **VIA EFS-WEB**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

INFORMATION DISCLOSURE STATEMENT UNDER 37 C.F.R. § 1.97(c)

A. Documents Listed in the Attached SB/08 Form

Pursuant to 37 C.F.R. §§ 1.56 and 1.97(c), Applicants brings to the attention of the Examiner the documents on the attached listing. This Information Disclosure Statement is being filed after the events recited in Section 1.97(b) but, to the undersigned's knowledge, before the mailing date of either a Final action, Quayle action, or a Notice of Allowance. Under the provisions of 37 C.F.R. § 1.97(c), this Information Disclosure Statement is accompanied by a fee of \$180.00 as specified by Section 1.17(p).

Applicants respectfully request that the Examiner consider the listed documents and indicate that they were considered by making appropriate notations on the attached form.

B. Teva's Paragraph IV Letter Dated November 25, 2009

The undersigned wishes to make of record the following information. Teva Parenteral Medicines, Inc., ("Teva") filed Abbreviated New Drug Application ("ANDA") No. 200479 with the FDA seeking approval of a generic 50 mg/mL Fulvestrant injection. In connection with ANDA No. 200479, Teva sent a letter to AstraZeneca Pharmaceuticals LP dated November 25, 2009, ("Teva's Letter") concerning U.S. Patent Nos. 6,774,122 and 7,456,160 ("the '122 and '160 patents").

The instant application claims the benefit of priority from each of the '122 and '160 patents. Teva's Letter alleges that the '122 and '160 patents are obvious in light of, *inter alia*, Howell *et al.* (cited in the Information Disclosure Statement filed on June 4, 2009) and McLeskey *et al.* (cited in this Information Disclosure Statement).

All documents cited in Teva's Letter are listed in the table below. To the extent Teva's Letter provided a pinpoint citation for any of the documents, the citation is also provided below. Otherwise, the phrase "generally" appears when Teva's Letter referred to the disclosure in the given document without a citation.

References cited in Teva's Paragraph IV Letter Dated November 25, 2009, Concerning AstraZeneca's U.S. Patent Nos. 6,774,122 and 7,456,160	
Reference	Citation
U.S. Patent No. 5,183,814 to Dukes et al., and its "European cognate," European Patent Application No. EP 0 346 014	Generally; Col. 3, I. 66 - Col. 4, I. 4; Col. 6, II. 20-26; Col. 9, II. 15-25; Example 3, col. 11, II. 1-11

References cited in Teva's Paragraph IV Letter Dated November 25, 2009, Concerning AstraZeneca's U.S. Patent Nos. 6,774,122 and 7,456,160	
Reference	Citation
U.S. Patent No.4,659,516 to Bowler et al. (and European Patent Application No. EP 0 138 504, which was termed an "equivalent" of U.S. Patent No.4,659,516 in Teva's Letter)	Generally
European Patent Application No. EP 0 346 014, which was termed the "European cognate" of U.S. Patent No. 5,183,814 in Teva's Letter	Generally
Howell <i>et al.</i> , "Pharmacokinetics, Pharmacological, and Anti-tumour Effects of the Specific Anti-oestrogen ICI 182780 in Women with Advanced Breast Cancer," <i>Brit J. Cancer</i> 74:300-308 (1996).	Generally; 300; 301; 302; 303; 305 Figure 2;
McLeskey <i>et al.</i> , "Tamoxifen-Resistant Fibroblast Growth Factor-Transfected MCF-7 Cells are Cross-Resistant <i>In Vivo</i> to the Antiestrogen ICI 182,780 and Two Aromatase Inhibitors," <i>Clin. Cancer Res.</i> 4:697-711 (1998).	Generally; 698
Wakeling <i>et al.</i> , "A Potent Specific Pure Antiestrogen with Clinical. Potential," <i>Cancer Res.</i> , 51:3867-73 (1991).	Generally; 3869
U.S. Patent No. 4,212,863	Col. 1, ll. 30-32
P.K. Gupta and GA. Brazeau (eds), <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999).	405 418

AstraZeneca Pharmaceuticals LP and other AstraZeneca related corporate entities brought suit against Teva and other Teva related corporate entities charging

them with infringement of the '122 and '160 patents. The suit was filed on January 7, 2010 in the U.S. District Court for the District of Delaware and was assigned Civil Action No. 10-18-JAP.

Subsequently, Teva withdrew ANDA No. 200479 and is no longer seeking approval of a generic 50 mg/mL Fulvestrant injection from the FDA. Civil Action No. 10-18-JAP was dismissed without prejudice on June 15, 2011.

Unless already of record, all documents cited in the preceding table are being submitted to the Office in the attached SB/08 form.

C. Documents from the prosecution of European Patent Applications member of the same family as the instant application

Applicants submitted documents from the opposition against European Patent Application No. 01900186.6 with the Information Disclosure Statement filed June 4, 2009. Applicants now supplement that submission with documents submitted after the June 4, 2009, Information Disclosure Statement. Applicants are also enclosing the search reports from European Patent Application Nos. 10180667.7 and 10180661.0. European Patent Application Nos. 01900186.6, 10180667.7, and 10180661.0 are European members of the same patent family as the instant application.

This submission does not represent that a search has been made or that no better art exists and does not constitute an admission that each or all of the documents listed in the attached SB/08 form or in the table above are material or constitute "prior art." If the Examiner applies any of the documents as prior art against any claims in the application and Applicant determines that the cited documents do not constitute "prior

art" under United States law, applicant reserves the right to present to the office the relevant facts and law regarding the appropriate status of such documents.


Applicants further reserve the right to take appropriate action to establish the patentability of the disclosed invention over the listed documents, should one or more of the documents be applied against the claims of the present application.

If there is any fee due in connection with the filing of this Statement not included herein, please charge the fee to Deposit Account 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated: June 20, 2011

By: 

Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known	
				Application Number	12/285,887
				Filing Date	October 15, 2008
				First Named Inventor	John R. EVANS
				Art Unit	1628
				Examiner Name	San Ming R. Hui
Sheet	1	of	1	Attorney Docket Number	11285.0056-00000

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS						
Examiner Initials ⁷	Cite No. ¹	Document Number		Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)				
		US-				
		US-				
		US-				
		US-				
		US-				

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS							
Examiner Initials ⁷	Cite No. ¹	Foreign Patent Document		Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁸
		Country Code ³	Number ⁴ Kind Code ⁵ (if known)				

NONPATENT LITERATURE DOCUMENTS			
Examiner Initials ⁷	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation ⁸
	1	McLeskey et al., "Tamoxifen-resistant fibroblast growth factor-transfected MCF-7 cells are cross-resistant <i>in vivo</i> to the antiestrogen ICI 182,780 and two aromatase inhibitors," Clin. Cancer Res., 4:697-711 (1998).	
	2	JRF Robertson, et al., "Fulvestrant: pharmacokinetics and pharmacology," British Journal of Cancer, 90(1):S7-S10 (2004).	
	3	John F. R. Robertson, "Fulvestrant (Faslodex®)--how to make a good drug better," The Oncologist, 12:774-784 (2007).	
	4	Search Report for European Patent Application No. 10180667.7 dated November 23, 2010.	
	5	Search Report for European Patent Application No. 10180661.0 dated January 19, 2011.	
	6	Documents from the Opposition against European Patent Application No. 01900186.6 from April 21, 2009 to September 7, 2009.	

Examiner Signature		Date Considered	
--------------------	--	-----------------	--

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Carlos M. Tellez/Delia Herring
Attorney Docket Number:	11285.0056-00000

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Pages:				
Claims:				
Miscellaneous-Filing:				
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				
Extension - 3 months with \$0 paid	1253	1	1110	1110

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Miscellaneous:				
Submission- Information Disclosure Stmt	1806	1	180	180
Total in USD (\$)				1290

Electronic Acknowledgement Receipt

EFS ID:	10343633
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Carlos M. Tellez
Filer Authorized By:	
Attorney Docket Number:	11285.0056-00000
Receipt Date:	20-JUN-2011
Filing Date:	15-OCT-2008
Time Stamp:	16:59:43
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Credit Card
Payment was successfully received in RAM	\$1290
RAM confirmation Number	4028
Deposit Account	
Authorized User	

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
-----------------	----------------------	-----------	----------------------------------	------------------	------------------

1		11285-0056-00000--20- JUN-2011-- ResponseandAmendment.pdf	587464 f0c60149d0e3c9cfe78972c101304eb349e299b	yes	17
Multipart Description/PDF files in .zip description					
		Document Description	Start	End	
		Amendment/Req. Reconsideration-After Non-Final Reject	1	1	
		Claims	2	8	
		Applicant Arguments/Remarks Made in an Amendment	9	17	
Warnings:					
Information:					
2		11285-0056-00000--20- JUN-2011--IDSandSB08.pdf	73727 27bc537a476efc3075103da02bd9fd7b2751a2f	yes	6
Multipart Description/PDF files in .zip description					
		Document Description	Start	End	
		Transmittal Letter	1	5	
		Information Disclosure Statement (IDS) Form (SB08)	6	6	
Warnings:					
Information:					
3	Non Patent Literature	11285-0056-00000--20- JUN-2011--MCLESKEY.pdf	552191 19a6a50aac4a78d8423d3acab80cb2225de83120	no	15
Warnings:					
Information:					
4	Non Patent Literature	11285-0056-00000--20- JUN-2011-- JFRRobertsonFulvestraint.pdf	118217 85e15a34de0ba428157506d8e148b124bce26f17	no	4
Warnings:					
Information:					
5	Non Patent Literature	11285-0056-00000--20- JUN-2011--JohnFRRobertson. pdf	291690 41898efd490c024a52c360ac37be99c00825e857	no	12
Warnings:					
Information:					
6	Foreign Reference	11285-0056-00000--20- JUN-2011--ISR10180667-7.pdf	65906 10fa5d3615cf47dba314a172adac66c1aa834914	no	5
Warnings:					

Information:					
7	Foreign Reference	11285-0056-00000--20- JUN-2011--ISR10180661-0.pdf	66093 <small>549156ea24a6e5820e5b114f514966558a0 1342b</small>	no	5
Warnings:					
Information:					
8	Non Patent Literature	11285-0056-00000--20- JUN-2011-- DocumentsfromOpposition.pdf	697437 <small>fcc44c176cbe2b76f5ea85a1b1162d8d5e30 0552</small>	no	51
Warnings:					
Information:					
9	Fee Worksheet (SB06)	fee-info.pdf	31886 <small>ff9996e9130d928a675262ef7a75a7e64bc1 1611</small>	no	2
Warnings:					
Information:					
Total Files Size (in bytes):				2484611	
<p>This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.</p> <p><u>New Applications Under 35 U.S.C. 111</u> If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.</p> <p><u>National Stage of an International Application under 35 U.S.C. 371</u> If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.</p> <p><u>New International Application Filed with the USPTO as a Receiving Office</u> If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.</p>					

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875	Application or Docket Number 12/285,887	Filing Date 10/15/2008	<input checked="" type="checkbox"/> To be Mailed
---	---	----------------------------------	--

APPLICATION AS FILED – PART I			OTHER THAN SMALL ENTITY				
(Column 1)		(Column 2)	SMALL ENTITY <input type="checkbox"/>		OR	SMALL ENTITY	
FOR	NUMBER FILED	NUMBER EXTRA	RATE (\$)	FEE (\$)		RATE (\$)	FEE (\$)
<input type="checkbox"/> BASIC FEE <small>(37 CFR 1.16(a), (b), or (c))</small>	N/A	N/A	N/A		OR	N/A	
<input type="checkbox"/> SEARCH FEE <small>(37 CFR 1.16(k), (l), or (m))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> EXAMINATION FEE <small>(37 CFR 1.16(o), (p), or (q))</small>	N/A	N/A	N/A			N/A	
TOTAL CLAIMS <small>(37 CFR 1.16(j))</small>	minus 20 =	*	X \$ =			X \$ =	
INDEPENDENT CLAIMS <small>(37 CFR 1.16(h))</small>	minus 3 =	*	X \$ =			X \$ =	
<input type="checkbox"/> APPLICATION SIZE FEE <small>(37 CFR 1.16(s))</small>	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).						
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENT <small>(37 CFR 1.16(j))</small>							
* If the difference in column 1 is less than zero, enter "0" in column 2.			TOTAL			TOTAL	

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY				
(Column 1)		(Column 2)	(Column 3)	SMALL ENTITY		OR	SMALL ENTITY		
AMENDMENT	06/20/2011	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	RATE (\$)	ADDITIONAL FEE (\$)	
	Total <small>(37 CFR 1.16(i))</small>	* 30	Minus	** 23	=	7	OR	X \$52=	364
	Independent <small>(37 CFR 1.16(h))</small>	* 8	Minus	***3	=	5	OR	X \$220=	1100
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>								
						TOTAL ADD'L FEE	OR	TOTAL ADD'L FEE	1464

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY				
(Column 1)		(Column 2)	(Column 3)	SMALL ENTITY		OR	SMALL ENTITY		
AMENDMENT		CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	RATE (\$)	ADDITIONAL FEE (\$)	
	Total <small>(37 CFR 1.16(i))</small>	*	Minus	**	=		OR	X \$ =	
	Independent <small>(37 CFR 1.16(h))</small>	*	Minus	***	=		OR	X \$ =	
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>								
						TOTAL ADD'L FEE	OR	TOTAL ADD'L FEE	

* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.
 ** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".
 *** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".
 The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.

Legal Instrument Examiner:
 /NICHELE PETERSON/

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**
 If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Document code: WFEE

United States Patent and Trademark Office
Sales Receipt for Accounting Date: 06/29/2011

NPETERSO SALE #00000001 Mailroom Dt: 06/20/2011 060916 12285887
01 FC : 1201 1,100.00 DA



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 5 columns: APPLICATION NO., FILING DATE, FIRST NAMED INVENTOR, ATTORNEY DOCKET NO., CONFIRMATION NO.
12/285,887 10/15/2008 John R. Evans 11285.0056-00000 1199

22852 7590 09/16/2011
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

EXAMINER

HUI, SAN MING R

ART UNIT PAPER NUMBER

1628

MAIL DATE DELIVERY MODE

09/16/2011

PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Office Action Summary	Application No. 12/285,887	Applicant(s) EVANS ET AL.	
	Examiner SAN-MING HUI	Art Unit 1628	

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) OR THIRTY (30) DAYS, WHICHEVER IS LONGER, FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) Responsive to communication(s) filed on 20 June 2011.
- 2a) This action is **FINAL**.
- 2b) This action is non-final.
- 3) An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.
- 4) Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

Disposition of Claims

- 5) Claim(s) 24-53 is/are pending in the application.
- 5a) Of the above claim(s) _____ is/are withdrawn from consideration.
- 6) Claim(s) _____ is/are allowed.
- 7) Claim(s) 24-53 is/are rejected.
- 8) Claim(s) _____ is/are objected to.
- 9) Claim(s) _____ are subject to restriction and/or election requirement.

Application Papers

- 10) The specification is objected to by the Examiner.
- 11) The drawing(s) filed on _____ is/are: a) accepted or b) objected to by the Examiner.
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
- 12) The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

Priority under 35 U.S.C. § 119

- 13) Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
a) All b) Some * c) None of:
 - 1. Certified copies of the priority documents have been received.
 - 2. Certified copies of the priority documents have been received in Application No. _____.
 - 3. Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

* See the attached detailed Office action for a list of the certified copies not received.

Attachment(s)

- 1) Notice of References Cited (PTO-892)
- 2) Notice of Draftsperson's Patent Drawing Review (PTO-948)
- 3) Information Disclosure Statement(s) (PTO/SB/08)
Paper No(s)/Mail Date _____.
- 4) Interview Summary (PTO-413)
Paper No(s)/Mail Date. _____.
- 5) Notice of Informal Patent Application
- 6) Other: _____.

DETAILED ACTION

This is a continuation of US 10/872,784, filed 6/22/2004, now patent 7,456,160, which is a continuation of US 09/756,291, filed 1/9/2001, now patent 6,774,122. The instant application also claims the benefit of UNITED KINGDOM 0000313.7, filed 01/10/2000 and UNITED KINGDOM 0008837.7, filed 04/12/2000.

Applicant's amendments filed 6/20/2011 have been entered. Claims 24-53 are pending.

The outstanding rejection under 35 USC 103(a) is withdrawn due to the cancellation of the claims.

Double Patenting

The nonstatutory double patenting rejection is based on a judicially created doctrine grounded in public policy (a policy reflected in the statute) so as to prevent the unjustified or improper timewise extension of the "right to exclude" granted by a patent and to prevent possible harassment by multiple assignees. A nonstatutory obviousness-type double patenting rejection is appropriate where the conflicting claims are not identical, but at least one examined application claim is not patentably distinct from the reference claim(s) because the examined application claim is either anticipated by, or would have been obvious over, the reference claim(s). See, e.g., *In re Berg*, 140 F.3d 1428, 46 USPQ2d 1226 (Fed. Cir. 1998); *In re Goodman*, 11 F.3d 1046, 29 USPQ2d 2010 (Fed. Cir. 1993); *In re Longi*, 759 F.2d 887, 225 USPQ 645 (Fed. Cir. 1985); *In re Van Ornum*, 686 F.2d 937, 214 USPQ 761 (CCPA 1982); *In re Vogel*, 422

Art Unit: 1628

F.2d 438, 164 USPQ 619 (CCPA 1970); and *In re Thorington*, 418 F.2d 528, 163 USPQ 644 (CCPA 1969).

A timely filed terminal disclaimer in compliance with 37 CFR 1.321(c) or 1.321(d) may be used to overcome an actual or provisional rejection based on a nonstatutory double patenting ground provided the conflicting application or patent either is shown to be commonly owned with this application, or claims an invention made as a result of activities undertaken within the scope of a joint research agreement.

Effective January 1, 1994, a registered attorney or agent of record may sign a terminal disclaimer. A terminal disclaimer signed by the assignee must fully comply with 37 CFR 3.73(b).

Claims 24-53 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-9 of U.S. Patent No. 6,774,122 ('122). Although the conflicting claims are not identical, they are not patentably distinct from each other because '122 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of the solvents and the excipients are within the range taught in '122. The optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition

Art Unit: 1628

in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

Claims 24-53 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-12 of U.S. Patent No. 7,456,160 ('160). Although the conflicting claims are not identical, they are not patentably distinct from each other because '160 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of the solvents and the excipients are within the range taught in '160. The optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

Claim Rejections - 35 USC § 103

The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

Claims 24-53 are rejected under 35 U.S.C. 103(a) as being unpatentable over McKeskey et al., from IDS filed 6/20/2011 in view of Dukes (EP 0 346 014), Osborne et al., Journal of National Cancer Institute, 1995;87(20):746-750, and the abstract of Wakeling et al., The Journal of Steroid Biochemistry and Molecular Biology, 1992;43(1-3):173-177.

McKeskey et al. teaches a studies employing subcutaneous injection of fulvestrant to nude mice. The fulvestrant formulation contains 50mg/ml in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol brought to volume with castor oil (see page 698, col. 2, Drugs section).

McKeskey et al. does not expressly teach the use of fulvestrant in treating hormonal dependent diseases of breast. It does not expressly teach the dosing regimen to be once a month, intramuscular administration, or the volume administered. McKeskey et al. does not expressly teach the herein claimed serum concentration of fulvestrant.

Dukes teaches antiestrogen agents, including fulvestrant, are useful in treating postmenopausal symptoms such as urogenital atrophy affecting the vagina (See page 3, lines 56-page 4, line 1; also page 7, line 28-29). Dukes teaches that antiestrogen

Art Unit: 1628

agent, including fulvestrant, via intramuscular route of administration may be used in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol (See page 7, line 20-24).

Osborne et al. teaches fulvestrant as useful in treating human breast cancer (See pages 747- 748, Result Section).

Wakeling et al. teaches the administration of fulvestrant (ICI 182780) demonstrating the antiestrogenic effect for over a 1 month period (see the abstract).

It would have been obvious to one of ordinary skill in the art at the time the invention was made to employ fulvestrant in McKeskey's, in the herein claimed dosing regimen and dosage, for treating hormonal dependent diseases such as breast cancer and postmenopausal symptoms.

One of ordinary skill in the art would have been motivated to employ fulvestrant in McKeskey's, in the herein claimed dosing regimen and dosage, for treating hormonal dependent diseases such as breast cancer and postmenopausal symptoms. It is known in the art that fulvestrant as being useful in treating hormonal dependent disease. It is also known in the art that fulvestrant can be administered intramuscularly and its antitumor effect can last for more than 1 month. Employing McKeskey's formulation of fulvestrant for intramuscular administration would be seen as obvious since administering a relative large volume of fulvestrant (5ml) would not be appropriate for subcutaneous administration. The examiner notes that in McKeskey's study, only 0.1ml was injected via subcutaneous administration. Furthermore, the optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is

Art Unit: 1628

obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

The examiner notes that the herein claimed serum concentration is considered to be an inherent effect of the formulation of fulvestrant.

Response to Arguments

Applicant's arguments with respect to claims 24-53 have been considered but are moot in view of the new ground(s) of rejection.

Applicant's amendment necessitated the new ground(s) of rejection presented in this Office action. Accordingly, **THIS ACTION IS MADE FINAL**. See MPEP § 706.07(a). Applicant is reminded of the extension of time policy as set forth in 37 CFR 1.136(a).

A shortened statutory period for reply to this final action is set to expire THREE MONTHS from the mailing date of this action. In the event a first reply is filed within TWO MONTHS of the mailing date of this final action and the advisory action is not

Art Unit: 1628

mailed until after the end of the THREE-MONTH shortened statutory period, then the shortened statutory period will expire on the date the advisory action is mailed, and any extension fee pursuant to 37 CFR 1.136(a) will be calculated from the mailing date of the advisory action. In no event, however, will the statutory period for reply expire later than SIX MONTHS from the date of this final action.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SAN-MING HUI whose telephone number is (571)272-0626. The examiner can normally be reached on Mon - Fri from 9:00 to 5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Brandon Fetterolf can be reached on (571) 272-2919. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

San-ming Hui

Application/Control Number: 12/285,887
Art Unit: 1628

Page 9

Primary Examiner
Art Unit 1628

/San-ming Hui/
Primary Examiner, Art Unit 1628

Notice of References Cited	Application/Control No. 12/285,887	Applicant(s)/Patent Under Reexamination EVANS ET AL.	
	Examiner SAN-MING HUI	Art Unit 1628	Page 1 of 1

U.S. PATENT DOCUMENTS

*	Document Number Country Code-Number-Kind Code	Date MM-YYYY	Name	Classification
	A US-			
	B US-			
	C US-			
	D US-			
	E US-			
	F US-			
	G US-			
	H US-			
	I US-			
	J US-			
	K US-			
	L US-			
	M US-			

FOREIGN PATENT DOCUMENTS

*	Document Number Country Code-Number-Kind Code	Date MM-YYYY	Country	Name	Classification
	N				
	O				
	P				
	Q				
	R				
	S				
	T				

NON-PATENT DOCUMENTS

*	Include as applicable: Author, Title Date, Publisher, Edition or Volume, Pertinent Pages)	
U	The abstract of Wakeling et al., The Journal of Steroid Biochemistry and Molecular Biology, 1992;43:1-3:173-177	
V		
W		
X		

*A copy of this reference is not being furnished with this Office action. (See MPEP § 707.05(a).)
Dates in MM-YYYY format are publication dates. Classifications may be US or foreign.

Index of Claims 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed

-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

<input type="checkbox"/> Claims renumbered in the same order as presented by applicant		<input type="checkbox"/> CPA		<input type="checkbox"/> T.D.		<input type="checkbox"/> R.1.47			
CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011						
	1	✓							
	2	✓							
	3	✓							
	4	✓							
	5	✓							
	6	✓							
	7	✓							
	8	✓							
	9	✓							
	10	✓							
	11	✓							
	12	✓							
	13	✓							
	14	✓							
	15	✓							
	16	✓							
	17	✓							
	18	✓							
	19	✓							
	20	✓							
	21	✓							
	22	✓							
	23	✓							
	24		✓						
	25		✓						
	26		✓						
	27		✓						
	28		✓						
	29		✓						
	30		✓						
	31		✓						
	32		✓						
	33		✓						
	34		✓						
	35		✓						
	36		✓						

<i>Index of Claims</i> 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed

-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011						
	37		✓						
	38		✓						
	39		✓						
	40		✓						
	41		✓						
	42		✓						
	43		✓						
	44		✓						
	45		✓						
	46		✓						
	47		✓						
	48		✓						
	49		✓						
	50		✓						
	51		✓						
	52		✓						
	53		✓						

EAST Search History

EAST Search History (Prior Art)

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	82307	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L2	387	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L3	2488	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L4	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L5	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L6	15161	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L7	1829323	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L8	7808	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L9	4	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L10	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L11	13	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58
L12	1810	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06: 18:58

L13	2	"6774122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L14	951	514/177.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L15	1378	514/178.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L16	2093489	castor oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L17	82307	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L18	387	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L19	2488	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L20	15161	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L21	1829323	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L22	7808	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L23	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L24	13	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L25	1810	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L26	82307	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06; 18:58
L27	4762	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT;	OR	ON	2011/09/06; 18:58

			IBM_TDB			
L28	2718	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L29	1411	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L30	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L31	3264	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L32	3264	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L33	82307	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L34	387	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L35	2488	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L36	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L37	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L38	15161	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L39	1829323	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L40	7808	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L41	4	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L42	7	((benzyl adj benzoate) or	US-PGPUB;	OR	ON	2011/09/06


		(phenylmethyl adj benzoate) same solvent) same (testosterone)	USPAT; EPO; JPO; DERWENT; IBM_TDB			18:58
L43	13	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L44	1810	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L45	82307	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L46	4762	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L47	2718	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L48	1411	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L49	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L50	3264	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L51	90122	breast adj cancer	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L52	2211	breast adj cancer and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L53	342	breast adj cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L54	1407	cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L55	2	"7456160".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58
L56	2	"6,774,122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2011/09/06 18:58

EAST Search History (Interference)

< This search history is empty >

9/ 6/ 2011 7:04:28 PM

C:\ Users\ shui\ Documents\ EAST\ Workspaces\ 12-285887.wsp

Search Notes 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

SEARCHED			
Class	Subclass	Date	Examiner
514	177, 178	12/19/10	SH
514	177, 178	9/6/11	SH

SEARCH NOTES		
Search Notes	Date	Examiner
EAST and inventor search in PALM	12/19/10	SH
update search in EAST and inventor search in PALM	9/6/11	SH

INTERFERENCE SEARCH			
Class	Subclass	Date	Examiner

--	--

REQUEST FOR CONTINUED EXAMINATION (RCE) TRANSMITTAL	Application Number: 12/285,887	Confirmation Number: 1199
	Filing Date: October 15, 2008	
	First Named Inventor: John E. EVANS	
	Group Art Unit: 1628	
	Examiner: HUI, San Ming R.	
Address to: Mail Stop RCE Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450	Attorney Docket Number: 11285.0056-00000	

This is a Request for Continued Examination (RCE) under 37 C.F.R. § 1.114 of the above-identified application.

Request for Continued Examination (RCE) practice under 37 C.F.R. § 1.114 does not apply to any utility or plant application filed prior to June 8, 1995, or to any design application.

1. Submission required under 37 C.F.R. § 1.114: Note: If the RCE is proper, any previously filed unentered amendments and amendments enclosed with the RCE will be entered in the order in which they were filed unless applicant instructs otherwise. If applicant does not wish to have any previously filed unentered amendment(s) entered, applicant must request non-entry of such amendment.

- a. Previously submitted. If a final Office action is outstanding, any amendments filed after the final Office action may be considered as a submission even if this box is not checked.
- i. Consider the arguments in the Appeal Brief or Reply Brief previously filed on _____.
- ii. Other _____
- b. **DO NOT ENTER** the amendment(s) previously filed on _____. An alternate submission is attached.
- c. Enclosed submission:
- i. Amendment/Reply
- ii. Affidavit(s)/Declaration(s)
- iii. Information Disclosure Statement
- iv. Other Terminal Disclaimer

2. Miscellaneous

- a. Suspension of action on the above-mentioned application is requested under 37 C.F.R. § 1.103(c) for a period of ____ months. (Period of suspension shall not exceed 3 months; fee under 37 C.F.R. § 1.17(i) required.)
- b. Other _____

3. Fees

- a. The filing fee is calculated as follows:
- i. \$930.00 RCE fee required under 37 C.F.R. § 1.17(e)
- ii. Petition for extension of time for (one (1) Months) **\$150.00**
- iii. Other Terminal Disclaimer (\$160.00)
- b. Payment in the amount of **\$1,240.00** enclosed.
- c. The Commissioner is authorized to charge any deficiencies in the filing fees, or credit any overpayments to Deposit Account No. 06-0916.

Signature of Applicant, Attorney, or Agent Required

Name: Carlos M. Téllez (202) 408-4000 Reg. No.: 48,638

Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P.

Signature: /Carlos M. Téllez/ Date: January 17, 2012

Certificate of Mailing or Transmission

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to Commissioner for Patents, MAIL STOP RCE, P.O. Box 1450, Alexandria, VA. 22313-1450, or facsimile transmitted to the U.S. Patent and Trademark Office on:

Name:

Signature: _____ Date: _____

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)	
)	
John R. Evans et al.)	Group Art Unit: 1628
)	
Application No.: 12/285,887)	Examiner: HUI, San Ming R.
)	
Filed: October 15, 2008)	Confirmation No.: 1199
)	
For: FORMULATION)	VIA EFS-WEB

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

INFORMATION DISCLOSURE STATEMENT UNDER 37 C.F.R. § 1.97(b)

Pursuant to 37 C.F.R. §§ 1.56 and 1.97(b), Applicant brings to the attention of the Examiner the listed documents on the attached listing. This Information Disclosure Statement is being filed before the mailing date of a first Office Action after the filing of a Request for Continued Examination in the above-referenced application.

Copies of the listed foreign and non-patent literature documents are attached.

Copies of the U.S. patent publications are not enclosed.

Documents from the prosecution of European Patent Applications members of the same family as the instant application

Applicants had submitted documents in previous Information Disclosure Statements from the prosecution histories of European Patent Application Nos. 01900186.6 (EP 1 250 138), 10180667.7 (EP 2 266 573), and 10180661.0 (EP 2 286 818), which are European members of the same patent family as the instant application. Applicants now supplement those submissions with documents made of

record in those European applications after the previous Information Disclosure Statement was filed.

Applicants respectfully request that the Examiner consider the listed documents and indicate they were considered by making appropriate notations on the attached form.

This submission does not represent that a search has been made or that no better art exists and does not constitute an admission that each or all of the listed documents are material or constitute "prior art." If the Examiner applies any of the documents as prior art against any claims in the application and Applicants determine that the cited documents do not constitute "prior art" under United States law, applicants reserve the right to present to the office the relevant facts and law regarding the appropriate status of such documents.

Applicants further reserve the right to take appropriate action to establish the patentability of the disclosed invention over the listed documents, should one or more of the documents be applied against the claims of the present application.

If there is any fee due in connection with the filing of this Statement, please charge the fee to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.



Dated: January 17, 2012

By: _____
Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
John R. Evans et al.) Group Art Unit: 1628
Application No.: 12/285,887) Examiner: HUI, San Ming R.
Filed: October 15, 2008) Confirmation No.: 1199
For: FORMULATION)

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

TERMINAL DISCLAIMER

Assignee, AstraZeneca AB, duly organized under the laws of Sweden and having its principal place of business at S-151 85 Sodertalje, Sweden, represents that is the assignee of the entire right, title, and interest in and to the above-identified application, Application No. 12/285,887 ("the '887 application"), filed on October 15, 2008, entitled "Formulation," in the names of John R. EVANS, and Rosalind U. GRUDY, as indicated by the assignment duly recorded in the United States Patent and Trademark Office for U.S. Application No. 10/872,784 ("the '784 application"), now U.S. Patent No. 7,456,160 (the '887 application being a Continuation of the '784 application) at Reel 015906, Frame 0402 on October 14, 2004.

Assignee, AstraZeneca AB, further represents that it is the assignee of the entire right, title, and interest in and to U.S. Patent Nos. 6,774,122, and 7,456,160 as indicated by the assignments duly recorded in the United States Patent and Trademark Office at

Reel 011635, Frame 0063 on March 27, 2001, and Reel 015906, Frame 0402 on October 18, 2004, respectively.

To obviate a double patenting rejection, Assignee hereby disclaims, except as provided below, the terminal part of the statutory term of any patents granted on the instant application that would extend beyond the expiration date of the full statutory term, as presently shortened by any terminal disclaimer, of prior U.S. Patent Nos. 7,456,160 and 6,774,122. Assignee hereby agrees that any patent so granted on the instant application shall be enforceable only for and during such period that it and the prior patent are commonly owned. This agreement runs with any patent granted on the instant application and is binding upon the grantee, its successors, or assigns.

In making the above disclaimer, Assignee does not disclaim the terminal part of any patent granted on the instant application that would extend to the expiration date of the full statutory term of the prior patents, as presently shortened by any terminal disclaimer, in the event that the prior patent later expires for failure to pay a maintenance fee, is held unenforceable, is found invalid by a court of competent jurisdiction, is statutorily disclaimed in whole or in part, is terminally disclaimed under 37 C.F.R. § 1.321, has all claims canceled by a reexamination certificate, is reissued, or is in any manner terminated before the expiration of its full statutory term as presently shortened by any terminal disclaimer.

In accordance with the fee schedule in 37 C.F.R. § 1.20(d), the required fee of \$160.00 is being filed with this disclaimer.


If a fee is required for an extension of time under 37 C.F.R. § 1.136 not accounted for above, such an extension is requested and the fee should also be charged to Deposit Account No. 06-0916.

The undersigned is authorized to act on behalf of assignee AstraZeneca AB.

I hereby declare that all statements made of my own knowledge and belief are true and that all statements made on information and belief are believed to be true and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Respectfully submitted,

Dated: 12th January 2012

By: 
Signature
Name: ALLEN FRANK GILES
Title: AUTHORIZED SIGNATORY
Assignee: AstraZeneca AB

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)	
)	
John R. Evans et al.)	Group Art Unit: 1628
)	
Application No.: 12/285,887)	Examiner: HUI, San Ming R.
)	
Filed: October 15, 2008)	Confirmation No.: 1199
)	
For: FORMULATION)	Mail Stop RCE
)	
)	VIA EFS-WEB

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

RESPONSE TO OFFICE ACTION, SUBMISSION UNDER 37 C.F.R. § 1.114,

AND PETITION FOR EXTENSION OF TIME

In reply to the Final Office Action mailed September 16, 2011 ("Office Action"), Applicants respectfully request reconsideration of the claimed invention in view of the following amendments and remarks. This paper fulfills the requirements of a submission under 37 C.F.R. § 1.114, and is filed together with a Request for Continued Examination (RCE).

Applicants hereby petition for a one-month extension of time to respond to the Office Action, extending the period for response to January 16, 2012. The requisite extension-of-time fee is being paid concurrently with this filing.

Amendments to the Claims are reflected in the listing of claims, which starts on page 2 of this paper. **Remarks** follow the amendment sections of this paper and start on page 7.

REMARKS

I. Status of the claims and amendments

Upon entry of the instant amendments, claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 will be pending in this application. Claims 25, 28, 31, 33, 37, 40, 43, 45, and 48-53 are cancelled in this Response without prejudice or disclaimer. New claims 54-57 are added in this Response and find support, for example, in the specification at ¶ [0053].¹

Applicants amended claim 24 to recite a formulation comprising “about 50 mg/ml-1 of fulvestrant; about 10% w/v of ethanol; about 10% w/v of benzyl alcohol; and about 15% w/v of benzyl benzoate.” Support for this amendment can be found, for example, in the specification at ¶¶ [0072]-[0075]. Applicants also amended claim 24 to recite that the method achieves a therapeutically significant blood plasma fulvestrant concentration “for at least four weeks.” Support for this amendment can be found, for example, in the specification at ¶ [0052]. Applicants amended claim 36 in a similar manner to claim 24, with support in the same portions of the specification as the amendments to claim 24 mentioned above. Applicants amended claims 32, 34, 44, and 46 to change their dependency because the claim from which each depended has been cancelled in this Response. None of the claim amendments introduce new matter.

Claims 24, 26, 27, 29, 30, 32, 34, 35, 54 and 55 are directed to methods for treating a hormonal dependent benign or malignant disease of the breast or

¹ Unless otherwise specified, all citations to the instant specification refer to the pagination in the published application, US 2010/0152149.

reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation *comprising* various components. Claims 36, 38, 39, 41, 42, 44, 46, 47, 56, and 57 are identical to claims 24, 26, 27, 29, 30, 32, 34, 35, 54 and 55 except that the phrase “formulation *consisting essentially of*” replaces the phrase “formulation *comprising*” the various components.

II. Statement of Substance of Interview under 37 C.F.R. § 1.133(b)

Applicants would like to thank Examiner San Ming Hui for granting a personal interview to Applicants on August 4, 2011. Applicants present this Statement of Substance of Interview in connection with that interview conducted between Examiner San Ming Hui, the undersigned, Dr. Paul R. Gellert (AstraZeneca Pharmaceuticals), and Mr. Allen F. Giles (AstraZeneca Pharmaceuticals).

During the interview, the undersigned and the Examiner discussed the then pending claims 24-53 and the disclosures of the following references: a) Howell et al., “Pharmacokinetics, Pharmacological, and Anti-tumour Effects of the Specific Anti-Estrogen ICI 182780 in Women with Advanced Breast Cancer,” *Brit J. Cancer* 74:300-308 (1996), b) European Patent Application No. EP 0 346 014, and McLeskey et al., “Tamoxifen-Resistant Fibroblast Growth Factor-Transfected MCF-7 Cells are Cross-Resistant In Vivo to the Antiestrogen ICI 182,780 and Two Aromatase Inhibitors,” *Clin. Cancer Res.* 4:697-711 (1998).

At the interview, the undersigned also mentioned the status of the lawsuit between AstraZeneca Pharmaceuticals and Teva Parenteral Medicines concerning a generic product containing 50 mg/ml of fulvestrant, which was also mentioned in the Information Disclosure Statement filed on June 20, 2011.

No agreement was reached and the Examiner indicated he would consider the information presented at the interview in the preparation of the next Office Action.

III. Double Patenting Rejection

The Office rejected claims 24-53 under the nonstatutory obviousness-type double patenting doctrine as being unpatentable over: (a) claims 1-9 of U.S. Patent No. 6,774,122 (“the ’122 patent”) and (b) claims 1-12 of U.S. Patent No. 7,456,160 (“the ’160 patent”).

With the sole purpose of expediting prosecution, Applicants submit a Terminal Disclaimer concurrently with this Response, which shows common ownership of the instant application and the ’122 and ’160 patents and should obviate this rejection. Accordingly, Applicants respectfully request that this rejection be withdrawn.

The filing of the Terminal Disclaimer is not an admission of the alleged obviousness of the instant claims in light of the claims in the ’122 and ’160 patents. *See, e.g.,* M.P.E.P. § 804.02.II; *Quad Environmental Technologies, Corp. v. Union Sanitary District*, 946 F.2d 870, 874 (Fed. Cir. 1991).

IV. Errors in the specification

Applicants would like to remind the Office of certain errors appearing in the instant specification. Applicants mentioned those errors in the Declaration Under 35 U.S.C §1.132 of Dr. Paul Gellert filed on August 2008 (“the Gellert Declaration”), in the parent application (Application No. 10/872,784). Applicants listed the Gellert Declaration in an Information Disclosure Statement being filed concurrently with this Response.

V. Rejections under 35 U.S.C. 103(a)

The Office rejected claims 24-53 under 35 U.S.C. 103(a) as being unpatentable over *McLeskey et al.*, *Clinical Cancer Research* 4:697-711 (1998) ("*McLeskey*"); in view of European Patent Specification No. EP 0 346 014, which names Michael Dukes as inventor ("*Dukes*"); *Osborne et al.*, *Journal of National Cancer Institute*, 87(20):746-750 (1995) ("*Osborne*"); and the abstract of *Wakeling et al.*, "ICI 182,780, *J. Steroid Biochemistry and Molecular Biology*, 43(1-3):173-177 (1992) ("*Wakeling*"). Office Action at 5.

According to the Office, *McLeskey* teaches "a stud[y] employing subcutaneous injection of fulvestrant to nude mice" and a "fulvestrant formulation contain[ing] 50mg/ml in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol brought to volume with castor oil." *Id.* The Office acknowledges that *McLeskey* does not expressly teach "the use of fulvestrant in treating hormonal dependent diseases of breast", "the dosing regimen to be once a month, intramuscular administration", "the volume administered", or "the herein claimed serum concentration of fulvestrant." *Id.*

In the Office's view, *Dukes* teaches that "antiestrogen agent[s], including fulvestrant, via intramuscular route of administration may be used in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol." *Id.* at 5-6.

The Office cites *Osborne* as teaching that fulvestrant is "useful in treating human breast cancer" (*id.* at 6) and *Wakeling* as teaching that "the administration of fulvestrant (ICI 182780) demonstrat[es] the antiestrogenic effect for over a 1 month period." *Id.*

According to the Office "[i]t would have been obvious to one of ordinary skill in the art at the time the invention was made to employ fulvestrant in [*McLeskey*], in the

herein claimed dosing regimen and dosage, for treating hormonal dependent diseases such as breast cancer and postmenopausal symptoms” because it is “known in the art that fulvestrant can be administered intramuscularly and its antitumor effect can last for more than 1 month.” *Id.*

The Office argues that “[e]mploying *McLeskey*’s formulation of fulvestrant for intramuscular administration would be seen as obvious since administering a relative large volume of fulvestrant (5ml) would not be appropriate for subcutaneous administration.” *Id.* The Office further argues that “the optimization of result effect[ive] parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan.” *Id.* Applicants respectfully traverse this rejection.

A. Declaration of Dr. Ronald J. Sawchuk

In support of Applicants’ statements regarding the state of the art and how one of ordinary skill in the art would have understood the references cited in the Office Action prior to the earliest priority date for the instant application (January 10, 2000), Applicants submit concurrently with this Response a declaration by Dr. Ronald J. Sawchuk (“Sawchuk Decl.”).

B. The Office has not made the necessary factual findings to support a conclusion of obviousness

Applicants understand that the Office’s rejection is based on at least the following two implicit assumptions: 1) that a person of ordinary skill in the art (“POSITA”) would have chosen the fulvestrant composition disclosed in *McLeskey*, from among all other known compositions in which fulvestrant had been dissolved, for the development of a

method of treating the diseases recited in the claims, and 2) that the POSITA would have had a reasonable expectation that such a composition would have been successful in those methods. Applicants respectfully submit that the Office has not provided support for those assumptions.

Applicants respectfully remind the Office that the focus in an obviousness rejection is not on what one of ordinary skill in the art *could have done*, but rather “on what a person of ordinary skill in the pertinent art would have known at the time of the invention, and on what such a person *would have reasonably expected to have been able to do* in view of that knowledge. M.P.E.P. § 2141.II (emphasis added). Thus, two of the relevant questions in this rejection are: (1) whether the knowledge in the art would have suggested to a POSITA that *McLeskey’s* composition had some advantages over other known fulvestrant compositions such that it would have been selected for the development of a method of human treatment, and (2) even assuming that a POSITA would have selected the fulvestrant *McLeskey’s* composition for the development of a method of treatment, whether in light of the knowledge in the art prior to January 10, 2000, one of ordinary skill in the art would have expected that the fulvestrant *McLeskey* composition cited by the Office would have been successful in a method of treating as recited in the instant claims.

Specifically, regarding the second question, even after *KSR*, an obviousness rejection in which the Office argues that the claimed invention would have been the result of a combination of references requires that the Office show that one of ordinary skill in the art would have had a reasonable expectation of success when combining the references. M.P.E.P. § 2143.02; *see also* M.P.E.P. § 2143.A (addressing the

requirements for a “combination of prior art elements” rationale). As will be explained below, a critical review and analysis of the state of the art at the time the instant application was filed leads to the conclusion that no such expectation existed.

Applicants will explain and discuss below the following *independent* reasons supporting withdrawal of the instant obviousness rejection:

- (1) *McLeskey* would not have suggested to a POSITA the specific %w/v composition recited in the claims;
- (2) None of the cited references would have provided a POSITA with information to select the fulvestrant composition disclosed in *McLeskey* over other known fulvestrant compositions;
- (3) The POSITA would not have had a reasonable expectation that the *McLeskey* composition would have been successful in such a method. Applicants present two independent arguments to support a lack of expectation of success:
 - a. One of ordinary skill in the art would have understood that results from subcutaneous administration, such as those in *McLeskey*, cannot be extrapolated to intramuscular administration and, thus, a POSITA would not have had an expectation as to whether the fulvestrant composition from *McLeskey* would have been effective for intramuscular delivery of fulvestrant.
 - b. Numerous variables affect the efficacy of an intramuscular formulation, among them the identity and proportion of each of its cosolvents, and a POSITA understands that the resulting variability precludes a POSITA from having an expectation a priori that a given formulation would be

successful in a given method of treatment until actual suitable in vivo experiments are performed.

1. **Independent Reason 1. McLeskey would not have suggested to a POSITA the specific %w/v composition recited in the claims**

McLeskey discloses two fulvestrant compositions. One composition was prepared by dissolving powdered drug in 100% ethanol and then spiking it into warmed peanut oil to give a final concentration of 50 mg/ml (“the *McLeskey* peanut oil composition”). *McLeskey* at 698, col. 2, under “Drugs”; Sawchuk Decl. at ¶ 16. The second composition is a 50 mg/ml fulvestrant composition “in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol, brought to volume with castor-oil” (“the *McLeskey* castor oil composition”). *Id.* The Office only refers to the *McLeskey* castor oil composition in the Office Action. Office Action at 5.

McLeskey does not specify whether the percentages in the *McLeskey* castor oil composition are in weight/volume units (%w/v, as recited in the instant claims) or in volume/volume units (%v/v). Sawchuk Decl. at ¶ 16. Dr. Sawchuk states that “[i]n a liquid composition, when a solute or cosolvent is a liquid, it is often convenient to express its concentration as a volume percent, i.e., % v/v.” *Id.* at ¶ 17.

Dr. Sawchuk provides various examples of references in which the concentration of liquid components in a composition is reported in terms of %v/v values, whereas the concentration of solid solutes is reported in terms of %w/v. *Id.* at ¶¶ 18-20.

Dr. Sawchuk concludes that “[b]ecause all of the components of the vehicle disclosed in *McLeskey* are liquids, one of ordinary skill in the art would have concluded that the

composition was described in terms of volume/volume percent units (% v/v).” *Id.* at ¶ 21.

Based on that information, Dr. Sawchuk states that “one of ordinary skill in the art would have concluded that the *McLeskey* castor oil composition on page 698 was reported in % v/v units and referred to a composition containing 10% v/v ethanol, 15% v/v benzyl benzoate, and 10% v/v benzyl alcohol in a castor oil vehicle. *Id.* at ¶ 22. This composition *is different* from a composition containing 10% w/v ethanol, 15% w/v benzyl benzoate, and 10% w/v benzyl alcohol in a castor oil vehicle. *Id.* The units of the fulvestrant composition recited in the instant claims are %w/v.

Dr. Sawchuk converted the %v/v values that *McLeskey* would have suggested to a POSITA into %w/v values, which would allow a direct comparison between the *McLeskey* castor oil composition and the composition recited in the instant claims. *Id.* at ¶¶ 23-27. Table 1 below shows the results of the calculation, with Column E having the final concentration values in %w/v units for each component in the *McLeskey* castor oil composition. *Id.* at ¶ 27.

Table 1. Information for 100 ml of the fulvestrant *McLeskey* castor oil composition

	A	B	C	D	E
Component	% v/v	Volume (ml)	Density (g/ml)	Weight (g)	% w/v
Ethanol	10	10	0.808	8.08	8.1
Benzyl Benzoate	15	15	1.118	16.77	16.8
Benzyl Alcohol	10	10	1.04156	10.42	10.4

According to Dr. Sawchuk, a POSITA “reading *McLeskey* would have concluded that *McLeskey* described a composition containing about 8.1% **w/v** ethanol, about

10.4% w/v benzyl alcohol, and about 16.8 % w/v benzyl benzoate in a castor oil vehicle.” *Id.* at ¶ 29. Therefore, *McLeskey* would not have suggested the fulvestrant composition recited in the claims comprising about 10% w/v of ethanol; about 10% w/v of benzyl alcohol; and about 15% w/v of benzyl benzoate.

In Dr. Sawchuk’s opinion, none of the references cited in the Office Action contain any disclosure “that would have suggested to one of ordinary skill in the art the modification of a composition containing about 8.1% w/v ethanol, about 16.8 % w/v benzyl benzoate, and about 10.4% w/v benzyl alcohol (i.e., the *McLeskey* castor oil composition) in an attempt to produce a composition as recited in the claims containing about 10% w/v ethanol, about 15% w/v benzyl benzoate, and about 10% v/v benzyl alcohol”. *Id.* at ¶ 30.

For at least this reason, the cited references, either alone or in combination, fail to meet all of the limitations of the claims, and Applicants respectfully request that this rejection be withdrawn.

2. **Independent Reason 2. The Office has not shown that a POSITA would have selected the *McLeskey* castor oil composition for the development of a method of treating involving intramuscular administration as instantly claimed**

Applicants remind the Office that “[o]bviousness requires more than a mere showing that the prior art includes separate references covering each separate limitation in a claim under examination.” *Unigene Laboratories, Inc. v. Apotex, Inc.*, No. 2010-1006, slip op. at 13 (Fed. Cir. Aug. 25, 2011) (internal citations omitted). Indeed, the Federal Circuit explained that:

[O]bviousness requires the additional showing that a person of ordinary skill at the time of the invention *would have selected and combined those prior art elements* in the normal course of research and development to yield the claimed invention.

Id. (internal citations omitted, italics added). Thus, in the instant rejection, the Office needs to identify reasons why a POSITA would have: a) selected and then b) combined the elements the Office argues are disclosed in the cited references. For example, the Office needs to explain why a POSITA would have selected the *McLeskey* castor oil composition (comprising ethanol, benzyl alcohol, and benzyl benzoate), from among the known fulvestrant formulations at the time of filing, to develop a method of treatment as instantly claimed.

Dr. Sawchuk explains that *McLeskey* provides no information that would have suggested to a POSITA the desirability of any of its two fulvestrant compositions over other known fulvestrant formulations. Sawchuk Decl. at ¶ 31. For example, Dr. Sawchuk points out that antitumor treatment with fulvestrant was ineffective in the *McLeskey* experiments. *Id.* at ¶ 33. In addition, with respect to the two formulations disclosed in *McLeskey*, Dr. Sawchuk highlights that *McLeskey* “did not provide any experimental data that would have allowed one of ordinary skill in the art to compare any aspect of the performance of the two fulvestrant compositions for the treatment of cancerous tumors.” *Id.* at ¶ 31.

Therefore, in Dr. Sawchuk’s opinion, “because of the lack of fulvestrant efficacy and the absence of pharmacokinetic data in *McLeskey*, one of ordinary skill in the art would have been unable to conclude whether either of the two fulvestrant *McLeskey* compositions (peanut oil or castor oil) was able to deliver a dose of fulvestrant that had

an antitumor therapeutic effect in the mice when administered subcutaneously.” *Id.* at ¶ 35.

In light of those circumstances, Dr. Sawchuk concludes that “*McLeskey* provides no information that would have led one of ordinary skill in the art to have a preference for either the peanut oil or the castor oil fulvestrant compositions over the other one, or even a preference for one of the two *McLeskey* fulvestrant compositions over other fulvestrant compositions known in the art prior to January 10, 2000.” *Id.* at ¶ 36.

Regarding fulvestrant compositions known in the art different from the *McLeskey* castor oil composition, Dr. Sawchuk lists various compositions disclosed in the references cited by the Office. *Id.* at ¶¶ 37-39. Among those formulations, Dr. Sawchuk mentions fulvestrant in an oil suspension (*Wakeling*), fulvestrant in a castor oil composition (*Osborne*), fulvestrant in a mixture of propylene glycol:ethanol:water:poloxamer 407 (*Dukes*), and fulvestrant in 400 mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml (*Dukes*), in addition to the peanut oil fulvestrant composition from *McLeskey*. Sawchuk Decl. at ¶ 37-39.

Therefore, Dr. Sawchuk concludes, “one of ordinary skill in the art had other choices besides the *McLeskey* castor oil composition with respect to potential fulvestrant formulations that could have been further investigated for the development of a method of treating humans with intramuscular fulvestrant.” *Id.* at ¶ 40. However, in Dr. Sawchuk’ opinion, “none of the references cited in the Office Action provides any information that would have guided one of ordinary skill in the art to select the *McLeskey* castor oil composition, over any of the other fulvestrant compositions

mentioned above,” for the potential development of a method of treatment as recited in the instant claims. *Id.*

In this regard, the Federal Circuit has explained that:

When a field is unreduced by direction of the prior art, and when prior art gives no indication of which parameters were critical or no direction as to which of many possible choices is likely to be successful, an invention is not obvious to try.

Unigene, slip op. at 15 (internal citation omitted).

In this case, the Office has not explained how the cited art directs a POSITA to select the *McLeskey* castor oil composition from among all other known fulvestrant compositions to develop a method as claimed. Indeed, according to Dr. Sawchuk, “none of the references cited by the Examiner provides any guidance as to the relevant factors to consider when selecting a formulation for the potential development of a method of treatment as recited in the instant claims.” Sawchuk Decl. at ¶ 41.

Nonetheless, in Dr. Sawchuk’s opinion, and judging solely on the basis of efficacy, “the *McLeskey* castor oil composition would have been among the least favored compositions to select for further development from among the fulvestrant compositions discussed above because one of ordinary skill in the art would not have been able to conclude from the information in *McLeskey* whether fulvestrant, using that composition, was sufficiently bioavailable to have an antitumor effect.” *Id.* Rather, according to Dr. Sawchuk and based on only efficacy, “the fulvestrant oil suspension from *Wakeling* would have been among the most favored formulations to select for further development from among those discussed above because at least that

formulation, when given as a single injection, showed a therapeutic antitumor effect in mice for over a one-month period.” *Id.*

Accordingly, at least because the Office has failed to explain why a POSITA would have selected the *McLeskey* castor oil composition from among those fulvestrant compositions known in the art at the time of filing to develop a method of treatment as claimed, the Office has not made a prima facie case of obviousness. Thus, Applicants respectfully request that this rejection be withdrawn.

3. **One of ordinary skill in the art would not have had an expectation that the formulation disclosed in *McLeskey*, administered subcutaneously to mice, would have been successful for intramuscular administration as instantly recited**

The POSITA would not have had a reasonable expectation that the *McLeskey* castor oil composition would have been effective to administer fulvestrant intramuscularly to achieve a therapeutic effect for at least four weeks, as instantly recited. Two independent reasons are set forth below supporting a lack of expectation of success for the combination of the references cited by the Office.

a) **Independent Reason 3. A POSITA would not have had an expectation that the results from subcutaneous injection in *McLeskey* would have been applicable to the intramuscular administration of fulvestrant**

One of ordinary skill in the art would have understood that results from subcutaneous administration, such as those reported in *McLeskey*, cannot be extrapolated to intramuscular administration. Sawchuk Decl. at ¶ 42. In Dr. Sawchuk’s view, “one of ordinary skill in the art would not have had an expectation as to whether the *McLeskey* castor oil composition would have had a therapeutic effect when

administered intramuscularly before actually performing suitable in vivo experiments.”
Id.

Dr. Sawchuk cites a few examples in which comparison of the results from subcutaneous administration yielded significant differences with respect to those from intramuscular administration. For example, a study of administration of probenecid in ewes showed that administration of the same dose of probenecid, a drug which may be used to prolong the half-life of some antibiotics in animals, resulted in significant differences in absorption and bioavailability of the drug when administered subcutaneously or intramuscularly. Guerrini V.H., Filippich L.J., English P.B., Schneider J., Cao G.R. and Bourne D.W.A., “Pharmacokinetics of probenecid in sheep”, *J Vet Pharmacol Ther.* 8(2):128-35 (1985) (“*Guerrini*”); Sawchuk Decl. at ¶ 44.

Dr. Sawchuk comments that in *Guerrini* the intramuscular dose was absorbed more rapidly and more completely than the subcutaneous dose, whereas the subcutaneous administration resulted in a “higher and more prolonged plasma probenecid concentration”. Sawchuk Decl. at ¶ 46. Due to the overall characteristics associated with subcutaneous administration, *Guerrini* reports that such a mode of administration is preferred over intramuscular administration under the conditions of its study. *Id.* at 46. Dr. Sawchuk concludes that “*this is an example where subcutaneous administration achieves a certain desired result but where intramuscular administration does not accomplish the same result.*” *Id.* (italics added).

In contrast to the pharmacokinetic profiles observed in *Guerrini*, in another study, subcutaneous administration of clindamycin, an antibiotic, resulted in faster absorption compared to intramuscular injection. Lavy E, Ziv G, Shem-Tov M, Glickman A, Dey A.,

“Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs”, *J Vet Pharmacol Ther.* 22(4):261-5 (1999) (“*Lavy*”); Sawchuk Decl. at ¶ 47. Nevertheless, the pharmacokinetic profiles in *Lavy* were such that subcutaneous administration maintained a therapeutic plasma concentration for a longer period of time than intramuscular administration. Sawchuk Decl. at ¶ 49. Based on the results from *Lavy*, Dr. Sawchuk concludes that in that case, “*one of ordinary skill in the art would not have been able to rely on data from subcutaneous administration to predict results of intramuscular administration because intramuscular administration would not have produced the same level of long-term efficacy achieved by subcutaneous administration.*” *Id.* (italics added).

In yet another study highlighting the lack of correlation between subcutaneous and intramuscular administration, Dr. Sawchuk gives an example where, in contrast to the results from *Lavy*, the absorption of the drug was more rapid and complete following intramuscular dosing than after subcutaneous injection. Ismail M., “Disposition kinetics of difloxacin after intravenous, intramuscular and subcutaneous administration in calves”, *Vet Res Commun.*, 31(4):467-76 (2007) (“*Ismail*”); Sawchuk Decl. at ¶ 50. Dr. Sawchuk states that for the purposes in *Ismail*, the intramuscular administration was preferred to subcutaneous administration. Sawchuk Decl. at ¶ 52. Dr. Sawchuk explains that “[i]n this case, contrary to the two examples above, the intramuscular administration was considered to be associated with greater clinical efficacy.” *Id.*

Dr. Sawchuk summarizes that “[t]hese three examples above show that there are significant differences in the rate and extent of absorption of a drug given by the intramuscular and subcutaneous route, even when given to the same animals in a

crossover study.” *Id.* at ¶ 53. Dr. Sawchuk concludes that “[a]s a result, it cannot be predicted a priori whether intramuscular or subcutaneous dosing will result in more rapid and/or complete drug absorption, as examples of both cases are found in the scientific literature.” *Id.* Dr. Sawchuk further explains that the examples above “*underscore the fact that efficacy of a given drug administered by a given route of dosing (e.g., intramuscular) cannot be known until appropriate comparative studies are performed in a suitable animal model.*” *Id.* at ¶ 54 (emphasis added). Dr. Sawchuk indicates that “[f]or some drugs, the desired effect might be achieved following a particular route of dosing, but for other drugs it might not,” which underlies the lack of expectation that results from subcutaneous administration could be indicative of results obtained from intramuscular administration. *Id.*

With respect to the specific results from *McLeskey*, Dr. Sawchuk concludes that “one of ordinary skill in the art having the very limited experimental subcutaneous data from *McLeskey* would not have had an expectation that the intramuscular administration of fulvestrant using the *McLeskey* castor oil composition would have been effective following intramuscular administration, such as in the method described in the claims.” *Id.* at ¶ 55.

For at least this additional reason, the instant claims are not obvious in light of the cited references and Applicants respectfully request that this rejection be withdrawn.

- b) **Independent Reason 4. Numerous variables affect the efficacy of an intramuscular formulation (e.g., identity and proportion of cosolvents) and a POSITA would have understood that the resulting variability precludes a POSITA from having an expectation a priori that a given formulation would be successful in a given method of treatment until actual suitable in vivo experiments are performed**

Dr. Sawchuk explains that “[t]ypically, during the development of an intramuscular dosage form for administration of a drug in humans, one would have carried out, among other tasks, formulation studies to determine suitable compositions in which the drug of interest is dissolved, as well as initial intramuscular dosing experiments in animals (e.g., mice, rabbits, and/or dogs) under various conditions (e.g., different compositions, different solvents, varying the proportion of the components of the composition, different drug concentrations, etc.) in order to gain an understanding of the pharmacokinetics of fulvestrant before attempting human administration.” Sawchuk Decl. at ¶ 58

Dr. Sawchuk highlights that by its very nature, the “existence of this generalized approach highlights the lack of expectation of success with respect to the extrapolation of the *McLeskey* disclosure of subcutaneous administration to mice, lacking any pharmacokinetic information, to human intramuscular administration.” *Id.*

With respect to formulation studies, Dr. Sawchuk cites to the Gellert Declaration as disclosing the importance of performing additional formulation studies in order to attempt to minimize potential side effects arising from the presence of co-solvents. *Id.* at ¶ 59-61.

Dr. Sawchuk explains, however, that “[r]egardless of how high or low the cosolvent concentrations are in a given formulation, the preparation of formulations in which a drug such as fulvestrant can be solubilized is not sufficient to ensure the desired therapeutic effect when such formulation is administered in vivo.” *Id.* at ¶ 62. Dr. Sawchuk cites to the instant specification as warning that “[s]imply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.” *Id.* (citing the specification at ¶ [0054]). Thus, Dr. Sawchuk concludes that “suitable experiments are needed to determine the pharmacokinetic performance of any candidate formulation(s).” *Id.*

According to Dr. Sawchuk, as part of that process to develop methods of human treatment, and in order to discriminate among the various formulations in development, “pharmacokinetic data could be used to characterize important variables” in the process of developing a suitable method of treatment. In the case of “drugs that are difficult to formulate, such as fulvestrant, the pharmacokinetic data could be useful to investigate the most promising formulation for the desired route of administration.” *Id.* at ¶ 63 (internal quotations omitted).

Dr. Sawchuk indicates that in a study testing seven different 100 mg/ml fulvestrant formulations intramuscularly in rabbits, the resulting pharmacokinetic data showed variability dependent on the proportion of the components in the formulations. *Id.* at ¶ 64 (citing data in PCT Application Publication No. WO 03/006064 (“WO 03/006064”). All of the fulvestrant formulations tested contained ethanol, benzyl alcohol, and benzyl benzoate in a castor oil vehicle, which are the same components of

the fulvestrant composition recited in the claims, but with different proportions for each component. *Id.*

Dr. Sawchuk explains that according to WO 03/006064, “[p]lasma levels were more variable than Control over the first 30 days following intramuscular administration of fulvestrant.” *Id.* at ¶ 65 (internal quotations omitted). Based on differences observed in the pharmacokinetic profiles, the formulations were divided into two groups, Group A, which “demonstrates “rapid release early time points, corresponding to formulations containing lower benzyl benzoate and low castor oil concentrations, while Group B shows a lower release, flatter profile corresponding to formulations containing lower benzyl benzoate and higher castor oil concentrations.” *Id.* (internal quotations omitted).

Summarizing the results in WO 03/006064, Dr. Sawchuk states that “higher benzyl benzoate concentrations in the formulation resulted in a more rapid initial release of fulvestrant, whereas lower benzyl benzoate concentrations resulted in a lower initial release, and a flatter plasma level profile.” *Id.* at ¶ 66. Dr. Sawchuk concludes that “[d]epending on the overall objective of the administration of fulvestrant, some of the fulvestrant formulations tested in WO 03/006064’s study would be more desirable than others for that given purpose and, based on the relevant pharmacokinetic profiles, one of ordinary skill in the art would be able to select one of those fulvestrant formulations for further development and/or testing.” *Id.*

Nonetheless, Dr. Sawchuk explains that “one of ordinary skill in the art would not have been able to determine whether a given fulvestrant formulation injected intramuscularly as in WO 03/006064 would have had the desired pharmacokinetic profile until such in vivo pharmacokinetic studies were carried out.” *Id.* at ¶ 67

Based on the differences in pharmacokinetic profiles from WO 03/006064, Dr. Sawchuk reiterates that “one of ordinary skill in the art knowing only the composition of a given formulation administered subcutaneously, but having no pharmacokinetic data from its intramuscular administration, would have had no expectation, one way or another, that the formulation would be effective when administered intramuscularly in a given method of treatment.” *Id.* at ¶ 68.

In particular, with respect to the disclosure in *McLeskey*, Dr. Sawchuk indicates that “one of ordinary skill in the art would not have had a reasonable expectation that the *McLeskey* castor oil composition would have been effective when given as an intramuscular injection, such as in the method of treatment recited in the claims” *Id.* at ¶ 69.

Thus, for this additional independent reason, the instant claims are not obvious over the references cited and Applicants respectfully request that this rejection be withdrawn.

VI. Conclusion

In view of the foregoing amendments and remarks, Applicants respectfully request reconsideration of this application and the timely allowance of the pending claims.

Please grant any extensions of time required to enter this response and charge any required fees not included with this Response to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.



Dated: January 17, 2012

By: _____
Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
)
John R. Evans et al.) Group Art Unit: 1628
)
Application No.: 12/285,887) Examiner: HUI, San Ming R.
)
Filed: October 15, 2008) Confirmation No.: 1199
)
For: FORMULATION) **VIA EFS-WEB**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

DECLARATION UNDER 37 C.F.R. § 1.132 OF RONALD J. SAWCHUK

I, **Ronald J. Sawchuk**, declare as follows:

Qualifications

1. My academic background and work experience are summarized in my curriculum vitae, which is attached as **Exhibit 1**.
2. Currently, I am a Professor of Pharmaceutics, Emeritus, and Morse Alumni Distinguished Teaching Professor. I am also the Director of the Bioanalytic and Pharmacokinetic Services Laboratory at the University of Minnesota.
3. I obtained a Bachelor of Science Degree in Pharmacy in 1963 from the University of Toronto. I also received a Masters of Science Degree in Pharmaceutics from the University of Toronto in 1966 and completed a Doctoral Degree (Ph. D.) in Pharmaceutical Chemistry (pharmacokinetics emphasis) at the University of California, San Francisco in 1972.

4. I joined the University of Minnesota in 1971 as an Instructor in Pharmaceutics, and served from 1972 to 1977 as an Assistant Professor of Pharmaceutics, from 1977 to 1983 as an Associate Professor of Pharmaceutics, and as a full Professor of Pharmaceutics from 1983 until my retirement in July of 2010.

5. At the University of Minnesota, I served as a member of the graduate programs in Pharmaceutics, Neurosciences, and Experimental and Clinical Pharmacology. From 1983 to 1989 and 1991 to 1994 I was the Director of Graduate Studies in Pharmaceutics at the University. From 1998 to 1999 I served as the Head of the Department of Pharmaceutics at the University of Minnesota.

6. Also, from 1982 to 1995, I served as Director of the Clinical Pharmacokinetics Laboratory at the College of Pharmacy at the University of Minnesota.

7. During my career, I received several honors, scholarships and awards, including the Weaver Medal of Honor in 2001, the Meritorious Manuscript Award from the American Association of Pharmaceutical Scientists in 1999 and the Hallie Bruce Memorial Lecture Award in 1996. In 2007, I received the American Pharmacists Association (APhA) Research Achievement Award in the Basic Pharmaceutical Sciences.

8. I am a member of numerous scientific and clinical societies. I am a Fellow of the American Association of Pharmaceutical Scientists and of the American Association for the Advancement of Science. I have been a member of the International Society of Anti-infective Pharmacology and the International Society for the Study of

Xenobiotics (ISSX). I recently served as a member-at-large on the American Association of Pharmaceutical Scientists (AAPS) Executive Council.

9. I have served on the editorial boards of scientific journals such as the Journal of Pharmaceutical Sciences and the Saudi Pharmaceutical Journal. I am currently on the Editorial Board of the AAPS Journal, and on the ISSX Journal, Xenobiotica. I have also served on numerous advisory committees and review panels.

10. I have participated in multiple research projects focused in the areas of preclinical and clinical pharmacokinetics, both publicly and privately funded. I am a named author on over 100 refereed scientific publications, in addition to several book chapters, a book that I co-edited on drug bioavailability, and over 170 abstracts which have been presented at scientific meetings. I have also given hundreds of invited lectures.

11. I have significant experience in the areas of pharmaceutical research, pharmacokinetics, and drug development. Therefore, I believe that I am qualified to render the opinions set forth in this declaration.

12. I have read the Office Action dated September 16, 2011 ("Office Action"), which is attached as **Exhibit 2**. Among other rejections, I understand that the Office Action rejects the claims pending in the captioned application as unpatentable over the following references:

- a. McLeskey et al., "Tamoxifen-resistant fibroblast growth factor-transfected MCF-7 cells are cross-resistant in vivo to the antiestrogen ICI 182,780 and

two aromatase inhibitors”, *Clinical Cancer Research* 4:697-711 (1998) (“*McLeskey*”, attached hereto as **Exhibit 3**);

- b. European Patent Specification No. EP 0 346 014, which names Michael Dukes as inventor (“*Dukes*”, attached hereto as **Exhibit 4**);
- c. Osborne et al., “Comparison of the effects of a pure steroidal antiestrogen with those of tamoxifen in a model of human breast cancer”, *J. National Cancer Institute*, 87(20):746-750 (1995) (“*Osborne*”, attached hereto as **Exhibit 5**); and
- d. the abstract of Wakeling et al., “ICI 182,780, a new antioestrogen with clinical potential”, *J. Steroid Biochemistry & Molecular Biology*, 43(1-3):173-177 (1992) (“*Wakeling*”, attached hereto as **Exhibit 6**);

13. I have read the instant application (“the ‘887 application”), which I believe corresponds to U.S. Application Publication No. US 2010/0152149 (attached hereto as **Exhibit 7**.)

14. I attach hereto **Exhibit 8**, which I believe is a copy of the pending claims in the ‘887 application with proposed amendments. I understand the claims in **Exhibit 8** will be filed in the Patent and Trademark Office as part of the response to the Office Action.

15. I understand that the earliest priority date for the ‘887 application is January 10, 2000. In the paragraphs below, I will refer to the state of the art in the areas of pharmaceutical research, pharmacokinetics, and drug development prior to January 10, 2000. I will also explain how a person of ordinary skill in that art at that time

would have understood the references cited in the Office Action and how such a person would have interpreted certain experimental results related to various fulvestrant formulations.

Disclosure in *McLeskey* regarding the castor oil fulvestrant composition

16. *McLeskey* discloses two fulvestrant compositions. One composition was prepared by dissolving powdered drug in 100% ethanol and then spiking it into warmed peanut oil to give a final concentration of 50 mg/ml (“the *McLeskey* peanut oil composition”). *McLeskey* at 698, col. 2, under “Drugs”. The second composition is a 50 mg/ml fulvestrant composition “in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol, brought to volume with castor oil” (“the *McLeskey* castor oil composition”). *Id.* *McLeskey* does not specify whether the percentages in the castor oil composition are in weight/volume units (% w/v, as recited in the claims of the ’887 application) or in volume/volume units (% v/v).

17. In a liquid composition, when a solute or cosolvent is a liquid, it is often convenient to express its concentration as a volume percent, i.e., % v/v. For the reasons that follow, I believe one of ordinary skill in the art would have concluded the *McLeskey* castor oil composition was described in volume/volume units (% v/v).

18. For example, U.S. Patent No. 3,164,520 (“the ’520 patent”, attached as **Exhibit 9**) entitled “Injectable Steroid Compositions Containing at least 75% Benzyl Benzoate” discloses the preparation of parenteral injections of steroid drugs in formulations containing benzyl benzoate, and often also containing castor oil or sesame oil. See, e.g., the working examples. The ’520 patent states: “The amount of benzyl

benzoate which may be employed in the compositions of this invention while still yielding satisfactory results has been found to range from about 75% to 100% by volume of the pharmaceutical vehicle employed.” The '520 patent at col. 2, ll. 10-14. In addition, each of the four claims of the '520 patent refers to a parenteral steroid formulation in a pharmaceutical vehicle or pharmaceutical carrier wherein at least 75% by volume of said vehicle is benzyl benzoate.

19. Raymond Huber, the named inventor of the '520 patent, is a co-author of a similar publication in which parenteral formulations of steroid hormones in castor oil are described. Riffkin, C., Huber, R., and Keysser, C.H., “Castor oil as a vehicle for parenteral administration of steroid hormones”, *J Pharm Sci*, 53(8): 891-95 (1964) (“*Riffkin*”, attached as **Exhibit 10**). *Riffkin* lists the compositions of various vehicles prepared in Tables IV to VI, which reference liquid components and their proportions in the overall composition in terms of percentage units (“%”). Although *Riffkin* does not specifically state that those compositions are % v/v, one would understand them to be % v/v because *Riffkin* refers to the concentrations of the solid solutes (the steroids) in terms of w/v, (e.g., mg./ml.), whereas the concentrations of the liquid components are simply reported in terms of “%” units. See, e.g., Tables V and VI. One would reasonably assume that, had *Riffkin* intended the concentration of the liquid components to be in terms of % w/v units, *Riffkin* would have explicitly indicated that fact, as it did for the solid components. Footnote 4 is another example of the use of the above nomenclature. Footnote 4 refers to the concentration of estradiol valerate in the injectable formulations, in terms of “mg./ml.”, but refers to a “%” value for the liquids—

castor oil, benzyl benzoate, and benzyl alcohol. Therefore, one would conclude that the composition of the solvents in *Riffkin's* vehicles is expressed as % v/v.

20. Other publications also describe the composition of injectable formulations comprising liquid solvents or co-solvents on a “by volume” basis. For example, a published review tabulates various excipients included in the formulation of injectable products marketed in the United States. Neema, S, Washkuhn, R.J., and Brendel, R.J., “Excipients and their use in injectable products”, *PDA J Pharm Sci Tech*, 51(4):166-171 (1997)(“*Neema*”, attached as **Exhibit 11**). *Neema* lists liquid solvents, co-solvents, and solubilizing agents, and identifies commercial products in which the content of such liquid agents is described on a % v/v basis (e.g., benzyl benzoate, 20% v/v; PEG 40 castor oil, 11.5% v/v; sorbitol, 50% v/v). See, e.g., Tables I and II.

21. Considering the above examples, and because all of the components of the vehicle disclosed in *McLeskey* are liquids, one of ordinary skill in the art would have concluded that the composition was described in terms of volume/volume percent units (% v/v).

22. Therefore, one of ordinary skill in the art would have concluded that the *McLeskey* castor oil composition on page 698 was reported in % v/v units and referred to a composition containing 10% v/v ethanol, 15% v/v benzyl benzoate, and 10% v/v benzyl alcohol in a castor oil vehicle. This composition *is different* from a composition containing 10% w/v ethanol, 15% w/v benzyl benzoate, and 10% w/v benzyl alcohol in a castor oil vehicle.

23. It is possible to convert % v/v values for a given component in a liquid composition into % w/v values by calculating the weight of the corresponding volume of that component in the composition. As a first approximation, the weight of the component can be calculated by multiplying the volume of the component by its density.

24. In order to facilitate this calculation, I assumed the preparation of 100 ml of the *McLeskey* castor oil composition and reported the associated volume and weight values in Table 1 below, using densities reported or calculated at 25°C. The resulting % w/v values are independent of the choice of a particular volume of the *McLeskey* castor oil composition for this calculation. However, a volume of 100 ml of the castor oil composition was selected for simplicity to show the corresponding volumes and weights. The differences between % v/v and % w/v compositions for each of the three components can be seen by comparing the values in Columns A and E. It should be noted that although these compositions are identical, they are described differently; in Column A, the composition is described on a percentage “by volume” (% v/v) basis, and in Column E, the composition is described on a percentage “by weight” (% w/v) basis.

Table 1. Information for 100 ml of the fulvestrant *McLeskey* castor oil composition¹

	A	B	C	D	E
Component	% v/v	Volume (ml)	Density (mg/ml)	Weight (g)	% w/v
Ethanol	10	10	0.808	8.08	8.1
Benzyl Benzoate	15	15	1.118	16.77	16.8
Benzyl Alcohol	10	10	1.04156	10.42	10.4

25. In Table 1, Column A represents the concentration of each component in the *McLeskey* castor oil composition in % v/v units (i.e., as one of ordinary skill in the art in would have understood the *McLeskey* disclosure). Column B represents the volume in milliliters (ml) of each component necessary to prepare 100 ml of the *McLeskey* castor oil composition.

26. Column C represents the density of each component in g/ml at 25°C, reported or calculated from published relative density data from the Merck Index, **Exhibit 12**. The Merck Index reports specific gravity values for liquid substances as the ratio of the density of the substance at a given temperature relative to the density of water at a reference temperature. **Exhibit 12** at p. xiv (entry for “d”). Regarding the benzyl benzoate and benzyl alcohol values, their densities were reported at 25°C and

¹ *McLeskey* does not indicate whether the ethanol used in its castor oil fulvestrant composition is dehydrated ethanol or the binary azeotropic ethanol composition containing about 96% ethanol by volume (see entry no. 3806 for ethanol in the Merck Index, 12th Ed., Merck & Co., Inc. (1996) at pp. 641-642 (“the Merck Index”, relevant copies attached as **Exhibit 8**)). The value in Table 1 for the density of ethanol corresponds to the density of the azeotropic ethanol composition. The density of dehydrated ethanol is 0.789 mg/ml at 20°C (**Exhibit 8**), which would produce an even lower w/v% value for ethanol than that reported in Table 1.

the density of water was reported at 4°C (**Exhibit 12** at entries no. 1159, 1162; pp. 189-190). Because the density of water at 3.98°C is 1.0000 g/ml (**Exhibit 12** at entry 10175; p. 1715), the values reported in the Merck Index for benzyl benzoate and benzyl alcohol were used in Table 1 as the corresponding densities in mg/ml (considering that 3.98°C is 4°C for purposes of this calculation). For ethanol, the Merck Index reports a specific gravity of 0.810 at 25°C with respect to the density of water at 25°C (**Exhibit 12** at entry no. 3806; p. 642). Thus, to obtain the density of ethanol (the binary azeotrope) at 25°C, I multiplied the density of water at 25°C, 0.997 mg/ml (**Exhibit 12** at entry no. 10175; p. 1715), by the specific gravity reported in the Merck Index (0.810) to produce a value of 0.808 mg/ml for the density of ethanol at 25°C.

27. Column D represents the weight of each component, obtained by multiplying the volume of each component (Column B) by its density (Column C). Column E represents the concentration of each component in the *McLeskey* castor oil composition in w/v% units, which is the weight of each component (Column D) in 100 ml of solution (the total volume of the composition) after rounding the value to a single decimal place.

28. Accordingly, based on the values in Table 1, a composition containing 10% v/v ethanol, 15% v/v benzyl benzoate, and 10% v/v benzyl alcohol translates into a composition containing about 8.1% w/v ethanol, about 16.8 % w/v benzyl benzoate, and about 10.4% w/v benzyl alcohol.

29. Thus, one of ordinary skill in the art reading *McLeskey* would have concluded that *McLeskey* described a composition containing about 8.1% w/v ethanol,

about 16.8 % w/v benzyl benzoate, and about 10.4% w/v benzyl alcohol in a castor oil vehicle.

30. Neither *McLeskey* nor any of the references cited in the Office Action contain any disclosure that would have suggested to one of ordinary skill in the art the modification of a composition containing about 8.1% w/v ethanol, about 16.8 % w/v benzyl benzoate, and about 10.4% w/v benzyl alcohol (i.e., the *McLeskey* castor oil composition) in an attempt to produce a composition as recited in the claims containing about 10% w/v ethanol, about 15% w/v benzyl benzoate, and about 10% w/v benzyl alcohol.

Disclosure in *McLeskey* regarding administration of fulvestrant compositions

31. As mentioned above, *McLeskey* disclosed two different fulvestrant compositions, the peanut oil composition and the castor oil composition. *McLeskey* at 698. *McLeskey*, however, did not provide any experimental data that would have allowed one of ordinary skill in the art to compare any aspect of the performance of the two fulvestrant compositions for the treatment of cancerous tumors. Therefore, *McLeskey* provided no information that would have suggested to one of ordinary skill in the art the desirability of either of its fulvestrant compositions over other known fulvestrant formulations.

32. *McLeskey* did not disclose plasma or blood levels of fulvestrant in mice after subcutaneous administration of either the peanut oil or the castor oil compositions. Thus, no information regarding the rate and/or extent of absorption of fulvestrant from

the subcutaneous injection site is available to one of ordinary skill in the art for either composition.

33. *McLeskey* concluded that treatment with fulvestrant (ICI 182,780), using either of the disclosed compositions was not effective in that it “did not slow estrogen-independent growth or prevent metastasis of tumors produced by FGF-transfected MCF-7 cells in ovariectomized nude mice.” *McLeskey* at Abstract. Thus, one of ordinary skill in the art would not have been informed about the usefulness of either fulvestrant formulation when administered subcutaneously to a mouse for the treatment of cancerous tumors.

34. *McLeskey* also reports that fulvestrant “retained activity” based on the results from injecting fulvestrant into “reproductively intact female mice for two weeks . . . at the same doses used in the above experiment” and the uteri subsequently harvested from those mice “weighed less than those from control mice and exhibited a complete lack of endometrial glandular structures (data not shown).” *Id.* at ¶¶ bridging 701-702. Unfortunately, *McLeskey* does not specify which of the two fulvestrant formulations, if any, (the peanut oil composition or the castor oil composition), was used in these experiments. *McLeskey* does not disclose the route of administration (subcutaneous, intramuscular, intraperitoneal, etc.) for the injection of fulvestrant into those “reproductively intact female mice.” Thus, one of ordinary skill in the art reading *McLeskey* cannot draw any conclusions regarding the extent to which fulvestrant administered subcutaneously became absorbed, if at all, when using the peanut oil or the castor oil compositions.

35. Indeed, because of the lack of fulvestrant efficacy and the absence of pharmacokinetic data in *McLeskey*, one of ordinary skill in the art would have been unable to conclude whether either of the two fulvestrant *McLeskey* compositions (peanut oil or castor oil) was able to deliver a dose of fulvestrant that had an antitumor therapeutic effect in the mice when administered subcutaneously, nor any insight about fulvestrant absorption characteristics (rate and extent) when administered via the *intramuscular route* in any species, including humans.

36. Thus, *McLeskey* provides no information that would have led one of ordinary skill in the art to have a preference for either the peanut oil or the castor oil fulvestrant compositions over the other one, or even a preference for one of the two *McLeskey* fulvestrant compositions over other fulvestrant compositions known in the art prior to January 10, 2000.

37. While I have not performed a search for fulvestrant compositions known in the art prior to January 10, 2000, I note that some of the references cited by the Examiner in the Office Action do disclose other fulvestrant compositions. For example, *Osborne* discloses experiments in which a composition of fulvestrant “in castor oil” was injected subcutaneously to female nude mice. *Osborne* (**Exhibit 5**) at 747, col. 1. Based on the positive results of those experiments, *Osborne* concludes that fulvestrant “is a more effective estrogen antagonist than tamoxifen in the MCF-7 tumor cell/nude mouse model system.” *Osborne* at Abstract.

38. The fulvestrant composition in *Wakeling* is described as having fulvestrant “in oil suspension” for parenteral administration to mice. *Wakeling* (**Exhibit 6**) at

Abstract. *Wakeling* reports that, “over a 1 month period, a single injection of [fulvestrant] in oil suspension achieved effects comparable with those of daily tamoxifen treatment.” *Id.*

39. *Dukes* discloses two different fulvestrant compositions for intramuscular injection, one containing fulvestrant dissolved “in a mixture of propylene glycol: ethanol: water: poloxamer 407” administered daily by intramuscular injection to rats. *Dukes* (**Exhibit 4**) at Example 2, p. 8. The second composition contained 50 mg of fulvestrant, “400 mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml.” *Id.* at Example 3, p. 9. For each composition, *Dukes* reports that “at all doses tested the compound [fulvestrant] selectively inhibits the action of the animals’ endogenous oestrogen on their uteri.” *Id.* at Examples 2 & 3, pp. 8-9.

40. Thus, it is clear that one of ordinary skill in the art had other choices besides the *McLeskey* castor oil composition with respect to potential fulvestrant formulations that could have been further investigated for the development of a method of treating humans with intramuscular fulvestrant. However, none of the references cited in the Office Action provides any information that would have guided one of ordinary skill in the art to select the *McLeskey* castor oil composition, over any of the other fulvestrant compositions mentioned above, for the potential development of such a method of treatment.

41. Moreover, none of the references cited by the Examiner provides any guidance as to the relevant factors to consider when selecting a formulation for the potential development of a method of treatment as recited in the instant claims.

However, judging solely on the basis of efficacy, the *McLeskey* castor oil composition would have been among the least favored compositions to select for further development from the fulvestrant compositions discussed above because the *McLeskey* experiments were ineffective and one of ordinary skill in the art would not have been able to conclude from the information in *McLeskey* whether fulvestrant, using that composition, was sufficiently bioavailable to have an antitumor effect. In this regard, and considering only efficacy, the fulvestrant oil suspension from *Wakeling* would have been among the most favored formulations to select for further development from among those discussed above because at least that formulation, when given as a single injection, showed a therapeutic antitumor effect in mice for over a one-month period.

Lack of disclosure in *McLeskey* regarding intramuscular efficacy of either fulvestrant composition disclosed therein

42. The mode of administration of a drug (e.g., oral, intramuscular, subcutaneous, etc.) and the dose administered affects the release profile of the drug. One of ordinary skill in the art would have understood that results from subcutaneous administration in general, and including those reported in *McLeskey*, cannot be extrapolated to intramuscular administration. As a result, one of ordinary skill in the art would not have had an expectation as to whether the *McLeskey* castor oil composition would have had a therapeutic effect when administered intramuscularly before actually performing suitable *in vivo* experiments.

43. There is abundant evidence in the scientific literature that the intramuscular and subcutaneous administration of a drug to the same animal or human

may produce very different plasma level curves, and therefore very different pharmacologic effects. These effects include the desired effects (efficacy) and those that are not desired (adverse events, or side effects). If a drug is poorly absorbed from the injected site, (e.g., too slowly, or to only a modest extent) the drug may show no effects whatsoever.

44. For example, a study in sheep using probenecid, a drug which may be used to prolong the half-life of some antibiotics in animals, demonstrates significant differences in the absorption of intramuscular and subcutaneous injections of probenecid. Guerrini V.H., Filippich L.J., English P.B., Schneider J., Cao G.R. and Bourne D.W.A., "Pharmacokinetics of probenecid in sheep", *J Vet Pharmacol Ther.* 8(2):128-35 (1985) ("*Guerrini*", attached as **Exhibit 13**).

45. Those investigators administered probenecid to ewes in doses of 1 gram by both intramuscular and subcutaneous injection. *Guerrini* at 129. The study shows that the absorption of probenecid is more rapid and complete following intramuscular injection, compared to subcutaneous injection. *Id.* at Abstract. *Guerrini* reports that the bioavailability of the intramuscular dose was 135% of that of the subcutaneous dose (corresponding to an average bioavailability of 46% for intramuscular injection compared with an average bioavailability of 34% for subcutaneous injection). *Id.* The subcutaneous dose was also absorbed more slowly, with average plasma levels of the drug peaking at 1.5 hr, compared to 0.67 hr for the intramuscular dose. *Id.* at 131. Because of this slower absorption following subcutaneous dosing, probenecid plasma concentrations remained higher after 2 hours when the drug was administered

subcutaneously than when it was administered intramuscularly. *Id.* at 135. Consistent with these observations, the rate constant for absorption for the intramuscular dose was 41% greater than for the subcutaneous dose (5.45 vs. 3.87 hr⁻¹). *Id.* at 133.

46. In this case, despite the overall higher bioavailability of intramuscular probenecid, the “higher and more prolonged plasma probenecid concentration” following subcutaneous administration resulted in “similar plasma concentrations to those found in man after oral administration.” *Id.* at 135. *Guerrini* concludes that “[t]he s.c. [subcutaneous] administration of probenecid in animals is preferred [to intramuscular administration] because muscle damage is avoided and it provided useful plasma concentrations.” *Id.* Thus, this is an example where subcutaneous administration achieves a certain desired result but where intramuscular administration does not accomplish the same result.

47. Another study shows that, contrary to the pharmacokinetic profiles observed in *Guerrini*, subcutaneous administration resulted in faster absorption compared to intramuscular injection. Lavy E, Ziv G, Shem-Tov M, Glickman A, Dey A., “Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs”, *J Vet Pharmacol Ther.* 22(4):261-5 (1999) (“Lavy”, attached as **Exhibit 14**).

48. *Lavy* reports that when a 10 mg/kg dose of clindamycin HCl, an antibiotic, was given subcutaneously to dogs, the average maximum blood serum concentration (C_{max}) of clindamycin was 20.8 µg/ml, and the time when this maximum occurred (T_{max}) averaged 46.7 min. *Lavy* at Table 3. When the same dose was given

intramuscularly to the same animals, the corresponding values for C_{max} and T_{max} were 4.4 µg/ml and 73 min, exhibiting a very much slower rate of absorption. *Id.* In addition, the exposure of the dogs to clindamycin, assessed through an analysis of the plasma serum area under the curve (AUC) was 2.9 times greater for the subcutaneous dose than for the intramuscular dose. *Id.* This means that the bioavailability of the subcutaneous dose of this drug is 2.9 times that of the intramuscular dose.

49. Based on the differences in pharmacokinetic profiles for subcutaneous and intramuscular administration, *Lavy* concludes that “it appears from the present study that the s.c. [subcutaneous] route is superior to the i.m. [intramuscular] in practical terms by permitting a longer treatment interval.” *Id.* at 265. This is another example in which subcutaneous administration is able to fulfill certain design criteria (maintain a therapeutic plasma concentration for a longer period of time) better than intramuscular administration. Therefore, under these circumstances, one of ordinary skill in the art would not have been able to rely on data from subcutaneous administration to predict results of intramuscular administration because intramuscular administration would not have produced the same level of long-term efficacy achieved by subcutaneous administration.

50. There are other reports in the literature that show that, in contrast to the results from *Lavy*, the absorption of a drug is more rapid and complete following intramuscular dosing than after subcutaneous injection. For example, when the fluoroquinolone antimicrobial agent difloxacin was given by these routes to the same calves in a crossover study, the rates of absorption differed greatly, with intramuscular

injection showing higher and earlier peak plasma concentrations, confirming much more rapid absorption. Ismail M., "Disposition kinetics of difloxacin after intravenous, intramuscular and subcutaneous administration in calves", *Vet Res Commun.*, 31(4):467-76 (2007) ("*Ismail*", attached as **Exhibit 15**).

51. After intramuscular and subcutaneous dosing, maximum plasma concentrations (C_{max}) of 3.38 and 2.18 µg/ml were observed after (T_{max}) 1.22 and 3.7 hr, respectively. *Ismail* at Abstract. The time for half of the dose to be absorbed when given by intramuscular injection was only 0.38 hr, whereas the corresponding time for absorption of the subcutaneously injected dose was 2.1 hr, over 5 times as long. *Id.* at 473.

52. Under the conditions of its study, *Ismail* concludes that "the doses of difloxacin used in this study are likely to involve better pharmacodynamic characteristics that are associated with greater clinical efficacy following i.m. [intramuscular] administration than following s.c. [subcutaneous] administration." *Id.* at Abstract. In this case, contrary to the two examples above, the intramuscular administration was considered to be associated with greater clinical efficacy.

53. These three examples above show that there are significant differences in the rate and extent of absorption of a drug given by the intramuscular and subcutaneous route, even when given to the same animals in a crossover study. As a result, it cannot be predicted a priori whether intramuscular or subcutaneous dosing will result in more rapid and/or complete drug absorption, as examples of both cases are found in the scientific literature.

54. Moreover, the examples above underscore the fact that efficacy of a given drug administered by a given route of dosing (e.g., intramuscular) cannot be known until appropriate comparative studies are performed in a suitable animal model. For some drugs, the desired effect might be achieved following a particular route of dosing, but for other drugs it might not. The rate and extent of drug absorption, and the associated pharmacodynamics (e.g., the achievement of a desired drug effect) may differ greatly depending on the properties of the drug, the choice of an animal model, and the site of drug administration.

55. Consequently, one of ordinary skill in the art having the very limited experimental subcutaneous data from *McLeskey* would not have had an expectation that the intramuscular administration of fulvestrant using the *McLeskey* castor oil composition would have been effective following intramuscular administration, such as in the method described in the claims. This is especially true because *McLeskey* did not disclose plasma or blood levels of fulvestrant in mice after subcutaneous administration of the formulation, nor any information regarding the rate and/or extent of absorption of fulvestrant from the subcutaneous injection site. Additionally, the claims recite achieving a given therapeutic plasma concentration for at least four weeks, and there is no information in any of the references cited in the Office Action that would have suggested that such long-term efficacy associated with a single dose would be exhibited using the *McLeskey* castor oil composition by any route of administration.

56. Thus, one of ordinary skill in the art would not have had an expectation that the castor oil composition disclosed in *McLeskey*, which was administered

subcutaneously to mice, would have been therapeutically effective upon intramuscular administration of fulvestrant, for example, by following the method described in the claims.

The composition of a formulation can have a significant effect on the efficacy observed when the formulation is administered

57. Where a dosage form of a drug is being developed for intramuscular administration in humans, one of ordinary skill in the art typically relies upon the results of intramuscular dosing studies in suitable animal models where pharmacokinetic data are collected to characterize the absorption of the drug from its dosage form.

58. Typically, during the development of an intramuscular dosage form for administration of a drug in humans, one would have carried out, among other tasks, formulation studies to determine suitable compositions in which the drug of interest is dissolved, as well as initial intramuscular dosing experiments in animals (e.g., mice, rabbits, and/or dogs) under various conditions (e.g., different compositions, different solvents, varying the proportion of the components of the composition, different drug concentrations, etc.) in order to gain an understanding of the pharmacokinetics of fulvestrant before attempting human administration. The very existence of this generalized approach highlights the lack of expectation of success with respect to the extrapolation of the *McLeskey* disclosure of subcutaneous administration to mice, lacking any pharmacokinetic information, to human intramuscular administration.

59. With respect to the importance of formulation studies, I have read the Declaration Under 35 U.S.C §1.132 of Dr. Paul Gellert filed on August 2008 (“the Gellert

Declaration”, cited as “Gellert Decl.” and enclosed here as **Exhibit 16**). I understand that the Gellert Declaration was submitted to the U.S. Patent and Trademark Office in Application No. 10/872,784 (as indicated by the caption on the first page of the declaration).

60. As part of the discussion of the development of methods of treatment involving the administration of fulvestrant, the Gellert Declaration states that “the experienced formulator would want to minimize the amount of co-solvents and excipients in any injectable formulation.” Gellert Decl. at ¶ 22.

61. Thus, even if the *McLeskey* castor oil composition had been considered as a potentially useful formulation in the development of a method of treatment for humans, one of ordinary skill in the art would have performed additional formulation studies to obtain a composition with suitable characteristics for the desired route of administration. The Gellert Declaration explains one of the rationales to perform those additional studies:

Ideally, it is best to select and use solvents that would maximize the solubility of the compound. Maximizing the solubility of a compound in a particular cosolvent system would result in lower total levels of the non-aqueous solvent(s) being administered to the patient, thereby lowering the chance for potential side effects.

Gellert Decl. at ¶ 22 (quoting directly from P.K. Gupta and G.A. Brazeau (eds), “Injectable Drug Development: Techniques to Reduce Pain and Irritation” Chapter 11, p. 217, Interpharm Press, Denver, Colorado (1999)).

62. Regardless of how high or low the cosolvent concentrations are in a given formulation, the preparation of formulations in which a drug such as fulvestrant can be

solubilized is not sufficient to ensure the desired therapeutic effect when such formulation is administered to patients. As explained in the '887 application “[s]imply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.”

Exhibit 7 at ¶ [0054]. Thus, suitable experiments are needed to determine the pharmacokinetic performance of any candidate formulation(s).

63. In that regard, it is understood that an animal model for drug dosage form performance may provide some discrimination among candidate dosage forms in development. Thus, the plasma concentration profile should reflect changes in the release characteristics of the drug from the formulation. That type of pharmacokinetic data could be used to characterize important variables in the development of a suitable method of treatment. For drugs that are difficult to formulate, such as fulvestrant (**Exhibit 7**, at ¶ [0014]), the pharmacokinetic data could be useful to investigate the most promising formulation for the desired route of administration.

64. For example, for fulvestrant, PCT Application Publication No. WO 03/006064 (“WO 03/006064”, attached here as **Exhibit 17**) shows pharmacokinetic results of intramuscular administration of fulvestrant to rabbits. Figure 1 shows differences in results when seven different formulations of fulvestrant, each containing 100 mg/ml of the drug, but having different co-solvent compositions, were dosed intramuscularly in rabbits. The table related to Example 4 on page 30 of WO 03/006064 reports the composition of each formulation, labeled F1 to F7. As can be seen, all of these fulvestrant formulations contained ethanol, benzyl alcohol, and benzyl benzoate in

a castor oil vehicle; these are the same components of the fulvestrant composition recited in the claims, but with different proportions for each component.²

65. WO 03/006064 reports that “[p]lasma levels were more variable than Control over the first 30 days” following intramuscular administration of fulvestrant. WO 03/006064 at 30, I. 23. WO 03/006064 explains that “some differences in profiles were noted over the first 30 days such that they were divided into 2 groups (with Formulation F7 showing intermediate behaviour).” *Id.* at 30, II. 29-30. According to WO 03/006064, Group A demonstrates “rapid release early time points”, corresponding to formulations containing high benzyl benzoate and low castor oil concentrations, while Group B shows a “lower release, flatter profile” corresponding to formulations containing lower benzyl benzoate and higher castor oil concentrations. *Id.* at 30, II. 31-34. WO 03/006064 replotted the data from Figure 1 corresponding to those formulations in Group A as part of Figure 2A and the data corresponding to those formulations in Group B as part of Figure 2B.

66. Therefore, based on WO 03/006064’s own characterization of the differences in the pharmacokinetic profile of different fulvestrant formulations, higher benzyl benzoate concentrations in the formulation resulted in a more rapid initial release of fulvestrant, whereas lower benzyl benzoate concentrations resulted in a lower initial release, and a flatter plasma level profile. Depending on the overall objective of the administration of fulvestrant, some of the fulvestrant formulations tested in

² The right-hand column in this table appears to indicate the % w/v composition of castor oil. All the entries in this column should more properly be “to 100%”, as they are in the Tables provided in the preceding Examples 2 and 3.

WO 03/006064's study would be more desirable than others for that given purpose and, based on the relevant pharmacokinetic profiles, one of ordinary skill in the art would be able to select one of those fulvestrant formulations for further development and/or testing.

67. However, one of ordinary skill in the art would not have been able to determine whether a given fulvestrant formulation injected intramuscularly as in WO 03/006064 would have had the desired pharmacokinetic profile until such *in vivo* pharmacokinetic studies were carried out. The testing of various formulations having different compositions, as portrayed in Figures 1, 2A and 2B, would typically be undertaken during the development of a dosage form in order to ensure an optimal method of treatment using a drug that is difficult to formulate. Such studies would be expected to demonstrate differences in the blood plasma concentrations of a test drug, and would allow the investigators to identify factors that would enhance the performance of the formulation.

68. Therefore, when considering the differences in pharmacokinetic profiles demonstrated in the example from WO 03/006064, it becomes clear that one of ordinary skill in the art knowing only the composition of a given formulation administered subcutaneously, but having no pharmacokinetic data following its intramuscular administration, would have had no expectation, one way or another, that the formulation would be effective when administered intramuscularly in a given method of treatment.

69. In particular, one of ordinary skill in the art would not have had a reasonable expectation that the *McLeskey* castor oil composition would have been

effective when given as an intramuscular injection, such as in the method of treatment recited in the claims.

70. I hereby declare that all the statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements are made with the knowledge that willful false statements so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

January 13, 2012
Date.

Ronald J. Sawchuk
Ronald J. Sawchuk

Exhibit List for Declaration Under 37 C.F.R. § 1.132 of Ronald J. Sawchuk

Exhibit No.	Description
1	Curriculum Vitae of Ronald J. Sawchuk
2	Office Action for U.S. Patent Application No. 12/285,887 dated September 16, 2011
3	McLeskey et al., "Tamoxifen-resistant fibroblast growth factor-transfected MCF-7 cells are cross-resistant in vivo to the antiestrogen ICI 182,780 and two aromatase inhibitors", <i>Clinical Cancer Research</i> 4:697-711 (1998) ("McLeskey")
4	European Patent Specification No. EP 0 346 014, naming Michael Dukes as inventor ("Dukes")
5	Osborne et al., "Comparison of the effects of a pure steroidal antiestrogen with those of tamoxifen in a model of human breast cancer", <i>J. National Cancer Institute</i> , 87(20):746-750 (1995) ("Osborne")
6	Abstract for Wakeling et al., "ICI 182,780, a new antioestrogen with clinical potential", <i>J. Steroid Biochemistry & Molecular Biology</i> , 43(1-3):173-177 (1992) ("Wakeling")
7	U.S. Patent Publication No. 2010/0152149
8	Pending claims in U.S. Application No. 12/285,887, with proposed amendments
9	U.S. Patent No. 3,164,520
10	Riffkin et al., "Castor oil as a vehicle for parental administration of steroid hormones", <i>J. Pharma Sci.</i> , 53(8):891-895 (1964) ("Riffkin")
11	Nema et al., "Excipients and their use in injectable products", <i>PDA J Pharma Sci Tech.</i> , 51(4):166-171 (1977) ("Nema")
12	The Merck Index, 12th Ed., Merck & Co., Inc. (1996) ("the Merck Index")
13	Guerrini et al., "Pharmacokinetics of probenecid in sheep", <i>J Vet Pharmacol Ther.</i> , 8:128-135 (1985) ("Guerrini")
14	Lavy, et al., "Pharmacokinetics of clindamycin HC1 administered intravenously, intramuscularly and subcutaneously to dogs:", <i>J Vet Pharmacol Ther.</i> , 22(4):261-265 (1999) ("Lavy")
15	Ismail, "Disposition kinetics of difloxacin after intravenous, intramuscular and subcutaneous administration in calves", <i>Vet Res Commun.</i> , 31(4):467-476 (2007) ("Ismail")
16	Declaration Under 35 U.S.C §1.132 of Dr. Paul Gellert filed on August 2008 in U.S. Application No. 10/872,784 ("the Gellert Declaration")
17	PCT Application Publication No. WO 03/006064

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known			
				Application Number		12/285,887	
				Filing Date		October 15,2008	
				First Named Inventor		John R. EVANS	
				Art Unit		1628	
				Examiner Name		HUI, San Ming R.	
Sheet	1	of	1	Attorney Docket Number		11285.0056-00000	

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS						
Examiner Initials	Cite No. ¹	Document Number		Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)				

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS							
Examiner Initials	Cite No. ¹	Foreign Patent Document		Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁶
		Country Code ³	Number ⁴ Kind Code ⁵ (if known)				
	1	WO	03/006064	23-JAN-2003	Astrazeneca AB		

NONPATENT LITERATURE DOCUMENTS			
Examiner Initials	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation ⁶
	2	The Merck Index, 12th Ed., Merck & Co., Inc., pgs. xiv, 189-190, 641-642 and 1715 (1996).	
	3	Guerrini, et al., "Pharmacokinetics of probenecid in sheep", J Vet Pharmacol Ther., 128-135 (1985).	
	4	Lavy, et al., "Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs", J Vet Pharmacol Ther., 22(4):261-265 (1999).	
	5	Ismail, "Disposition kinetics of difloxacin after intravenous, intramuscular and subcutaneous administration in calves", Vet Res Commun., 31(4):467-476 (2007).	
	6	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) from August 27, 2009 to December 15, 2011.	
	7	Documents from the prosecution of European Application No. 10180667.7 (EP 2 266 573) from November 23, 2010 to December 19, 2011.	
	8	Documents from the prosecution of European Application No. 10180661.0 (EP 2 286 818) from January 19, 2011 to December 19, 2011.	
	9	Declaration Under 35 U.S.C §1.132 of Dr. Paul Gellert filed in August 2008 in U.S. Application No. 10/872,784.	
Examiner Signature			Date Considered

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 2093
)	
EVANS et al.)	
)	
Application No.: 10/872,784)	Group Art Unit: 1617
)	
Filed: June 22, 2004)	Examiner: Hui, San-Ming R
)	
FOR: FORMULATION)	
)	

DECLARATION UNDER 35 U.S.C. § 1.132
OF PAUL RICHARD GELLERT

PAUL RICHARD GELLERT of AstraZeneca, Alderley Park, Macclesfield, Cheshire, UK
declares:

1. I graduated from the University of Oxford in Chemistry in 1984. I undertook postgraduate research with Professor Brian Howard in the Physical Chemistry Laboratory at the University of Oxford leading to the award of a D.Phil in 1988. From February 1988 until the present I have been employed by AstraZeneca, (formerly Zeneca and ICI) initially as a Senior Research Scientist and subsequently as a Team Leader/Manager, Principal Scientist and, since 2004, a Senior Principal Scientist.
2. I have worked in the formulation and drug delivery area throughout my career with AstraZeneca, where my research and development work has covered a range of formulation types including sustained released injections, including fulvestrant.
3. During the course of my study of the subject application (hereinafter "the Evans Application") and the underlying data, I have become aware of several transcription or other errors between certain disclosures of the subject application and the underlying laboratory notebook data. One purpose of this Declaration is to point out the existence

and nature of these errors and to report further testing that has been carried out under my guidance to obtain additional data (paragraphs 4-10 below and Attachments A-D). A further purpose of this Declaration is to set out and document the manner in which an experienced formulator would likely have approached the task of developing a sustained release injectable formulation suitable for human use for a steroidal compound such as fulvestrant in about early 2000, which I understand is when the priority applications supporting the Evans Application were filed (paragraphs 11 - 25 below and Attachment E). Citations to literature and patent references in this Declaration will be in the format Lead Author (Date), and the full citations are given in the Table of References at the end of this Declaration. A copy of each cited reference (or cited portions of the longer references) is included in Attachment F under the Tab number noted in the Table of References.

4. In Table 2 of the Evans Application, the solubility of fulvestrant in castor oil appears to have been transcribed incorrectly from the original source, the laboratory notebook. The value in the latter is 24.5 mg/ml and not 20 mg/ml. In other experiments to determine the solubility of fulvestrant in castor oil and also in benzyl benzoate, some variability was observed.
5. In Table 3 of the Evans Application, the given solubility values were generated at 4°C and not at 25°C as is stated in the title of Table 3. For fulvestrant formulations, it is preferable that the fulvestrant remains completely in solution at both 4°C and 25°C. The 4°C temperature corresponds to the storage temperature (2°C to 8°C in the FDA approved label for Faslodex), and the 25°C temperature corresponds to the administration temperature (ambient temperature). In addition, the specified solubility values on this Table 3 are mean values calculated from analysis of replicate samples from one or more trials. The individual values are shown in handwriting in the amended version of Table 3 in Attachment A. In addition, it appears that the mean values for the last three compositions have been incorrectly calculated. The corrected mean values, together with the correction of the temperature from "25°C" to read "4°C", are also shown in handwriting in the amended version of Table 3 in Attachment A.

6. I have evaluated the transcription and other errors against the original application disclosures and conclude that these do not change the ultimate conclusions made from the data as originally reported. The addition of 15% w/v benzyl benzoate to compositions having total alcohol concentrations in castor oil of 10%, 15%, 20% and 30% w/v unexpectedly provides a positive effect on fulvestrant solubility, significantly increasing the solubility of fulvestrant in the compositions despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.
7. An additional set of experiments has been conducted at 25°C under my guidance to obtain consistent data with reduced variability from a single set of rigorously controlled solubility experiments and to demonstrate that the unexpected increase of solubility of fulvestrant by adding benzyl benzoate into compositions containing ethanol, benzyl alcohol and castor oil, is present across the broader range of composition encompassed by the claims being presented with this Declaration. The solubility of fulvestrant in benzyl benzoate and in castor oil was also measured in the same set of experiments using the same batch of benzyl benzoate and the same batch of castor oil as were used to make up the compositions. The Experimental Test Procedure is described in Attachment B.
8. The results from these solubility experiments are shown in the table in Attachment C. These results show that the solubility of fulvestrant in castor oil alone (21.4 mg/ml) is significantly greater than the solubility of fulvestrant in benzyl benzoate alone (3.8 mg/ml) and demonstrate the unexpected increase in fulvestrant solubility on the addition of 10, 15 and 25% w/v benzyl benzoate, in place of an equivalent amount of castor oil, to compositions having total alcohol concentrations in castor oil of 10%, 15%, 20%, 25% and 30% w/v.
9. Thus, the results that were obtained from experiments conducted under rigorously controlled conditions and with an expanded range of compositions, as shown in Attachment C, confirm the ultimate conclusions drawn from the results shown in Table 3 of the original application disclosure, namely that the addition of 10% to 25% w/v benzyl

benzoate to compositions having total alcohol concentrations in castor oil of between 10% to 30% w/v unexpectedly provides a positive effect on fulvestrant solubility, significantly increasing the solubility of fulvestrant in the compositions despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

10. During the course of my study of the Evans Application and the underlying source materials it was drawn to my attention that some of the composition data given for Delestrogen and Delalutin somehow had been shifted one column to the right. Thus, for Delestrogen, the 78% and 58% figures shown under the BzBz column should have been under the OIL column; the 20% and 40% figures shown under the BzOH column should have been under the BzBz column; and the 2% figures shown under EtOH should have been under the BzOH column. Similarly for Delalutin, the “up to 2%” shown under the EtOH column should have been under the BzOH column. This table reports that the source of this data was J.Pharm.Sci (1964) 53(8) 891, which is Riffkin (1964) elsewhere referred to in this Declaration, and I have also verified the corrected data from the entries for Delalutin and Delestrogen in PDR (1973). A copy of Table 1 from the Evans Application is reproduced as Attachment D, on which these corrections have been made in handwriting, and I have additionally more correctly noted that Delalutin is 17-hydroxy progesterone *caproate*, and that the “COMP” designation for Delalutin should be “BMS” (Bristol-Myers Squibb). Attachment D also includes a one page explanation of the corrections to this Table 1.

11. In about early 2000, a person responsible for developing a sustained release injectable formulation suitable for administration to humans for a new steroidal compound such as fulvestrant, would have had specialized training and experience in developing pharmaceutical formulations and methods for their administration. In developing such a formulation for fulvestrant, the objective would have been to formulate an intramuscular (IM) injection that would provide for the satisfactory sustained release of fulvestrant over a period of at least two weeks and preferably over a period of at least four weeks to reduce the frequency of administration, and would have a target fulvestrant content of at

least 45 mg/mL so as to provide a fulvestrant dose of at least 250 mg in a single 5-6 mL injection. From my personal experience and knowledge of the literature at about that time, I believe that such an experienced formulator would likely have approached the task of developing a formulation for fulvestrant in about the following manner.

12. Given the foregoing objective, the experienced formulator would have appreciated that the traditional administration options to explore were intramuscular (IM) injection of a sustained release aqueous or oil suspension or an oil-based solution (depot) containing at least 250 mg of fulvestrant in a volume of vehicle that is tolerable for injection, *i.e.*, no more than 5 or 6 mL.
13. Because of the extremely low solubility of fulvestrant in water, a reasonable starting point would have been to investigate intramuscular injection of an aqueous or oil suspension of fulvestrant. However, the formulator would have found that injection of an aqueous suspension of fulvestrant resulted in extensive local tissue irritation at the injection site as well as a poor release profile, such as reported in paragraph [0042] of the Evans Application. Since suspensions thus were not an acceptable option for fulvestrant, the experienced formulator would have moved on to further explore whether 250 mg of fulvestrant could be solubilised in no more than 5-6 mL of an oil-based vehicle, *i.e.*, to achieve the target fulvestrant concentration of at least 45 mg/mL.
14. In the preformulation phase, the experienced formulator would have conducted a literature review or otherwise would have become familiar with commercially marketed injectable formulations, particularly injectable sustained release formulations of steroids or other relatively insoluble compounds such as those listed in Table I of the Evans Application, with the objective of identifying potential oil vehicles, co-solvents and other excipients that already had been found to be tolerated and/or to have passed through regulatory review, and which might be candidates for further consideration and testing for the fulvestrant formulation. This review also would have provided guidance with respect to concentration levels of such co-solvents and other excipients that generally had been found acceptable in sustained release oil-based intramuscular injections administered to

humans. This objective is confirmed, for example, in Nema (1997) at page 166:

Generally, a knowledge of which excipients have been deemed safe by the FDA or are already present in a marketed product provides increased assurance to the formulator that these excipients will probably be safe for their new drug product. ... Regulatory bodies may view an excipient previously approved in an injectable dosage form favorably, and will frequently require less safety data.

The purpose of this Nema paper was thus “to present the various excipients that have been included in the formulation of injectable products marketed in the USA.”¹ Similar objectives were intended to be served by the compilations of commercial formulations in Strickley I (1999), Strickley II (2000) and Strickley III (2000):

This compilation will also be useful for those interested in knowing what additives are currently used in injectable products and at what concentrations they are administered in practice. This compilation only focuses on marketed formulations and does not delve into the subject of preclinical or drug discovery formulations associated with early-stages pharmacokinetics or proof-of-concept pharmacodynamics, where the formulation scientist is not bound by regulatory constraints.

(Strickley I (1999) at 324).

Powell (1998) similarly states at page 238 with respect to its compilation of commercially used excipients:

Thus, the formulation scientist is often faced with a dilemma --- which excipients are truly available for use (based on what has been used previously), and which are not? ... And at what concentrations, and by what route? ...

Herein are listed the excipients found in most of the approved and marketed parenteral formulations, given systematically by excipient name. In this format it is easy to determine what concentrations were used, the route of administration, the main rationale for addition of that excipient, the drug that was formulated, the manufacturer, brand name, etc.

15. From the literature review, the formulator would have noted reference to a number of intramuscular injectable sustained release oil-based steroidal formulations that had been

¹ Nema (1997) does caution, however, that there is no guarantee that the new drug product will be safe as excipients are combined with other additives and/or with a new drug, creating unforeseen potentiation or synergistic toxic effects.

commercially marketed:

- Strickley I (1999), Table VII:
 - Haloperidol Decanoate/Haldol decanoate (50-100 mg/mL in sesame oil, benzyl alcohol 1.2%);
 - Testosterone Enanthate/Delatestryl (200 mg/mL in sesame oil, chlorobutanol 5 mg/mL);
- PDR (1973) at pages 1277-1278
 - Proluton/progesterone (50 mg/mL in sesame oil, 150 mg/ml benzyl benzoate, 5 mg/ml benzyl alcohol, 1 mg/ml propylparaben);
- PDR (1973) at pages 1349-1354
 - Deladumone/Testosterone Enanthate & Estradiol Valerate (90 & 4 mg/mL in sesame oil, 0.5% chlorobutanol);
 - Deladumone OB/Testosterone Enanthate & Estradiol Valerate (180 & 8 mg/mL in sesame oil, 2% benzyl alcohol);
 - Delalutin/hydroxyprogesterone caproate (250 mg/mL in 52% castor oil, 46% benzyl benzoate, 2% benzyl alcohol);
 - Delestrogen/estradiol valerate (20 mg/mL in 78% castor oil, 20% benzyl benzoate, 2% benzyl alcohol and 40 mg/mL in 58% castor oil, 40% benzyl benzoate, 2% benzyl alcohol);
 - Delatestryl/Testosterone Enanthate (200 mg/mL in sesame oil, 0.5% chlorobutanol);
 - Delalutecval 2X/hydroxyprogesterone caproate & estradiol valerate (250 mg/mL & 5 mg/mL in castor oil, 45% benzyl benzoate, 1.6% benzyl alcohol);
- PDR (1973) at pages 1391-1392
 - Prolixin Enanthate/FluphenazineEnanthate (25 mg/mL in sesame oil, 1.5% benzyl alcohol);
- Wang (1980):
 - Depo-Testosterone/testosterone cypionate (100 mg/mL in 87.4% cottonseed oil, 0.1 mL benzyl benzoate, 9.45 mg benzyl alcohol as a preservative);
- Mackey (1995):
 - Testoviron Depot/testosterone enanthate (250 mg/mL in castor oil and benzyl

benzoate);

as well as a number of other commercialized oil based long-acting IM injectable formulations reported on Table 1 of the Evans Application.

16. As a further part of the preformulation phase, the experienced formulator would have conducted a preformulation solubility screen, separately measuring the solubility of fulvestrant in a range of pure solvents, including the potential oil and co-solvent candidates that had been identified in the above literature review as being suitable for inclusion in intramuscular injection formulations. See, for example, Gupta (1999), Chapter 17 at page 402, under the heading “Formulation Development”:

The activities necessary to develop a parenteral product can be placed into the following three broad areas: preformulation, formulation, and scale-up. While there are alternative development perspectives, all development ultimately needs to accomplish the same activities. Preformulation includes the characteristics of the bulk drug plus initial screening for excipient compatibility with the drug.

“Preformulation studies” are said to “provide fundamental data and experience necessary to develop formulations for a specific compound” including, as item 8.1 in the outline of areas of specific interest, a determination of “solubility” in “selected solvents” (at 403). “Significant formulation activities begin with initial preformulation data and knowledge of the specific route of administration” (at 405), which “formulation activities include the identification and selection of a suitable vehicle (aqueous, nonaqueous or co-solvent system) ...” (at 404). It is further noted that “injection volume is one of the most important considerations in the formulation development of a commercial product” (at 405). When carrying out such a preformulation solubility screen with fulvestrant, the formulator would have found that fulvestrant had extremely low solubility in water, low solubility in most oils (but highest in castor oil), low solubility in benzyl benzoate, and the highest solubility in ethanol and benzyl alcohol, such as reported in Table 2 of the Evans Application.

17. With the information on prior commercialized formulations and the fulvestrant solubility data from the preformulation screen (such as reported in Table 2 of the Evans

Application), the experienced formulator would have selected castor oil as the oil vehicle because of the higher solubility of fulvestrant in castor oil relative to the other oils tested. Nevertheless, he would have appreciated that the target fulvestrant concentration of at least 45 mg/mL could not be achieved with castor oil alone, and that a co-solvent would be required.

18. A number of the commercialized formulations that would have been identified in the literature review (including the castor oil-based formulations) have a substantial benzyl benzoate component, which may be present as a co-solvent. See, for example, Delalutin noted in paragraph 15 above, which is reported in PDR (1973) and noted in Table I of the Evans Application, and is one of the formulations discussed in Riffkin (1964), "Castor Oil as a Vehicle for Parenteral Administration of Steroid Hormones" (see Riffkin n. 6). Delalutin is 250 mg/mL 17-hydroxyprogesterone caproate dissolved in 52% castor oil, 46% benzyl benzoate and 2% benzyl alcohol. However, Riffkin Table II reports that the solubility of 17-hydroxyprogesterone caproate in castor oil alone is only 55.6 mg/mL, but the solubility of 17-hydroxyprogesterone caproate in benzyl benzoate is substantially higher, being at least 250 mg/mL (see example 4 of Huber (US '520) and Attachment E discussed below). Even if not needed as a cosolvent, Riffkin (1964) notes that "the addition of benzyl alcohol or benzyl benzoate to castor oil resulted in a lower and more favorable viscosity, making it easier to inject" (paragraph bridging pages 893-894).

19. However, the skilled formulator would have appreciated from the fulvestrant solubility data generated in the preformulation screen that fulvestrant had very different solubility characteristics relative to the steroids of previous commercial formulations. Attachment E is a compilation showing the chemical structures and relative solubilities in castor oil and sesame oil of the compounds named in Riffkin (1964) Table II compared to the structure and the solubility of fulvestrant in these oils. It can be seen that the solubility of fulvestrant in castor oil and in sesame oil (20 mg/mL and 0.58 mg/mL, respectively, from Table 2 of the Evans Application) is appreciably lower than the solubility of the other steroids in these oils (taken from Table II of Riffkin (1964)). The second page of Attachment E tabulates the concentration in benzyl benzoate of five named steroids, taken

from Examples 1-5 of Huber (US '520), ranging from 200 to 400 mg/ml.² By comparison, the solubility of fulvestrant in benzyl benzoate is reported in Table 2 of the Evans Application as being only 6.15 mg/mL, and only 3.8 mg/mL as determined in the recently conducted tests reported in Attachment C.

20. The experienced formulator thus would have expected that benzyl benzoate would *not* act as a co-solvent for fulvestrant in castor oil because the solubility of fulvestrant in benzyl benzoate was significantly lower than its solubility in castor oil. The addition of benzyl benzoate to castor oil, for whatever reason, would have been expected to *decrease, rather than increase*, the solubility of fulvestrant in the resulting castor oil/benzyl benzoate mixture. This is confirmed in Table 4 of the Evans Application, which reports a fulvestrant solubility of only 12.6 mg/mL in the castor oil vehicle containing only 15% benzyl benzoate, compared to the 20 mg/mL solubility of fulvestrant in castor oil alone as reported in Table 2.³
21. Based on the solubility data determined in the preformulation screen (such as reported in Table 2 of the Evans Application), ethanol and/or benzyl alcohol would have been seen as the best co-solvent candidates for raising the fulvestrant solubility to the 45 mg/mL target in the castor oil vehicle, and would also function to lower the viscosity of the resulting formulation and make it easier to inject. Consistent with this solubility data, Dukes (US '814) added 40% w/v benzyl alcohol in order to dissolve 50 mg/mL fulvestrant in the castor oil-based formulation used in the experimental rat studies of his Example 3. It thus would have been apparent that 40% w/v benzyl alcohol could function as a co-solvent in castor oil to achieve the target fulvestrant concentration. Nevertheless, the skilled formulator would have been concerned with using such a high alcohol content in intramuscular injectable formulations for administration to a human.

² Data taken from the Examples of Huber (US '520); these are concentrations used in the examples and not necessarily the actual maximum solubility of each steroid in benzyl benzoate, which may be higher. Huber was a co-author on Riffkin (1964).

³ It should be noted that in the further tests that were recently conducted under my guidance (paragraphs 7-9 above and Attachments B and C hereto), the solubility of fulvestrant in castor oil alone was again tested and found to be 21.4 mg/mL, and the solubility of fulvestrant in benzyl benzoate alone was again tested and found to be only 3.8 mg/mL, which further confirms that benzyl benzoate would not be expected to act as a cosolvent for fulvestrant in castor oil.

22. First of all, the experienced formulator would want to minimize the amount of co-solvents and excipients in any injectable formulation. For example, as stated in Gupta (1999), Chapter 17, "Formulation and Administration Techniques to Minimize Injection Pain and Tissue Damage Associated with Parental Products" at page 414:

Cosolvents are commonly used to enhance drug solubility and stability. Cosolvents may include ethanol, propylene glycol, polyethylene glycols, and glycerine. These components have intrinsic effects on biologic tissue and can alter the properties of other excipients, thus influencing the tissue damage or pain caused by a product. There is a dearth of literature on the pain caused by cosolvents, but there is also a growing body of knowledge on the tissue damage that they can cause. It is not certain that tissue damage is always directly correlated with the injection pain, but minimization of both pain on injection and potential for tissue damage should be included in the product development plan.

See also Gupta (1999), Chapter 11, titled Cosolvent Use in Injectable Formulations, page 217:

Ideally, it is best to select and use solvents that would maximize the solubility of the compound. Maximizing the solubility of a compound in a particular cosolvent system would result in lower total levels of the non-aqueous solvent(s) being administered to the patient, thereby lowering the chance for potential side effects.

This objective would have applied to aqueous and oil-based systems alike, in that the precedent of commercialized formulations identified in the literature review would have confirmed that fixed oils, such as castor oil, have long been commercially used and accepted as the major component of oil-based sustained release intramuscular injectable steroidal formulations. On the other hand, co-solvents such as ethanol or benzyl alcohol have generally been used only in far lesser concentrations, as discussed in the following paragraph.

23. Thus, use of such a high content of either benzyl alcohol or ethanol would have been contrary to precedent as shown from the review of commercialized oil-based intramuscular injectable sustained release formulations. The literature review as of early 2000 would have shown that any benzyl alcohol in such formulations was almost always

present as a preservative in a concentration of about 2% or less, occasionally at a concentration of up to 5%, but only rarely at higher concentrations. With respect to benzyl alcohol see, for example:

- Gupta (1999), Chapter 11 at page 229 stating that benzyl alcohol “is typically used in concentrations of up to 2 percent as a preservative and up to 5 percent as a solvent,” and then discussing reported toxicities.
- Nema (1997), Table V at page 168, reporting that benzyl alcohol was present as an antimicrobial preservative in 74 injectable formulations (not limited to oil-based IM formulations) at concentrations of from 0.75-5% (note that benzyl alcohol is not included at all in Nema Table I, “Solvents and Co-solvents”);
- Powell (1998), the benzyl alcohol listing at pages 244-246, particularly those indicated as being used in IM formulations;
- Strickley I (1999) at page 329 notes the inclusion of 2% benzyl alcohol in an IM lorazepam formulation in a propylene glycol vehicle, but does not include benzyl alcohol at all in Table VI listing “Cosolvents Used in Parenteral Formulations;”
- Lopatin (1972) noting in Table 3 at page 727 opposite Benzyl alcohol, “Toxic. Used in concentration of not over 3%. Has irritant action in concentration of 5%;”
- Cornelius (US ‘863), col. 1, lines 30-35 stating, “It is known that the solubility of steroids in vegetable or animal oils can be increased by the addition of excipients such as benzyl alcohol and benzyl benzoate. An objection to the use of such excipients, and specifically benzyl alcohol in somewhat higher concentrations, is that these agents may irritate the tissues.”

The literature review as of early 2000 also would have shown that, with few exceptions, ethanol was not included in such formulations in excess of about 10%. See, for example:

- Gupta (1999), Chapter 11 at page 225 noting that ethanol has been used at levels up to 50 percent, but these levels typically are associated with pain on injection;
- Strickley I (1999), Table VI, “List of Cosolvents Used in Parenteral Formulations” more specifically lists the ethanol content in IM formulations for specifically identified drugs, which concentrations range only from 2.5 to 10%; an IM/IV lorazepam formulation in a propylene glycol vehicle is noted at page 329 as having 18% alcohol, but is not included with the IM formulations in Table VI;

- Nema (1997), Table I, “Solvents and Co-solvents” at page 167, lists ethanol as being in 24 formulations with a concentration range of 0.6-80% (for Prograf); note that this is misleading, however, since Prograf is a *concentrate* for intravenous infusion only, and is to be diluted 250 to 1000 times before administration;
- Powell (1998), lists “alcohol” at page 242 and “ethyl alcohol” at page 255, wherein the ethanol concentration for IM formulations ranges from 0.61-10%.

24. Thus, even though Dukes (US '814) had demonstrated that the target 45 mg/mL fulvestrant concentration could be achieved by adding 40% benzyl alcohol to the castor oil vehicle, the precedent of commercialized IM oil-based systems would have motivated the experienced formulator to substantially reduce the benzyl alcohol content of the formulation intended for human use, and this commercial precedent would have made him very reluctant to replace benzyl alcohol with the substantial amount of ethanol that would be needed to maintain the target fulvestrant concentration. Benzyl benzoate clearly would not be considered to solve this dilemma, but rather would be expected to have a negative effect on fulvestrant solubility since fulvestrant was even less soluble in benzyl benzoate than in castor oil, that is, one would have expected that adding benzyl benzoate would require still *more* alcohol to maintain the target fulvestrant concentration.⁴

25. Under these circumstances, the discovery by Evans *et al.*, that the addition of benzyl benzoate to the castor oil/alcohol mixture actually increases the solubility of fulvestrant such that more fulvestrant could be dissolved in a given volume of formulation, was unexpected and truly surprising. This positive benzyl benzoate effect on fulvestrant solubility in the resulting formulation is shown in Table 3 of the specification (and is not changed by the above-noted corrections), and is confirmed and demonstrated over a broader range of formulation composition by the additional set of experiments conducted under my guidance and discussed in paragraphs 7-9 above, the results of which are reported in Attachments C.

⁴ It should be noted that even apart from this solubility issue, there would have been no motivation to add benzyl benzoate for viscosity reduction since the significant quantity of alcohol would serve the dual function of acting as a co-solvent as well as reducing the injection viscosity and making it easier to inject, whereas the benzyl benzoate would be expected to have a negative effect on the fulvestrant solubility.

The undersigned declares further that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punished by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardise the validity of the application or any patent issuing thereon.

Signature: P. R. Markel.

Date: 8th August 2008.

TABLE OF REFERENCES

Tab	Author/Inventor	Reference Citation/Patent
1	Cornelius (US '863)	US Patent 4,212,863
2	Dukes (EP '014)	EP 0 346 014 A1 (corresponds to US Patent 5,183,814)
3	Dukes (US '814)	US Patent 5,183,814 (corresponds to EP 0 346 013 A1)
4	Gupta (1999)	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
5	Huber (US '520)	US Patent 3,164,520
6	Lopatin (1972)	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
7	Mackey (1995)	M.A. Mackey, A.J. Conway and D.J. Handelsman. Tolerability of intramuscular injections of testosterone ester in oil vehicle. <i>Hum. Reprod.</i> 10 : 862-865 (1995)
8	Nema (1997)	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
9	PDR (1973)	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
10	Powell (1998)	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 (1998)
11	Riffkin (1964)	C. Riffkin, R. Huber and C.H. Keysser. Castor oil as a vehicle for parenteral administration of steroid hormones. <i>J. Pharm. Sci.</i> 53 : 891-5 (1964)
12	Strickley I (1999)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
13	Strickley II (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
14	Strickley III (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
15	Wang (1980)	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

ATTACHMENT A

TABLE 3

		EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C							
		4°C							
		% w/v							
	Ethanol (96%)	5	5	10	10	10	10	15	15
	Benzyl Alcohol	5	5	5	5	10	10	15	15
	Benzyl Benzoate		15		15		15		15
Mean	Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
	Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	68 64	76 77	102 103
Individual values		27.8	35.5	54.4	64.1	48.4	68.9	65.8	80.6
		25.8	36.1	38.6	47.3	61.3	60.2	76.9	101.9
					53.0		63.2	90.0	121.6
					80.3		72.9	73.4	107.4

ATTACHMENT B:

Experimental Test Procedure for measuring the solubility of fulvestrant in different solvent vehicles at 25°C

1. Solvent vehicles for the solubility experiments were prepared by weighing the required amount of benzyl benzoate, benzyl alcohol and ethanol into a 20 ml volumetric flask and then diluting to volume with castor oil.
2. For each solvent vehicle in which the solubility of fulvestrant was to be determined, 1.0-1.5g of fulvestrant was weighed into each of 3 separate vials (2 dram size) and 5mls of the solvent vehicle was added to each vial, except for the pure castor oil vehicle, where 80mg of fulvestrant were weighed into each of the 3 separate vials and 2mls of the castor oil added to each vial. The reduced amount of fulvestrant and lower volume of solvent vehicle was needed to maintain stirring and achieve adequate mixing with the pure castor oil vehicle due to the combination of its higher viscosity and lower fulvestrant solubility/higher undissolved fulvestrant levels compared to the other solvent vehicles.
3. A magnetic stirrer bar was placed into each vial and the vials were capped and then placed on a magnetic stirrer block maintained at $25 \pm 0.5^{\circ}\text{C}$.
4. After 5 days of stirring at $25 \pm 0.5^{\circ}\text{C}$, an aliquot of each fulvestrant/solvent vehicle mixture was removed from each vial and placed into an Eppendorf tube which was then centrifuged at 12000 rpm for 5 minutes at ambient temperature.
5. For all but the fulvestrant/castor oil mixture, 1 ml of the supernatant was then removed from the Eppendorf tube and pipetted into a 10ml or 20ml volumetric flask and then diluted to volume with methanol and mixed to give a sample for analysis. The choice of whether to use a 10ml or 20ml volumetric flask for a particular sample was dependent on the likely concentration of fulvestrant in the sample and the quantifiable concentration range of the HPLC assay method used. For the fulvestrant/castor oil mixture, 100 μl of the supernatant was removed from the Eppendorf tube and pipetted into a 1ml volumetric flask and then diluted to volume with methanol and mixed to give a sample for analysis.
6. Step 5 was repeated to give a duplicate sample for analysis. Thus, this gave 2 samples for each of the 3 vials, giving a total of 6 samples for analysis for each solvent vehicle tested.
7. The resultant samples were analysed for fulvestrant content by reverse phase High

Performance Liquid Chromatography (HPLC). The HPLC method that was used is described below at point 9. The fulvestrant content obtained for each sample was used to calculate a value for the concentration of fulvestrant dissolved in the corresponding solvent vehicle after stirring for 5 days at 25°C.

8. The mean solubility of fulvestrant for each different solvent vehicle tested was calculated as the arithmetic mean of the 6 individual values for the concentration of fulvestrant dissolved in the corresponding solvent vehicle.

9. HPLC Method details:

Gradient HPLC Method

Eluent A : 27% Methanol / 32% Acetonitrile / 41% Water

Eluent B : 41% Methanol / 49% Acetonitrile / 10% Water

Column : 15cm 3.5µm Symmetry C8 4.6mm i.d.

Detection wavelength : 225 nm

Flow rate : 2 mL min⁻¹

Temperature : 40°C

Injection volume : 10 µL

Gradient programme :

Time (min)	Eluent A (%)	Eluent B (%)
0	100	0
25	100	0
55	0	100
65	0	100
66	100	0
70	100	0

Retention time of fulvestrant: 21 minutes approximately

ATTACHMENT C:

EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C

		% w/v																				
Ethanol (96%)	0	0	5	5	5	5	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Benzyl Alcohol	0	0	5	5	5	5	5	5	5	5	5	10	10	10	10	10	10	10	10	10	10	10
Benzyl Benzoate	0	100		10	15	25		10	15	25			10	15	25			10	15	25		
Castor oil	100	0	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Mean Fulvestrant solubility [mgml ⁻¹]	21.4	3.8	27.6	29.2	43.3	47.5	64.6	71.6	84.2	94.0	68.1	87.2	93.4	118.9	96.6	107.7	116.1	139.6	121.3	144.6	143.8	166.2
Individual values [mgml ⁻¹]	23.2	3.9	29.5	31.2	43.9	48.3	64.2	76.2	83.8	95.2	68.6	90.0	92.5	122.1	104.1	106.1	115.5	138.9	110.0	129.8	148.2	163.3
	17.8	4.0	28.3	26.3	45.1	50.7	66.8	72.1	81.9	97.8	68.9	84.9	92.1	120.3	74.0	86.6	117.9	141.0	120.0	133.5	147.1	164.8
	21.5	3.9	24.5	31.5	44.3	45.4	61.2	66.2	93.2	95.6	71.6	87.6	93.9	120.4	102.0	112.6	118.8	139.4	124.4	150.1	144.4	168.5
	21.8	3.8	26.6	29.3	45.4	45.2	66.0	65.7	84.6	96.1	67.6	88.1	93.0	118.3	98.6	117.9	116.1	142.1	125.6	151.7	144.4	169.7
	22.2	4.0	27.0	29.1	36.9	47.6	65.8	75.4	82.4	88.2	67.0	90.7	93.8	116.8	102.1	107.9	117.0	138.7	123.3	151.2	139.5	165.5
	22.0	3.2	29.6	27.8	44.3	47.6	63.6	73.9	79.1	91.0	64.8	82.1	95.3	115.7	98.4	115.1	111.5	137.9	124.6	151.1	139.1	165.5

ATTACHMENT D

TABLE 1

OIL-BASED LONG-ACTING STEROID/PEPTIDE INJECTIONS										
PRODUCT NAME	STEROID	DOSE	TYPE	CLASS ¹	SOURCE	OH ²	Safe	Safe	Exact	Other Details
METHANDIENOL	Dexamethasone propionate	20 mg	Androgen	Organic	ABPI Data	Anabolic	-	-	-	1 ml 3 weeks
	Fluoxymesterone phenylpropionate	60 mg								
	Dexamethasone isocaproate	60 mg								
	Fluoxymesterone decanoate	100 mg								
PROGESTIN DEPOT	Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	Schering BC	ABPI Data	Cortic	up to 40%	-	-	1 or 2 ml 2 week
TOSTERON	Hydroxyprogesterone succinate	300 mg	Progestin	Diamant	Dist. Vidal 1999	Steroid	-	-	-	2 ml 4-6 weeks
	Progesterone	50 mg or 150 mg								
TROPICOLONE	Testosterone Enanthate	1.5 mg	Androgen	Thermax	Dist. Vidal 1997	OH ²	-	-	-	1 ml 10 to 30 days
	Mestosterone undecanoate	60 mg								
	Hydroxyprogesterone laurate	80 mg								
NOXONEST	Mestosterone undecanoate	200 mg	Contraceptive	Schering BC	ABPI Data	Cortic	YES	-	1 ml 8 weeks	
BENZOTESTOSTERONE	Testosterone benzoate	5 mg	Androgen	Bioss	Dist. Vidal 1998	Anabolic	-	-	-	1 ml 3 week
PROGESTERONE-RETARD	Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	Fisher	Dist. Vidal 1999	Cortic	YES	-	1 or 2 ml 1 week	
ORAVIRMAN	Testosterone 17- β -valerate	5 mg/ml ³	Androgen	Schering BC	Dist. Vidal 1998	Cortic	YES	-	-	1 or 2-4
	Hydroxyprogesterone caproate	250 mg/ml ³								
PARABOLAN	Decaborone	76 mg	Androgen	Negroni	Dist. Vidal 1997	Anabolic	72 mg	40 mg	1.5 ml 2 weeks	
OIL ESTROGEN	Testosterone valerate	20 mg/ml ³ 40 mg/ml ³	Androgen	Bioss	J Pharm. Sci (1964) J Pharm. Sci (1964)	Cortic	74 80	10%	7%	-
OBLAATTEN	17-Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	BMS	J Pharm. Sci (1964) J Pharm. Sci (1964)	Cortic	YES	YES	-	up to 2 ml

OH² = hydroxylation
 BC = benzyl alcohol
 OH¹ = other OH
 Vidal = Dist. Vidal % are w/w and
³ concentration as measured directly from a single sample

Corrections to Table 1

In Table 1, the given values for the benzyl benzoate, benzyl alcohol and ethanol levels for the Delestrogen and Delalutin products have been incorrectly entered into the wrong columns. The entries are shown in their correct form in the attached corrected version of Table 1. The error is apparent from a review of the reference J.Pharm Sci (1964) 53 (8) 891 (Riffkin) which is stated in Table 1 as being the Source of the information for the Delestrogen and Delalutin products:

- In the Summary on page 895 of Riffkin, Delestrogen and Delalutin are identified as castor oil based commercially available products containing estradiol valerate at 20 & 40 mg/ml and 17-hydroxy-progesterone caproate at 250 mg/ml respectively.
- Furthermore, details of particular vehicle compositions for estradiol valerate and 17-hydroxy-progesterone caproate are given in Tables V and VI
 - In Table VI, the only 20 mg/ml formulation of estradiol valerate, also referred to as commercially available, has the composition castor oil 78%, benzyl benzoate 20% and benzyl alcohol 2%.
 - In Table VI, the only 40 mg/ml castor oil based formulation of estradiol valerate, has the composition castor oil 58%, benzyl benzoate 40% and benzyl alcohol 2%.
 - In Table V, there are three 250/mg/ml castor oil based formulations of 17-hydroxy-progesterone caproate that all contain benzyl benzoate. Two of these formulations also contain 2% benzyl alcohol and the other formulation does not contain benzyl alcohol ie they all contain up to 2% benzyl alcohol.
- None of the vehicle compositions disclosed in Tables V and VI in Riffkin contain ethanol. Therefore the entries in the Ethanol column of Table 1 for the Delestrogen and Delalutin products must have been incorrectly entered in the wrong column and should have been entered into the Benzyl Alcohol column.
- It is also apparent from Table VI that the 78% and 58% entries in the Benzyl Benzoate column of Table 1 for the Delestrogen products should have been entered into the Oil column and the 20% and 40% entries in the Benzyl Alcohol column should have been entered into the Benzyl Benzoate column
- The exact compositions for the Delestrogen and Delalutin products are confirmed in the Physicians Desk Reference (Edition 27, 1973) on page 1352.

In addition, the name of the steroid given in Table 1 for the Delalutin product should have been 17-hydroxy-progesterone caproate and not just 17-hydroxy-progesterone. Also the entry under the Company column for the same product should read BMS rather than DMS.

ATTACHMENT E

Structure of compounds disclosed in Riffkin et al.

17-Hydroxypregesterone caproate:




Testosterone:




Estradiol valerate:



Pregesterone:



On the other hand, fulvestrant has the following structure:



From Riffkin et al. Table II:

Steroid	Solubility [mg/ml] at 25°C	
	Castor oil	Sesame oil
Fulvestrant	28	0.38
17-Hydroxypregesterone caproate	55.5	23.4
Testosterone	38.6	5.4
Estradiol valerate	60.6	16.1
Pregesterone	52.0	22.9

Tabulation of data from Examples of Huber, 3,164,520:

Example	Steroid	Steroid concentration in benzyl benzoate (mg/ml)
1	16,17-dihydroxyprogesterone	200
2	testosterone palmitate	200
3	progesterone	250
4	Progesterone + 17-hydroxyprogesterone caproate	250 + 250
5	Testosterone enanthate	400

ATTACHMENT F

TABLE OF REFERENCES

Tab	Author/Inventor	Reference Citation/Patent
1	Cornelius (US '863)	US Patent 4,212,863
2	Dukes (EP '014)	EP 0 346 014 A1 (corresponds to US Patent 5,183,814)
3	Dukes (US '814)	US Patent 5,183,814 (corresponds to EP 0 346 013 A1)
4	Gupta (1999)	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
5	Huber (US '520)	US Patent 3,164,520
6	Lopatin (1972)	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
7	Mackey (1995)	M.A. Mackey, A.J. Conway and D.J. Handelsman. Tolerability of intramuscular injections of testosterone ester in oil vehicle. <i>Hum. Reprod.</i> 10 : 862-865 (1995)
8	Nema (1997)	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
9	PDR (1973)	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
10	Powell (1998)	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 [pages 238-255 provided] (1998)
11	Riffkin (1964)	C. Riffkin, R. Huber and C.H. Keysser. Castor oil as a vehicle for parenteral administration of steroid hormones. <i>J.Pharm.Sci.</i> 53 : 891-5 (1964)
12	Strickley I (1999)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
13	Strickley II (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
14	Strickley III (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
15	Wang (1980)	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

ATTACHMENT F

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Carlos M. Tellez
Attorney Docket Number:	11285.0056-00000

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Pages:				
Claims:				
Miscellaneous-Filing:				
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				
Extension - 1 month with \$0 paid	1251	1	150	150

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Miscellaneous:				
Request for continued examination	1801	1	930	930
Statutory or terminal disclaimer	1814	1	160	160
Total in USD (\$)				1240

Electronic Acknowledgement Receipt

EFS ID:	11825286
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Carlos M. Tellez
Filer Authorized By:	
Attorney Docket Number:	11285.0056-00000
Receipt Date:	17-JAN-2012
Filing Date:	15-OCT-2008
Time Stamp:	14:22:14
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Credit Card
Payment was successfully received in RAM	\$1240
RAM confirmation Number	710
Deposit Account	
Authorized User	

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
-----------------	----------------------	-----------	-------------------------------------	------------------	------------------

1	Rule 130, 131 or 132 Affidavits	Exhibit_1--SawchukCV.pdf	573897 0db3336ec7b725b7384203972fb4278a02935cb7	no	33
Warnings:					
Information:					
2	Rule 130, 131 or 132 Affidavits	Exhibit_2--Office_Action_16-Sep-2011.PDF	731211 4f15e37d81f3921bc18cf2849f9c70a49b6f50c3	no	11
Warnings:					
Information:					
3	Rule 130, 131 or 132 Affidavits	Exhibit_5--Osborne.PDF	272030 4e32db3fb5737723f50c8e449ea0179da1e80b8d	no	5
Warnings:					
Information:					
4	Rule 130, 131 or 132 Affidavits	Exhibit_8--Faslox_claims_with_proposed_amendments--09-Dec-2011.pdf	124576 45495791c328a7d8966a65508cf87928a54cd90	no	5
Warnings:					
Information:					
5	Rule 130, 131 or 132 Affidavits	Exhibit_9--US3164520.pdf	255520 89671a0136d2ae2d42cfce35a327341d1599cb6	no	2
Warnings:					
Information:					
6	Rule 130, 131 or 132 Affidavits	Exhibit_12--Merck_Index.PDF	283056 733b2ad6666064111e4d0f6e5f95aaf07a1191a7	no	8
Warnings:					
Information:					
7	Rule 130, 131 or 132 Affidavits	Exhibit_14--Lavy.pdf	145860 822312df838d475bdcca4823dde494952d18adbb	no	5
Warnings:					
Information:					
8	Rule 130, 131 or 132 Affidavits	Exhibit_15--Ismail.pdf	196801 d6457a120eb4a440923c0055fb92ab89a7365e4f	no	10
Warnings:					
Information:					
9	Non Patent Literature	Pages_from_File_History--23-Nov-2010_to_19-Dec-2011--EP_2_266_573--10180667.pdf	1273117 c2ebafe4125da30ba99dcca4b9b69e2b07d394ec6	no	46
Warnings:					
Information:					

10	Non Patent Literature	Pages_from_File_History--19-Jan-2011_to_19-Dec-2011--EP2286818--10180661.pdf	1199990 1c4606766d904accf7efedc7cafc03fa094e1e31	no	42
Warnings:					
Information:					
11	Rule 130, 131 or 132 Affidavits	Exhibit_3--McLeskey-.pdf	2883384 679455e1c82a25a4cbd04f88ef0c5bbea905ffad	no	15
Warnings:					
Information:					
12	Rule 130, 131 or 132 Affidavits	Exhibit_4--EP0346014-B1--Dukes-.pdf	1621917 a4de1af888d43ce8e2f9293549f04868eb2e018	no	14
Warnings:					
Information:					
13	Rule 130, 131 or 132 Affidavits	Exhibit_6--Wakeling-.pdf	138081 bde6a21a2cdce0a6dc4f595e036bd85a31ba7708	no	2
Warnings:					
Information:					
14	Rule 130, 131 or 132 Affidavits	Exhibit_7--US_2010-0152149A1-.pdf	711671 018c92234b63aecdff82b816f1e4bf8fb65bb19	no	10
Warnings:					
Information:					
15	Rule 130, 131 or 132 Affidavits	Exhibit_10--Riffkin-.pdf	421271 6358b8b493e6d354b7c8e75be125f8ff2f9626ec	no	5
Warnings:					
Information:					
16	Rule 130, 131 or 132 Affidavits	Exhibit_11--Nema-.pdf	656507 6b87d551e40378b970c2500b66aae11414fe376	no	6
Warnings:					
Information:					
17	Rule 130, 131 or 132 Affidavits	Exhibit_13--Guerrini-.pdf	388452 3aaebbcc6b379de3323f881d4626ad737ff5458	no	8
Warnings:					
Information:					
18	Rule 130, 131 or 132 Affidavits	Exhibit_17--WO_03-006064-.pdf	2208801 2ccc7cedbb97c919135349a171b4629417d96fa3	no	60
Warnings:					
Information:					

19	Non Patent Literature	Pages_from_file_history--07-Sep-2009_to_15-Dec-2011--EP_1_250_138--01900186.pdf	273713 62d9d727f5df88f5912b60f6bb78d071d86871d9	no	14
Warnings:					
Information:					
20	Non Patent Literature	11285-0056--Ismail.pdf	196708 0764a889d53ad854199c5f607db58bc5c4f04c94	no	10
Warnings:					
Information:					
21	Non Patent Literature	11285-0056--Lavy.pdf	145800 5f924db76257b91b39716191b33ed33fc78d3853	no	5
Warnings:					
Information:					
22	Non Patent Literature	11285-0056--MerckIndex.pdf	286744 0f74063ace9c86c0b93120415c7982f69d1e6db	no	8
Warnings:					
Information:					
23	Non Patent Literature	11285--0056--Guerrini-.pdf	388489 02126ff82853baa264be55a03882e90192463c1c	no	8
Warnings:					
Information:					
24	Rule 130, 131 or 132 Affidavits	Exhibit_16--Gellert_Declaration.pdf	26188243 e9d11fb259f0f64f886e275013489e678af70eeb	no	25
Warnings:					
Information:					
25	Terminal Disclaimer Filed	11285-0056--Terminal_Disclaimer.pdf	117851 cf75f4f4fc88c4b4e6d0640323124e0274331b7	no	4
Warnings:					
Information:					
26		Response_to_16-Sep-2011_OA--Final_version_17-Jan-2012.pdf	156764 1f0800313b0a0d6ca584d647ee4efd2b0c29e3ef	yes	28
	Multipart Description/PDF files in .zip description				
	Document Description		Start	End	
	Amendment Submitted/Entered with Filing of CPA/RCE		1	1	
	Claims		2	6	

	Applicant Arguments/Remarks Made in an Amendment		7		28
Warnings:					
Information:					
27	Rule 130, 131 or 132 Affidavits	Sawchuk_Declaration-- Final_Executed_Version--13- Jan-2012-.pdf	227943 c28d18fc2d36703e478af54c1bdf52e48b72f77e	no	27
Warnings:					
Information:					
28	Request for Continued Examination (RCE)	11285-0056-RCE.pdf	58137 a4fcc2e01f43253b596af8013e379e9018c3de6f	no	1
Warnings:					
This is not a USPTO supplied RCE SB30 form.					
Information:					
29	Transmittal Letter	11285-0056--IDS--2764723_1.pdf	62840 cd8e8453319af93099d7537ea119509e8ac4fe49	no	2
Warnings:					
Information:					
30	Information Disclosure Statement (IDS) Form (SB08)	11285-0056--FormSB-08.pdf	60677 b0dab4c887c0778c5ee913f3a393d430d34c34bc	no	1
Warnings:					
Information:					
This is not an USPTO supplied IDS fillable form					
31	Non Patent Literature	11285-0056--Gellert.pdf	26190826 dea521c94d1f3fa816f16a3e32905517a11fc4bdc	no	25
Warnings:					
Information:					
32	Fee Worksheet (SB06)	fee-info.pdf	33259 8e6b3c56a1c465297b62998668e3c51581ca3b37	no	2
Warnings:					
Information:					
Total Files Size (in bytes):			68474136		

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

RONALD J. SAWCHUK

PERSONAL DATA

Present Address: Department of Pharmaceutics
College of Pharmacy
Room 9-149A Weaver-Densford Hall
University of Minnesota
308 Harvard Street S.E.
Minneapolis, MN 55455
Telephone: (612) 624-0646
E-mail: sawch001@umn.edu

Home Address: 14934 Pixie Point Circle SE
Prior Lake, MN 55372
Telephone: (952) 226-6507

Born: May 29, 1940, Toronto, Ontario, Canada
Marital Status: Married, three children
Citizenship: Dual: U.S. and Canadian

EDUCATION

1959	(High School)	Oakwood Collegiate Institute, Toronto Secondary School Grade XIII)
1963	B.Sc. Phm.	University of Toronto, Toronto Ontario College of Pharmacy Licentiate No. 10748
1966	M. Sc. Phm.	University of Toronto, Toronto
1972	Ph.D.	University of California, San Francisco Pharmaceutical Chemistry (Pharmacokinetics)

PROFESSIONAL AND ACADEMIC EXPERIENCE

1963 - 1965	Teaching Assistant, University of Toronto
1966	Community Pharmacist (part-time), Toronto
1966 - 1968	Teaching Assistant, University of California
1971 - 1972	Instructor in Pharmaceutics, University of Minnesota
1972 - 1977	Assistant Professor of Pharmaceutics, University of Minnesota
1977 - 1983	Associate Professor of Pharmaceutics, University of Minnesota
1974 - 1982	Associate Director, Clinical Pharmacokinetics Laboratory, U of Minnesota
1982 - 1995	Director, Clinical Pharmacokinetics Laboratory, College of Pharmacy, U of Minnesota
1983 - 2010	Professor of Pharmaceutics, University of Minnesota
1983 - 1989	Director of Graduate Studies in Pharmaceutics, University of Minnesota
1983 - 1986	Acting Head, Department of Pharmaceutics, University of Minnesota
1984 (summer)	Quarter Leave, Sandoz Pharma, Pharmacokinetics and Drug Metabolism Dept., Basel, Switzerland (M. Lemaire)
1991 - 1994	Director of Graduate Studies in Pharmaceutics, University of Minnesota
1992 (Summer)	Quarter Leave, Sandoz Pharma, Drug Safety, Basel, Switzerland (W. Niederberger)
1996 - 1999	Member, Board of Directors, Century Mortar Club
1997 (Spring)	Semi-Quarter Leave, Toyama Medical and Pharmaceutical University, Japan (H. Sato)
1997 (Summer)	Semi-Quarter Leave, Novartis AG, PKDM, Basel, Switzerland (J. Vonderscher)
1998 - 1999	Head, Department of Pharmaceutics, University of Minnesota
2001 (Summer)	Faculty Development Leave, Novartis AG, PKDM, Basel, Switzerland (M. Lemaire)
2010 - present	Professor Emeritus of Pharmaceutics, University of Minnesota

APPOINTMENTS AND PROFESSIONAL RESPONSIBILITIES

1972 - present	Member, Graduate Program in Pharmaceutics, University of Minnesota
1982 - present	Consultant to the pharmaceutical industry
1995 - present	Director, Bioanalytic and Pharmacokinetic Services, University of Minnesota
1995 - present	Editorial Board, <i>Saudi Pharmaceutical Journal</i>
1996 - 2007	Editorial Board, <i>Journal of Pharmaceutical Sciences</i>
1996 - present	Member, Graduate Program in Neurosciences, University of Minnesota
2001 - present	Member, Graduate Program in Experimental and Clinical Pharmacology, U of M
2002 - present	Member, Graduate Program in Social, Administrative and Clinical Pharmacy, U of M
2008 - present	Editorial Advisory Board, <i>AAPS Journal</i>
2009 - present	Editorial Board, <i>Xenobiotica</i>

OTHER PROFESSIONAL ACTIVITIES

Prepared two videotapes on "Pharmacokinetics" for undergraduate instruction, 1974
Co-editor of a book with James Blanchard, Ph.D. and B.B. Brodie, Ph.D., entitled "Principles and Perspectives in Drug Bioavailability." S. Karger, Publisher, 1979
Assistant Director, Clinical Pharmacokinetics Laboratory, 1974-82
Consultant in the Establishment and Implementation of the Drug Quality Assurance Program, United Hospitals, St. Luke's Division, St. Paul, 1975
Participant in Critical Incidents Workshop, PDI - College of Pharmacy, 1977
Assessor in the Pharmacy Assessment Exercises, 1978
Coordinator for Continuing Education in Pharmacy, TV Series 1978, 1980
Expert, Bureau of Drugs and Biologics, Food and Drug Administration, 1982-84
Screening Committee, Abstracts, Basic Pharmaceutics Section, APS, APhA, 1981
Review of Grants, Medical Research Council (Canada) 1980-86
Review of Grants, British Columbia Health Care Foundation, 1981-84
Advisory Consultant, Site Visit Team NIH (NINCDS) Yale University School of Medicine, October 1979
Member, Site Visit Team NIH (NINCDS) University of Utah School of Medicine, January 1983
Member, Special Pharmacology Study Section NIH, April-June 1988
Review of Grants, Idaho State Board of Education, 1989-91
Review of Grants, Greater Minnesota Corporation, 1990-91
Organizer and Symposium Co-Chair, "Microdialysis in Drug Metabolism and Disposition Studies", for the Annual AAPS Meeting, San Antonio TX, 1992
Symposium Co-Chair, "Kinetic and Dynamic Challenges of the 90's", for the Annual AAPS Meeting, San Diego, CA, 1994
Organizing Committee Member for the NATO Advanced Study Institute, "Pharmacokinetics: From Theory to Practice", Erice, Italy, April 5-16, 1994
Co-organizer and Participating Instructor, "Pharmacokinetics for the Pharmacist and Pharmaceutical Scientist" University of Milan, Varese, September 10 -15, 1995.
Member, Board of Directors, Century Mortar Club, 1996-present.
National Advisory Committee, FAMU RCMi Program, Tallahassee, FL 1996-present
Co-organizer and Participating Instructor, "Pharmacokinetics for the Biomedical and Pharmaceutical Scientist", University of Milan, Varese, September 7 -12, 1997.
Scientific Advisory Committee, 1st Symposium on Microdialysis and Pharmacokinetics, Leiden, The Netherlands April 1998
Organizer and Participating Instructor, "Pharmacokinetics for the Biomedical and Pharmaceutical Scientist", University of Malta, Msida, September 6 -15, 1998.
Founder, Microdialysis Focus Group, American Association of Pharmaceutical Scientists, 1998.
Scientific Advisory Committee, 2nd International Symposium on Microdialysis in Drug Research and Development, Stockholm, Sweden, June 2000
Chair, Microdialysis Focus Group, American Association of Pharmaceutical Scientists, 1998-2000.
Co-Chair, Organizing Committee, 3rd International Symposium on Microdialysis in Drug Research and Development, Minneapolis, MN, USA, June 2002
Visiting Professor, Guilin Medical College, Guilin PRC (2002-2007)
Scientific Advisory Committee, 4th International Symposium on Microdialysis in Drug Research and Development, Vienna, Austria, June 2004

Scientific Advisory Committee, Abbott Laboratories, for the FDA Critical Path Initiative, September 2004
Scientific Advisory Committee, 5th International Symposium on Microdialysis in Drug Research and Development, Leiden,
The Netherlands, June 2006
GLP-1 Scientific Advisory Panel, Medtronic, Minneapolis, MN, April 2009-present

CURRENT AND PAST MEMBERSHIP IN PROFESSIONAL AND SCIENTIFIC SOCIETIES

American Association of Pharmaceutical Scientists (Fellow)
American Association for the Advancement of Sciences (Fellow)
American Pharmacists Association (APhA)
American Society for Pharmacology and Experimental Therapeutics
International Society of Anti-Infective Pharmacology
International Society for the Study of Xenobiotics
Technology Park, Heidelberg, Germany
Century Mortar Club (Board of Directors, 1996-98)
Rho Chi Honor Society

SCHOLARSHIPS, HONORS AND AWARDS

1964 Scholarship, Canadian Foundation for the Advancement of Pharmacy
1965-66 National Research Council of Canada
1965 Warner-Lambert Research Fellowship
1968-70 National Institute of Health (NIH) Training Grant
1981-82 Teacher of the Year, College of Pharmacy, University of Minnesota
1986 Recipient of Horace T. Morse-Amoco Foundation Award
1988 Fellow, American Association of Pharmaceutical Scientists
1990 Fellow, American Association for the Advancement of Sciences
1996 Hallie Bruce Memorial Lecture Award
1997 Fellowship, Japanese Society for the Promotion of Science
1999 Meritorious Manuscript Award, American Association of Pharmaceutical Scientists
2001 Weaver Medal of Honor
2004 Distinguished Lecture, Creighton University School of Pharmacy and Health Professions
2005 Academy of Distinguished Teachers, University of Minnesota
2006 Distinguished Lecture, Temple University School of Pharmacy
2007 APhA Research Achievement Award in the Basic Pharmaceutical Sciences

COMMITTEE APPOINTMENTS

COLLEGE OF PHARMACY

1972-73, 1973-74 Student American Pharmaceutical Association Minnesota Chapter (Faculty Advisor)
1972-75 Student Admissions and Academic Standing Committee, College of Pharmacy
1972-73 Task Force on College of Pharmacy Organization
1973-74 Continuing Education Committee
1972-78 Admissions Committee for Pharm.D. Program, College of Pharmacy (Chair 1973-74; 1977-78)
1974-75 University of Minnesota Health Sciences B/C Implementation Committee
1974-77 Constitution and By-laws Committee
1974-75 Unit K Committee, Graduate School
1975-76 Task Force on Pharm.D. Admissions
1976-78 Professional Education Committee
1977-78 Task Force on Travel
1977-78 Anatomy, Physiology, Pathology Study Group
1977-78 Drug Product Design and Evaluation Study Group
1976-78 Search Committee for Biopharmaceutics Faculty Member
1977-78 Search Committee for Assistant Director HCMC
1977-78 Search Committee for Research Associate, CEP Project D-1 (Chairman)
1978-79 Pharm.D. Program Planning Committee (Chairman)
1978-79, 1979-80 Computer Systems Committee (Chairman)

1979-80 Professional Education Committee (Chairman)
 1980-81 Educational Policy Committee (Chairman)
 1980-82 Externship Committee
 1981-82 Academic Standing Committee
 1981-83 Health Sciences Policy and Review Council
 1981-82 Graduate Faculty Nominations and Course Proposals Committee
 1982-83 Academic Standing Committee (Chairman)
 1982-83 Advisory Committee on Animal Care Facilities
 1983-85 Council of Directors of Graduate Studies
 1982-83 Task Force on Computers
 1983 Search Committee for Department Chairman (Chairman)
 1983 Search Committee for Clinical Faculty at HCMC
 1984 Ad Hoc Committee on External Pharm.D. Program
 1984 Executive Committee (Chairman)
 1984 Search Committee for Dean of College of Pharmacy
 1984 Search Committee for Psychiatry Position, St. Paul-Ramsey Medical Center
 1984 Search Committee for Clinical Faculty at Hennepin County Medical Center
 1985 Endowed Chair in Pharmaceutics Search Committee (Chair)
 1985 Assistant Professor in Pharmaceutics Search Committee (Chair)
 1985 Appointments, Promotion and Tenure Committee
 1985 Space Committee
 1985 Clinical Assistant Professor (MMC) Search Committee
 1985-89 Executive Committee
 1986-87 Appointments, Promotion and Tenure Committee (Chair)
 1986-90 Educational Policy Committee
 1986-87 Subcommittee of Educational Policy Committee
 1986 Search Committee for Endowed Chair (Chair)
 1986-87 College of Pharmacy Strategic Planning Committee
 1986-87 Subcommittee of Strategic Planning Committee to Develop College Goals and Objectives
 1987-90 Continuing Pharmacy Education Advisory Committee (Chair)
 1987-88 Admissions Committee
 1988-89 Admissions Committee (Chair)
 1989-91 Promotion and Tenure Committee
 1991-92 Promotion and Tenure Committee (Chair-Elect)
 1991-92 General Research Support Committee
 1992-93 Promotion and Tenure Committee (Chair)
 1992-93 General Research Support Committee
 1993-94 Academic Standing Committee (Chair-Elect)
 1994-95 Academic Standing Committee (Chair)
 1994-98 College Computer Committee
 1995-96 Promotion and Tenure Committee
 1995-96 Internal Organization and Leadership Task Force
 1996-97 Nontraditional Pharm.D. Task Force
 1997-98 Search Committee for Endowed Chair in Geriatric Pharmacotherapy
 1997-98 Admissions Committee
 1997-98 Search Committee for Immunotherapy Faculty Position (Chair)
 1998-2000 Search Committee for Pharmaceutics Faculty Position
 2000-2001 Educational Policy Committee
 2001-2002 Search Committee for ECP Faculty Position
 2001-2002 Educational Policy Committee (Chair)
 2001-2002 Search Committee for Pharmaceutics Faculty Position
 2001-2002 College of Pharmacy Phar. Sci. 2020 Committee, Capital Campaign (Co-Chair)
 2001-2004 College of Pharmacy Faculty Consultative Committee
 2002-2003 Educational Policy Committee (Past Chair)
 2002-2003 College of Pharmacy Collegiate Review Committee (Chair)
 2002-2003 College of Pharmacy Central Council (Faculty Representative)
 2002-2003 College of Pharmacy Instructional Development Working Group for the Duluth Expansion
 2003-2005 Search Committee for Pharmaceutics Faculty Position at UMD (Chair)
 2004-2007 College of Pharmacy Assessment Committee

2005-2006 Search Committee for Endowed Chair in Geriatric Pharmacotherapy
2006-2007 Search Committee for Pharmaceutics Faculty Position

UNIVERSITY COMMITTEE APPOINTMENTS

1974-78 Subcommittee on Academic-Industrial Interface, Academic Relations Committee, 3M Technical Forum
1975-76 Health Sciences Primary Health Care Program Committee (Alternate),
Solicitor for the University of Minnesota Consolidated Fund Drive
1977-78 Alternate Senator (U. of Minnesota)
1978-81 Senator (U. of Minnesota)
1984-85 Health Sciences Learning Resources Committee
1986 College Delegate to All-University Single Quarter Leave Working Group, Academic Affairs
1989 Health Sciences Policy and Review Council, Graduate School
1989-91; 1991-93 Biological Sciences (formerly Plant and Animal Sciences) Policy and Review Council, Graduate School
1991-93 Graduate Faculty Nominations Subcommittee, Biological Sciences Policy and Review Council, Graduate School
1992-93 Graduate Faculty Nominations Subcommittee (Chair), Biological Sciences Policy and Review Council, Graduate School
1995-1998 Biological Sciences Policy and Review Council, Graduate School
1997-98 Faculty Research Development Proposal Review Committee for the Academic Health Center
2001-2004 Academic Health Center Faculty Consultative Committee
2001-2002 SCFP Subcommittee on Twin Cities Facilities and Support Services (STCFSS)
2003 AHC Seed Grant Review Committee
2003 AHC FCC Internal Screening Committee for Academy of Excellence Nominees
2004-2007 All-University Honors Committee, University of Minnesota

STATE, NATIONAL, AND INTERNATIONAL COMMITTEE APPOINTMENTS

1974-76 Representative to AACP Council of Faculties
1977-78 AACP Task Force on Guidelines for Pharm.D. Accreditation
1980-82 Academic Advisory Committee, Kellogg Pharmaceutical Scientist Program
1981 Screening Committee for Academy of Pharmaceutical Sciences, Basic Pharmaceutics Section
1989-present Member, Scientific Committee, International Pharmaceutical Technology Symposium (FIP)
1990 Academic Affairs Committee, AACP (Member)
1990 Program Committee, Controlled Release Society Annual Meeting (Member)
1989-91 Continuing Education Committee, State Board of Pharmacy (Member)
1990-95 USP Committee of Revision (Member)
1991-93 NIH/NINDS Antiepileptic Drug Development Program (Consultant)
1995 Fellows Nominations Committee for AAPS, PPDM Section
1995 Screening Committee for AAPS PPDM Section Abstracts
1997-2000 Fellows Nominations Committee for AAPS, PPDM Section
1999-2000 Committee on AAPS Section Structure and Procedure Guideline
2000-2001 PPDM Vice Chair, American Association of Pharmaceutical Scientists
2000-2002 Co-Chair, Organizing Committee, 3rd International Symposium on Microdialysis in Drug Research and Development
2001-2002 PPDM Chair Elect, American Association of Pharmaceutical Scientists
2001-2002 Annual Program Planning Committee, American Association of Pharmaceutical Scientists
2001-2002 Program Coordinating Committee, American Association of Pharmaceutical Scientists
2002-2003 PPDM Section Chair, American Association of Pharmaceutical Scientists
2002-2003 PPDM Committee for Graduate Student Symposium Awardees, American Association of Pharmaceutical Scientists
2002-2003 Short Course Program Review Team, American Association of Pharmaceutical Scientists
2003-2004 PPDM Section Past-Chair, American Association of Pharmaceutical Scientists
2004-2007 Member-at-Large, American Association of Pharmaceutical Scientists Executive Council
2004-2006 Clinical and Operational Working Group (CORWG), NASA
2004-2005 AAPS Executive Council Liaison to the Clinical Sciences section of AAPS
2005-2006 AAPS Executive Council Liaison to the DDD section of AAPS
2005-2006 AAPS Executive Council Liaison to the PDD section of AAPS
2005-2006 AAPS Executive Council Liaison to the 2006 Annual Meeting Program Committee

2005-2006	AAPS Executive Council Liaison to the 2006 Annual Meeting Screeners
2005-2006	AAPS Executive Council Liaison to the 2006 Program Coordination Committee
2006	AAPS Reference Resources Task Force
2006-2007	AAPS Executive Council Liaison to the APQ section of AAPS
2006-2007	AAPS Executive Council Liaison to the PT section of AAPS
2006-2007	AAPS Executive Council Liaison to the International Affairs Committee
2009-2011	Epilepsy NINDS Steering Committee
2009-2011	NINDS Consortium to Study Bioequivalence of AED Products

INVITED PRESENTATIONS

Continuing Education Program (6 hours) Minneapolis, MN, 1973.
Upper Midwest Hospital Conference, 1974.
Continuing Education Program (6 hours) Rochester, MN, 1974.
University of Illinois, Chicago, IL, 1974.
Department of Clinical Pharmacology, University of Minnesota, 1974.
AACP Annual Meeting and Teachers' Seminar (Workshop Leader), Lake Kiamasha, NY, 1975.
Debate Symposium, "Drug Product Selection," St. Paul, MN, 1977.
Continuing Education for Minneapolis Veteran Pharmacists (2 hours), Minneapolis, MN, 1978.
Continuing Education in Pharmacy (2 hours), Mankato, MN, 1978
Continuing Education in Pharmacy "Seminar at Sea" (4 hours of instruction), 1978.
HPLC Workshop, Invited Lecturer, Bloomington, MN, 1978.
University of Kentucky, Lexington, KY, 1979.
American Association of Clinical Chemists, Midwest Section, Minneapolis, MN, 1979.
University of Illinois, Chicago, IL, 1979.
Smith Kline Corp., Philadelphia, PA, 1979.
Department of Pathology, St. Cloud Hospital, St. Cloud, MN, 1979.
Comprehensive Epilepsy Program, Minneapolis, MN, 1979.
University of North Carolina, Chapel Hill, NC, 1979.
Burrroughs Wellcome Co., Research Triangle Park, NC, 1979.
St. Paul-Ramsey Medical Center, St. Paul, MN, 1989.
Continuing Education in Pharmacy (4 hours) Minneapolis, MN, September-October, 1981.
Medical Research Council of Canada, Visiting Professor, University of British Columbia, Vancouver, 1982.
Invited Lecturer, National Institutes of Health, Epilepsy Branch, Bethesda, MD, 1982.
Geriatric Research, Education and Clinical Center, Bloomington, MN, September, 1982.
Continuing Education in Pharmacy (6 hours), Duluth, MN, September, 1982.
Ciba-Geigy, Pharmaceuticals Division, Ardsley, December 2, 1982.
Swiss Federal Institute of Technology, Zurich, Switzerland, June 19, 1984.
Biopharmacy Division, Sandoz AG, Basel, Switzerland, June 22, 1984.
Biopharmacy Division, Sandoz AG, Basel, Switzerland, July 24, 1984.
"Cyclosporine Pharmacokinetics in the Rabbit: In Vivo Disposition and In Situ Absorption Studies," Rhone-Poulenc Visiting Professor, University of Toronto, Ontario, February 5, 1985.
"Pharmacokinetics and Pharmacodynamics," Drug Therapy Symposium VI, St. Paul, MN, February 27, 1985.
"Absorption and Disposition Studies with Cyclosporine," Sandoz, AG, Basel, Switzerland, July 15, 1985.
"Absorption of Cyclosporine from Rabbit Small Intestine Using an In Situ Perfusion Model," Vorstand des Instituts fur Pharmazie U. Lebensmittelchemie der Ludwig-Maximilians-Universitat, Munich, West Germany, July 17, 1985.
"Analytic considerations in the Investigation of the Pharmacokinetics of Cyclosporine," Medizinischen Hochschule, Hanover, West Germany, September 11, 1985.
"Mixed-Order Absorption of a Sustained Release Carbamazepine Tablet in Humans," Institut fur Pharmazeutische Technologie der Johann Wolfgang Goethe-Universitat, Frankfurt am Main, West Germany, May 15, 1986.
"Simultaneous First- and Zero-order Absorption of Commercial Carbamazepine Tablets," 5th Symposium on Biopharmaceutics and Pharmacokinetics, Piestany, Czechoslovakia, May 22, 1986.
"Simultaneous First- and Zero-order Absorption of Tegretol in Human Volunteers," National Institutes of Health, Epilepsy Branch, NINCDS, Bethesda, MD, November 6, 1986.
"Comparison of Plasma AUCs using the Traditional Point-by-Point and Pooled Sample Methods: Application in the Analysis of Human Pharmacokinetics of Carbamazepine and its metabolites," Food and Drug Administration, Rockville, MD, July 20, 1987.

"Pharmacokinetics in Contemporary Pharmacy Practice," Minneapolis Veteran Pharmacists Association, Richfield, MN, September 15, 1987.

"The Absorption and Disposition Kinetics of Carbamazepine and its Metabolites in Humans," Ciba-Geigy, Summit, NJ, July 23, 1987.

The following four lectures were given in Beijing, Chengdu, and Guilin, China during a visit sponsored by the Chinese Academy of Medical Sciences in late October/early November 1987:

1. "Theory and Application of a Pharmacokinetic Model in Individualizing Dosing Regimens for the Aminoglycosides."
2. "First- and Zero-order Absorption of Carbamazepine from Commercial Tablets in Epileptic Patients and Normal Volunteers."
3. "Significance of Nonlinear Disposition Kinetics in the Adjustment of Dosing Regimens."
4. "Relative Bioavailability of Phenytoin Formulations: Problems in Assessment Due to Michaelis-Menten Elimination Kinetics."

"Does Tegretol need to be Dosed TID?" Comprehensive Epilepsy Program, Minneapolis, MN, March 21, 1988.

"The Kinetics of Absorption of Carbamazepine (Tegretol) and its Metabolism in Humans," Vorstand des Instituts der Pharmazie, Ludwig-Maximilians Universität, Munich FRG, June 8, 1988.

"Pharmacokinetic and Physiologic Considerations in Oral Controlled Drug Delivery," Novel Drug Delivery Symposium, Minneapolis, MN, September 20, 1988.

"Clinical Applications of the Two-Compartment Open Model," Regional Kidney Disease Program, Hennepin County Medical Center, Minneapolis, MN, November 16, 1988.

The following five lectures were presented in a Continuing Education in Pharmacy Program: "Concepts and Applications in Pharmacokinetics, Parts I and II"; "Therapeutic Response and Toxicity"; "Monitoring Drug Therapy"; and "Bioavailability and Bioequivalence", St. Thomas, Virgin Islands, March 8-13, 1989.

"The Pharmacokinetics of Zidovudine (AZT) with Some Observations on the Interaction with Probenecid," Queen's University of Belfast, Belfast, North Ireland, June 15, 1989.

"Pharmacokinetic and Analytical Considerations in Monitoring Zidovudine (AZT) Levels in Children with Aids," Fourth International Congress on Pediatric Laboratory Medicine, Washington, DC, August 23, 1989.

"Inhibition of Zidovudine Metabolism and Excretory Transport," Department of Pharmacodynamics, Semmelweis University of Medicine, Budapest, Hungary, September 13, 1989.

"Evaluating Bioequivalence," Western Michigan Society of Hospital Pharmacists, Grand Rapids, MI, March 2, 1990.

"Effect of Temperature and Medium of Analysis on Cyclosporine Concentration," Canadian Consensus Meeting on Cyclosporine Monitoring, Minaki Lodge, Canada, May 11, 1990.

"Studies of the Interaction between Zidovudine (AZT) and Probenecid in Animals and Humans." Pharmaceuticals and Process R & D, Ayerst Laboratories Inc., Rouse's Point, NY, August 17, 1990.

"Mechanistic Studies to Examine the Effect of Probenecid on the Brain Uptake of Zidovudine," Shanghai Medical University, Shanghai, P.R.C., October 13, 1990.

A lecture series (16 hrs) on the topic of "Clinical Pharmacokinetics and Therapeutic Drug Monitoring" was given to staff members of the Chinese Academy of Medical Sciences and Hospital Pharmacists, Beijing, P.R.C., October 15-20, 1990.

"Comparative Intestinal Absorption of Compounds of Varying Lipophilicity, and the Effect of Absorptive Water Flux." Lederle Laboratories, Pearl River, NY, September 12, 1991.

"Analysis of Zidovudine Distribution into Specific Brain Regions Utilizing Microdialysis," Bristol Myers-Squibb Research Institute, Princeton, NJ, September 17, 1991.

"Distribution of AZT Into Specific Brain Regions in the Rabbit Utilizing Microdialysis," University of Illinois College of Medicine, Peoria, IL, October 9, 1991.

"Studies on the Transport of Nucleosides into Specific Brain Regions Using Microdialysis with *In Vivo* Calibration." University of Florida, College of Pharmacy, Gainesville, FL, December 6, 1991.

"Analysis of Zidovudine Distribution into Specific Brain Regions Utilizing Microdialysis," University of Arizona College of Pharmacy, Tucson, AZ, February 17, 1992.

"Regional Considerations in the In Situ Intestinal Absorption of Glycylcycline and Minocycline, and the Effect of Solvent Drag," Lederle Laboratories, Pearl River, NY, May 11, 1992.

"Comparative Absorption of Fluorothymidine and Related Nucleosides in Different Anatomic Intestinal Regions," Lederle Laboratories, Pearl River, NY, May 11, 1992.

"Microdialysis Techniques for the Study of Drug Distribution, and the Problem of Recovery *In Vivo*," Europhor Toulouse, France, June 19, 1992.

"The Use of Microdialysis in Studying the Distribution of Exogenous Substances in Biological Tissues," Sandoz Pharma, Basel Switzerland, June 24, 1992.

"Inhibition of Brain Distribution and Systemic Clearance of AZT by Probenecid," Sandoz Pharma, Basel Switzerland, June 30, 1992.

- "Uptake of Zidovudine (AZT) into Rabbit Brain Using Microdialysis with *In Vivo* Calibration," Knoll AG, Ludwigshafen, Germany, July 1, 1992.
- "Microdialysis in the Study of the Distribution and Metabolism of Exogenous Substances," Pharmaceutical Chemical Institute, University of Heidelberg, Heidelberg, Germany, July 2, 1992.
- "The Relationship Between Urine and Plasma Concentrations of Lipophilic Drugs: Implications for Therapeutic Drug Monitoring," Sandoz Pharma, Basel Switzerland, July 8, 1992.
- "Estimation of the Elimination Rate Constant for Metabolites which Exhibit Formation-Rate Limited Disappearance," Sandoz Pharma, Basel Switzerland, July 23, 1992.
- "Experimental Determination of Free Tissue Levels Using Microdialysis," 4th Biennial Conference on Chemotherapy of Infectious Diseases and Malignancies, Prague, Czechoslovakia, August 31, 1992.
- "*In Situ* Intestinal Absorption of Tetracycline Derivatives and the Effect of Absorptive Water Flux," Lederle Laboratories, Pearl River, NY, November 13, 1992.
- "Reversibility of Carbamazepine Autoinduction upon Dose Termination in Normal Volunteers," Abbott Laboratories, Abbott Park, IL, December 2, 1992.
- "Barriers to the Oral Delivery of Drugs," Wyeth-Ayerst Research, Radnor, PA, February 23, 1993.
- "Preliminary Results of Studies which Examine the Distribution of the NMDA Antagonist, EAB 515, to Rat Brain," Sandoz Pharma, Basel Switzerland, April 26, 1993.
- "Microdialysis Calibration Using the Zero-Net Flux Method and Retrodialysis in Studying the Distribution of Exogenous Substances to Rat Brain," Sandoz Pharma, Basel Switzerland, April 26, 1993.
- "Investigation of the Pharmacodynamics of the NMDA Antagonist, EAB 515, in the Rat During Intravenous and Intracerebroventricular Administration." Sandoz Research Institute, Berne, Switzerland, April 28, 1993.
- "Comparative Distribution of AZT to Brain Tissue Extracellular Fluid During Intravenous and Intracerebroventricular Infusion." Food and Drug Administration, Rockville, MD, May 21, 1993.
- "Interspecies Scaling of Pharmacokinetics in the Evaluation and Development of New Antiepileptic Drugs." Natural Resources Research Institute, University of Minnesota—Duluth, Duluth, MN, August 11, 1993.
- "Application of Pharmacokinetic Principles in Practice." Minneapolis Veteran Pharmacists Association, St. Louis Park, MN, September 21, 1993.
- "Microdialysis as a Tool to Study Drug Delivery to the Brain." North Jersey American Chemical Society Drug Metabolism Discussion Group, Somerset, NJ, October 7, 1993.
- "Graduate Studies and Research Careers in Pharmaceutics." University of Minnesota—Duluth Department of Chemistry, Duluth, MN, December 3, 1993.
- "Microdialysis in Pharmacokinetic and Drug Metabolism Studies." 95th Annual Meeting, American Society for Clinical Pharmacology and Therapeutics, New Orleans, LA, April 1, 1994.
- "Modeling and Simulation of Complex Pharmacokinetic Systems." NATO Advanced Study Institute, Erice, Italy, April 12, 1994.
- "Microdialysis in the Study of Drug Distribution." NATO Advanced Study Institute, Erice, Italy, April 13, 1994.
- "Pharmacokinetic Studies Utilizing Microdialysis." Department of Pharmaceutical Chemistry, University of Kansas, Lawrence, KS, May 2, 1994.
- "Pharmacokinetic Studies Utilizing Microdialysis and On-Line HPLC." 4th International Workshop in Bioanalysis, Lawrence, KS, July 12, 1994.
- "Application of Microdialysis in Pharmacokinetic Studies." Gordon Research Conference in Drug Metabolism, Holderness School, Plymouth, NH, July 20, 1994.
- "Microdialysis and its Application in Pharmacokinetic Studies." Ciba-Geigy, Pharmacokinetics and Bioanalytics Division, Ardsley, NY, July 25, 1994.
- "Assessing Drug Transport in the Brain with Microdialysis." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
- "Applications of Microdialysis in Preclinical Pharmacokinetic Studies." 3M Pharmaceuticals, 3M Center, St. Paul, MN, November 29, 1994.
- "Problems in Assessing the Absorption of Carbamazepine from Sustained Release Dosage Forms in Epileptic Patients." Pharmavene, Inc., Gaithersburg, MD, February 23, 1995.
- "Selected Preclinical Pharmacokinetic Studies with Tacrine." Parke-Davis Pharmaceuticals, Ann Arbor, MI, May 5, 1995.
- "Brain Distribution and Metabolism Studies with Tacrine and Two Hydroxylated Metabolites." Department of Pharmaceutics and Pharmacodynamics, University of Illinois, Chicago, IL, July 28, 1995.
- "Microdialysis and its Application in Preclinical Drug Distribution and Absorption Studies." Chiron Corporation, Emeryville, CA, August 18, 1995.
- "The Principle of Quantitative Microdialysis and its Application in Preclinical Drug Distribution Studies." Genentech, Inc., South San Francisco, CA, October 9, 1995.
- "Graduate Programs and Research Opportunities in Pharmaceutics." 13th Annual Symposium on Pharmaceutical Sciences Graduate Programs, Merrillville, IN, October 21, 1995.

- "Principles of Microdialysis and Applications in Preclinical Drug Distribution and Absorption Studies," Wyeth-Ayerst, Pearl River, NY, December 6, 1995.
- "Microdialysis in Preclinical Drug Distribution Studies." Dupont Merck, Newark, DE, December 8, 1995.
- "Microdialysis and its Application to the Study of Drug/Metabolite Distribution in the Central Nervous System," University of Pittsburgh, Pittsburgh, PA, January 25, 1996.
- "Therapeutic Drug Monitoring: A Fodor's Guide." Hallie Bruce Memorial Lecture Award, Minnesota Society of Health Services Pharmacists, Minneapolis, MN, April 13, 1996"
- "Preclinical Studies of Drug Distribution to the Brain using Microdialysis." Pharmaceutical Peptides Inc, Cambridge, MA, May 2, 1996.
- "Microdialysis and its Application in Nonclinical Studies of Drug Distribution and Absorption." Bristol-Myers Squibb, Princeton, NJ, June 24, 1996.
- "Continuous Monitoring by Microdialysis in Neuropharmacokinetic Investigations." Faculty of Pharmacy, University of Tanta, Tanta, Egypt, March 5, 1997.
- "Preclinical Studies of Drug Distribution to the Brain using Microdialysis," Toyama Medical and Pharmaceutical University, Toyama, Japan, April 11, 1997.
- "Application of Pharmacokinetic Principles in Individualizing Aminoglycoside Dosing," Toyama Medical and Pharmaceutical University, Toyama, Japan, April 11, 1997.
- "Preclinical Studies of Drug Distribution to the Brain using Microdialysis," Meiji College of Pharmacy, Japan, April 18, 1997.
- "Individualizing Aminoglycoside Dosing and Once-a-Day Aminoglycosides," Meiji College of Pharmacy, Japan, April 18, 1997.
- "Education of Pharmacists and Pharmaceutical Scientists at the University of Minnesota," 260th Meeting on Continuing Education of Pharmacists, Okuda-Shinmachi, Toyama, Japan, April 26, 1997.
- "Pharmacokinetic Basis of Drug-drug Interactions," Novartis Workshop on Metabolic Drug-Drug Interactions, Schluchsee, Germany, October 14, 1997.
- "Microdialysis and its Application in Preclinical Pharmacokinetic Studies," Merck Research Laboratories, West Point PA, December 16, 1997.
- "Microdialysis and its Application in Preclinical Pharmacokinetic Studies," Merck and Co, Inc. Rahway NJ, December 17, 1997.
- "Brain Distribution Studies employing Microdialysis and Crossover Designs," *1st International Symposium in Drug Research and Development*, Noorwijkerhout, Netherlands, April 3, 1998.
- "Application of Sample Pooling in the Time Domain to Estimate CL, Vss and MRT in the Search for Lead Compounds." Chiron Corporation, Emeryville, CA, May 5, 1998.
- "Microdialysis as a Sampling Technique in Preclinical Pharmacokinetic Studies." Pfizer Inc, Groton CT, June 18, 1998
- "Assessing Drug Delivery to the CNS Using Microdialysis Sampling." School of Medicine, University of Minnesota, Duluth, October 19, 1998.
- "Pharmacokinetic Studies Using Microdialysis Sampling." American Association of Pharmaceutical Scientists Annual Meeting, San Francisco CA, November 18, 1998.
- "Applications of Microdialysis in Pharmacokinetics: Brain, Blood, and Middle Ear Fluid." Bristol-Myers Squibb, Wallingford CT, May 14, 1999.
- "Applications of Microdialysis in Preclinical Pharmacokinetics: Brain, Blood and Middle Ear Fluid." Parke-Davis, Ann Arbor, MI, May 21, 1999.
- "Blood Sample Pooling and the Determination of Mean Residence Times in High-Throughput Pharmacokinetic Screening." Parke-Davis, Ann Arbor, MI, May 21, 1999.
- "Role of controlled release formulations in the steady-state pharmacokinetics and pharmacodynamics of anticonvulsants" Impax Pharmaceuticals, Inc, Hayward CA, June 9, 1999.
- "Investigating Neuropharmacokinetics and Drug Delivery to the CNS using Microdialysis" *8th International Conference on In Vivo Methods: Monitoring Molecules in Neuroscience*. Stony Brook NY, June 19-23, 1999.
- "Use of Microdialysis in Pharmacokinetics" at the 8th BMSR Workshop on *Advanced Methods of Pharmacokinetic and Pharmacodynamic System Analysis*, Marina del Rey, CA June 25-26, 1999.
- "Applications of Microdialysis in Preclinical Pharmacokinetics." Amgen, Inc., Thousand Oaks, CA, June 28, 1999.
- "Pharmacokinetic -Pharmacodynamic Principles in Drug Development." Chiron Corporation, Emeryville, CA, August 20, 1999.
- "Microdialysis and its Application in Pharmacokinetics: Brain, Blood, and Middle Ear Fluid." Abbott Labs, Abbott Park IL Aug 27, 1999.
- "Distribution kinetics of antibiotics to the chinchilla middle ear" Department of Biopharmaceutical Sciences, Uppsala University, Uppsala, Sweden, March 16, 2000.
- "In Vivo Microdialysis as a Tool to Study Site Specific Drug Delivery" *Millennial World Congress of Pharmaceutical Sciences*. San Francisco CA, April 17, 2000.

- "In Vivo Microdialysis as a Tool to Study Site Specific Drug Delivery" *Engebretson Symposium on Drug Discovery and Development*. Minneapolis, MN. May 18, 2000.
- "In Vivo Microdialysis as a Tool to Study Drug Delivery". *19th Annual Robert S. Rozman Memorial Symposium*, Langhorne PA, May 25, 2000.
- "Basic Principles of Microdialysis, Experimental Setup". *Course on Basic and Advanced Aspects of In Vivo Microdialysis*", Stockholm, Sweden, June 14, 2000.
- "Recovery: Basic Idea and Practical Methods". *Course on Basic and Advanced Aspects of In Vivo Microdialysis*", Stockholm, Sweden, June 14, 2000.
- "Studies of Distribution of Antibiotics to the Middle Ear by Microdialysis" *2nd International Symposium on Microdialysis in Drug Research and Development*, Stockholm, Sweden, June 15, 2000.
- "Basic Concepts in Clinical Pharmacokinetics" A 2-Day Course. Abbott Laboratories, Abbott Park IL and Victory Hospital, Waukegan, IL, July 18-19, 2000
- "Microdialysis and its Application in Preclinical Pharmacokinetics: Brain, Blood, and Middle Ear Fluid." Dupont Pharmaceuticals, Wilmington, DE July 12, 2000.
- "Pharmacokinetic-Pharmacodynamic Principles in Drug Development." Abbott Labs, Abbott Park IL Jan 9, 2001
- "Biopharmaceutical and Pharmacokinetic Considerations in Delivering Drug to the CNS" Medtronic Neuro Division, Minneapolis. January 25, 2001
- "Clinical Pharmacokinetic Principles in Drug Development." Novartis Pharma, Tokyo, April 12, 2001
- "In Vivo Microdialysis as a Tool to Study Site Specific Drug Delivery" Showa University, Tokyo, Japan, April 13, 2001
- "In Vivo Microdialysis as a Tool to Study Drug Delivery in Preclinical Studies". Xi'an Medical College, Xi'an, PRC. April 25, 2001
- "Principles of Pharmacokinetics and their Application in Drug Development" Novartis Pharma, Basel, Switzerland, July 3, 2001.
- "Microdialysis and its Application in Preclinical Studies of Drug Delivery to Target Tissues" Boehringer-Ingelheim Pharma KG, Dept. of Pharmacokinetics & Drug Metabolism, Biberach, Germany, July 5, 2001.
- "Estimation of Intrinsic Clearances and Organ Partition Coefficients in an Organ Perfusion Model" Novartis Pharma, Basel, Switzerland, July 26, 2001.
- "Pharmacodynamic Modeling of the Sigmoid Emax Model" Novartis Pharma, Basel, Switzerland, July 31, 2001.
- "Prediction of the Pharmacokinetics of Cefdinir in Children from the Results of Animal Studies. Omnicef® Clinical Advisory Meeting, Dallas, TX, February 9, 2002.
- "Applications of Microdialysis in Studying Drug Delivery to Specific Targets". Guilin Medical School, Guilin PRC, March 28, 2002
- "Microdialysis: A Tool to Study Brain Uptake?" Gordon Research Conference on the Barriers of the CNS, Tilton School, Tilton NH, June 25, 2002
- "A Model for the Distribution of Drugs between Plasma, CSF and Parenchyma", Workshop on Microdialysis Techniques in the CNS, Gordon Research Conference on the Barriers of the CNS, Tilton School, Tilton NH, June 26, 2002
- "Microdialysis in the Study of Drug Delivery to the Central Nervous System", Department of Pharmaceutics, Seoul National University, Seoul, South Korea, November 25, 2002.
- "Investigating Antibiotic Delivery to the Middle Ear". Chong Kun Dang Pharma, Cheonan, South Korea, November 27, 2002.
- "Microdialysis and its Application in Preclinical Pharmacokinetic and Drug Delivery Investigations", 32nd Annual Meeting of the Korean Pharmaceutical Society, Seoul, South Korea, November 28, 2002.
- "Applications of Pharmacokinetic Principles in Drug Development". Schering-Plough Research Institute. Kenilworth, NJ. December 19, 2002
- "A Course in Pharmacokinetics in Pharmaceutical Development". Abbott Laboratories. Harrison Conference Center, Lake Bluff, IL. May 15-16, 2003
- "Characterizing Antibiotic Delivery to the Middle Ear for the Treatment of Otitis Media. Biomedical Simulations Resource Workshop: Advanced Methods of PK/PD Systems Analysis. Marina del Rey, CA. June 20-21, 2003.
- "Cerebrospinal Fluid Distribution of Intrathecally Administered Antiviral Nucleosides". Monitoring Molecules in Neuroscience. 10th International Conference on In Vivo Methods. Department of Neuroscience, Karolinska Institutet Stockholm, Sweden. June 24-27, 2003
- "Microdialysis Sampling in Drug Development: Applications in Preclinical Research." Sunrise School, American Association of Pharmaceutical Scientists Annual Meeting, Salt Lake City, UT, October 26, 2003.
- "Clinical Pharmacokinetics in Pharmaceutical Development." Abbott Laboratories. Harrison Conference Center, Lake Bluff, IL. July 23-24, 2003.
- "Microdialysis Sampling in Drug Development: Applications in Preclinical Research." Sunrise School, American Association of Pharmaceutical Scientists Annual Meeting, Salt Lake City, UT. October 26, 2003.
- "The Role of Pharmacokinetics in Drug Discovery." Abbott Laboratories. Harrison Conference Center, Lake Bluff, IL. March 18, 2004.

- “Microdialysis and its Application in Preclinical Pharmacokinetic and Drug Delivery Investigations.” CDER, Food and Drug Administration, Rockville, MD. March 29, 2004.
- “Interspecies Scaling, PB-PK modeling and Microdialysis in Antibiotic Drug Development.” Novartis Institute for Biomedical Research, Cambridge, MA. April 9, 2004.
- “Does it get to the Target Site? Microdialysis as a Tool to Study Preclinical Drug Distribution and Delivery” Amgen Inc., Thousand Oaks, CA. April 30, 2004.
- “Microdialysis of Antibiotics.” 4th International Symposium on Microdialysis in Drug Research and Development, Vienna, Austria, June 19, 2004.
- “The Chinchilla Microdialysis AOM Model” Pfizer Global Pharmaceuticals, New York, NY. June 25, 2004.
- “Advantages of the Chinchilla Microdialysis Model” Scientific Basis for Tissue-Directed Antimicrobial Therapy Symposium, Boston MA, July 21-22, 2004.
- “Evaluating Drug Distribution to the Target Site and Predicting Tissue Exposure in Humans from Animal Data” Scientific Advisory Committee, Abbott Laboratories. The FDA Critical Path Initiative and the Role of Modeling/Simulation in Improving the Efficiency of Drug Development. Lake Forest, IL. September 89, 2004.
- “Assessing Drug Delivery to the Target Site: The Role of Microdialysis in Measuring Tissue Exposure in Animals and Humans.” Distinguished Lecture, Creighton University School of Pharmacy and Health Professions, Omaha NE, November 30, 2004.
- “Microdialysis—Introduction to Basic Principles and Applications”. AAPS Workshop on Microdialysis Principles, Application, and Regulatory Perspectives, Nashville TN, November 4, 2005.
- “A Phase I Open-Label, Dose-Ranging Study to Investigate the Safety and Tolerability of Gabapentin Injection Administered Intrathecally in Individuals with Chronic, Intractable Pain: A Pharmacokinetic Report”. Medtronic WHQ, Fridley, MN, February 16, 2006.
- “Public Outreach and AAPS: Students are the Future of Our Association”. Temple University School of Pharmacy, Philadelphia, PA. February 20, 2006.
- “Assessing Drug Delivery: Using Microdialysis to Measure Target Site Exposure in Animals and Humans”. Wyeth Distinguished Lecture Series, Temple University School of Pharmacy, Philadelphia, PA. February 20, 2006.
- “Pharmacokinetics for Scientists Engaged in Drug Discovery”. Lundbeck Research, USA. Paramus NJ. February 24, 2006.
- “Pharmacokinetic Issues related to Intrathecal Drug Dosing”. Medtronic WHQ, Fridley, MN, March 15, 2006.
- “TTM Technology: Antibiotic Distribution to Middle Ear Fluid” Abbott Laboratories, Abbott Park, IL. May 16, 2006
- “Trans-tympanic Membrane (TTM) Drug Delivery to the Middle Ear” Alcon Laboratories, Fort Worth TX. Feb 2, 2007.
- “Bugs and Drugs: Does the Anti-infective Agent get to the Target Site?”. Science Luncheon Presentation. APHA Annual Meeting. Atlanta, GA. March 18, 2007
- “Future Perspectives on the Contributions of Microdialysis in Drug Research and Development” Keynote Address. Fifth International Symposium on Microdialysis in Drug Research and Development. Leiden, NE. April 25, 2007.
- “Drug Delivery to the Middle Ear across the Tympanic Membrane for Therapy of Acute Otitis Media”. Global Gators 6th Symposium on Clinical Pharmacy and Clinical Pharmacology. Munich, Germany. June 9, 2007.
- “The Pharmacokinetics of Hydrophilic Drugs during Intrathecal Infusion: the Concept of a Targeted Delivery Advantage”. Novartis Pharma AG, Basel, Switzerland. June 13, 2007.
- “Trans-tympanic Membrane Delivery of an Antibiotic into Chinchilla Middle Ear” Alcon Laboratories, Fort Worth TX. October 15, 2008.
- “A Phase I Study to Investigate the Safety and Pharmacokinetics of Intrathecal Gabapentin Injection in Individuals with Chronic Pain”. University of Poitiers, Poitiers, France, April 29, 2009.
- “Cerebrospinal fluid flow, and the convective/diffusive transport of drug in the CSF” Abbott GmbH and Co., Ludwigshafen, Germany. Oct 15, 2010.
- “The Neuropharmacokinetics of Hydrophilic Drugs during Intrathecal Infusion: the Concept of a Targeted Brain Delivery Advantage” Abbott GmbH and Co., Ludwigshafen, Germany. Oct 15, 2010.
- “A Brief Introduction to Pharmacokinetics” Upsher-Smith Laboratories, Inc., Maple Grove, MN. December 2, 2010.
- “CSF flow, and convective/diffusive transport of drugs” Upsher-Smith Laboratories, Inc., Maple Grove, MN. December 2, 2010.
- “Modeling the delivery of drugs to target sites in the CNS” Upsher-Smith Laboratories, Inc., Maple Grove, MN. December 2, 2010.

TEACHING AT THE UNIVERSITY OF MINNESOTA

Undergraduate

1971 - 1972	Co-instructor in Phar 5680 "Pharmacokinetics"
1971 - 1975	Discussant in Pharm.D. Conferences
1972 - 1973	Participating instructor in Phar 5670
1972 - 1978	Discussion leader in Pharm.D. I conferences
1972 - 1985	Course director, Phar 5680 "Pharmacokinetics"
1975 - 1995	Course director, Phar 5685 "Clinical Pharmacokinetics"
1991 - 1999	Course director, Phar 5681 "Basic Pharmacokinetic Modeling"
1996 - 1998	Course director and Participating instructor, Phmc 5460 "Pharmacokinetics"
1998 - 2003	Course director and instructor, Phar 6216 "Pharmacokinetic Simulation and Data Analysis using SAAM"
1999 - 2004	Course director and Participating instructor, Phar 6163 "Pharmacokinetics"
1998 - 2004	Participating instructor in Phar 6164 "Biopharmaceutics"
2004 - 2010	Participating instructor, Phar 6163 "Pharmacokinetics"

Graduate

1972 - 1999	Course director in Phm 8420 "Modeling Approaches in Pharmacokinetics" participating instructor in Phm 8421, Phm 8425
1972 - 2005	Participating instructor in Phm 8100 (Seminar) and Phm 8101 (Pharmaceutics Readings)
1984 - 1999	Participating instructor in Phm 8425 "Advanced Topics in Pharmacokinetics"
1986 - 1999	Course co-director in Phm 8105 "Pharmacokinetics Research Seminar"
2000 - 2006	Course co-director in Phm 8150 "Pharmacokinetics Research Seminar"
2000 - 2006	Course Co-director and Participating instructor in Phm 8421 "Advanced Pharmacokinetics"
2004 - 2010	Participating instructor in Phm 8481 "Advanced Neuropharmaceutics"
2006 - 2010	Participating instructor in Phm 8421 "Advanced Pharmacokinetics"

TEACHING AT OTHER SITES

- "An Introduction to Clinical Pharmacokinetics" Abbott Laboratories. Abbott Park, IL. January 9-10, 2001.
- "An Introduction to Clinical Pharmacokinetics" Abbott Laboratories. Abbott Park, IL. March 29-30, 2001.
- "An Introduction to Clinical Pharmacokinetics" Abbott Laboratories. Abbott Park, IL. May 17-18, 2001.
- "An Introduction to Clinical Pharmacokinetics" Abbott Laboratories. Abbott Park, IL. May 17-18, 2001.
- "An Introduction to Clinical Pharmacokinetics" Abbott Laboratories. Lake Bluff, IL. November 8-9, 2001.
- "An Introduction to Pharmacokinetics" Abbott Laboratories. Lake Bluff, IL. March 14-15, 2002.
- "An Introduction to Pharmacokinetics" Abbott Laboratories. Abbott Park, IL. July 22-23, 2002.
- "An Introduction to Pharmacokinetics" Abbott Laboratories. Parsippany, NJ. Aug 26-27, 2002.
- "An Introduction to Pharmacokinetics" Bristol-Myers Squibb. Wilmington, DE. September 19-20, 2002.
- "An Introduction to Pharmacokinetics" Schering Plough Corp. Kenilworth, NJ. December 19-20, 2002.
- "An Introduction to Pharmacokinetics" Abbott Laboratories. Lake Bluff, IL. July 23-24, 2003.
- "Short Course in Pharmacokinetics for Drug Discovery" Abbott Laboratories. Lake Bluff, IL. March 17, 2004.
- "Preclinical Pharmacokinetics in Pharmaceutical Discovery." Bristol-Myers Squibb, Princeton, NJ. May 6-7, 2004.
- "Introduction to Pharmacokinetics" Abbott Laboratories. Abbott Park, IL, July 27-28, 2004.
- "Introduction to Clinical Pharmacokinetics" Millennium Pharmaceuticals, Inc. Cambridge, MA, December 23, 2004.
- "Introduction to Clinical Pharmacokinetics" Gilead Sciences, Foster City, CA. December 8-9, 2005.
- "Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist" Co-instructor. Boehringer-Ingelheim Pharmaceuticals, Inc., USA. Ridgefield, CT, April 13-14, 2006
- "An Introduction to Pharmacokinetics" Lundbeck Research, USA, Inc. Paramus, NJ, February 24, 2006.
- "Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist" Co-instructor. Abbott Laboratories. Abbott Park, IL. June 4-5, 2007.
- "Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist" Co-instructor. Theravance, Inc. South San Francisco, CA. August 20-21, 2007.
- "Basic Pharmacokinetic Concepts" Co-instructor. US Patent and Trademark Office. Alexandria, VA. October 4, 2007.
- "Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist" Co-instructor. Allergan, Inc. Irvine, CA. July 24-25, 2008.

- “Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist”. Co-instructor. Abbott Laboratories, Abbott Park, IL. August 19-20, 2008.
- “Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist”. Co-instructor. Gilead Sciences, Foster City, CA. October 9-10, 2008.
- “Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist”. Co-instructor. Genentech, South San Francisco, CA. July 23-24, 2009.
- “Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist”. Co-instructor. Abbott Laboratories. Abbott Park, IL. July 30-31, 2009.
- “Neuropharmacokinetic Concepts for CNS Drug Delivery”. Co-instructor. Abbott Laboratories. Abbott Park, IL. January 8, 2010.
- “Basic Pharmacokinetic Concepts for the Pharmaceutical Scientist”. Co-instructor. Abbott Laboratories. Abbott Park, IL. August 4-5, 2010.
- “Basic Pharmacokinetic Concepts for the Upsher-Smith Pharmaceutical Scientist”. Co-instructor. Upsher-Smith Laboratories, Maple Grove, MN. September 21-23, 2011.
- “Basic Pharmacokinetic Short Course for Pharmaceutical Scientists”. Co-instructor. Novartis Pharma, Florham Park, NJ. November 17-18, 2011.

GRADUATE STUDENTS SUPERVISED

Graduate Students supervised as Primary Advisor:

1978	Wargin, W.A.	Ph.D.
1978	El-Yazigi, A.	Ph.D.
1980	Mugure Pyron	M.S.
1981	Sue-Chi Wu	M.S.
1983	Hsuehling Su	M.S.
1984	Dale Yu	Ph.D.
1984	Walid Awni	Ph.D.
1985	Lillian Riad	M.S.
1985	Rose Eggerth	Ph.D.
1987	Hisham Abou-Auda	Ph.D.
1989	Mohsen Hedaya	Ph.D.
1989	Ajit K. Shah	Ph.D.
1989	Lillian Riad	Ph.D.
1991	Helen Chan	Ph.D.
1992	William Elmquist	Ph.D.
1992	Shekman Wong	Ph.D.
1994	Yanfeng Wang	Ph.D.
1994	Bimal Malhotra	Ph.D.
1996	Richard Brundage	Ph.D.
1997	Zheng Yang	Ph.D.
1998	Belinda Cheung	Ph.D.
2001	Yue Huang	Ph.D.
2001	Guanfa Gan	Ph.D.
2002	Joanna Peng	Ph.D.
2002	Tong Zhu	Ph.D.
2004	Ji Ping	Ph.D.
2004	Wei Liu	Ph.D.
2005	Yan Song	Ph.D.
2007	Nael Mostafa	Ph.D.
2007	Zhihong Li	Ph.D.

GRANTS, CONTRACTS, and OTHER SUPPORT

1972-73	University of Minnesota Graduate School
1973-74	University of Minnesota Media Production Fund
1975-78; 1978-80	NIH/NINCDS Comprehensive Epilepsy Program Contract (Principal Investigator, Project D-1)
1976; 1977	Medical Education and Research Foundation Grant (Co-investigator with John W. McBride, M.D.)
1976-78	FDA Contract to Study the Pharmacokinetics and Toxicology of Phenytoin Sodium Products in Clinical Patients
1980-81; 1981-83	Grant to Support Research Involving the Analysis of Cyclosporin A in Biological Fluids (Sandoz, Inc.)
1982-83; 1983-84	"Pharmacokinetics and Biopharmaceutic Studies of Cyclosporin A in Selected Animal and <i>In Vitro</i> Systems" (NIH; Principal Investigator; Co-investigator, R.P. Enever)
1984	Comparative Bioavailability of Sodium Phenytoin in Normal Volunteers (ZenithLabs)
1984	Relative Bioavailability of Carbamazepine in Chewable and Conventional Tablets (Ciba-Geigy)
1984	Transdermal Delivery of Propranolol (Medtronics)
11/84 - 1/85	Absorption and Metabolism of Carbamazepine in Normal Volunteers (Ciba-Geigy)
1/85 - 4/85	Transdermal Absorption of α -Blockers (Medtronics, Inc.)
11/85 - 4/86	Relative Bioavailability of Sustained Release Oral Dosage Forms of Carbamazepine (Ciba-Geigy)
1/86 - 6/86	Analysis of Analgesics in Receptor Media (Medtronics)
8/86 - 12/86	Bioequivalence of Carbamazepine Oral Dosage Forms (Ciba-Geigy)
1/86 - 12/88	Pharmacokinetics of Diltiazem in the Rabbit (Marion)
2/87 - 9/87	Bioequivalence of Carbamazepine Dosage Forms Demonstrating Varying Dissolution Rates (Ciba-Geigy)
6/1/87 - 10/15/87	Effect of Urine Flow on the Renal Clearance of Carbamazepine and its Metabolites in Humans (Ciba-Geigy)
1/88 - 6/88	Effect of Fasting on the Absorption of Diclofenac Sodium in Normal Human Volunteers (Ciba-Geigy)
4/1/89 - 3/31/92	Enhancing Brain Uptake of AZT by Transport Inhibition, (NINCDS / NIH)
7/1/89 - 6/30/90	Induction of Carbamazepine Metabolism as a Function of Dosing Rate in Normal Volunteers (Ciba-Geigy)
9/91 - 6/92	Brain Distribution of EAB-515 in the Rabbit (Sandoz, Ltd.)
11/91 - 5/92	<i>In Situ</i> Absorption from Rabbit Intestine (Lederle Laboratories)
3/92 - 8/92	Clinical Studies of the Absorption of an Oral Immunosuppressant (Apotex Laboratories)
11/92 - 10/93	Brain Distribution of an NMDA-Receptor Antagonist in the Rat (Sandoz, Ltd.)
10/92 - 5/93	Brain Uptake of a CNS-Active Agent (Warner-Lambert)
11/92 - 3/93	<i>In Situ</i> Absorption from Rabbit Intestine (Lederle Laboratories)
9/94 - 5/95	Population Pharmacokinetic Analysis of A General Anesthetic in Man (Abbott Laboratories)
7/94 - 5/95	Brain Uptake of a Cholinesterase Inhibitor and its Metabolites (Warner-Lambert)
10/94 - 9/95	Distribution of Antiviral Nucleosides into Rat Cortex (Bristol-Myers Squibb)
9/95 - 8/96	Bioanalytical Methods Development of Selected Drugs and Metabolites (MedTox)
1/96 - 9/96	Pharmacokinetic Analysis of IL-2 in the Pig (Chiron)
1/96 - 6/98	Analysis of Selected Macrolides by High-pressure Liquid Chromatography (TAP)
4/96 - 10/97	Brain Penetration of Fosphenytoin and Phenytoin in the Rabbit (Warner-Lambert)
4/96 - 9/96	Analysis and Brain Uptake of PPI-457 (Pharmaceutical Peptides, Inc)
7/97 - 3/98	Regional Intestinal Absorption of Anti-CMV agents (Bristol-Myers Squibb)
8/97 - 6/98	EM574 absorption in the rabbit in situ (TAP)
7/97 - 9/97	Pharmacokinetics of macrolides in Protocol EM-97-006 (TAP)
11/97 - 12/97	Drug Interaction Pharmacokinetic Analysis (McNeil)
8/97 - 3/98	Analysis and Pharmacokinetics of Macrolides in EM-97-008 (TAP)
10/97 - 12/97	Pharmacokinetics of Slow Release Agents in the CNS (Chiron)
1/98	LC/MS/MS Equipment Grant (TAP)
2/98 - 12/99	Analysis of Macrolides and Metabolites in EM-97-013 (TAP)
3/98 - 12/99	Chemical Stability of Selected Agents (Medtronic)
5/98	Validation of Analysis of Macrolides in Dog Plasma (TAP)
8/98	Validation of Analysis of Macrolides in Rabbit Plasma (TAP)
8/98	Stability of anticancer drugs in solution (Medtronic)
8/98	EM574 toxicokinetics in the dog (TAP)
12/98	EM574 toxicokinetics in the rabbit (TAP)
1/99	Pharmacodynamics of EM574 on LES Pressure (protocol 004)(TAP)
3/99	Effect of Time of Dosing on Absorption of EM574 (protocol 007)(TAP)
4/99	Effect of gastric emptying on the pharmacokinetics of EM574 and its metabolites (protocol 002)(TAP)

2/9	Stability of FUdR and heparin in solution (Medtronic)
2/99	Pharmacodynamics and PKs of EM574 and its metabolites during chronic dosing (protocol 029)(TAP)
3/00 - 8/01	Pharmacokinetics of CDTR and distribution to middle ear fluid (TAP)
8/00 - 6-01	Distribution of ketolides to middle ear fluid (Abbott)
10/01 - 09/03	Pharmacokinetics of Ketolides (Abbott)
12/01 - 11/03	Pharmacokinetics and Distribution of cefdinir (Abbott)
12/02 – 12/03	Effect of a P-Glycoprotein Inhibitor on the Middle Ear Distribution of Clarithromycin (Abbott)
12/02 – 6/04	Distribution a Cephalosporin into Middle Ear Fluid in Children with Otitis Media (H LaRoche)
12/02 – 12/04	Development and Testing of Formulations for Delivery of Antibiotics to the Middle Ear (Abbott)
5/03 - 4/05	A New Approach for the Therapy of Otitis Media (Abbott)
8/04 - 7/05	Distribution of Macrolide antibiotics to tissue sites (Pfizer)
5/05 - 11/05	Testing the Distribution of Amoxicillin into Middle Ear Fluid in the Chinchilla following Pulsatile Dose Administration (Advancis)
1/06 – 9/06	Distribution of Macrolide antibiotics to Pulmonary Tissue and Skeletal Muscle (Pfizer)
9/07 – 10/08	Transtympanic Membrane Delivery of an Antibiotic to the Middle Ear (Alcon)
11/07 – 12/08	Development of an Acute Otitis Media Middle Ear Microdialysis Model in the Chinchilla with Implanted Tympanostomy Tube (Alcon)
1/10 – 12/10	Testing the Penetration of an Antibiotic into Chinchilla Middle Ear using Transtympanic Membrane Delivery Formulations – Phase II (Alcon)
1/11 – 12/11	Testing the Penetration of an Antibiotic into Chinchilla Middle Ear using Transtympanic Membrane Delivery Formulations – Phase II B (Alcon)
1/12 – 6/12	Testing the Penetration of Moxifloxacin into Chinchilla Middle Ear– Phase II, Supplement II (Alcon)

PATENTS

United States Patent. Number 7,220,431 “METHODS AND COMPOSITIONS FOR APPLYING PHARMACOLOGIC AGENTS TO THE EAR.” UMN Docket # Z01159. RJ Sawchuk and BW Cheung. Issue Date: May 22, 2007. Filing Date: November 27, 2002: #06,306,517

PUBLICATIONS

BOOKS AND CHAPTERS

J. Blanchard, R.J. Sawchuk and B.B. Brodie, Editors. Principles and Perspectives in Drug Bioavailability. S. Karger, Basel, 1979.

J. Blanchard and R.J. Sawchuk, "Drug Bioavailability: An Overview" in Principles and Perspectives of Drug Bioavailability, J. Blanchard, R.J. Sawchuk and B.B. Brodie (Editors), S. Karger, Basel, 1979.

I.E. Leppik, J. Shope, R.J. Sawchuk, W.A. Hauser and B. Van Dyne, "Variability of Antiepileptic Drug Levels During Chronic Therapy" in Epileptology, M. Dam, L. Gram and K. Penry (Editors), Raven Press, NY, 1981.

R.J. Sawchuk, "Drug Absorption and Disposition in Burn Patients" in The Pharmacokinetic Basis of Drug Treatment, N. Massoud, L.Z. Benet, and J.G. Gambertoglio (Editors), Raven Press, NY, 1984.

"Use of Microdialysis in Drug Delivery Studies" Theme Issue. W.F. Elmquist and R.J. Sawchuk (Editors) *Advanced Drug Delivery Reviews* 45, Nos. 2-3 (2000).

R. J. Sawchuk and B.W.Y. Cheung, "Application of Microdialysis in Pharmacokinetic Studies" in Handbook of Microdialysis: Methods, Applications and Clinical Aspects, B.H.C. Westerink and T.I.F.H. Cremers (Editors), Academic Press, Amsterdam, 2007.

PROFESSIONAL PUBLICATIONS

R.J. Sawchuk, "The Plateau Principle in Drug Therapy. Part 1: The Principle and Its Application." *Minn. Pharmacist* 27(6): 19 (1973).

R.J. Sawchuk, "The Plateau Principle in Drug Therapy. Part 2: Factors Governing Plateau Levels During Chronic Drug Therapy." *Minn. Pharmacist* 27(7): 8 (1973).

M.C. Meyer, A.B. Straughn, L.J. Leeson, R.H. Levy and R.J. Sawchuk, "Meprobamate Bioavailability Monograph." *JAPhA* NS17, 173 (1977).

R.J. Sawchuk and T.S. Rector, "Burn-Induced Alterations in Drug Absorption and Disposition." *Minn. Pharmacist* 35: 6-9 (1981).

SCIENTIFIC PUBLICATIONS (NON-REFEREED)

I.E. Leppik, J. Cloyd and R.J. Sawchuk, "Coefficient of Variation as a Measure of Compliance" Letter. *Lancet*, October 14, 1978.

SCIENTIFIC PUBLICATIONS (REFEREED)

1. R.J. Sawchuk, J.M. Anderson and J.G. Nairn. "Stirring apparatus for the investigation of unstable strongly adsorbing chemicals." *J. Pharm. Sci.* 55: 1463 (1966).
2. R.J. Sawchuk and J.G. Nairn. "Rate studies on the binding of bilirubin by ion-exchange resins." *J. Pharm. Sci.* 57: 1896 (1968).
3. R.J. Sawchuk, J. Robayo and K.W. Miller. "The distribution of propranolol between blood and plasma in hypertensive patients." *Br. J. Clin. Pharmacol.* 1: 440 (1974).
4. R.J. Sawchuk and D.E. Zaske. "Pharmacokinetics of dosing regimens which utilize multiple intravenous infusions: gentamicin in burn patients." *J. Pharmacokin. Biopharm.* 4: 183 (1976).

5. D.E. Zaske, R.J. Sawchuk, D.N. Gerding and R.G. Strate. "Increased dosage requirements of gentamicin in burn patients." *J. Trauma* 16: (1976).
6. R.J. Sawchuk, D.E. Zaske, R.J. Cipolle, W.A. Wargin and R.G. Strate. "Kinetic model for gentamicin dosing with the use of individual patient parameters." *Clin. Pharmacol. Therap.* 21: 362 (1977).
7. J.C. Cloyd, D.E. Bosch and R.J. Sawchuk. "Concentration-time profile of phenytoin after admixture with small volumes of intravenous fluids." *Am. J. Hosp. Pharm.* 35: 45-48 (1978).
8. J.D. Wirtschafter, C.R. Volk and R.J. Sawchuk. "Trans-aqueous diffusion of acetylcholine to denervated iris sphincter muscle: A hypothetical mechanism for the tonic pupil syndrome (Adie's Syndrome). *Annals of Neurology* 4: 1-5 (1978).
9. D.E. Zaske, R.J. Sawchuk and R.G. Strate. "The necessity of increased doses of amikacin in burn patients." *Surgery* 84: 603-608 (1978).
10. S.M. Ehlers, D.E. Zaske and R.J. Sawchuk. "Massive theophylline overdose: rapid removal by charcoal memoperfusion. *JAMA* 240: 474 (1978).
11. I.E. Leppik, V. Ramani, R.J. Sawchuk and R.J. Gumnit. "Increased clearance of phenytoin during mononucleosis." *NEJM* 300: (1979).
12. H.G. McCoy, R.J. Cipolle, S.M. Ehlers, R.J. Sawchuk and D.E. Zaske. "Severe methanol poisoning: application of a pharmacokinetic model for ethanol therapy and hemodialysis." *Am. J. Med.* 67: 804-807 (1979).
13. R.J. Sawchuk, T.S. Rector, J.J. Fordice and I.E. Leppik. "Effect of influenza vaccination on plasma phenytoin concentrations." *Therap. Drug Monitoring* 1: 285-288 (1979).
14. R.J. Sawchuk and T.S. Rector. "Steady-state plasma concentrations as a function of the absorption rate and dosing interval for drugs exhibiting concentration-dependent clearance: consequences for phenytoin therapy." *J. Pharmacokin. Biopharm.* 7: 543-555 (1979).
15. S. Pancorbo, R.J. Sawchuk, C. Dashe and M. Schallock. "Use of a pharmacokinetic model for individualizing intravenous doses of aminophylline." *Eur. J. Clin. Pharmacol.* 16: 251-254 (1979).
16. I.E. Leppik, J.C. Cloyd, R.J. Sawchuk and S.M. Pepin. "Compliance and variability of plasma phenytoin levels." *Therap. Drug Monitoring* 1: 475-483 (1979).
17. R.L. Kriel, J.C. Cloyd, K.H. Green, R.J. Sawchuk, L.A. Lockman and R. Eggerth. "The pharmacokinetics of valproic acid in children." *Ann. Neurology* 6: 179 (1979).
18. R.J. Sawchuk and L.L. Cartier. "Liquid-chromatographic method for simultaneous determination of phenytoin and 5-(4-hydroxyphenyl)-5-phenylhydantoin in plasma and urine." *Clin. Chem.* 26: 835-839 (1980).
19. N.K. Kouchenour, M. Emery and R.J. Sawchuk. "Phenytoin absorption and metabolism in pregnancy." *Obstetrics and Gynecology* 56: 577-582 (1980).
20. S.E. Chen, R.J. Sawchuk and E.J. Staba. "American ginseng. III. Pharmacokinetics of ginsenosides in the rabbit." *Eur. J. Drug Metab. Pharmacokin.* 5: 161-168 (1980).
21. D. Baker, J.C. Rotschafer, R.J. Sawchuk, K.B. Crossley and L.C. Solem. "Vancomycin pharmacokinetics." *J. Pediatr.* 97: 502-503 (1980).
22. R.J. Sawchuk* and T.S. Rector. "Drug kinetics in burn patients." *Clin. Pharmacokinetics* 5: 548-556 (1980).
23. G.R. Matzke, J.C. Cloyd and R.J. Sawchuk. "Acute phenytoin and primidone intoxication, a pharmacokinetic analysis." *J. Clin. Pharmacol.* 21: 92-99 (1981).

24. El-Yazigi and R.J. Sawchuk. "Theophylline absorption and disposition in the rabbit: oral, intravenous, and concentration-dependent kinetic studies." *J. Pharm. Sci.* 70: 452-456 (1981).
25. J.H. Fischer, J.C. Cloyd, R.L. Kriel, R. Eggerth and R.J. Sawchuk. "The effect of concomitant antiepileptic therapy on valproic acid pharmacokinetics." *Epilepsia* 22: 237 (1981).
26. R.J. Sawchuk and L.L. Cartier. "Liquid-chromatographic determination of cyclosporin A in blood and plasma." *Clinical Chemistry* 27: 1368-1371 (1981).
27. G.R. Matzke and R.J. Sawchuk. "Elevated serum phenytoin concentrations in a uremic patient when measured by enzyme-multiplied immunoassay." *Drug Intell. Clin. Pharm.* 15: 386-387 (1981).
28. R.J. Sawchuk, S.M. Pepin, I.E. Leppik and R.J. Gumnit. "Rapid and slow release phenytoin in epileptic patients at steady state: comparative plasma levels and toxicity." *J. Pharmacokin. Biopharm.* 10: 365-382 (1982).
29. R.J. Sawchuk, S.M. Pepin, I.E. Leppik and R.J. Gumnit. "Rapid and slow release phenytoin in epileptic patients at steady state: assessment of relative bioavailability utilizing Michaelis-Menten parameters." *J. Pharmacokin. Biopharm.* 10: 383-391 (1982).
30. J. Rotschafer, K. Crossley, D. Zaske, K. Mead, R.J. Sawchuk and L.D. Solem. "Pharmacokinetics of vancomycin: observations in 28 patients and dosage recommendation." *Antimicrob. Ag. Chemother.* 22: 391-394 (1982).
31. R.J. Sawchuk and L.L. Cartier. "Liquid-chromatographic method for simultaneous determination of carbamazepine and its epoxide metabolite in plasma." *Clin. Chem.* 28: 2127-2130 (1982).
32. W.A. Wargin, R.J. Sawchuk, J.W. McBride, H.G. McCoy and M.L. Rylander. "Variable first-pass elimination of propranolol following single and multiple oral doses in hypertensive patients." *Eur. J. Drug Metab. Pharmacokin.* 7: 183-189 (1982).
33. J.C. Cloyd, R.L. Kriel, J.H. Fischer, R.J. Sawchuk and R.M. Eggerth. "Pharmacokinetics of valproic acid in children. I. Multiple antiepileptic drug therapy." *Neurology* 33: 185-191 (1983).
34. R.M. Ferguson, R.K. Fidelus, R.J. Sawchuk and K. Gajl. "The mechanism of action of cyclosporine in man." *Transpl. Proc.* 15(1): (1983).
35. D.K. Yu and R.J. Sawchuk. "Gas-liquid chromatographic determination of propylene glycol and plasma and urine." *Clin. Chem.* 29: 2088-2090 (1983).
36. R.J. Sawchuk and G.R. Matzke. "Contribution of 5-(4-hydroxyphenyl)-5-phenylhydantoin to the discrepancy between phenytoin analyses by EMIT and high pressure liquid chromatography." *Ther. Drug Monit.* 6: 97-103, (1984).
37. R.K. Sylvester, B. Lewis, K. Caldwell, M. Lobell, R. Perri and R.J. Sawchuk. "Phenytoin malabsorption secondary to cisplatin, vinblastine and bleomycin." *Ther. Drug Monit.* 6: 302-305 (1984).
38. W. Awni and R.J. Sawchuk. "The pharmacokinetics of cyclosporine. I. Single-dose and constant-rate infusion studies in the rabbit." *Drug Metab. Disposition* 13(2): 127-132 (1984).
39. W. Awni and R.J. Sawchuk. "The pharmacokinetics of cyclosporine. II. Blood-plasma distribution and binding studies." *Drug Metab. Disposition* 13(2): 133-138 (1984).
40. K.H. Chan, R.J. Sawchuk, T.A. Thompson, E. Redalieu, W.E. Wagner, Jr., A.R. LeSher, B.J. Weeks, N.R. Hall and A. Gerardin. "Bioequivalence of carbamazepine chewable and conventional tablets: single dose of steady-state studies." *J. Pharm. Sci.* 74(8): 866-870 (1985).
41. A. El-Yazigi and R.J. Sawchuk. "In vitro - in vivo correlation and dissolution studies with oral theophylline dosage forms." *J. Pharm. Sci.* 74(2): 161-164 (1985).

42. D.K. Yu, W.F. Elmquist and R.J. Sawchuk. "Pharmacokinetics of propylene glycol in humans during multiple dosing regimens." *J. Pharm. Sci.* 74(8): 876-878 (1985).
43. R.J. Sawchuk and W. Awni. "Absorption of cyclosporine from rabbit small intestine *in situ*." *J. Pharm. Sci.* 75(12): 1151-1156 (1986).
44. G.R. Matzke, R.C. Brundage and R.J. Sawchuk. "Protein binding of phenytoin, p-hydroxyphenytoin, and p-hydroxy phenytoin glucuronide." *J. Clin. Pharmacol.* 26: 677-679 (1986).
45. R.M. Ferguson, D.M. Canafax, R.J. Sawchuk and Simmons R.L. Cyclosporine blood level monitoring: the early posttransplant period. *Transplantation Proceedings.* 18(2 Suppl 1):113-22, (1986).
46. C. Fletcher, R.J. Sawchuk, B. Chinnock, P. de Miranda and H.H. Balfour. "Human pharmacokinetics of the antiviral drug DHPG." *Clin. Pharmacol. Therap.* 40: 281-286 (1986).
47. L.E. Riad, K.K.H. Chan, W.E. Wagner and R.J. Sawchuk. "Simultaneous first- and zero-order absorption of carbamazepine tablets in humans." *J. Pharm. Sci.* 75(9): 897-900 (1986).
48. D.K. Yu and R.J. Sawchuk. "Pharmacokinetics of propylene glycol in the rabbit" *J. Pharmacok. Biopharm.* 15: 1-8 (1987).
49. A.K. Shah and R.J. Sawchuk. "Liquid chromatographic determination of cyclosporine and its metabolites in blood." *Clinical Chemistry* 34: 1467-1471 (1988).
50. M.A. Hedaya and R.J. Sawchuk. "A sensitive liquid chromatographic method for the determination of 3'-azido-3'-deoxythymidine (AZT) in plasma and urine." *Clinical Chemistry* 34:1565-1568 (1988).
51. L.E. Riad and R.J. Sawchuk. "A sensitive method for the simultaneous determination of carbamazepine, its epoxide and transdiol metabolites in plasma by microbore liquid chromatography." *Clinical Chemistry* 34: 1863-1866 (1988).
52. M.A. Hedaya and R.J. Sawchuk. "Effect of probenecid on the renal clearance of zidovudine (AZT) and its distribution into cerebrospinal fluid." *J. Pharm. Sci.* 78: 716-722 (1989).
53. R.J. Sawchuk and M.A. Hedaya. "Modeling the enhanced uptake of zidovudine (AZT) into cerebrospinal fluid: the effect of probenecid." *Pharmaceutical Research* 7: 332-338 (1990).
54. M.A. Hedaya, W.F. Elmquist and R.J. Sawchuk. "Probenecid inhibits the metabolic and renal clearance of zidovudine (AZT) in human volunteers." *Pharmaceutical Research* 7: 411-417 (1990).
55. R. Padmanabhan, J.B. Phipps, G.A. Lattin, and R.J. Sawchuk. "*In vitro* and *in vivo* evaluation of transdermal iontophoretic delivery of hydromorphone." *J. Contr. Rel.* 11: 123-135, (1990).
56. M.A. Hedaya and R.J. Sawchuk. "A sensitive and specific liquid-chromatographic assay for determination of ganciclovir in plasma and urine and its application to pharmacokinetic studies in the rabbit." *Pharmaceutical Research* 7: 1113-1118 (1990).
57. W.F. Elmquist, L.E. Riad, I.E. Leppik and R.J. Sawchuk. "The relationship between carbamazepine urine and plasma concentrations: implications for therapeutic drug monitoring." *Pharmaceutical Research* 8: 282-284 (1991).
58. L.E. Riad and R.J. Sawchuk. "Effect of polyethylene glycol 400 on the intestinal permeability of carbamazepine in the rabbit." *Pharmaceutical Research* 8: 491-497 (1991).
59. L.E. Riad, K.K. Chan, and R.J. Sawchuk. "Determination of the relative formation and elimination clearance of two major carbamazepine metabolites in humans: a comparison between traditional and pooled sample analysis." *Pharmaceutical Research* 8: 541-543 (1991).

60. S.L. Wong and R.J. Sawchuk. "High performance liquid chromatographic determination of 2',3,-didehydro-3'-deoxythymidine (D4T) in human and rabbit plasma and urine and its application to pharmacokinetic studies in the rabbit." *Pharmaceutical Research* 8: 619-623 (1991).
61. A.K. Shah and R.J. Sawchuk. "Effect of coadministration of intralipid on the pharmacokinetics of cyclosporine in the rabbit." *Biopharm. Drug Disp.* 12: 457-466 (1991).
62. L.E. Riad and R.J. Sawchuk. "Absorptive clearance of carbamazepine and selected metabolites in rabbit intestine." *Pharmaceutical Research* 8: 1050-1055 (1991).
63. A.K. Shah, R. C. Brundage, A. Gratwohl and R.J. Sawchuk. "Pharmacokinetic model for the subcutaneous absorption of cyclosporine during chronic dosing in the rabbit." *J. Pharm. Sci.* 81: 491-495 (1992).
64. S.L. Wong, M.A. Hedaya and R.J. Sawchuk. "Competitive inhibition of zidovudine clearance by probenecid during continuous co-administration." *Pharmaceutical Research* 9: 228-235 (1992).
65. S.L. Wong, Y. Wang and R.J. Sawchuk. "Analysis of zidovudine distribution to specific regions in rabbit brain using microdialysis." *Pharmaceutical Research* 9: 332-338 (1992).
66. S.L. Wong, K.S. Van Belle and R.J. Sawchuk. "Distributional transport kinetics of zidovudine between plasma and brain extracellular fluid/cerebrospinal fluid in the rabbit: investigation of the inhibitory effect of probenecid utilizing microdialysis." *J. Pharmacol. Exp. Therap.* 264: 899-909 (1993).
67. Y. Wang, S.L. Wong, and R.J. Sawchuk. "Microdialysis calibration using retrodialysis and zero-net flux: application to a study of the distribution of zidovudine to rabbit cerebrospinal fluid and thalamus." *Pharmaceutical Research* 10: 1411-1419 (1993).
68. L.E. Riad and R.J. Sawchuk. "Transient steady state analysis: application in the determination of the relative formation and elimination clearances of two major carbamazepine metabolites in humans." *Pharmaceutical Research* 10: 1090-1092 (1993).
69. E.H. Abdennebi, N. Khales, R.J. Sawchuk and C.M. Stowe. "Thiamphenicol pharmacokinetics in sheep." *J. Vet. Pharmacol. Therap.* 17: 12-16 (1994).
70. R.L. Oberle, H. Das, S.L. Wong, K.K.H. Chan, and R.J. Sawchuk. "Pharmacokinetics and metabolism of diclofenac sodium in Yucatan miniature pigs." *Pharmaceutical Research* 11: 698-703 (1994).
71. B.K. Malhotra, M. Lemaire, and R.J. Sawchuk. "Investigation of the distribution of EAB 515 to cortical ECF and CSF in freely moving rats utilizing microdialysis." *Pharmaceutical Research* 11: 1223-1232 (1994).
72. E.H. Abdennebi, R.J. Sawchuk and C.M. Stowe. "Thiamphenicol pharmacokinetics in beef and dairy cattle." *J. Vet. Pharmacol. Therap.* 17: 365-368 (1994).
73. W.F. Elmquist, K.K.H. Chan, V.A. John and R.J. Sawchuk. "Transsynovial distribution: synovial mean transit time of diclofenac and other non-steroidal antiinflammatory drugs." *Pharmaceutical Research* 12: 1689-1697 (1994).
74. L.E. Riad, R.J. Sawchuk, M.M. MacAlary, and K.K.H. Chan. "Effect of food on the multiple peak behavior of diclofenac sodium slow release tablets in humans." *Amer. J. Therap.* 2: 1-6 (1995).
75. A.K. Shah, R.C. Brundage, K.D. Lake and R.J. Sawchuk. "The estimation of the plasma free fraction of cyclosporine in rabbits and heart transplant patients: the application of a physiologic model of renal clearance." *Biopharm. Drug Disp.* 16: 59-70 (1995).
76. R.J. Sawchuk, J.A. Maloney, L.L. Cartier, R.J. Rackley, K.K.H. Chan and H.S.L. Lau. "Analysis of diclofenac and four of its metabolites in human urine by HPLC." *Pharmaceutical Research* 12: 756-752 (1995).
77. Y. Wang and R.J. Sawchuk. "Zidovudine transport in the rabbit brain during intravenous and intracerebroventricular infusion." *J. Pharm. Sci.* , 84: 871-876 (1995).

78. E.D. Kharasch, M.D. Karol, C. Lanni and R.J. Sawchuk. "Clinical sevoflurane metabolism and disposition; I. Sevoflurane and metabolite pharmacokinetics." *Anesthesiology* 82: 1369-1378 (1995).
79. B.K. Malhotra, M. Lemaire, J.F. Brouillard, and R.J. Sawchuk. "High-performance liquid chromatographic Analysis of (S)- β -amino-5-phosphonomethyl[1,1'-biphenyl]-3-propanoic acid, EAB 515) in brain and blood microdialysate (on-line) and in plasma ultrafiltrate of freely moving rats" *J. Chromatog B: Biomed Appl.* 679: 167-176 (1996).
80. B.W.Y. Cheung, Y. Wang, M. Brewster, and R.J. Sawchuk. "Brain delivery of carbamazepine during intravenous administration of polyethylene glycol and hydroxypropyl- β -cyclodextrin formulations. *s.t.p. pharma sciences* 7: 78-84 (1997).
81. Z. Yang, RC Brundage, R.H. Barbhaiya, and R.J. Sawchuk "Microdialysis studies of the distribution of stavudine into the central nervous system in the freely-moving rat. *Pharmaceutical Research.* 14: 865-72 (1997)
82. J.W. Bartges, C.A. Osborne, L.J. Felice, L.A. Koehler, L.K. Ulrich, K.A. Bird, M. Chen, and R.J. Sawchuk. "Bioavailability and pharmacokinetics of intravenously and orally administered allopurinol in healthy beagles" *American Journal of Veterinary Research.* 58: 504-10 (1997)
83. R. Dvorsky, S. Balaz, and R.J. Sawchuk. "Kinetics of subcellular distribution of compounds in simple biosystems and its use in QSAR. *J. Theoretical Biology.* 185: 213-22 (1997).
84. W.F. Elmquist and R.J. Sawchuk. "Application of microdialysis in pharmacokinetics. (Invited Review) *Pharmaceutical Research* 14: 267-288 (1997).
85. M. Brewster, W.R. Anderson, D. Meinsma, D. Moreno, A.I. Webb, L. Pablo, K.S. Estes, H. Derendorf, N. Bodor, R.J. Sawchuk, B.W.Y. Cheung, and E. Pop. "Intravenous and Oral Pharmacokinetic Evaluation of a 2-hydroxypropyl- β -cyclodextrin-based Formulation of Carbamazepine in the dog: Comparison with Commercially Available Tablets and Suspensions. *J. Pharm. Sci.* 86: 335-339 (1997)
86. B.K. Malhotra, R.C. Brundage, M. Lemaire, and R.J. Sawchuk. "Modeling the route of administration-based enhancement in the brain delivery of EAB 515, studied by microdialysis" *.J Drug Targetting* 4: 277-288 (1997).
87. L.E. Riad and R.J. Sawchuk. "A partial area difference method for estimating elimination rate constants and distribution volumes of metabolites." *J. Pharm Sci.* 87: 769-773 (1998).
88. Z. Yang and R.J. Sawchuk "Investigation of distribution, transport and uptake of anti-HIV Drugs to the central nervous system " *Advanced Drug Delivery Reviews* 39: 5-31 (1999).
89. S.C. Chen, R. J. Sawchuk, R.C. Brundage, C Horvath, H.V. Mendenhall, R.A. Gunther, and R.A. Braeckman. "Plasma and lymph pharmacokinetics of recombinant human interleukin-2 and PEG-modified interleukin-2 in pigs." *JPET* 293: 248-259 (2000).
90. M. Davidson, A. Marwah, R.J. Sawchuk, K. Maki, P. Marwah, C. Weeks, and H. Lardy,. "Safety and pharmacokinetics of escalating doses of 3-acetyl-7-oxo-dehydroepianrosterone in healthy male volunteers" *Clin Invest. Med.* 23: 300-310 (2000).
91. W.F. Elmquist and R.J. Sawchuk. "Use of microdialysis in drug delivery studies." *Advanced Drug Delivery Reviews.* 45: 123-124 (2000).
92. R.J. Sawchuk and W.F. Elmquist. "Microdialysis in the study of drug transporters in the CNS." *Advanced Drug Delivery Reviews.* 45: 295-307 (2000).
93. R.J. Sawchuk D.J. Mulford, and M.D. Mayer. "Pharmacokinetics of a new cephalosporin." *J. Resp. Dis.* 22: No. 8 Suppl. S43-S51 (2001).
94. Y. Huang, P. Ji, A. Inano, Z. Yang, G.S. Giebink, and R.J. Sawchuk. "Microdialysis studies of the middle ear distribution kinetics of amoxicillin in the awake chinchilla." *J. Pharm. Sci.*, 90: 2088-2098 (2001).

95. G. Gan, L.L. Cartier, Y. Huang, Z. Yang, and R.J. Sawchuk. "Intestinal absorption and presystemic elimination of the prokinetic agent, EM574, in the rabbit." *J. Pharm. Sci.*, 91: 218-228 (2002).
96. T. Zhu, B.W.Y. Cheung, L.L. Cartier, G.S. Giebink, and R.J. Sawchuk. "Simultaneous Intravenous and Intramiddle-Ear Dosing to Determine Cefditoren Influx and Efflux Clearances in Middle Ear Fluid in Freely Moving Chinchillas." *J. Pharm. Sci.*, 92: 1947-1956 (2003).
97. J.Z. Peng, R.P. Rimmel, and R.J. Sawchuk. "Inhibition of murine cytochrome P4501A by tacrine: in vitro studies." *Drug Metab Dispos* 32: 805-812 (2004).
98. B.W.Y. Cheung, L.L. Cartier, H.Q. Russlie and R.J. Sawchuk. "The Application of Sample Pooling Methods for Determining AUC, AUMC and Mean Residence Times in Pharmacokinetic Studies" *Fundamental & Clinical Pharmacology*, 19, 347-354 (2005).
99. Z. Yang, Y. Huang, G. Gan, and R. J. Sawchuk. "Microdialysis Evaluation of the Brain Distribution of Stavudine Following Intranasal and Intravenous Administration to Rats." *J Pharm. Sci.* 94, 1577-88 (2005).
100. P. Jacobson, J. Rogosheske, J. N. Barker, K. Green, J. Ng, D. Weisdorf, Y. Tan, J. Long, R. Rimmel, R. Sawchuk, P. McGlave. Relationship of mycophenolic acid exposure to clinical outcome after hematopoietic cell transplantation. *Clin Pharmacol Ther.* 78: 486-500 (2005).
101. R.J. Sawchuk, B.W.Y. Cheung, P. Ji, and L. L. Cartier. "Microdialysis Studies of the Distribution of Antibiotics to Chinchilla Middle Ear Fluid." *Pharmacotherapy* 25 (12): Part 2 140S-145S (2005)
102. B. W. Y. Cheung, W. Liu, P. Ji, L. L. Cartier, Z. Li, N. Mostafa, and R. J. Sawchuk. "The Chinchilla Microdialysis Model for the Study of Antibiotic Distribution to Middle Ear Fluid." *AAPSJ* 8 (1), E41-E47 (2006)
103. Z. Yang, P. Manitpisitkul and R. J. Sawchuk. "In Situ Studies of Regional Absorption of Lobucavir and Ganciclovir from Rabbit Intestine and Predictions of Dose-Limited Absorption and Associated Variability in Humans *J. Pharm. Sci.*, 95: 2276-2292 (2006).
104. Z. Yang, G. Gan and R. J. Sawchuk. "Correlation between Net Water Flux and Absorptive Clearance Determined from In Situ Intestinal Perfusion Studies Does Not Necessarily Indicate a Solvent Drag Effect." *J. Pharm. Sci.*, 96: 517-21 (2007).
105. Chaurasia CS. Muller M. Bashaw ED. Benfeldt E. Bolinder J. Bullock R. Bungay PM. DeLange EC. Derendorf H. Elmquist WF. Hammarlund-Udenaes M. Joukhadar C. Kellogg DL Jr. Lunte CE. Nordstrom CH. Rollema H. Sawchuk RJ. Cheung BW. Shah VP. Stahle L. Ungerstedt U. Welty DF. Yeo H. AAPS-FDA Workshop White Paper: microdialysis principles, application, and regulatory perspectives. *Journal of Clinical Pharmacology*. 47(5): 589-603 (2007).
106. Y. Huang, Z. Yang,, L. Cartier, B. Cheung, and R. J. Sawchuk "Estimating Amoxicillin Influx/Efflux in Chinchilla Middle Ear Fluid and Simultaneous Measurement of Antibacterial Effect" *Antimicrobial Agents and Chemotherapy*, 51 (12): 4336-4341 (2007).
107. J. Karjagin, S. Lefevre, K. Oselin, K. Kipper, S. Marchand, A. Tikkerberi, J. Starkopf, W. Couet and R. J. Sawchuk. "Pharmacokinetics of Meropenem Determined by Microdialysis in the Peritoneal Fluid of Patients With Severe Peritonitis Associated With Septic Shock" *Clin Pharmacol Ther.* 83 (3): 452-459 (2008).
108. D.P. Scher, R. J. Sawchuk, B. H. Alexander, and J. L. Adgate. "Estimating Absorbed Dose of Pesticides in a Field Setting using Biomonitoring Data and Pharmacokinetic Models. *J Toxicol Environ Health A.* 71 (6): 373-83 (2008).
109. R. J. Sawchuk. Liquid-Chromatographic Determination of Cyclosporin A in Blood and Plasma. Citation Classic. *Clin Chem* 54: 2076-2077 (2008).
110. C. Dahyot-Fizelier, S. Lefevre, L. Laksiri, S. Marchand, R. J. Sawchuk, W. Couet, and O. Mimoz. Kinetics of Imipenem Distribution into the Peritoneal Fluid of Patients with Severe Peritonitis Studied by Microdialysis. *Clin Pharmacokinetics* 49(5): 323-334 (2010).

ABSTRACTS

1. R.J. Sawchuk and T.N. Tozer. "Mixed first-order and capacity-limited elimination of 4-aminoantipyrine in the rabbit under steady-state conditions." Fifth International Congress of Pharmacology, Abstracts of Volunteer Papers, p. 202, 1972.
2. D.E. Zaske, R.B. Johnson, M.T. Spiline, R.J. Sawchuk and K.W. Miller. "Utilization of total body clearance calculations in the determination of I.V. dose therapy of theophylline in asthmatic patients." *The Pharmacologist* 15, 206 (1973)
3. R.J. Sawchuk, O.A. Olusanya, D.E. Zaske and K.W. Miller. "Evaluation of the bioequivalence of three theophylline preparations." *Proceedings of the Minnesota Society of Internal Medicine*, p. 2, October 1974.
4. K.W. Miller, P.B. Johnson, D.E. Zaske and R.J. Sawchuk. "Clinical usefulness of measuring serum theophylline levels." *Proceedings of the Minnesota Society of Internal Medicine*, p. 2, October 1974.
5. D. Schuppan, R.J. Sawchuk, C.S. Hansen, M.L. Funk, R.E. Ober and L.A. Fernandez. "Comparative study of plasma levels produced by two formulations of R-802, a new synthetic urinary tract anti-bacterial agent: pharmacokinetics during single and multiple dosing." *APhA Academy of Pharmaceutical Sciences. Abstract. Atlanta, GA, November 1975.*
6. D.E. Zaske, D.N. Gerding, R.G. Strate, R.J. Sawchuk and R.M. Jager. "Rapid elimination of gentamicin in burn patients." *Annual Meeting of the American Burn Association. Abstract. Denver, CO 1975.*
7. M.L. Rylander, W.A. Wargin, R.J. Sawchuk, J.W. McBride, E. Haus and D.E. Zaske. "The clinical efficacy and pharmacokinetics of propranolol in hypertensive patients." *Eleventh Mid-Year Clinical Meeting of the American Society of Hospital Pharmacists, December 1976.*
8. D.L. Uden, D.E. Zaske, R.J. Sawchuk and R.G. Strate. "Gentamicin kinetics in burn patients." *Eleventh Mid-Year Meeting of the American Society of Hospital Pharmacists, December 1976.*
9. D.E. Zaske, R.G. Strate and R.J. Sawchuk. "The necessity of increased doses of amikacin in burn patients." *Eleventh Mid-Year Meeting of the American Society of Hospital Pharmacists, December 1976.*
10. W.A. Wargin, R.J. Sawchuk, R.J. Cipolle, D.E. Zaske and R.G. Strate. "Application of a kinetic model for establishing gentamicin dosing regimens based on individual patient parameters." *American Pharmaceutical Association Academy of Pharmaceutical Services, November 1976.*
11. R.J. Cipolle, R.J. Sawchuk, R.G. Strate and D.E. Zaske. "The clinical application of a kinetic model for individualizing gentamicin dose and dosing interval." *Interscience Conference on Antimicrobial Agents and Chemotherapy, October 1976.*
12. K.W. Miller, R.J. Sawchuk, J.J. Fordice and D.E. Zaske. "The clinical pharmacokinetics laboratory of the University of Minnesota College of Pharmacy." *American Association of Colleges of Pharmacy, July 1976.*
13. R.J. Cipolle, R.J. Sawchuk and D.E. Zaske. "Computer-assisted individualized gentamicin dosing." *The Second Annual Mid-Western Conference for Clinical Pharmacy Residents, Preceptors and Teachers, 1976.*
14. J.W. McBride, M.L. Rylander, W.A. Wargin, R.J. Sawchuk and D.E. Zaske. "Propranolol kinetics in hypertension: a single dose study." *Minnesota Society of Internal Medicine, October 1976.*
15. R.A. Whyte, D.L. Larson, R.J. Sawchuk and P.S. Portoghese. "Relationships of mouse brain-plasma transfer constants of N-substituted normeperidines to their partition coefficients." *Fourteenth Annual Medicinal Chemistry Meeting-in-Miniature, Minneapolis, MN, 1976.*

16. D.E. Zaske, R.G. Strate and R.J. Sawchuk. "Increased dosage requirements of amikacin in burn patients." Ninth Annual American Burn Association, Anaheim, CA, March 1977.
17. H.G. McCoy, R.J. Cipolle, S.M. Ehlers, R.J. Sawchuk and D.E. Zaske. "Severe methanol poisoning: pharmacokinetic basis for successful treatment with ethanol." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
18. J.C. Cloyd, D.E. Bosch and R.J. Sawchuk. "Concentration-time profile of phenytoin after admixture with small volumes of intravenous fluids." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
19. R.J. Sawchuk, S.M. Pepin, J.C. Cloyd, I.E. Leppik and R.J. Gumnit. "Steady-state analysis of phenytoin kinetics applied to dose adjustments in epileptic patients with comparison to nomogram." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
20. G.R. Matzke, J.C. Cloyd, R.J. Sawchuk. "Acute phenytoin and primidone intoxication: A pharmacokinetic analysis." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
21. J.C. Rotschafer, C. Foley, D. Zaske and R. Sawchuk. "Glutethimide overdose: clinical and kinetic observations of glutethimide and a metabolite." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
22. R. Carter, D. Zaske, S. Ehlers and R. Sawchuk. "Charcoal hemoperfusion: an effective method for removal of theophylline in a severe overdose." 12th Annual ASHP Midyear Clinical Meeting, Atlanta, GA, December 1977.
23. R.J. Sawchuk, I.E. Leppik and R.E. Cranford. "Pharmacokinetics of large dose intravenous phenytoin therapy." 11th World Congress of Neurology Meeting, Amsterdam, The Netherlands, September 11-16, 1977.
24. J.D. Wirtschafter, C.R. Volk and R.J. Sawchuk. "Trans-aqueous diffusion of acetylcholine to denervated iris sphincter muscle: a hypothetical mechanism for the tonic pupil syndrome (Adie's Syndrome)." Association of Research in Vision and Ophthalmology, Sarasota, FL, May 1978.
25. W.A. Wargin, M.L. Rylander, R.J. Sawchuk and J.W. McBride. "The pharmacokinetics of orally administered propranolol in hypertensive patients." APhA Academy of Pharmaceutical Sciences, Phoenix, AZ, 1977.
26. R.J. Sawchuk, J.C. Cloyd, I.E. Leppik, S.M. Pepin and R.J. Gumnit. "Steady-state analysis of phenytoin kinetics applied to dose adjustments in epileptic patients." APhA Academy of Pharmaceutical Sciences, Phoenix, AZ, 1977.
27. R.J. Cipolle, R.G. Strate, D.E. Zaske and R.J. Sawchuk. "Amikacin pharmacokinetics: effect of age on elimination." Inter-Science Conference on Antimicrobial Agents and Chemotherapy, New York, 1977.
28. D.E. Zaske, K.B. Crossley, R.J. Sawchuk, D.L. Uden and K.E. Mead. "Vancomycin kinetics: excessive steady-state serum levels resulting from recommended dosages." Interscience Conference on Antimicrobial Agents and Chemotherapy, New York, 1977.
29. D.E. Zaske, R.J. Sawchuk and R.G. Strate. "Increased dosage requirements of amikacin in burn patients." American Burn Assoc., 9th Annual Meeting, Anaheim, CA, April 1977.
30. S.E. Chen, R.J. Sawchuk and E.J. Staba. "Pharmacokinetics of ginseng compounds." 2nd International Ginseng Symposium, Seoul, Korea, 1978.
31. J.C. Cloyd, R.J. Sawchuk, I.E. Leppik and S.M. Pepin. "The direct linear plot: use in estimating Michaelis-Menten parameters and individualizing phenytoin dosage regimens in epileptic patients." Epilepsy International Symposium, Vancouver, Canada, September 10-14, 1978.
32. I.E. Leppik, V. Ramani, R.J. Sawchuk and R.J. Gumnit. "Seizures and altered phenytoin metabolism in mononucleosis." Epilepsy International Symposium, Vancouver, Canada, September 10-14, 1978.
33. I.E. Leppik, J. Cloyd, R. Sawchuk and D. Fryd. "Coefficient of variation as an objective measure of compliance." Epilepsy International Symposium, Vancouver, Canada, September 10-14, 1978.

34. E. Ramsay, I. Leppik, R. Strauss, B.J. Wilder and R.J. Sawchuk. "Plasma clearance and volume of distribution of phenytoin during pregnancy." Epilepsy International Symposium, Vancouver, Canada, September 10-14, 1978.
35. R.L. Kriel, J.C. Cloyd, K.H. Green, R.J. Sawchuk, L.A. Lockman and R.M. Eggerth. "The pharmacokinetics of valproic acid in children." Child Neurology Society, Hanover, NH, September 1979.
36. P. Subronto, C.M. Stowe and R. Sawchuk. "Pharmacokinetics of aminophylline in dairy cattle." Sixtieth Conference of Research Workers in Animal Diseases, Chicago, IL, November 26-27, 1979.
37. I.E. Leppik, J. Shope, R.J. Sawchuk, W.A. Hauser and B. Van Dyne. "Variability of antiepileptic drug levels during chronic therapy." American Academy of Neurology, New Orleans, April 28-May 3, 1980.
38. R.M. Eggerth, J.C. Cloyd, R.L. Kriel and R.J. Sawchuk. "Valproic acid pharmacokinetics in epileptic children." 12th Annual Graduate Student Pharmaceutics Research Meeting, Columbus, OH, June 25, 1980.
39. R. Sylvester, R.J. Sawchuk and R. Perri. "Decreased phenytoin absorption with cis-platinum, vinblastine and bleomycin therapy." American Society of Clinical Oncology Abstracts, 1981.
40. I.E. Leppik, J. Fischer, R. Kriel and R.J. Sawchuk. "Altered phenytoin clearance during febrile illness." American Academy of Neurology, Toronto, Ontario, April 1981.
41. W.M. Awni and R.J. Sawchuk. "Preliminary pharmacokinetic studies of cyclosporin A in animals." Fourteenth Annual Graduate Student Pharmaceutics Research Meeting, Morgantown, WV, June 1982.
42. R.M. Ferguson, R. Fidelus-Gort, J.J. Rynasiewicz, K. Gajl-Peczalska, and R.J. Sawchuk. "The immunosuppressive action of cyclosporin A (CSA) in man." Annual Meeting of the American Society of Transplant Surgery, London, August 1982.
43. J.H. Fischer, I.E. Leppik, R.L. Kriel, J.D. MacDonald and R.J. Sawchuk. "Clinical significance of absorption differences between carbamazepine tablet and suspension formulations." Fourteenth Epilepsy International Symposium, London, August 1982.
44. L.E. Riad and R.J. Sawchuk. "The pharmacokinetics of carbamazepine in the rabbit." The Second University of Minnesota/3M Research Poster Session, Minneapolis, MN, May 1984.
45. W.M. Awni and R.J. Sawchuk. "Cyclosporine pharmacokinetics in the rabbit." The Second University of Minnesota/3M Research Poster Session, Minneapolis, MN, May 1984.
46. W.F. Elmquist, D.K. Yu and R.J. Sawchuk. "Propylene glycol pharmacokinetics." The Second University of Minnesota/3M Research Poster Session, Minneapolis, MN, May 1984.
47. C. Fletcher, R. Sawchuk, B. Chinnock, C. Vicary, L.E. Kirk, P. De Miranda, and H.H. Balfour, Jr. "Human pharmacokinetics of 9-[2-Hydroxy-1-(hydroxymethyl)ethoxymethyl]guanine." Interscience Conference on Antimicrobial Agents and Chemotherapy, Minneapolis, MN, September 28, 1985.
48. R.J. Sawchuk, L. Riad and K.K.H. Chan. "Simultaneous first- and zero-order absorption of carbamazepine (Tegretol) tablets in humans." 16th Epilepsy International Congress, Hamburg, West Germany, September 6, 1985.
49. W.M. Awni and R.J. Sawchuk. "Differential absorption of cyclosporine from two consecutive anatomic regions of the small intestine using an *in situ* perfusion model." Academy of Pharmaceutical Sciences Meeting, Minneapolis, MN, October 20, 1985.
50. H.A. Abou-Auda and R.J. Sawchuk. "Absorption and metabolism of salicylate esters in rabbit small intestine." Academy of Pharmaceutical Sciences Meeting, Minneapolis, MN, October 18-24, 1985.
51. L.E. Riad, K.K.H. Chan and R.J. Sawchuk. "Contribution of epoxide pathway to induction of carbamazepine metabolism during multiple dosing in humans." Academy of Pharmaceutical Sciences meeting, Minneapolis, MN, October 18-24, 1985.

52. L.E. Riad and R.J. Sawchuk. "Contribution of epoxide formation to the auto-induction of carbamazepine in the rabbit." Academy of Pharmaceutical Sciences Meeting, Minneapolis, MN, October 18-24, 1985.
53. R.M. Eggerth and R.J. Sawchuk. "Investigation of the dose-dependent elimination of phenytoin in the rabbit." Academy of Pharmaceutical Sciences Meeting, Minneapolis, MN, October 18-24, 1985.
54. L.E. Riad and R.J. Sawchuk. "Determination of the relative formation of elimination clearance of carbamazepine metabolites in humans: a comparison of traditional and pooled sample analysis." Third European Congress of Biopharmaceutics and Pharmacokinetics, Freiburg, Germany, April 21, 1987.
55. W.F. Elmquist, D.K. Yu, G. Solis and R.J. Sawchuk. "Physiological modelling of the dependence of renal clearance on urine flow: propylene glycol pharmacokinetics in man." Third European Congress of Biopharmaceutics and Pharmacokinetics, Freiburg, Germany, April 21, 1987.
56. A.K. Shah, K.D. Lake and R.J. Sawchuk. "Blood levels of cyclosporine A and its metabolites in rabbit and man." American Association of Pharmaceutical Scientists Annual Meeting, Boston, MA, June 1987.
57. K.D. Lake, Pharm.D. *, A.K. Shah, Ph.D., R. Emery, M.D. and R.J. Sawchuk, Ph.D. "Measurement of cyclosporine metabolites in the blood of heart transplant recipients." American College of Clinical Pharmacy Annual Meeting, May 1987.
58. A.K. Shah, A. Gratwohl and R.J. Sawchuk. "Subcutaneous absorption of cyclosporine in rabbits." 2nd International Congress on Cyclosporine, Washington, DC, November 4-7, 1987.
59. M.A. Hedaya and R.J. Sawchuk. "The effect of probenecid on the renal excretion of azidothymidine (AZT) in the rabbit." Third Annual Meeting, AAPS, Orlando, FL, October 29-November 4, 1988.
60. W.F. Elmquist, K.K.H. Chan, V.A. John and R.J. Sawchuk. "Transsynovial distribution of diclofenac." Third Annual Meeting, AAPS, Orlando, FL, October 29-November 4, 1988.
61. L.E. Riad and R.J. Sawchuk. "Administration of site-related differences in carbamazepine disposition in the rabbit." Third Annual Meeting, AAPS, Orlando, FL, October 29-November 4, 1988.
62. A.K. Shah and R.J. Sawchuk. "Effect of urine flow on the renal clearance of cyclosporine in rabbit." Third Annual Meeting, AAPS, Orlando, FL, October 29-November 4, 1988.
63. W.F. Elmquist, L.E. Riad, I.E. Leppik and R.J. Sawchuk. "Physiological modelling of the dependence of renal clearance on urine flow: carbamazepine and two metabolites in man." Third Annual Meeting, AAPS, Orlando, FL, October 29-November 4, 1988.
64. M.A. Hedaya and R.J. Sawchuk. "Pharmacokinetic analysis of the enhanced distribution of zidovudine into rabbit cerebrospinal fluid caused by probenecid." V. International Conference on AIDS, Montreal, June 4-9, 1989.
65. Y. Wang, M.A. Hedaya and R.J. Sawchuk. "Effect of mannitol IV infusion on distribution of zidovudine between plasma and cerebrospinal fluid in rabbits." V. International Conference on AIDS, Montreal, June 4-9, 1989.
66. M.A. Hedaya, W.F. Elmquist and R.J. Sawchuk. University of Minnesota, Minneapolis, Minnesota, USA. "Probenecid inhibits the metabolic and renal clearances of zidovudine in human volunteers." V. International Conference on AIDS, Montreal, June 4-9, 1989.
67. M.A. Hedaya and R.J. Sawchuk. "A sensitive and specific HPLC method for the determination of zidovudine and zidovudine glucuronide in plasma and urine." V. International Conference on AIDS, Montreal, June 4-9, 1989.
68. M.A. Hedaya and R.J. Sawchuk. "A simple and specific liquid-chromatographic method for determination of ganciclovir (DHPG) in plasma and urine." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
69. L.E. Riad and R.J. Sawchuk. "Intestinal absorption of carbamazepine in the rabbit." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.

70. A.J. Shah and R.J. Sawchuk. "The effect of altering blood lipids on the pharmacokinetics of cyclosporine in the rabbit." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
71. M.A. Hedaya, W.F. Elmquist and R.J. Sawchuk. "Mechanism of zidovudine (AZT) dose-sparing by probenecid in humans." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
72. W.F. Elmquist and R.J. Sawchuk. "Tissue mean transit time determination from steady-state partition coefficients and tissue/plasma concentration ratios in the post-distributive phase." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
73. L.E. Riad and R.J. Sawchuk. "Relative bioavailability and effects of food on plasma levels of diclofenac sodium following a single oral dose of Voltaren 100-mg SR tablet." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
74. O.H. Chan and R.J. Sawchuk. "Intestinal absorption of diltiazem in the rabbit." Fourth Annual Meeting, AAPS, Atlanta, GA, October 21-26, 1989.
75. R.J. Sawchuk. "Pharmacokinetic and analytical considerations in monitoring zidovudine (AZT) levels in children with AIDS." The Fourth International Congress on Pediatric Laboratory Medicine, Washington, DC, August 23, 1989.
76. K.D. Lake, A.K. Shah, R.W. Emery, D.R. Holder and R.J. Sawchuk. "The effect of urine flow rate on the renal clearance of cyclosporine in heart transplant recipients." International Society for Heart Transplantation, Tenth Annual Meeting, Fort Lauderdale, FL, April 4-6, 1990.
77. R.J. Sawchuk. "Effect of temperature and medium of analysis on CsA concentration." Canadian Consensus Meeting on Cyclosporine Monitoring, Minaki Lodge, Ontario, May 10-13, 1990.
78. H. Das, S.W. Wong, K. Chan, R.J. Sawchuk and R.L. Oberle. "Pharmacokinetics and metabolism of diclofenac sodium (Voltaren) after IV and oral administration to Yucatan miniswine." Eastern Regional AAPS meeting, New Brunswick, NJ, June 3-5, 1990.
79. S.M. Wong and R.J. Sawchuk. "Interaction of zidovudine and probenecid at steady state." NATO Advanced Study Institute Symposium "New Trends in Pharmacokinetics." Erice, Sicily, September 4-15, 1990.
80. W.F. Elmquist and R.J. Sawchuk. "Tissue mean transit time determination from steady-state partition coefficients and tissue/plasma concentration ratios in the postdistributive phase." NATO Advanced Study Institute Symposium "New Trends in Pharmacokinetics," Erice, Sicily, September 4-15, 1990.
81. A.K. Shah, R.J. Sawchuk, A. Gratwohl and R.C. Brundage. "Modeling subcutaneous absorption of cyclosporine A in rabbits." NATO Advanced Study Institute Symposium "New Trends in Pharmacokinetics," Erice, Sicily, September 4-15, 1990.
82. L.E. Riad, K.K.H. Chan and R.J. Sawchuk. "Effects of dosing rate on carbamazepine enzymatic induction in normal volunteers." NATO Advanced Study Institute Symposium "New Trends in Pharmacokinetics," Erice, Sicily, September 4-15, 1990.
83. W.F. Elmquist, L.E. Riad, I.E. Leppik and R.J. Sawchuk. "The relationship between carbamazepine urine and plasma concentrations: implications for therapeutic drug monitoring." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
84. S.M. Wong, L.E. Riad, P. Degen, V. John, K. Chan and R.J. Sawchuk. "Comparison of plasma concentrations of diclofenac sodium and its metabolites during single and multiple dosing of Voltaren SR tablet in humans." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
85. O.H. Chan and R.J. Sawchuk. "Saturable first-pass elimination of diltiazem during intestinal absorption in the rabbit." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.

86. L.E. Riad and R.J. Sawchuk. "Carbamazepine enzymatic induction: extent and effects of dosing rate in normal volunteers." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
87. Y.-F. Wang, M.A. Hedaya and R.J. Sawchuk. "Comparative absorption and disposition pharmacokinetics of AZT and AZddU in rabbits." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
88. S.L. Wong and R.J. Sawchuk. "Investigation of distribution of zidovudine into the brain by microdialysis." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
89. S.L. Wong and R.J. Sawchuk. "Interaction of zidovudine and probenecid at steady state." AAPS Fifth Annual Meeting, Las Vegas, NV, November 4-8, 1990.
90. S.L. Wong, Y.-F. Wang and R.J. Sawchuk. "Brain/plasma distribution kinetics of zidovudine (AZT) in the rabbit using microdialysis." 1991 Symposium on Microdialysis and Allied Analytical Techniques, Indianapolis, IN, May 16-17, 1991.
91. K. Van Belle, S.L. Wong, Y.-F. Wang and R.J. Sawchuk. "Comparative delivery and distribution kinetics of zidovudine (AZT) and AZddU into the rabbit brain by microdialysis." 1991 Symposium on Microdialysis and Allied Analytical Techniques, Indianapolis, IN, May 16-17, 1991.
92. Y.-F. Wang, S.L. Wong and R.J. Sawchuk. "*In vitro* and *in vivo* calibration of microdialysis probes using retrodialysis." 1991 Symposium on Microdialysis and Allied Analytical Techniques, Indianapolis, IN, May 16-17, 1991.
93. B.W. Cheung, L.E. Riad and R.J. Sawchuk. "Comparative uptake of selected anticonvulsant drugs into rabbit brain." 1991 Symposium on Microdialysis and Allied Analytical Techniques, Indianapolis, IN May 16-17, 1991.
94. L.E. Riad, K.K. Chan, J.A. Maloney and R.J. Sawchuk. "Reversibility of carbamazepine autoinduction upon dose termination in normal volunteers." AAPS Sixth Annual Meeting, Washington DC, November 17-21, 1991.
95. S.L. Wong, M.A. Hedaya and R.J. Sawchuk. "Modeling the competitive inhibition of renal and nonrenal clearance of zidovudine by probenecid." AAPS Sixth Annual Meeting, Washington, DC, November 17-21, 1991.
96. K. Van Belle, S.L. Wong, Y.-F. Wang and R.J. Sawchuk. "Comparison of pharmacokinetics and CNS distribution of coadministered AZT and AZddU by microdialysis." AAPS Sixth Annual Meeting, Washington, DC, November 17-21, 1991.
97. Y.-F. Wang, S.L. Wong and R.J. Sawchuk. "*In vitro* and *in vivo* calibration of microdialysis probes using retrodialysis: application to the study of brain/plasma distribution of zidovudine." AAPS Sixth Annual Meeting, Washington, DC, November 17-21, 1991.
98. W.F. Elmquist, L.E. Riad, I.E. Leppik and R.J. Sawchuk. "The relationship between urine and plasma concentrations of carbamazepine and phenytoin in epileptic patients on chronic therapy." AAPS Sixth Annual Meeting, Washington, DC, November 17-21, 1991.
99. S.L. Wong, Y.-F. Wang, and R.J. Sawchuk. "Effect of dose on distribution of zidovudine (AZT) into rabbit brain using microdialysis with *in vivo* calibration." AAPS Sixth Annual Meeting, Washington, DC, November 17-21, 1991.
100. W.F. Elmquist, I.E. Leppik and R.J. Sawchuk. "Physiological modeling of the dependence of renal clearance on urine flow II: phenytoin and HPPH in humans." American Association of Pharmaceutical Scientists Annual Meeting, Washington, DC, November 17-21, 1991.
101. R.J. Sawchuk. "Study of zidovudine distribution into the CNS utilizing microdialysis." 25th Annual Higuchi Research Seminar, Lake of the Ozarks, MO, March 8-11, 1992.
102. Y.F. Wang and R.J. Sawchuk. "Microdialysis studies of brain/plasma distribution of AZT during intraventricular and intravenous infusion." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.

103. S.L. Wong, K. Van Belle, and R.J. Sawchuk. "A microdialysis study of probenecid's effect on the transport kinetics of zidovudine in rabbit brain." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
104. S.L. Wong, and R.J. Sawchuk. "Pharmacokinetic interaction of zidovudine and salicylic acid during continuous infusion." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
105. S.L. Wong, and R.J. Sawchuk. "Effect of salicylic acid on the distribution of zidovudine between plasma and cerebrospinal fluid." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
106. R.C. Brundage, K.K.H. Chan, and R.J. Sawchuk. "Population pharmacokinetic modeling of nicotine following transdermal drug administration." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
107. R.C. Brundage, K.K.H. Chan, and R.J. Sawchuk. "Population pharmacokinetics of diclofenac potassium using routinely collected experimental data." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
108. W.F. Elmquist and R.J. Sawchuk. "Simulation of the effect that the time delay in sampling from the bladder has on urine concentrations: implications for therapeutic drug monitoring using urine specimens." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
109. J.P. Zhong, Y.F. Wang and R.J. Sawchuk. "Absorption of three antiviral nucleosides from different anatomic regions of rabbit intestine." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
110. Y.F. Wang, S.L. Wong, and R.J. Sawchuk. "On-line microdialysis calibration using retrodialysis and the zero-net flux method: application to a study of the distribution of AZT to rabbit CSF and thalamus." AAPS Seventh Annual Meeting, San Antonio, TX, November 15-19, 1992.
111. R.J. Sawchuk. "Comparative distribution of AZT into brain tissue extracellular fluid during intravenous and intraventricular infusion using microdialysis." 26th Annual Higuchi Research Seminar, Lake of the Ozarks, MO, March 14-17, 1993.
112. Y. Wang and R.J. Sawchuk, "Comparison of renal clearance of AZdU following IV bolus and constant-rate infusion." 8th Annual Meeting, American Association of Pharmaceutical Scientists, Orlando, FL, November 14-18, 1993.
113. Y. Wang and R.J. Sawchuk. "Assessment of oral absorption of AZT and AZdU in the rabbit using deconvolution." 8th Annual Meeting, American Association of Pharmaceutical Scientists, Orlando, FL, November 14-18, 1993.
114. Y. Wang, Y. Wei and R.J. Sawchuk. "Microdialysis studies of carrier-mediated transport of AZT from brain to plasma during intracerebroventricular infusion." 8th Annual Meeting, American Association of Pharmaceutical Scientists, Orlando, FL, November 14-18, 1993.
115. B.K. Malhotra, M. Lemaire, and R.J. Sawchuk. "Investigation of the CNS distribution of EAB 515 in freely moving rats by microdialysis." 8th Annual Meeting, American Association of Pharmaceutical Scientists, Orlando, FL, November 14-18, 1993.
116. A.K. Shah, R.C. Brundage, K.D. Lake and R.J. Sawchuk. "Estimation of plasma free fraction (fu) of cyclosporine (CYA) in the rabbit and heart transplant (HT) patients by NONMEM using a physiological model of renal clearance (CLr)." 8th Annual Meeting, American Association of Pharmaceutical Scientists, Orlando, FL, November 14-18, 1993.
117. B.K. Malhotra, R.C. Brundage, M. Lemaire, and R.J. Sawchuk. "Modeling of the CNS distribution of EAB 515 following IV and ICV administration." 5th Symposium: Frontiers of Pharmacokinetics and Pharmacodynamics, Baltimore, MD, April 18-20, 1994.
118. R.C. Brundage, B.K. Malhotra, J.A. Maloney and R.J. Sawchuk. "Brain distribution of tacrine and the 1-hydroxy and 2-hydroxy tacrine metabolites determined by microdialysis in freely-moving rats." 5th Symposium: Frontiers of Pharmacokinetics and Pharmacodynamics, Baltimore, MD, April 18-20, 1994.

119. Y. Wang and R.J. Sawchuk. "CNS Distribution of inulin-[¹⁴C]-carboxylic acid in rabbits." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
120. B.K. Malhotra, R.C. Brundage and R.J. Sawchuk. "Simultaneous microdialysis of portal and jugular blood following IV bolus and oral lavage in freely-moving rats." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
121. R.C. Brundage, B.K. Malhotra, J.A. Maloney and R.J. Sawchuk. "Brain distribution of tacrine and its 1- and 2-hydroxylated metabolites determined by microdialysis in freely-moving rats." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
122. B.K. Malhotra, R.C. Brundage, Y. Wang and R.J. Sawchuk. "Dialysis membrane-limited and aqueous boundary layer-limited *in vitro* microdialysis recovery." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
123. B.W.Y. Cheung, Y. Wang and R.J. Sawchuk. "Preliminary studies of effects of hydroxy-propyl-beta-cyclodextrin on carbamazepine distribution into rabbit brain." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
124. R.J. Sawchuk, J.A. Maloney, L.L. Cartier, R.J. Rackley, K.K.H. Chan and H.S.H. Lau. "Analysis of diclofenac and four of its metabolites in human urine by HPLC." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
125. R.C. Brundage, M. Lemaire and R.J. Sawchuk. "Modeling of the CNS distribution of EAB 515 following IV and ICV administration." 9th Annual Meeting, American Association of Pharmaceutical Scientists, San Diego, CA, November 6-10, 1994.
126. B.K. Malhotra, M. Lemaire, J.F. Brouillard and R.J. Sawchuk. "High-performance liquid chromatographic (HPLC) analysis of EAB 515 in brain and blood microdialysate (on-line) and in plasma ultrafiltrate of freely-moving rats." 10th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, November 5-9, 1995.
127. R.J. Sawchuk, R.C. Brundage, E.D. Kharasch and M.D. Karol. "Physiological-based pharmacokinetic (PBPK) modeling of sevoflurane, a volatile anesthetic." 10th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, November 5-9, 1995.
128. R.C. Brundage, S. Thomas Forgue and R.J. Sawchuk. "Comparative distribution of a series of aminoacridines of varying polarities into rat cortex using microdialysis." 10th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, November 5-9, 1995.
129. Z. Yang, R.C. Brundage, L.L. Cartier, J.A. Maloney and R.J. Sawchuk. "Development of a microdialysis method to study brain distribution of stavudine (d4t) in freely-moving rats." 10th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, November 5-9, 1995.
130. B.K. Malhotra, R.C. Brundage and R.J. Sawchuk. "Estimation of presystemic disposition of drugs based upon combination of area ratios and deconvolution." 10th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, November 5-9, 1995.
131. M.A. Osman, R. J. Sawchuk, and M.K. Youssef. "In Situ Absorption of the Antiviral drug, stavudine, from the rabbit intestine." European Symposium on Formulation of Poorly Available Drugs for Oral Administration, Paris, February 5- 6, 1996.
132. M.A. Osman, R.J. Sawchuk, and M.K. Youssef. "Ganciclovir (DHPG), an antiviral nucleoside that exhibits high absorptive clearance in the rabbit colon *in situ*." European Symposium on Formulation of Poorly-available Drugs for Oral Administration, Paris, February 5- 6, 1996.
133. B. K. Malhotra, M. Lemaire, J.F. Brouillard, and R.J. Sawchuk. "High-performance liquid chromatographic (HPLC) analysis of EAB 515 in brain and blood microdialysate (on-line) and in plasma ultrafiltrate of freely-moving rats." 11th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, October 29-31, 1996.

134. R.J. Sawchuk, R.C. Brundage, E.D. Kharasch, and M.D. Karol. "Physiologically-based pharmacokinetic (PBPK) modeling of sevoflurane, a volatile anesthetic." 11th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, October 29-31, 1996.
135. R.C. Brundage, S.T. Fogue, and R.J. Sawchuk. "Comparative distribution of a series of aminoacridines of varying polarities into rat cortex using microdialysis." 11th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, October 29-31, 1996.
136. Z. Yang, R.C. Brundage, L.L. Cartier, J.A. Maloney, and R.J. Sawchuk. "Development of a microdialysis method to study brain distribution of stavudine (d4T) in freely-moving rats." 11th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, October 29-31, 1996.
137. B.K. Malhotra, R. C. Brundage, and R. J. Sawchuk. "Estimation of presystemic disposition of drugs based upon combination of area ratios and deconvolution." 11th Annual Meeting, American Association of Pharmaceutical Scientists, Miami Beach, FL, October 29-31, 1996.
138. Z. Yang and R.J. Sawchuk. "A modified solvent drag model and its application in studying intestinal absorption of polar drugs." 12th Annual Meeting, American Association of Pharmaceutical Scientists, Boston, MA, November 2-6, 1997.
139. Z. Yang and R.J. Sawchuk. "Estimating the intestinal absorptive clearance of drugs: consideration of water absorption during in situ single-pass perfusion studies." 12th Annual Meeting, American Association of Pharmaceutical Scientists, Boston, MA, November 2-6, 1997.
140. Z. Yang, P. Manitpisitkul, F.P. LaCreta, C.K. Knupp, R.H. Barbhैया, and R.J. Sawchuk. "In situ studies of the regional absorption of lobucavir and ganciclovir from rabbit intestine." American Association of Pharmaceutical Scientists Annual Meeting, San Francisco, CA, November 15-19, 1998.
141. Z. Yang, Y. Huang, G. Gan, and R.J. Sawchuk. "Microdialysis evaluation of the brain distribution of stavudine following intranasal administration." American Association of Pharmaceutical Scientists Annual Meeting, San Francisco, CA, November 15-19, 1998.
142. Z Yuan, P. Ji, S. Giebink, and R.J. Sawchuk. "Antibiotic Middle Ear Pharmacokinetics by Microdialysis." American Association of Pharmaceutical Scientists Annual Meeting, San Francisco, CA, November 15-19, 1998.
143. R.J. Sawchuk, P. Ji, Y. Huang. "Distribution of Amoxicillin to Middle Ear Fluid Using Microdialysis Sampling." Higuchi Research Seminar, Lake of the Ozarks, MO, March 14-17, 1999.
144. Y. Huang, and R.J. Sawchuk. "Antibiotic Pharmacokinetics in Chinchilla Middle Ear using Microdialysis." Association of Pharmaceutical Scientists Midwest Regional Meeting, Chicago, IL, May 17, 1999
145. R.J. Sawchuk, P. Ji, Y. Huang, and S. Giebink. "Kinetics of Transport of Antibiotics to Middle Ear Fluid Using Microdialysis Sampling." Seventh International Symposium on Recent Advances in Otitis Media, Fort Lauderdale, FL, June 1-5, 1999.
146. G. Gan, and R.J. Sawchuk. "Intestinal Absorption and Pre-Systemic First Pass Elimination of Minocycline and Propranolol in Rabbits" *American Association of Pharmaceutical Scientists Annual Meeting*, New Orleans, LA, November 14-18, 1999.
147. B.W.Y. Cheung, L.L. Cartier, H. Q. Russlie, and R.J. Sawchuk. "Using Sample Pooling Methods in the Determination of AUC and AUMC in Pharmacokinetic Studies" *American Association of Pharmaceutical Scientists Annual Meeting*, New Orleans, LA, November 14-18, 1999.
148. Y. Huang, L.L. Cartier, and R.J. Sawchuk. "Analysis of Clarithromycin by Chemiluminescence: In Vitro/InVivo Microdialysis Studies" *American Association of Pharmaceutical Scientists Annual Meeting*, New Orleans, LA, November 14-18, 1999.

149. P. Guo, P. Ji, B.W.Y. Cheung, J.B. McCarthy, and R.J. Sawchuk. "Fibronectin Peptide (FN C/H V-Y) Assay and Stability in Human and Rat Plasma" *American Association of Pharmaceutical Scientists Annual Meeting*, New Orleans, LA, November 14-18, 1999.
150. R. J. Sawchuk, B. W. Y. Cheung, L. L. Cartier, H. Q. Russlie, T. Zhu, Y. Huang, G. S. Giebink, D. Mulford and M. Mayer. "Kinetics of Cefditoren Distribution to Middle Ear Fluid in The Unanesthetized Chinchilla" *40th Interscience Conference on Antimicrobial Agents and Chemotherapy*. Toronto, ON, September 17-20, 2000
151. G.S. Giebink, T.M. Sheehy, M. Quartey, R.J. Sawchuk, M. Mayer Cefditoren Pharmacodynamics for Streptococcus Pneumoniae (Pnc) Acute Otitis Media (AOM) in the Chinchilla Model" *40th Interscience Conference on Antimicrobial Agents and Chemotherapy*. Toronto, ON, September 17-20, 2000.
152. Y. Huang, R. J. Sawchuk. "Studies of the Middle Ear Distribution Kinetics of Amoxicillin in the Awake Chinchilla Using Microdialysis". *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
153. J.Z. Peng, R.C. Brundage, and R.J. Sawchuk. "Study of Presystemic Elimination of 4-mono-methylamino-antipyrine (MAAP) after Consecutive Doses in Freely Moving Rats Using On-line Microdialysis". *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
154. T. Zhu, Y. Huang, L.L. Cartier, R. J. Sawchuk "In Vitro Microdialysis and Protein Binding Studies of Cefditoren" *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
155. G. Gan, L. L. Cartier, Y. Huang, Z. Yang, R. J. Sawchuk "Intestinal Absorption and Presystemic Elimination of the Prokinetic Agent, EM574, in Rabbits" *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
156. L. C. Musib, J. C. Cloyd, A.K. Birnbaum, T. J. Hietpas, R. J. Sawchuk, I. E. Leppik, T. R. Browne, S. F. Holloway, G. S. Holden, J. O. Rarig. "Preliminary Report on Phenytoin Bioavailability in the Elderly Using an Intravenous Stable-Labeled Isotope". *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
157. Z. Yang, L. M. Zadjura, C. J. D'Arienzo, D. B. Wang-Iverson, R. J. Sawchuk "Use of Sample Pooling in Drug Discovery to Screen for Pharmacokinetic Properties of Compounds in Rats" *American Association of Pharmaceutical Scientists Annual Meeting*, Indianapolis, IN, October 29- November 2, 2000.
158. R.J. Sawchuk, Y. Huang, P. Ji, M. Quartey, G.S. Giebink. "Influx/Efflux Penetration Clearance of Amoxicillin between Plasma and Middle Ear Fluid in Freely Moving Chinchillas using Microdialysis" *4th Extraordinary International Symposium on Recent Advances in Otitis Media*, Sendai, Japan, April 16-20, 2001.
159. T. Zhu, B. W. Cheung, L.L. Cartier, G. S. Giebink, D.J. Mulford, M.D. Mayer, R.J. Sawchuk. "Study of Cefditoren Distribution Kinetics to Middle Ear Fluid in Freely-moving Chinchillas Using Microdialysis." *American Association of Pharmaceutical Scientists Annual Meeting*, Denver, CO, October 21-25, 2001.
160. W. Liu, B.W. Cheung, L.L. Cartier, T. Zhu, M.M. Paris, R.J. Sawchuk. "In vitro/In vivo Microdialysis and Protein Binding Studies of the ketolide antibiotic, ABT-773." *American Association of Pharmaceutical Scientists Annual Meeting*, Denver, CO, October 21-25, 2001.
161. Y. Huang, R.J. Sawchuk. "Estimation of Amoxicillin Influx and Efflux Clearance between Plasma and Middle Ear Fluid Following Simultaneous Systemic and Local Ear Doses in Awake Chinchilla Using Microdialysis." *American Association of Pharmaceutical Scientists Annual Meeting*, Denver, CO, October 21-25, 2001.
162. J.Z. Peng, R.C. Brundage, R.J. Sawchuk. "The Influence of Drug Pre-exposure on First-pass Metabolism of Tacrine in Rats." *American Association of Pharmaceutical Scientists Annual Meeting*, Denver, CO, October 21-25, 2001.
163. J.Peng, R.J.Sawchuk, and R.P.Remmel "Mechanism-based inactivation of CYP1A2 by tacrine" *11th North American Meeting of the International Society for the Study of Xenobiotics*, Orlando, FL. October 27-31, 2002.

164. Y.Song, L.L.Cartier, B.W Cheung, R J Sawchuk. "An Animal Model for Multi-site CSF Disposition Studies of Intrathecally Administered Agents". *American Association of Pharmaceutical Scientists Annual Meeting*, Toronto, Ontario, Canada. November 10-14, 2002.
165. J.Z.Peng, R.Rommel, R.J.Sawchuk. "Modeling And Simulation Of In Vivo PK Profiles Based On Mechanism-based Inhibition From In Vitro Studies: Inactivation Of CYP1A2 by Tacrine" *American Association of Pharmaceutical Scientists Annual Meeting*, Toronto, Ontario, Canada. November 10-14, 2002.
166. Y.Song, L.L.Cartier, B. Cheung, R.J. Sawchuk. "Multi-site CSF Disposition Studies of Intrathecally Administered Antiviral Nucleosides in a Novel Animal Model". *8th International Meeting of the International Society for the Study of Xenobiotics*, Dijon France. April 27-31, 2003
167. W. Liu, B. W. Y. Cheung, R. J. Sawchuk. "Distribution Kinetics of Cethromycin in the Chinchilla Middle Ear". *8th International Symposium on Recent Advances in Otitis Media*. Fort Lauderdale, FL. June 3 - 7, 2003
168. P Ji, L Cartier, B W Y Cheung, R J Sawchuk. "Distribution Kinetics Of Cefdinir Between Plasma And Middle Ear Fluid In The Freely Moving Chinchillas". *American Association of Pharmaceutical Scientists Annual Meeting*, Salt Lake City, UT. October 26-30, 2003.
169. W. Liu, B. W. Y. Cheung, R J Sawchuk. "Efflux Transport Of Cethromycin Following Direct Intra-bulla Dosing In The Chinchilla Middle Ear". *American Association of Pharmaceutical Scientists Annual Meeting*, Salt Lake City, UT. October 26-30, 2003.
170. Y. Song and R. J. Sawchuk. "Pharmacokinetics of Zidovudine and Stavudine in the Cerebrospinal Fluid after Intrathecal Administration in a Novel Rabbit Model". *American Association of Pharmaceutical Scientists Annual Meeting*, Baltimore, MD. November 8, 2004.
171. N. Mostafa and R. J. Sawchuk. "Determination of Ofloxacin Clearance from the Middle Ear Fluid using Microdialysis". *American Association of Pharmaceutical Scientists Annual Meeting*, Nashville, TN. November 7, 2005.
172. Z. Li, B. W. Y. Cheung, L. Cartier, and R. J. Sawchuk. "The Possible Role of P-glycoprotein in the Distribution of Clarithromycin to the Middle Ear". *American Association of Pharmaceutical Scientists Annual Meeting*, Nashville, TN. November 7, 2005.
173. R. J. Sawchuk, L. M. Page, and R. L. Rauck. "Pharmacokinetics of Gabapentin Injection in Cerebrospinal Fluid and Plasma with Intrathecal Administration". *FDA Science Forum*, Washington, DC. April 18, 2006.
174. R. J. Sawchuk. "Bugs and Drugs: Does the Anti-infective Agent get to the Target Site?" *American Pharmacists Association Annual Meeting*, Atlanta, GA. March 18, 2007.
175. R. J. Sawchuk. "Future Perspectives on the Contributions of Microdialysis in Drug Research and Development" *Fifth International Symposium on Microdialysis in Drug Research and Development*. Leiden, NE. April 25, 2007.
176. R.J. Sawchuk. "Drug Delivery to the Middle Ear across the Tympanic Membrane for Therapy of Acute Otitis Media". *Global Gators 6th Symposium on Clinical Pharmacy and Clinical Pharmacology*. Munich, Germany. June 9, 2007.
177. T. Wang, R. J. Sawchuk and W. F. Elmquist. "Modeling Distributional Kinetics: Comparison of Two Methods Based on a Partial-Areas Analysis". *American Association of Pharmaceutical Scientists Annual Meeting*, San Diego, CA, November 13, 2007.
178. L. Laksiri, C. Dahyot-Fizelier, S. Lefeuvre, S. Marchand, O. Mimoz, R. J. Sawchuk, and W. Couet. "Microdialysis Study of Imipenem Distribution in the Peritoneal Fluid of Patients with Peritonitis". *Interscience Conference on Antimicrobial Agents and Chemotherapy*, Washington, DC. October 26, 2008.
179. R. J. Sawchuk, G. M. Wall, B. W. Y. Cheung, L. L. Cartier, and D. B. Madhura. "Trans-tympanic Membrane Delivery of an Antibiotic into Chinchilla Middle Ear: a Model for the Treatment of Otitis Media". *European Society of Pediatric Infectious Diseases Meeting 2010*, Nice, FR, May 4-8, 2010.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
12/285,887	10/15/2008	John R. Evans	11285.0056-00000	1199

22852 7590 09/16/2011
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

EXAMINER

HUI, SAN MING R

ART UNIT	PAPER NUMBER
1628	

MAIL DATE	DELIVERY MODE
09/16/2011	PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Office Action Summary	Application No. 12/285,887	Applicant(s) EVANS ET AL.	
	Examiner SAN-MING HUI	Art Unit 1628	

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) OR THIRTY (30) DAYS, WHICHEVER IS LONGER, FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) Responsive to communication(s) filed on 20 June 2011.
- 2a) This action is **FINAL**. 2b) This action is non-final.
- 3) An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.
- 4) Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

Disposition of Claims

- 5) Claim(s) 24-53 is/are pending in the application.
- 5a) Of the above claim(s) _____ is/are withdrawn from consideration.
- 6) Claim(s) _____ is/are allowed.
- 7) Claim(s) 24-53 is/are rejected.
- 8) Claim(s) _____ is/are objected to.
- 9) Claim(s) _____ are subject to restriction and/or election requirement.

Application Papers

- 10) The specification is objected to by the Examiner.
- 11) The drawing(s) filed on _____ is/are: a) accepted or b) objected to by the Examiner.
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
- 12) The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

Priority under 35 U.S.C. § 119

- 13) Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
- a) All b) Some * c) None of:
1. Certified copies of the priority documents have been received.
2. Certified copies of the priority documents have been received in Application No. _____.
3. Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

* See the attached detailed Office action for a list of the certified copies not received.

Attachment(s)

- | | |
|--|---|
| 1) <input checked="" type="checkbox"/> Notice of References Cited (PTO-892) | 4) <input type="checkbox"/> Interview Summary (PTO-413)
Paper No(s)/Mail Date. _____ |
| 2) <input type="checkbox"/> Notice of Draftsperson's Patent Drawing Review (PTO-948) | 5) <input type="checkbox"/> Notice of Informal Patent Application |
| 3) <input type="checkbox"/> Information Disclosure Statement(s) (PTO/SB/08)
Paper No(s)/Mail Date _____ | 6) <input type="checkbox"/> Other: _____ |

DETAILED ACTION

This is a continuation of US 10/872,784, filed 6/22/2004, now patent 7,456,160, which is a continuation of US 09/756,291, filed 1/9/2001, now patent 6,774,122. The instant application also claims the benefit of UNITED KINGDOM 0000313.7, filed 01/10/2000 and UNITED KINGDOM 0008837.7, filed 04/12/2000.

Applicant's amendments filed 6/20/2011 have been entered. Claims 24-53 are pending.

The outstanding rejection under 35 USC 103(a) is withdrawn due to the cancellation of the claims.

Double Patenting

The nonstatutory double patenting rejection is based on a judicially created doctrine grounded in public policy (a policy reflected in the statute) so as to prevent the unjustified or improper timewise extension of the "right to exclude" granted by a patent and to prevent possible harassment by multiple assignees. A nonstatutory obviousness-type double patenting rejection is appropriate where the conflicting claims are not identical, but at least one examined application claim is not patentably distinct from the reference claim(s) because the examined application claim is either anticipated by, or would have been obvious over, the reference claim(s). See, e.g., *In re Berg*, 140 F.3d 1428, 46 USPQ2d 1226 (Fed. Cir. 1998); *In re Goodman*, 11 F.3d 1046, 29 USPQ2d 2010 (Fed. Cir. 1993); *In re Longi*, 759 F.2d 887, 225 USPQ 645 (Fed. Cir. 1985); *In re Van Ornum*, 686 F.2d 937, 214 USPQ 761 (CCPA 1982); *In re Vogel*, 422

Art Unit: 1628

F.2d 438, 164 USPQ 619 (CCPA 1970); and *In re Thorington*, 418 F.2d 528, 163 USPQ 644 (CCPA 1969).

A timely filed terminal disclaimer in compliance with 37 CFR 1.321(c) or 1.321(d) may be used to overcome an actual or provisional rejection based on a nonstatutory double patenting ground provided the conflicting application or patent either is shown to be commonly owned with this application, or claims an invention made as a result of activities undertaken within the scope of a joint research agreement.

Effective January 1, 1994, a registered attorney or agent of record may sign a terminal disclaimer. A terminal disclaimer signed by the assignee must fully comply with 37 CFR 3.73(b).

Claims 24-53 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-9 of U.S. Patent No. 6,774,122 ('122). Although the conflicting claims are not identical, they are not patentably distinct from each other because '122 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of the solvents and the excipients are within the range taught in '122. The optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition

Art Unit: 1628

in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980).

It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

Claims 24-53 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-12 of U.S. Patent No. 7,456,160 ('160). Although the conflicting claims are not identical, they are not patentably distinct from each other because '160 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of the solvents and the excipients are within the range taught in '160. The optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

Claim Rejections - 35 USC § 103

The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

Claims 24-53 are rejected under 35 U.S.C. 103(a) as being unpatentable over McKeskey et al., from IDS filed 6/20/2011 in view of Dukes (EP 0 346 014), Osborne et al., Journal of National Cancer Institute, 1995;87(20):746-750, and the abstract of Wakeling et al., The Journal of Steroid Biochemistry and Molecular Biology, 1992;43(1-3):173-177.

McKeskey et al. teaches a studies employing subcutaneous injection of fulvestrant to nude mice. The fulvestrant formulation contains 50mg/ml in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol brought to volume with castor oil (see page 698, col. 2, Drugs section).

McKeskey et al. does not expressly teach the use of fulvestrant in treating hormonal dependent diseases of breast. It does not expressly teach the dosing regimen to be once a month, intramuscular administration, or the volume administered. McKeskey et al. does not expressly teach the herein claimed serum concentration of fulvestrant.

Dukes teaches antiestrogen agents, including fulvestrant, are useful in treating postmenopausal symptoms such as urogenital atrophy affecting the vagina (See page 3, lines 56-page 4, line 1; also page 7, line 28-29). Dukes teaches that antiestrogen

Art Unit: 1628

agent, including fulvestrant, via intramuscular route of administration may be used in a dosage of 50mg to 5g in vehicle comprising castor oil and benzyl alcohol (See page 7, line 20-24).

Osborne et al. teaches fulvestrant as useful in treating human breast cancer (See pages 747- 748, Result Section).

Wakeling et al. teaches the administration of fulvestrant (ICI 182780) demonstrating the antiestrogenic effect for over a 1 month period (see the abstract).

It would have been obvious to one of ordinary skill in the art at the time the invention was made to employ fulvestrant in McKeskey's, in the herein claimed dosing regimen and dosage, for treating hormonal dependent diseases such as breast cancer and postmenopausal symptoms.

One of ordinary skill in the art would have been motivated to employ fulvestrant in McKeskey's, in the herein claimed dosing regimen and dosage, for treating hormonal dependent diseases such as breast cancer and postmenopausal symptoms. It is known in the art that fulvestrant as being useful in treating hormonal dependent disease. It is also known in the art that fulvestrant can be administered intramuscularly and its antitumor effect can last for more than 1 month. Employing McKeskey's formulation of fulvestrant for intramuscular administration would be seen as obvious since administering a relative large volume of fulvestrant (5ml) would not be appropriate for subcutaneous administration. The examiner notes that in McKeskey's study, only 0.1ml was injected via subcutaneous administration. Furthermore, the optimization of result effect parameters (e.g., dosing regimen, weight ratio of the actives and the excipients) is

obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

The examiner notes that the herein claimed serum concentration is considered to be an inherent effect of the formulation of fulvestrant.

Response to Arguments

Applicant's arguments with respect to claims 24-53 have been considered but are moot in view of the new ground(s) of rejection.

Applicant's amendment necessitated the new ground(s) of rejection presented in this Office action. Accordingly, **THIS ACTION IS MADE FINAL**. See MPEP § 706.07(a). Applicant is reminded of the extension of time policy as set forth in 37 CFR 1.136(a).

A shortened statutory period for reply to this final action is set to expire **THREE MONTHS** from the mailing date of this action. In the event a first reply is filed within **TWO MONTHS** of the mailing date of this final action and the advisory action is not

mailed until after the end of the THREE-MONTH shortened statutory period, then the shortened statutory period will expire on the date the advisory action is mailed, and any extension fee pursuant to 37 CFR 1.136(a) will be calculated from the mailing date of the advisory action. In no event, however, will the statutory period for reply expire later than SIX MONTHS from the date of this final action.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SAN-MING HUI whose telephone number is (571)272-0626. The examiner can normally be reached on Mon - Fri from 9:00 to 5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Brandon Fetterolf can be reached on (571) 272-2919. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

San-ming Hui

Application/Control Number: 12/285,887
Art Unit: 1628

Page 9

Primary Examiner
Art Unit 1628

/San-ming Hui/
Primary Examiner, Art Unit 1628

Notice of References Cited	Application/Control No. 12/285,887	Applicant(s)/Patent Under Reexamination EVANS ET AL.	
	Examiner SAN-MING HUI	Art Unit 1628	Page 1 of 1

U.S. PATENT DOCUMENTS

*	Document Number Country Code-Number-Kind Code	Date MM-YYYY	Name	Classification
	A US-			
	B US-			
	C US-			
	D US-			
	E US-			
	F US-			
	G US-			
	H US-			
	I US-			
	J US-			
	K US-			
	L US-			
	M US-			

FOREIGN PATENT DOCUMENTS

*	Document Number Country Code-Number-Kind Code	Date MM-YYYY	Country	Name	Classification
	N				
	O				
	P				
	Q				
	R				
	S				
	T				

NON-PATENT DOCUMENTS

*	Include as applicable: Author, Title Date, Publisher, Edition or Volume, Pertinent Pages)
U	The abstract of Wakeling et al., The Journal of Steroid Biochemistry and Molecular Biology, 1992;43:1-3:173-177
V	
W	
X	

*A copy of this reference is not being furnished with this Office action. (See MPEP § 707.05(a).)
Dates in MM-YYYY format are publication dates. Classifications may be US or foreign.

Comparison of the Effects of a Pure Steroidal Antiestrogen With Those of Tamoxifen in a Model of Human Breast Cancer

C. Kent Osborne, Ester B. Coronado-Heinsohn, Susan G. Hilsenbeck, Bryant L. McCue, Alan E. Wakeling, Richard A. McClelland, David L. Manning, Robert I. Nicholson*

Background: Tamoxifen, a nonsteroidal estrogen antagonist, is the most prescribed drug for the treatment of breast cancer. The use of tamoxifen is limited, however, by the development of resistance to this compound in most patients. Although tamoxifen behaves primarily as an estrogen antagonist, it has agonist (or growth-stimulatory) activity as well. ICI 182,780 is a 7 α -alkylsulfinyl analogue of estradiol lacking agonist activity. The absence of agonist activity may make this steroidal antiestrogen superior to tamoxifen in suppressing tumor cell growth. **Purpose:** We compared the inhibitory effects of ICI 182,780, tamoxifen, and estrogen withdrawal on the growth of established tumors and on tumorigenesis in a model system that uses estrogen-dependent, human MCF-7 breast tumor cells growing in athymic nude mice. We also studied the hormonal responsiveness of tumors that became resistant to the two estrogen antagonists and the effects of these drugs on estrogen-regulated gene expression. **Methods:** MCF-7 cells were injected subcutaneously into the flanks of castrated, female nude mice. The effects of repeated doses of tamoxifen and ICI 182,780 (500 μ g and 5 mg, respectively) on the growth of established tumors (8-10 mm in size) were determined after supplemental estrogen was removed. The effects of antiestrogen treatments on the process of tumorigenesis, in the absence of estrogen supplementation, were determined by initiating drug administration on

the same day as tumor cell inoculation. To evaluate the hormonal responsiveness of tumors resistant to tamoxifen and ICI 182,780, 1-mm³ segments of the tumors were transplanted onto the flanks of new recipient mice, which were then treated with estrogen or the antiestrogens—alone or in combination. Tumor growth was monitored by measuring tumor volumes twice a week. Expression of the estrogen-responsive genes, pLIV1 and pS2, in the tumors of treated animals was analyzed using blots of total cellular RNA and complementary DNA probes. **Results:** Treatment with ICI 182,780 suppressed the growth of established tumors twice as long as treatment with tamoxifen or estrogen withdrawal. Tumorigenesis, in the absence of supplemental estrogen, was delayed to a greater extent in ICI 182,780-treated mice than in tamoxifen-treated mice. ICI 182,780 was found to be more effective than tamoxifen in reducing the expression of estrogen-regulated genes. Most tumors eventually became resistant to ICI 182,780 and grew independently of estrogen. **Conclusions:** ICI 182,780 is a more effective estrogen antagonist than tamoxifen in the MCF-7 tumor cell/nude mouse model system. [J Natl Cancer Inst 87:746-750, 1995]

Tamoxifen, a nonsteroidal antiestrogen, is the most prescribed drug for the treatment of breast cancer. When used in the adjuvant setting after surgery for primary breast cancer, about one fifth of the deaths at 10 years are avoided by 2 years or more of treatment (1). Tamoxifen is also effective in inducing remissions in women with estrogen receptor (ER)-positive metastatic breast cancer. Invariably, however, tumors become resistant to tamoxifen, and tumor progression and death ensue. The evolution to tamoxifen resistance in metastatic breast cancer occurs after an average treatment duration of only 10-12 months, severely limiting the usefulness of this approach.

The mechanisms by which tumors acquire resistance to tamoxifen are poorly understood. Loss of ER from the tumor can occur by selection of an ER-negative clone or by suppression of receptor expression, but this loss explains only a minority of cases (2). Growing experi-

mental and clinical evidence suggests that resistance in some patients may be caused by the intrinsic estrogen agonist properties of tamoxifen. Although tamoxifen is predominantly an estrogen antagonist in breast cancer cells, acquisition of increasingly dominant agonist activity over time may result in clinical resistance because of the acquired ability of the drug to stimulate, rather than to inhibit, tumor growth (3-7). The mechanisms for tamoxifen-stimulated tumor growth are not clear, but these data suggest that antiestrogens with pure antagonist properties might have superior antitumor activity.

ICI 182,780 is a 7 α -alkylsulfinyl analogue of estradiol that differs substantially from tamoxifen in terms of its chemical, pharmacologic, and biologic properties. This agent has no intrinsic estrogen-agonist activity and, thus, is considered a "pure" antiestrogen (8,9). It has potent antiestrogenic activity in preclinical in vitro and in vivo model systems (10). We recently reported (7) that treating nude mice with ICI 182,780 inhibits the growth of MCF-7 human breast tumor implants that had acquired tamoxifen resistance through the mechanism of tamoxifen-stimulated growth. Similar results were obtained with another analogue, ICI 164,384, studied earlier (11). These data suggest the possibility that pure steroidal antiestrogens may be effective in some tamoxifen-resistant patients.

In the present study, we have investigated the preclinical activity of ICI 182,780 in more detail. We compared the inhibitory effects of ICI 182,780, tamoxifen, and estrogen withdrawal on the

*Affiliations of authors: C. K. Osborne, E. B. Coronado-Heinsohn, S. G. Hilsenbeck, B. L. McCue, Department of Medicine, Division of Medical Oncology, The University of Texas Health Science Center at San Antonio, San Antonio, Tex.

A. E. Wakeling, Zeneca Pharmaceuticals, Mereside Alderley Park, Macclesfield, Cheshire, England, U.K.

R. A. McClelland, D. L. Manning, R. I. Nicholson, Tenovus Cancer Research Center, University of Wales College of Medicine, Heath Park, Cardiff, Wales, U.K.

Correspondence to: C. Kent Osborne, M.D., Department of Medicine, Division of Medical Oncology, The University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr., San Antonio, TX 78284-7884.

See "Notes" section following "References."

growth of established tumors and on tumorigenesis in a model system that uses estrogen-dependent, human MCF-7 breast tumor cells growing in athymic nude mice. We also studied the hormonal responsiveness of tumors that had become resistant to the two estrogen antagonists and the effects of these drugs on estrogen-regulated gene expression.

Materials and Methods

Nude Mouse Model System

ER-positive MCF-7 human breast cancer cells (passage 100-200) were cultured as described previously (12). The athymic nude mice used in these experiments were 4- to 5-week-old female castrated BALB/c-nu⁺/nu⁺ mice purchased from Harlan Sprague-Dawley, Inc. (Madison, Wis.). The methods for maintenance and housing of the mice and for growing MCF-7 tumors from cell suspensions and from tumor transplants have been published in detail (3,7). Animal care was in accordance with institutional guidelines.

Approximately 5×10^5 MCF-7 cells were injected subcutaneously into the flanks, just under the forelimb, of female nude mice to initiate tumor formation. Estrogen supplementation was provided in the form of a 0.25-mg estradiol (E₂) pellet (Innovative Research, Rockville, Md.) placed subcutaneously in the interscapular region of the mice. The effects of tamoxifen and ICI 182,780 on the growth of established tumors were studied after the tumors had reached a size of 8-10 mm (3-5 weeks). At this time, the animals were randomly allocated into four treatment groups: 1) continued estrogen supplementation, 2) removal of the E₂ pellet, 3) removal of the E₂ pellet plus treatment with 500- μ g tamoxifen citrate (Zeneca Pharmaceuticals, Wilmington, Del.) in peanut oil (injected subcutaneously each day, Monday through Friday), or 4) removal of the E₂ pellet and treatment with the indicated doses of ICI 182,780 (Zeneca Pharmaceuticals, Macclesfield, England) in castor oil (subcutaneous injections once a week). Initial dose-response studies with ICI 182,780 were performed in the presence of continued estrogen supplementation. Tumor growth was assessed, and tumor volumes were measured twice a week as described previously (12).

In tumorigenesis experiments, various treatments were begun on the same day tumor cells were injected. Inoculated mice were randomly allocated immediately into four treatment groups: 1) estrogen supplementation, 2) 500 μ g tamoxifen once a day, Monday through Friday, 3) 5 mg ICI 182,780 once a week, or 4) drug vehicle (peanut oil and/or castor oil). Tumor volumes were measured twice a week.

To investigate the hormonal responsiveness of tumors that had become resistant to ICI 182,780, mice with resistant tumors were killed by cervical dislocation, and the tumors were resected and cut into 1-mm³ fragments. The fragments were then transplanted subcutaneously on the flank just under the forelimb of new 4- to 5-week-old recipient mice that were then treated with estrogen, tamoxifen, ICI 182,780, or vehicle alone.

Estrogen and Progesterone Receptor Assays

ER content was determined in tumors homogenized in 0.4 M KCl-Tris buffer, using the ER antibody kit (ER-ELA; Abbott Laboratories, North Chicago, Ill.). Progesterone receptor (PgR) levels were measured by a ligand-binding, dextran-coated charcoal method (3).

Estrogen-Regulated Gene Expression

Expression of the estrogen-responsive genes, pLIV1 and pS2, was determined by northern blot analysis, using complementary DNA (cDNA) probes labeled with [³²P]deoxycytidine triphosphate (3000 Ci/mmol; Amersham Ltd., Amersham, England, U.K.) by the random-priming method as described previously (13). Briefly, total RNA was obtained from the tumors of treated mice by cell lysis in 4 M guanidinium thiocyanate and 1% 2-mercaptoethanol and centrifugation through 5.7 M caesium chloride (Beckman L-80 ultracentrifuge, SW50 rotor, 34 000 rpm at 20 °C for 17 hours). Purified samples were stored in RNase-free water at -70 °C before electrophoresis (10 μ g/lane), blotting, and hybridization. Densitometric analysis of autoradiographs was performed using a model 620 video densitometer (Bio-Rad Laboratories, Richmond, Calif.), and values obtained were corrected for equivalence of RNA loading by comparison with the signals generated using a cDNA probe to human glyceraldehyde 3-phosphate dehydrogenase (G3PDH) (Clontech Laboratories, Inc., Palo Alto, Calif.).

Recorded densitometry values represent the area of peak values obtained, following background subtraction, from equivalently exposed autoradiographs (where x = band width in mm and y = optical density value). Hybridizations of each set of filters in the study were carried out simultaneously with the same labeled probes. The reported values represent means of groups, and at least two separate hybridizations of different filters were performed for each probe (stripping the previous probe with high-stringency washes and checking for clearance by autoradiography).

Statistical Analysis

Analyses were performed using either the Kruskal-Wallis one-way analysis of variance (when there were more than two groups) or the Wilcoxon signed rank test for two samples. All statistical tests were two-sided.

Results

ICI 182,780 Dose-Response

ICI 182,780 inhibited estrogen-induced growth of MCF-7 tumors in a dose-dependent manner. Estrogen-supplemented mice with established MCF-7 tumors were randomly allocated to receive either continued estrogen treatment or estrogen treatment plus injections of ICI 182,780 once a week in doses ranging from 0.5

mg to 10.0 mg. Inhibitory activity was modest with doses of 0.5 mg or 1.0 mg, while more dramatic—but approximately equivalent—inhibitory effects were observed with 5.0-mg and 10.0-mg doses (data not shown). For subsequent experiments, a dose of 5.0 mg per mouse, given once a week, was used.

Effect of Estrogen Withdrawal, Tamoxifen, and ICI 182,780 on MCF-7 Tumor Growth

Treatment of mice by removal of the E₂ pellet alone or with tamoxifen or ICI 182,780 significantly inhibited MCF-7 tumor growth (Fig. 1). In this experiment, tumor volumes remained stable for nearly 100 days after estrogen withdrawal before progression ensued. In contrast, tumor volumes decreased slightly with tamoxifen and ICI 182,780 treatment, and tumor size remained stable for variable periods of time. A consistent observation was the delayed time to progression that was evident in mice treated with ICI 182,780. With estrogen withdrawal alone or with tamoxifen, tumors developed resistance, and progression was evident in all mice after 3-4 months of treatment (median, 97 and 104 days, respectively). However, the median time to progression was nearly twice as long with ICI 182,780, and the growth of some tumors remained controlled for extended periods of time (median, 200 days). In fact, two of the 10 tumors from ICI 182,780-treated mice still had not progressed after 11 months and one small tumor (4 mm diameter) completely regressed and did not reappear during the course of the experiment (data not shown).

Effect of ICI 182,780 on Tumorigenesis

ICI 182,780 also had a greater impact on tumor formation in mice in which drug treatments were begun on the day of tumor cell inoculation (Fig. 2). Tumors grew rapidly in mice treated with estrogen. Tumor growth was substantially delayed in mice treated with tamoxifen, but after 2 months, the growth rate increased. Tumors grew very slowly, or not at all, in mice treated with ICI 182,780—similar to the growth pattern observed in estrogen-deprived mice (12). By day 70, barely measurable tumors were present in the majority of mice. In another experiment (data not shown), three of six mice




Fig. 1. Effects of estrogen (estradiol [E₂]) withdrawal, tamoxifen, and ICI 182,780 on MCF-7 tumor growth. Estrogen-supplemented mice were inoculated with MCF-7 cells. On day 36 when tumors had formed, mice were randomly allocated to treatment by withdrawal of estrogen (-E₂; -○-); withdrawal of estrogen and treatment with 500 μg tamoxifen given once a day, Monday through Friday (-◆-); or 5 mg ICI 182,780 given once a week (-■-). Tumor volumes were determined at the times shown. n = 10 mice per group; means ±SE.




Fig. 2. Effect of estrogen, tamoxifen, and ICI 182,780 on MCF-7 tumorigenesis. Mice were inoculated with MCF-7 cells on day 0 and randomly allocated immediately to receive treatment with a 17β estradiol (E₂) pellet (+E₂; -○-); 500 μg tamoxifen given once a day, Monday through Friday (-◆-); or 5 mg ICI 182,780 given once a week (-■-). Tumor volumes were determined at the times shown. n = 8 mice per group; means ±SE.

treated with ICI 182,780 failed to grow measurable tumors even after 6 months of treatment.

ICI 182,780-Resistant Tumors

As indicated above, tumor resistance eventually occurred in most, but not all,

mice treated with ICI 182,780. This resistance was manifested by regrowth of tumors, usually after many months of treatment. To investigate the hormonal sensitivity of these resistant tumors, fragments of a tumor that had progressed after months of treatment with

ICI 182,780 were transplanted into new castrated, recipient mice that were then treated with estrogen, tamoxifen, ICI 182,780, tamoxifen plus ICI 182,780, or vehicle alone. This experiment was conducted five times with different tumor transplants, and a representative result is shown in Fig. 3. Transplanted tumor fragments grew well in all mice, even those treated with vehicle alone (-E₂), suggesting estrogen independence. However, in four of five experiments, tumor growth was slightly increased by estrogen treatment (+E₂), indicating continued sensitivity to the hormone. As expected, growth of these transplanted ICI 182,780-resistant tumors was also observed in recipient mice treated with ICI 182,780.

Interestingly, in four of the five experiments, treatment of recipient mice with tamoxifen alone or tamoxifen plus ICI 182,780 resulted in a slight retardation of tumor growth compared with treatment using vehicle alone or ICI 182,780 alone, although the observed differences in the individual experiments were modest and not statistically significant. A total of six of the 25 mice in these experiments showed slower tumor growth with tamoxifen treatment, indicating some heterogeneity among the transplanted fragments in response to tamoxifen. However, most mice resistant to ICI 182,780 showed cross-resistance to tamoxifen.

Resistance to ICI 182,780 was not due to a complete loss of tumor ER, although treatment with this drug reduced expression of both ER and PgR. Tumors harvested 4 weeks after initiating treatment with ICI 182,780 (ER = 37 ± 3 fmol/mg protein; PgR = 27 ± 7 fmol/mg protein) as well as those harvested at the time of resistance to ICI 182,780 (ER = 16 ± 4 fmol/mg protein; PgR = 17 ± 8 fmol/mg protein) expressed both ER and PgR at markedly reduced levels compared with estrogen-treated controls (ER = 208 ± 81 fmol/mg protein; PgR = 103 ± 20 fmol/mg protein) (*P* = .024).

Expression of two estrogen-responsive genes, pS2 and pLIV1, was also measured in these tumors (Table 1). pS2 and pLIV1 messenger RNA (mRNA) expression was reduced by 20%-74% in tumors from tamoxifen-treated mice (*P* = .013). It is interesting that pS2 and pLIV1 expression remained suppressed even after




Fig. 3. Hormonal sensitivity of ICI 182,780-resistant tumors. Fragments of a tumor that had developed resistance in a mouse treated with ICI 182,780 were transplanted into new recipient female castrated nude mice. The recipient mice were then randomly allocated to receive vehicle alone (-E₂; -□-), an estradiol (E₂) pellet (+E₂; -◇-); tamoxifen alone (-▲-); ICI 182,780 alone (-◆-), or a combination of tamoxifen plus ICI 182,780 (-■-). Tumor volumes were calculated on the days shown. n = 6 mice per group; means ± SE.

evolution to tamoxifen resistance when the drug was stimulating tumor growth (3). In fact, pS2 was significantly lower in tamoxifen-resistant tumors than in tamoxifen-sensitive tumors ($P = .012$). This finding suggests that the agonist activity of tamoxifen, if responsible for tamoxifen-stimulated tumor growth, may be specific to genes associated with cell proliferation, while its antagonist activity continues to suppress the activity of genes less crucial for tumor survival. In contrast to the results obtained with tamoxifen, mRNA expression was nearly

abrogated by treatment with ICI 182,780 ($P < .004$), and there was no difference between sensitive and resistant tumors. It is unlikely, therefore, that ICI 182,780 resistance is caused by metabolic conversion of the drug to E₂, since expression of these estrogen-regulated genes remained low.

Discussion

Clinical data demonstrate that, in some patients, the current endocrine therapies for breast cancer result in temporary tumor regression or growth stabilization, followed by tumor regrowth, usually within 6-18 months of treatment. We have developed an experimental in vivo model that mimics this clinical scenario. Our data suggest that, in this experimental model system, ICI 182,780 possesses a greater ability to suppress estrogen-sensitive gene expression and greater antitumor activity than the partial estrogen antagonist tamoxifen. In addition, MCF-7 tumorigenesis was significantly delayed by ICI 182,780 when compared with tamoxifen. Moreover, a proportion of treated mice failed to develop tumors even after prolonged follow-up, an event rarely encountered in our experience treating mice with tamoxifen. ICI 182,780 also suppressed growth of established tumors for a significantly longer duration

than treatment by estrogen withdrawal alone or with tamoxifen. Finally, expression of the estrogen-regulated genes pS2 and pLIV1 was nearly abolished by treatment with ICI 182,780.

Previous reports by us and by other investigators (7,11,14-16) have also shown that the growth of tumors with acquired tamoxifen resistance can be inhibited or blocked by treatment with a pure antiestrogen such as ICI 182,780, suggesting that the pure antiestrogens work by a different mechanism of action than tamoxifen and other similar antiestrogens. Tamoxifen resistance in our model system is associated with drug-induced tumor growth stimulation that occurs after an initial period of growth suppression (3). The ability of tamoxifen alone to stimulate the growth of these tumors is less than that of estrogen. Interestingly, when combined with estrogen, tamoxifen can still inhibit estrogen-stimulated growth, indicating that it continues to possess both estrogen-agonist and antagonist properties (7). The increasingly dominant agonist properties of tamoxifen that develop after prolonged treatment can be blocked by the addition of pure antiestrogens (7,11). Evidence for tamoxifen-stimulated tumor growth as a mechanism for acquired tamoxifen resistance in patients has also been presented (5,6,17). On the basis of these preclinical studies, it has been suggested that treatment with ICI 182,780 might induce tumor regression in some patients who have developed tamoxifen resistance. One recent study (18) has shown that short-term ICI 182,780 treatment of patients who have ER-positive tumors causes statistically significant reductions in the Ki67 labeling index and reductions in the expression of estrogen-regulated genes such as PgR and pS2. In addition, remissions have now been reported in tamoxifen-resistant patients treated with this drug (19).

Although ICI 182,780 controls MCF-7 tumor growth for longer durations than tamoxifen, eventual resistance to this agent is common. MCF-7 tumors that progress after prolonged treatment are estrogen-independent (grow in the absence of estrogen supplementation) although they are still estrogen-sensitive (growth is enhanced by estrogen). The mechanisms by which resistance to

Table 1. Expression of estrogen-sensitive genes*

Treatment group (No. of blots analyzed)	Gene, relative mRNA level	
	pS2	pLIV1
Estrogen (4)	12.2 ± 0.7	12.2 ± 0.6
Tamoxifen-sensitive (5)	9.8 ± 0.5	6.0 ± 1.5
Tamoxifen-resistant (5)	3.2 ± 0.4	7.5 ± 1.8
ICI-sensitive (5)	0.3 ± 0.05	0 ± 0
ICI-resistant (8)	0.6 ± 0.23	2.3 ± 1.3

*mRNA expression was measured by northern blot analysis of total RNA extracted from MCF-7 tumors taken from mice treated with estrogen (controls), tamoxifen for 3 weeks (tamoxifen-sensitive), tamoxifen until the time of tumor progression (tamoxifen-resistant), ICI 182,780 for 4 weeks (ICI-sensitive), or ICI 182,780 until tumor progression (ICI-resistant). Values shown are the means ± SE of scanning densitometry units corrected for RNA loading.

ICI 182,780 develops are not clear, but reduced levels of ER and reduced expression of estrogen-regulated genes (compared with tamoxifen-sensitive or with tamoxifen-resistant tumors) are evident. Reduced ER levels have also been seen in tumors from patients treated with ICI 182,780, in cultured breast cancer cells, and in mouse uterine tissue following the administration of the prototype pure antiestrogen ICI 164,384 (18-20). Other data suggest that the pure antiestrogen-ER complex may be more fragile and more susceptible to receptor degradative pathways (16). In contrast, ER levels are high in tamoxifen-resistant tumors obtained with our model system (3). On the basis of our data, we would predict that most patients with ICI 182,780-resistant tumors would not respond well to subsequent treatment with tamoxifen.

Even if pure antiestrogens are shown to have superior antitumor activity in women with breast cancer, they may not be the optimal antiestrogens for clinical use. The estrogenic properties of tamoxifen in bone and on blood lipids may help to reduce bone loss and prevent cardiovascular disease, which are added benefits when treating breast cancer patients for prolonged periods after surgery for primary tumors or for breast cancer prevention (21,22). The effect of ICI 182,780 on these parameters is not yet known, but it might be deleterious given its lack of estrogenic qualities. However, treatment with ICI 182,780 might not be associated with the increased risk of endometrial cancer recently attributed to tamoxifen (23). Further clinical study of pure antiestrogens in tamoxifen-resistant and in tamoxifen-naïve patients is clearly indicated.

References

- (1) Systemic treatment of early breast cancer by hormonal, cytotoxic, or immune therapy. 133 randomised trials involving 31,000 recurrences and 24,000 deaths among 75,000 women. Early Breast Cancer Trialists' Collaborative Group [see comment citations in Medline]. *Lancet* 339:1-15, 71-85, 1992
- (2) Encarnacion CA, Ciocca DR, McGuire WL, et al: Measurement of steroid hormone receptors in breast cancer patients on tamoxifen. *Breast Cancer Res Treat* 26:237-246, 1993
- (3) Osborne CK, Coronado E, Allred DC, et al: Acquired tamoxifen resistance: correlation with reduced breast tumor levels of tamoxifen and isomerization of trans-4-hydroxytamoxifen [see comment citation in Medline]. *J Natl Cancer Inst* 83:1477-1482, 1991
- (4) Gottardis MM, Jordan VC: Development of tamoxifen-stimulated growth of MCF-7 tumors in athymic mice after long-term antiestrogen administration. *Cancer Res* 48:5183-5187, 1988
- (5) Pritchard KI, Thomson DB, Myers RE, et al: Tamoxifen therapy in premenopausal patients with metastatic breast cancer. *Cancer Treat Rep* 64:787-796, 1980
- (6) Hoogstraten B, Gad-el-Mawla N, Maloney TR, et al: Combined modality therapy for first recurrence of breast cancer. A Southwest Oncology Group Study. *Cancer* 54:2248-2256, 1984
- (7) Osborne CK, Jarman M, McCague R, et al: The importance of tamoxifen metabolism in tamoxifen-stimulated breast tumor growth. *Cancer Chemother Pharmacol* 34:89-95, 1994
- (8) Wakeling AE, Bowler J: Novel antioestrogens without partial agonist activity. *J Steroid Biochem* 31:645-653, 1988
- (9) Wakeling AE, Dukes M, Bowler J: A potent specific pure antiestrogen with clinical potential. *Cancer Res* 51:3867-3873, 1991
- (10) Wakeling AE: Steroidal pure antiestrogens. In *Regulatory Mechanisms in Breast Cancer* (Lippman M, Dickson R, eds). Boston: Kluwer Acad Publ, 1991, pp 239-257
- (11) Gottardis MM, Jiang SY, Jeng MH, et al: Inhibition of tamoxifen-stimulated growth of an MCF-7 tumor variant in athymic mice by novel steroidal antiestrogens. *Cancer Res* 49:4090-4093, 1989
- (12) Osborne CK, Hobbs K, Clark GM: Effect of estrogens and antiestrogens on growth of human breast cancer cells in athymic nude mice. *Cancer Res* 45:584-590, 1985
- (13) Manning DL, McClelland RA, Gee JM, et al: The role of four oestrogen-responsive genes, pLIV1, pS2, pSYD3 and pSYD8, in predicting responsiveness to endocrine therapy in primary breast cancer. *Eur J Cancer* 29A:1462-1468, 1993
- (14) Brüner N, Frandsen TL, Holst-Hansen C, et al: MCF7/LCC2: a 4-hydroxytamoxifen resistant human breast cancer variant that retains sensitivity to the steroidal antiestrogen ICI 182,780. *Cancer Res* 53:3229-3232, 1993
- (15) Lykkesfeldt AE, Sorensen EK: Effect of estrogen and antiestrogens on cell proliferation and synthesis of secreted proteins in the human breast cancer cell lines MCF-7 and a tamoxifen resistant variant subline, AL-1. *Acta Oncol* 31:131-138, 1992
- (16) Parker MG: Action of pure antiestrogens in inhibiting estrogen receptor action. *Breast Cancer Res Treat* 26:131-137, 1993
- (17) Wiebe VJ, Osborne CK, Fuqua SA, et al: Tamoxifen resistance in breast cancer. *Crit Rev Oncol Hematol* 14:173-188, 1993
- (18) DeFriend DJ, Howell A, Nicholson RI, et al: Investigation of a new pure antiestrogen (ICI 182780) in women with primary breast cancer. *Cancer Res* 54:408-414, 1994
- (19) DeFriend DJ, Blamey RW, Robertson JF, et al: Response to the pure antiestrogen ICI 182780 after tamoxifen failure in advanced breast cancer. *Breast Cancer Res Treat* 17:136, 1993
- (20) Gibson MK, Nemmers LA, Beckman WC Jr, et al: The mechanism of ICI 164,384 antiestrogenicity involves rapid loss of estrogen receptor in uterine tissue. *Endocrinology* 129:2000-2010, 1991
- (21) Love RR, Mazess RB, Barden HW, et al: Effects of tamoxifen on bone mineral density in postmenopausal women with breast cancer [see comment citation in Medline]. *N Engl J Med* 326:852-856, 1992
- (22) Love RR, Newcombe PA, Wiebe DA, et al: Effects of tamoxifen therapy on lipid and lipoprotein levels in postmenopausal patients with node-negative breast cancer [see comment citation in Medline]. *J Natl Cancer Inst* 82:1327-1332, 1990
- (23) Fomander T, Hellström AC, Moberger B: Descriptive clinicopathologic study of 17 patients with endometrial cancer during or after adjuvant tamoxifen in early breast cancer. *J Natl Cancer Inst* 85:1850-1855, 1993

Notes

Supported in part by Public Health Service grants CA30251, CA30195, CA58183, and CA54174 from the National Cancer Institute, National Institutes of Health, Department of Health and Human Services.

Manuscript received September 19, 1994; revised December 27, 1994; accepted March 6, 1995.

Need the latest information on clinical trials for HIV and AIDS?

Call the AIDS Clinical Trials Information Service:

1-800-TRIALS-A

(1-800-874-2572) Monday through Friday, 9 a.m. to 7 p.m. eastern time.

Pending Claims in U.S. Application No. 12/285,887 as of December 9, 2011,
with proposed amendments

Claims 24, 32, 34, 36, 44, and 46 are proposed to be amended. New claims 54-57 are proposed to be added. Claims 25, 28, 31, 33, 37, 40, 43, 45, and 48-53 are proposed to be cancelled. Deletions appear in ~~strikethrough font~~ and/or ~~[[inside double brackets]]~~ and additions are underlined.

Claims 1-23 (Cancelled)

24. (Currently amended) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation comprising:

~~at least 45 mgml⁻¹ of fulvestrant;~~

~~a mixture of from 17—23% w/v of ethanol and benzyl alcohol;~~

~~12—18% w/v of benzyl benzoate; and~~

about 50 mgml⁻¹ of fulvestrant;

about 10% w/v of ethanol;

about 10% w/v of benzyl alcohol;

about 15% w/v of benzyl benzoate; and

a sufficient amount of castor oil vehicle;

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least ~~two~~four weeks.

25. (Cancelled)

26. (Previously presented) The method of claim 24, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml^{-1} .
27. (Previously presented) The method of claim 24, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
28. (Cancelled)
29. (Previously presented) The method of claim 24, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
30. (Previously presented) The method of claim 24, wherein the method further comprises once monthly administration of the formulation.
31. (Cancelled)
32. (Currently amended) The method of ~~claim 31~~ claim 26, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
33. (Cancelled)
34. (Currently amended) The method of ~~claim 33~~ claim 32, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.

35. (Previously presented) The method of claim 34, wherein the method further comprises once monthly administration of the formulation.
36. (Currently amended) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation consisting essentially of:
- ~~at least 45 mgml⁻¹ of fulvestrant;~~
 - ~~a mixture of from 17–23% w/v of ethanol and benzyl alcohol;~~
 - ~~12–18% w/v of benzyl benzoate; and~~
 - ~~a sufficient amount of castor oil vehicle;~~
 - about 50 mgml⁻¹ of fulvestrant;
 - about 10% w/v of ethanol;
 - about 10% w/v of benzyl alcohol;
 - about 15% w/v of benzyl benzoate; and
- wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least ~~two~~ four weeks.
37. (Cancelled)
38. (Previously presented) The method of claim 36, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.

39. (Previously presented) The method of claim 36, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
40. (Cancelled)
41. (Previously presented) The method of claim 36, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
42. (Previously presented) The method of claim 36, wherein the method further comprises once monthly administration of the formulation.
43. (Cancelled)
44. (Currently amended) The method of ~~claim 43~~ claim 38, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
45. (Cancelled)
46. (Currently amended) The method of ~~claim 45~~ claim 44, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
47. (Previously presented) The method of claim 46, wherein the method further comprises once monthly administration of the formulation.

Claims 48-53 (Cancelled)

54. (New) The method according to claim 24, wherein the formulation is administered in a divided dose.
55. (New) The method according to claim 35, wherein the formulation is administered in a divided dose.
56. (New) The method according to claim 36, wherein the formulation is administered in a divided dose.
57. (New) The method according to claim 47, wherein the formulation is administered in a divided dose.

1

3,164,520

INJECTABLE STEROID COMPOSITIONS CONTAINING AT LEAST 75% BENZYL BENZOATE

Raymond Charles Huber, Martinsville, N.J., assignor to Olin Mathieson Chemical Corporation, New York, N.Y., a corporation of Virginia

No Drawing. Filed Oct. 29, 1962, Ser. No. 233,931
4 Claims. (Cl. 167-58)

This invention relates to compositions of matter and more particularly to new parenterally administrable pharmaceutical compositions comprising one or more active medicaments and a physiologically acceptable non-toxic pharmaceutical vehicle, comprised essentially of benzyl benzoate.

The active medicament which may be incorporated in the novel compositions of this invention may be any one which is administered for use in comparatively large unit dosages, for example, 10 mg./ml. to 500 mg./ml. and which is soluble in benzyl benzoate. Examples of the medicaments which may be employed in this invention include inter alia, steroid hormones, especially those steroid hormones which exhibit anabolic, estrogenic, androgenic and progestational activity, for example, 17-hydroxyprogesterone and the esters thereof, testosterone, estradiol and the acid esters thereof, progesterone and its derivatives and Δ^1 -testololactone and its derivatives. In the most preferable embodiment of this invention the active medicament is a steroid hormone although other pharmaceutically active compounds may also be employed, with satisfactory results.

Heretofore it has been well recognized in the preparation of parenterally administrable pharmaceutical compositions that a suitable solvent must be employed to render the composition injectable. However, as the science of medicine has progressed it has been found that increasingly higher dosages of certain medicaments must be employed in the treatment of certain ailments in order to achieve several advantages. Among these advantages can be numbered the prolongation of activity of the medicaments involved and the lessening of the total number of individual injections which are needed to obtain the same results.

Additionally, it has been found that new chemical modifications of medicaments are continually being discovered and the solubility of these modified medicaments in the solvents commonly employed, appears to be more and more limited and it has therefore become increasingly difficult to dissolve these new modified medicaments in parenterally acceptable vehicles. It is well-known that certain pharmaceutical vehicles yield satisfactory results at low level medicament concentrations when employed in compositions for parenteral administration. Such vehicles are the vegetable oils such as cotton seed oil, peanut oil, sesame oil, or corn oil, in combination with small amounts of benzyl benzoate. However, when an increased dosage level of the medicaments is employed, along with a correspondingly necessary increased amount of pharmaceutical vehicle it has been found that certain undesirable disadvantages exist.

The undesirable disadvantages which are present when the prior art vehicles are employed with a high dosage level of medicaments, are many. In addition to the prior art vehicles being incapable of solubilizing any great quantities of the medicaments, it has been found that the compositions heretofore employed produce an undue amount of irritation at the site of injection, when parenterally administered into the animal being treated.

It has now been found that the disadvantages encountered in the parenteral administration of high dosage levels of the medicaments of this invention can be avoided by employing the novel pharmaceutical compositions of

2

this invention. It has been found that these disadvantages can be overcome by employing benzyl benzoate as the essential component of the pharmaceutical vehicle of parenterally administrable compositions. The benzyl benzoate has been found to be capable of dissolving great quantities of the medicaments of this invention and the resulting parenterally administrable composition employing this vehicle does not produce undue irritation when injected into the animals being treated.

The amount of benzyl benzoate which may be employed in the compositions of this invention while still yielding satisfactory results has been found to range from about 75% to 100% by volume of the pharmaceutical vehicle employed. Thus the ratio of benzyl benzoate present in the pharmaceutical vehicle as compared to any other ingredients therein must be at least 3 to 1. In the most preferable embodiment of this invention it has been found that a pharmaceutical vehicle consisting essentially of pure benzyl benzoate yields the best results although at lower levels satisfactory results are also obtained.

As is common in the art of preparing parenterally administrable pharmaceutical compositions other additives such as preservatives, antioxidants or anesthetics, such as benzyl alcohol and the other like well known additives may also be included in the pharmaceutical compositions of this invention. However, their use herein is permissive and not mandatory as their incorporation or omission in the final product of this invention does not substantially affect the results herein obtained.

The compositions of this invention are easily prepared by merely taking the desired amount of medicament and dissolving it in the pharmaceutical vehicle of this invention by any means known in the art, for example, by mere stirring.

The final compositions of this invention are parenterally administrable to the animal being treated. The administration of the composition may be accomplished intramuscularly, subcutaneously or in any other manner known to the art as may be determined in the individual cases wherein this invention is employed. It has been generally found that the most preferable results are obtained when an intramuscular route of administration is employed, although other methods of administration will also give satisfactory results.

The invention is more particularly illustrated by the following examples:

Example 1

Two g. of the acetophenone derivative of 16,17-dihydroxyprogesterone are dissolved in 10 ml. of benzyl benzoate with stirring and warming. The resultant solution is then filled in vials of 5 ml. each and sterilized by autoclaving at 121° C. for two hours.

0.25 ml. of the resulting solution is then injected into the vastus lateralis muscle of a rabbit producing a lesion at the site of the injection having the size of about 640 cubic millimeters after two days.

When 2 g. of the acetophenone derivative of 16,17-dihydroxyprogesterone are dissolved in 4.5 ml. of benzyl benzoate and 5.5 ml. of castor oil in accordance with the procedure of Example 1 and 0.25 ml. of the resultant solution is injected intramuscularly into the rabbit a lesion at the site of injection having a size of 967 cubic millimeters after two days.

Example 2

The procedure of Example 1 is followed except that 2 g. of testosterone palmitate are substituted for the acetophenone derivative of 16,17-dihydroxyprogesterone of Example 1.

0.25 ml. of the resultant solution is injected intramus-

3

cularly into a rabbit producing a lesion at the site of injection having the size of about 420 cubic millimeters after two days. When 2 g. of testosterone palmitate are dissolved in a vehicle consisting of 40% castor oil and 60% benzyl benzoate and the resultant solution is injected intramuscularly into the rabbit, a lesion at the site of injection having a size of 610 cubic millimeters is produced after two days.

Example 3

A 25% solution of progesterone is prepared by dissolving 2.5 g. of progesterone in benzyl benzoate to make 10 ml. Sterilization is obtained by autoclaving the solution at 121° C. for 2 hours. When 0.25 mg. of this solution is injected into the vastus lateralis muscle of the rabbit, a lesion is produced which, after 2 days, measures 672 cubic millimeters.

When 2.5 g. of progesterone are dissolved to make 10 ml. in a mixture of 50% benzyl benzoate and 50% castor oil as the vehicle, and 0.25 ml. of this solution is injected into the rabbit muscle, a lesion size of 898 cubic millimeters is produced after two days.

Example 4

A 50% solution of hormones is prepared by dissolving 2.5 g. of progesterone and 2.5 g. of 17-hydroxyprogesterone caproate in benzyl benzoate to make 10 ml. of final product. After autoclaving at 121° C. for 2 hours to sterilize, 0.25 ml. of the solution is injected into a rabbit muscle and the lesion size is measured after 2 days. A lesion consisting of 572 cubic millimeters was produced. When this same hormone combination in the same proportions was dissolved in a vehicle consisting of 46% benzyl benzoate and 54% castor oil, a rabbit muscle lesion size of 1047 cubic millimeters is produced 2 days after injection of 0.25 ml. of test material.

Example 5

A 40% solution of testosterone enanthate is prepared by dissolving 4.0 g. in benzyl benzoate to make 10 ml. of final volume. After autoclaving at 121° C. for 2 hours to sterilize, 0.25 ml. of the solution is injected into the vastus lateralis muscle of the rabbit and the lesion size is measured after 2 days. A lesion consisting of 847 cubic millimeters is produced.

When this same quantity of hormone is dissolved in a vehicle consisting of 20% benzyl benzoate and 80% sesame oil and 0.25 ml. is injected a lesion size of 1441 cubic millimeters is produced.

Example 6

A 5% solution of Δ^1 -testololactone is prepared by dis-

4

solving 50 mg./ml. in benzyl benzoate and after autoclaving to sterilize, 0.25 ml. of the solution is injected into a rabbit muscle. After 2 days a lesion size of only 483 cubic millimeters is produced.

Example 7

15 mg. of Δ^1 -testololactone is dissolved in a solution comprised of 7.5 ml. of benzyl benzoate and 2.5 ml. of castor oil. The resultant solution is sterilized, then filled in vials of 5 ml. each and sterilized by autoclaving at 121° C. for 2 hours. The injectable solution may then be administered to the patient being treated.

This invention may be variously otherwise embodied within the scope of the appended claims.

What is claimed is:

1. A parenterally administrable pharmaceutical composition comprising the acetophenonide of 16,17-dihydroxyprogesterone and a physiologically acceptable non-toxic pharmaceutical vehicle wherein at least 75% by volume of said vehicle is benzyl benzoate.

2. A parenterally administrable pharmaceutical composition comprising testosterone palmitate and a physiologically acceptable non-toxic pharmaceutical vehicle wherein at least 75% by volume of said vehicle is benzyl benzoate.

3. A parenterally administrable pharmaceutical composition comprising testosterone enanthate and a physiologically acceptable non-toxic pharmaceutical vehicle wherein at least 75% by volume of said vehicle is benzyl benzoate.

4. A method of administering a large single dosage of a steroid which comprises parenterally administering to the patient being treated a composition comprising a steroid selected from the group consisting of 17-hydroxyprogesterone, the caproate ester of 17-hydroxyprogesterone, testosterone, the enanthate ester of testosterone, the palmitate ester of testosterone, estradiol, progesterone, and Δ^1 -testololactone, and a pharmaceutical carrier, said carrier being at least 75% by volume of benzyl benzoate.

References Cited in the file of this patent

Chemical Abstracts, vol. 52, p. 7620b, 1958 (abstr. of Gerosa et al., Ann. Chim., Rome, 47, pp. 1388-1393 (1957)).

Chemical Abstracts, vol. 42, p. 9084g, 1948.

Chemical Abstracts, vol. 47, p. 6611d, 1953.

Merck Index, 7th ed., 1960, p. 137.

U.S. Dispensatory, 25th ed., 1955, p. 160.

Sax: Handbook of Dangerous Materials, p. 45, Reinhold, New York, 1951.

THE MERCCK INDEX

AN ENCYCLOPEDIA OF
CHEMICALS, DRUGS, AND BIOLOGICALS

TWELFTH EDITION

Susan Budavari, *Editor*
Maryadele J. O'Neil, *Senior Associate Editor*
Ann Smith, *Associate Editor*
Patricia E. Heckelman, *Assistant Editor*
Joanne F. Kinneary, *Assistant Editor*

Published by
Merck Research Laboratories
Division of

MERCK & CO., INC.

Whitehouse Station, NJ

1996

MERCK & CO., INC.
Whitehouse Station, N.J.
U.S.A.


1st Edition—1889
2nd Edition—1896
3rd Edition—1907
4th Edition—1930
5th Edition—1940
6th Edition—1952
7th Edition—1960
8th Edition—1968
9th Edition—1976
10th Edition—1983
11th Edition—1989

Library of Congress Catalog
Card Number 89-60001
ISBN Number 0911910-12-3

Copyright © 1996 by MERCK & CO., INC.
All rights reserved. Copyright under the Universal Copyright Convention
and the International Copyright Convention.
Copyright reserved under the Pan-American Copyright Convention.


Printed in the U.S.A.

CM-cellulose	O-(carboxymethyl) cellulose	D.R.P.	(Deutsches Reichs-Patent)—German patent
CMI	cell-mediated immunity	ds	double stranded
CML	cell-mediated lymphocytotoxicity	DTT	dithiothreitol
CMP	cytidine 5'-monophosphate (cytidylic acid)	dyn	dynes
CNS	central nervous system	(E)-	<i>entgegen</i> , stereodescriptor, <i>see Glossary</i>
CoA or CoASH	coenzyme A	E _{1cm} ^{1%}	the absorbance of a solution containing one gram per 100 ml contained in a cell having an absorption path of one cm
coll. vol.	collective volume	E _m	molar extinction coefficient (concn in g-moles/l)
compd	compound	EAA	excitatory amino acid
compn	composition	EAC	erythrocyte coated by antibody and complement
Con A	concanavalin A	EAE	experimental allergic encephalomyelitis
conc(d)	concentrated	EC	electron capture
concn	concentration	ECF-A	eosinophil chemotactic factor of anaphylaxis
config	configuration	ECG	electrocardiogram
constit	constituent	E.C. No.	Enzyme Commission Number
contd	continued	ed.	edition
contg	containing	ED	effective dose
CoQ	coenzyme Q (ubiquinone)	Ed(s).	editor(s)
cor(r)	corrected	EDTA	ethylenediaminetetraacetic acid
corresp	corresponding, corresponds	EEG	electroencephalogram
cp	centipoise	e.g.	(<i>exempli gratia</i>) for example
C.P.	chemically pure	EGF	epidermal growth factor
cpd	compound	<i>idem</i>	the same (author), plural of <i>idem</i>
crit press	critical pressure	EINECS	European Inventory of Existing Chemical Substances
crit temp	critical temperature	EKG	electrocardiogram
cryst	crystalline, crystals	ELISA	enzyme-linked immunosorbent assay
crystn	crystallization	emf	electromotive force
CSF	colony stimulating factor, cerebral spinal fluid	en	ethylenediamine (in formulas) stereochemical descriptor, <i>see Glossary</i>
CTFA	Cosmetic, Toiletary and Fragrance Assoc.	endo-	
CTP	cytidine triphosphate	EPA	Environmental Protection Agency
Cys	cysteine	EPO	erythropoietin; European Patent Office
Cyt	cytosine	ε (epsilon)	molar extinction coefficient (concn in g-moles/l); dielectric constant
d	density; specific gravity (d ₄ ¹⁹ specific gravity at 19° referred to water at 4°)	eq	equation
d-	<i>dextro</i> (rotatory), the opposite of <i>l dextro</i> (in configurational sense only), the opposite of <i>L</i>	equilib	equilibrium
Da	daltons	equiv	equivalent
DEAE cellulose	O-(diethylaminoethyl)cellulose	esp	especially
dec, decomp	decompose(s), decomposition	esu	electrostatic units of electrical charge
decompn	decomposition	Et	ethyl C ₂ H ₅ —
deg	degree	η (eta)	viscosity
deliquesc	deliquescent	<i>et al.</i>	(<i>et alii</i>) and others
Δ (delta)	indicates the locant of the double bond	etc.	<i>et cetera</i>
deriv	derivative	Et ₂ O	ether
determn	determination	EtOH	ethyl alcohol
DFP, DIFP, DIPP	diisopropyl fluorophosphate or diisopropyl phosphofluoridate	Eur. pat. Appl.	European patent application
Dha	dihydroalanine	ev	electron volt
Dhb	dehydrobutyrine, β-methyldehydroalanine	evac	evacuated
diff	difference	evapn	evaporation
dil(d), (n)	dilute, diluted, dilution	exo-	stereochemical descriptor, <i>see Glossary</i>
distln	distillation	expt(ly)	experimental(ly)
dl-	racemic	ext(d)	extract, extracted
DL-	optically inactive by external compensation as contrasted with <i>meso-</i>	extern	externally
dm	decimeter(s)	°F	Fahrenheit degrees; also Fourneau
DMA	dimethylacetamide	F-1-P	fructose 1-phosphate
DMARD	disease modifying antirheumatic drug	F-6-P	fructose 6-phosphate
DMF	dimethylformamide	FA	fatty acid
DMSO	dimethylsulfoxide	FAB	fast atom bombardment
DNA	deoxyribonucleic acid	FAD (FADH ₂)	flavin adenine dinucleotide (reduced form)
cDNA	complementary DNA	FCA	Freund's complete adjuvant (same as CFA)
mtDNA	mitochondrial DNA	Fd	ferredoxin
DNAase	deoxyribonuclease	F.D.A.	Food and Drug Administration (U.S.A.)
DNFB	2,4-dinitro-1-fluorobenzene	FD & C	Food, Drug and Cosmetic (U.S.A.)
DNFB	2,4-dinitrophenyl or 2,4-dinitrophenol		
DNP	dihydroxyphenylalanine		
Dopa	degree of polymerization (number of monomeric units in the polymer)		
dp, DP			



Crystals from acetone + ether, mp 143°. uv max: 259 nm ($E_{1\%}^{1\text{cm}} = 437$). Soluble in water. pH about 6.
THERAP CAT: Anticholinergic.

1157. Benzylamine. *N,N*-Dimethyl-3-[[1-(phenylmethyl)-1*H*-indazol-3-yl]oxy]-1-propanamine; 1-benzyl-3-[3-(dimethylamino)propoxy]-1*H*-indazole; 1-benzyl-1*H*-indazol-3-yl 3-(dimethylamino)propyl ether; benzindamine. $C_{19}H_{23}N_3O$; mol wt 309.41. C 73.76%, H 7.49%, N 13.58%, O 5.17%. Prepn: Fr. pat. 1,382,855; Palazzo, U.S. pat. 3,318,905 (1964, 1967 both to Angelini Francesco); Palazzo *et al.*, *J. Med. Chem.* 9, 38 (1966). Pharmacology: Lisciani *et al.*, *Eur. J. Pharmacol.* 3, 157 (1968). Metabolism: Catanese *et al.*, *Arzneimittel-Forsch.* 16, 1354 (1966); Kataoka *et al.*, *Chem. Pharm. Bull.* 19, 1511 (1971). Toxicology: B. Silvestrini *et al.*, *Toxicol. Appl. Pharmacol.* 10, 148 (1967). Series of articles on pharmacology: *Arzneimittel-Forsch.* 37, 587-646 (1987).



bp_{0.55} 160°.

Hydrochloride, $C_{19}H_{23}N_3O.HCl$, *Afloben*, *Andolex*, *Benalgin*, *Benzyrin*, *Difflam*, *Dorinamin*, *Enzamin*, *Imotryl*, *Ririlim*, *Riripen*, *Salyzoron*, *Saniflor*, *Tamas*, *Tantum*, *Verax*. Crystals, mp 160°. uv max: 306 nm ($E_{1\%}^{1\text{cm}} = 160$). Very sol in water; rather sol in ethanol, chloroform, *n*-butanol. LD₅₀ in mice, rats (mg/kg): 110, 100 i.p.; 515, 1050 orally (Silvestrini).

THERAP CAT: Analgesic; anti-inflammatory; antipyretic.

THERAP CAT (VET): Anti-inflammatory.

1158. Benzyl Acetate. *Acetic acid phenylmethyl ester*; *acetic acid benzyl ester*. $C_9H_{10}O_2$; mol wt 150.18. C 71.98%, H 6.71%, O 21.31%. $C_6H_5CH_2OOCCH_3$. Occurs in a number of plants, particularly jasmine: S. Arctander, *Perfume and Flavor Materials of Natural Origin* (Elizabeth, N.J., 1960) pp 313-314. Prepd from benzyl chloride, acetic acid or sodium acetate and triethylamine: Merker, Scott, *J. Org. Chem.* 26, 5180 (1961); Hennis *et al.*, *Ind. Eng. Chem., Prod. Res. Develop.* 6, 193 (1967). Toxicity study: P. M. Jenner *et al.*, *Food Cosmet. Toxicol.* 2, 327 (1964).

Liquid; pear-like odor. bp 213°, bp₁₀₂ 134°. mp -51°. d_4^{25} 1.050. n_D^{20} 1.5232, n_D^{25} 1.4998. Flash pt, closed cup: 216° F (102° C). Practically insol in water. Misc with alcohol, ether. LD₅₀ orally in rats: 2490 mg/kg (Jenner).

Caution: If ingested can cause G.I. irritation with vomiting and diarrhea. Also irritating to skin, eyes, respiratory tract.

USE: In perfumery, solvent for cellulose acetate and nitrate.

1159. Benzyl Alcohol. *Benzenemethanol*; phenylcarbinol; phenylmethanol; α -hydroxytoluene. C_7H_8O ; mol wt 108.14. C 77.75%, H 7.46%, O 14.80%. $C_6H_5CH_2OH$. Constituent of jasmine, hyacinth, ylang-ylang oils, Peru and Tolu balsams, storax, where it occurs in ester form also.


Originally prepd by the Cannizzaro reaction from benzaldehyde + KOH: Cannizzaro, *Ann.* 88, 129 (1853); cf. Hickinbottom, *Reactions of Organic Compds.* (Longmans, London, 3rd ed., 1957) p 251; A. I. Vogel, *Practical Organic Chemistry* (Longmans, London, 3rd ed., 1959) p 711; Gattermann-Wieland, *Praxis des organischen Chemikers* (de Gruyter, Berlin, 40th ed., 1961) p 193. Produced on a large scale by the action of sodium or potassium carbonate on benzyl chloride: Ger. pat. 484,662; *Chem. Zentr.* 1930, I, 1052; *Frdl.* 16, 426; cf. Kirk-Othmer *Encyclopedia of Chemical Technology* vol. 3 (Interscience, New York, 1964) pp 442-449. Toxicity: Smyth *et al.*, *Arch. Ind. Hyg. Occup. Med.* 4, 119 (1951).

Liquid. Faint aromatic odor. Sharp burning taste. d_4^{20} 1.04535; d_4^{25} 1.04156. mp -15.19°. bp₇₆₀ 204.7°; bp₆₀₀ 183.0°; bp₂₀₀ 160.0°; bp₁₀₀ 141.7°; bp₆₀ 129.3°; bp₄₀ 119.8°; bp₂₀ 105.8°; bp₁₀ 92.6°; bp₅ 80.8°; bp_{1.0} 58.0°. n_D^{20} 1.54035; n_D^{25} 1.53837; Dreisbach, Martin, *Ind. Eng. Chem.* 41, 2875 (1941). Absorption spectrum: Brode, *J. Phys. Chem.* 30, 61 (1926). Vapor density 3.72 (air = 1.00). Flash pt, closed cup 213°F, open cup 220°F. Autoignition temp 817°F. One gram dissolves in about 25 ml water. One volume dissolves in 1.5 vols of 50% ethyl alcohol. Misc with abs and 94% alcohol, ether, chloroform. LD₅₀ orally in rats: 3.1 g/kg (Smyth).

USE: Manuf other benzyl compds. Pharmaceutical aid (antimicrobial). Solvent for gelatin, casein (when hot), solvent for cellulose acetate, shellac. Used in perfumery and in flavoring (mostly in form of its aliphatic esters). In microscopy as embedding material.

THERAP CAT (VET): Has been used for relief from pruritis.

1160. Benzylamine. *Benzenemethanamine*; aminotoluene; phenylmethylamine; moringine. C_7H_9N ; mol wt 107.16. C 78.46%, H 8.47%, N 13.07%. Prepn from benzylchloride and ammonia: Mason, *J. Chem. Soc.* 63, 1311 (1893); by redn of benzonitrile: Carothers, Jones, *J. Am. Chem. Soc.* 47, 3051 (1925); from benzyl bromide + acetamide: Erikson, *Ber.* 59, 2665 (1926); from *N*-benzylphthalimide + hydrazine hydrate: Ing, Manske, *J. Chem. Soc.* 129, 2348 (1926). Identity with moringine: Chakravarti, *Bull. Calcutta School Trop. Med.* 3, 162 (1955); C.A. 50, 16891e (1956).



Liquid; strongly alkaline reaction. bp 185°; bp₁₂ 90°. d_4^{20} 0.983. n_D^{20} 1.5401. Miscible with water, alcohol, ether.

Hydrochloride, $C_7H_9N.HCl$, crystals, mp 253°.

Hydroiodide, $C_7H_9N.HI$, leaflets, mp 162°.

Caution: Highly irritating to skin, mucous membranes.

USE: In organic synthesis.

1161. Benzylaniline. *N-Phenylbenzenemethanamine*; *N-phenylbenzylamine*; benzylphenylamine. $C_{13}H_{13}N$; mol wt 183.25. C 85.21%, H 7.15%, N 7.64%. $C_6H_5CH_2NHC_6H_5$. Prepn from benzyl alc and aniline in the presence of KOH: Sprinzak, *J. Am. Chem. Soc.* 78, 3207 (1956); from benzaldehyde and aniline in the presence of $NaBH_4$: Schellenberg, *J. Org. Chem.* 28, 3259 (1963).

Prisms, mp 37-38°. bp 306-307°. Practically insol in water; sol in alcohol, chloroform, ether.

1162. Benzyl Benzoate. *Benzoic acid phenylmethyl ester*; benzoic acid benzyl ester; benzylbenzenecarboxylate; Ascabin; Venzonate; Ascabiol. $C_{14}H_{12}O_2$; mol wt 212.25. C 79.23%, H 5.70%, O 15.08%. $C_6H_5COOCH_2C_6H_5$. Contained in Peru and Tolu balsams. Prepd by the action of sodium benzoate on benzaldehyde: Kamm, Kamm, *Org. Syn. coll. vol. I*, 104 (2nd ed., 1941); by the dry esterification of sodium benzoate and benzyl chloride in the presence of triethylamine: Thorp, Nottorf, *Ind. Eng. Chem.* 39, 1300 (1947). Toxicity studies: Graham, Kuizenga, *J. Pharmacol. Exp. Ther.* 84, 358 (1945); Draize *et al.*, *J. Pharmacol. Exp. Ther.* 93, 26 (1948). Comprehensive description: M. M. A.

Hassan, J. S. Mossa, *Anal. Profiles Drug Subs.* 10, 55-74 (1981).

Leaflets or oily liq; faint, pleasant, aromatic odor; sharp burning taste. mp 21°. d_4^{25} 1.118. bp 323-324°. bp₁₆ 189-191°. bp_{4.5} 156°. Sparingly volatile with steam. n_D^{25} 1.5681. Insol in water or glycerol. Miscible with alc, chloroform, ether, oils. LD₅₀ in rats, mice, rabbits, guinea pigs (g/kg): 1.7, 1.4, 1.8, 1.0 orally (Draize).

Caution: In exptl animals, ingestion causes progressive incoordination, excitation, convulsions, death. May cause skin irritation in humans. Avoid contact with eyes. *Clinical Toxicology of Commercial Products*, R. E. Gosselin et al., Eds. (Williams & Wilkins, Baltimore, 4th ed., 1976) Section II, p 137.

USE: As solvent of cellulose acetate, nitrocellulose and artificial musk; substitute for camphor in celluloid and plastic pyroxylin compds; perfume fixative; in confectionery and chewing gum flavors.

THERAP CAT: Scabicide, pediculicide.

THERAP CAT (VET): Acaricide, pediculicide. *Contraindicated* in cats.

1163. Benzyl Bromide. (*Bromomethyl*)benzene; α -bromotoluene; ω -bromotoluene. C₇H₇Br; mol wt 171.04. C 49.16%, H 4.13%, Br 46.72%. C₆H₅CH₂Br. Prep'd by the action of bromine on toluene in ultraviolet light: v. Konek, *Loczka, Ber.* 57, 679 (1924); Zelinsky, Ger. pat. 478,084; *Chem. Zentr.* 1929 II, 1216; *Frdl.* 16, 335; by the action of bromine on dibenzyl ether: Lachman, *J. Am. Chem. Soc.* 45, 2359 (1923).

Lacrimatory liquid. mp -3.9°. bp 198-199°. bp₁₀₀ 127°. d_4^{25} 1.4380; d_4^{17} 1.443; d_4^{15} 1.3886. Slowly decomp by water.

Caution: Intensely irritating to skin, eyes, mucous membranes. Large doses cause CNS depression.

1164. Benzyl Chloride. (*Chloromethyl*)benzene; α -chlorotoluene. C₇H₇Cl; mol wt 126.59. C 66.42%, H 5.57%, Cl 28.01%. C₆H₅CH₂Cl. Made by cautious chlorination of toluene: A. I. Vogel, *Practical Organic Chemistry* (Longmans, London, 3rd ed., 1959) p 538; Gattermann-Wieland, *Praxis des organischen Chemikers* (de Gruyter, Berlin, 40th ed., 1961) p 92. Manuf: Faith, Keyes & Clark's *Industrial Chemicals*, F. A. Lowenheim, M. K. Moran, Eds. (Wiley-Interscience, New York, 4th ed., 1975) pp 145-148.

Very refractive liquid; rather unpleasant, irritating odor. d_4^{25} 1.100. bp 179°. mp -48° to -43°. n_D^{25} 1.5415. Insol in water. Miscible with alcohol, chloroform, ether. Rapidly dec when heated in the presence of iron.

Caution: Potential symptoms of overexposure are irritation of eyes and nose; weakness; irritability; headache; skin eruption; pulmonary edema. See *NIOSH Pocket Guide to Chemical Hazards* (DHHS/NIOSH 90-117, 1990) p 46.

USE: Manuf benzyl compds, perfumes, pharmaceutical products, dyes, synthetic tannins, artificial resins.

1165. Benzyl Cinnamate. 3-Phenyl-2-propenoic acid phenylmethyl ester; *trans-cinnamic acid benzyl ester*; cinnamin. C₁₇H₁₆O₂; mol wt 238.29. C 80.65%, H 5.92%, O 13.43%. C₆H₅CH=CHCOOCH₂C₆H₅. Constituent of storax, Peru and Tolu balsams: Tschirch, *Trog, Arch. Pharm.* 232, 70 (1894); Tschirch, Oberländer, *ibid.* 559. Prep'n: Volwiler, Vliet, *J. Am. Chem. Soc.* 43, 1672 (1921); Eliel, Anderson, *ibid.* 74, 547 (1952); Bender, Zerner, *ibid.* 84, 2550 (1962). Toxicity study: P. M. Jenner et al., *Food Cosmet. Toxicol.* 2, 327 (1964).

Crystals from 95% ethanol; sweet odor of balsam. mp 39°; also reported as mp 33-34° (Volwiler, Vliet). Dec on distillation at ordinary pressure; bp_{0.5} 154-157°, bp₃ 195-200°, bp₂₂ 228-230°. Practically insol in water, propylene glycol and glycerin. Sol in alc, ether, oils. LD₅₀ orally in rats: 5530 mg/kg (Jenner).

USE: In artificial flavors, in perfumes, mainly as a fixative.

1166. Benzyl Cyanide. *Benzeneacetonitrile*; phenylacetonitrile; α -tolunitrile; ω -cyanotoluene. C₈H₇N; mol wt 117.15. C 82.02%, H 6.02%, N 11.96%. C₆H₅CH₂CN. Occurs in garden cress and other plants; made from benzyl chloride, and NaCN: Adams, Thal, *Org. Syn.* vol. 2, 9 (1922); coll. vol. I, 101 (107 in 2nd ed.).

Oily liquid, aromatic odor. d_4^{25} 1.0214. mp -23.8°. bp₇₀₀ 233.5°; bp₁₀₀ 161.8°; bp₂₀ 119.4°; bp_{1.9} 60°. n_D^{25} 1.52105. Insoluble in water, miscible with alc, ether.

1167. Benzyl Ether. 1,1'-[Oxybis(methylene)]bis(benzene); dibenzyl ether. C₁₄H₁₄O; mol wt 198.26. C 84.81%, H 7.12%, O 8.07%. (C₆H₅CH₂)₂O. Prep'd: Lachman, *J. Am. Chem. Soc.* 45, 2356 (1923); Staab, Wendel, *Ber.* 93, 2902 (1960); Lichtenberger, Tritsch, *Bull. Soc. Chim. France* 1961, 363. Manuf by reduction of benzaldehyde in the presence of [Co(CO)₄]₂: Wender, Orchin, U.S. pat. 2,614,107 (1952 to U.S.A. as represented by the Secy. of Agr.). Physical properties: Svirbely et al., *J. Am. Chem. Soc.* 71, 507 (1949); Dreisbach, Martin, *Ind. Eng. Chem.* 41, 2875 (1949). Miscibility: Jackson, Drury, *ibid.* 51, 1491 (1959).

Unstable liquid, bp 295-298° (with dec), bp₂₁ 173-174°; bp₁ 125.5-126.5°. Appears to dec slowly at ordinary temps. d_4^{25} 1.0341; d_4^{25} 0.99735; d_4^{20} 1.00142; d_4^{15} 1.0482. n_D^{25} 1.5601 (Svirbely et al.), 1.53851 (Dreisbach, Martin); n_D^{20} 1.54057 (Dreisbach, Martin), 1.566 (Lichtenberger, Tritsch). Practically insol in water; miscible with ethanol, ether, chloroform, acetone.

USE: Plasticizer for nitrocellulose; solvent in perfumery.

1168. Benzyl Ethyl Ether. (*Ethoxymethyl*)benzene. C₉H₁₂O; mol wt 136.19. C 79.37%, H 8.88%, O 11.75%. C₆H₅CH₂OC₂H₅. Preparation from sodium ethoxide and benzyl bromide: Letsinger, Pollart, *J. Am. Chem. Soc.* 78, 6079 (1956); by reduction of benzaldehyde diethyl acetal with LiAlH₄-AlCl₃: Eliel, Rerick, *J. Org. Chem.* 23, 1088 (1958).

Oily liquid, aromatic odor. bp 186°; bp₁₀ 65°. d 0.949, n_D^{25} 1.4955. Volatile with steam. Practically insol in water; miscible with alcohol, ether.

1169. Benzyl Formate. *Formic acid phenylmethyl ester*; *formic acid benzyl ester*. C₈H₈O₂; mol wt 136.15. C 70.57%, H 5.92%, O 23.50%. HCOOCH₂C₆H₅. Prep'n from formic acid and benzyl alcohol: Mailhe, *Chem. Ztg.* 35, 508 (1911).

Liquid; pleasant fruity odor. d 1.081. bp 203°. Practically insol in water. Sol in alcohol.


USE: Solvent for cellulose esters; in perfumery.

1170. Benzyl Fumarate. (*E*)-2-Butenedioic acid bis(phenylmethyl) ester; *fumaric acid dibenzyl ester*; dibenzyl fumarate. C₁₈H₁₆O₄; mol wt 296.32. C 72.96%, H 5.44%, O 21.60%. C₆H₅CH₂OOCCH=CHCOOCH₂C₆H₅. Prep'n from fumaric acid and benzyl alcohol: Volwiler, Vliet, *J. Am. Chem. Soc.* 43, 1672 (1921).

Cryst powder, mp 58.5-59.5°. bp₃ 210-211°. Practically insol in water. Sol in alcohol, chloroform, ether, oils.

USE: In room spray deodorant: Kulka, U.S. pat. 3,077,457 (1963 to Fritzsche Bros.).


1171. Benzylhydrochlorothiazide. 6-Chloro-3,4-dihydro-3-(phenylmethyl)-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide; 3-benzyl-6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide; 6-chloro-7-sulfamoyl-3-benzyl-3,4-dihydro-1,2,4-benzothiadiazine 1,1-dioxide; 3-benzyl-6-chloro-3,4-dihydro-7-sulfamoyl-1,2,4-benzothiadiazine 1,1-dioxide; Behdy. C₁₆H₁₄ClN₂O₂S₂; mol wt 387.87. C 43.35%, H 3.64%, Cl 9.14%, N 10.83%, O 16.50%, S 16.53%. Prep'n: Werner et al., *J. Am. Chem. Soc.* 82, 1161 (1960); Novello et al., *J. Org. Chem.* 25, 970 (1960); Ugi, U.S. pat. 3,108,097 (1963).



Crystals from acetic acid + water, mp 260-262°. Also reported as crystals from water, mp 269°.

THERAP CAT: Antihypertensive; diuretic.


1172. Benzylideneacetone. 4-Phenyl-3-buten-2-one; benzalacetone; methyl styryl ketone; cinnamyl methyl ketone; acetocinnamone. C₁₀H₁₀O; mol wt 146.19. C 82.16%, H 6.89%, O 10.94%. C₆H₅CH=CHCOCH₃. Prep'd by con-



Yellow liquid. bp_2 123-125°. n_D^{25} 1.569-1.672. d_4^{25} 1.029-1.031. LD_{50} orally in rats, mice: 1920, 1730 mg/kg (Piul'skaya).

USE: Antioxidant in feed and food; antidegradation agent for rubber.

3801. Ethoxzolamide. *6-Ethoxy-2-benzothiazolesulfonamide*; ethoxzolamide; Cardrase; Ethamide; Glaucotensil; Redupresin. $C_9H_{10}N_2O_3S_2$; mol wt 258.32. C 41.85%, H 3.90%, N 10.84%, O 18.58%, S 24.83%. Carbonic anhydrase inhibitor. Prepn: Brit. pat. 795,174; J. Korman, U.S. pat. 2,868,800 (1958, 1959 both to Upjohn).




Crystals from ethyl acetate + Skellysolve B, mp 188-190.5°.

THERAP CAT: Diuretic.

THERAP CAT (VET): Diuretic.

3802. Ethylbenzotropine. *endo-3-(Diphenylmethoxy)-8-ethyl-8-azabicyclo[3.2.1]octane*; *3a-(diphenylmethoxy)-8-ethyl-1aH,5aH-nortropine*; *N-ethylnortropine benzhydryl ether*; tropethydrylin; *N-ethyl-8-aza-3-bicyclo[3.2.1]octyl benzhydryl ether*; *N-ethylbenzotropine*; ethylbenzotropine. $C_{22}H_{27}NO$; mol wt 321.46. C 82.20%, H 8.47%, N 4.36%, O 4.98%. Prepn: Brit. pat. 804,837 (1958 to Sandoz); Boehringer *et al.*, Brit. pat. 824,875 (1959 to Boehringer, Ing.). Used as hydrochloride or hydrobromide salts.



Hydrochloride, $C_{22}H_{27}NO.HCl$, *Ponalid*. Crystals from acetone, mp 190-191°.

Hydrobromide, $C_{22}H_{27}NO.HBr$, *Panalid*. Crystals from methanol + ether, mp 226-228°.

THERAP CAT: Anticholinergic.

3803. Ethyl Acetate. *Acetic acid ethyl ester*; acetic ether; vinegar naphtha. $C_4H_8O_2$; mol wt 88.11. C 54.53%, H 9.15%, O 36.32%. $CH_3COOC_2H_5$. Obtained by the slow distillation of a mixture of acetic acid, ethyl alc, and sulfuric acid: Alheritiere, Mercier, U.S. pat. 2,787,636 (1957 to Usines de Melle); Faith, Keyes, & Clark's *Industrial Chemicals*, F. A. Lowenheim, M. K. Moran, Eds. (Wiley-Interscience, New York, 4th ed., 1975) pp 350-354. Toxicity: H. F. Smyth *et al.*, *Am. Ind. Hyg. Assoc. J.* 23, 95 (1962).

Clear, volatile, flammable liq; characteristic fruity odor; pleasant taste when diluted. Slowly dec by moisture, then acquires an acid reaction. Absorbs water (up to 3.3% w/w). d_4^{20} 0.902; d_4^{25} 0.898. bp 77°. mp -83°. Flash pt +7.2° (open cup). Ignition temp 800°F. Explosive limits (% vol in air): 2.2 to 11.5. n_D^{20} 1.3719. Vapor density 3.04 (air = 1). One ml dissolves in 10 ml water at 25°; more sol at lower and less sol at higher temps. Misc with alc, acetone, chloroform, ether. Azeotropic mixture with water (6.1% w/w) bp

70.4°. Azeotropic mixture with water (7.8% w/w) and alc (9.0% w/w) bp 70.3°. Keep tightly closed in a cool place and away from fire. LD_{50} orally in rats: 11.3 ml/kg (Smyth).

Caution: Potential symptoms of overexposure are irritation of eyes, nose and throat; narcosis; dermatitis. See *NIOSH Pocket Guide to Chemical Hazards* (DHHS/NIOSH 90-117, 1990) p 104.

USE: Pharmaceutical aid (flavor); artificial fruit essences; solvent for nitrocellulose, varnishes, lacquers, and aeroplane dopes; manuf smokeless powder, artificial leather, photographic films and plates, artificial silk, perfumes; cleaning textiles, etc.

3804. Ethyl Acetoacetate. *3-Oxobutanoic acid ethyl ester*; acetoacetic acid ethyl ester; acetoacetic ester; ethyl 3-oxobutanoate. $C_6H_{10}O_3$; mol wt 130.14. C 55.37%, H 7.74%, O 36.88%. $CH_3COCH_2COOC_2H_5$. Only the equilibrium mixture of the keto and enol forms is described here. Prepd from ethyl acetate by the action of sodium, sodium ethoxide, sodamide, or calcium: Inglis, Roberts, *Org. Syn. coll. vol. 1*, 235 (2nd ed., 1941); Hansley, Schott, U.S. pat. 2,843,623 (1958 to Natl. Distillers); Scheibler, *Ann.* 565, 176 (1949); Gattermann-Wieland, *Praxis des Organischen Chemikers* (de Gruyter, Berlin, 40th ed., 1961) p 218. Discussion of keto-enol tautomerism: Ward, *J. Chem. Ed.* 39, 95 (1962). Toxicity study: H. F. Smyth *et al.*, *J. Ind. Hyg. Toxicol.* 31, 60 (1949).

Liq. Agreeable odor. d_4^{20} 1.0357; d_4^{25} 1.0288; d_4^{25} 1.0213; d_4^{25} 0.9924; d_4^{25} 0.9703. mp -45°. bp_{760} 180.8°; bp_{400} 158.2°; bp_{200} 138.0°; bp_{60} 106°; bp_{20} 81.1°; bp_3 54.0°; $bp_{1.0}$ 28.5°. n_D^{20} 1.41937. Absorption spectrum: Morton, Rosney, *J. Chem. Soc.* 1926, 711. Flash pt, closed cup: 184°F. Sol in about 35 parts water; misc with the usual organic solvents. LD_{50} orally in rats: 3.98 g/kg (Smyth).

Caution: Moderately irritating to skin, mucous membranes.

3805. Ethyl Acrylate. *2-Propenoic acid ethyl ester*; acrylic acid ethyl ester. $C_5H_8O_2$; mol wt 100.12. C 59.98%, H 8.05%, O 31.96%. $CH_2=CHCOOCH_2CH_3$. Prepd from ethylene chlorohydrin or acrylonitrile, ethanol, and sulfuric acid; also by an oxo reaction from acetylene, carbon monoxide, and ethanol in the presence of suitable catalysts. See the refs under Methyl Acrylate.

Monomer, liquid, acid, penetrating odor, retained by clothing. *Lacrimator*. d_4^{20} 0.9405. fp below -72°. bp_{760} 99.4°; $bp_{30.2}$ 20° (polymerizes on distn). n_D^{20} 1.404. Specific heat at -60°: 0.442 cal/g/°C. Heat of vaporization 8.27 kcal/mol; heat of combustion 655.49 kcal/mol. Flash pt, open cup: 60°F (15°C). Vapor density 3.45 (air = 1). Soly in water at 20°: 2 g/100 ml. Soly of water in ethyl acrylate at 20°: 1.5 g/100 g. Sol in alcohol, ether. Azeotropes: 45.0% water = bp 81°; 56.8% ethanol = bp 76°. Easily polymerizes on standing; polymerization process speeded up by heat, light, and peroxides. If pure, the monomer can be stored below +10° without incurring polymerization.

Polymer, transparent, elastic substance. Practically no odor. Little adhesive power. Resists the usual solvents.

Caution: Potential symptoms of overexposure to the monomer are irritation of eyes, respiratory system and skin. See *NIOSH Pocket Guide to Chemical Hazards* (DHHS/NIOSH 90-117, 1990) p 106. See also *Patty's Industrial Hygiene and Toxicology* vol. 2A, G. D. Clayton, F. E. Clayton, Eds. (Wiley-Interscience, New York, 3rd ed., 1981) p 2292-2296. This substance may reasonably be anticipated to be a carcinogen: *Seventh Annual Report on Carcinogens* (PB95-109781, 1994) p 203.

USE: The monomer in the manuf of water emulsion paint vehicles; in production of emulsion-based polymers used in textile and paper coatings, leather finish resins and adhesives. Imparts flexibility to hard films.

3806. Ethyl Alcohol. *Ethanol*; absolute alcohol; anhydrous alcohol; dehydrated alcohol; ethyl hydrate; ethyl hydroxide. C_2H_6O ; mol wt 46.07. C 52.14%, H 13.13%, O 34.73%. C_2H_5OH . Manuf: by fermentation of starch, sugar, and other carbohydrates; from ethylene, acetylene, sulfite waste liquors, and synthesis gas (CO + H₂); by hydrolysis of ethyl sulfate, and oxidation of methane. Toxicity: G. S. Wiberg *et al.*, *Toxicol. Appl. Pharmacol.* 16, 718 (1970). Embryotoxicity in mammals: N. A. Brown *et al.*,

Science 206, 573 (1979). Possible mechanism for actions of ethanol on the brain: G. Aston-Jones *et al.*, *Nature* 296, 857 (1982). Ethanol-induced chromosomal abnormalities in mice: M. H. Kaufman, *ibid.* 302, 258 (1983). Disruption of reproductive function in female primates following alcohol self-administration: N. K. Mello *et al.*, *Science* 221, 677 (1983). Review of metabolism and toxicity: C. S. Lieber in *Reviews in Biochemical Toxicology* vol. 5, E. Hodgson *et al.*, Eds. (Elsevier, New York, 1983) pp 267-312; of pharmacology: L. Pohorecky, J. Brick, *Pharmacol. Ther.* 36, 335-427 (1988); of hepatotoxicity: C. S. Lieber, L. M. DeCarli, *J. Hepatol.* 12, 394-401 (1991). General reviews: P. Baud, "Ethyl Alcohol Industry" in Grignard, *Traité de Chimie Organique* vol. 5 (Masson, 1937) pp 841-975; Zabel, *Chem. Inds.* (now *Chem. Week*) 64, 212 (1949); Faith, Keyes & Clark's *Industrial Chemicals*, F. A. Lowenheim, M. K. Moran, Eds. (Wiley-Interscience, New York, 4th ed., 1975) pp 355-364; P. D. Sherman, P. R. Kavasmancik, "Ethanol" in Kirk-Othmer *Encyclopedia of Chemical Technology* vol. 9 (Interscience, New York, 3rd ed., 1980) pp 338-380.

Clear, colorless, very mobile, flammable liquid; pleasant odor; burning taste. Absorbs water rapidly from air. d_4^{20} 0.789. bp 78.5°. mp -114.1°. n_D^{20} 1.361. Flash pt, closed cup: 13°C. Miscible with water and with many organic liquids. *Keep tightly closed, cool, and away from flame!* LD₅₀ in young, old rats (g/kg): 10.6, 7.06 orally (Wiberg).

The terms *95% alcohol* and *alcohol* (when used alone) refer to a binary azeotrope having a distillate composition of 95.57% ethyl alcohol (by wt) and bp 78.15°. *Alcohol, USP* is specified as containing not less than 92.3% and not more than 93.8% by weight, corresponding to not less than 94.9% and not more than 96.0% by vol of C₂H₅OH at 15.56°. d_{20}^{25} 0.810; d 0.816 at 15.56° (60°F). *Diluted alcohol*, prepd from equal vols 95% alcohol and water, contains about 41.5% by wt or about 48.9% by vol of C₂H₅OH. d_{20}^{25} 0.931; d 0.936 at 15.56° (60°F). See U.S.P. XXI, 22, 1530 (1985).

Caution: Nausea, vomiting, flushing, mental excitement or depression, drowsiness, impaired perception, incoordination, stupor, coma, death may occur, *cf.* *Clinical Toxicology of Commercial Products*, R. E. Gosselin *et al.*, Eds. (Williams & Wilkins, Baltimore, 5th ed., 1984) Section III, pp 166-171.

USE: Most ethyl alcohol is used in alcoholic beverages in suitable dilutions. Other uses are as solvent in laboratory and industry, in the manufacture of denatured alcohol, pharmaceuticals (rubbing compds, lotions, tonics, colognes), in perfumery, in organic synthesis. Octane booster in gasoline. Pharmaceutical aid (solvent).

THERAP CAT: Antiseptic.

THERAP CAT (VET): Antiseptic. To destroy nerve tissue. Solvent and dehydrating agent.

3807. Ethyl Alcohol, Denatured. Denatured alcohol. Ethyl alcohol to which has been added some substance or substances which, while allowing the use of the alcohol in the most varied industries and arts, renders it entirely unfit for consumption as a beverage. The most commonly used denaturants, either alone or in combination, are the following: Methanol, camphor, Aldehol, amyl alcohol, gasoline, isopropanol, terpineol, benzene, castor oil, acetone, nicotine, aniline dyes, ether, cadmium iodide, pyridine bases, sulfuric acid, kerosene, diethyl phthalate. *Formula 1* is 5 gallons approved wood alcohol added to 100 gal of 95% ethanol. *Formula 2B* is 0.5 gal benzene added to 100 gal of 95% ethanol. Similarly *formula 3A* contains 5 gal commercial methanol, *formula 6B* contains 0.5 gal pyridine bases, *formula 12A* 5 gal benzene, *formula 13A* 10 gal ethyl ether, *formula 19 4* gal methyl isobutyl ketone and 1 gal kerosene, *formula 20 5* gal crude chloroform, *formula 23A* 10 gal acetone, *formula 28 10* gal benzene, *formula 28A* 1 gal gasoline, *formula 30 10* gal methanol, *formula 32 5* gal ethyl ether, *formula 33 30* lbs methyl violet, *formula 35A* 5 gal ethyl acetate, *formula 39C* 1 gal diethyl phthalate; *formula 44* contains 20 gal *n*-butanol. Additional permissible formulas are given in *Appendix to Regulations No. 3, Formulae for Completely and Specially Denatured Alcohol*, published by the U.S. Treasury Dept., Bureau of Industrial Alcohol. Reprinted in N. A. Lange, *Handbook of Chemistry*.

Caution: Denaturants, particularly methanol, may modify and increase toxic symptoms caused by ingestion and exposure to fumes.

3808. Ethylamine. *Ethanamine*; monoethylamine; aminoethane. C₂H₇N; mol wt 45.08. C 53.28%, H 15.65%, N 31.07%. C₂H₅NH₂. Prepn from ethyl iodide + liq ammonia: Watt, Otto, *J. Am. Chem. Soc.* 69, 836 (1947); from ethanol + ammonia: Davies *et al.*, U.S. pat. 2,609,394 (1952 to ICI); Lemon, Myerly, U.S. pat. 3,022,349 (1962 to Union Carbide). Toxicity study: H. F. Smyth *et al.*, *Arch. Ind. Hyg. Occup. Med.* 10, 61 (1954).

Flammable, liq; ammonia odor; strong alkaline reaction. d_{20}^{25} 0.689. bp 16.6°. Solidif -80°. Miscible with water, alcohol, ether. *Keep tightly closed and in cold place.* LD₅₀ orally in rats: 0.40 g/kg (Smyth).

Hydrochloride, C₂H₇N.HCl, crystals from ethanol + water, mp 110°. d 1.22. Soluble in 0.4 part water; freely sol in alcohol; slightly sol in chloroform or acetone. Practically insol in ether. *Keep well closed.*

Hydriodide, C₂H₇N.HI, hygroscopic crystals, mp 188°. d 2.10. Freely sol in water or alcohol. Practically insol in chloroform, ether. *Keep well closed and protected from light.*


Oleate, C₂H₇N.C₁₈H₃₄O₂, *ethanamine (Z)-9-octadecenoate, Etalate*. Commercial prepn is a 5% soln with 2% benzyl alcohol as anodyne.

Caution: Potential symptoms of overexposure are irritation of eyes; skin burns; respiratory irritation; dermatitis. See *NIOSH Pocket Guide to Chemical Hazards* (DHHS/NIOSH 90-117, 1990) p 106.

USE: In resin chemistry; stabilizer for rubber latex; intermediate for dyestuffs, medicinals; in oil refining; in organic syntheses.

THERAP CAT: Oleate as a sclerosing agent.

3809. N-Ethylamphetamine. *N-Ethyl- α -methylbenzene-ethanamine*; *N-ethyl- α -methylphenethylamine*; *N-ethyl- ω -phenylisopropylamine*; 2-ethylamino-1-phenylpropane; Adiparhol; Apetinin. C₁₁H₁₇N; mol wt 163.26. C 80.93%, H 10.50%, N 8.58%. Prepn: Keil, Dobbe, Ger. pat. 767,263 (1952 to Theodor H. Temmler); Leonard *et al.*, *J. Am. Chem. Soc.* 80, 4858 (1958). Separation of isomers: *Brit. Pat.* 814,339 (1959 to Sterling Drug).




bp₁₄ 104.5-106°. n_D^{25} 1.4986. *d*-Form hydrochloride, C₁₁H₁₇N.HCl, mp 154-156°. [α]_D²⁵ +17.2° (c = 2 in water).

l-Form hydrochloride, C₁₁H₁₇N.HCl, mp 155-156°. [α]_D²⁵ -17.3° (c = 2 in water).

Note: This is a controlled substance (stimulant) listed in the U.S. Code of Federal Regulations, Title 21 Part 1308.11 (1995).

THERAP CAT: Anorexic.

3810. Ethyl Amyl Ketone. *5-Methyl-3-heptanone*; amyl ethyl ketone; EAK. C₈H₁₆O; mol wt 128.21. C 74.94%, H 12.58%, O 12.48%. *Review:* Buller, *Ind. & Eng. Chem.* 48, 1323 (1956).




Liquid. Mild fruity odor. d_{20}^{25} 0.820-0.824. One gallon weighs 6.83 lbs at 20°. bp₇₆₀ 157-162°. Flash pt 59° (138°F). Evaporation rate 0.3 (*n*-butyl acetate = 1.0). n_D^{25} 1.4195. Slightly miscible with water. Compatible with alcohols, ketones, ethers, many other organic solvents.

Caution: Narcotic in high concns. **USE:** Solvent for nitrocellulose-alkyd, nitrocellulose-maleic, and vinyl resins.

3811. Ethylaniline. *N-Ethylbenzenamine*; ethylphenylamine. C₉H₁₁N; mol wt 121.18. C 79.29%, H 9.15%, N


W

10173. Warburganal. *[1S-(1 α ,4 α ,8 α \beta)]-1,4,4a,5,6,7,8,8a-Octahydro-1-hydroxy-5,5,8a-trimethyl-1,2-naphthalenedicarboxaldehyde.* C₁₅H₂₂O₃; mol wt 250.34. C 71.97%, H 8.86%, O 19.17%. Drimane sesquiterpene with antifeedant activity against the African army worm. Biological activity includes plant growth regulation, cytotoxic, antimicrobial and molluscicidal properties. Isoln from *Warburgia ugandensis*, *Canellaceae* and structure: I. Kubo *et al.*, *Chem. Commun.* 1976, 1013. Relationship between structure and antifeedant activity: K. Nakanishi, I. Kubo, *Isr. J. Chem.* 16, 28 (1977). Synthesis of (\pm)-warburganal: S. P. Tanis, K. Nakanishi, *J. Am. Chem. Soc.* 101, 4398 (1979); T. Nakata *et al.*, *ibid.* 4400; A. S. Kende, T. J. Blacklock, *Tetrahedron Letters* 1980, 3119; P. A. Wender, S. L. Eck, *ibid.* 1982, 1871; D. M. Hollinshead *et al.*, *J. Chem. Soc. Perkin Trans. I* 1983, 1579. See also: Japan. Kokai 80 136,238, and 80 136,240 (both 1980 to Inst. Phys. Chem. Res.); Japan. Kokai 81 43,236 (1981 to Suntory Ltd.); Japan. Kokai 83 38,232 (1983 to Teikoku Zoki). Synthesis of (-)-warburganal: H. Okawara *et al.*, *Tetrahedron Letters* 1982, 1087.



mp 98-99°. uv max (methanol): 224 nm (ϵ 6300). $[\alpha]_D^{25}$ -260° (c = 0.350 in CHCl₃).

10174. Warfarin. *4-Hydroxy-3-(3-oxo-1-phenylbutyl)-2H-1-benzopyran-2-one*; *3-(α -acetylbenzyl)-4-hydroxycoumarin*; *1-(4'-hydroxy-3'-coumarinyl)-1-phenyl-3-butanone*; *3- α -phenyl- β -acetyl-4-hydroxycoumarin*; compound 42; WARF compound 42; Co-Rax; Rodex. C₁₉H₁₆O₃; mol wt 308.33. C 74.01%, H 5.23%, O 20.76%. The commercial product is the racemic mixture; the *S*(-)-form is more active than the *R*-isomer. Prepd by the Michael condensation of benzylidene-acetone with 4-hydroxycoumarin: Stahmann *et al.*, U.S. pat. 2,427,578 (1947); Schroeder, Link, U.S. pat. 2,765,321 (1956 to Wisconsin Alumni Res. Found.); Link, U.S. pat. 2,777,859 (1957). Resolution and abs configuration: West *et al.*, *J. Am. Chem. Soc.* 83, 2676 (1961); Preis, *Dissertation Abstr.* 18, 793 (1958); Preis *et al.*, U.S. pat. 3,239,529 (1966 to Wisconsin Alumni Res. Found.). Mechanism of action: Bell *et al.*, *Biochemistry* 11, 1959 (1972). Conformation in soln: E. J. Valente *et al.*, *J. Med. Chem.* 20, 1849 (1977); 21, 141, 231 (1978). Human metabolism: R. J. Lewis, W. F. Trager, *Ann. N.Y. Acad. Sci.* 179, 205 (1971). Stereospecific HPLC determ in plasma: C. Banfield, M. Rowland, *J. Pharm. Sci.* 72, 921 (1983). Antimetastatic effect in lung cancer: L. R. Zacharski *et al.*, *Cancer* 53, 2046 (1984); in rat adenocarcinoma: B. L. Neubauer *et al.*, *J. Urol.* 135, 163 (1986). Toxicity studies: E. C. Hagan, J. L. Radomski, *J. Am. Pharm. Assoc.-Sci. Ed.* 42, 379 (1953); N. Back *et al.*, *Pharmacol. Res. Commun.* 10, 445 (1978). Review of therapeutic uses: J. V. Lloyd, *Med. J. Aust.* 142, 197-201 (1985); and pharmacology: J. Hirsh *et al.*, *Chest* 102, Suppl., 312S-326S



(1992). Comprehensive description: S. A. Babhair *et al.*, *Anal. Profiles Drug Subs.* 14, 423-452 (1985).

Crystals from alc, mp 161°. uv max (water, pH 10): 308 nm (ϵ 13610). Soluble in acetone, dioxane. Moderately sol in methanol, ethanol, isopropanol, some oils. Freely sol in alkaline aq solns (forms a water-soluble sodium salt). Practically insol in water, benzene, cyclohexane, Skellysolves A and B. Warfarin has an acidic enol which forms metallic salts and an acetate, mp 117-118°, and a ketone which forms an oxime, mp 182-183° and a 2,4-dinitrophenylhydrazone, mp 215-216°.

Sodium salt, C₁₉H₁₅NaO₄, *Coumadin*, *Marevan*, *Panwarfin*, *Prothromadin*, *Tintorane*, *Warfilone*, *Waran*. Slightly bitter, crystalline powder. Discolored by light. Very sol in water; freely sol in alcohol; very slightly sol in chloroform, ether. LD₅₀ in male rats, female rats, mice, rabbits (mg/kg): 323, 58, 374, ~800 orally (Hagen); also reported as LD₅₀ in male, female rats (mg/kg): 100.3, 8.7 orally (Back).

Potassium salt, C₁₉H₁₅KO₄, *Athrombin-K*.

Compound with 2-(dimethylamino)ethanol, C₂₃H₂₇NO₃, *warfarin-deanol*, *MD-6134*, *Adoisine*.

Caution: Potential symptoms of overexposure are hematuria, back pain; hematoma of arms and legs; epistaxis, bleeding lips and mucous membrane hemorrhage; abdominal pain, vomiting and fecal blood; petechial rash; abnormal hematologic indices. See *NIOSH Pocket Guide to Chemical Hazards* (DHHS/NIOSH 90-117, 1990) p 224. See also *Clinical Toxicology of Commercial Products*, R. E. Gosselin *et al.*, Eds. (Williams & Wilkins, Baltimore, 5th ed., 1984) Section III, pp 395-397.

USE: Rodenticide.

THERAP CAT: Anticoagulant.

10175. Water. Hydrogen oxide. H₂O; mol wt 18.02. H 11.19%, O 88.81%. Reviews: N. E. Dorsey, *Properties of Ordinary Water-Substance*, A.C.S. Monograph Series no. 81, (Reinhold, New York, 1940) 673 pp; D. Eisenberg, W. Kauzmann, *The Structure and Properties of Water* (Oxford University Press, New York, 1969) 296 pp; Ebsworth *et al.*, in *Comprehensive Inorganic Chemistry* vol. 2, J. C. Bailar, Jr. *et al.*, Eds. (Pergamon Press, Oxford, 1973) pp 741-747.

Pyrogen-free water (water for injection) is distilled water rendered free of fever-producing proteins (bacteria and their metabolic products). Method of prepn: Ishizuka *et al.*, *C.A.* 49, 15177 (1955).

Liquid. Temp of max density 3.98°. d₄^{3.98} 1.000000 g/ml (0.999972 g/cc). d₄⁰ 0.997. d₀⁰ 0.917 g/cc; d₄⁰ (liq) 0.999868. Density tables: Bigg, *Brit. J. Appl. Phys.* 18, 521 (1967); Kell, *J. Chem. Eng. Data* 12, 66 (1967). Expands on freezing. mp 0°. bp 100°. One liter satd vapor weighs 0.5974 g at 100° and 760 mm. Crit temp 374.2°; crit pressure 218 atm. Sp. heat (liq; 14°) 1.000 cal/g/°C. Latent heat of fusion: 1.436 kcal/mole. Latent heat of vaporization: 9.717 kcal/mole. n_D²⁰ 1.3330. Dielectric const (0°) 87.740. Dipole moment (25°) in benzene 1.76; in dioxane 1.86. Ionization const for pure water only: K (25°) 1.008 × 10⁻¹⁴; at moderate concn of solutes (e.g. 1.0M KOH): K (25°) 0.971 × 10⁻¹⁴. The most universal solvent known.

10176. Water Gas. Blue gas. Obtained by blowing steam through incandescent coke. *Composition:* 6% CO₂; 42% CO; 51% H₂; 1% N₂.

Caution: Asphyxiant. USE: In the manuf of ammonia as source of hydrogen. Cf. Producer Gas.

10177. Watermelon. Arbuse. *Citrullus vulgaris* Schrad., *Cucurbitaceae*, cultivated in hot and temperate zones the world over. Contains diuretic principles: Bliss *et al.*, *Am. J. Pharm.* 105, 53 (1933); Roby *et al.*, *ibid.* 111, 68 (1939).

10178. Wheat Germ Oil. Cav-Ecol; Myopone; Denamone. Obtained by hydraulic expression or solvent extraction of wheat germ which constitutes ~2% of a wheat grain, the seed of *Triticum aestivum* L. (*T. sativum* Lam., *T. vulgare* Vill.), *Gramineae*. *Constit.* (of the oil): Linoleic acid 44.1%, oleic acid 30.0%, satd acids 15.1%, linolenic acid 10.8%, unsaponifiable matter 4.7%. The unsaponifiable matter contains vitamin E-active tocopherols (reported as 0.5% of the oil and as 2 international vitamin E units per

Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs

E. LAVY*

G. ZIV†

M. SHEM-TOV‡

A. GLICKMAN§ &

A. DEY¶

*Koret School of Veterinary Medicine, The Hebrew University of Jerusalem, P.O. Box 12, Rehovot 76100, Israel, ‡Vetgenerics Research G. Ziv Ltd., P.O. Box 2473 Rehovot 76124, Israel, §Ministry of Agriculture, Kimron Veterinary Institute, P.O. Box 12, Bet-Dagan 50250, Israel, and ¶Avind Chemie GmbH, Hannover, Germany, †passed away November 1997

Lavy, E., Ziv, G., Shem-Tov, M., Glickman, A., Dey, A. Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs *J. vet. Pharmacol. Therap.* 22, 261–265.

A buffered aqueous solution of clindamycin HCl (200 mg/mL) was injected intravenously (i.v.) intramuscularly (i.m.) and subcutaneously (s.c.) in a non-randomized, partial cross-over trial involving six male and six female dogs. Blood samples were collected at conventional, predetermined time periods and serum drug concentrations were determined by microbiological assay. Dogs were observed clinically for signs of pain, and activity of serum creatine phosphokinase (CPK) was monitored after i.m. dosing.

The i.v. data from five of the dogs best fitted a two-compartment open-system pharmacokinetic model whereas a non-compartment model was most suitable for analysis of the data from the remaining seven dogs. The mean i.v. elimination half-life ($t_{1/2\beta}$) and the mean residence time (MRT) were 124 and 143 min, respectively. The mean volume of distribution at steady state (V_{ss}) was 0.86 L/kg. Little pain was recorded upon i.m. injection; mean peak serum drug concentration (C_{max}) was 4.4 µg/mL, the elimination half-life ($t_{1/2el}$) was 247 min and the calculated bioavailability (F) was 115% of the i.v. dose. Serum CPK activity was elevated to 25-fold the pretreatment level in samples collected 4, 8 and 12 h after i.m. injection. Pain was not recorded after s.c. drug administration; the mean C_{max} of 20.8 µg/mL was significantly greater than the corresponding value for the i.m. route, and F was 310%. The s.c. route appears to be superior to the i.m. route in terms of local tolerance and serum drug level; a 10 mg/kg SID treatment regimen is suggested for treatment of canine infections due to clindamycin sensitive bacteria.

(Paper received 16 December 1998; accepted for publication 11 May 1999)

E. Lavy, Koret School of Veterinary Medicine, The Hebrew University, P.O. Box 12, Rehovot 76100, Israel.

INTRODUCTION

A semisynthetic derivative of lincomycin, clindamycin has been shown to be clinically effective and is recommended for treatment of staphylococcal and anaerobic infections of skin, soft tissue and bone in dogs (Berg *et al.*, 1984; Greene, 1989; Braden *et al.*, 1988; Braden *et al.*, 1987). Clindamycin is available for parenteral administration as the 2-phosphate and hydrochloride. Clindamycin 2-phosphate is microbiologically inactive, but is hydrolyzed *in vivo* to clindamycin (Webber *et al.*, 1980). The pharmacokinetics of clindamycin phosphate in dogs were studied after single intravenous (i.v.) and intramuscular (i.m.) administrations at 11 mg/kg clindamycin (Webber *et al.*, 1980) and after single subcutaneous (s.c.) injections at 2.75, 5.5, 11 and 21 mg/kg clindamycin (Webber *et al.*, 1980). Based on the pharmacokinetics of the drug, s.c. dosage regimen of 11 mg/kg of clindamycin free base as clindamycin-2-phosphate/kg body weight every 24 h was recommended (Budsberg *et al.*, 1992). The i.m. administration of clindamycin-2-phosphate solution (50 mg/mL) induced signs of

pain and other side-effects and, therefore, this route was not recommended (Budsberg *et al.*, 1992). A buffered 20% aqueous solution of clindamycin hydrochloride (200 mg/mL) is available for pharmacokinetic and clinical testing. The purpose of this study was to determine the concentrations of clindamycin in normal canine serum after single i.v., i.m. and s.c. administrations of clindamycin HCl and compare the derived kinetic variables with those obtained earlier in dogs injected with equal doses of clindamycin phosphate (Webber *et al.*, 1980; Budsberg *et al.*, 1992). Local tolerance and appearance of side-effects following i.m. and s.c. administrations were particularly examined.

MATERIALS AND METHODS

Animals

Twelve adult mixed breed dogs, six males and six females (4–13 kg b.w.) were used in the study. All dogs were housed in the test facility for 3 weeks prior to the study. Dogs had free access to

water and commercial dry dog ration before and during the study. Inclusion criteria included normal findings on physical examination, complete blood count (CBC) serum concentrations of urea nitrogen, creatinine, albumin, total protein glucose, bilirubin, triglycerides, cholesterol, calcium, magnesium, sodium, potassium, chlorides, inorganic phosphorus and serum activities of alkaline phosphatase, alanine aminotransferase, aspartate aminotransferase and amylase were determined. Dogs selected for the study exhibited normal CBC and blood biochemistry test results.

Experimental design

Intravenous protocol

Jugular vein catheters were placed in each dog; patency of each catheter was maintained with heparinized saline. A blood sample was taken prior to the beginning of the trial. Each dog was given the 20% buffered aqueous clindamycin HCl i.v. at 10 mg/kg. Blood samples were obtained at 10, 20, 30, 40, 50, 60, 80, 90, 120, 180, 240, 360, 480 and 600 min post injection. Blood was allowed to clot at 20°C for 2 h and was then centrifuged at 1000 × g; the serum was collected and stored at -20°C until it was assayed.

Intramuscular and subcutaneous protocols

A 2 week rest period was allowed for all dogs. All indwelling jugular venous catheter was placed and maintained as for the i.v. protocol, and a baseline blood sample was taken. The injection site (4 × 4 cm² area) on the dorsal aspect of the left and right sites were then shaved to remove short hair. Nine dogs received a single i.m. injection of the 20% clindamycin HCl at 10 mg/kg in the left-side of the neck and the remaining three dogs received a single i.m. injection of 3–5 mL sterile physiological saline in the neck. Two weeks later, nine dogs were prepared by procedures identical to those used before i.m. drug administration. Six dogs received a single s.c. injection of 20% clindamycin HCl at 10 mg/kg in the right-side of the neck. All dogs were observed immediately following i.m. and s.c. injections for evidence of pain, itching or irritation. The injection sites on both sides of the neck were palpated at each blood sampling time and at least two-times per day on the following 2 days and any abnormal finding such as pain, swelling and discoloration were recorded.

Blood samples were obtained at 15, 30, 60, 90, 150, 210, 270, 390, 510, 630, 720 and 1440 min post i.m. injection. Blood samples were collected at 30, 45, 60, 90, 120, 180, 240, 360, 480, 600, 720 and 1440 min post s.c. injection. Blood samples were processed as for after i.v. injection.

Clindamycin analysis

Clindamycin concentrations were measured by microbiological well/agar plate diffusion assay as previously described (Bennett *et al.*, 1966). The assay organism *S. lutea* ATCC 9341, was inoculated into antibiotic Medium No. 1 (Difco, Detroit, MI, USA) and a 7.0 mm layer seeded medium was added to each Petri Plates. Six wells, 8 mm in diameter, were cut into the agar at equal distances. A 50.0 µL aliquot of samples (and standard clindamycin HCl solution) was alternately added to each well and

the plates were incubated at 37°C for 14–16 h. The concentration of drug in each sample was calculated from zone of inhibition diameters using polynomial regression techniques. Sensitivity limit of assay method was 0.1 µg/mL. Standard curves were derived using clindamycin HCl (Sigma Chemical Co. St. Louis, MO, USA) in dog serum. The correlation coefficient of the standard curve from 0.10 to 6.0 µg/mL was 0.99 ($P < 0.001$). Samples with concentrations > 6.0 µg/mL were diluted with antibiotic-free dog serum to bring clindamycin concentrations within the range of the standard curve. The coefficients of variation of repeatedly assayed samples at concentrations ranging between 1–6 µg/mL and 0.1–1.0 µg/mL were 7.5% and 12.5%, respectively. Samples were assayed in duplicate and data are reported as mean ± SD. It was recognized that this assay fails to distinguish between clindamycin and its putative active metabolites and, therefore, results were expressed as serum clindamycin antimicrobial equivalent activity. Thus the term 'clindamycin concentration' where used throughout this report is rather clindamycin antimicrobial equivalent activity.

Serum creatine phosphokinase (CPK)

Serum CPK values were determined, using as enzymatic method (CK-NAC-active creatine kinase EC2.7.3.2, Randox Laboratories Ltd., Crumlin, Northern Ireland), in blood samples collected at 0, 4, 8, 12, 24, 32, 48 and 72 h after nine dogs were injected i.m. with 20% clindamycin HCl solution, three dogs were injected with saline, three dogs were injected s.c. with 20% clindamycin HCl and three dogs were administered saline s.c. As large differences in pretreatment CPK values were found among the dogs examined, serum CPK data were converted to percentage by dividing each post treatment value by the pretreatment value for the corresponding animal. The post treatment CPK data are presented as mean ± SD-fold rise from pretreatment (baseline) CPK value.

Data analysis

Estimates of first-order rate constants and volumes were initially obtained by subjecting mean data to analysis, using iterative least squares regression analysis (Brown & Manno, 1978). The concentrations vs. time data from each dog were then analysed, using a microcomputer program for nonlinear weighted least square regression (Bourne, 1986).

The most appropriate pharmacokinetic model was selected on the basis of the lowest weighted sum of squares and the lowest Akaike's information criterion (AIC) value (Yamaoka *et al.*, 1978) for data from each dog. The i.v., i.m. and s.c. areas under the curves (AUCs) were calculated using trapezoidal approximations between time of drug administration and 1440 min afterwards. Differential calculus methods (Edwards & Penney, 1982) were used to estimate peak serum drug concentrations (C_{max}) and time of C_{max} (t_{max}) after i.m. and s.c. administrations. Kinetic values are presented as mean ± SD; half lives, however, are presented as harmonic mean ± pseudo-SD (Lam *et al.*, 1985). The paired Student's *t*-test was used for calculating the significance of the differences in the mean kinetic values for the i.m. and s.c. routes; $P < 0.05$ value was considered significant.

RESULTS

Clinical signs indicative of slight pain were noticed in four to five of the dogs immediately following i.m. injection; the remaining dogs did not exhibit any pain reaction. Signs suggesting pain or discomfort were not shown by any dog after s.c. injection. Palpation of the injection site did not elicit any pain reaction. Local changes could not be felt at the injection site. Thus, the neck side injected with clindamycin HCl could not be differentiated from the side injection with sterile saline solution.

Mean serum clindamycin concentrations after i.v., i.m. and s.c. administrations are presented (Table 1). Data are also presented graphically as mean log₁₀ serum concentrations vs. time plot (Fig. 1). The i.v. data from five of the dogs best fitted a two-compartment open system pharmacokinetic whereas a one-compartment model was most suitable for analysis of data from the remaining seven dogs. Thus, the kinetic values C_p^0 , A , and $t_{1/2\alpha}$ presented (Table 2) represent data from these five dogs; it shows a rapid rate of drug distribution from the central to the peripheral body compartment. The elimination half-life ($t_{1/2\beta}$) and the mean residence time (MRT) were 124.0 ± 57.0 min and 143.0 ± 34.0 min, respectively and the steady state volume of distribution (V_{ss}) was 0.86 ± 0.35 L/kg. After i.m. drug administration, the mean C_{max} (4.4 ± 0.5 µg/mL) was significantly ($P < 0.05$) lower than the corresponding value for the s.c. administration (20.8 ± 6.2 µg/mL).

The mean absorption time (MAT) of clindamycin HCl solution injected s.c. was significantly shorter than after i.m. administration. The mean $t_{1/2el}$ i.m. value (427.0 ± 209.0 min) was not significantly different from the mean $t_{1/2el}$ for the s.c. route (310.2 ± 190.4 min) but these values were significantly longer than the mean i.v. $t_{1/2\beta}$. The mean s.c. AUC was significantly larger than the mean i.m. AUC and the resulting calculated bioavailability (F) values which were 1.15 and 3.1 for the i.m. and s.c. routes, respectively (Table 3).

Serum CPK activity rose sharply within 8 h after i.m. injection of clindamycin HCl; activity returned to pretreatment level by 48 h post treatment. A minimal rise in serum CPK activity was observed after s.c. clindamycin injection. The i.m. and s.c. administration of saline did not affect serum CPK activity.

DISCUSSION

The clinical manifestations of pain described (Budberg *et al.*, 1992) following i.m. administration of 5% solution of clindamycin phosphate to dogs were not seen at all after i.m. injection of more concentrated (20%) buffered aqueous solution of clindamycin HCl. We can only speculate on the causes for these differences in local tolerance; they could be due to the type of clindamycin salt, the presence of buffer or a four-fold smaller volume injected using the 20% clindamycin HCl solution. The transient rise in serum CPK activity observed after i.m. injection of clindamycin HCl to dogs (Fig. 2) indicates some degree of muscle tissue damage at the injection site (Steinnes *et al.*, 1978). However, a similar or even higher and more persistent rise has been documented for a long

Table 1. Mean serum clindamycin concentrations (µg/mL) after intravenous, intramuscular and subcutaneous injection of clindamycin HCl to dogs at 10 mg/kg body weight

Time (min)	Treatment					
	Intravenous		Intramuscular		Subcutaneous	
	Mean	SD	Mean	SD	Mean	SD
10	13.4	2.3	NS		NS	
15	NS		2.6	0.76	NS	
20	11.3	2.2	NS		NS	
30	10.2	2.3	3.6	0.90	16.8	12.8
40	8.9	2.5	NS		NS	
45	NS		NS		19.5	3.8
50	7.7	1.8	NS		NS	
60	6.8	1.4	4.0	0.60	17.4	6.7
80	5.75	0.6	NS		NS	
90	5.6	1.3	4.0	0.70	13.5	4.1
120	4.1	0.87	NS		12.6	4.6
150	NS		3.5	0.50	NS	
180	3.0	0.7	NS		10.7	3.2
210	NS		3.0	0.40	NS	
240	2.1	0.45	NS		6.9	2.2
270	NS		2.3	0.40	NS	
360	0.82	0.22	NS		4.6	1.4
390	NS		1.7	0.40	NS	
480	0.46	0.10	NS		3.3	1.8
510	NS		1.1	0.40	NS	
600	0.31	0.11	NS		2.7	1.6
630	NS		0.70	0.30	NS	
720	NS		0.60	0.30	1.7	1.2
1440	NS		0.30	0.10	0.3	0.1




Fig. 1. Serum clindamycin concentrations (µg/mL, log₁₀ scale) after intravenous, intramuscular and subcutaneous administration of clindamycin HCl to dogs at 10 mg/kg.

list of approved veterinary injectable products which are very commonly used in small and large animal practice without any observable pain reactions (Rasmussen, 1980; Svendsen, 1983). A better safety evaluation of i.m. clindamycin HCl therapy must wait until data from multiple injections are available. The present

Table 2. Selected pharmacokinetic values for clindamycin HCl administered intravenously to 12 dogs at 10 mg/kg


Kenetic value & unit	Mean	SD
Cp°, µg/mL	18.75	3.71
A, µg/mL	11.06	3.35
B, µg/mL	7.54	3.35
$t_{1/2\alpha}$, min	11.00	13.30
$t_{1/2\beta}$, min	124.00	57.00
MRT, min	143.00	34.00
V_c , L/kg	0.56	0.11
V_{ss} , L/kg	0.86	0.35
AUC, µg/mL.min	1457.00	280.00
Cl_t , mL/min/kg	6.10	1.10
r^2	0.967	0.031

A = ordinal intercept of fastest disposition slope minus the intercept of the slowest disposition slope; B = ordinal intercept of the slowest disposition slope; Cp° = initial serum concentration; $t_{1/2\alpha}$ = distribution half-life; $t_{1/2\beta}$ = elimination half-life; MRT = mean residence time; V_c = volume of the central compartment; V_{ss} = volume of distribution at steady state; AUC = area under the concentration-time curve from zero to 24 h post-treatment; Cl_t = total body clearance; r^2 = correlation coefficient to the line of best fit for a two-compartment open system pharmacokinetic.

Table 3. Selected pharmacokinetic values for clindamycin administered intramuscularly and subcutaneously to dogs at 10 mg/kg

Kinetic value and unit	Treatment			
	Intramuscular <i>n</i> = 9		Subcutaneous <i>n</i> = 6	
	Mean	SD	Mean	SD
C_{max} , µg/mL	4.4	0.5	20.8	6.2
t_{max} , min	73.0	16.0	46.7	20.1
MAT, min	546.0	226.0	224.5	163.5
$t_{1/2el}$, min	427.0	209.0	310.2	190.4
MRT, min	700.0	246.0	364.2	147.3
AUC, µg/mL min	1806.0	346.0	5258.0	2161.0
F^*	1.15	0.19	3.10	0.22

C_{max} , peak maximal serum concentration; t_{max} , time to peak serum concentration; MAT = mean absorption time, calculated as $MRT_{non-i.v.}$; $MRT_{i.v.}$; F^* = bioavailability, calculated as $AUC_{non-i.v.}/AUC_{i.v.}$.

**Fig. 2.** Serum CPK activity in dogs after intramuscular administration of clindamycin HCl at 10 mg/kg.

findings clearly indicate that the s.c. route is superior to the i.m. route in terms of local tolerance.

Interpretation of data gathered in the course of the present study must take into consideration the assay method used (microbiological). Although a good agreement was shown between microbiological and chemical (GC) test results in dog serum for clindamycin (Ziv & Shem-Tov, unpublished data), there is a slight chance that there are some putative active metabolites. The disposition curves after i.v. administration of clindamycin HCl was best represented as a two-compartment open model in only five of the dogs. The i.v. study protocol we used called for collecting the first post treatment blood sample at 10 min.

Because of the rapid distribution rate of the drug in the dog ($t_{1/2\alpha}$ of 3.5 ± 1.1 min) according to (Budberg *et al.*, 1992), we probably missed observing the distribution phase. Values for the other major kinetic parameters found in the present study were also different from the values calculated in dogs injected i.v. with clindamycin phosphate (Budberg *et al.*, 1992). Thus, mean $t_{1/2\beta}$, MRT and AUC after clindamycin phosphate administration were 194.6 min, 263.4 min and 2009.5 µg·mL, respectively. Such differences in kinetic values, although small, could result from the rate of appearance of bioactive antibiotic in the serum after i.v. administration of the microbiologically inactive clindamycin phosphate, differences in body-weight (clindamycin phosphate was injected to dogs weighting 20 to 30 kg), (Budberg *et al.*, 1992) or slightly different methods used for calculating the kinetic variable. On the other hand, mean Cl_t , V_c , and V_{ss} for clindamycin phosphate were very close to the corresponding values for clindamycin HCl estimated in the present study. Regardless of these small differences, the large V_{ss} of clindamycin indicates possible wide distribution in the body fluids and tissues. Direct measurements of tissue clindamycin concentrations in humans (Panzer *et al.*, 1972; Dhawan & Thadepalli, 1982) and cats (Brown *et al.*, 1990) confirmed these assumptions.

The kinetic variables calculated from the i.m. serum drug level data for clindamycin HCl and clindamycin phosphate were in good agreement. The short t_{max} (1h) and average bioavailability of nearly 100% support rapid and complete absorption of the drug from the site of i.m. injection, as was remarked earlier (Budberg *et al.*, 1992). The kinetic profile of the drug in serum of all dogs after s.c. administration of clindamycin HCl is rather unique; mean C_{max} (20.8 µg/mL) was nearly 4.5 times greater than the mean i.m. C_{max} . Moreover, mean serum concentrations during the first 12 h post treatment by the s.c. route were two to three-fold higher than the concentrations found after i.m. drug administration (Fig. 1). After a nearly equivalent dose of the drug (11 mg/kg) was injected s.c. to dogs as clindamycin phosphate (Webber *et al.*, 1980) a mean C_{max} of 6.1 ± 0.3 µg/mL was recorded at t_{max} of 40–60 min. We found that the terminal elimination rate of the drug from serum ($t_{1/2el}$) after s.c. administration of 20% clindamycin HCl solution (310.2 ± 190.4 min) was considerably longer than the reported (Budberg *et al.*, 1992) $t_{1/2el}$ of 234.8 ± 27.3 min after an equivalent dose was given to dogs i.m. as clindamycin phosphate. A $t_{1/2el}$ of 13.9 h was calculated (Webber *et al.*, 1980) from the serum clindamycin data of dogs injected s.c. with clindamycin phosphate.

It appears therefore, that s.c. administration of clindamycin HCl allows for rapid, complete drug absorption and, at the same time, acts as a depot which limits the rate of drug elimination from serum. A entero-hepatic circulation effect was suggested to operate in dogs treated orally with clindamycin HCl (Lavy *et al.*, 1999) contributing to the prolongation of $t_{1/2el}$ to nearly 6 h, and for calculated oral bioavailability values exceeding 100%. Whatever the pharmacokinetic processes involved, it appears from the present study that the s.c. route is superior to the i.m. in practical terms by permitting a longer treatment interval.

Earlier studies (Braden *et al.*, 1987; Budsberg *et al.*, 1992) attempted to establish i.v. and i.m. dosing recommendations using average serum concentrations at steady state with the accompanying peak (C_{pmax}) and trough (C_{pmin}) concentrations as means for calculating (Gibaldi, 1982; Riviere, 1988) dosage regimens for clindamycin phosphate in dogs. The calculated dosage schedule was eventually found to be in agreement with the currently recommended oral dosage schedule of 11 mg/kg, q 12 h (Budsberg *et al.*, 1992). We have tried to use a similar approach for selecting desirable, potentially antibacterial effective, serum drug concentrations in dogs given clindamycin HCl by s.c. route. In relating the minimal inhibitory concentration (MIC) of clindamycin to its pharmacokinetic properties it has been assumed (Webber *et al.*, 1980; Budsberg *et al.*, 1992; Brown *et al.*, 1990; Riviere, 1988) that: (a) tissue drug concentration at least equal to the MIC is maintained throughout the entire dose interval; (b) the drug is minimally bound to serum protein and serum concentrations are equal to, or even slightly lower than, the concentration in major target sites of the body (excluding bone); (c) the kinetic profiles of the drug in serum and the target tissue on multiple dosing are very similar; and (d) the MIC for *Staphylococcus aureus/intermedius* ranges from 0.04 to 0.4 µg/mL and for most anaerobic bacteria, the MIC ranges from 0.1 to 3.1 µg/mL but the MIC 90 is in effect > 1.6 µg/mL (Greene, 1989; Budsberg *et al.*, 1992; Brown *et al.*, 1990). Using the mean serum concentration values, we observed that a single s.c. 10 mg/kg SID dosage regimen appears to be appropriate for clindamycin HCl for the treatment of staphylococcal soft tissue infections. For anaerobic infections, however, this treatment should be given BID. A more intensive course of clindamycin therapy is apparently required for the treatment of staphylococcal bone infections in the dog (Braden *et al.*, 1987; Braden *et al.*, 1988; Budsberg *et al.*, 1991).

REFERENCES

Bennett, J.V., Brodie, J.L., Benner, E.J. & Kirby W.M.M. (1996) Simplified, accurate method for antibiotic assay of clinical specimens. *Applied Microbiology*, **14**, 170–175.

Berg, J.N., Scanlan, C.M. & Buening, G.M. (1984) Clinical models for anaerobic bacterial infections in dogs and their use in testing the efficacy of clindamycin and lincomycin. *American Journal of Veterinary Research*, **45**, 1299–1306.

Braden, T.D., Johnson, C.A., Gabel, C.L. *et al.* (1987) Posologic evaluation of clindamycin, using a canine model of posttraumatic osteomyelitis. *American Journal of Veterinary Research*, **48**, 1101–1105.

Braden, T.D., Johnson, C.A., Wakenell, P. *et al.* (1988) Efficacy of clindamycin in the treatment of staphylococcus aureus osteomyelitis in dogs. *Journal of American Veterinary Medical Association*, **192**, 1721–1725.

Bourne, D.W.A. (1986) Multi-forte: a microcomputer program for modeling and simulation of pharmacokinetic data. *Computer Methods Programs Biomedical*, **23**, 277–281.

Brown, R.D. & Manno, J.E. (1978) ESTRIP, a basic computer program for obtaining initial polyexponential parameter estimates. *Journal of Pharmacological Science*, **67**, 1687–1691.

Brown, S.A., Zaya, M.J., Dieringer, T.M. *et al.* (1990) Tissue concentrations of clindamycin after multiple oral doses in normal cats. *Journal of Veterinary Pharmacology and Therapeutics*, **13**, 270–277.

Budsberg, S.C., Gallo, J.M., Starliper, C.E. *et al.* (1991) Comparison of cortical bone local infusion and intravenous administration. *Journal of Orthopaedic Research*, **9**, 594–599.

Budsberg, S.C., Kemp, D.T. & Wolski, N. (1992) Pharmacokinetics of clindamycin phosphate in dogs after single intravenous and intramuscular administration. *American Journal of Veterinary Research*, **53**, 2333–2336.

Dhawan, V.K. & Thadepalli, H. (1982) Clindamycin: a review of fifteen year of experience. *Review of Infectious Diseases*, **4**, 1133–1153.

Edwards, C.H. & Penney, D.E. (1982) *Calculus and analytical geometry*, Englewood Cliffs, Prentice-Hall Inc, NJ.

Gibaldi, M. (1982) Noncompartmental pharmacokinetics. In *Pharmacokinetics*, 2nd ed. Marcel Dekker, New York. pp. 409–417.

Greene, C.E. (1989) Antibacterial chemotherapy. In *Clinical microbiology and infectious diseases in dog and cat*. 2nd ed. WB Saunders Co, Philadelphia. pp. 479–482.

Lam, F.C., Hung, C.T., Perrier, D.G. (1985) Estimation of variance for harmonic mean half-lives. *Journal of Pharmacological Science*, **74**, 229–231.

Lavy, E., Ziv, G., Shem-Tov, M. *et al.* (1999) Oral bioequivalence of clindamycin hydrochloride tablets and clindamycin monohydrate capsules in normal and fasted dogs. *American Journal of Veterinary Research*, (submitted).

Panzer, J.D., Brown, D.C., Epstein, W.L. *et al.* (1972) Clindamycin levels in various body tissues and fluids. *Journal of Clinical Pharmacology*, **12**, 259–262.

Rasmussen, F. (1980) Tissue damage at the injection site after intramuscular injection of drugs in food-producing animals. In *Trends in veterinary pharmacology and toxicology*. Eds. van Miert, A.S.J.P.A.M., Frens, J., van der Kreek, F.W. Elsevier Scientific Publishers, Amsterdam. pp. 27–330.

Riviere, J.E. (1988) Veterinary clinical pharmacokinetics modelling. *Compendium for Continuing Education for Practicing Veterinarians*, **10**, 314–328.

Stennes, E., Rasmussen, F., Svendsen, O. *et al.* (1978) A comparative study of serum creatine phosphokinase (CPK) activity in rabbits, pigs and humans after intramuscular injection of local damaging drugs. *Acta Pharmacologica Toxicologia*, **42**, 357–364.

Svendsen, O. (1983) Intramuscular injections and local muscle: an experimental study of the effect of injection speed. *Acta Pharmacologica Toxicologia*, **52**, 305–309.

Webber, D.J., Barbiere, A.R., Lallinger, A.J. *et al.* (1980) Pharmacokinetics of clindamycin following subcutaneous administration of clindamycin phosphate in the canine. *Journal of Veterinary Pharmacology and Therapeutics*, **3**, 133–143.

Yamaoka, K., Nakagawa, T., Uno, T. (1978) Application of Akaike's information criterion (AIC) in the evaluation of linear pharmacokinetic equation. *Journal of Pharmacokinetics and Biopharmaceutics*, **6**, 165–175.

Disposition Kinetics of Difloxacin After Intravenous, Intramuscular and Subcutaneous Administration in Calves

M. Ismail

Department of Pharmacology, Faculty of Veterinary Medicine, Cairo University,
Giza, Egypt

*Correspondence: E-mail: m_ism@hotmail.com

ABSTRACT

The pharmacokinetics of difloxacin (Dicural) was studied in a crossover study using three groups ($n = 4$) of male and female Friesian calves after intravenous (i.v.), intramuscular (i.m.) and subcutaneous (s.c.) administrations of 5 mg/kg body weight. Drug concentration in plasma was determined by high-performance liquid chromatography using fluorescence detection. The plasma concentration–time data following i.v. administration were best fitted to a two-compartment open model and those following i.m. and s.c. routes were best fitted using one-compartment open model. The collected data were subjected to a computerized kinetic analysis. The mean i.v., i.m. and s.c. elimination half-lives ($t_{1/2\beta}$) were 5.56 ± 0.33 h, 6.12 ± 0.42 h and 7.26 ± 0.6 h, respectively. The steady-state volume of distribution (V_{dss}) was 1.12 ± 0.09 L/kg and total body clearance (Cl_{B}) was 2.19 ± 0.1 ml/(min. kg). The absorption half-lives ($t_{1/2\text{ab}}$) were 0.38 ± 0.027 h and 2.1 ± 0.09 h, with systemic bioavailabilities (F) of $96.5\% \pm 6.4\%$ and $84\% \pm 5.5\%$ after i.m. and s.c. administration, respectively. After i.m. and s.c. dosing, peak plasma concentrations (C_{max}) of 3.38 ± 0.13 $\mu\text{g/ml}$ and 2.18 ± 0.12 $\mu\text{g/ml}$ were attained after (t_{max}) 1.22 ± 0.20 h and 3.7 ± 0.52 h. The MIC_{90} of difloxacin for *Mannheimia haemolytica* was 0.29 ± 0.04 $\mu\text{g/ml}$. The $\text{AUC}/\text{MIC}_{90}$ and $C_{\text{max}}/\text{MIC}_{90}$ ratios for difloxacin following i.m. administration were 120 and 11.65, respectively and following s.c. administration were 97.58 and 7.51, respectively. Difloxacin was 31.7–36.8% bound to calf plasma protein. Since fluoroquinolones display concentration-dependent activities, the doses of difloxacin used in this study are likely to involve better pharmacodynamic characteristics that are associated with greater clinical efficacy following i.m. administration than following s.c. administration.

Keywords: difloxacin, Dicural, female calves, male calves, minimum inhibitory concentration, pharmacodynamics, pharmacokinetics, *Mannheimia haemolytica*

Abbreviations: $t_{1/2\alpha}$, distribution half-life; $t_{1/2\beta}$, elimination half-life (i.v.); k_{12} , first-order rate constant for transfer from central to peripheral compartment; k_{21} , first-order rate constant for transfer from peripheral to central compartment; V_c , apparent volume of the central compartment; V_{dss} , volume of distribution at steady state; Cl_{B} , total body clearance; $\text{AUC}_{0-\infty}$, area under curve from zero time to infinity; MRT, mean residence time; k_{ab} , first-order absorption rate constant; $t_{1/2\text{ab}}$, absorption half-life; k_{el} , first-order elimination rate constant; $t_{1/2\text{el}}$, elimination half-life (i.m. or s.c.); MAT, mean absorption time; C_{max} , maximum plasma concentration; t_{max} , time to peak plasma concentration; F , systemic bioavailability; MIC_{90} , minimum inhibitory concentration; cfu, colony-forming unit

INTRODUCTION

Difloxacin is a difluoroquinolone antimicrobial agent with high *in vitro* activity against a wide range of Gram-positive bacteria, Gram-negative bacteria and mycoplasmas (Digranes and Dibb, 1986; Mader *et al.*, 1987; Fernandes, 1988; Brown, 1996). Similarly to that of other fluoroquinolones, the bactericidal activity of difloxacin is mediated by inhibition of subunit A of DNA topoisomerases II (gyrase), an enzyme that is essential for DNA synthesis and repair (Wolfson and Hooper, 1989; Drlica and Zhao, 1997). Difloxacin differs in particular from other fluoroquinolones on account of its *p*-fluorophenyl ring at position

N-1 of the quinoline nucleus, which reportedly gives it enhanced activity against Gram-positive bacteria (Walker, 2000).

Difloxacin is rapidly absorbed and has high systemic bioavailabilities following oral administration in chickens, pigs (Inui *et al.*, 1998) and dogs (Frazier *et al.*, 2000) as well as i.m. administration in rabbits (Abd el Aty *et al.*, 2005). In addition, it has good distribution characteristics and long elimination half-life in goats (Atef *et al.*, 2002).

Difloxacin is indicated for treatments of bovine respiratory disease caused by *Pasteurella* spp. and or *Mycoplasma* spp. Although the pharmacokinetics of difloxacin has been investigated in a number of species, including chickens, pigs, dogs and goats (Inui *et al.*, 1998; Frazier *et al.*, 2000; Atef *et al.*, 2002; Heinen, 2002; Abd El Aty *et al.*, 2005), there are no reports evaluating the pharmacokinetics of the drug in calves.

The purpose of this study was to investigate the pharmacokinetic and pharmacodynamic correlation following intravenous (i.v.), intramuscular (i.m.) and subcutaneous (s.c.) administration of difloxacin in calves at the dose rate (5 mg/kg body weight) recommended by the manufacturer in many countries.

MATERIAL AND METHODS

Experimental design

A three-period crossover study was undertaken in 6 male and 6 female healthy Friesian calves (10.5 months of age with a mean weight \pm SEM of 257.8 ± 18.7 kg), such that each calf received difloxacin (Dicural 10% injectable solution, Fort Dodge Animal Health, Holland) at a dose of 5 mg/kg body weight by i.v., i.m. or s.c. route. Each calf was housed in an individual pen and fed on antibiotic-free pelleted concentrates (Zagazig Ration, Zagazig Company, El Sharqia, Egypt), hay and alfalfa. Food and water were provided *ad libitum*.

Calves were randomly assigned into three groups, with each group containing four animals, two of each sex. In period 1 of the study, animals of group 1 received 5 mg/kg difloxacin intravenously into the right jugular vein and animals of group 2 received the same dose into the thigh muscle; animals of group 3 received the same dose subcutaneously in the neck region. Subsequent treatments in periods 2 and 3 were administered according to a Latin square design so that each calf received each treatment in sequence.

Blood samples (5 ml/per sample) were obtained by venepuncture of the jugular vein into heparinized tubes just before administration of the drug by different routes and at 5, 10, 15 and 30 min and at 1, 2, 4, 6, 8, 10, 12, 24 and 36 h after i.v., i.m. or s.c. dosing. Samples were allowed to stand protected from light at room temperature (approximately 24°C) for 20 min and then centrifuged at 1500g for 10 min to harvest plasma. Plasma was aliquoted and stored at -70°C pending analysis.

Difloxacin assay

Instrumentation. Drug concentration in plasma was determined using reversed-phase high-performance liquid chromatography (HPLC) according to the method previously described

by Frazier and colleagues (2000). The HPLC system consisted of a Model 616 solvent delivery pump (Waters, Milford, MA, USA), a Waters Model 600 S controller, a Model 717-plus autosampler equipped with a temperature-controlled rack (Waters), and a variable-wavelength fluorescence UV detector (Spectrofluorometric detector Rf 10, Shimadzu)

Chromatographic conditions. The chromatographic conditions included a mobile phase of acetonitrile in 0.05 mol/L sodium phosphate buffer (pH \approx 3.5) (25:75 v/v) at a flow rate of 1 ml/min through a reversed-phase C₁₈ column (Discovery, Supelco, 5 μ m, 4.6 \times 150 mm). Fluorescence detection was at an excitation wavelength of 295 nm and emission wavelength of 500 nm.

Calibration curve. For preparation of the calibration curves, plasma of antibiotic-naive calves was spiked with 0.02, 0.05, 0.1, 0.5, 5 and 10 μ g/ml difloxacin. Quality control samples were prepared in large volume from an independent weighing of reference standards, aliquoted into the same type of vials used for storage of plasma samples, and frozen until the day of the assay. A calibration curve was obtained by plotting the peak height ratio versus the nominal concentrations. The equation was calculated by the least-squares method using linear regression. The minimum quantitative limit of the assay was 0.02 μ g/ml. The standard curve of difloxacin in calf plasma was linear between 0.02 and 5 μ g/ml, the value of correlation coefficients (r) was >0.97 . The peak height ratio of an unknown specimen (peak height of difloxacin/peak height of internal standard) was compared with that of the standard; over the concentration range 0.02–5 μ g/ml, the concentration of difloxacin was directly related to the peak height ratio. Any sample concentrations of difloxacin greater than 5 μ g/ml were diluted 1:1 or 1:10 and the analysis was repeated. Dilution of quality control samples that exceeded the ranges of the standard curve with drug-free plasma was confirmed not to interfere with accurate drug quantification.

Sample extraction. The plasma samples or calibration standards to be assayed (500 μ l) were placed in centrifuge tube and spiked with 50 μ l of internal standard (ofloxacin 5 μ g/ml in 0.05 mol/L phosphate buffer) and vortexed. Dichloromethane (6 ml) was added, samples were vortexed, the aqueous layer was removed by aspiration, and the organic layer was evaporated to dryness. The residue was reconstituted using HPLC mobile phase (250 μ l) and transferred to an autosampler vial for injection.

Validation of the assay method. The precision and accuracy of the method were evaluated by repetitive analysis of the plasma samples ($n = 12$) spiked with 0.02, 0.05, 0.1, 0.5 and 5 μ g/ml difloxacin. Intra-day precision and accuracy were obtained by analysis of these samples on one day by the same operator. Inter-day precision and accuracy were obtained by assay of these samples ($n = 12$) on different days by two operators. Stability of analytes was determined by comparing peak heights in quality control samples with those in freshly prepared standards in plasma. The recoveries were calculated by comparison of plasma and aqueous samples ($n = 6$).

The intra-assay coefficient of variation for plasma was <3.2% and the intra-assay accuracy was >94.7%. The inter-assay coefficient of variation for plasma was <4.1% and the inter-assay accuracy was >95%. Recovery of difloxacin from plasma was found to be 94%.

Estimation of protein binding

The extent of plasma protein binding was determined *in vitro* by a method previously described (Singhvi *et al.*, 1977). Plasma from each calf was spiked with 0.05, 0.1, 0.5, 1, 5 and 10 µg/ml difloxacin and 1 ml was added to a commercial ultrafiltration device (Centrifree 4104, Amicon Corp., Danvers, MA, USA). The ultrafiltration device was centrifuged at a fixed angle (28°) (Sorvall, RC-5B Refrigerated super speed centrifuge, GSA rotor, DuPont Instruments, Newtown, Connecticut) at 1200g for 30 min at 37°C. This resulted in an ultrafiltrate volume of at least 200 µl. The ultrafiltrate was frozen until assayed for difloxacin. The percentage of protein-bound fraction (*B*) was calculated according to the equation $B = \frac{[\text{initial plasma (difloxacin)} - \text{ultrafiltrate (difloxacin)}]}{[\text{initial plasma (difloxacin)}]} \times 100$. The coefficients of variation for this method were <4.7%.

Minimum inhibitory concentration

The minimum inhibitory concentrations (MIC₉₀) of difloxacin for *Mannheimia haemolytica* isolated from diseased calves (12 isolates) were determined by broth microdilution technique (Jones *et al.*, 1985). Ten replicates of twofold dilutions of difloxacin (0.062, 0.125, 0.25, 0.5, 1.0, 2.0 µg/ml) were used. Fifty microlitres of each concentration was added to each well. One well in each row contained only Muller–Hinton (MH) broth to serve as an inoculation and growth control. Clinical isolates (24-hour-old cultures) were subcultured to MH broth at a density of approximately 10⁸ colony-forming units (cfu)/ml, compared with density of a 0.5 McFarland standard. This suspension was further diluted to 10⁵ cfu/ml in MH broth. Fifty microlitres of the suspension was delivered to each well. The MIC was the lowest concentration of difloxacin for which no visible growth was observed after 18 h of incubation at 37°C. The coefficients of variation for this method were <3.6%.

Pharmacokinetic analysis

Following intravenous administration, the plasma concentration–time data were fitted to a two-compartment open model system (Baggot, 1978) according to the biexponential equation $C_t = Ae^{-\alpha t} + Be^{-\beta t}$, where C_t is the plasma concentration of difloxacin; t is time after intravenous administration; A and α are the intercept and slope, respectively, of the distribution phase; B and β are the intercept and slope of the elimination phase; and e is the base of natural logarithms. Pharmacokinetic variables were obtained by use of a computer program (R Strip, Micromath, UT, USA). The distribution and elimination half-lives ($t_{1/2\alpha}$ and $t_{1/2\beta}$), the volume of distribution at steady state (V_{dss}), the volume of the central compartment (V_c), the total body clearance (Cl_B) and the two-compartment

microconstants k_{12} , k_{21} were computed according to standard equations (Gibaldi and Perrier, 1982).

Following i.m. and s.c. administration, plasma concentration data were analysed by compartmental and non-compartmental methods based on statistical moment theory (Gibaldi and Perrier, 1982). In compartmental analysis, data were found to be best fitted using a one-compartment open model and first-order absorption rate constant according to the equation $C_t = Ee^{-k_{ab}t} - Ce^{-k_{el}t}$, where C_t is the plasma concentration of difloxacin; t is time after i.m. or s.c. administration, k_{el} is the elimination rate constant and k_{ab} is the first-order absorption rate constant. The terminal elimination half-life ($t_{1/2el}$) and absorption half life ($t_{1/2ab}$) were calculated as $(\ln 2)/k_{el}$ or $(\ln 2)/k_{ab}$, respectively. The area under the plasma concentration–time curve ($AUC_{0-\infty}$) and the area under the first moment curve ($AUMC_{0-\infty}$) were calculated by the trapezoidal rule for all measured data with extrapolation to infinity using C_{36}/k_{el} or C_{36}/β , where C_{36} is the plasma concentration at 36 h divided by the β (elimination rate constant for two-compartment model) or k_{el} in the case of the one-compartment open model. The mean residence time (MRT) was calculated as $MRT = AUMC_{0-\infty}/AUC_{0-\infty}$. The mean absorption time (MAT) was calculated as $MAT = MRT_{(s.c.cri.m.)} - MRT_{(i.v.)}$. The peak plasma concentration (C_{max}) and time to maximum concentration (t_{max}) were taken from the plot of each calf's concentration–time curve. Bioavailability (F ; fraction of drug absorbed systemically) was calculated as $F = AUC_{(i.mors.c.)}/AUC_{i.v.} \times 100$.

Pharmacodynamic efficacy of difloxacin was determined by calculation of C_{max}/MIC_{90} and $AUC_{0-\infty}/MIC$ ratios following i.m and s.c. administration of the drug using MIC_{90} for *Mannheimia haemolytica* determined in the present study.

Statistical analysis

Results are presented as mean \pm standard error (SE). All data were subjected to analysis of variance with a significant level of 0.05 and the homogeneity of variances was tested by Bartlett test. Testing for homogeneity of the variances suggested the use of the non-parametric Mann–Whitney test for comparing the pharmacokinetic parameters after i.m. and s.c. administration, p -values <0.05 were considered significant. All the previously mentioned tests were performed using the SPSS 6.1.3 software package (SAS Inc., Cary, NC, USA)

RESULTS

Following i.v administration of difloxacin, the plasma concentration–time curve of the drug was decreased in a biphasic manner, as characterized by a two-compartment open model. Figure 1 illustrates the mean plasma concentrations as a function of time following all routes of administration. Table I summarizes the pharmacokinetic parameters for difloxacin following i.v. administration. The distribution half-life ($t_{1/2\alpha}$) was 0.24 ± 0.021 h and the elimination half life ($t_{1/2\beta}$) was 5.56 ± 0.33 h. The V_{dss} was 1.12 ± 0.09 L/kg and MRT was 7.32 ± 0.56 h.




Figure 1. Semilogarithmic graph depicting the time-plasma concentration course of difloxacin in calves following intravenous, intramuscular and subcutaneous administration of 5 mg/kg body weight ($n = 12$)

TABLE I
Kinetic parameters (mean \pm SE) of difloxacin following a single i.v. injection of 5 mg/kg body weight in calves ($n = 12$)

Parameter ^a	Unit	Value
α	h^{-1}	2.82 ± 0.16
$t_{1/2\alpha}$	h	0.24 ± 0.021
β	h^{-1}	0.12 ± 0.01
$t_{1/2\beta}$	h	5.56 ± 0.33
k_{12}	h^{-1}	1.51 ± 0.07
k_{21}	h^{-1}	1.21 ± 0.08
V_c	L/kg	0.47 ± 0.034
V_{dss}	L/kg	1.12 ± 0.09
Cl_B	$\text{ml}/(\text{min} \cdot \text{kg})$	2.19 ± 0.1
$AUC_{0-\infty}$	$\mu\text{g}/(\text{ml} \cdot \text{h})$	36.0 ± 2.6
MRT	h	7.32 ± 0.56

^a α , β , hybrid rate constants representing the slopes of distribution and elimination phases, respectively; $t_{1/2\alpha}$, distribution half-life; $t_{1/2\beta}$, elimination half-life; k_{12} , first-order rate constant for transfer from central to peripheral compartment; k_{21} , first-order rate constant for transfer from peripheral to central compartment; V_c , apparent volume of the central compartment; V_{dss} , volume of distribution at steady state; Cl_B , total body clearance; $AUC_{0-\infty}$, area under curve from zero time to infinity; MRT, mean residence time

TABLE II
Mean \pm SE kinetic parameters of difloxacin following single i.m. and s.c. injections of 5 mg/kg body weight in calves ($n = 12$)

Parameter ^a	Unit	Value	
		i.m.	s.c.
k_{ab}	h^{-1}	1.78 ± 0.09	$0.33 \pm 0.01^{**}$
$t_{1/2ab}$	h	0.38 ± 0.027	$2.1 \pm 0.09^{**}$
k_{el}	h^{-1}	0.11 ± 0.009	0.09 ± 0.006
$t_{1/2el}$	h	6.12 ± 0.42	7.26 ± 0.6
$AUC_{0-\infty}$	$\mu g/(ml \cdot h)$	34.82 ± 1.9	$28.3 \pm 1.5^{**}$
MRT	h	8.71 ± 0.62	$13.65 \pm 1.1^{**}$
MAT	h	1.38 ± 0.11	$6.33 \pm 0.52^{**}$
C_{max}	$\mu g/mL$	3.38 ± 0.13	$2.18 \pm 0.12^{**}$
t_{max}	h	1.22 ± 0.2	$3.7 \pm 0.52^{**}$
F	%	96.5 ± 6.4	84 ± 5.5

^a k_{ab} , first-order absorption rate constant; $t_{1/2ab}$, absorption half-life; k_{el} , first-order elimination rate constant; $t_{1/2el}$, elimination half-life; $AUC_{0-\infty}$, area under curve from zero time to infinity; MRT, mean residence time; MAT, mean absorption time; C_{max} , maximum plasma concentration; t_{max} , time to peak plasma concentration; F , systemic bioavailability

* $p < 0.01$; ** $p < 0.001$

Following i.m. and s.c. administration, the plasma concentration – time data were best fitted by a one-compartment open model and the drug was detected in plasma for 36 h following administration. Table II displays the pharmacokinetic parameters following i.m. and s.c. administration. The absorption half-life, mean absorption time and mean residence time following i.m. administration ($t_{1/2ab} = 0.38 \pm 0.027$ h; MAT = 1.38 ± 0.11 h; MRT = 8.71 ± 0.62 h) were significantly shorter than those following s.c. administration ($t_{1/2ab} = 2.1 \pm 0.09$ h; MAT = 6.33 ± 0.52 h; MRT = 13.65 ± 1.1 h). A significantly higher maximum plasma concentration was attained earlier following i.m. administration than that following the s.c. route. The elimination half-life ($t_{1/2el}$) and systemic bioavailability were not significantly different between the two routes. The extent of plasma protein binding varied between 31.7% and 36.8% of difloxacin spiked into plasma of antimicrobial-naive calves. The MIC₉₀ for *Mannheimia haemolytica* was 0.29 ± 0.04 $\mu g/ml$. The AUC/MIC₉₀ and C_{max}/MIC_{90} ratios for difloxacin following i.m. administration were 120 and 11.65, respectively, and following s.c. administration were 97.58 and 7.51, respectively.

DISCUSSION

In the present study, serum concentration of difloxacin was determined by HPLC method. Sarafloxacin, the active metabolite of difloxacin in most animal species (Heinen, 2002), was not quantified in the present study because previous investigation had indicated that

sarafloxacin constitutes a very small proportion in serum and its peak (0.033 µg/ml) was close to the lower quantification level in the earlier study as well as in the present study (Chu *et al.*, 1985).

Difloxacin is a well-tolerated drug in calves; no adverse effect was reported following i.v., i.m. or s.c. administration of the drug at a dose of 5 mg/kg body weight. Following i.v administration of difloxacin in calves, the disposition kinetic curve declined in a biphasic manner, suggesting that the drug disposition followed a two-compartment open model. This finding is in agreement with previous reports in goats (Atef *et al.*, 2002). Difloxacin has good distribution characteristics represented by short distribution half-life and large volume of distribution ($t_{1/2\alpha} = 0.24$ hour; $V_{dss} = 1.12$ L/kg). Similar findings have been reported for difloxacin in goats (Atef *et al.*, 2002) and for other fluoroquinolones (enrofloxacin and danofloxacin) in calves (Davidson *et al.*, 1986; Mann and Frame, 1992). The elimination half-life in calves (5.56 h) was slightly shorter than that reported in goats (Atef *et al.*, 2002). The total body clearance of difloxacin (2.19 ml/(min.kg)) in calves was very similar to that in goats (2.16 ml/(min.kg)) (Atef *et al.*, 2002). In contrast, a much higher clearance of difloxacin was reported in rabbits (9.83 ml/(min.kg)) (Abd el Aty *et al.*, 2005). Such differences could be attributable to interspecies variations in drug metabolism and elimination.

The extent of plasma protein binding varied between 31.7% and 36.8% of difloxacin spiked into plasma of antimicrobial-naive calves; a higher plasma protein binding has been reported for difloxacin (46–52%) in humans (Granneman *et al.*, 1986).

Following i.m administration, difloxacin was rapidly absorbed with a short $t_{1/2ab}$ of 0.38 h; rapid absorption is also reflected by short MAT of 1.38 h. Rapid absorption ($t_{1/2ab} = 0.37$ h) following i.m. administration has been reported for difloxacin in goats (Atef *et al.*, 2002). Conversely, difloxacin was slowly absorbed following s.c. administration and MAT was significantly longer as a result of continued absorption from the site of injection. Following i.m. injection, maximum plasma concentration of the drug ($C_{max} = 3.38$ µg/ml) was significantly higher and was achieved earlier ($t_{max} = 1.22$ h) than with s.c. injection ($C_{max} = 2.18$ µg/ml, $t_{max} = 3.7$ h); these values are comparable to that of 3.7 µg/ml attained after 1.25 h following i.m. administration to goats at a dose rate similar to that used in the present study (Atef *et al.*, 2002).

A longer elimination half life for difloxacin was recorded following s.c. administration than following the i.m. route. However, this difference was not significant and could be attributed mainly to the slow release of the drug from the injection site following s.c. injection.

The systemic bioavailability reported in the current study indicates excellent absorption of the drug (96.5%) following i.m. injection. These values are similar to that reported following i.m. administration of the drug in goats (Atef *et al.*, 2002). A smaller proportion of the drug (84%) reached the systemic circulation following administration by the s.c. route.

With i.m. and s.c. administration of difloxacin in calves, plasma concentration was above the MIC₉₀ for *Mannheimia haemolytica* for 12 h following administration. From previous studies with fluoroquinolone and ciprofloxacin, it has been proposed that treatment should be optimized by providing an AUC/MIC ratio of at least 125 (Forrest *et al.*, 1993; Sullivan *et al.*, 1993) and a C_{max} /MIC ratio of 10 or greater. These two breakpoints are essential for

clinical efficacy of fluoroquinolones (Vogelman *et al.*, 1988; Dalhoff and Ullmann, 1990). In the present study, by incorporating C_{\max} and AUC data following i.m. and s.c. administration of difloxacin together with the MIC_{90} of 0.29 for *Mannheimia haemolytica*, the following values were obtained. The AUC/MIC_{90} and C_{\max}/MIC_{90} ratios for difloxacin following i.m. administration were 120 and 11.65, respectively, and following s.c. administration were 97.58 and 7.51, respectively. The reported ratios following i.m. administration are close to the recommended breakpoints, whereas those following s.c. administration fall short of these breakpoints.

Since fluoroquinolones display concentration-dependent activities, the dose of difloxacin used in this study are likely to involve better pharmacodynamic characteristics that are associated with greater clinical efficacy following i.m. administration rather than following s.c. route. Additionally, there is a need for further investigation that correlates the pharmacokinetics and pharmacodynamics of difloxacin in plasma, exudates, transudate and bronchial secretion following s.c. administration in calves naturally infected with *Mannheimia haemolytica* to extend the *ex vivo* data obtained in this study.

REFERENCES

- Abd el Aty, A.M., Goudah, A., Ismail, M. and Shimoda, M., 2005. Disposition kinetics of difloxacin in rabbits after intravenous and intramuscular injection of Dicural. *Veterinary Research Communications*, **29**, 297–304
- Atef, M., El Banna, H.A., Abd El Aty, A.M. and Goudah, A., 2002. Pharmacokinetics of difloxacin in goats. *Deutsche Tierärztliche Wochenschrift*, **109**, 320–323
- Baggot, J.D., 1978. Some aspects of clinical pharmacokinetics in veterinary medicine. *Journal of Veterinary Pharmacology and Therapeutics*, **1**, 5–18
- Brown, S.A., 1996. Fluoroquinolones in animal health. *Journal of Veterinary Pharmacology and Therapeutics*, **19**, 1–14
- Chu, D.T., Granneman, G.R. and Fernandes, P.B., 1985. Abbott-56619. *Drug Future*, **10**, 543–545
- Dalhoff, A. and Ullmann, U., 1990. Correlation between pharmacokinetics, pharmacodynamics and efficacy of antibacterial agents in animal models. *European Journal of Clinical Microbiology and Infectious Diseases*, **9**, 479–487
- Davidson, J.N., Conzleman, G.M. and Baggot, J.D., 1986. Pharmacokinetics of enrofloxacin in preruminant and ruminant calves. *Proceedings of the Western Pharmacological Society*, **29**, 129–132
- Digranes, A. and Dibb, W., 1986. *In vitro* activities of A-56619 (Difloxacin) and A 65520, two aryl fluoroquinolones. *Chemotherapy*, **34**, 298–307
- Drlica, K. and Zhao, X., 1997. DNA gyrase, topoisomerase IV, and the 4-quinolones. *Microbiology and Molecular Biology Review*, **61**, 377–392
- Fernandes, P.B., 1988. Mode of action, and *in vitro* and *in vivo* activities of the fluoroquinolones. *Journal of Clinical Pharmacology*, **28**, 156–168
- Forrest, A., Nix, D.F. and Ballow, A., 1993. Pharmacodynamics of intravenous ciprofloxacin in seriously ill patients. *Antimicrobial Agents and Chemotherapy*, **37**, 1073–1081
- Frazier, D.L., Thompson, L., Terettien, A. and Evans, E.I., 2000. Comparison of fluoroquinolone pharmacokinetic parameters after treatment with marbofloxacin, enrofloxacin and difloxacin in dogs. *Journal of Veterinary Pharmacology and Therapeutics*, **23**, 293–302
- Gibaldi, M. and Perrier, D., 1982. Non compartmental analysis based on statistical moment theory. *Pharmacokinetics*, 2nd edn, (Marcel Dekker, New York), 409–424
- Granneman, G.R., Synder, K.M and Shu, V.S., 1986. Difloxacin metabolism and pharmacokinetics in human after oral doses. *Antimicrobial Agents and Chemotherapy*, **30**, 689–693
- Heinen, F., 2002. Comparative serum pharmacokinetics of enrofloxacin and difloxacin as well as their main metabolites in dogs. *Journal of Veterinary Pharmacology and Therapeutics*, **25**, 1–5
- Inui, T., Taira, T., Matsushita, T. and Endo, T., 1998. Pharmacokinetic properties and oral bioavailabilities of difloxacin in pig and chicken. *Xenobiotica*, **28**, 887–893

- Jones, R.N., Barry, A.L., Govan, T.L. and Washington, J.A., 1985. Susceptibility tests, microdilution and macrodilution broth procedures. In: Lannette, E.H., Alaws, A.B., Hausler, W.J. and Sholdomy, H.J. (eds.), *Manual of Clinical Microbiology* (American Society for Microbiology, Washington, DC), 972–977
- Mader, I.T., Morrison, L.T. and Admas, K.R., 1987. Comparative evaluation of A-56619 (Difloxacin) and Nafcillin in the treatment of experimental *Staphylococcus aureus* osteomyelitis. *Antimicrobial Agents and Chemotherapy*, **34**, 310–316
- Mann, D.D. and Frame, G.M., 1992. Pharmacokinetic study of danofloxacin in cattle and swine. *American Journal of Veterinary Research*, **53**, 1022–1026
- Singhvi, S.M., Heald, A.F., Gadebusch, H.H., Resnick, M.E., Ditazio, L.T. and Leitz, M.A., 1977. Human serum protein binding of cephalosporin antibiotics *in vitro*. *Journal of Laboratory and Clinical Medicine*, **89**, 414–420
- Sullivan, M.C., Cooper, B.W., Nightingale, C.H., Quintiliani, R. and Lawlor, M.T., 1993. Evaluation of the efficacy of ciprofloxacin against *Streptococcus pneumoniae* by using a mouse protection model. *Antimicrobial Agents and Chemotherapy*, **37**, 234–239
- Vogelman, B., Gudmundsson, S. and Leggett, J., 1988. Correlation of antibacterial pharmacokinetic parameters with therapeutic efficacy in animal models. *Journal of Infectious Diseases*, **158**, 831–847
- Walker, R.D., 2000. Fluoroquinolones. In: Prescott, J.F., Baggot, J.D. and Walker, R.D. (eds), *Antimicrobial Therapy in Veterinary Medicine*, 3rd edn (Iowa State University Press, Ames, IA)
- Wolfson, J.S. and Hooper D.C., 1989. Fluoroquinolone antimicrobial agents. *Clinical Microbiology Reviews*, **2**, 378–424

(Accepted: 12 November 2005; Published online: 23 December 2006)

Tamoxifen-resistant Fibroblast Growth Factor-transfected MCF-7 Cells Are Cross-Resistant *in Vivo* to the Antiestrogen ICI 182,780 and Two Aromatase Inhibitors¹

Sandra W. McLeskey, Lurong Zhang, Dorraya El-Ashry, Bruce J. Trock, Cecilia A. Lopez, Samir Kharbanda, Christopher A. Tobias, Lori A. Lorant, Rachel S. Hannum, Robert B. Dickson, and Francis G. Kern²

Lombardi Cancer Center [S. W. M., L. Z., B. J. T., C. A. L., S. K., C. A. T., R. S. H., D. E.-A., R. B. D., F. G. K.], Departments of Biochemistry and Molecular Biology [D. E.-A., F. G. K.], Cell Biology [R. B. D., L. Z.], Medicine [B. J. T.], and Pharmacology [S. W. M.], and the School of Nursing [S. W. M.], Georgetown University Medical Center, Washington, D. C. 20007

ABSTRACT

Although the antiestrogen tamoxifen has been the mainstay of therapy for estrogen receptor (ER)-positive breast cancer, successful treatment of responsive tumors is often followed by the acquisition of tamoxifen resistance. Subsequently, only 30–40% of patients have a positive response to second hormonal therapies. This lack of response might be explained by mechanisms for tamoxifen resistance that sensitize ER pathways to small amounts of estrogenic activity present in tamoxifen or that bypass ER pathways completely. To elucidate one possible mechanism of tamoxifen resistance, we treated ovariectomized tumor-bearing mice injected with fibroblast growth factor (FGF)-transfected MCF-7 breast carcinoma cells with the steroidal antiestrogen ICI 182,780 or one of two aromatase inhibitors, 4-OHA or letrozole. These treatments did not slow estrogen-independent growth or prevent metastasis of tumors produced by FGF-transfected MCF-7 cells in ovariectomized nude mice. FGF-transfected cells had diminished responses to ICI 182,780 *in vitro*, suggesting that autocrine activity of the transfected FGF may be replacing estrogen as a mitogenic stimulus for tumor growth.

ER levels in FGF transfectants were not down-regulated, and basal levels of transcripts for estrogen-induced genes or of ER-mediated transcription of estrogen response element (ERE) luciferase reporter constructs in the FGF expressing cells were not higher than parental cells, implying that altered hormonal responses are not due to down-regulation of ER or to FGF-mediated activation of ER. These studies indicate that estrogen independence may be achieved through FGF signaling pathways independent of ER pathways. If so, therapies directed at the operative mechanism might produce a therapeutic response or allow a response to a second course of antiestrogen treatment.

INTRODUCTION

Because conventional therapy is not usually curative in clinical breast cancer, development of tamoxifen resistance, in which breast tumors previously growth-inhibited by tamoxifen become refractory, represents an important therapeutic dilemma. However, the development of tamoxifen resistance is not necessarily associated with progression to an ER³-negative phenotype. In many cases of clinical tamoxifen resistance, ER expression may be retained (1–4), implying that the resistance is due an alteration in activity of the tamoxifen/ER complex. Tamoxifen resistance in such a case could result from three possible mechanisms that, according to present knowledge, would not preclude successful treatment with an alternative hormonal therapy. First, alterations in the ER could arise, which might diminish or extinguish inhibitory responses to tamoxifen, leaving only its partial agonist effects to predominate (5–8). Second, tamoxifen resistance arising in the setting of an intact ER could be a result of altered intratumoral tamoxifen metabolism, which might produce more estrogenic metabolites locally (7, 9–11). Third, available tamoxifen could be sequestered by an increase in antiestrogen binding sites not associated with ERs (12). As mentioned, in each of these three instances, substitution of a hormonal therapy different from tamoxifen might result in a clinical response. Two such alternative therapies used in this report are steroidal estrogen antagonists, such as ICI 182,780, which lack the partial agonist activity of tamoxifen, and aromatase inhibitors, which inhibit endogenous estrogen production by all tissues, depriving the ER of its ligand.

Although the mechanisms of tamoxifen resistance de-

Received 7/3/97; revised 11/26/97; accepted 12/10/97.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹This work was supported by NIH Grants CA50376 (to F. G. K.), CA09218 (to F. G. K. and S. W. M.), CA53185 (to F. G. K. and R. B. D.), CA66154 (to S. W. M.), CA71465 (to D. E.-A.), and Cancer Center Grant CA51008; American Cancer Society Grant IRG-193 (to S. W. M.); U.S. Army Medical Research and Material Command Grants DAMD 17-94-4172 (to D. E.-A.) and DAMD 17-94-J-4173 (to S. W. M.); and a Susan Komen Foundation Fellowship (to L. Z.).

²To whom requests for reprints should be addressed, at Southern Research Institute, P. O. Box 55305, 2000 Ninth Avenue South, Birmingham, AL 35255-5305. Phone: (205) 581-2480; Fax: (205) 581-2877; E-mail: kern@sri.org.

³The abbreviations used are: ER, estrogen receptor; FGF, fibroblast growth factor; IMEM, improved minimal essential medium; X-gal, 5-bromo-4-chloro-3-indoyl-β-D-galactopyranoside; FBS, fetal bovine serum; 4-OHA, 4-hydroxyandrostenedione; NK, natural killer; CCS, charcoal-stripped calf serum; ERE, estrogen response element; CAT, chloramphenicol acetyltransferase; RT, reverse transcription.

scribed above should be amenable to alternative hormonal therapy, early results for small numbers of tamoxifen-resistant patients have shown that only about 30–40% of such patients have a positive response to subsequent ICI 182,780 or aromatase inhibitor therapy (13–20). These data imply alternative mechanisms for tamoxifen resistance. Constitutive production of autocrine growth factor(s) or growth factor receptors by tumor cells has been proposed as a mechanism for tamoxifen resistance that may or may not involve ER pathways. Evidence supporting this hypothesis is gained from the acquisition of estrogen-independent growth in tumor models, including the one used in this report, in which growth factors or growth factor receptors have been overexpressed in estrogen-dependent breast carcinoma cell lines (21–26). In addition, recent clinical data showing decreased efficacy of tamoxifen in treating tumors overexpressing *c-erbB2* (27) supports a role for growth factor signaling in clinical tamoxifen resistance. Because some growth factor signaling pathways, including the ERB-B pathway, have been shown to interact with ER signaling pathways (25, 28–32), increased growth factor signaling could be one mechanism by which cells could become sensitive to previously ineffective amounts of estrogenic stimulation produced by the partial agonist activity of tamoxifen itself or its estrogenic metabolites, above. In cases in which such interactions have been demonstrated, the growth factor and ER pathways may act collaboratively (25), making the final outcome susceptible to pharmacological manipulations of either pathway and implying that second line hormonal therapies might have an effect. However, increased autocrine or intracrine growth factor signaling might also bypass the need for ER-mediated growth stimulation in tumor cells or affect stromal components of the tumor, such as endothelial or immune cells (33–36), to alter the tumor environment in ways conducive to tumor growth. In either case, alternative hormonal therapies might not be effective.

Recently, cell-specific coactivators and corepressors have been identified for steroid hormone receptors, including the ER, which may influence steroid receptor-induced transcription positively or negatively (37, 38). Thus, the activity of tamoxifen in inhibiting or even stimulating tumor growth might depend on the relative expression of various stimulatory or inhibitory cofactors in a particular tumor (39, 40). However, transient transfection experiments suggest that tamoxifen-resistant tumors produced by such mechanisms should still be sensitive to pure antiestrogens (40).

FGFs and their receptors have been shown to be present with high frequency in breast cancer specimens (41–50). Evidence for a possible role for FGF signaling in the estrogen-independent growth of breast tumors is gained from study of clonal and polyclonal FGF-transfected MCF-7 cell lines, which are capable of forming large, progressively growing tumors in ovariectomized or tamoxifen-treated nude mice. Moreover, the FGF-transfected cells are metastatic, forming micrometastases in lymph nodes, lungs, and other organs (21, 22, 51). The estrogen-independent and tamoxifen-resistant growth of FGF-transfected MCF-7 cells suggests an interaction between FGF signaling pathways and ER-activated pathways that could occur at the level of the ER itself or at the end point of both pathways, where they impinge on growth mechanisms. If FGF-mediated growth pathways bypass the ER pathway to affect growth di-

rectly, we would expect that growth would be unaffected by hormonal treatments devoid of agonist activity. We therefore sought to determine the sensitivity of the estrogen-independent tumor growth of FGF-transfected MCF-7 cells to ICI 182,780 or aromatase inhibitors. In contrast to what was seen with ERB-B signaling pathways, we report that FGF-mediated pathways appear to provide an alternative growth stimulatory signal that is not dependent on ER activation.

MATERIALS AND METHODS

Cell Lines. FGF-transfected MCF-7 cell lines have been described previously (21, 22, 51, 52). Briefly, the ML-20 clonal cell line is a MCF-7-derived cell line that is stably transfected with a *lacZ* expression vector. The *in vitro* and *in vivo* growth characteristics of ML-20 cells are indistinguishable from wild-type MCF-7 cells (51), and >90% of the cells routinely stain positive for β -galactosidase expression by X-gal staining (52). MKL-F (FGF-4-transfected; Ref. 52) and FGF-1 clone 18 (FGF-1-transfected) cells (22) resulted from the stable transfection of the ML-20 clonal cell line with expression vectors for FGF-4 (also known as hst-1/K-FGF) and FGF-1 (also known as acidic FGF or aFGF), respectively. Both cell lines continue to stably express β -galactosidase, allowing effects of FGF overexpression on metastatic capability to be assessed by X-gal staining of organs and tissues of tumor-bearing mice. The MKL-4 cell line was derived by transfecting wild-type MCF-7 cells (of similar passage number used for the ML-20 transfection) with an expression vector for FGF-4, which produced the clonal MKS-1 cells (21). These cells were then retransfected with an expression vector for *lacZ*, yielding MKL-4 cells (51). Cells were maintained in IMEM (Biofluids, Rockville, MD) supplemented with 5% FBS in a humidified, 37°C, 5% CO₂ incubator in routine culture until used for tumor cell injection.

Drugs. ICI 182,780 was kindly donated by Dr. Alan Wakeling of Zeneca Pharmaceuticals (Macclesfield, England), and was administered s.c. at a dose of 5 mg in 0.1 ml of vehicle every week. For the experiment depicted in Fig. 1, powdered drug was first dissolved in 100% ethanol and spiked into warmed peanut oil (Eastman Kodak, Rochester, NY) to give a final concentration of 50 mg/ml. For the experiments depicted in Fig. 1, B and C, 50 mg/ml preformulated drug in a vehicle of 10% ethanol, 15% benzyl benzoate, 10% benzyl alcohol, brought to volume with castor oil, was supplied by B. M. Vose (Zeneca Pharmaceuticals). 4-OHA was donated by Angela Brodie (University of Maryland, Baltimore, MD) and was administered s.c. at a dose of 1 mg/mouse/day 6 days of the week in a vehicle of 0.3% hydroxypropylcellulose. Letrozole was donated by Dr. Ajay Bhatnagar (Novartis, Ltd., Basel, Switzerland) and was administered via gavage at a dose of 1 mg/mouse/day 6 days of the week in a vehicle of 0.3% hydroxypropylcellulose. Sustained-release (60 day) pellets containing 5 mg of tamoxifen were obtained from Innovative Research of America (Sarasota, FL) and implanted s.c. in the interscapular area at the time of tumor cell injection.

Tumor Cell Injection. The procedure for tumor cell injection has been described previously (21). Briefly, tumor cells were scraped into their normal growth medium, and viable cells were quantified using trypan blue exclusion. The cells were

resuspended in their normal growth medium at a density of 66.7×10^6 cells/ml, and 0.15 ml (containing 10 million cells) were used to inject ovariectomized mice (nude or *beige/nude/xid*) into the mammary fat pad. For the experiment involving MKL-4 cells and nude mice (Fig. 1A), each mouse was injected bilaterally into the thoracic mammary fat pads (two injections per mouse). There were seven mice in the vehicle group and five mice in each treatment group. For the experiments involving MKL-4 cells and *beige/nude/xid* mice (Fig. 2), four tumor cell injections were given, two on each side in the thoracic mammary fat pad and two in the inguinal mammary fat pad; treatment groups consisted of four mice. For the experiments involving MKL-F and FGF-1, clone 18 cells (Fig. 1, B and C), each mouse was injected once in the right thoracic mammary fat pad. There were seven mice in the each vehicle group, and treatment groups consisted of five or six mice each. Tumors resulting from the injections were measured twice weekly in three dimensions using calipers. Tumor volume is the product of the largest dimension, the orthogonal measurement, and the tumor depth, as described previously (21). Because the FGF-1-transfected clone 18 cell line produces tumors that in some cases are surrounded by a fluid-filled sac that confounds tumor measurements (22), these tumors were measured postmortem by weighing them.

Determination of Metastasis. Organs were harvested from tumor-bearing animals, fixed briefly, and stained with X-gal as reported previously (51) and viewed through a dissecting microscope (Olympus SZH). Clusters of blue-staining cells were identified as micrometastases. In accordance with previous results, no macrometastases were identified (21, 22, 51, 53).

Growth Assays. Anchorage-dependent and anchorage-independent growth assays were performed as described (21). Briefly, for anchorage-dependent growth, cells were plated in 24-well culture dishes at a density of 10,000 cells/well for the time course experiments (Fig. 4) and 20,000 or 30,000 cells/well for the concentration-response experiments (Fig. 5). For growth in FBS, following overnight attachment, treatments were added at the indicated concentrations, and cells were counted on the indicated days. For growth assays under estrogen-depleted conditions, cells were stripped of estrogens during a 24-h period the day following plating by changing the medium four times to phenol red-free IMEM supplemented with 5% CCS (21). We have found that this stripping procedure allows complete removal of estrogens without substantial proliferation of cells before treatments are added. Following the stripping procedure, on day 0, treatments were added, and counting of cells was done as above.

Doubling times were determined according to the following equation: doubling time = $t_2 - t_1 / 3.32 \log(N_2/N_1)$, where N_2 and N_1 are the number of cells at times t_2 and t_1 , respectively. N_1 and N_2 are the means of quadruplicate determinations.

Anchorage-independent assays in FBS-containing medium were done as described previously (21). For experiments using estrogen-depleted conditions, cells were stripped of estrogens over a 24-h period as described above before being plated in soft agar. Colonies greater than 60 μm were counted using an Omnicon 3600 Image Analysis system.

ER Assays. [^3H]Estradiol binding has been described previously (54, 55). Briefly, cells grown to 70% confluence were stripped with twice daily medium changes over 4 days

with 5% CCS in phenol red-free IMEM. The prolonged stripping method allows ERs to become up-regulated to maximal levels. Cells were harvested, washed sequentially at 4°C with serum-free, phenol red-free IMEM followed by TEG (10 mM Tris, pH 7.4, 1 mM EDTA, 10% glycerol), and resuspended in 1 ml of TEG plus 1 mM DTT, 0.5 M NaCl and a cocktail of protease inhibitors (1 mg/ml leupeptin, 77 $\mu\text{g}/\text{ml}$ aprotinin, 1 $\mu\text{g}/\text{ml}$ pepstatin A). A whole-cell extract was prepared by homogenization with 40 strokes in a Teflon-glass Dounce homogenizer followed by centrifugation at $105,000 \times g$ for 30 min. Protein content of the supernatant was determined by the method of Bradford (56), and protein concentrations were adjusted to 2 mg/ml. Extracts were incubated with 10 nM [^3H]17 β -estradiol with or without a 100-fold excess of unlabeled estradiol for 16 h at 4°C. Unbound ligand was removed by absorption with dextran-coated charcoal followed by centrifugation. Aliquots of the supernatant were counted in a Beckman liquid scintillation counter.

Northern Blots. Cells were grown to 50% confluence in IMEM supplemented with 5% FBS and then stripped of estrogens as described for the growth assays, above. Treatments of 0.1% ethanol (vehicle) or 10^{-8} M 17 β -estradiol in the same medium were added. Cultures were harvested after 3 days of treatment, and RNA was extracted using RNeasy (Qiagen, Inc.) according to the manufacturer's directions. Thirty μg of each RNA were subjected to electrophoresis in a 1.2% formaldehyde/agarose gel and transferred to nylon (Hybond-N, Amersham Corp., Arlington Heights, IL) by capillarity. ^{32}P -labeled antisense riboprobes for pS-2, GAPDH, and cathepsin D were prepared and sequentially hybridized to the membrane overnight at 65°C [hybridization buffer was 50% formamide, 50 mM Na_2HPO_4 , 0.8 M NaCl, 10 mM EDTA, 2.5 \times Denhardt's solution (1 \times Denhardt's = 0.02% polyvinylpyrrolidone, 0.02% BSA), 0.2% SDS, 400 $\mu\text{g}/\text{ml}$ yeast tRNA, and 400 $\mu\text{g}/\text{ml}$ sonicated salmon sperm DNA with 10^6 DPM/ml of the appropriate probe]. The membrane was washed three times in 0.1% SDS/0.1 \times SSC at 80°C for the PS-2 and cathepsin D probes, and 75°C for the GAPDH probe. Autoradiograms and PhosphorImager (Molecular Dynamics Model 445SI) quantitation of individual hybridization signals were obtained between the sequential hybridizations. For the results depicted in Fig. 7, A and B, PhosphorImager values obtained for PS-2 or cathepsin were normalized to those obtained for GAPDH.

Progesterone Receptor mRNA Determination by RT-PCR. The primers for human progesterone receptor that produce a 205-bp PCR product have been described previously (57). The human GAPDH primers that produce a 437-bp PCR product are as follows: 5'-AAG GTC GGT GTG AAC GGA TTT G-3' (sense) and 5'-TGG TGC AGG ATG CAT TGC TG-3' (antisense). RT-PCR was performed with 0.1 μg of test RNAs, except T47D cells, where 0.02 μg was used, using the GeneAmp RNA PCR kit (PE Applied Biosystems, Foster City, CA) according to the manufacturer's instructions with the following modifications: the RT reaction was primed with 0.0625 μM random hexamers in a volume of 40 μl , with 2 μl each of ^{35}S -labeled UTP and ^{35}S -labeled ATP (each 3000 Ci/mmol, 10 $\mu\text{Ci}/\mu\text{l}$, Amersham Corp.) substituted for water in the reaction. Then, 20 μl of each RT reaction were transferred into two tubes for separate GAPDH and progesterone receptor PCR reactions.

Cycle analyses using RNA from ML-20, estradiol-treated cells (the highest expressors of progesterone receptor) revealed that amplification remained logarithmic at 35 cycles for the GAPDH reaction and 40 cycles for the progesterone receptor reaction, making these assays semiquantitative. The GAPDH PCR reaction was performed using standard reagent conditions recommended by the manufacturer and cycles of 95°C for 45 s and 50°C for 45 s for 35 cycles. For the progesterone receptor PCR reaction, final MgCl₂ concentrations were adjusted to 1.25 mM, and 0.25 M acetamide was included. Cycles were of 95°C for 45 s and 50°C for 45 s for 40 cycles. GAPDH and progesterone receptor reaction products were first visualized by ethidium bromide staining following electrophoresis in a 2% agarose gel. Products were then electrophoresed on a 4–20% acrylamide gel that was subjected to both autoradiography and PhosphorImager quantitation as described above.

Transient Transfection, Luciferase, and CAT Reporter Assays. ML-20 and clone 18 cells were plated in 6-well plates, allowed to attach overnight, and stripped of estrogens in a procedure similar to that for the growth assays (see above). Following stripping, cells were transfected by the calcium phosphate, low-CO₂ method (58). The luciferase plasmids pGLB-MERE or pGLB-MNON were obtained by inserting an approximately 1.48-kb fragment containing a glucocorticoid response element-deleted mouse mammary tumor virus promoter with either a substituted double consensus ERE (MERE) or the same sequence with the ERE palindromes scrambled (MNON) (59) into the *Hind*III site of pGLB (Promega, Madison, WI). Each dish received 2.5 µg of either pGLB-MERE or pGLB-MNON and 1.0 µg pCMV-CAT, which directs constitutive expression of CAT, cotransfected as a control for transfection efficiency. Following transfection, each well was washed twice with PBS and incubated for 48 h in medium containing vehicle (0.01% ethanol), 10⁻⁹ M estradiol, 10⁻⁷ M ICI 182,780, a combination of E₂ and ICI, 10 ng/ml FGF-1 plus 10 µg/ml heparin, or a combination of FGF, heparin, and ICI 182,780. (Duplicate samples of each treatment were used.) Cells were lysed and assayed for luciferase activity using the Luciferase Reporter Gene Assay (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's instructions. Luciferase values, expressed as relative light units, for each sample were corrected for background by subtracting the value of lysates of untransfected cells prepared in parallel. CAT expression was assayed using the CAT ELISA (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's instructions. Protein content of the lysates was determined using the BCA Protein Assay Reagent (Pierce, Rockford, IL). Luciferase and CAT values, normalized for protein, were used to calculate mean specific relative light units/ng CAT.

Statistical Analyses. Statistical methods used for tumor growth have been described previously (53, 60). For Figs. 1 and 2, only mice surviving at the end of the experiment were included in the analysis. When no tumor developed from a particular injection, tumor volume was recorded as zero. The repeated measures ANOVA (60) was used to compare tumor volumes among the treatment groups using measurements taken over the entire time course of the experiment. In addition, final tumor volumes (or weights in the case of clone 18) were compared among treatment groups at the end of each experiment using ANOVA. For analysis of metastasis in Table 1, for

each transfectant, analysis of covariance was used to compare the effects of treatment on total metastases, total distant metastases (lung metastases plus other metastases), lymph node metastases, lung metastases, and other metastases. The analyses were all conducted with final tumor volume (or weight for the clone 18 cells) included in the model as a covariate. The analyses considered the effects of all treatments simultaneously, as well as the effects of individual treatment comparisons (which were adjusted for multiple comparisons using Dunnett's method). For each transfectant, the effect of final tumor volume (or weight for clone 18) on the number of metastases was evaluated using linear regression (for each of the categories of metastasis described above). In Fig. 3, paired *t* tests were performed comparing control and transfected cells under different conditions of treatment. For the anchorage-dependent growth assays depicted in Fig. 4, we examined the effect of treatment on the rate of cell growth, using linear regression with an interaction between time and treatment. To compare cell growth rates and doubling times among the cell lines under specific treatment conditions, nested linear regression models were used. For Fig. 6, ANOVA was used to determine significant differences in ER binding among cell lines.

RESULTS

Estrogen-independent Growth of Tumors Produced by FGF-transfected MCF-7 Cells Is Not Inhibited by Treatment with a Pure Antiestrogen or with Aromatase Inhibitors. We have previously shown that both FGF-1- and FGF-4-transfected MCF-7 cells form progressively growing tumors in ovariectomized nude mice, as well as in similar mice treated with tamoxifen (21, 22, 53). Although ovariectomized mice could be expected to have substantially lower levels of estrogenic compounds than reproductively intact mice, some estrogens are synthesized at extraovarian sites, such as adrenal gland, liver, fat, or possibly the tumor itself. The transfected cells evidently still possess ERs, because they respond to estrogen and tamoxifen administered to the mice, as well as to these compounds used in tissue culture (21, 22). To test the hypothesis that growth of the FGF-transfected cells in ovariectomized or tamoxifen-treated nude mice is due to increased sensitivity to the small amounts of estrogens still present in ovariectomized nude mice, we tested the ability of a pure antiestrogen, ICI 182,780, and two aromatase inhibitors, 4-OHA and letrozole, to inhibit the estrogen-independent tumor growth produced by these FGF-transfected cell lines.

In a first experiment to test the above hypothesis, FGF-4-transfected MKL-4 cells were injected as before, and the mice were treated with vehicle, tamoxifen, or ICI 182,780. There were no significant differences in tumor volume among the treatment groups considered over the entire time course of the experiment (*P* = 0.72) or at the final time point (Fig. 1A; *P* = 0.72). Treatment with ICI 182,780 did not inhibit tumor growth below that achieved in vehicle-treated mice (*P* = 0.675). Thus, the failure of ICI 182,780 to inhibit the estrogen-independent growth exhibited by this cell line supports the hypothesis that such growth does not result from small amounts of estrogenic




Fig. 1 Growth of FGF-transfected MCF-7 cells in ovariectomized nude mice is not inhibited by treatment with ICI 182,780, 4-OHA, or letrozole. Ten million cells from the indicated cell lines were injected into the mammary fat pads of ovariectomized nude mice treated with vehicle (VEH); a 5-mg, 60-day-release tamoxifen pellet (TAM); ICI 182,780, 5 mg s.c. every week (ICI); 1 mg of 4-OHA s.c. per day 6 days of the week (4-OHA); or 1 mg of letrozole per day via gavage 6 days of the week (LET). Columns, group mean; bars SE. Numbers above each column are the percentages of injections resulting in measurable tumors at that time point. A, volumes of tumors produced by one clonal FGF-4-transfected MCF-7 cell line, MKL-4, at the indicated number of days following tumor cell injection. B, volumes of tumors produced by a second clonal FGF-4-transfected MCF-7 cell line, MKL-F, at the indicated number of days following tumor cell injection. C, weights of tumors produced by a clonal FGF-1-transfected MCF-7 cell line, FGF-1, clone 18, weighed after sacrifice of the animals 28 days after tumor cell injection. (Because the FGF-1 producing MCF-7 cells may form fluid-filled sacs around the tumor, confounding tumor measurements before sacrifice, only postmortem weights are presented here.)

growth stimulation achieved by extraovarian estrogen production.

We wished to assess the effect of ICI 182,780 on metastasis as well as on tumor growth. In spite of its retention of the transfected *lacZ* expression plasmid, the MKL-4 cell line becomes heterogeneous over time with respect to β -galactosidase expression, such that a few cells have high expression, but most are negative (52). We therefore used a second clonal FGF-4-transfected MCF-7 cell line, MKL-F, the β -galactosidase expression of which is stable, for a subsequent experiment involving FGF-4-transfected MCF-7 cells. Because FGF-1 has also been shown to produce estrogen-independent *in vivo* growth when transfected into MCF-7 cells (22), we also included a clone of FGF-1-transfected cells designated clone 18, the β -galactosidase expression of which is also stable. For these experiments, two aromatase inhibitors, 4-OHA (61, 62) and letrozole (63), were also used to inhibit extraovarian synthesis of estrogens.

In agreement with the experiment using MKL-4 cells depicted in Fig. 1A, when the FGF-4-transfected MKL-F cells were used, there were no differences in tumor volume among treatment groups over all time points ($P = 0.382$), and ICI 182,780 did not decrease tumor growth below that obtained in vehicle-treated animals (Fig. 1B; $P = 0.837$ for the last time point). In addition, neither 4-OHA nor letrozole decreased tumor growth below vehicle-treated levels ($P = 0.571$ and 0.931 for the last time point, respectively).

FGF-1-transfected clone 18 cells form tumors that are sometimes surrounded by a fluid-filled sac (22, 53), preventing

accurate tumor volume measurements during the course of the experiment. Consequently, when these cells were used (Fig. 1C), only terminal tumor weights were analyzed with ANOVA. As with the MKL-4 and MKL-F cells, ICI 182,780 did not inhibit estrogen-independent tumor growth in the clone 18 cells ($P = 0.977$). Administration of ICI 182,780 to animals injected with ML-20 cells, a clonal line of β -galactosidase-transfected wild-type MCF-7 cells (51), also produced no effect when compared with vehicle-treated animals [*i.e.*, no progressively growing tumors were obtained in either case (data not shown)]. In other, separate experiments, a polyclonal population of control vector-transfected ML-20 cells that forms progressively growing tumors in estrogen-supplemented mice (22) did not form tumors in either untreated or ICI 182,780-treated animals.⁴ Thus, the continued progressive *in vivo* growth of FGF-transfected cells in ovariectomized animals treated with either a pure antiestrogen or aromatase inhibitors demonstrates that the estrogen-independent growth of these cells in untreated ovariectomized nude mice is not due to estrogenic activity produced at extraovarian sites.

Because ICI 182,780, 4-OHA, and letrozole were without effect in the experiments described above, we injected reproductively intact female mice for 2 weeks with these compounds at the same doses used in the above experiments to observe for activity in preventing effects of endogenous estrogens on the

⁴ Unpublished results.

Table 1 Metastasis of FGF-transfected MCF-7 cells is not inhibited by treatment with ICI 182,780 or aromatase inhibitors

Mice were sacrificed and tumors and organs were subjected to X-gal staining as described previously (51). Mice bearing tumors produced by injection of MKL-4 cells were sacrificed at 61 days; for MKL-F tumors, mice were sacrificed after 64 days; and for FGF-1 clone 18 tumors, mice were sacrificed after 28 days.

Injected cells/ treatment	No. of tumor- bearing mice	Metastatic site		
		Positive lymph nodes/ lymph nodes examined	Lung	Other
MKL-4				
Vehicle	3	3/10	3	7
TAM ^a	4	5/18	2	2
ICI 182,780	5	4/23	3	4
MKL-F				
Vehicle	6	0/27	3	1
TAM	5	4/20	3	0
ICI 182,780	3	0/14	1	0
4-OHA	3	0/13	0	0
LET	3	1/12	0	0
FGF-1 clone 18				
Vehicle	6	5/24	2	0
TAM	6	3/23	3	3
ICI 182,780	4	2/13	3	1
4-OHA	5	5/18	2	1
LET	5	4/22	3	0

^aTAM, tamoxifen; LET, letrozole.

endometrium. Uteri harvested from mice injected with either ICI 182,780, 4-OHA, and letrozole weighed less than those from control mice and exhibited a complete lack of endometrial glandular structures (data not shown). Thus, these compounds retained activity, although they had no effect on tumor growth in our experiments.

Metastatic Frequency of Tumors Produced by FGF-transfected MCF-7 Cells in Mice Treated with ICI 182,780 or Aromatase Inhibitors Is Not Affected by Treatment. Because the FGF-4-transfected MKL-F cells and the FGF-1-transfected clone 18 cells stably express bacterial β -galactosidase by virtue of *lacZ* transfection, we were able to detect distant micrometastases from tumors produced by these cells. Although the MKL-4 cells become heterogeneous over time with respect to β -galactosidase expression, some high-expressing cells do remain (52), so animals from the experiment depicted in Fig. 1A were also analyzed. Table 1 shows the frequency of metastasis detected for each organ examined. However, there were no significant effects of treatment on reduction of total metastases, distant metastases, lymph node, lung, or other metastases for tumors produced by any of the cell lines. Because we have previously shown that the degree of metastasis in this tumor system is correlated with tumor size in tumors produced by both the MKL-4 and MKL-F cells (51, 53), we evaluated the correlation of individual tumor size with frequency of metastasis in individual mice for the clone 18 cells and found that tumor size

and incidence of metastasis were indeed significantly correlated ($P = 0.014$).

Effects of FGF and/or Estrogen on the Residual Immune System of Nude Mice Is Not Responsible for the Estrogen-independent Growth of FGF-transfected MCF-7 Cells. Although nude mice have a T-cell defect, they retain NK cell activity. It has been postulated that the residual NK activity in nude mice is responsible for some xenograft rejection and poor metastatic ability of xenografts (35). Estrogen and tamoxifen have been shown to decrease NK cell activity in nude mice (36), but estrogen increases the ability of NK cells to kill MCF-7 cells in cytotoxicity assays (64). In addition, transforming growth factor β , which might be secreted by MCF-7 cells in response to tamoxifen treatment (65), can decrease NK cell activity (33, 34). FGF-2 (also known as basic FGF or bFGF) has been shown to negatively affect NK activity by decreasing endothelial cell adhesion molecule expression (66), raising the possibility that FGF-1 and/or FGF-4 might have the same effect. In addition, B-cell maturation of nude mice is defective because it lacks appropriate help from T-cells (35). Because of the complexity of possible interactions of estrogen, FGFs, and MCF-7 cells with the immune system, and because of the possibility that the estrogen-independent or tamoxifen-stimulated *in vivo* phenotype of the MKL-4 cells is due to an effect of the transfected FGF and/or estrogen on the remaining immunocompetence in the nude mouse, we injected the MKL-4 clonal line of FGF-4-transfected MCF-7 cells into triply deficient *beige/nude/xid* mice. These mice exhibit intermediate NK activity coupled with defects in maturation of both B and T lymphocytes (35). Tumor growth in this host was somewhat slower than in the athymic nude mice because tumors measured at 74 days (Fig. 2A) were smaller than tumors using the same cells in nude mice measured at 61 days (Fig. 1A). However, estrogen-independent and tamoxifen-resistant growth was again seen in these animals. Pulmonary and lymphatic micrometastases were present in two of two tumor-bearing mice examined (data not shown). Injection of the clonal MCF-7 cell line ML-20 into this host produced much smaller tumors in estrogen-treated animals, as depicted in Fig. 2B. Tumor nodules produced by ML-20 cells in animals treated with tamoxifen were quite small and static, and ultimately they regressed, as has been previously shown in nude mice (21, 67). Although tumor growth was slower and the differences between treatment groups did not reach significance in this host for either cell line, the tumorigenic, tamoxifen-resistant, metastatic phenotype of MKL-4 cells was not altered in this host, and estrogen-independent growth of control ML-20 cells did not occur. We therefore conclude that the residual immunocompetence remaining in nude mice is not important in the estrogen-independent, tamoxifen-resistant *in vivo* growth of these transfectants.

FGF-transfected MCF-7 Clonal Cell Lines Have Diminished *In Vitro* Responses to ICI 182,780. Because ICI 182,780 did not affect the estrogen-independent growth of the FGF-transfected MCF-7 cells *in vivo* and because we have previously shown that the FGF transfectants do not respond to 4-hydroxytamoxifen in estrogen-containing medium to the same extent as the parental cells (21, 22), we determined their growth responses to ICI 182,780 *in vitro*.

In anchorage-independent growth assays using FBS-con-

Fig. 2 Tumorigenicity of FGF-4-transfected MCF-7 cells is not increased by injection into *beige/nude/xid* mice. Ten million MKL-4 cells were injected into the mammary fat pads of ovariectomized *beige/nude/xid* mice (four sites per mouse). Tumors were measured as in Fig. 1. The measurements shown are from 74 days after tumor cell injection **A**, volumes of tumors produced by FGF-4-transfected MKL-4 cells; **B**, volumes of tumors produced by ML-20 cells (a clonal line of MCF-7 cells).





Fig. 3 Effect of ICI 182,780 on anchorage-independent growth of FGF-transfected cells. **A**, for assay of anchorage-independent growth in FBS-containing soft agar, 20,000 cells from each cell line were plated in top agar in 35-mm dishes as described (21). After 8 days of growth, colonies greater than 60 μm in diameter were quantitated using an Omnicon Image Analysis system. *Columns*, mean of triplicate dishes; *bars*, SE. **B**, for assay of anchorage-independent growth in estrogen-depleted medium, cells from each cell line were subjected to a 24-h stripping procedure using 5% CCS in IMEM as described. Twenty thousand stripped cells were plated into top agar in 35-mm dishes, and after 14 days of growth, colonies were quantitated as above. *Columns*, mean of triplicate dishes; *bars*, SE.

taining medium (Fig. 3A), as previously reported (21, 22, 74), the baseline colony formation of the FGF transfectants is higher than that of the parental cells ($P < 0.03$). Moreover, when 10^{-7} M ICI 182,780 was added to this medium, the control ML-20 cells and the FGF transfectants were all growth inhibited, but the FGF transfectants still exhibited a higher rate of colony formation than the control ML-20 cells ($P < 0.008$). Whereas colony formation by control transfected cells was inhibited by 99% by ICI 182,780 treatment, the inhibition of colony formation of the FGF-transfected cells ranged from 67 to 84%. For all cell lines tested, addition of 10^{-8} M estradiol to the ICI 182,780-containing medium reversed the inhibition produced by ICI 182,780 (data not shown). Thus, in this assay, the FGF transfectants retained an increased ability for anchorage-independent growth in spite of treatment with ICI 182,780.

In agreement with what we have previously reported (21,

22, 74), in estrogen-depleted medium (Fig. 3B), FGF-transfected clonal lines again had significantly greater baseline colony formation than ML-20 cells ($P < 0.05$), with the exception of the MKL-4 line, which just missed significance ($P = 0.06$). As in FBS-containing medium, when ICI 182,780 was added to the medium, the FGF transfectants had significantly increased colony formation when compared with the control ML-20 cells ($P < 0.015$), indicating that the increased colony formation in estrogen-depleted medium is not due to increased sensitivity to residual estrogens remaining after the stripping process. Colony formation in the presence of ICI 182,780 was variably increased by the addition of 10^{-8} M estradiol. Thus, in estrogen-depleted medium, the FGF transfectants again had increased ability for anchorage-independent growth in the presence of ICI 182,780.

Although the anchorage-dependent growth rate of the FGF transfectants did not differ substantially from ML-20 cells in




Fig. 4 Effect of ICI-182,780 on anchorage-dependent growth of FGF-transfected MCF-7 cells in 10% FBS. A, ML-20 (parental) cells; B, FGF-4-transfected MKL-4 cells; C, FGF-4-transfected MKL-F cells; D, FGF-1-transfected clone 18 cells. E, doubling times calculated from the experiments depicted in A–D. Cells were plated in 24-well plates at 10,000 cells/well and allowed to attach overnight. The following day (day 0), media were changed to the indicated treatments. Treatment concentrations were as follows: vehicle, 0.1% ethanol; ICI 182,780, 10^{-7} M; estradiol, 10^{-8} M. Cells were harvested and counted at the indicated time points. Linear regression was performed on the data points for each treatment and the lines obtained are shown as indicated. This is a representative experiment of two.

FBS-containing medium (doubling time for ML-20 cells was 36.3 h, versus 24.4–30.2 h for the FGF transfectants), in medium supplemented with 10^{-7} M ICI 182,780, their growth rate was slowed to a much lesser extent than the control cells (Fig. 4). The doubling time for ML-20 cells in ICI 182,780-containing medium (83.6 h) was more than twice the doubling times for the FGF transfectants (29.9–39.7 h; Fig. 4E), and all of the FGF transfectants had significantly higher growth rates in the presence of ICI 182,780 than ML-20 cells ($P = 0.001$ for MKL-4, 0.007 for MKL-F, and 0.0001 for clone 18). The effect of ICI 182,780 was partially reversed by 10^{-8} M 17β -estradiol in all cell lines tested. Thus, in this assay, as in the anchorage-independent growth assay, the FGF transfectants grew better in the presence of ICI 182,780 than the control ML-20 cells.

Because others have shown that MCF-7 cells that have acquired estrogen independence exhibit increased sensitivity to estradiol or to the partial agonist properties of tamoxifen (68, 69), we determined the concentration-response relationships for 17β -estradiol, ICI 182,780, and 4-hydroxytamoxifen for the control ML-20 cells and the three FGF transfectants. In estrogen-depleted medium, estradiol stimulated growth with approximately the same potency (Table 2 and Fig. 5A) in all four cell

Table 2 Potencies of 17β -estradiol in stimulation of growth and ICI 182,780 in inhibition of estradiol stimulation of growth

Potencies were determined graphically from the concentration-response relationships depicted in Fig. 5.

Cell line	EC ₅₀ , 17β -Estradiol (M)	IC ₅₀ , ICI 182,780 (M)
ML-20	2×10^{-11}	2×10^{-9}
MKL-4	0.5×10^{-11}	5×10^{-9}
MKL-F	2×10^{-11}	8×10^{-9}
Clone 18	2×10^{-11}	10×10^{-9}

lines, in agreement with published results for MCF-7 cells (70). As previously shown (Ref. 22 and this report), the maximal effect of estradiol is diminished in stimulating growth of the FGF transfectants, which had a maximal response about 70% of that of the control ML-20 cells (Fig. 5A). When ICI 182,780 was added to estrogen-depleted medium supplemented with 10^{-10} M estradiol, its potency was slightly lower for the FGF-transfected cells than for the control cells, with IC₅₀ values ranging from 2.5 to 5 times less than that of the parental cells (Table 2 and Fig. 5B). In accordance with our previous results for 4-hydroxyta-




Fig. 5 Concentration-response relationships for 17 β -estradiol and ICI 182,780. **A**, 30,000 cells/well of the indicated cell lines were plated in 24-well dishes. After overnight attachment, the cells were stripped with four changes of estrogen-depleted medium over 24 h, after which the indicated concentrations of estradiol were added in fresh estrogen-depleted medium. Cells were harvested after 5 days of growth and counted on a Coulter counter. *Data points* (●, ML-20; ■, MKL-F; ▲, MKL-4; ◆, clone 18), mean of quadruplicate determinations; *bars*, SE. **B**, 20,000 cells/well were plated and stripped as for **A**. Treatments consisted of 10^{-10} M 17 β -estradiol alone or with the addition of the indicated concentrations of ICI 182,780. Cells were harvested and counted after 5 days. *Data points* (●, ML-20; ■, MKL-F; ▲, MKL-4; ◆, clone 18), mean of quadruplicate determinations; *bars*, SE.

moxifen (21, 22), the maximal growth-inhibitory effect of ICI 182,780 was less for the FGF transfectants than for the parental cells. All four cell lines exhibited similar small growth stimulation when treated with varying concentrations of 4-hydroxytamoxifen in estrogen-depleted medium (data not shown), in agreement with published reports (68). We conclude that the FGF transfectants do not exhibit substantially increased sensitivity to ER agonists but may be slightly less sensitive to ICI 182,780 when compared with the control ML-20 cells.

FGF-transfected MCF-7 Cells Have Numbers of ERs Similar to the Parental Cells. Others have shown that heregulin-induced growth factor signaling in MCF-7 cells results in down-regulated ERs (25, 26). Because FGF-transfected MCF-7 cells still respond to some extent to estrogen, tamoxifen, and ICI 182,780 *in vivo* and *in vitro* (Figs. 2–5 and Refs. 21 and 22), it seems obvious that they still have ERs. Nonetheless, we measured ER binding on the four cell lines used in these experiments to see whether there were differences between cell lines. Fig. 6 shows ER binding data observed for ML-20, MKL-4, MKL-F, and clone 18 cell lines. ANOVA used with these data revealed no significant differences among cell lines in numbers of binding sites for [3 H]estradiol ($P = 0.566$). Moreover, each cell line contains ample numbers of ERs that are functional, at least with respect to estrogen binding. Although it is difficult to make a direct comparison of these results with those obtained in other laboratories, it would seem that transfections with FGF-1 and FGF-4 produce a different result than transfection of MCF-7 cells with heregulin, which down-regulated ER number.

ERs Are Not Constitutively Activated in FGF-transfected MCF-7 Cells and Remain Capable of Inducing Transcription When Activated with Estrogen. As mentioned, others have reported that growth factor signaling by IGF, EGF, and heregulin can activate ER (25, 71–73). We therefore sought to determine whether ER was constitutively activated in FGF-




Fig. 6 ER levels by ligand binding assay. Ligand binding was performed using [3 H]17 β -estradiol on cells stripped of estrogens prior to the assay. *Columns*, means of four separate determinations; *bars*, SE.

transfected MCF-7 cells by determining whether basal levels of expression of estrogen-inducible genes, encoding pS2, cathepsin D, and progesterone receptor were elevated. We also evaluated the capability of the ER expressed in these cells to induce increased levels of these genes when activated by estrogen (Fig. 7). Although basal levels of expression of pS2 and cathepsin D (Fig. 7B) or progesterone receptor mRNA (Fig. 7D) varied between the cell lines, the FGF-transfected lines did not have consistently elevated levels of expression, which would be ex-




Fig. 7 Basal and estrogen-induced levels of transcripts for estrogen-responsive genes. RNA (0.1 μ g) extracted from cells growing in phenol red-free IMEM supplemented with 5% CCS and either 0.1% ethanol or 10^{-8} M 17β -estradiol was subjected to Northern blot analysis for pS2, cathepsin D, and GAPDH transcripts using 30 μ g of total RNA (A) or semiquantitative RT-PCR for progesterone receptor (PGR) and GAPDH transcripts using 0.1 μ g of total RNA as template for RT (C). RNA from T47D cells (0.02 μ g), which express high levels of progesterone receptor, was used as a positive control, and 0.1 μ g of RNA from MDA-MB-231 cells, which do not express progesterone receptor, was used as a negative control. Reactions that contained no RNA or no reverse transcriptase yielded no amplified bands (data not shown). Transcript/GAPDH ratios obtained by Phosphorimager analysis were analyzed for fold induction produced by 17β -estradiol (A and C) or comparison of basal expression with that of control ML-20 cells (B and D).

pected if the ER were constitutively activated by virtue of the FGF transfection. Similarly, the degree of induction for pS2, cathepsin D, and progesterone receptor (Fig. 7, A and C) attained by estrogen treatment was variable between cell lines, but the transfected cells did not exhibit consistently increased or decreased sensitivity to estrogen when compared with controls. Thus, the differences in basal expression or degree of estrogen induction for these estrogen-induced genes between the different cell lines is probably due to clonal or experimental variability, rather than being an effect of transfection.


Transient transfection of control and an FGF-overexpressing cell line with an ERE-containing luciferase reporter construct also indicated that ER was not constitutively activated by virtue of FGF transfection. We measured the ability of ERs expressed by the control ML-20 and one FGF-transfected cell line to direct transcription of the luciferase reporter gene driven by an ERE-containing promoter under basal and estrogen or FGF-1 stimulated conditions (Fig. 8). Neither the clone 18 nor the control ML-20 cells exhibited transcriptional activity greater than background, as determined by a similar reporter plasmid in which the ERE sequences were scrambled, under any experimental conditions other than estrogen treatment. In particular, transcription of the reporter was not elevated in the FGF-1-transfected clone 18 cells under estrogen-depleted conditions or decreased when treated with ICI 182,780. Moreover, treatment of the control ML-20 cell line with FGF-1 did not induce transcription of the reporter. We conclude that ERs in the FGF-transfected clone 18 line do not exhibit constitutive transcriptional activity and that activation of FGF receptors of

untransfected cells by FGF-1 does not induce ER-mediated transcriptional activity. Thus, taken together, our results indicate that the transfected FGFs are stimulating growth by a mechanism that bypasses the ER-mediated growth-stimulatory pathway.

DISCUSSION

In this report, we have shown that the estrogen-independent *in vivo* growth of FGF-transfected MCF-7 cells is not affected by ICI 182,780 or by either of two aromatase inhibitors. This treatment failure cannot be attributed to an estrogen-, tamoxifen-, or FGF-induced decrease in the immunocompetence remaining in nude mice. The persistence of estrogen-independent growth despite pharmacological strategies to abrogate all estrogenic activity supports the hypothesis that the effect of FGF transfection in promoting such growth is due to a direct effect of the transfected FGF. These findings are supported by our data showing normal numbers of ERs present in the FGF transfectants, which are able to direct expression of known estrogen-induced genes and interact with an ERE-containing promoter in a reporter plasmid but which are not constitutively activated in the FGF-transfected cell lines. The direct effect of FGF transfection on tumor growth may be to promote mitogenesis of the transfected cells by autocrine or intracrine FGF receptor activation. This viewpoint is supported by the generally increased proliferation rate and colony-forming ability of the FGF transfectants under estrogen-depleted tissue culture conditions. However, in addition to having a mitogenic effect on tumor cells, the

Fig. 8 ER is not constitutively activated in FGF-1-overexpressing or FGF-1-treated MCF-7 cells. Cells stripped of estrogens were transiently transfected with a luciferase reporter plasmid (pGLB, Promega) driven by a mouse mammary tumor virus promoter that contained two ERE sequences (pGLB-MERE) or the same plasmid with the ERE sequences scrambled (pGLB-MNON). Luciferase results were corrected for protein content of lysates, transfection efficiency by comparison with a cotransfected constitutively expressed CAT reporter plasmid (pCMV-CAT), and background luciferase activity.



transfected FGF may also stimulate tumor growth via effects on stromal components of the tumor, such as fibroblasts or endothelial cells. Investigations concerning the relative contributions of autocrine and/or paracrine effects of the transfected FGF-1 to tumor growth of the clone 18 cells have revealed that autocrine or intracrine effects are important for estrogen-independent tumor growth but do not seem to be necessary for either estrogen-stimulated or tamoxifen-resistant tumor growth (74). Moreover, the insensitivity of the estrogen-independent *in vivo* growth of the FGF transfectants to ICI 182,780 or the aromatase inhibitors implies that clinical tamoxifen resistance due to FGF receptor-mediated signaling may not respond to a second hormonal therapy.

The mechanism(s) determining whether a given clinical case of antiestrogen resistance will be responsive to a second hormonal manipulation has not been elucidated and may be multifactorial. Because only 30–40% of patients with acquired tamoxifen resistance have a positive response to a second hormonal therapy, with an additional 30% showing no immediate disease progression after switching therapies (13, 14, 19, 20), ER alterations that render the receptor unresponsive or differentially responsive to hormones or utilization of alternative pathways for growth that do not involve the ER may be responsible for 30–70% of acquired tamoxifen resistance. ER mutations and/or splice variants have been shown to be present in only a low percentage of clinical breast cancer cases (2, 11, 75, 76). Only a subset of these alterations results in receptors capable of producing tamoxifen resistance, and one of these, an exon 5 deletion, is not present more frequently in tamoxifen-resistant patients (77). Therefore, this mechanism is probably not a common mode of tamoxifen resistance.

The recent discovery of the *ERβ* gene (78) has raised the question of whether responses to antiestrogens for this gene product differ from those of the previously studied *ERα*. To date, transcription driven by *ERβ* at a consensus ERE in response to various antiestrogens does not seem to be different from that of *ERα* (79). However, AP-1-mediated transcription can be influenced by ER activation independent of ER-ERE

interactions (71, 80). Both ICI 182,780 and tamoxifen were shown to activate transfected *ERβ*-induced transcription at AP-1 sites in a transient transfection assay using MCF-7 cells (79). However, native MCF-7 cells have not been shown to express substantial levels of *ERβ* (81). Therefore, if our ML-20 cells are representative of native MCF-7 cells described in the literature, we would not expect that the effects of FGF transfection on *in vivo* and *in vitro* growth are due to *ERβ*-mediated stimulation of transcription at AP-1 sites.

Ligand-independent activation of the ER by growth factors has been shown to occur when activated mitogen-activated protein kinase phosphorylates a serine residue within the AFI domain of the ER (30–32). This phosphorylation also appears to increase the agonistic effects of tamoxifen but does not alter the antagonistic properties of pure antiestrogens, such as ICI 182,780 (30). Thus, growth factor pathways that result in activation of the ER might be expected to produce a situation in which tumor growth becomes supersensitive to low concentrations of hormonal agonists. Under such circumstances, the partial agonist activity of tamoxifen in promoting growth might be enhanced, whereas a pure antiestrogen would remain growth inhibitory. In one such example, overexpression of *ERB-B2* in MCF-7 cells has been shown to result in estrogen-independent and tamoxifen-insensitive growth *in vitro* and *in vivo* (23, 30). Moreover, clinical studies show decreased benefit of tamoxifen treatment in node-negative *ERB-B2*-overexpressing breast tumors (27). In support of the possibility that activation of this particular signaling pathway results in down-regulation of ERs, similar to the effect of agonist activation, it has been found that activation of *ERB-B2* signaling pathways in MCF-7 cells by the ligand, heregulin, activates the ER by phosphorylation (25) and down-regulates ER number (25, 26). This implies that an interaction between the activated *ERB-B2* signal transduction pathway and the ER is responsible for the estrogen independence and decreased tamoxifen sensitivity of the *ERB-B2* transfectants. This interpretation is further supported by the observation that the effects of added heregulin on ER activation in parental MCF-7 cells can be blocked with ICI 182,780, which also

blocks activation resulting from ERB-B2 overexpression (25). The results using these transfected cell systems, therefore, support the view that interactions between these particular growth factor pathways and the ER can produce tamoxifen resistance but may still be at least partially sensitive to ICI 182,780. Our data in this report suggest that this is not the case with FGF signaling, further suggesting that there are alternative growth-stimulating pathways that bypass the ER.

In vitro growth assays with the FGF transfectants demonstrate an increased estrogen-independent growth and reduced effectiveness of a pure antiestrogen, under both anchorage-dependent and anchorage-independent conditions, and suggest that increased growth is not due to increased potency of residual low levels of estrogen. Because separate experiments using pooled FGF-1 transfectants, as compared with pooled control transfectants, also demonstrate reduced sensitivity to ICI 182,780 (data not shown) similar to that seen with the clonal cell lines used in this study, this effect is unlikely to be due to clonal variation. Moreover, when autocrine FGF-1 signaling in the FGF-1-transfected clone 18 cells is abrogated by subsequent transfection with a dominant negative FGF receptor, sensitivity to ICI 182,780 is restored (74), implying that the reduced sensitivity seen in these experiments is due to FGF receptor activation by the transfected FGF.

Despite the activation of endogenous FGF receptors (82) by the transfected ligand, we did not observe a down-regulation of ERs in these cells, as was reported for the ERB-B2 transfectants, above. Although our data showing a slightly decreased potency of ICI 182,780 in inhibiting estradiol-stimulated growth could be interpreted as showing a slight effect of FGF receptor pathway activation on the affinity of the ER for ICI 182,780, the similar potency of 17β -estradiol in all cell lines argues against sensitization of the ER to small amounts of estradiol being responsible for the estrogen-independent growth of these cells and suggests that FGF overexpression does not alter the affinity of ER for 17β -estradiol. In addition, ICI 182,780 did not reduce anchorage-independent growth to levels of the parental cells, as one would expect if such growth were due to ligand-independent activation of the ER by the transfected FGF (Fig. 3). Moreover, we do not observe enhanced levels of mRNA estrogen-responsive genes, such as *pS2*, *cathepsin D*, or *progesterone receptor* under estrogen-depleted conditions in our transfectants. Finally, transcriptional assays using an ERE-containing reporter did not show high basal levels of transcriptional activity in the FGF transfectants. When taken together, these data provide evidence for a mechanism by which FGF-stimulated estrogen-independent growth bypasses the ER signal transduction pathway. Moreover, the algebraically additive effects of tamoxifen and estrogen to the estrogen-independent *in vivo* growth of some of the FGF transfectants (21, 22) and continued high frequency of colony formation in ICI 182,780-containing medium argues for an additive effect of ER signaling to that produced by the FGF. Studies to further investigate interactions between ER and FGF receptor signaling pathways in these transfectants are under way in our laboratory.

Previous laboratory attempts to mimic tamoxifen resistance have produced varied results with respect to cross-resistance to steroidal antiestrogens and evidence of interaction of growth factor receptor-activated and ER-activated growth pathways.

MCF-7 cells selected for growth in estrogen-depleted medium have acquired supersensitivity to estrogen *in vitro* and *in vivo* and remain sensitive to steroidal antiestrogens (69). When the LCC1 cell line, derived from MCF-7 cells by progressive *in vivo* and *in vitro* selection under estrogen-depleted conditions (83), was subjected to a subsequent *in vitro* selection in tamoxifen, the resulting LCC2 cell line remained sensitive to ICI 182,780 (84). However, a second cell line, designated LCC9, derived from the same LCC1 parent but selected instead with ICI 182,780, is cross-resistant to tamoxifen (85). Other cell lines selected for resistance to the steroidal pure antiestrogens ICI 164,384 or ICI 182,780 are cross-resistant to the other steroidal antiestrogen but not to tamoxifen (86). Additionally, a MCF-7-derived cell line selected for estrogen-independent growth in nude mice exhibits decreased numbers of ERs, is growth stimulated *in vivo* by tamoxifen, and exhibits increased AP-1-mediated transcriptional activation independent of ER activation but retains sensitivity to ICI 182,780 (87). However, tumors produced by MCF-7 cells selected *in vivo* for resistance to ICI 182,780 have shown only weak responses to tamoxifen (88). MCF-7 or T47D cells that inducibly overexpress cyclin D have been found to exhibit resistance to both tamoxifen and steroidal antiestrogens (89). Because cyclin D has been shown to be at the convergence of growth factor and ER pathways that stimulate growth (90), these results could be pertinent to our model system. Together, these diverse data imply heterogeneity for the mechanism of antiestrogen resistance and predict that clinical response to a second hormonal therapy in a given case of breast cancer will depend on the characteristics of that particular tumor.

In summary, our studies implicate direct action by FGFs in the estrogen-independent growth produced by transfection of either FGF-4 or FGF-1 into MCF-7 cells, and they rule out effects resulting from increased sensitivity of the transfectants to small amounts of extraovarian estrogen production. Our data also imply that effects of the transfected FGFs do not involve a direct interaction with the ER itself or ER signal transduction pathways, which ultimately stimulate growth, although the two pathways may still converge or interact at common downstream targets (90). We demonstrate that FGF activity at its receptor is capable of producing an increased proliferation rate of the transfectants under estrogen-depleted conditions *in vitro*, and this effect may be partly responsible for estrogen-independent growth *in vivo*. We and others have found FGF family members to be expressed in breast tissue and/or breast tumors (41–48). Moreover, FGF receptors are rather ubiquitously expressed, have been shown to be present in clinical breast cancer (49, 50), and can be activated by multiple FGF family members as well as heparin, cell adhesion molecules, or activating mutations (91). Thus, it is likely that FGF receptor-mediated signaling is operative in a significant proportion of ER-positive breast tumors. Therefore, the model described in this report might be pertinent to a number of clinical cases of tumor growth that is refractory to therapy with antiestrogens. In contrast to some of the models mentioned above, which may mimic tamoxifen-resistant breast tumors that would respond to a second hormonal therapy, we predict that tumors in which FGF receptor-mediated signaling drives autonomous growth would be refractory to alternative hormonal therapies, as well as to tamoxifen. Therapy of such tumors with agents directed against the autocrine or paracrine effects of

FGFs (53) might result in beneficial effects in such cases and might result in the restoration of antiestrogen sensitivity.

ACKNOWLEDGMENTS

The authors are indebted to A. Brodie, A. Bhatnagar, and A. Wakeling for providing 4-OHA, letrozole, and ICI 182,780, respectively. B. M. Vose supplied formulated ICI 182,780. A. Murray and the Lombardi Cancer Center Animal Research Resource assisted with animal experiments. P. Chambon and B. R. Westley provided the pS2 and cathepsin D riboprobes, respectively. S. Thomas and J. Alexander provided technical assistance. We thank A. Wellstein and R. Clarke for thoughtfully critiquing the manuscript. Animal protocols for this work were approved by the Georgetown University Animal Care and Use Committee.

REFERENCES

- Robertson, J. F. R. Oestrogen receptor: a stable phenotype in breast cancer. *Br. J. Cancer*, *73*: 5–12, 1996.
- Karnik, P. S., Kulkarni, S., Liu, X-P., Budd, G. T., and Bukowski, R. M. Estrogen receptor mutations in tamoxifen-resistant breast cancer. *Cancer Res.*, *54*: 349–353, 1994.
- Johnston, S. R. D., Sacconi-Jotti, G., Smith, I. E., Salter, J., Newby, J., Copen, M., Ebbs, S. R., and Dowsett, M. Changes in estrogen receptor, progesterone receptor, and pS2 expression in tamoxifen-resistant human breast cancer. *Cancer Res.*, *55*: 3331–3338, 1995.
- Johnston, S. R. D., Lu, B., Dowsett, M., Liang, X., Kaufman, M., Scott, G. K., Osborne, C. K., and Benz, C. C. Comparison of estrogen receptor DNA binding in untreated and acquired antiestrogen-resistant human breast tumors. *Cancer Res.*, *57*: 3723–3727, 1997.
- Jiang, S-Y., Langan-Fahey, S. M., Stella, A. L., McCague, R., and Jordan, V. C. Point mutation of estrogen receptor (ER) in the ligand-binding domain changes the pharmacology of antiestrogens in ER-negative breast cancer cells stably expressing complementary DNAs for ER. *Mol. Endocrinol.*, *6*: 2167–2174, 1992.
- Catherino, W. H., Wolf, D. M., and Jordan, V. C. A naturally occurring estrogen receptor mutation results in increased estrogenicity of a tamoxifen analog. *Mol. Endocrinol.*, *9*: 1053–1063, 1995.
- Wolf, D. M., and Fuqua, S. A. W. Mechanisms of action of antiestrogens. *Cancer Treat. Rev.*, *21*: 247–271, 1995.
- Mahfoudi, A., Roulet, E., Dauvois, S., Parker, M. G., and Wahli, W. Specific mutations in the estrogen receptor change the properties of antiestrogens to full agonists. *Biochemistry*, *92*: 4206–4210, 1995.
- Osborne, C. K., Coronado, E., Allred, D. C., Wiebe, V., and DeGregorio, M. Acquired tamoxifen resistance: correlation with reduced breast tumor levels of tamoxifen and isomerization of *trans*-4-hydroxytamoxifen. *J. Natl. Cancer Inst.*, *83*: 1477–1482, 1991.
- Tonetti, D. A., and Jordan, V. C. Possible mechanisms in the emergence of tamoxifen-resistant breast cancer. *Anticancer Drugs*, *6*: 498–507, 1995.
- Osborne, C. K., and Fuqua, S. A. W. Mechanisms of tamoxifen resistance. *Breast Cancer Res. Treat.*, *32*: 49–55, 1994.
- Pavlik, E. J., Nelson, K., Srinivasan, S., Powell, D. E., Kenady, D. E., DePriest, P., Gallion, H. H., and van Nagell, J. R., Jr. Resistance to tamoxifen with persisting sensitivity to estrogen: possible mediation by excessive antiestrogen binding site activity. *Cancer Res.*, *52*: 4106–4112, 1992.
- Howell, A., DeFriend, D., Robertson, J., Blamey, R., and Walton, P. Response to a specific antiestrogen (ICI 182780) in tamoxifen-resistant breast cancer. *Lancet*, *345*: 29–30, 1995.
- Howell, A., Downey, S., and Anderson, E. New endocrine therapies for breast cancer. *Eur. J. Cancer*, *32A*: 576–588, 1996.
- Miller, W. R. Aromatase inhibitors: where are we now? *Br. J. Cancer*, *73*: 415–417, 1996.
- Buzdar, A., Jonat, W., Howell, A., Jones, S. E., Blomqvist, C., Vogel, C. L., Eiermann, W., Wolter, J. M., Azab, M., Webster, A., and Plourde, P. V. Anastrozole, a potent and selective aromatase inhibitor, versus megestrol acetate in postmenopausal women with advanced breast cancer: results of overview analysis of two phase III trials. *J. Clin. Oncol.*, *14*: 2000–2011, 1996.
- Bonnefoi, H. R., Smith, I. E., Dowsett, M., Trunet, P. F., Houston, S. J., da Luz, R. J., Rubens, R. D., Coombes, R. C., and Powles, T. J. Therapeutic effects of the aromatase inhibitor fadrozole hydrochloride in advanced breast cancer. *Br. J. Cancer*, *73*: 539–542, 1996.
- Buzdar, A. U., Smith, R., Vogel, C., Bonomi, P., Keller, A. M., Favis, G., Mulagha, M., and Cooper, J. Fadrozole HCL (CGS-16949A) versus megestrol acetate treatment of postmenopausal patients with metastatic breast carcinoma. *Cancer (Phila.)*, *77*: 2503–2513, 1996.
- Howell, A., DeFriend, D. J., Robertson, J. F. R., Blamey, R. W., Anderson, L., Anderson, E., Sutcliffe, F. A., and Walton, P. Pharmacokinetics, pharmacological and anti-tumour effects of the specific antiestrogen ICI 182780 in women with advanced breast cancer. *Br. J. Cancer*, *74*: 300–308, 1996.
- Castiglione-Gertsch, M. New aromatase inhibitors: more selectivity, less toxicity, unfortunately, the same activity. *Eur. J. Cancer*, *3*: 393–395, 1996.
- McLeskey, S. W., Kurebayashi, J., Honig, S. F., Zwiebel, J., Lippman, M. E., Dickson, R. B., and Kern, F. G. Fibroblast growth factor 4 transfection of MCF-7 cells produces cell lines that are tumorigenic and metastatic in ovariectomized or tamoxifen-treated athymic nude mice. *Cancer Res.*, *53*: 2168–2177, 1993.
- Zhang, L., Kharbanda, S., Chen, D., Bullocks, J., Miller, D. L., Ding, I. Y. F., Hanfelt, J., McLeskey, S. W., and Kern, F. G. MCF-7 breast carcinoma cells overexpressing FGF-1 form vascularized metastatic tumors in ovariectomized or tamoxifen-treated nude mice. *Oncogene*, *15*: 2093–2108, 1997.
- Liu, Y., El-Ashry, D., Chen, D., Ding, I. Y. F., and Kern, F. G. MCF-7 breast cancer cells overexpressing transfected *c-erbB-2* have an *in vitro* growth advantage in estrogen-depleted conditions and reduced estrogen dependence and tamoxifen sensitivity *in vivo*. *Breast Cancer Res. Treat.*, *34*: 97–117, 1995.
- Benz, C. C., Scott, G. K., Sarup, J. C., Johnson, R. M., Tripathy, D., Coronado, E., Shepard, H. M., and Osborne, C. K. Estrogen-dependent, tamoxifen-resistant tumorigenic growth of MCF-7 cells transfected with *HER2/neu*. *Breast Cancer Res. Treat.*, *24*: 85–95, 1992.
- Pietras, R. J., Arboleda, J., Reese, D. M., Wongvipat, N., Pegram, M. D., Ramos, L., Gorman, C. M., Parker, M. G., Sliwkowski, M. X., and Slamon, D. J. HER-2 tyrosine kinase pathway targets estrogen receptor and promotes hormone-independent growth in human breast cancer cells. *Oncogene*, *10*: 2435–2446, 1995.
- Tang, C. K., Perez, C., Grunt, T., Waibel, C., Cho, C., and Lupu, R. Involvement of heregulin- β 2 in the acquisition of the hormone-independent phenotype of breast cancer cells. *Cancer Res.*, *56*: 3350–3358, 1996.
- Carlomagno, C., Perrone, F., Gallo, C., De Laurentiis, M., Lauria, R., Morabito, A., Pettinato, G., Panico, L., D'Antonio, A., Bianco, A. R., and De Placido, S. *c-erbB2* overexpression decreases the benefit of adjuvant tamoxifen in early stage breast cancer without axillary lymph node metastases. *J. Clin. Oncol.*, *14*: 2702–2708, 1996.
- Arnold, S. F., Obourn, J. D., Jaffe, H., and Notides, A. C. Phosphorylation of human estrogen receptor by mitogen-activated protein kinase and casein kinase II: consequences on DNA binding. *J. Steroid Biochem. Mol. Biol.*, *55*: 163–172, 1995.
- Katzenellenbogen, B. S., Montano, M. M., Le Goff, P., Schodin, D. J., Kraus, W. L., Bhardwaj, B., and Fujimoto, N. Antiestrogens. Mechanisms and actions in target cells. *J. Steroid Biochem. Mol. Biol.*, *53*: 387–393, 1995.
- Kato, S., Endoh, H., Masuhiro, Y., Kitamoto, T., Uchiyama, S., Sasaki, H., Masushige, S., Gotoh, Y., Nishida, E., Kawashima, H., Metzger, D., and Chambon, P. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science (Washington DC)*, *270*: 1491–1494, 1995.
- Bunone, G., Briand, P. A., Miksicek, R. J., and Picard, D. Activation of the unliganded estrogen receptor by EGF involves the MAP

- kinase pathway and direct phosphorylation. *EMBO J.*, *15*: 2174–2183, 1996.
32. Ignar-Trowbridge, D. M., Pimentel, M., Parker, M. G., McLachlan, J. A., and Korach, K. S. Peptide growth factor cross-talk with the estrogen receptor requires the A/B domain and occurs independently of protein kinase C or estradiol. *Endocrinology*, *137*: 1735–1744, 1996.
 33. Arteaga, C. L., Carty-Dugger, T., Moses, H. L., Hurd, S. D., and Pietsch, J. A. Transforming growth factor β , can induce estrogen-independent tumorigenicity of human breast cancer cells in athymic mice. *Cell Growth Differ.*, *4*: 193–201, 1993.
 34. Arteaga, C. L., Hurd, S. D., Winnier, A. R., Johnson, M. D., Fendly, B. M., and Forbes, J. T. Anti-transforming growth factor (TGF)- β antibodies inhibit breast cancer cell tumorigenicity and increase mouse spleen natural killer cell activity. *J. Clin. Invest.*, *92*: 2569–2576, 1993.
 35. Clarke, R. Human breast cancer cell line xenografts as models of breast cancer: the immunobiologies of recipient mice and the characteristics of several tumorigenic cell lines. *Breast Cancer Res. Treat.*, *39*: 69–86, 1996.
 36. Gottardis, M. M., Wagner, R. J., Borden, E. C., and Jordan, V. C. Differential ability of antiestrogens to stimulate breast cancer (MCF-7) growth *in vivo* and *in vitro*. *Cancer Res.*, *49*: 4765–4769, 1989.
 37. Katzenellenbogen, J. A., O'Malley, B. W., and Katzenellenbogen, B. S. Tripartite steroid hormone receptor pharmacology: interaction with multiple effector sites as a basis for the cell- and promoter-specific action of these hormones. *Mol. Endocrinol.*, *10*: 119–131, 1996.
 38. Horwitz, K. B., Jackson, T. A., Bain, D. L., Richer, J. K., Takimoto, G. S., and Tung, L. Nuclear receptor coactivators and corepressors. *Mol. Endocrinol.*, *10*: 1167–1177, 1996.
 39. Smith, C. L., Nawaz, Z., and O'Malley, B. W. Coactivator and corepressor regulation of the agonist/antagonist activity of the mixed antiestrogen, 4-hydroxytamoxifen. *Mol. Endocrinol.*, *11*: 657–666, 1997.
 40. Jackson, T. A., Richer, J. K., Bain, D. L., Takimoto, G. S., Tung, L., and Horwitz, K. B. The partial agonist activity of antagonist-occupied steroid receptors is controlled by a novel hinge domain-binding coactivator L7/SPA and the corepressors N-CoR or SMRT. *Mol. Endocrinol.*, *11*: 693–705, 1997.
 41. Ding, I. Y. F., McLeskey, S. W., Chang, K., Fu, Y. M., Acol, J. C., Shou, M. T., Alitalo, K., and Kern, F. G. Expression of fibroblast growth factors (FGFs) and receptors (FGFRs) in human breast carcinomas. *Proc. Am. Assoc. Cancer Res.*, *33*: 269, 1992.
 42. Ulich, T. R., Yi, E. S., Cardiff, R., Yin, S., Bikhazi, N., Biltz, R., Morris, C. F., and Pierce, G. F. Keratinocyte growth factor is a growth factor for mammary epithelium *in vivo*. *Am. J. Pathol.*, *144*: 862–868, 1994.
 43. Visscher, D. W., DeMattia, F., Ottosen, S., Sarkar, F. H., and Crissman, J. D. Biologic and clinical significance of basic fibroblast growth factor immunostaining in breast carcinoma. *Mod. Pathol.*, *8*: 665–670, 1995.
 44. Koos, R. D., Banks, P. K., Inkster, S. E., Yue, W., and Brodie, A. M. H. Detection of aromatase and keratinocyte growth factor expression in breast tumors using reverse transcription-polymerase chain reaction. *J. Steroid Biochem. Mol. Biol.*, *45*: 217–225, 1993.
 45. Yiangou, C., Gomm, J. J., Coope, R. C., Law, M., Luqmani, Y. A., Shousha, S., Coombes, R. C., and Johnston, C. L. Fibroblast growth factor 2 in breast cancer: occurrence and prognostic significance. *Br. J. Cancer*, *75*: 28–33, 1997.
 46. Gomm, J. J., Smith, J., Ryall, G. K., Baillie, R., Turnbull, L., and Coombes, R. C. Localization of basic fibroblast growth factor and transforming growth factor β 1 in the human mammary gland. *Cancer Res.*, *51*: 4685–4692, 1991.
 47. Bansai, G. S., Yiangou, C., Gomm, J. J., Luqmani, Y. A., Coombes, R. C., and Johnston, C. L. Expression of fibroblast growth factor 1 is lower in breast cancer than in normal breast. *Br. J. Cancer*, *72*: 1420–1426, 1995.
 48. Smith, J., Yelland, A., Baillie, R., and Coombes, R. C. Acidic and basic fibroblast growth factors in human breast tissue. *Eur. J. Cancer*, *30A*: 496–503, 1994.
 49. Luqmani, Y. A., Mortimer, C., Yiangou, C., Johnston, C. L., Bansai, G. S., Sinnett, D., Law, M., and Coombes, R. C. Expression of two variant forms of fibroblast growth factor receptor type 1 in human breast. *Int. J. Cancer*, *64*: 274–279, 1995.
 50. Penault-Llorca, F., Bertucci, F., Adélaïde, J., Parc, P., Coulier, F., Jacquemier, J., Birnbaum, D., and deLapeyrière, O. Expression of FGF and FGF receptor genes in human breast cancer. *Int. J. Cancer*, *61*: 170–176, 1995.
 51. Kurebayashi, J., McLeskey, S. W., Johnson, M. D., Lippman, M. E., Dickson, R. B., and Kern, F. G. Quantitative demonstration of spontaneous metastasis by MCF-7 human breast cancer cells cotransfected with fibroblast growth factor 4 and LacZ. *Cancer Res.*, *53*: 2178–2187, 1993.
 52. McLeskey, S. W., Zhang, L., Kharbanda, S., Kurebayashi, J., Lippman, M. E., Dickson, R. B., and Kern, F. G. Fibroblast growth factor overexpressing breast carcinoma cells as models of angiogenesis and metastasis. *Breast Cancer Res. Treat.*, *39*: 103–117, 1996.
 53. McLeskey, S. W., Zhang, L., Trock, B. J., Kharbanda, S., Liu, Y., Gottardis, M. M., Lippman, M. E., and Kern, F. G. Effects of AGM-1470 and pentosan polysulphate on tumorigenicity and metastasis of FGF-transfected MCF-7 cells. *Br. J. Cancer*, *73*: 1053–1062, 1996.
 54. Edwards, D. P., and McGuire, W. L. 17- β -estradiol is a biologically active estrogen in human breast cancer cells in tissue culture. *Endocrinology*, *107*: 884–891, 1980.
 55. Miller, D. L., El-Ashry, D., Cheville, A. L., Liu, Y., McLeskey, S. W., and Kern, F. G. Emergence of MCF-7 cells overexpressing a transfected epidermal growth factor receptor (EGFR) under estrogen-depleted conditions: evidence for a role of EGFR in breast cancer growth and progression. *Cell Growth Differ.*, *5*: 1263–1274, 1994.
 56. Bradford, M. A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, *72*: 248–254, 1976.
 57. Chevillard, S., Müller, A., Levalois, C., Lainé-Bidron, C., Vielh, P., and Magdelénat, H. Reverse transcription-polymerase chain reaction (RT-PCR) assays of estrogen and progesterone receptors in breast cancer. *Breast Cancer Res. Treat.*, *41*: 81–89, 1996.
 58. Chen, C., and Okayama, H. High-efficiency transformation of mammalian cells by plasmid DNA. *Mol. Cell. Biol.*, *7*: 2745–2752, 1987.
 59. El-Ashry, D., Chrysolgelos, S. A., Lippman, M. E., and Kern, F. G. Estrogen induction of TGF- α is mediated by an estrogen response element composed of two imperfect palindromes. *J. Steroid Biochem. Mol. Biol.*, *59*: 261–269, 1996.
 60. Heitjan, D. F., Manni, A., and Santen, R. J. Statistical analysis of *in vivo* tumor growth experiments. *Cancer Res.*, *53*: 6042–6050, 1993.
 61. Brodie, A. M. H., and Santen, R. J. Aromatase and its inhibitors in breast cancer treatment: overview and perspective. *Breast Cancer Res. Treat.*, *30*: 1–6, 1994.
 62. Dowsett, M., and Coombes, R. C. Second generation aromatase inhibitor: 4-hydroxyandrostenedione. *Breast Cancer Res. Treat.*, *30*: 81–87, 1994.
 63. Demers, L. M., Lipton, A., Harvey, H. A., Kambic, K. B., Grossberg, H., Brady, C., and Santen, R. J. The efficacy of CGS 20267 in suppressing estrogen biosynthesis in patients with advanced stage breast cancer. *J. Steroid Biochem. Mol. Biol.*, *44*: 687–691, 1993.
 64. Screpanti, I., Felli, M. P., Toniato, E., Meco, D., Martinotti, S., Frati, L., Santoni, A., and Gulino, A. Enhancement of natural-killer-cell susceptibility of human breast-cancer cells by estradiol and v-Ha-ras oncogene. *Int. J. Cancer*, *47*: 445–449, 1991.
 65. Knabbe, C., Lippman, M. E., Wakefield, L. M., Flanders, K. C., Kasid, A., Derynck, R., and Dickson, R. B. Evidence that transforming growth factor β is a hormonally regulated negative growth factor in human breast cancer cells. *Cell*, *48*: 417–428, 1987.
 66. Melder, R. J., Koenig, G. C., Witwer, B. P., Safabakhsh, N., Munn, L. L., and Jain, R. K. During angiogenesis, vascular endothelial growth

- factor and basic fibroblast growth factor regulate natural killer cell adhesion to tumor endothelium. *Nat. Med.*, 2: 992-997, 1996.
67. Osborne, C. K., Hobbs, K., and Clark, G. M. Effect of estrogens and antiestrogens on growth of human breast cancer cells in athymic nude mice. *Cancer Res.*, 45: 584-590, 1985.
68. Katzenellenbogen, B. S., Kendra, K. L., Norman, M. J., and Berthois, Y. Proliferation, hormonal responsiveness, and estrogen receptor content of MCF-7 human breast cancer cells grown in the short-term and long-term absence of estrogens. *Cancer Res.*, 47: 4355-4360, 1987.
69. Masamura, S., Santner, S. J., Heitjan, D. F., and Santen, R. J. Estrogen deprivation causes estradiol hypersensitivity in human breast cancer cells. *J. Clin. Endocrinol. Metab.*, 80: 2918-2925, 1995.
70. Wakeling, A. E. Comparative studies on the effects of steroidal and nonsteroidal oestrogen antagonists on the proliferation of human breast cancer cells. *J. Steroid Biochem.*, 34: 183-188, 1989.
71. Umayahara, Y., Kawamori, R., Watada, H., Imano, E., Iwama, N., Morishima, T., Yamasaki, Y., Kajimoto, Y., and Kamada, T. Estrogen regulation of insulin-like growth factor I gene transcription involves an AP-1 enhancer. *J. Biol. Chem.*, 269: 16433-16442, 1994.
72. Ignar-Trowbridge, D. M., Teng, C. T., Ross, K. A., Parker, M. G., and Korach, K. S. Peptide growth factors elicit estrogen receptor-dependent transcriptional activation of an estrogen-response element. *Mol. Endocrinol.*, 7: 992-998, 1993.
73. El-Tanani, M. K. K., and Green, C. D. Two separate mechanisms for ligand-independent activation of the estrogen receptor. *Mol. Endocrinol.*, 11: 928-937, 1997.
74. Zhang, L., Kharbanda, S., Hanfelt, J., and Kern, F. G. Both autocrine and paracrine effects of transfected acidic fibroblast growth factor are involved in the estrogen-independent and antiestrogen-resistant growth of MCF-7 breast cancer cells. *Cancer Res.*, 58: 352-361, 1998.
75. Leygue, E., Huang, A., Murphy, L. C., and Watson, P. H. Prevalence of estrogen receptor variant messenger RNAs in human breast cancer. *Cancer Res.*, 56: 4324-4327, 1996.
76. Roodi, N., Bailey, L. R., Kao, W.-Y., Verrier, C. S., Yee, C. J., Dupont, W. D., and Parl, F. F. Estrogen receptor gene analysis in estrogen receptor-positive and receptor-negative primary breast cancer. *J. Natl. Cancer Inst.*, 87: 446-451, 1995.
77. Daffada, A. A. I., Johnston, S. R. D., Smith, I. E., Detre, S., King, N., and Dowsett, M. Exon 5 deletion variant estrogen receptor messenger RNA expression in relation to tamoxifen resistance and progesterone receptor/pS2 status in human breast cancer. *Cancer Res.*, 55: 288-293, 1995.
78. Kuiper, G. G. J. M., Enmark, E., Peltö-Huikko, M., Nilsson, S., and Gustafsson, J. A. Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc. Natl. Acad. Sci. USA*, 93: 5925-5930, 1997.
79. Paech, K., Webb, P., Kuiper, G. G. J. M., Nilsson, S., Gustafsson, J. A., Kushner, P. J., and Scanlon, T. S. Differential ligand activation of estrogen receptors ER α and ER β at AP1 sites. *Science (Washington DC)*, 277: 1508-1510, 1997.
80. Gaub, M.-P., Bellard, M., Scheuer, I., Chambon, P., and Sassone-Corsi, P. Activation of the ovalbumin gene by the estrogen receptor involves the fos-jun complex. *Cell*, 63: 1267-1276, 1990.
81. Kuiper, G. G. J. M., Carlsson, B., Grandien, K., Enmark, E., Häggblad, J., Nilsson, S., and Gustafsson, J. A. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors α and β . *Endocrinology*, 138: 863-870, 1997.
82. McLeskey, S. W., Ding, I. Y. F., Lippman, M. E., and Kern, F. G. MDA-MB-134 breast carcinoma cells overexpress FGF receptors and are growth-inhibited by FGF ligands. *Cancer Res.*, 54: 523-529, 1994.
83. Brünner, N., Boulay, V., Fojo, A., Freter, C. E., Lippman, M. E., and Clarke, R. Acquisition of hormone-independent growth in MCF-7 cells is accompanied by increased expression of estrogen-regulated genes, but without detectable DNA amplifications. *Cancer Res.*, 53: 283-290, 1993.
84. Brünner, N., Frandsen, T. L., Holst-Hansen, C., Bei, M., Thompson, E. W., Wakeling, A. E., Lippman, M. E., and Clarke, R. MCF7/LCC2: a 4-hydroxytamoxifen resistant human breast cancer variant that retains sensitivity to the steroidal antiestrogen ICI 182,780. *Cancer Res.*, 53: 3229-3232, 1993.
85. Brünner, N., Boysen, B., Jirus, S., Skaar, T. C., Holst-Hansen, C., Lippman, J., Frandsen, T., Spang-Thomsen, M., Fuqua, S. A. W., and Clarke, R. MCF7/LCC9: an antiestrogen-resistant MCF-7 variant in which acquired resistance to the steroidal antiestrogen ICI 182,780 confers an early cross-resistance to the nonsteroidal antiestrogen tamoxifen. *Cancer Res.*, 57: 3486-3493, 1997.
86. Lykkesfeldt, A. E., Larsen, S. S., and Briand, P. Human breast cancer cell lines resistant to pure anti-estrogens are sensitive to tamoxifen treatment. *Int. J. Cancer*, 61: 529-534, 1995.
87. Dumont, J. A., Bitonti, A. J., Wallace, C. D., Baumann, R. J., Cashman, E. A., and Gross-Doersen, D. E. Progression of MCF-7 breast cancer cells to antiestrogen-resistant phenotype is accompanied by elevated levels of AP-1 DNA binding activity. *Cell Growth Differ.*, 7: 351-359, 1996.
88. Osborne, C. K., Coronado-Heinsohn, E. B., Hilsenbeck, S. G., McCue, B. L., Wakeling, A. E., McClelland, R. A., Manning, D. L., and Nicholson, R. I. Comparison of the effects of a pure steroidal antiestrogen with those of tamoxifen in a model of human breast cancer. *J. Natl. Cancer Inst.*, 87: 746-750, 1995.
89. Wilcken, N. R. C., Prall, O. W. J., Musgrove, E. A., and Sutherland, R. L. Inducible overexpression of cyclin D1 in breast cancer cells reverses the growth-inhibitory effects of antiestrogens. *Clin. Cancer Res.*, 3: 849-854, 1997.
90. Lukas, J., Bartkova, J., and Bartek, J. Convergence of mitogenic signalling cascades from diverse classes of receptors at the cyclin D-cyclin-dependent kinase-pRb-controlled G₁ checkpoint. *Mol. Cell. Biol.*, 16: 6917-6925, 1996.
91. Green, P. J., Walsh, F. S., and Doherty, P. Promiscuity of fibroblast growth factor receptors. *BioEssays*, 18: 639-646, 1996.

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11 Publication number: **0 346 014 B1**

12

EUROPEAN PATENT SPECIFICATION

- 45 Date of publication of patent specification: **03.08.94** 51 Int. Cl.⁵: **A61K 31/565, //(A61K31/565, 31:165)**
- 21 Application number: **89305563.2**
- 22 Date of filing: **02.06.89**

54 Therapeutic product for the treatment of peri- or postmenopausal conditions.

30 Priority: **06.06.88 GB 8813353**

43 Date of publication of application:
13.12.89 Bulletin 89/50

45 Publication of the grant of the patent:
03.08.94 Bulletin 94/31

54 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

56 References cited:
EP-A- 0 124 369
EP-A- 0 138 504

CHEMICAL ABSTRACTS, vol. 109, no. 3, 18th July 1988, page 73, abstract no. 17199p, Columbus, Ohio, US; N. FROELANDER et al.: "Growth hormone and somatomedin C during post-menopausal replacement therapy with estrogen alone and in combination with an antiestrogen", & MATURITAS 1988, 9(4), 297-302

73 Proprietor: **ZENECA LIMITED**
Imperial Chemical House, 9 Millbank
London SW1P 3JF(GB)

72 Inventor: **Dukes, Michael**
54 Styal Road
Wilmslow Cheshire, SK9 4AQ(GB)

74 Representative: **Tait, Brian Steele et al**
ICI Group Patents Service Dept.
PO Box 6
Shire Park
Bessemer Road
Welwyn Garden City Herts AL7 1HD (GB)

EP 0 346 014 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

This invention relates to a therapeutic product for use in a new method of medical treatment and, more particularly, it relates to a product comprising an oestrogen and a pure antioestrogen for use in a new method for the treatment or prophylaxis of perimenopausal or postmenopausal conditions, particularly perimenopausal or postmenopausal osteoporosis. The invention also relates to a pharmaceutical composition comprising an oestrogen and a pure antioestrogen and to the use thereof in the manufacture of a new medicament for use in the treatment or prophylaxis of perimenopausal or postmenopausal conditions.

When a female animal, particularly a human female, enters the perimenopausal stage the animal's ovaries begin to secrete less of the female sex hormones, particularly oestradiol. Symptoms in women at this stage include the following: vasomotor disturbances (hot flushes), urogenital atrophy (particularly affecting the vagina and distal urethra), psychosomatic complaints, changes in lipid metabolism and osteoporosis. The rate of decline of ovarian function and the severity of the above-mentioned symptoms are highly variable between individual women but in a substantial number of individuals the symptoms are sufficiently severe that treatment is required. Oestrogen replacement therapy has been used in women and it is generally recognised to be effective in combatting the typical perimenopausal and post-menopausal symptoms (British Medical Journal, 1987, 295, 914; American Journal of Obstet. and Gynecol., 1987, 156, 1298 and 1347). However oestrogen replacement therapy can also cause uterine hyperplasia, irregular vaginal menstruation and, in a small proportion of women, endometrial cancer (American Journal of Obstet. and Gynecol., 1987, 156, 1313).

To combat the continuous unopposed stimulation of oestrogen-responsive tissues an oestrogen and a progestogen are normally co-administered for part of each treatment period thereby causing regular vaginal menstruation. (American Journal of Obstet. and Gynecol., 1987, 156, 1304). However the continuation of menstrual periods is unattractive to many postmenopausal women and, in addition, progestogens can cause side effects, for example oedema, premenstrual irritability and breast tenderness.

Alternative therapies are therefore required.

It has recently been shown that compounds demonstrating a mixture of oestrogenic and antioestrogenic properties in warm-blooded animals, including humans, may be of use in the treatment of postmenopausal conditions (European Patent Specification No.0178862). Particular compounds stated to have such activity include clomiphene and tamoxifen. Comprehensive reviews of the clinical usage of these compounds are available, for example a review of clomiphene by Clark et al. in Pharmacology and Therapeutics, 1982, Volume 15, pages 467 to 519, and a review of tamoxifen by Furr et al. in Pharmacology and Therapeutics, 1984, Volume 25, pages 127-205.

It has also recently been shown that a treatment regime comprising the dosing to postmenopausal women of the oestrogen ethinyloestradiol led to an increase in serum growth hormone (GH) levels whereas the periodic dosing of both ethinyloestradiol and the antioestrogen tamoxifen led to a reduction of GH levels to pre-treatment levels [N. Froehlander et al., Maturitas, 1988, 9(4), 297 (Chem. Abstracts, 109, 17199p)].

It has also recently been shown that a treatment regime comprising the dosing of a small amount of an oestrogen, for example oestrone sulphate or natural conjugated oestrogens, followed by the dosing of an antioestrogen, for example tamoxifen or clomiphene led to the partial inhibition of the maximum oestrogen-induced stimulation of uterine endometrial tissue (A. Kauppila et al., Gynecol. obstet. Invest., 1988, 25, 58 and Arch. Gynecol., 1983, 234, 49).

It has now been found that administration of an oestrogen and a pure antioestrogen, whether simultaneously, sequentially or separately, results in the oestrogen being selectively effective in some oestrogen-responsive tissues, for example bone, and being selectively opposed in other oestrogen-responsive tissues, for example the endometrium of the uterus, and this is the basis of the present invention.

A pure antioestrogen is a compound which possesses antioestrogenic activity and no oestrogenic activity. This may be demonstrated in rats by the effect of the compound in antagonising the increase in weight of the uterus of an immature female rat produced by administering oestradiol benzoate to said rat. Thus, when each of a pure antioestrogen and oestradiol benzoate are administered for 3 days to such a rat, a smaller increase in uterine weight is produced than the substantial increase which would be produced by the administration of oestradiol benzoate alone. Unlike the known antioestrogens tamoxifen and clomiphene, when a pure antioestrogen is administered alone to a rat no increase in uterine weight whatsoever is observed.

It is disclosed in European Patent Specification No. 138504 that certain preferred steroidal antioestrogens are pure antioestrogens. It is also disclosed in European Patent No. 124369 that certain preferred non-steroidal antioestrogens are pure antioestrogens.

According to the present invention there is provided a product comprising an oestrogen and a pure antioestrogen as a combined preparation for simultaneous, sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions, the pure antioestrogen being selected from

N-n-butyl-N-methyl-, N-1H,1H-heptafluorobutyl-N-methyl- or N,N-(3-methylpentamethylene)-11-(3,17 β -
5 dihydroxyoestra-1,3,5(10)-trien-7 α -yl)undecanamide;

N-n-butyl- or N-1H,1H-heptafluorobutyl-3-p-[4-(3,17 β -dihydroxyoestra-1,3,5(10)-trien-7 α -yl)butyl]-
phenylpropionamide;

7 α -(10-p-chlorophenylthiododecyl)-, 7 α -(10-p-chlorophenylsulphinyldecyl)-, 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]-, 7 α -[10-(4,4,4-trifluorobutylsulphinyl)decyl]- or 7 α -[10-(p-chlorobenzylsulphinyl)decyl]-
10 oestra-1,3,5(10)-triene-3,17 β -diol;

7 α -(9-n-heptylsulphinyl)noyl)oestra-1,3,5(10)-triene-3,17 β -diol; and
a compound of the formula:-

NU-A-X-R¹

15 wherein NU is 6-hydroxy-2-p-hydroxyphenylnaphth-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₅-(1,4-phenylene)-(CH₂)₂-;

or NU is 1,2,3,4-tetrahydro-6-hydroxy-2-p-hydroxyphenylnaphth-1-yl (either 1RS,2RS or 1RS,2SR isomer), or 1,2,3, 4-tetrahydro-6-hydroxy-2-p-hydroxyphenyl-2-methylnaphth-1-yl (either the 1RS,2RS or
20 1RS,2SR isomer), and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;

or NU is (1RS,2RS)-5-hydroxy-2-p-hydroxyphenylindan-1-yl or (1RS,2RS)-5-hydroxy-2-p-hydroxyphenyl-2-
methylindan-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;

and wherein XR¹ is -CONR¹R² wherein R² is hydrogen or methyl and R¹ is n-butyl, 1H,1H-heptafluorobutyl,
n-pentyl or n-hexyl, or XR¹ is -SR¹, SOR¹ or -SO₂R¹ wherein R¹ is n-pentyl, n-hexyl, 4,4,5,5,5-pen-
25 tafluoropentyl or 1H,1H,2H,2H,3H,3H-heptafluorohexyl.

In a particular product of the invention the oestrogen component of a product of the invention is oestradiol, ethinyloestradiol, oestriol, oestrone, natural conjugated oestrogens, piperazine oestrone sulphate, mestranol, chlorotrianisene, dienoestrol, stilboestrol or hexoestrol or a pharmaceutically-acceptable ester thereof.

30 A pharmaceutically-acceptable ester of the oestrogen component of a product of the invention is, for example, an alkyl or aryl ester each of up to 12 carbon atoms. It will be appreciated that an ester of a steroidal oestrogen may be formed at the 3-position, the 17-position or at both of these positions. It will also be appreciated that an ester may be formed at one or both of the phenolic groups in some non-steroidal oestrogens, for example stilboestrol and hexoestrol. A suitable alkyl ester of up to 12 carbon atoms is, for
35 example, an acetate, propionate, butyrate, valerate, hexanoate, heptanoate, octanoate, cyclopentylpropionate, nonanoate, decanoate, undecanoate or dodecanoate. A suitable aryl ester of up to 12 carbon atoms is, for example, a benzoate, toluate or naphthoate. A preferred pharmaceutically-acceptable ester of the oestrogen component of a product of the invention includes, for example, oestradiol cyclopentylpropionate, oestradiol dipropionate, oestradiol heptanoate, oestradiol undecanoate, oestradiol
40 valerate and stilboestrol dipropionate.

In a further particular product of the invention the pure antioestrogen is

N-n-butyl-, N-n-butyl-N-methyl-, N-n-pentyl, N-(1H,1H-heptafluorobutyl)-or N-(1H,1H-heptafluorobutyl)-N-
methyl-3-p-[5-(6-hydroxy-2-p-hydroxyphenylnaphth-1-yl)pentyl]phenylpropionamide;

N-methyl-N-(1H,1H-heptafluorobutyl)-p-[4-[(1RS,2RS)-6-hydroxy-2-p-hydroxyphenyl-2-methyl-1,2,3,4-
45 tetrahydronaphth-1-yl]-butyl] phenylpropionamide;

(1RS,2RS)-1-[4-[p-(2-n-hexylthioethyl)phenyl]butyl]-2-p-hydroxyphenyl-1,2,3,4-tetrahydronaphth-6-ol or the
corresponding 4,4,5,5,5-pentafluoropentylthio derivative, or the corresponding hexylsulphinyl, hexylsulphonyl
or pentafluoropentylsulphinyl derivatives;

2-p-hydroxyphenyl-1-[5-[p-(2-n-hexylthioethyl)phenyl]pentyl]naphth-6-ol or the corresponding hexylsulphinyl
50 derivative; or

(1RS, 2RS)-1-[4[p-(2-n-hexylthioethyl)phenyl]butyl]-2-p-hydroxyphenyl-2-methyl-1,2,3,4-tetrahydronaphth-6-
ol or the corresponding 4,4,5,5,5-pentafluoropentylthio derivative, or the corresponding hexylsulphinyl or
pentafluoropentylsulphinyl derivative, or the corresponding (1RS,2SR) isomers of both the hexylthio and
hexylsulphinyl derivatives.

55 A preferred product of the invention comprises an oestrogen and a pure antioestrogen for use as stated above wherein the oestrogen is oestradiol or ethinyloestradiol, or a pharmaceutically-acceptable ester thereof, and the pure antioestrogen is 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol or (1RS,2RS)-2-p-hydroxyphenyl-2-methyl-1-[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]-

1,2,3,4-tetrahydronaphth-6-ol.

A particularly preferred product of the invention comprises an oestrogen and a pure antioestrogen for use as stated above wherein the oestrogen is oestradiol, oestradiol benzoate, oestradiol valerate or oestradiol undecanoate and the pure antioestrogen is 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]-
5 oestra-1,3,5(10)-triene-3,17 β -diol.

According to a further feature of the invention there is provided a process for the manufacture of a product comprising an oestrogen and a pure antioestrogen as a combined preparation for simultaneous, sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions, which process comprises bringing together said oestrogen and said pure antioestrogen.

10 In a further feature of the invention there is provided a process for the manufacture of a product comprising an oestrogen and a pure antioestrogen for simultaneous use in selective oestrogen therapy of perimenopausal or postmenopausal conditions, which process comprises bringing into admixture said oestrogen and said pure antioestrogen.

A product of the invention may be administered to a warm-blooded animal, including a human, in the
15 form of a pharmaceutical composition. Thus according to a further feature of the present invention there is provided a pharmaceutical composition which comprises the product of the invention together with a pharmaceutically-acceptable diluent or carrier.

As mentioned above a product of the invention is useful for selective oestrogen therapy of peri-
menopausal or postmenopausal conditions. It will be understood that there is no absolute requirement that
20 the oestrogen and pure antioestrogen components of the product of the invention must be dosed simultaneously. Sequential or separate use of these components may also provide selective oestrogen therapy and such use is to be understood to fall within the definition of a product of the invention. Thus it will be appreciated that a pharmaceutical composition according to the present invention includes a composition comprising an oestrogen, a pure antioestrogen and a pharmaceutically-acceptable diluent or
25 carrier. Such a composition conveniently provides the product of the invention for simultaneous use in selective oestrogen therapy of perimenopausal or postmenopausal conditions. A pharmaceutical composition according to the present invention also includes separate compositions comprising a first composition comprising an oestrogen and a pharmaceutically-acceptable diluent or carrier, and a second composition comprising a pure antioestrogen and a pharmaceutically-acceptable diluent or carrier. Such a composition
30 conveniently provides the product of the invention for sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, capsules, aqueous or oily suspensions, emulsions or dispersible powders or granules), for topical use (for
35 example as creams, ointments, gels, or aqueous or oily solutions or suspensions; for example for use within a transdermal patch), for parenteral administration (for example as a sterile aqueous or oily solution or suspension for intravenous, subcutaneous, intramuscular or intravascular dosing), or as a suppository for rectal dosing or as a pessary for vaginal dosing.

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art.

40 Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or alginic acid; binding agents such as gelatin or starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl *p*-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated
45 or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft
50 gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl-
55 methylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation

products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

5 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, castor oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as
10 ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, such as sweetening, flavouring and colouring
15 agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as castor oil, soya bean oil or arachis oil, or a mineral oil, such as, for example, liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides
20 such as lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

The pharmaceutical compositions may also be in the form of sterile injectable aqueous or oily
25 suspensions, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol, in a vegetable oil (such as arachis oil, castor oil or coconut oil) or in a mineral oil (such as liquid paraffin).

30 Conveniently the subcutaneous or intramuscular injection of an aqueous suspension or an oily solution or suspension of a pharmaceutical composition of the invention provides a depot of the active ingredients at the injection site from which those ingredients may leach out over a period of time to provide the sustained release thereof.

Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating
35 excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable,
40 vehicle or diluent using conventional procedure well known in the art.

According to a further feature of the invention there is provided a process for the manufacture of a pharmaceutical composition as defined above which comprises bringing into admixture a product as defined above together with a pharmaceutically-acceptable diluent or carrier.

45 The invention also provides the use of a product as defined above for the manufacture of a combined preparation for use simultaneously, sequentially or separately in selective oestrogen therapy of perimenopausal or postmenopausal conditions.

It will be appreciated that the definition of the product of the invention and the pharmaceutical composition of the invention includes only those products or compositions which are useful in a new method for the treatment or prophylaxis of perimenopausal or postmenopausal condition. Pharmaceutical
50 compositions comprising an oestrogen and a pure antioestrogen, together with a pharmaceutically-acceptable diluent or carrier, are novel. In European Patent Specifications Nos. 138504 and 124369 it is disclosed that the antioestrogenic activity of the compounds disclosed therein may be demonstrated by the co-administration of a test compound and oestradiol benzoate to an immature female rat. Antioestrogenic activity is demonstrated by antagonism of the increase in weight of the uterus of the rat which is produced
55 when oestradiol benzoate alone is administered to said rat. It is to be noted that, during those tests, the oestradiol benzoate was given by subcutaneous injection whereas the test compound was given separately either orally or subcutaneously.

According to a further aspect of the invention there is provided a pharmaceutical composition comprising an oestrogen and a pure antioestrogen together with a pharmaceutically-acceptable diluent or carrier.

The pharmaceutical compositions of this feature of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients well known in the art such as, for example, those disclosed above.

This aspect of the invention also provides a process for the manufacture of a pharmaceutical composition as defined immediately above which comprises bringing into admixture an oestrogen and a pure antioestrogen together with a pharmaceutically-acceptable diluent or carrier.

The invention also provides the use of a pharmaceutical composition as defined immediately above for the manufacture of a new medicament for use in selective oestrogen therapy of perimenopausal or postmenopausal conditions.

As stated above a product of the invention is of use in selective oestrogen therapy of perimenopausal or postmenopausal conditions. Selective oestrogen therapy may be demonstrated using the standard procedure set out below:-

a) an *in vivo* assay measuring the antioestrogenic activity of a compound and any oestrogenic activity possessed by that compound. This may be demonstrated in rats by the effect of the compound in antagonising the increase in weight of the uterus of an immature female rat produced by administering oestradiol benzoate to said rat. Thus, when each of a pure antioestrogen and oestradiol benzoate are administered for 3 days to such a rat, a smaller increase in uterine weight is produced than the substantial increase which would be produced by the administration of oestradiol benzoate without the pure antioestrogen. Unlike the known antioestrogens tamoxifen and clomiphene, when a pure antioestrogen is administered alone to a rat no increase in uterine weight whatsoever is observed.

The oestrogenic activity of a compound may be demonstrated in rats by the effect of the compound when it is administered alone to said rat on the uterine weight of the animal.

b) An *in vivo* assay in mature rats measuring the antioestrogenic activity of a compound by the effect of the compound when dosed during a test period of 28 days in antagonising the protective effect on the animals' bone density of their endogenous oestrogens. The bone density of a group of ovariectomised rats in which endogenous oestrogen levels are much reduced serves as a control for the effect expected to be produced by a fully effective antioestrogen.

The antioestrogenic activity of the compound in mature rats can also be measured in the same assay by measuring the effect of the compound in antagonising the effect of the animals' endogenous oestrogens which serve to increase the weight of their uteri.

A comparison of the potencies of the antioestrogenic effects of a compound as measured by its effects on the animals' bone density and uterine weights allows the selectivity of the antioestrogenic effects of the compound to be measured.

Although the pharmacological properties of a product of the invention vary with the structures of the oestrogenic and antioestrogenic components and with the route of administration, in general a product of the invention comprises:-

(i) an oestrogen which possesses oestrogenic activity in the above test (a) at doses in the range, for example, 0.002-2.0 mg/kg orally or in the range, for example, 0.0001-0.1 mg/kg subcutaneously;

(ii) a pure antioestrogen which possesses antioestrogenic activity in the above tests (a) and (b) at doses in the range, for example, in test (a): ED₅₀ 0.05-5 mg/kg orally or ED₅₀ 0.01-1.0 mg/kg subcutaneously; in test (b): antiuterotrophic effect:- ED₅₀ < 20 mg/kg/day orally, < 2 mg/kg/day subcutaneously or intramuscularly and < 10 mg/kg/injection when dosed as an intramuscular depot injection; reduction in bone density:- ED₅₀ > 20 mg/kg/day orally, > 5 mg/kg/day subcutaneously or intramuscularly and > 10 mg/kg/injection when dosed as an intramuscular depot injection.

A product of the invention is thereby seen to be surprisingly selective as the activity of the pure antioestrogen component is expressed to a high degree within uterine tissue but to a lesser degree on bone.

The size of the dose, for therapeutic or prophylactic purposes, of a product of the invention as defined above will naturally vary according to the nature and severity of the conditions presented, the age and menopausal state of the animal and the route of administration.

In general the minimum quantity of the oestrogenic component of a product of the invention as defined above will be chosen so as to provide a beneficial effect with regard to the nature and severity of the conditions presented. The quantity of the pure antioestrogenic component is then chosen to antagonise to a substantial degree the effect of the oestrogenic component on the uterine tissue. Methods of evaluating the condition of uterine tissue are well known to the man skilled in the art, for example, by examination of a

specimen of endometrial tissue taken by, for example, suction or, for example, by way of a biopsy.

So far as the oestrogenic component of a product of the invention as defined above is concerned the size of the dose and routes of administration conventionally utilised in oestrogen replacement therapy may be used. Thus, for example, a tablet containing, for example, 0.5 to 2 mg of oestradiol, oestradiol benzoate, natural conjugated oestrogens or oestradiol valerate may be administered daily. Alternatively a tablet containing 10 to 100 µg of ethinyloestradiol may be administered daily. Alternatively the oestrogenic component may be administered by, for example, intramuscular injection utilising, for example, 1 to 10 mg of oestradiol benzoate dissolved in an oil such as ethyl oleate; for example, transdermal means utilising, for example, 10-100 µg of oestradiol contained within a transdermal patch; or, for example, vaginal application utilising, for example, daily application of 0.5 to 2 mg of natural conjugated oestrogens contained within 0.5 to 5 ml of a cream.

So far as the antioestrogenic component of a product of the invention as defined above is concerned the size of the dose is chosen such that the effect of the oestrogenic component on uterine tissue is antagonised to a substantial degree whereas the beneficial effect of the oestrogenic component on bone is substantially unopposed. Thus, for example, the antioestrogenic component may be formulated in like manner to the oestrogenic component, for example as a tablet, an oily solution suitable for intramuscular injection, within a transdermal patch, or within a cream suitable for vaginal application.

The daily administration of one or more tablets containing conveniently 50 mg to 5 g, and preferably 50 mg to 500 mg, of a pure antioestrogen may be used. Preferably the pure antioestrogen may be administered by the periodic intramuscular injection of, for example, an aqueous suspension or an oily solution or suspension containing 50 mg to 5 g of the pure antioestrogen. Preferably an oily solution, for example a solution containing arachis or castor oil, an alcohol such as benzyl alcohol and 50 mg to 500 mg of the pure antioestrogen is employed. Such an injection provides a depot of the pure antioestrogen which thereafter leaches out from the injection site to provide a selective antioestrogenic effect for a period of, for example, one to six weeks.

As mentioned above a product of the invention is useful for selective oestrogen therapy of perimenopausal or postmenopausal conditions. As previously mentioned perimenopausal and postmenopausal conditions include, for example, vasomotor disturbances (hot flushes), urogenital atrophy (particularly affecting the vagina and the distal urethra), psychosomatic complaints, changes in the lipid metabolism and osteoporosis. The selective antioestrogenic effect of the pure antioestrogenic component of a product of the invention, as demonstrated by a greater antioestrogenic effect on the uterus of a rat than on the bone of the rat, allows the beneficial effect of the oestrogenic component of the product of the invention to be selectively applied to the bone and prevents the detrimental effect of an unopposed oestrogenic effect on the uterus. The utero-selective effect of the pure antioestrogenic component of a product of the invention will allow the beneficial effect of the oestrogenic component of a product of the invention to be applied to other oestrogen-responsive tissues, for example those causing vasomotor disturbances, psychosomatic complaints and changes in lipid metabolism.

The invention will now be illustrated in the following non-limiting Examples.

40 **Example 1**

Assay in Mature Rats of the Selective Antioestrogenic Activity of a Pure Antioestrogen

The pure antioestrogen used was (1RS,2RS)-2-p-hydroxyphenyl-2-methyl-1-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]-1,2,3,4-tetrahydronaphth-6-ol.

The compound was given subcutaneously as a solution in arachis oil at doses of 2 mg/kg/day and 10 mg/kg/day to two groups of 5 mature rats for a total of 28 days. Further groups of 5 mature rats served as an untreated control group. A further group of 5 mature rats was ovariectomised to serve as another control group. At the end of the treatment period the weights of the uteri of the test and control groups of rats were determined. In addition the femurs were dissected, weighed and their volumes were determined using Archimedes Principle. The femurs were then burned and the residual ash was weighed. From these data, gross femur density and bone mineral density were calculated as follows:-

Gross Femur Density = Femur Weight/Femur Volume
55 Bone Mineral Density = Femur Ash Weight/Femur Volume

The results shown below in Tables I and II demonstrate that at a dose of 2 mg/kg/day subcutaneously the test compound selectively inhibits the action of the animals' endogenous oestrogen on their uteri (90%

inhibition of uterine weight) whereas there was no significant inhibition of either bone mineral density or of gross femur density.

TABLE I

5

Treatment	Uterine Weight (mg)	Calculated Inhibition
Untreated Controls	382 ± 34	
Ovariectomised Controls	111 ± 14	
Test Compound at 2 mg/kg/day s.c.	135 ± 8	91%
Untreated Controls	369 ± 47	
Ovariectomised Controls	99 ± 5	
Test Compound at 10 mg/kg/day s.c.	125 ± 4	90%

10

15

TABLE II

Treatment	Gross Femur Density (g/ml)	Calculated Inhibition	Bone Mineral Density (g/ml)	Calculated Inhibition
Untreated Controls	1.612 ± 0.010		0.742 ± 0.009	
Ovariectomised Controls	1.569 ± 0.010		0.685 ± 0.010	
Test Compound at 2 mg/kg/day s.c.	1.604 ± 0.006	19%*	0.730 ± 0.007	21% *
Untreated Controls	1.629 ± 0.014		0.766 ± 0.005	
Ovariectomised Controls	1.571 ± 0.007		0.704 ± 0.005	
Test Compound at 10 mg/kg/day s.c.	1.580 ± 0.004	84%	0.727 ± 0.005	63%

20

25

* This level of inhibition was not statistically significant.

30

Example 2

The experiment described in Example 1 was repeated except that the pure antioestrogen used was 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol. This compound was given at a series of doses as a daily intramuscular injection, the compound having been dissolved in a mixture of propylene glycol: ethanol: water: poloxamer 407. The formulation contained 25 mg of test compound, 100 mg of ethanol (96%), 100 mg of water, 20 mg of poloxamer 407 and sufficient propylene glycol to bring the solution to a volume of 1 ml.

35

The results shown below in Tables III and IV demonstrate that at all doses tested the compound selectively inhibits the action of the animals' endogenous oestrogen on their uteri whereas there was no significant inhibition of gross femur density.

40

TABLE III

45

Treatment	Uterine Weight (mg)	Calculated Inhibition
Untreated Controls	302 ± 36	
Ovariectomised Controls	70 ± 1.3	
Test Compound (mg/kg)		
0.1	208 ± 17	41
0.3	174 ± 16	55
1	94 ± 9	90
3	103 ± 2	86

50

55

TABLE IV

Treatment	Gross Femur Density (g/ml)	Calculated Inhibition
Untreated Controls	1.523 ± 0.008	
Ovariectomised Controls	1.491 ± 0.006	
Test Compound at (mg/kg)		
0.1	1.528 ± 0.005	0%
0.3	1.528 ± 0.008	0%
1	1.532 ± 0.005	0%
3	1.533 ± 0.005	0%

15 **Example 3**

The pure antioestrogen used was 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol.

20 Each of a series of selected doses of this compound was dissolved in a mixture of castor oil and benzyl alcohol and given by intramuscular injection to a group of 5 mature rats. The formulation contained 50 mg of the test compound, 400 mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. In each case a second dose was administered two weeks after the first dose. Two weeks after the second dose the weights of the uteri of the test groups of rats were determined. In addition the femurs were

25 A further group of rats, given two injections of castor oil separated by a two week period, served as an intact control group. A further group of rats was ovariectomised to serve as another control group.

The results shown below in Tables V and VI demonstrate that at all doses tested the compound selectively inhibits the action of the animals' endogenous oestrogen on their uteri whereas at the two higher test doses there was no significant inhibition of gross femur density.

TABLE V

Treatment	Uterine Weight (mg)	Calculated Inhibition
Intact Controls	318 ± 31	
Ovariectomised Controls	76 ± 4	
Test Compound (mg/rat/dose)		
0.75	202 ± 23	48
1.25	180 ± 15	57
2.5	123 ± 12	81

TABLE VI

Treatment	Gross Femur Density (g/ml)	Calculated Inhibition
Intact Controls	1.584 ± 0.007	
Ovariectomised Controls	1.521 ± 0.005	
Test Compound (mg/rat/dose)		
0.75	1.562 ± 0.004	35
1.25	1.576 ± 0.004	13*
2.5	1.569 ± 0.007	23*

* This level of inhibition was not statistically significant.

15

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. A product comprising an oestrogen and a pure antioestrogen as a combined preparation for simultaneous, sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions, the pure antioestrogen being selected from

N-n-butyl-N-methyl-, N-1H,1H-heptafluorobutyl-N-methyl- or N,N-(3-methylpentamethylene)-11-(3,17β-dihydroxyoestra-1,3,5(10)-trien-7α-yl)undecanamide;

- N-n-butyl- or N-1H,1H-heptafluorobutyl-3-p-[4-(3,17β-dihydroxyoestra-1,3,5(10)-trien-7α-yl)butyl]-phenylpropionamide;

7α-(10-p-chlorophenylthiodecyl)-, 7α-(10-p-chlorophenylsulphonyldecyl)-, 7α-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]-, 7α-[10-(4,4,4-trifluorobutylsulphonyl)decyl]- or 7α-[10-(p-chlorobenzylsulphonyl)decyl]-oestra-1,3,5(10)-triene-3,17β-diol;

7α-(9-n-heptylsulphonylnonyl)oestra-1,3,5(10)-triene-3,17β-diol; and

- a compound of the formula:-

NU-A-X-R¹

wherein NU is 6-hydroxy-2-p-hydroxyphenylnaphth-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₅-(1,4-phenylene)-(CH₂)₂-;

or NU is 1,2,3,4-tetrahydro-6-hydroxy-2-p-hydroxyphenylnaphth-1-yl (either 1RS,2RS or 1RS,2SR isomer), or 1,2,3,4-tetrahydro-6-hydroxy-2-p-hydroxyphenyl-2-methylnaphth-1-yl (either the 1RS,2RS or 1RS,2SR isomer), and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;

or NU is (1RS,2RS)-5-hydroxy-2-p-hydroxyphenylindan-1-yl or (1RS,2RS)-5-hydroxy-2-p-hydroxyphenyl-2-methylindan-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;

and wherein XR¹ is -CONR¹R² wherein R² is hydrogen or methyl and R¹ is n-butyl, 1H,1H-heptafluorobutyl, n-pentyl or n-hexyl, or XR¹ is -SR¹, -SOR¹ or -SO₂R¹ wherein R¹ is n-pentyl, n-hexyl, 4,4,5,5,5-pentafluoropentyl or 1H,1H,2H,2H,3H,3H-heptafluorohexyl.

2. A product as claimed in claim 1 wherein the oestrogen is oestradiol, oestradiol benzoate, oestradiol valerate or oestradiol undecanoate and the pure antioestrogen is 7α-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17β-diol.

3. A process for the manufacture of a product comprising an oestrogen and a pure antioestrogen as a combined preparation for simultaneous, sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions as claimed in any one of claims 1 and 2, which process comprises bringing together said oestrogen and said pure antioestrogen.

4. A pharmaceutical composition comprising a product as claimed in any one of claims 1 and 2 together with a pharmaceutically-acceptable diluent or carrier.

5. The use of a product as claimed in any one of claims 1 and 2 for the manufacture of a combined preparation for use simultaneously, sequentially or separately in selective oestrogen therapy of

perimenopausal or postmenopausal conditions.

Claims for the following Contracting States : ES, GR

- 5 1. A process for the manufacture of a product comprising an oestrogen and a pure antioestrogen as a combined preparation for simultaneous, sequential or separate use in selective oestrogen therapy of perimenopausal or postmenopausal conditions, the pure antioestrogen being selected from
- N-n-butyl-N-methyl-, N-1H, 1H-heptafluorobutyl-N-methyl- or N,N-(3-methylpentamethylene)-11-(3,17β-dihydroxyoestra-1,3,5(10)-trien-7α-yl)undecanamide;
- 10 N-n-butyl- or N-1H,1H-heptafluorobutyl-3-p-[4-(3, 17β-dihydroxyoestra-1,3,5(10)-trien-7α-yl)butyl]-phenylpropionamide;
- 7α-(10-p-chlorophenylthiodecyl)-, 7α-(10-p-chlorophenylsulphonyldecyl)-, 7α-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]-, 7α-[10-(4,4,4-trifluorobutylsulphonyl)decyl]- or 7α-[10-(p-chlorobenzylsulphonyl)decyl]-oestra-1,3,5(10)-triene-3,17β-diol;
- 15 7α-(9-n-heptylsulphonylnonyl)oestra-1,3,5(10)-triene-3,17β-diol; and
a compound of the formula:-
- NU-A-X-R¹
- 20 wherein NU is 6-hydroxy-2-p-hydroxyphenylnaphth-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₅-(1,4-phenylene)-(CH₂)₂-;
or NU is 1,2,3,4-tetrahydro-6-hydroxy-2-p-hydroxyphenylnaphth-1-yl (either 1RS,2RS or 1RS,2SR isomer), or 1,2,3, 4-tetrahydro-6-hydroxy-2-p-hydroxyphenyl-2-methylnaphth-1-yl (either the 1RS,2RS or 1RS,2SR isomer), and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;
- 25 or NU is (1RS,2RS)-5-hydroxy-2-p-hydroxyphenylindan-1-yl or (1RS,2RS)-5-hydroxy-2-p-hydroxyphenyl-2-methylindan-1-yl and A is -(CH₂)₁₀-, -(CH₂)₁₁- or -(CH₂)₄-(1,4-phenylene)-(CH₂)₂-;
and wherein XR¹ is -CONR¹R² wherein R² is hydrogen or methyl and R¹ is n-butyl, 1H,1H-heptafluorobutyl, n-pentyl or n-hexyl, or XR¹ is -SR¹, SOR¹ or -SO₂R¹ wherein R¹ is n-pentyl, n-hexyl, 4,4,5,5,5-pentafluoropentyl or 1H,1H,2H,2H,3H,3H-heptafluorohexyl,
- 30 which process is characterised by bringing together said oestrogen and said pure antioestrogen.
2. A process as claimed in claim 1 wherein the oestrogen is oestradiol, oestradiol benzoate, oestradiol valerate or oestradiol undecanoate and the pure antioestrogen is 7α-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17β-diol.
- 35 3. A process for the manufacture of a pharmaceutical composition which comprises bringing into admixture a product as defined in any one of claims 1 and 2 together with a pharmaceutically-acceptable diluent or carrier.

40 **Patentansprüche**

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Produkt, das ein Östrogen und ein reines Antiöstrogen als Kombinationspräparat zur gleichzeitigen, aufeinanderfolgenden oder voneinander getrennten Verwendung zur selektiven Östrogen-Therapie von
- 45 Beschwerden während der oder nach den Wechseljahren enthält, wobei das reine Antiöstrogen aus folgendem ausgewählt ist:
- N-n-Butyl-N-methyl-, N-1H,1H-Heptafluorbutyl-N-methyl- oder N,N-(3-Methylpentamethylen)-11-(3,17β-dihydroxyöstra-1,3,5(10)-trien-7α-yl)undecanamid;
- 50 N-n-Butyl- oder N-1H,1H-Heptafluorbutyl-3-p-[4-(3,17β-dihydroxyöstra-1,3,5(10)-trien-7α-yl)butyl]-phenylpropionamid;
- 7α-(10p-Chlorphenylthiodecyl)-, 7α-(10p-Chlorphenylsulfinyldecyl)-, 7α-[9-(4,4,5,5,5-Pentafluoropentylsulfinyl)nonyl]-, 7α-[10-(4,4,4-Trifluorbutylsulfinyl)decyl]- oder 7α-[10-(p-Chlorbenzylsulfinyl)decyl]-östra-1,3,5 (10) -trien-3,17β-diol;
- 7α-(9-n-Heptylsulfinylnonyl)östra-1,3,5 (10)-trien-3,17β-diol; und
- 55 einer Verbindung mit der Formel: -

NU-A-X-R¹

- in der NU für 6-Hydroxy-2-p-hydroxyphenylnaphth-1-yl steht und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_5$ -(1,4-Phenylen)-(CH₂)₂- steht;
 oder in der NU für 1,2,3,4-Tetrahydro-6-hydroxy-2-p-hydroxyphenylnaphth-1-yl (entweder das 1RS,2RS- oder 1RS,2SR-Isomer) oder für 1,2,3,4-Tetrahydro-6-hydroxy-2-p-hydroxyphenyl-2-methylnaphth-1-yl (entweder das 1RS,2RS- oder 1RS,2SR-Isomer) steht, und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_4$ -(1,4-Phenylen)-(CH₂)₂- steht;
 oder in der NU für (1RS,2RS)-5-Hydroxy-2-p-hydroxyphenylindan-1-yl oder (1RS,2RS)-5-Hydroxy-2-p-hydroxyphenyl-2-methylindan-1-yl steht und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_4$ -(1,4-Phenylen)-(CH₂)₂- steht;
 und in der XR¹ für -CONR¹R² steht, wobei R² für Wasserstoff oder Methyl steht und R¹ für n-Butyl, 1H,1H-Heptafluorbutyl, n-Pentyl oder n-Hexyl steht, oder in der XR¹ für -SR¹, SOR¹ oder -SO₂R¹ steht, wobei R¹ für n-Pentyl, n-Hexyl, 4,4,5,5,5-Pentafluorpentyl oder 1H,1H,2H,2H,3H,3H-Heptafluorhexyl steht.
- 15 2. Produkt nach Anspruch 1, wobei es sich bei dem Östrogen um Östradiol, Benzoessäureöstradiolester, Valeriansäureöstradiolester oder Undecensäureöstradiolester handelt und bei dem reinen Antiöstrogen um 7 α -[9-(4,4,5,5,5-Pentafluorpentylsulfanyl)nonyl]östra-1,3,5(10)-trien-3,17 β -diol.
- 20 3. Verfahren zur Herstellung eines Produkts, das ein Östrogen und ein reines Antiöstrogen als Kombinationspräparat zur gleichzeitigen, aufeinander folgenden oder voneinander getrennten Verwendung zur selektiven Östrogen-Therapie von Beschwerden während der oder nach den Wechseljahren enthält, nämlich nach einem der Ansprüche 1 und 2, wobei bei dem Verfahren das Östrogen und das reine Antiöstrogen zusammengebracht werden.
- 25 4. Pharmazeutische Zusammensetzung, die ein Produkt nach einem der Ansprüche 1 und 2 zusammen mit einem pharmazeutisch geeigneten Verdünnungsmittel oder Träger enthält.
- 30 5. Verwendung eines Produkts nach einem der Ansprüche 1 und 2 zur Herstellung eines Kombinationspräparats zur gleichzeitigen, aufeinander folgenden oder voneinander getrennten Verwendung in der selektiven Östrogen-Therapie von Beschwerden während der oder nach den Wechseljahren.

Patentansprüche für folgende Vertragsstaaten : ES, GR

- 35 1. Verfahren zur Herstellung eines Produkts, das ein Östrogen und ein reines Antiöstrogen als Kombinationspräparat zur gleichzeitigen, aufeinanderfolgenden oder voneinander getrennten Verwendung in der selektiven Östrogen-Therapie von Beschwerden während der oder nach den Wechseljahren enthält, wobei das reine Antiöstrogen aus folgendem ausgewählt ist:
 N-n-Butyl-N-methyl-, N-1H,1H-Heptafluorbutyl-N-methyl- oder N,N-(3-Methylpentamethylen)-11-(3,17 β -dihydroxyöstra-1,3,5(10)-trien-7 α -yl)undecanamid;
 40 N-n-Butyl- oder N-1H,1H-Heptafluorbutyl-3-p-[4-(3,17 β -dihydroxyöstra-1,3,5(10)-trien-7 α -yl)butyl]-phenylpropionamid;
 7 α -(10p-Chlorphenylthiodecyl)-, 7 α -(10-p-Chlorphenylsulfanyldecyl)-, 7 α -[9-(4,4,5,5,5-Pentafluorpentylsulfanyl)nonyl]-, 7 α -[10-(4,4,4-Trifluorbutylsulfanyl)decyl]- oder 7 α -[10-(p-Chlorbenzylsulfanyl)decyl]östra-1,3,5(10)-trien-3,17 β -diol;
 45 7 α -(9-n-Heptylsulfanyl)östra-1,3,5(10)-trien-3,17 β -diol; und
 einer Verbindung mit der Formel: -

NU-A-X-R¹

- 50 in der NU für 6-Hydroxy-2-p-hydroxyphenylnaphth-1-yl steht und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_5$ -(1,4-Phenylen)-(CH₂)₂- steht;
 oder in der NU für 1,2,3,4-Tetrahydro-6-hydroxy-2-p-hydroxyphenylnaphth-1-yl (entweder das 1RS,2RS- oder 1R2,2SR-Isomer) oder für 1,2,3,4-Tetrahydro-6-hydroxy-2-p-hydroxyphenyl-2-methylnaphth-1-yl (entweder das 1RS,2RS- oder 1RS,2SR-Isomer) steht, und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_4$ -(1,4-phenylen)-(CH₂)₂- steht;
 55 oder in der NU für (1RS,2RS)-5-Hydroxy-2-p-hydroxyphenylindan-1-yl oder (1RS,2RS)-5-Hydroxy-2-p-hydroxyphenyl-2-methylindan-1-yl steht und A für $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - oder $-(CH_2)_4$ -(1,4-Phenylen)-(CH₂)₂- steht;

und in der XR^1 für $-CONR^1R^2$ steht, wobei R^2 für Wasserstoff oder Methyl steht und R^1 für n-Butyl, 1H,1H-Heptafluorbutyl, n-Pentyl oder n-Hexyl steht, oder in der XR^1 für $-SR^1$, SOR^1 oder $-SO_2R^1$ steht, wobei R^1 für n-Pentyl, n-Hexyl, 4,4,5,5,5-Pentafluoropentyl oder 1H,1H,2H,2H,3H,3H-Heptafluorhexyl steht,

5 wobei das Verfahren dadurch gekennzeichnet ist, daß das Östrogen und das reine Antiöstrogen zusammengebracht werden.

2. Verfahren nach Anspruch 1, wobei es sich bei dem Östrogen um Östradiol, Benzoessäureöstradiolester, Valeriansäureöstradiolester oder Undecansäureöstradiolester handelt und bei dem reinen Antiöstrogen
10 um 7α -[9-(4,4,5,5,5-Pentafluoropentylsulfinyl)nonyl]östra-1,3,5(10)-trien-3,17 β -diol.

3. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, bei dem ein wie in einem der Ansprüche 1 und 2 definiertes Produkt mit einem pharmazeutisch geeigneten Verdünnungsmittel oder Träger gemischt wird.

15

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Produit comprenant un oestrogène et un antioestrogène pur sous forme d'une préparation mixte pour
20 une utilisation simultanée, séquentielle ou distincte dans une thérapie sélective par oestrogènes de troubles périménopausiques ou postménopausiques, l'anti-oestrogène pur étant choisi entre

le N-n-butyl-N-méthyl-, N-1H,1H-heptafluorobutyl-N-méthyl- ou N,N-(3-méthylpentaméthylène)-11-(3,17 β -dihydroxyoestra-1,3,5(10)-triène-7 α -yl)undécaneamide ;

le N-n-butyl- ou N-1H,1H-heptafluorobutyl-3-p-[4-(3,17 β -dihydroxyoestra-1,3,5(10)-triène-7 α -yl)-butyl]phénylpropionamide ;
25

le 7α -(10-p-chlorophénylthiodécyl)-, 7α -(10-p-chlorophénylsulfinyldécyl)-, 7α -[9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl]-, 7α [10-(4,4,4-trifluorobutylsulfinyl)-décyl]- ou 7α -[10-(p-chlorobenzylsulfinyl)-décyl]-oestra-1,3,5(10)-triène-3,17 β -diol ;

le 7α -(9-n-heptylsulfinyl)nonyl]oestra-1,3,5(10)-triène-3,17 β -diol ; et

30 un composé de formule :

NU-A-X-R¹

dans laquelle NU représente un groupe 6-hydroxy-2-p-hydroxyphénylnapht-1-yl et A représente un
35 groupe $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - ou $-(CH_2)_5$ -(1,4-phénylène)-(CH₂)₂- ;

ou bien NU représente un groupe 1,2,3,4-, tétrahydro-6-hydroxy-2-p-hydroxyphénylnapht-1-yle (isomère 1RS, 2RS ou bien 1RS, 2SR), ou un groupe 1,2,3,4-tétrahydro-6-hydroxy-2-p-hydroxyphényl-2-méthylnapht-1-yle (isomère 1RS, 2RS ou bien 1RS, 2SR), et A représente un groupe $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - ou $-(CH_2)_4$ -(1,4-phénylène)-(CH₂)₂- ;

40 ou bien NU représente un groupe (1RS, 2RS)-5-hydroxy-2-p-hydroxyphénylindane-1-yle ou (1RS, 2RS)-5-hydroxy-2-p-hydroxyphényl-2-méthylindane-1-yle et A représente un groupe $-(CH_2)_{10}$ -, $-(CH_2)_{11}$ - ou $-(CH_2)_4$ -(1,4-phénylène)-(CH₂)₂- ; et dans laquelle XR^1 représente un groupe $-CONR^1R^2$ dans lequel R^2 représente l'hydrogène ou un groupe méthyle et R^1 représente un groupe n-butyle, 1H,1H-heptafluorobutyle, n-pentyle ou n-hexyle, ou bien XR^1 représente un groupe $-SR^1$, SOR^1 ou $-SO_2R^1$ dans lequel R^1
45 représente un groupe n-pentyle, n-hexyle, 4,4,5,5,5-pentafluoropentyle ou 1H,1H,2H,2H,3H,3H-heptafluorohexyle.

2. Produit suivant la revendication 1, dans lequel l'oestrogène est l'oestradiol, le benzoate d'oestradiol, le valérate d'oestradiol ou l'undécanoate d'oestradiol et l'anti-oestrogène pur est le 7α -[9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl]oestra-1,3,5(10)-triène-3,17 β -diol.
50

3. Procédé de préparation d'un produit comprenant un oestrogène et un anti-oestrogène pur sous forme d'une préparation mixte pour une utilisation de manière simultanée, séquentielle ou distincte dans une
55 thérapie sélective par oestrogènes de troubles périménopausiques ou postménopausiques, suivant l'une quelconque des revendications 1 et 2, procédé qui comprend l'association dudit oestrogène et dudit anti-oestrogène pur.

4. Composition pharmaceutique comprenant un produit suivant l'une quelconque des revendications 1 et 2, en association avec un dilant ou support pharmaceutiquement acceptable.
5. Utilisation d'un produit suivant l'une quelconque des revendications 1 et 2 pour la production d'une préparation mixte pour une utilisation de manière simultanée, séquentielle ou distincte dans une thérapie sélective par oestrogènes de troubles périménopausiques ou postménopausiques.

Revendications pour les Etats contractants suivants : ES, GR

- 10 1. Procédé de préparation d'un produit comprenant un oestrogène et un anti-oestrogène pur sous forme d'une préparation mixte pour une utilisation simultanée, séquentielle ou distincte dans une thérapie sélective par oestrogènes de troubles périménopausiques ou postménopausiques, l'anti-oestrogène pur étant choisi entre

15 le N-n-butyl-N-méthyl-, N-1H,1H-heptafluorobutylN-méthyl- ou N,N-(3-méthylpentaméthylène)-11-(3,17β-dihydroxyoestra-1,3,5(10)-triène-7α-yl)undécaneamide ;

le N-n-butyl- ou N-1H,1H-heptafluorobutyl-3-p-[4-(3,17β-dihydroxyoestra-1,3,5(10)-triène-7α-yl)-butyl]phénylpropionamide ;

20 le 7α-(10-p-chlorophénylthiodécyl)-, 7α-(10-p-chlorophénylsulfinyldécyl)-, 7α-[9-(4,4,5,5,5-pentafluoropentylsulfinylnonyl)-, 7α[10-(4,4,4-trifluorobutylsulfinyldécyl)- ou 7α-[10-(p-chlorobenzylsulfinyldécyl)]-oestra-1,3,5(10)-triène-3,17β-diol ;

le 7α-(9-n-heptylsulfinylnonyl)oestra-1,3,5(10)-triène-3,17β-diol ; et un composé de formule :

NU-A-X-R¹

25 dans laquelle NU représente un groupe 6-hydroxy-2-p-hydroxyphénylnapht-1-yl et A représente un groupe -(CH₂)₁₀-, -(CH₂)₁₁- ou -(CH₂)₅-(1,4-phénylène)-(CH₂)₂- ;

30 ou NU représente un groupe 1,2,3,4-tétrahydro-6-hydroxy-2-p-hydroxyphénylnapht-1-yle (isomère 1RS, 2RS, ou 1RS, 2SR), ou un groupe 1,2,3,4-tétrahydro-6-hydroxy-2-p-hydroxyphényl-2-méthylapht-1-yle (isomère 1RS, 2RS ou bien 1RS, 2SR), et A représente un groupe -(CH₂)₁₀-, -(CH₂)₁₁- ou -(CH₂)₄-(1,4-phénylène)-(CH₂)₂- ;

ou bien NU représente un groupe (1RS, 2RS)-5-hydroxy-2-p-hydroxyphénylindane-1-yle ou (1RS, 2RS)-5-hydroxy-2-p-hydroxyphényl-2-méthylindane-1-yle et A représente un groupe -(CH₂)₁₀-, -(CH₂)₁₁- ou -(CH₂)₄-(1,4-phénylène)-(CH₂)₂- ;

35 et dans laquelle XR¹ représente un groupe -CONR¹R² dans lequel R² représente l'hydrogène ou un groupe méthyle et R¹ représente un groupe n-butyle, 1H,1H-heptafluorobutyle, n-pentyle ou n-hexyle, ou bien XR¹ représente un groupe -SR¹, -SOR¹ ou -SO₂R¹ dans lequel R¹ représente un groupe n-pentyle, n-hexyle, 4,4,5,5,5-pentafluoropentyle ou 1H,1H,2H,2H,3H,3H-heptafluorohexyle, procédé qui est caractérisé par l'association dudit oestrogène et dudit anti-oestrogène pur.

- 40 2. Procédé suivant la revendication 1, dans lequel l'oestrogène est l'oestradiol, le benzoate d'oestradiol, le valérate d'oestradiol ou l'undécanoate d'oestradiol et l'anti-oestrogène pur est le 7α-[9-(4,4,5,5,5-pentafluoropentylsulfinylnonyl)]oestra-1,3,5(10)-triène-3,17β-diol.

- 45 3. Procédé de production d'une composition pharmaceutique, qui comprend le mélange d'un produit tel que défini dans l'une quelconque des revendications 1 et 2, avec un dilant ou support pharmaceutiquement acceptable.

50

55



Email article

ScienceDirect

Scopus

SciTopics

Applications

Register

Login

Go to SciVal Suite

Home Browse Search My settings My alerts

Help

Articles

All fields

Author

Advanced search

Images

Journal/Book title

Volume

Issue

Page

Search ScienceDirect

Search tips

Accelerate your research career.

Back to results | < Previous 27 of 37 Next >

Abstract

Related Articles

- Are breast tumours resistant to tamoxifen also resistan...
The Journal of Steroid Biochemistry and Molecular Biolo...
 - Therapeutic potential of pure antiestrogens in the tre...
The Journal of Steroid Biochemistry and Molecular Biolo...
 - Oestrogen/growth factor cross talk in breast carcinoma...
Trends in Pharmaceutical Sciences
 - Short-term effects of pure anti-oestrogen ICI 182780 o...
European Journal of Cancer
 - Use of pure antiestrogens to elucidate the mode of act...
Biochemical Pharmacology
- ▶ View more related articles

My Applications

Add

Table Download

Find HTML data tables from the current article to download.

Find Tables

About Table Download

MostDownloaded

Most downloaded articles from the last three months in this journal:

Endocrine disrupting chemicals and other substances of concern in food contact materials. An updated review of exposure, effect and risk assessment

Mechanisms and resistance in glucocorticoid control of inflammation

The effects of vitamin D on skeletal muscle function and cellular signaling

View the most downloaded articles for The Journal of Steroid Biochemistry and Molecular Biology

Provided by ScienceDirect Top25

Hide Applications

Related reference work articles e.g. encyclopedias

The Journal of Steroid Biochemistry and Molecular Biology
Volume 43, Issues 1-3, September 1992, Pages 173-177

doi:10.1016/0960-0760(92)90204-V | How to Cite or Link Using DOI
Permissions & Reprints

ICI 182,780, a new antioestrogen with clinical potential

Alan E. Wakeling^a and Jean Bowler^a

^aBioscience I and Chemistry I, ICI Pharmaceuticals, Mereside Laboratories, Alderley Park, Macclesfield, Cheshire SK10 4TG, England

Available online 17 January 2003.

Abstract

Previous studies in this laboratory identified a series of 7 α -alkylamide analogues of 17 β -oestradiol which are pure antioestrogens. Among this initial lead series of compounds, exemplified by ICI 164,384, none was of sufficient *in vivo* potency to merit serious consideration as a candidate for clinical evaluation. Further structure-activity studies identified a new compound, ICI 182,780, 7 α -[9-(4,4,5,5,5-pentafluoro-pentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, with significantly increased antioestrogenic potency. The antiuterotrophic potency of ICI 182,780 is more than 10-fold greater than that of ICI 164,384. ICI 182,780 has no oestrogen-like trophic activity and, like ICI 164,384 is peripherally selective in its antioestrogenic effects. The increased *in vivo* potency of ICI 182,780 was also reflected, in part, by intrinsic activity at the oestrogen receptor and in the growth inhibitory potency of ICI 182,780 in MCF-7 human breast cancer cells. ICI 182,780 was a more effective inhibitor of MCF-7 growth than 4'-hydroxytamoxifen, producing an 80% reduction of cell number under conditions where 4'-hydroxytamoxifen achieved a maximum of 50% inhibition. Sustained antioestrogenic effects of ICI 182,780, following a single parenteral dose of ICI 182,780 in oil suspension, were apparent in both rats and pigtail monkeys. *In vivo*, the antitumour activity of ICI 182,780 was demonstrated with xenografts of MCF-7 and Br10 human breast cancers in athymic mice where, over a 1 month period, a single injection of ICI 182,780 in oil suspension achieved effects comparable with those of daily tamoxifen treatment. Thus, ICI 182,780 provides the opportunity to evaluate clinically the potential therapeutic benefits of complete blockade of oestrogen effects in endocrine-responsive human breast cancer.


 To whom correspondence should be addressed.

Copyright © 1992 Published by Elsevier Ltd.

The Journal of Steroid Biochemistry and Molecular Biology
Volume 43, Issues 1-3, September 1992, Pages 173-177

- Personal Care Products and Breast Cancer
Encyclopedia of Environmental Health
- Estrogen and Progesterone Receptors in Breast Cancer
Encyclopedia of Hormones
- Steroid Hormones and Hormone Receptors
Encyclopedia of Cancer
- Breast Cancer
xPharm: The Comprehensive Pharmacology Reference
- Phytoestrogens
Encyclopedia of Hormones

▶ More related reference work articles



View Record in Scopus



LabChip XT

Accurately & reproducibly perform fast, automated nucleic acid fractionation, learn how here! [Learn More](#)

Nanomechanical Test Instruments Company

Learn more about the global leaders in the field of nanomechanical test instruments now! [Learn More](#)



Cell Symposia: Stem Cell Programming & Reprogramming

December 8 – 10, 2011, Lisbon, Portugal
Register before Oct 8 to save \$100 Register Now [Learn More](#)



Proactive Support Plan

Learn how our plan can help increase uptime & eliminate issues before they arise. Visit us now! [Learn More](#)



Back to results | < Previous 27 of 37 Next >

Home Browse Search My settings My alerts

Help

About ScienceDirect
What is ScienceDirect
Content details
Set up
How to use
Subscriptions
Developers

Contact and Support
Contact and Support

About Elsevier
About Elsevier
About SciVerse
About SciVal
Terms and Conditions
Privacy policy
Information for advertisers



Copyright © 2011 Elsevier B.V. All rights reserved. SciVerse® is a registered trademark of Elsevier Properties S.A., used under license. ScienceDirect® is a registered trademark of Elsevier B.V.



US 20100152149A1

(19) **United States**

(12) **Patent Application Publication**

Evans et al.

(10) **Pub. No.: US 2010/0152149 A1**

(43) **Pub. Date: Jun. 17, 2010**

(54) **FORMULATION**

(76) Inventors: **John R. Evans**, Macclesfield (GB);
Rosalind U. Grundy, Macclesfield (GB)

Correspondence Address:
MORGAN LEWIS & BOCKIUS LLP
1111 PENNSYLVANIA AVENUE NW
WASHINGTON, DC 20004 (US)

(21) Appl. No.: **12/285,887**

(22) Filed: **Oct. 15, 2008**

Related U.S. Application Data

(63) Continuation of application No. 10/872,784, filed on Jun. 22, 2004, now Pat. No. 7,456,160.

(30) **Foreign Application Priority Data**

Jan. 10, 2000 (GB) 0000313.7
Apr. 12, 2000 (GB) 0008837.7

Publication Classification

(51) **Int. Cl.**
A61K 31/566 (2006.01)
A61P 35/00 (2006.01)

(52) **U.S. Cl.** **514/182**

(57) **ABSTRACT**

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphanyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.




Figure 1

FLOW DIAGRAM OF MANUFACTURING




Figure 2

FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a Continuation Application of copending U.S. patent application Ser. No. 10/872,784, filed Jun. 22, 2004, which claims benefit of U.S. patent application Ser. No. 09/756,291, filed Jan. 9, 2001 which claims the benefit of Great Britain Application No. 0008837.7 filed Apr. 12, 2000 and Great Britain Application No. 0000313.7, filed Jan. 10, 2000, all of which are incorporated herein by reference in their entireties.

BACKGROUND OF THE INVENTION

[0002] 1. Field of the Invention

[0003] The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol.

[0004] 2. Description of the Related Art

[0005] Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

[0006] An alternative approach to oestrogen withdrawal is to antagonise oestrogens with antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Farr and Jordan 1984, May and Westley 1987).

[0007] The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for the design and testing of novel pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c, Wakeling and Bowler 1987, 1988.

[0008] Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989). One of these, 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. In vitro findings and early clinical experience with 7α [9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

[0009] 7α -[9-(4,4,5,5,5-Pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

[0010] Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells in vitro; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

[0011] Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

[0012] In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

[0013] European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

[0014] Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml^{-1} (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

[0015] Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

[0016] In the formulations within Table 1 a number of different oils are used to solubilise the compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

TABLE 1

OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS							
PRODUCT NAME	STEROID	DOSE	TYPE	COMP ¹	SOURCE	OIL	BzBz
SUSTANON 100	Testosterone propionate	30 mg	Androgen	Organon	ABPI Data Sheet Comp. 1999	Arachis	
	Testosterone phenylpropionate	60 mg					
	Testosterone isocaproate	60 mg					
	Testosterone decanoate	100 mg					
PROLUTON DEPOT	Hydroxy progesterone hexanoate	250 mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet Comp. 1999	Castor	up to 46%
TOCOGESTAN	Hydroxy progesterone enantate	200 mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%
	Progesterone	50 mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%
	α-Tocopherol	250 mg					
Estrapronicate	1.3 mg						
TROPHOBOLINE	Nandrolone undecanoate	50 mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%
	Hydroxyprogesterone heptanoate	80 mg					
	Norethisterone oenanthoate	200 mg					
NORISTERAT	Norethisterone oenanthoate	200 mg	Contraceptive	Schering HC	ABPI Data Sheet Comp. 1999	Castor	YES
BENZO-GYNOESTRYL	Estradiol hexahydrobenzoate	5 mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis	
	Hydroxy progesterone caproate	250 mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES
PROGESTERONE-RETARD	Estradiol 17-β-valerate	5 mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES
	Hydroxyprogesterone caproate	250 mgml ⁻¹					
GRAVIBINAN	Estradiol 17-β-valerate	5 mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES
PARABOLAN	Hydroxyprogesterone caproate	250 mgml ⁻¹					
DELESTROGEN	Trenbolone	76 mg	Androgen	Negma	Dict. Vidal 1997	Arachis	
	Estradiol valerate	20 mgml ⁻¹ 40 mgml ⁻¹	Estradiol	BMS	J. Pharm. Sci (1964) 53(8) 891	Castor	78% 58%
DELALUTIN	17-Hydroxy progesterone	250 mgml ⁻¹	Progestogen	DMS	J. Pharm. Sci. (1964) 53(8) 891	Castor	YES

PRODUCT NAME	STEROID	BzOH	EtOH	DOSE	DOSING
SUSTANON 100	Testosterone propionate	0.1 ml		1 ml	3 weeks
	Testosterone phenylpropionate				
	Testosterone isocaproate				
	Testosterone decanoate				
PROLUTON DEPOT	Hydroxy progesterone hexanoate			1 or 2 ml	1 week
TOCOGESTAN	Hydroxy progesterone enantate			2 ml	<1 week
	Progesterone				
	α-Tocopherol				
TROPHOBOLINE	Estrapronicate			1 ml	15 to 30 days
	Nandrolone undecanoate				
	Hydroxyprogesterone heptanoate				
NORISTERAT	Norethisterone oenanthoate			1 ml	8 weeks
BENZO-GYNOESTRYL	Estradiol hexahydrobenzoate			1 ml	1 week
	Hydroxy progesterone caproate			1 or 2 ml	1 week
PROGESTERONE-RETARD	Estradiol 17-β-valerate			1 or 2 ml	1-2 weeks
	Hydroxyprogesterone caproate				
GRAVIBINAN	Estradiol 17-β-valerate			1 or 2 ml	1-2 weeks
PARABOLAN	Trenbolone	75 mg	45 mg	1.5 ml	2 weeks
	Estradiol valerate	20% 40%	2% 2%		
	17-Hydroxy progesterone	YES	up to 2%		

BzBz = benzylbenzoate

BzOH = benzylalcohol

EtOH = ethanol

Dict. Vidal = Dictionnaire Vidal

% are w/v and

¹approximate as measured directly from a single sample

[text missing or illegible when filed]described which comprises 50 mg of fulvestrant, 400 mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml.

Manufacture at a commercial scale of a formulation as described in U.S. Pat. No. 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to

lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

[0017] The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound 7α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

BRIEF DESCRIPTION OF THE DRAWING

[0018] FIG. 1 shows the release profile in vivo of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

[0019] FIG. 2 shows a process flow diagram associated with the Formulation Example.

DETAILED DESCRIPTION OF THE INVENTION

[0020] Table 2 shows the solubility of fulvestrant in a number of different solvents.

TABLE 2

SOLUBILITY OF FULVESTRANT	
SOLVENT	SOLUBILITY (mgml ⁻¹ at 25° C.)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

[0021] As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil—see (Riffkin et. al. J. Pharm. Sci., (1964), 53, 891).

[0022] However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the

doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

[0023] Currently guidelines recommend that no more than 5 mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250 mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10 ml of castor oil.

[0024] The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of >50 mgml⁻¹ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of <5 ml—see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly causes the solubilisation of fulvestrant into a concentration of at least 50 mgml⁻¹—see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents—see Table 2 above—is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

[0025] Therefore, we present as a feature of the invention a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10% w/v, 4-9% w/v, 4-8% w/v, 4-7% w/v, 4-6% w/v and most preferably at about 5% w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0026] Another feature of the invention is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0027] Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0028] Further features of the invention include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45 mgml⁻¹ of fulvestrant.

[0029] For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight

will be present, for example a 1% weight per volume formulation will contain within a 100 ml volume of formulation 1 g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1 ml of formulation
30%	300 mg
20%	200 mg
10%	100 mg
5%	50 mg
1%	10 mg

[0030] Preferred pharmaceutical formulations of the invention are as described above wherein:

[0031] 1. The total volume of the formulation is 6 ml, or less, and the concentration of fulvestrant is at least 45 mgml⁻¹.

[0032] 2. The total amount of fulvestrant in the formulation is 250 mg, or more, and the total volume of the formulation is 6 ml, or less.

[0033] 3. The total amount of fulvestrant in the formulation is 250 mg and the total volume of the formulation is 5-5.25 ml.

[0034] It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5 ml dose is required it would be appreciated that an excess of up to 0.25 ml, preferably up to 0.15 ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

[0035] Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3% w/v, at least 5% w/v, at least 7% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and 3-35% w/v, 4-35% w/v, 5-35% w/v, 5-32% w/v, 7-32% w/v, 10-30% w/v, 12-28% w/v, 15-25% w/v, 17-23% w/v, 18-22% w/v and ideally 19-21% w/v.

[0036] The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

[0037] The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

[0038] The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

[0039] It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopoeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56° C. Dehydrated alcohol in the US Pharmacopoeia contains not less than 99.5% ethanol by volume when measured at 15.56° C.

[0040] Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v, at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less, 50% w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 5-60% w/v, 7-55% w/v, 8-50% w/v, 10-50% w/v, 10-45% w/v, 10-40% w/v, 10-35% w/v, 10-30% w/v, 10-25% w/v, 12-25% w/v, 12-22% w/v, 12-20% w/v, 12-18% w/v, 13-17% w/v and ideally 14-16% w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15% w/v.

[0041] It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

[0042] Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(% w/v)	Pharmaceutically-acceptable non-aqueous ester (% w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19- 21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

[0043] By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

[0044] We have surprisingly found that the above formulations of the invention provide, after intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

[0045] This finding is indeed surprising for the following reasons.

[0046] 1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.

[0052] By use of the term "extended release" we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2-5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

[0053] It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5 ml formulation is sequentially administered in two separate injections of 2.5 ml, this is a further feature of the invention

[0054] Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

[0055] Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

TABLE 3

Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25° C.

	% w/v							
Ethanol (96%)	5	5	10	10	10	15	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

[0047] 2. Our findings from studies using ¹⁴C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

[0048] It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

[0049] It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine—Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

[0050] We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

[0051] By use of the term "therapeutically significant levels" we mean that blood plasma concentrations of at least 2.5 ngml⁻¹, ideally at least 3 ngml⁻¹, at least 8.5 ngml⁻¹, and up to 12 ngml⁻¹ of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 1.5 ngml⁻¹.

[0056] The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

TABLE 4

Solubility comparisons of fulvestrant in oil based formulations with and without alcohols

Formulation ^(a)	Fulvestrant Solubility mg ml ⁻¹ @ 25° C.	
	Complete vehicle	Vehicle minus alcohols
Castor oil based	81.2	12.6
Miglyol 812-N based	86.8	1.7
Sesame seed/Castor oil (1:1) based	70.1	4.4
Sesame seed oil based	45.7	0.7
Arachis oil based	40.2	<0.2

^(a) Complete Vehicle Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined.

Effect of Formulation on Precipitation of Fulvestrant at the Injection Site

[0057]

Formulation ^a	Days						
	2	3	4	7	10	30	51
Formulation F1 castor oil based	0	0	0	0	0	0	0
Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
Formulation F3 sesame seed oil/ castor oil based	+ ^c	++	++	+++	++	+	+

0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)

^a Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.^b Mainly large needle shaped crystals^c Small needles and/or spherulites of crystals

[0058] Precipitation of fulvestrant and the release profile was determined with the above formulations in an in vivo rabbit study.

[0059] FIG. 1 shows the release profile in vivo of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits (data normalised to 50 mg per 3 kg; mean given; number of animals per timepoint=8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

[0060] Therefore we present as a further feature of the invention an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45 mgml⁻¹ of fulvestrant.

[0061] A further feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in medical therapy.

[0062] A further feature of the invention is a method of treating a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45 mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

[0063] Preferably 5 ml of the intramuscular injection is administered.

[0064] A further feature of the invention is use of fulvestrant in the preparation of a pharmaceutical formulation as

describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

[0065] Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

[0066] As described above fulvestrant is useful in the treatment of oestrogen-dependent indications such as breast cancer and gynaecological conditions, such as endometriosis.

[0067] In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

[0068] A further feature of the invention is a pharmaceutical formulation adapted for intra-muscular injection comprising 11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol; 35% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45 mgml⁻¹ of 11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol.

[0069] Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

FORMULATION EXAMPLE

[0070] Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2 μm porosity. The sterile filtrate is kept under a nitrogen overlay as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram of FIG. 2.

[0071] Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains

[0072] 10% weight per volume of benzyl alcohol

[0073] 10% weight per volume of ethanol

[0074] 15% weight per volume of benzyl benzoate

[0075] 250 mg of fulvestrant for each 5 ml of finished formulation

[0076] and the remaining amount as castor oil

REFERENCES

- [0077] 1. Bowler J, Lilley T J, Pittam J D, Wakeling A E. Novel steroidal pure antiestrogens. *Steroids* 1989; 54:71-99.
- [0078] 2. Wakeling A E. Novel pure antiestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
- [0079] 3. Wakeling A E. Steroidal pure antiestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
- [0080] 4. Wakeling A E. Therapeutic potential of pure antiestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- [0081] 5. Wakeling A E, Bowler J. Steroidal pure antiestrogens. *Journal Endocrinology* 1987; 112: R7-10.
- [0082] 6. Wakeling A E, Bowler J. Biology and mode of action of pure antiestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.
1. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least 2 weeks.
2. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for at least 4 weeks.
3. A pharmaceutical formulation as claimed in claim 1 wherein the blood plasma fulvestrant concentration is attained for 2 to 5 weeks.
4. A pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45 mgml⁻¹ of fulvestrant.
5. A pharmaceutical formulation as claimed in claim 1 or claim 4 which contains 25% w/v or less of a pharmaceutically-acceptable alcohol.
6. A pharmaceutical formulation as claimed in claim 5 which contains 20% w/v or less of a pharmaceutically-acceptable alcohol.
7. A pharmaceutical formulation as claimed in claim 1 or claim 4 which contains 60% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
8. A pharmaceutical formulation as claimed in claim 7 which contains 50% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
9. A pharmaceutical formulation as claimed in claim 7 which contains 45% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
10. A pharmaceutical formulation as claimed in claim 7 which contains 40% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
11. A pharmaceutical formulation as claimed in claim 7 which contains 35% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
12. A pharmaceutical formulation as claimed in claim 7 which contains 30% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
13. A pharmaceutical formulation as claimed in claim 7 which contains 25% w/v or less of a pharmaceutically-acceptable non-aqueous ester solvent.
14. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
15. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
16. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
17. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the total volume of the formulation is 6 ml, or less, and the concentration of fulvestrant is at least 45 mgml⁻¹.
18. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the total amount of fulvestrant in the formulation is 250 mg, or more, and the total volume of the formulation is 6 ml, or less.
19. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the total amount of fulvestrant in the formulation is 250 mg and the total volume of the formulation is 5 to 5.25 ml.
20. A pharmaceutical formulation as claimed in claim 1 or claim 4 wherein the pharmaceutically-acceptable alcohol is a mixture of 10% weight of ethanol per volume of formulation, 10% weight of benzyl alcohol per volume of formulation and 15% weight of benzyl benzoate per volume of formulation and the ricinoleate vehicle is castor oil.
21. A method of treating a benign or malignant diseases of the breast or reproductive tract by administration to a human in need of such treatment by intramuscular a pharmaceutical formulation as claimed in claim 1 or claim 4.
22. A method as claimed in claim 21 for treating breast cancer.
23. A syringe or vial containing a pharmaceutical formulation as defined in claim 20.

* * * * *

definite bearing on the usefulness of any column packing prepared. The performances of the seven supports mentioned previously were examined under the same operating conditions. The supports that can be used for lightly loaded packings are: glass beads, Gas Chrom-P, and Chromosorb W-HMDS. The other four supports cannot be used for lightly loaded column packing since their interaction with the antihistamines causes excessive peak tailing.

The hydrogen flame detector in conjunction with the 0.010-in. stainless capillary column would not respond to compounds with boiling points above 330°. This limitation prevented evaluation of this column for the analysis of these antihistamines.

The 100-ft. 0.065-in. copper open tubular column was coated with XF-1150 and evaluated using the above group of antihistamines. The Sr⁹⁰ ionization detector was used with a column flow of 36 ml./minute. The retention times obtained were comparable to the 6-ft.-XF-1150 packed column, but the peak base widths were considerably wider. Because of this increase in base width, the 0.065-in. column was less efficient than the 6-ft. packed column.

A 250-ft. 0.065-in. column wound on a 1¹/₄-in. diameter mandrel has been reported to be more efficient than a packed column (15). There are two possible reasons why efficiency was less than previously reported: (a) the column was shorter (100 ft.), and (b) the winding configuration was markedly different. The column was wound on a 1¹/₄ × 1/4-in. bar which resulted in a definite flattening of the tube around the edge of the bar.

CONCLUSIONS

The antihistamines investigated, except for meclizine, can be separated, identified, and concentration estimated using the Carbowax 20M, PDEAS, and XF-1150 columns described. The PDEAS column is the most efficient of the three for the analysis of antihistamines.

The usefulness of the 0.010-in. capillary and the 0.065-in. open tubular columns cannot be properly evaluated until the mentioned limitations are removed.

REFERENCES

- (1) Brochmann-Hanssen, E., and Svendsen, A. B., *THIS JOURNAL*, 51, 318(1962).
- (2) Cieplinski, E. W., *Anal. Chem.*, 35, 256(1963).
- (3) Parker, K. D., and Kirk, P. L., *ibid.*, 33, 428(1963).
- (4) *Ibid.*, 33, 1378(1961).
- (5) Brochmann-Hanssen, E., and Svendsen, A. B., *THIS JOURNAL*, 51, 938(1962).
- (6) Parker, K. D., Fontan, C. R., and Kirk, P. L., *Anal. Chem.*, 34, 1345(1962).
- (7) Fales, H. M., and Pisano, J. J., *Anal. Biochem.*, 3, 337(1962).
- (8) Parker, K. D., Fontan, C. R., and Kirk, P. L., *Anal. Chem.*, 34, 757(1962).
- (9) Anders, M. W., and Mannering, G. J., *J. Chromatog.*, 7, 258(1962).
- (10) Parker, K. D., Fontan, C. R., and Kirk, P. L., *Anal. Chem.*, 35, 356(1963).
- (11) Fontan, C. R., Smith, W. C., and Kirk, P. L., *ibid.*, 35, 591(1963).
- (12) Zubyk, W. J., and Conner, A. Z., *ibid.*, 32, 912(1960).
- (13) Averill, W., "Progress in Industrial Gas Chromatography," edited by Szymanski, H. A., Plenum Press, New York, N. Y., 1961, p. 225.
- (14) MacDonald, A., Jr., and Pfau, R. T., *THIS JOURNAL*, 52, 816(1963).
- (15) Quiram, E. R., *Anal. Chem.*, 35, 593(1963).

Castor Oil as a Vehicle for Parenteral Administration of Steroid Hormones

By C. RIFFKIN, R. HUBER, and C. H. KEYSER

Steroid hormones may be administered parenterally in high concentrations as oil solutions. In this form they exhibit a prolonged action and reduce the number of injections required. To accommodate the demand for increasingly greater concentrations of hormones in solution, castor oil in combination with other suitable oil-miscible solvents, has been found to fulfill a need. The development of several formulations together with the results of animal testing, as well as clinical trials in humans, attest to the acceptability of this oil for the purposes intended.

FIXED OILS are included in the "United States Pharmacopeia XVI" as nonaqueous vehicles for injection and are characterized as being of vegetable origin, essentially odorless, and without suggestion of rancidity. They must also comply with certain measurable physical limits specified for the saponification, acid, and iodine values.

Received September 9, 1963, from the Pharmacology and Pharmaceutical Research & Development Sections, Squibb Institute for Medical Research, New Brunswick, N. J.

Accepted for publication November 19, 1963.

The authors are indebted to the Chemical Control Laboratories for their assistance with the assays; to Dr. N. Coy and H. Roberts of the Physical and Analytical Section, for their development of special assays; to G. Lockwood of the Pharmacology Section for the animal muscle irritation tests; and especially to Dr. E. C. Reifstein, Jr., of the medical staff, for the information concerning the clinical trials.

After subcutaneous injection, Deanesly and Parkes (1) observed the persistence of olive oil and castor oil in animal tissue. Comparing other oils Brown, *et al.* (2), reported that sesame and corn oils were superior to cottonseed and peanut oils because they were less irritating, less antigenic, more quickly released from tissue, and possessed superior physical properties.

More recently the use of steroid hormone medication has expanded considerably. Due to limited water solubility, hormones have been administered as aqueous suspensions or solutions in oil. It has been claimed that the latter provided the slow release preferred in cyclical

TABLE I.—ANALYSIS OF COMMERCIAL OILS AND COMPARISON TO U.S.P. XVI SPECIFICATIONS

Oil	Lot No.	ml. 0.02 N NaOH Equiv. to Free Fatty Acid in 10-Gm. Sample	Sapon. Value	Iodine Value
Castor Oil	U.S.P. specs.	35.0 ^a	179-185	83-88
	23946	14.0	183.3	84.8
	25589	4.6	179.8	87.0
	23463	7.9	182.7	84.5
	33742	9.2	180.4	84.2
Sesame Oil	U.S.P. specs.	3.0	188-195	103-116
	23549A	0.5	189.6	106.9
	26953	1.4	194.0	111.8
	33646	0.75	189.6	104.7
	29981	0.45	191.7	108.2
Cottonseed Oil	U.S.P. specs.	2.0	190-198	109-116
	49684	...	195.9	111.8
	44441	...	196.3	113.1
Corn Oil	U.S.P. specs.	2.0	187-193	102-128
	52148	1.0	194.5	119.1
	36716	1.2	191.4	124.4
	33436	1.2	189.3	125.0
	33715	1.0	189.3	123.0
Peanut Oil	U.S.P. specs.	2.0	185-195	84-100
	22160	1.2	192.0	94.4
	20993	1.4	191.7	93.2
	33622	0.8	193.1	87.8
	26147	1.2	190.4	93.9

^a The U.S.P. specifies that the titration of free fatty acids in oral grade castor oil shall not exceed 7 ml. of 0.1 N NaOH which is equal to 35.0 ml. of 0.02 N NaOH.

TABLE II.—SOLUBILITY OF STEROIDS IN U.S.P. OILS AT 25°

Steroid	mg./ml.		
	Castor Oil	Sesame Oil	Peanut Oil
17-Hydroxyprogesterone caproate	55.6	23.4	27.9
Testosterone	38.6	5.4	8.1
Estradiol valerate	60.6	16.1	18.8
Progesterone	52.0	22.9	23.5

therapy (3). Using withdrawal bleeding in human females as the criterion, Master, *et al.* (4), compared the duration of action of an aqueous suspension of progesterone with an oil solution, and confirmed the superiority of the latter. The prolongation of activity was generally related to storage in the fatty depots of the body (5).

In 1952 Junkmann (6) determined that a testosterone ester dissolved in sesame oil prolonged the androgenic effects in castrated rats. Davis and Wied (7) demonstrated that prolonged activity was also obtained in humans when oil solutions of a progesterone derivative were injected. There was still a limiting factor, however, in that only a relatively small amount of hormone could be dissolved in the traditional oils. To increase the solvent power of the oil it was necessary to add compatible and non-irritating cosolvents. Such additions consisted of benzyl benzoate, benzyl alcohol, ethyl lactate, ethyl oleate, etc. The U.S.P. recognized the need for such "other vehicles," with the restrictions that they must be safe in the volume of injection administered, and that they should

not interfere with the therapeutic efficacy of the preparation or its testing.

Demand for increased hormone concentrations per dose, furthered the search for an acceptable oil with greater solubilizing power *per se*. Boschann (8) in 1954, observed that 17-hydroxyprogesterone caproate in a castor oil-ethyl lactate vehicle was well tolerated. In addition, private communications from clinicians in West Germany¹ reported good tolerance to Proluton-Depot containing a castor oil-benzyl benzoate vehicle. Since then other hormones have been used as solutions in ricinoleic acid esters, as well as in castor oil (9-11). Accordingly, an investigation was undertaken into the suitability of castor oil as a vehicle for parenteral administration of steroid hormones.

METHODS AND RESULTS

Representative samples of U.S.P. oils obtained from commercial sources were tested in accordance with the official method for free fatty acid content, saponification, and iodine values. The results are listed in Table I along with the U.S.P. XVI specifications for these oils.

Solubility of selected steroids in various oils was determined in the following manner. An excess of steroid was stirred for 4 hours at room temperature (25°) in the test oil, after which the undissolved solids were removed by filtration, and the clear solution assayed for steroid content. Table II shows the results obtained.

An attempt was made to reduce the free fatty acids in castor oil by treatment with alumina and anhy-

¹ Dr. Napp, Universitäts-Krankenhaus, Hamburg; Dr. Pots, Humboldt-Universität-Charité Frauenklinik, Berlin; Dr. Prill, Universitäts-Frauenklinik, Würzburg; and Dr. Rauscher, Universitäts-Frauenklinik, Vienna.

TABLE III.—ABSORPTION OF OIL FROM ANIMAL MUSCLE^a

Days after Injection	Oil	ml. 0.02 N NaOH Equiv.	Residual Oil in Muscle (estd.)
1-3	Castor oil (aged)	50	1 day —50% 3 days—20%
1-3	Castor oil U.S.P.	13	1 day —30% 3 days—10%
1-3	Sesame oil U.S.P.	1.4	1 day —30% 3 days—30%
7-60	All oils	...	Declining 10 to 2%

^a 1 ml. injected into back muscle of rabbit.

drous sodium sulfate. Three grams of dried, powdered, amorphous aluminum oxide (Merck No. 1097) and 6 Gm. of anhydrous sodium sulfate, reagent grade, were suspended in 120 ml. of oil and heated at 80° under a blanket of nitrogen for 1.5 hours. After allowing the oil to cool to room temperature, the solids were filtered off and the acids titrated in the usual manner. A significant reduction in free fatty acid was not obtained.

The absorption characteristics of oils with varying fatty acid content were examined and compared on a biological basis. Aged castor oil with a high free fatty acid content was compared to fresh U.S.P. castor oil with a low acid content and U.S.P. sesame oil by injecting 1 ml. of oil into the back muscles of rabbits, approximately 2 in. from the iliac crest. A rotational pattern of injection was used and the oil samples were stained to aid visibility in the tissues. The animals were sacrificed and the muscles excised and examined grossly. The results were averaged and appear in Table III.

The test disclosed that oil migrated or was carried to the fascia, and very small amounts remained for 60 days. Localized degeneration produced by the high acid value castor oil was essentially healed in 7 days, and the low acid value castor oil appeared to be no more irritating than sesame oil.³

In a specific test for irritation 0.25 ml. of the above oil samples were also injected into the *vastus lateralis* muscles of rabbits. After 2 days the animals were sacrificed and the injected muscles examined grossly for evidence of irritation. It was found that the castor oil containing a high level of free fatty acid produced a lesion size measuring approximately 121 mm.³. The lesion itself was characterized mainly by degeneration of local tissue without necrosis. Castor oil with low free fatty acid and sesame oil, on the other hand, produced no measurable lesion at the injection site.

Combinations of benzyl alcohol and benzyl benzoate with both castor oil and sesame oil were also injected into the *vastus lateralis* muscles of rabbits and Table IV lists the lesion sizes produced.

Solutions which were formulated for clinical trials in humans were prepared by dissolving the steroid hormones in appropriate vehicles at 60° under nitrogen. The solutions were then filtered through a coarse sintered-glass filter with the aid of nitrogen pressure, filled into vials, and sterilized by autoclaving for 2 hours at 121° (15 lb. steam pressure). The products were then submitted for assay, safety, and

³ Due to the apparent increase in free fatty acids with aging, subsequent work utilized fresh oils which required for neutralization less than 3 ml. of 0.1 N NaOH (15 ml. of 0.2 N NaOH) per 10 Gm. of sample.

animal muscle irritation testing prior to release for clinical investigation.

DISCUSSION

Throughout the investigation it was desirable to have a reference oil to serve as a basis for comparison. Since sesame oil is universally accepted as a parenteral oil vehicle, it was chosen as the "standard" vegetable oil to be compared to castor oil, with and without other cosolvents. The physical, chemical, and biological properties of sesame oil are well documented and require no comments here.

Chemically, castor oil consists of the triglycerides of ricinoleic acid, together with small quantities of glycerides of other acids. The quantitative composition is given by Eckey (12) as follows: ricinoleic acid 87%, oleic acid 7.4%, linoleic acid 3.1%, dihydroxyricinoleic acid 0.6%, and miscellaneous acids 2.4%. Two grades are commonly recognized in this country—U.S. No. 1 which is cold pressed oil, and U.S. No. 3 which is oil extracted from the pressed cake. Only the former is used for medicinal purposes.

The high viscosity of castor oil compared to other vegetable oils is undoubtedly related to hydrogen bonding and it is probably the hydroxy groups which contribute to the greater polarity and superior solvent power of the oil. As indicated in Table I, the saponification and iodine values of commercial castor oil appear to be slightly lower than the U.S.P. XVI limits for oils used for injection. On the other hand, the content of free fatty acids even in fresh oil, varies considerably and exceeds the traditional limits for injectable oils. The significance of this is somewhat obscure, although "Remington's Practice of Pharmacy, 12th edition," page 387, states "a low free fatty acid content is essential since it indicates a fresh and pure product and not one that is likely to have become old and heavily contaminated with bacterial products."

Despite better solubility of steroids in castor oil, other cosolvents were necessary to dissolve the

TABLE IV.—LOCAL IRRITATION PRODUCED IN RABBIT MUSCLE BY INJECTION OF VARIOUS OIL VEHICLES^a

Identification	Composition	Lesion size, mm. ³
SHY-47-2	Sesame oil 98% Benzyl alcohol 2%	61
SHY-47-4	Castor oil 98% Benzyl alcohol 2%	Too small to measure
SHY-47-3	Sesame oil 95% Benzyl alcohol 5%	506
SHY-47-5	Castor oil 95% Benzyl alcohol 5%	106
SHY-14-2	Sesame oil 65% Benzyl benzoate 35%	291
SHY-14-5	Castor oil 65% Benzyl benzoate 35%	184
SHY-47-6	Sesame oil 63% Benzyl benzoate 35% Benzyl alcohol 2%	207
SHY-47-7	Castor oil 63% Benzyl benzoate 35% Benzyl alcohol 2%	262
SHY-14-3	Sesame oil 50% Benzyl benzoate 50%	291
SHY-14-6	Castor oil 50% Benzyl benzoate 50%	158

^a A 0.25-ml. quantity of the oil vehicle was injected into the *vastus lateralis* muscle of the rabbit. Two days later the muscle was excised and the lesion size measured in mm.³.

increasingly higher concentrations required by therapeutic regimens. Often these materials contributed additional advantages. For example, the addition of benzyl alcohol or benzyl benzoate to castor oil resulted in a lower and more favorable viscosity, making it easier to inject. Also, benzyl alcohol was an effective preservative and local anesthetic.

The nature of the irritative response depended on the particular hormone, its concentration in the formulations, and/or the composition of the vehicle. Although rabbit muscles are more sensitive than human muscles, they were selected primarily because local changes in the muscle were observed easily. It was not always possible, however, to correlate muscle irritation in animals to that of humans.

A numerical assignment to lesion size was used solely as a convenience for grading response. The numbers alone do not adequately describe the nature of the response, however. More completely it is characterized by the amount of hemorrhage and edema and the incidence, degree, and extent of local degeneration produced by the injection. A slight, reversible irritative response may cover a large area and a severe irreversible one may be comparatively small. A decrease in the size of the degenerated area indicates a reversible condition. The presence of necrosis, which is the most damaging situation, means that the cellular structure was destroyed and repair must take place. The debris must be removed and the original cellular mass in the area replaced with fibrous connective tissue. The extent of this fibrosis or formation of scar tissue gives an index of the amount of irreversible damage. Fortunately necrosis was not encountered, indicating the lack of permanent muscle damage. Since these changes take time, final assessment of the effects of an injection in the muscle frequently required observation for 7 days or longer.

It is unfortunate that pain cannot be measured by any known method of animal testing. The animal usually does not respond unless the painful stimulus is marked. Furthermore, the pain caused by injection into human muscle is not usually proportionate to the irritation produced either in animal muscle or in human muscle. Realizing that these limitations are inherent in animal test methods, it remained for final acceptability to be determined in man.

When it was discovered that 17-hydroxyprogesterone caproate possessed high progestational activity, potencies of the order of 65 mg./ml. were used. By increasing the dose, additional prolongation of action was obtained, and eventually concentrations of the order of 250 mg./ml. were required. Such a solution in sesame oil produced acceptable animal muscle tolerance, but the pain and local reaction in humans was so great as to prohibit the adoption of the formulation as a commercial product (see Table V, Lot Pr. 142-53/15-10).³ Solutions were also prepared using castor oil as the vehicle, and Table V lists the formulations tested and the results obtained. Information obtained from the clinical trials (14-21) attested to the acceptability and safety of the adopted formulations.

Inherent in the development of an acceptable formulation of 17-hydroxyprogesterone caproate was

³ Reactions in excess of 5-6% were considered unacceptable.

TABLE V.—EVALUATION OF 250 mg./ml. 17-HYDROXYPROGESTERONE CAPROATE SOLUTIONS IN VARIOUS OIL VEHICLES

Vehicle Composition	Animal Muscle Lesion Size, mm. ^a	Lot Number and Remarks on Clinical Testing
Sesame oil 50% Benzyl benzoate 50%	1049	Pr.142-53/15-7—238 injections, 20.6% reactions, rejected
Castor oil 58% Benzyl benzoate 40%	691	Pr.142-53/15-8—270 injections, 23.2% reactions, rejected
Benzyl alcohol 2% Sesame oil 60% Benzyl benzoate 35%	697	Pr.142-53/15-10—189 injections, 10.7% reactions, rejected
Benzyl alcohol 5% Castor oil 54% Benzyl benzoate 46%	258	Pr.142-53/15-11—503 injections, 4.2% reactions, accepted
Castor oil 52% Benzyl benzoate 46% Benzyl alcohol 2%	633	Pr.142-53/15-13—924 injections, 1.3% reactions, accepted

^a Injection of 0.25 ml. into *vastus lateralis* muscle of rabbits and lesion size determined 2 days after injection.

TABLE VI.—EVALUATION OF ESTRADIOL VALERATE IN VARIOUS OIL VEHICLES

Composition	Animal Muscle Lesion Size, mm. ^a	Lot Number and Remarks
20 mg./ml. in Castor oil 78%, Benzyl benzoate 20%, Benzyl alcohol 2%	197	Es.31-53/15-B—Commercially available
30 mg./ml. in Sesame oil 60%, Benzyl benzoate 40%	306	DEK-98-2—Not tested clinically; dosage increased to 40 mg./ml.
30 mg./ml. in Castor oil 80%, Benzyl benzoate 20%	194	Es.31-53-V—Not tested clinically; dosage increased to 40 mg./ml.
40 mg./ml. in Sesame oil 65%, Benzyl benzoate 30%, Benzyl alcohol 5%	803	SHX-94-4—Too irritating; not tested clinically
40 mg./ml. in Sesame oil 58%, Benzyl benzoate 40%, Benzyl alcohol 2%	496	Es.31-53-8—201 injections, 23.2% reactions, rejected
40 mg./ml. in Castor oil 58%, Benzyl benzoate 40%, Benzyl alcohol 2%	250	Es.31-53-A—826 injections, 2.67% reactions (all mild), accepted

^a Injection of 0.25 ml. into *vastus lateralis* muscle of rabbits and lesion size determined 2 days after injection.

the required development of a suitable assay method. This was accomplished by Roberts and Florey (13) using paper-strip chromatography.

Since estrogens are more potent than progestogens and require less per dose, an acceptable formulation of estradiol valerate was easier to prepare. Besides use in estrogen therapy, estradiol valerate has found utility in the treatment of carcinoma, and for that purpose high dosages were required. Concentrations were increased from 10 to 40 mg./ml. and

again formulations containing castor oil in the vehicle proved to be less irritating than similar preparations containing sesame oil. Physically and chemically both oil solutions were stable. Based on acceptable preliminary data, formulations such as those listed in Table VI were prepared and tested. Acceptability in humans was confirmed by clinicians and described in the literature (22, 23) and in case reports.⁴

SUMMARY

1. The development and testing of parenteral steroid hormone formulations has been described, using castor oil as a vehicle.

2. After ascertaining stability and animal muscle irritation, selected formulations were evaluated in humans. They exhibited a prolonged action, were effective and well tolerated.

3. Examples of commercially available products are the estrogen, estradiol valerate⁵ at 20 mg./ml. and 40 mg./ml., and the progestogen, 17-hydroxyprogesterone caproate⁶ at 250 mg./ml.

⁴ Case reports: estradiol valerate, 20 mg./ml. in castor oil 78%, benzyl benzoate 20%, benzyl alcohol 2%—90 injections in 46 patients. Two mild local reactions. Estradiol valerate 40 mg./ml. in castor oil 58%, benzyl benzoate 40%, benzyl alcohol 2%—51 patients. Number of injections not completely tabulated. One report is in press.

⁵ Marketed as Delastrogen by E. R. Squibb & Sons, New York, N. Y.

⁶ Marketed as Delalutin by E. R. Squibb & Sons, New York, N. Y.

REFERENCES

- (1) Deanesly, R., and Parkes, A. S., *J. Physiol.*, **78**, 155 (1933).
- (2) Brown, W. E., Wilder, V. M., Schwartz, P., *J. Lab. Clin. Med.*, **29**, 259(1944).
- (3) "Hormone Therapy in Practice," 2nd ed., Schering, A. G., Erich Blaschker, Berlin, 1954, p. 109.
- (4) Master, W. H., Grody, M. H., and Magallon, D. T., *J. Clin. Endocrinol.*, **12**, 1445(1952).
- (5) "Modern Trends in Endocrinology," H. Gardiner-Hill, ed., P. B. Hoeber, 1958, p. 233.
- (6) Junkmann, K., *Arch. Exptl. Pathol. Pharmacol.*, **215**, 85(1952).
- (7) Davis, M. E., and Wied, G. L., *J. Clin. Endocrinol. Metab.*, **15**, 923(1955).
- (8) Boschmann, H. W., *Arch. Wochenschr.*, **9** (25) 591 (1954).
- (9) Richter, U. S. pat. 2,822,316.
- (10) Brit. pat. 817,241.
- (11) Ercoli, A., U. S. pat. 2,983,649.
- (12) Eckey, E. W., "Vegetable Fats and Oils," A.C.S. Monograph, Reinhold Publishing Co., New York, N. Y., Series No. 123, pp. 587-597.
- (13) Roberts, H. R., and Florey, K., *THIS JOURNAL* **51**, 794 (1962).
- (14) Short, C. L., *Am. Practitioner Dig. Treatment*, **11**, 149(1960).
- (15) Greenblatt, R. B., and Dutta, S. N., *Ariz. Med.*, **18** (4), 107(1961).
- (16) Kelley R. M., and Baker, W. H., *New Engl. J. Med.*, **264**, 216(1961).
- (17) Kennedy, B. J., *J. Am. Med. Assoc.*, **184**, 758(1963).
- (18) Danforth, D. N., and Buckingham, J. C., *Postgrad. Med.*, **32**, 345(Oct. 1962).
- (19) Kistner, R. W., *Clin. Obstet. Gynecol.*, **5**, 1166(1962).
- (20) Siegel, I., *Obstet. Gynecol.*, **21**, 666(1963).
- (21) Pellegrino, L., *Current Therap. Res.*, **4**, 301(1962).
- (22) MacDonal, I., and Yettra, M., *Med. Clin. N. Am.*, **43**, 971(1959).
- (23) Eichner, E., Brown, M., and Sable, M., *J. Intern. Coll. Surgeons*, **32**, 394(1959).

Isolation of Marrubiin, a Sterol, and a Sesquiterpene from *Marrubium vulgare*

By HAROLD J. NICHOLAS*

A simple column chromatographic method for isolating the bicyclic diterpene marrubiin from acetone and ethanol extracts of *Marrubium vulgare* L. is described. An unsaturated sterol of the stigmastanol series, present in esterified form, and a sesquiterpene (C₁₅H₂₂O₂) have been isolated from the extracts.

IN PREPARATION for radioactive tracer work on the biosynthesis of marrubiin it was necessary to examine extracts of the plant for associated terpenoid substances. A convenient column chromatographic method was therefore devised for separating relatively pure marrubiin from crude acetone extracts. Two new terpenoid substances were detected in the extracts.

EXPERIMENTAL

Materials and Methods.—Ground *M. vulgare* L. was obtained from the Wunderlich-Diez Corp.,

Received August 15, 1963, from the Department of Obstetrics and Gynecology, University of Kansas Medical Center, Kansas City.

Accepted for publication November 6, 1963.

This investigation was supported by a grant from the National Institutes of Health, U. S. Public Health Service, Bethesda, Md.

The author is indebted to Fuad Jarjoura and Sharon Moriarity for their technical assistance.

* Present address: Institute of Medical Education and Research, St. Louis, Mo., and Department of Biochemistry, St. Louis University School of Medicine, St. Louis, Mo.

Hasbrouck Heights, N. J.¹ This material was exhaustively extracted with hot acetone or hot ethanol. Either solution on removal of solvent by distillation (the last stages *in vacuo*) yielded black, viscous material which was used for further examination. Melting points were determined on a Fisher-Johns melting point apparatus. Optical rotations (in CHCl₃) and C—H analyses were determined by Drs. G. Weiler and F. B. Strauss, Microanalytical Laboratory, Oxford, England. An infrared spectrum of the unidentified diterpene was determined on a Perkin-Elmer spectrophotometer by the KBr disk method.² An infrared spectrum of the sterol was determined in chloroform solution in a 0.1-mm. sealed cell, compensated with CHCl₃, on a Beckman IR-4 recording infrared spectrophotometer,³ and by the KBr disk method. The

¹ This firm has given assurance that the material investigated was *M. vulgare* or white horehound, not *Ballota hirsuta* (black horehound).

² We are indebted to the Department of Pathology, University of Kansas, for this determination.

³ Determined by Sadtler Research Laboratories, Philadelphia, Pa.

Excipients and Their Use in Injectable Products

SANDEEP NEMA*, R. J. WASHKUH, and R. J. BRENDEL

Mallinckrodt Medical, Incorporated, Saint Louis, Missouri

ABSTRACT: Formulation of a new drug product with excipients, that have been previously added to an approved injectable product, may save pharmaceutical companies developmental time and cost. The Physicians' Desk Reference (PDR) and Handbook on Injectable Drugs were reviewed, extracting all information on excipients. The information was consolidated into eight tables, categorizing excipients as 1) Solvents and Co-solvents, 2) Solubilizing, Wetting, Suspending, Emulsifying or Thickening agents, 3) Chelating Agents, 4) Antioxidants and Reducing Agents, 5) Antimicrobial Preservatives, 6) Buffers and pH Adjusting Agents, 7) Bulking Agents, Protectants, and Tonicity Adjustors, and 8) Special Additives. Where applicable, tables list frequency of use, concentration, and an example of a commercial product containing the excipient. Excipients which are included in the 1996 FDA 'Inactive Ingredient Guide,' but do not appear in the PDR or Handbook on Injectable Drugs, were included as a separate list.

Introduction

Injectable products require a unique formulation strategy. The formulated product has to be sterile, pyrogen free and, in the case of solutions, free of particulate matter. Preferably, the formulation will be isotonic, and depending on the route of administration (for instance, for intra-spinal or intra-cisternal routes), antioxidants and preservatives may not be allowed. For a given drug, the risk of adverse events is higher if it is administered as an injection versus a non-parenteral route. The requirement for sterility demands that the excipients be able to withstand autoclaving or other sterilization processes. These factors limit the choice of excipients available to the formulators.

Generally, a knowledge of which excipients have been deemed safe by the FDA or are already present in a marketed product provides increased assurance to the formulator that these excipients will probably be safe for their new drug product. However, there is no guarantee that the new drug product will be safe as excipients are combined with other additives and/or with a new drug, creating unforeseen potentiation or synergistic toxic effects. Regulatory bodies may view an excipient previously approved in an injectable dosage form favorably, and will frequently require less safety data. A new additive in a formulated product will always require additional studies adding to the cost and timeline of product development.

The purpose of this paper is to present the various excipients that have been included in the formulation of injectable products marketed in the USA. This information is not readily available. A literature search indicates that the last paper dealing with this was published in 1980 (1). Products approved outside the US are not covered in this

review. Also, sterile dosage forms not administered parenterally, such as solutions for irrigation, ophthalmic or otic drops, and ointments were excluded.

Methodology

Physicians' Desk Reference published in 1994 & 1996 (2, 3), and Handbook on Injectable Drugs (4) were used as the primary source of information. Entries on all injectable drugs were summarized in an Excel worksheet. Each product was classified by Manufacturer, Trade name, Drug name, Route of Administration, SVP/LVP, pH of Product, Solvent Used, Solubilizing/Suspending Agent, Preservative, Antioxidant, Chelator and Other Formulation Additives.

The resulting Excel sheet had information on more than 700 products. This information was condensed into easy-to-read tables. Each table has been categorized based on the primary function of excipient in the formulation. For example, citrates are classified as buffers and not as chelating agents, and ascorbates are categorized as antioxidants, although they can serve as buffers. This classification system was based on our experience in formulation development and on the published literature. Such simplification avoids duplication of entries and provides the audience with easy-to-read tables.

Some duplication was unavoidable. Tables VII and VIII contain some excipients which may have also been listed in the first six tables. Whenever the reference specifically designated a specific function to an ingredient it was re-listed in Tables VII and VIII. For example, glycine can be used as a buffer or as a stabilizing (protecting) agent. Therefore, glycine is listed in Tables VI and VII. Methyl paraben is a preservative (Table V) but also has a special function in Adriamycin RDI[®] formulation (Table VIII).

The concentration of excipients is listed as percentages weight by volume (w/v) or volume by volume (v/v). If the product was listed as lyophilized or powder, these percent-

Received October 1, 1996. Accepted for publication May 16, 1997.

*Author to whom correspondence should be addressed: P.O. Box 5840, St. Louis, MO, 63134

TABLE I
Solvents and Co-solvents

Excipient	Frequency	Range	Example
Benzyl Benzoate	2	20% v/v	Depo-Testosterone [®] (Upjohn) 20% v/v
Cottonseed Oil	1	73.6% w/v	Depo-Testosterone [®] (Upjohn) 73.6% w/v
N,N Dimethylacetamide	1	6% w/v	Vumon [®] (Bristol Myers) 6% w/v
Ethanol	24	0.6-80%	Prograf [®] (Fujisawa) 80% w/v
Glycerin (Glycerol)	9	1.6-70% w/v	Multitest CMI [®] (Connaught) 70% w/v
Peanut oil	1	*	Bal In Oil [®] (Becton Dickinson)
Polyethylene glycol			
PEG	4	0.15-50%	Secobarbital sodium (Wyeth-Ayerst) 50%
PEG 300	2	50-65%	VePesid [®] (Bristol Myers) 65% w/v
PEG 400	2	*	Ativan [®] (Wyeth-Ayerst)
PEG 3350	5	0.3-3%	Depo-Medrol [®] (Upjohn) 2.95% w/v
Poppyseed oil	1	1%	Ethiodol [®] (Savage) 1%
Propylene Glycol	25	0.2-75.2%	Terramycin Solution (Roerig) 75.2%
Safflower oil	2	5-10%	Liposyn II [®] (Abbott) 10%
Sesame oil	6	*	Solganal Inj. [®] (Schering)
Soybean oil	4	5-20% w/v	Intralipid [®] (Clintec) 20%
Vegetable oil	2	*	Virilon IM Inj. [®] (Star Pharmaceuticals)

* No data available.

ages were derived based on the reconstitution volume commonly used. The tables list the range of concentration used, typical or most common concentration employed, and examples of products containing the excipient, specifically those which use extremely low or high concentrations.

Discussions

Table I list solvents and co-solvents used in parenteral products. Water for injection is the most common solvent but may be combined or substituted with a co-solvent to improve the solubility or stability of drugs. Oils like safflower and soybean are used in total parenteral nutrition products where they serve as a fat source and as carriers for fat-soluble vitamins. Ethanol and propylene glycol are used, either alone or in combination with other solvents, in more than 50% of parenteral co-solvent systems. It is surprising to see propylene glycol used more often than polyethylene

glycols (PEGs) in spite of its higher myotoxicity and hemolyzing effects (5, 6). Probably, the presence or generation of peroxides in PEGs is a major limitation.

Table II includes a broad category of excipients whose function in formulation could be—(1) Viscosity imparting or suspending agents like carboxy methyl cellulose, sodium carboxy methyl cellulose, sorbitol, acacia, Povidone, hydrolyzed gelatin; (2) Solubilizing, wetting or emulsifying agents like Cremophore EL, sodium desoxycholate, Polysorbate 20 or 80, PEG 40 castor oil, PEG 60 castor oil, sodium dodecyl sulfate, lecithin or egg yolk phospholipid; (3) Aluminum monostearate which is added to fixed oil to form viscous or gel-like suspending medium. Polysorbate 80 is the most common and versatile solubilizing, wetting and emulsifying agent.

Only a limited number of chelating agents are used in parenteral products (Table III). They serve to complex heavy

TABLE II
Solubilizing, Wetting, Suspending, Emulsifying or Thickening Agents

Excipient	Frequency	Range	Example
Acacia	2	7%	Tuberculin Old Test [®] (Lederle) 7%
Aluminum monostearate	1	2%	Solganal Inj. [®] (Schering) 2%
Carboxy methyl cellulose	4	1%	Bicillin [®] (Wyeth-Ayerst) 0.55%
Carboxy methyl cellulose, sodium	9	0.1-0.75%	Lupron Depot [®] (TAP) 0.75% w/v
Cremophore EL [*]	3	50-65% w/v	Sandimmune [®] (Sandoz) 65% w/v
Desoxycholate sodium	1	0.4% w/v	Fungizone [®] (Bristol Myers) 0.41% w/v
Egg yolk phospholipid	3	1.2%	Intralipid [®] (Clintec) 1.2%
Gelatin, Hydrolyzed	1	16% w/v	Cortone [®] (Merck) 16% w/v
Lecithin	7	0.4-1.2% w/v	Diprivan [®] (Zeneca) 1.2% w/v
Polyoxyethylated fatty acid	1	7% w/v	AquaMephyton [®] (Merck) 7% w/v
Polysorbate 80 (Tween 80)	31	0.01-12%	Cardarone X L.v. [®] (Wyeth-Ayerst) 10%
Polysorbate 20 (Tween 20)	5	0.01-0.4%	Calcijex [®] (Abbott) 0.4% w/v
PEG 40 castor oil**	1	11.5% v/v	Monistat [®] (Janssen) 11.5% v/v
PEG 60 castor oil***	1	20% w/v	Prograf [®] (Fujisawa) 20% w/v
Povidone (Polyvinyl pyrrolidone)	6	0.5-0.6% w/v	Bicillin [®] (Wyeth-Ayerst) 0.6% w/v
Sodium dodecyl sulfate (Na lauryl sulfate)	1	0.018% w/v	Proleukin [®] (Cetus) 0.018% w/v
Sorbitol	3	25-50%	Aristrospan [®] (Fujisawa) 50% v/v

* Cremophor EL: Etoacs 35, polyethoxylated castor oil, polyoxyethylene 35 castor oil.

** PEG 40 castor oil; polyoxyl 40 castor oil, castor oil POE-40, Croduret 40, polyoxyethylene 40 castor oil, Protachem CA-40.

*** PEG 60 hydrogenated castor oil; Cremophor RH 60, hydrogenated castor oil POE-60, Protachem CAH-60.

TABLE III
Chelating Agents

Excipient	Frequency	Range	Example
Calcium disodium EDTA*	9	0.01-0.1%	Wydase® (Wyeth-Ayerst) 0.1% w/v
Disodium EDTA	34	0.01-0.1%	Calcijex® (Abbott) 0.11% w/v
Sodium EDTA	1	0.20%	Folvite® (Lederle) 0.2%
DTPA**	1	0.04%	Magnevist® (Berlex) 0.04%

* EDTA = Ethylenediaminetetraacetic acid.

** DTPA = Diethylenetriaminopentaacetic acid; Pentetic acid.

metals and therefore can improve the efficacy of antioxidants or preservatives. In our opinion, calcium EDTA has an advantage over tetrasodium salt by not contributing sodium and not chelating calcium from the blood.

An antioxidant as a class is defined as those compounds that can act as reducing agents or may serve as free radical scavengers. Table IV summarizes the antioxidants, their frequency of use, concentration range and examples of products containing them. Sulfite, bisulfite, and metabisulfite constitute the majority of antioxidants used in parenteral products despite several reports of incompatibilities and

toxicity (7, 8). Butylated hydroxy anisole, butylated hydroxy toluene and propyl gallate are primarily used in semi/non-aqueous vehicles because of their low aqueous solubility. Ascorbic acid/sodium ascorbate may serve as an antioxidant, buffer, and chelating agent in the same formulation.

Benzyl alcohol was the most common antimicrobial preservative present in parenteral formulations (Table V). This is consistent with other surveys (9). Parabens are the next most common preservatives. Thirty-nine products had a combination of methyl and propyl parabens; eleven had only methyl, and one had only propyl paraben. Thimerosal was surprisingly common, especially in vaccines, even though some individuals have sensitivity to mercurics. Chlorocresol is purported to be a good preservative for parenterals, but our survey did not find any examples of commercial products containing chlorocresol.

Table VI lists buffers and chemicals used to adjust the pH of formulations. Phosphate, citrate, and acetate are the most common buffers used in parenteral products. Mono and diethanolamine are added to adjust pH and form corresponding salts. Hydrogen bromide, sulfuric acid, benzene sulfonic acid and methane sulfonic acids are added to drugs which are bromide (Scopolamine HBr, Hyoscine HBr, UDL), sulfate (Nebcin, Tobramycin sulfate, Lilly), besylate

TABLE IV
Antioxidants and Reducing Agents

Excipient	Frequency	Range	Example
Acetone sodium bisulfite	4	0.2-0.4% w/v	Novocaine® (Sanoft-Winthrop) 0.4% w/v
Ascorbate (sodium/acid)	7	0.1-4.8% w/v	Vibramycin® (Roerig) 4.8% w/v
Bisulfite sodium	28	0.02-0.66% w/v	Amikin® (Bristol Myers) 0.66% w/v
Butylated hydroxy anisole (BHA)	3	0.0028-0.03% w/v	Aquasol® (Astra) 0.03%
Butylated hydroxy toluene (BHT)	3	0.00116-0.03% w/v	Aquasol® (Astra) 0.03%
Cystein/Cysteinate HCl	2	0.07-0.10% w/v	Acthar Gel® (Rhône-Poulanc) 0.1% w/v
Dithionite sodium (Na hydrosulfite, Na sulf-oxylate)	1	0.10%	Numorphan® (DuPont) 0.10%
Gentisic acid	1	0.02% w/v	OctreoScan® (Mallinckrodt)
Gentisic acid ethanolamine	1	2%	M.V.I. 12® (Astra) 2%
Glutamate monosodium	2	0.1% w/v	Varivas® (Merck) 0.1% w/v
Formaldehyde sulfoxylate sodium	9	0.075-0.5% w/v	Terramycin Solution (Roerig) 0.5% w/v
Metabisulfite potassium	1	0.10%	Vasoxyl® (Glaxo-Wellcome) 0.10%
Metabisulfite sodium	29	0.02-1% w/v	Intropin® (DuPont) 1% w/v
Monothioglycerol (Thioglycerol)	6	0.1-1%	Terramycin Solution (Roerig) 1%
Propyl gallate	2	0.02%	Navane® (Roerig)
Sulfite sodium	7	0.05-0.2% w/v	Enion® (Ohmeda) 0.2% w/v
Thioglycolate sodium	1	0.66% w/v	Sus-Phrine® (Forest) 0.66% w/v

TABLE V
Antimicrobial Preservatives

Excipient	Frequency	Range	Example
Benzalkonium chloride	1	0.02% w/v	Celestone Soluspan® (Schering) 0.02% w/v
Benzethonium chloride	4	0.01%	Benadryl® (Parke-Davis) 0.01% w/v
Benzyl alcohol	74	0.75-5%	Dimenhydrinate® (Steris) 5%
Chlorobutanol	17	0.25-0.5%	Codine phosphate (Wyeth-Ayerst) 0.5%
m-Cresol	3	0.1-0.3%	Humatrope® (Lilly) 0.30%
Myristyl gamma-picolinium chloride	2	0.0195-0.169% w/v	Depo-Provera® (Upjohn) 0.169% w/v
Paraben methyl	50	0.05-0.18%	Inapsine® (Janssen) 0.18% w/v
Paraben propyl	40	0.01-0.1%	Xylocaine w/Epinephrine (Astra) 0.1% w/v
Phenol	48	0.2-0.5%	Calcimar® (Rhône Poulanc) 0.5% w/v
2-Phenoxyethanol	3	0.50%	Havrix® (SmithKline Beecham) 0.50% w/v
Phenyl mercuric nitrate	3	0.001%	Antivenin® (Wyeth-Ayerst) 0.001%
Thimerosal	46	0.003-0.01%	Atgam® (Upjohn) 0.01%

TABLE VI
Buffers and pH Adjusting Agents

Excipient	Example
Acetate	
Sodium	Miacalcin Injection® (Sandoz)
Acetic acid	Miacalcin Injection® (Sandoz)
Glacial acetic acid	Brevibloc Injection® (Ohmeda)
Ammonium	Bumex Injection® (Roche)
Ammonium hydroxide	Triostat Injection® (SmithKline Beecham)
Benzene sulfonic acid	Tracrium Injection® (Glaxo-Wellcome)
Benzoate Sodium/acid	Valium Injection® (Roche)
Bicarbonate Sodium	Cefotan Injection® (Zeneca)
Carbonate Sodium	HypoRho-D® (Bayer)
Citrate	
Acid	DTIC-Dome® (Bayer)
Sodium	Ceredase® (Genzyme)
Disodium	Cerezyme® (Genzyme)
Trisodium	Cerezyme® (Genzyme)
Diethanolamine	Bactrim IV® (Roche)
Glucono delta lactone	Quinidine® (Lilly)
Glycine	Hep-B Gammegee® (Merck)
Hydrochloric acid	Amicar® (Immunex)
Hydrogen bromide	Scopolamine (UDL)
Lactate acid/Sodium	Fentanyl citrate & Droperidol (Astra)
Lysine	Eminase Injection® (Roberts)
Maleic acid	Librium Injection® (Roche)
Methanesulfonic acid	DHE-45 Injection® (Sandoz)
Monoethanolamine	Terramycin Solution (Roerig)
Phosphate	
Acid (phosphoric)	Humegon® (Organon)
Monobasic potassium	Zantac Injection® (Glaxo-Wellcome)
Monobasic sodium*	Pregnyl® (Organon)
Dibasic sodium**	Prolastin® (Bayer)
Tribasic sodium	Synthroid® (Knoll)
Sodium hydroxide	Optiray® (Mallinckrodt)
Sulfuric acid	Nebcin® (Lilly)
Tartrate acid/sodium	Methergine Injection® (Sandoz)
Tromethamine	Optiray® (Mallinckrodt)

* Sodium biphosphate, Sodium dihydrogen phosphate or Na dihydrogen orthophosphate.

** Sodium phosphate, Disodium hydrogen phosphate.

(Tracrium Inj., Atracurium besylate) or mesylate (DHE 45 Injection, Dihydroergotamine mesylate) salts. Glucono delta lactone is used to adjust the pH of Quinidine gluconate (Lilly). Benzoate buffer, at a concentration of 5%, is used in Valium Injection. Citrates are common buffers that can have a dual role as chelating agents. Lysine and glycine are amino acids which function as buffers and stabilize protein and peptide formulations. These amino acids are also used as lyo-additives and may prevent cold denaturation. Lactate and tartrate are occasionally used as buffer systems.

Table VII lists additives which are used to modify osmolality, and as bulking or lyo-cryo protective agents. Dextrose and sodium chloride are used to adjust tonicity in the majority of formulations. Some amino acids, glycine, alanine, histidine, imidazole, arginine, asparagine, aspartic acid, are used as bulking agents for lyophilization and may serve as stabilizers for proteins or peptides and as buffers. Monosaccharides (dextrose, glucose, lactose), disaccharide (sucrose), polyhydric alcohols (inositol, mannitol, sorbitol), glycol (PEG 3350), Povidone (polyvinylpyrrolidone), and proteins (albumin, gelatin) are commonly used as lyo-additives.

TABLE VII
Bulking Agents, Protectants, and Tonicity Adjustors

Excipient	Example
Alanine	Thrombate III® (Bayer)
Albumin	Bioclate® (Arco)
Albumin human	Botox® (Allergan)
Amino acids	Havrix® (SmithKline Beecham)
L-Arginine	Activase® (Genentech)
Asparagine	Tice BCG® (Oganon)
L-Aspartic acid	Pepcid® (Merck)
Calcium chloride	Phenergan Injection® (Wyeth-Ayerst)
Citric acid	Sensorcaine-MPF® (Astra)
Dextrose	Betaseron® (Berlex)
Gelatin hydrolyzed	Acthar® (Rhône-Poulanc Rorer)
Glucose	Iveegan® (Immuno-US)
Glycerin	Tice BCG® (Oganon)
Glycine	Atgam Injection® (Upjohn)
Histidine	Antihemophilic Factor, human (Am. Red Cross)
Imidazole	Helixate® (Armour)
Inositol	OctreoScan® (Mallinckrodt)
Lactose	Caverject® (Upjohn)
Magnesium chloride	Terramycin Solution® (Roerig)
Magnesium sulfate	Tice BCG® (Oganon)
Mannitol	Elispar® (Merck)
Polyethylene glycol 3350	Bioclate® (Arco)
Polysorbate 80	Helixate® (Armour)
Potassium chloride	Varivax® (Merck)
Povidone	Alkeran® (Glaxo-Wellcome)
Sodium chloride	WinRho SD® (Univax)
Sodium succinate	Actimmune® (Genentech)
Sodium sulfate	Depo-Provera® (Upjohn)
Sorbitol	Panhematin® (Abbott)
Sucrose	Prolastin® (Bayer)

Special Additives

These additives have been included in pharmaceutical formulation to serve specific functions (Table VIII). Below is a summary of the special additives along with their intended use—

- (1) Calcium gluconate injection (American Regent) is a saturated solution of 10% w/v; calcium d-saccharate tetrahydrate 0.46% w/v is added to prevent crystallization during temperature fluctuations.
- (2) Cipro IV® (Ciprofloxacin, Bayer) contains lactic acid as a solubilizing agent for the antibiotic.
- (3) Premarin Injection® (Conjugated Estrogens, Wyeth-Ayerst Labs) is a lyophilized product that contains simethicone to prevent formation of foam during reconstitution.
- (4) Dexamethasone acetate (Dalalone DP, Forest, Decadron-LA, Merck, Dalalone DP Injection, UAD Labs) and Dexamethasone Na phosphate (Merck) are available as suspension or solution. These dexamethasone formulations contain creatine or creatinine as an additive.
- (5) Adriamycin RDF® (Doxorubicin hydrochloride, Pharmacia) contains methyl paraben, 0.2 mg/mL, to increase dissolution (10).
- (6) Ergotrate maleate (Ergonovine maleate, Lilly) contains 0.1% ethyl lactate as a solubilizing agent.
- (7) Estradurin Injection® (Polyestradiol phosphate, Wyeth-Ayerst Labs) uses Niacinamide (12.5 mg/ml)

TABLE VIII
Special Additives

Excipient	Example
Acetyl tryptophanate	Human Albumin (American Red Cross)
Aluminum hydroxide	Recombinant HB [®] (Merck)
Aluminum phosphate	Tetanus Toxoid Adsorbed [®] (Lederle)
Aluminum potassium sulfate	TD Adsorbed Adult [®] (Connaught)
E-Aminocaproic acid	Eminase [®] (Roberts)
Calcium D-saccharate	Calcium Gluconate (American Regent)
Caprylate sodium	Human Albumin (American Red Cross)
8-Chlorotheophylline	Dimenhydrinate (Steris)
Creatine	Dalalone DP [®] (Forest)
Creatinine	Hydrocortone Phosphate (Merck)
Diazotric acid	Conray (Mallinckrodt)
Gamma Cyclodextrin	Cardiotec (Squibb)
Ethyl lactate	Ergotrate maleate [®] (Lilly)
Ethylenediamine	Aminophylline [®] (Abbott)
L-Glutamate sodium	Kabikinase [®] (Pharmacia)
Iron ammonium citrate	Tice BCG [®] (Oganon)
Lactic acid	Cipro IV [®] (Bayer)
D,L-Lactic and Glycolic acid copolymer	Zoladex [®] (Zeneca)
Maltose	Gamimune [®] (Bayer)
Meglumine	Magnevist [®] (Berlex)
Niacinamide	Estradurin [®] (Wyeth-Ayerst)
Paraben methyl	Adriamycin RDF [®] (Pharmacia)
Protamine	Insulatard NPH [®] (Novo Nordisk)
Simethicone	Premarin Injection [®] (Wyeth-Ayerst)
Sodium saccharin	Compazine Injection [®] (Smith-Kline Beecham)
Tri-n-butyl phosphate	Venoglobulin [®] (Alpha Therapeutic)
von Willebrand factor	Bioclote [®] (Arco)
Zinc	Lente Insulin [®] (Novo Nordisk)

as a solubilizing agent. Hydetrastol[®] (Merck) also contains niacinamide.

- (8) Aluminum in the form of aluminum hydroxide, aluminum phosphate or aluminum potassium sulfate is used as adjuvant in various vaccine formulations to elicit an increased immunogenic response.
- (9) Zoladex[®] (Goserelin acetate, Zeneca) is administered subcutaneously as microspheres. These spheres are made of D,L-lactic and glycolic acid copolymer. Lupron Depot Injection[®] (TAP) are lyophilized microspheres of gelatin and glycolic-lactic acid for intramuscular injection.
- (10) Gamma cyclodextrin is used as a stabilizer in Cardiotec[®] at a concentration of 50 mg/mL.
- (11) Sodium caprylate (sodium octoate) has antifungal properties, but it is also used to improve the stability of albumin solution against effects of heat. Albumin solution can be heat pasteurized by heating at 60°C for 10 hours in the presence of sodium caprylate. Acetyl tryptophanate sodium is also added to albumin formulations.
- (12) Meglumine (N-methylglucamine) is used as an ex-

TABLE IX
List of Excipient from 1996 FDA 'Inactive Ingredient Guide'

Ammonium sulfate	Pentetate (DTPA) calcium trisodium
Benzyl chloride	Poloxamer 165
Butyl paraben	PEG 4000
Calcium chloride sodium	PEG 600
Calcitriol calcium	Polyglactin
Castor oil	Poly lactide
Cellulose (microcrystalline)	Polyoxyethylene fatty acid esters
Cholesterol	Polyoxyethylene sorbitan monosterate
Deoxycholic acid	Polyoxyl 35 Castor oil
Diazotric acid	Polysorbate 40
Dicyclohexyl carbodiimide	Polysorbate 85
Diethyl amine	Potassium hydroxide
Dimyristoyl lecithin	Potassium phosphate, dibasic
Dimyristoyl phosphatidyl-glycerol	Sodium bisulfate
Disofenin	Sodium chlorate
Docosate sodium	Sodium hypochloride
Edamine	Sodium iodide
Exametazime	Sodium pyrophosphate
Gluceptate sodium	Sodium thiosulfate, anhydrous
Gluceptate calcium	Sodium trimetaphosphate
Glucuronic acid	Sorbitan monopalmitate
Guanidine HCl	Stannous chloride
Iofetamine HCl	Stannous fluoride
Lactobionic acid	Stannous tartrate
Lecithin hydrogenated soy	Starch
Lidofenin	Succimer
Medrofenin	Succinic acid
Medronate disodium	Sulfurous acid
Medronic acid	Tetrakis (1-isocyno-2-methoxy-2-methyl-propante) copper (I) Te
Methyl boronic acid	Thiazoximic acid
Methyl cellulose	Trithiazoximic acid
Methylene blue	Urea
N-(carbamoyl-methoxy polyethylene-glycol 2000)-1,2-distearoyl	Zinc acetate
N-2-hydroxyethyl piperazine N'-2' ethane sulphonate acid	Zinc chloride
Nioxime	Zinc oxide
Nitric acid	2-ethyl hexanoic acid
Oxyquinoline	PEG vegetable oil

ipient and to form in-situ salt. For example, diazotric acid, an X-ray contrast agent, is more stable when autoclaved as meglumine salt than as sodium salt (11). Meglumine is also added to Magnevist[®], a magnetic resonance contrast agent, formulation.

- (13) Surprisingly, sodium saccharine is used in Stelazine[®] and Compazine[®] formulations; our guess is that it serves as a stabilizer and tonicity adjuster.
- (14) Tri-n-butyl phosphate is present as an excipient in human immune globulin solution (Venoglobulin[®]). Its exact function in the formulation is not known, but it may serve as a scavenging agent.
- (15) von Willebrand factor is used to stabilize recombinant antihemophilic factor (Bioclote[®]).
- (16) Maltose serves as a tonicity adjuster and stabilizer in immune globulin formulation (Gamimune N[®]).
- (17) Epsilon amino caproic acid (6-amino hexanoic acid) is used as a stabilizer in anistreplase (Eminase injection[®]).
- (18) Zinc and protamine have been added to insulin to form complexes and control the duration of action.

Recently, FDA has published 'Inactive Ingredient Guide' which lists all the excipients in alphabetical order. Each ingredient is followed by the route of administration (for example, iv, oral) and, in some cases, the range of concentration used in the approved drug product. However, this list does not provide the name of commercial product(s) corresponding to each excipient. Table IX is a summary of all the excipients which are included in the 'Inactive Ingredient Guide,' but do not appear in PDR or Handbook on Injectable Drugs.

References

1. Y. J. Wang and R. R. Kowal, "Review of excipients and pH's for parenteral products used in the United States," *J. Parenter. Sci. Technol.*, 34(6), 432 (1980).
2. Physicians' Desk Reference, ed. 48, 1994.
3. Physicians' Desk Reference, ed. 50, 1996.
4. L. A. Trissel, "Handbook on Injectable Drugs," ed. 8, American Society of Hospital Pharmacists, Inc., 1994.
5. K. W. Reed and S. Yalkowsky, "Lysis of human red blood cells in the presence of various cosolvents," *J. Parenter. Sci. Technol.*, 39(2), 64 (1985).
6. G. A. Bazeau and Ho-Leung Fung, "Use of an in-vitro model for the assessment of muscle damage from intramuscular injections: In-vitro-in-vivo correlation and predictability with mixed solvent systems," *Pharm. Res.*, 6(9), 766 (1989).
7. J. W. Munson, A. Hussain, and R. Bilous, "Precautionary note for use of bisulfite in pharmaceutical formulation," *J. Pharm. Sci.*, 66(12), 1775 (1977).
8. L. C. Schroeter, "Sulfurous acid salts as pharmaceutical antioxidants," *J. Pharm. Sci.*, 50(11), 891 (1961).
9. R. Dabbah, "The use of preservatives in compendial articles," *Pharmaceutical Forum*, 22(4), 2696 (1996).
10. T. J. Baumann, M. A. Smythe, K. Kaufmann, Z. Miloboszewski, J. O'Malley, and R. R. Fudge, "Dissolution times of adriamycin and adriamycin RDR," *Am. J. Hosp. Pharm.*, 45, 1667 (1988).
11. Y. J. Wang, T. C. Dahl, G. D. Leesman, and D. C. Monkhouse, "Optimization of autoclave cycles and selection of formulation for parenteral product, Part II: Effect of counter-ion on pH and stability of diatrizoic acid at autoclave temperatures," *J. Parenter. Sci. Technol.*, 38(2), 72 (1984).
12. 'Inactive Ingredient Guide,' Division of Drug Information Resources, FDA, CDER, January 1996.

Pharmacokinetics of probenecid in sheep

V. H. GUERRINI, L. J. FILIPPICH,* P. B. ENGLISH,* JENNY SCHNEIDER,
G. R. CAO* & D. W. A. BOURNE

Departments of Pharmacy and *Veterinary Medicine, University of Queensland, St Lucia,
Australia

Guerrini, V.H., Filippich, L.J., English, P.B., Schneider, J., Cao, G.R. & Bourne, D.W.A. Pharmacokinetics of probenecid in sheep. *J. vet. Pharmacol. Therap.* 8, 128–135.

Six Merino ewes were given 1 g (27 g/kg) probenecid by the intravenous (i.v.), intramuscular (i.m.) and subcutaneous (s.c.) routes. After i.v. injection, the biological half-life was 1.55 h and apparent volume of distribution at the steady state (Vd_{ss}) 0.18 l/kg. Body clearance (Cl_B) and renal clearance (Cl_R) were 0.12 l/h/kg and 0.03 l/h/kg, respectively. Approximately 28% of unchanged probenecid was excreted in urine. Plasma probenecid concentrations after i.v., i.m. and s.c. injections were 133, 37, and 31 $\mu\text{g/ml}$, respectively, at 15 min; 76, 36, and 34 $\mu\text{g/ml}$ at 1 h; and 43, 23 and 34 $\mu\text{g/ml}$ at 2 h. The average bioavailability of probenecid given by i.m. and s.c. injection was 46% and 34%, respectively. However, after 2 h, probenecid plasma concentrations remained higher when it was given subcutaneously than when it was given intramuscularly.

Urine output was correlated positively ($P < 0.05$) with k_{el} and Cl_B . Urine pH increased significantly ($P < 0.01$) for the first 2 h, and then steadily declined over the subsequent 6 h. The results suggested that probenecid in sheep was rapidly eliminated because it was rapidly excreted in the normal but alkaline urine. Subcutaneous administration of probenecid in animals may be a useful alternative to oral or i.v. administration.

Dr V. H. Guerrini, Department of Pharmacy, University of Queensland, St Lucia, Queensland 4067, Australia.

INTRODUCTION

Probenecid, also used as a uricosuric agent in man, has mainly been used to prolong the biological half-life and to alter the distribution characteristics of some penicillins and third-generation cephalosporins (Gibaldi & Scharz, 1968; Welling *et al.*, 1979). The third-generation cephalosporins have a relatively short plasma half-life, and this has impaired their use in veterinary medicine (Guerrini *et al.*, 1983; Kalager *et al.*, 1982). The pharmacokinetics of probenecid must be well known if this drug is to be used to improve the disposition of the third-generation cephalo-

sporins. Since the oral administration of probenecid to ruminants is not practical, the pharmacokinetics of this drug had to be determined after parenteral administration.

Urinary pH and urine flow rate are known to affect the elimination of probenecid, and these parameters should be measured in conjunction with the plasma concentration of this drug (Melethil & Conway, 1976; Dayton *et al.*, 1963). The purpose of this study was to determine the pharmacokinetics of probenecid after i.v., i.m. and s.c. administration to sheep. The effects of urine flow rate and urinary pH on the pharmacokinetics of probenecid were also investigated.

MATERIALS AND METHODS

Animals

Six adult Merino ewes (average body weight 36.8 ± 4.8 kg) were acclimatized for two weeks before the study. Each sheep was kept in an individual metabolism cage with water and lucerne chaff freely available (ambient temperature $25 \pm 2^\circ\text{C}$; ambient humidity $55 \pm 10\%$). A jugular vein was cannulated and the urinary bladder was catheterized for the collection of blood and urine samples, respectively.

Drug administration

Each sheep was given 1 g of probenecid (Merck Sharp and Dohme, Granville South, NSW) as a solution in phosphate buffer (pH 7.4) (Martindale, 1982). The drug was given by rapid i.v., deep i.m. (gluteal muscle) or s.c. (pre-scapular region) injection in three separate experiments each separated by a 2-week rest period.

Specimen collection

Blood samples were collected from the jugular vein via an indwelling cannula. The patency of cannula was maintained by injecting a 1/1000 sodium heparin solution into cannulae before each sampling. The initial portion of each blood sample was discarded. Urine was collected in plastic vials via indwelling catheters using a method described previously (Guerrini *et al.*, 1983). After the drug was given by i.v., i.m. and s.c. administration, blood samples (3–5 ml) were collected at 5, 10, 15, 20, 40 minutes and 1, 2, 3, 4, 5, 6, 8, 12, 14, and 18 h. Urine (total volume) was collected at 0, 30 min and 1, 1.5, 2, 3, 4, 5, 6, 8, 10, 12, 14 and 16 h after dosing.

Analytical methods

Plasma and urine concentrations of probenecid were measured by high-pressure liquid chromatography (HPLC) with u.v. detection (Model 440 absorbance detector, Waters

Associates, Milford, MA). Plasma and urine samples (250 μl each) were vortexed with acetonitrile (250 and 1500 μl , respectively) and centrifuged for 3 minutes to remove protein material. Twenty μl of the supernatant was then injected onto the HPLC column (RP-8, 10 μm , Brownlee Lab., Santa Clara, CA). The mobile phase (65% sodium dihydrogen phosphate buffer [0.02 M, pH 5] 30% methanol and 5% acetonitrile) was pumped (Model M-45, Waters Associates) through the column at a flow rate of 1.5 ml/min. Probenecid was quantitated at 254 nm with a retention time of 3 min. Peak heights were used to calculate plasma probenecid concentrations and all determinations were performed in duplicate. Standard curves were prepared in the range 0.2–400 $\mu\text{g/ml}$ for plasma, and 2–2000 $\mu\text{g/ml}$ for urine.

Renal function tests

The glomerular filtration rate and effective renal plasma flow were estimated using ^{125}I -labeled sodium iothalamate and sodium *p*-aminohippurate (PAH) as previously described (Filippich, 1982; Guerrini *et al.*, 1983).

Statistical analysis

The individual animal plasma and urine data, and the averaged plasma and urine data, were fitted with appropriate pharmacokinetic models using the digital computer program NONLIN (Metzler *et al.*, 1974) modified to operate on a minicomputer (Bourne & Wright, 1981). Mean data points with standard deviation values greater than the mean values were not included for analysis. These points were those measured 2.5 h after i.v., 4.0 h after i.m. and 8.0 h after s.c. administration. Initial estimates obtained for the pharmacokinetic parameters, apparent volume of the central compartment (V_C), elimination rate constant (k_{e1}), urinary excretion rate constant (k_e), rate constant for diffusion into tissue (k_{12}) and out of tissue (k_{21}) were subjected to a weighted iterative least-squares analysis using NONLIN. In the case of i.m. and s.c. data, V_C/F (where F was the bioavailability expressed as a fraction of 1) was

used as an initial parameter. The harmonic mean plasma half-life value was calculated as $0.693/\beta$. Renal clearance (Cl_R) was calculated as $k_e \cdot V_C$, and body clearance (Cl_B) as $k_{e1} \cdot V_C$. The apparent volume of distribution at the steady state ($V_{d,ss}$) was calculated as $(k_{12} + k_{21}) \cdot V_C$ divided by k_{21} . The area under the plasma concentration vs time curve (AUC) was calculated by the trapezoidal rule. Linear regression was used to estimate the degree of correlation between urine volumes voided and k_e , terminal plasma half-life, or Cl_B . A paired *t*-test was used to test the significance of the difference between mean pH values and mean urine flow rate values.

RESULTS

Averaged plasma concentrations of probenecid and averaged cumulative amounts of probenecid excreted into urine measured in six sheep given probenecid by the i.v. route are shown in Fig. 1. Similar data determined in the same sheep given probenecid by the i.m. and s.c. routes are presented in Figs 2 and 3, respectively.

After i.v. injection of the drug, plasma probenecid concentrations were 133 ± 18 $\mu\text{g/ml}$ at 15 minutes, 76 ± 26 $\mu\text{g/ml}$ at 1 h, and

42.6 ± 32.8 $\mu\text{g/ml}$ at 2 h. The plasma and urine data were subsequently fitted by a two-compartment pharmacokinetic model and the pharmacokinetic parameters obtained with NONLIN are shown in Table I. Urine volume voided over 12 h correlated significantly ($P < 0.05$) with the excretion rate constant, the renal clearance, and the biological half-life (negative correlation) of probenecid.

After i.m. administration, the average plasma probenecid concentration was 37.4 ± 26.8 $\mu\text{g/ml}$ at 15 minutes, 35.5 ± 29.0 $\mu\text{g/ml}$ at 1 h, and 22.7 ± 18.6 $\mu\text{g/ml}$ at 2 h. The average AUC value 118 ± 69 $\mu\text{g}\cdot\text{h/ml}$. The individual and average data were fitted by a two-compartment pharmacokinetic model (NONLIN) with an added first-order absorption step (k_a) and are shown in Table II. The values of the distribution rate constants obtained in the analysis of the i.v. data were used in the i.m. analysis as fixed constants. The bioavailability of the i.m. dosage form was calculated from the ratio of V_C (i.v.) to V_C/F (i.m.) and the ratio of the AUC values corrected for k_{e1} . Calculated by these two methods, the bioavailability was $48 \pm 36\%$ and $43 \pm 33\%$, respectively, giving an average value of 46%. The average total amount of unchanged probenecid found in the urine was $39.5 \pm 4.1\%$ of the dose. The




FIG. 1. Linear plot of average plasma concentration (●) and cumulative amount of drug excreted into urine (▲) vs time in six sheep after i.v. administration of 1 g (27 mg/kg) of probenecid. The points are the experimentally determined average values (with ± 1 SE shown as vertical bars) and the solid line was calculated using the 'best fit' values (as calculated with NONLIN).




FIG. 2. Linear plot of average plasma concentration (●) and cumulative amount of drug excreted into urine (▲) vs time in six sheep after i.m. administration of 1 g (27 mg/kg) of probenecid. The points are the experimentally determined average values (with \pm SE shown as vertical bars) and the solid line was calculated using the 'best fit' values (as calculated with NONLIN). Standard errors for values taken at 0.08, 0.17, 0.25 and 0.33 h were 13.1, 21.2, 26.7 and 30.1, respectively.




FIG. 3. Linear plot of average plasma concentration (●) and cumulative amount of drug excreted into urine (▲) vs time in six sheep after s.c. administration of 1 g (27 mg/kg) of probenecid. The points are the experimentally determined average values (with \pm 1 SE shown as vertical bars) and the solid line was calculated using the 'best fit' values (as calculated with NONLIN).

averaged parameter values were used to calculate the solid lines in Fig. 2.

After s.c. administration, the average plasma probenecid concentration was $31.2 \pm 19.7 \mu\text{g/ml}$ at 15 minutes, $34.3 \pm 11.8 \mu\text{g/ml}$ at 1 h, and $33.7 \pm 13.7 \mu\text{g/ml}$ at 2 h. Plasma probenecid concentration remained higher

($P < 0.05$) after 2 h by s.c. injection than by i.m. administration. The individual and average plasma and urine data were fitted by a two-compartment pharmacokinetic model with an added first-order absorption step as described for the i.m. data. The pharmacokinetic parameters obtained by this method are

TABLE I. Pharmacokinetic parameters measured after i.v. administration of 1 g probenecid to each of six sheep

Values	Sheep						$\bar{x} \pm \text{SD}$ (37 \pm 5)	Average
	No. 1 (37 kg)	No. 2 (32 kg)	No. 3 (41 kg)	No. 4 (34 kg)	No. 5 (33 kg)	No. 6 (44 kg)		
k_c (h^{-1})	0.072	0.385	0.399	0.295	0.060	0.485	0.373 \pm 0.180	0.215
k_{misc} (h^{-1})	0.479	0.958	1.240	1.025	0.343	0.442	0.748 \pm 0.372	0.586
k_{cl} (h^{-1})	0.55	1.34	1.64	1.32	0.40	0.93	1.03 \pm 0.49	0.80
k_{21} (h^{-1})	3.95	1.79	5.94	0.84	2.57	2.41	2.92 \pm 1.80	1.57
k_{21} (h^{-1})	10.2	3.07	9.10	6.51	4.34	5.26	6.42 \pm 2.78	3.26
V_c (l/kg)	0.10	0.15	0.08	0.15	0.15	0.10	0.12 \pm 0.03	0.13
V_{ss} (l/kg)	0.18	0.23	0.14	0.17	0.23	0.14	0.18 \pm 0.04	0.19
$t_{1/2}^*$ (h)	1.76	0.22	0.73	0.61	2.80	1.14	1.55 [†]	1.36
AUC ($\mu\text{g}\cdot\text{h}/\text{ml}$)	437	141	178	148	466	221	265 \pm 147	241
Cl_B (l/h/kg)	0.06	0.20	0.13	0.20	0.06	0.09	0.12 \pm 0.006	0.10
Cl_R (l/h/kg)	0.01	0.06	0.04	0.06	0.02	0.03	0.03 \pm 0.002	0.03
GFR (ml/min/kg)	1.97	2.89	2.20	1.74	2.21	2.10	2.19 \pm 0.39	
RPF (ml/min/kg)	12.4	17.4	13.4	10.5	13.1	11.0	13.0 \pm 2.84	
Unchanged drug in urine (% of dose)	16	31	28	24	16	54	28 \pm 14	
Urine voided at 12 h (ml)	106	308	428	627	136	584	348 \pm 100	

*Terminal half-life.

[†]Harmonic mean calculated as $0.693/\beta$.

shown in Table III. The bioavailability of the s.c. dosage form was $35 \pm 12\%$ and $32 \pm 14\%$, respectively (calculated by the two methods described above), giving an average value of 34% of the administered dose. The average total amount of unchanged drug found in urine was $36.3 \pm 13.6\%$ of the dose. The average values were used to calculate the solid lines in Fig. 3.

Urine flow rate after i.v., i.m. and s.c. administration of probenecid is shown in Table IV. Average urinary pH values measured after IM administration are also shown on Table 4. The pH values increased significantly from 8.08 ± 0.37 at 0 minutes to 8.53 ± 0.24 ($P < 0.01$) at 2 h, and then steadily declined to 7.51 ± 1.19 (not significantly different to the zero time value, $P > 0.05$) over the subsequent 6 h. Increased urine output was correlated ($P < 0.05$) with increased k_e ($r = 0.55$ for 18 pairs) and increased Cl_B ($r = 0.46$ for 18 pairs). The glomerular

filtration rate (GFR) and renal plasma flow rate (RPF) for individual sheep are shown in Table I. The average values for GFR and RPF were 2.19 ± 0.39 and 13 ± 2.5 ml/min/kg body weight, respectively.

DISCUSSION

The average plasma half-life or body clearance for probenecid in sheep after i.v., i.m. and s.c. administration in the present study was shorter than that found in man and in other mammalian species. Cunningham *et al.* (1981) reported that the plasma half-life of probenecid in man after doses of 0.5–1.0 g intravenously was 2–6 h, whereas, in the present study, the average biological half-life in sheep was 0.6–2.8 h. The body clearance for probenecid in man (70 kg body weight) was 0.02 l/h/kg (Perel *et al.*, 1971), compared with 0.12 l/h/kg in sheep. The normally

TABLE II. Pharmacokinetic parameters measured after i.m. administration of 1 g probenecid to each of six sheep

Values	Sheep						$\bar{x} \pm SD$	Average
	No. 1	No. 2	No. 3	No. 4	No. 5	No. 6		
k_c (h^{-1})	0.119	0.715	0.151	0.346	0.364	0.396	0.349 ± 0.214	0.282
k_{misc} (h^{-1})	0.150	1.034	0.315	0.607	0.473	0.546	0.519 ± 0.298	0.445
k_{cl} (h^{-1})	0.27	1.74	0.47	0.95	0.84	0.94	0.87 ± 0.51	0.73
Cl_B (l/h/kg)	0.43	0.95	0.77	0.30	0.14	0.13	0.37 ± 0.31	0.25
k_a (h^{-1})	8.75	1.14	10.9	6.51	2.64	2.70	5.45 ± 3.90	3.74
Bioava* (%)	8.3	37.9	16.2	44.2	99.1	82.2	48.0 ± 36.0	42.7
Bioava† (%)	6.5	27.6	14.0	46.8	85.5	77.5	43.0 ± 32.9	37.6
V_C/F (l/kg)	1.61	0.54	0.57	0.31	0.17	0.13	0.56 ± 0.55	0.34
AUC ($\mu g \cdot h/ml$)	73.7	41.4	101.0	90.6	222.0	179.0	118.0 ± 68.5	113.0
Cl_R (l/h)	7.1	12.5	3.5	3.7	2.0	2.3	7.2 ± 4.4	3.5
Drug in urine (% of dose)	41	41	33	37	44	41	40 ± 4	
Urine voided at 12 h (ml)	160	632	819	547	582	654	565 ± 220	

*Bioavailability calculated as Auc (i.m.)/AUC (i.v.) $\times k_{cl}$ (i.m.)/ k_{cl} (i.v.).

†Bioavailability calculated as V_C (i.v.)/ V_C (i.m.).

TABLE III. Pharmacokinetic parameters measured after s.c. administration of 1 g of probenecid to each of six sheep

Values	Sheep						$\bar{x} \pm SD$	Average
	No. 1	No. 2	No. 3	No. 4	No. 5	No. 6		
k_c (h^{-1})	0.191	0.116	0.125	0.020	0.145	0.088	0.114 ± 0.057	0.097
k_{misc} (h^{-1})	0.160	0.314	0.118	0.114	0.213	0.122	0.173 ± 0.078	0.127
k_{cl} (h^{-1})	0.35	0.43	0.24	0.13	0.36	0.21	0.29 ± 0.11	0.22
Cl_B (l/h/kg)	0.18	0.12	0.11	0.05	0.13	0.09	0.11 ± 0.04	0.09
k_a (h^{-1})	2.56	9.43	4.87	1.61	2.48	2.29	3.87 ± 2.94	3.87
Bioava* (%)	21.5	51.9	26.7	38.6	43.6	28.8	35.2 ± 11.5	28.8
Bioava† (%)	20.6	54.0	17.6	38.0	40.1	23.0	32.2 ± 14.2	30.7
AUC ($\mu g \cdot h/min$)	148	229	322	561	228	282	295 ± 143	249
V_C/F (l/kg)	0.51	0.28	0.46	0.39	0.36	0.44	0.41 ± 0.08	0.42
Cl_R (l/h/kg)	0.33	0.29	0.17	0.05	0.35	0.15	0.22 ± 0.12	0.21
Drug in urine (%) of dose)	52	28	46	14	40	38	36 ± 14	
Urine voided at 12 h (ml)	185	201	555	279	286	394	316 ± 138	

*Bioavailability calculated as $[AUC$ (s.c./AUC i.v.) $\times [k_{cl}$ (s.c.)/ k_{cl} (s.c.)/ k_{cl} (i.v.)].

†Bioavailability calculated as V_C (i.v.)/(V_C/F) (s.c.).

TABLE IV. Average (\pm SD) urinary pH and urine volume after i.v., i.m. and probenecid in six Merino Ewes

Time (h)	pH* (units)	Urine volume (ml/min)	Urine volume (ml/min)	Urine volume (ml/min)
		i.v.	i.m.	s.c.
0	8.08 (0.37)			
0.5	8.45 (0.22)	0.62 (0.37)	0.93 (0.48)	0.61 (0.54)
1.0	8.50 (0.24)	0.58 (0.37)	1.20 (1.25)	0.71 (0.52)
2.0	8.53 (0.24)	0.38† (0.18)	0.80 (0.71)	1.08 (1.10)
3.0	8.52 (0.28)	0.53 (0.47)	0.64 (0.40)	0.47 (0.27)
4.0	8.48 (0.30)	0.46 (0.27)	0.41‡ (0.22)	0.32 (0.11)
5.0	8.06 (0.37)	0.40 (0.28)	0.34‡ (0.16)	0.33 (0.12)
6.0	7.76 (0.82)	0.65 (0.71)	0.53 (0.36)	0.26 (0.09)
8.0	7.51 (1.19)	0.35 (0.24)	0.26† (0.21)	0.32 (0.11)

*pH measured during i.m. study.

†Significantly different from the 0.5 h value ($P < 0.05$).

‡Significantly different from the 0.5 h value ($P < 0.01$).

higher urine pH value found in sheep should increase the ionized fraction of probenecid in the renal tubules, thereby decreasing the drug's tubular reabsorption and enhancing its renal excretion (Goodman & Gilman, 1980). The Cl_R value for probenecid in man (based on 70 kg body weight) was approximately 0.001 l/h/kg (Dayton *et al.*, 1963; Perel *et al.*, 1971), whereas in the present experiment the Cl_R value was 0.03 l/h/kg. The higher proportion of the dose excreted as unchanged drug in sheep (25–40%) compared with the value found in man (5–11%) further supports the suggestion that high urinary pH in sheep retards the reabsorption of the drug from the kidneys.

In sheep with higher urine output over 12 h, the biological half-life of probenecid was shorter, and the rate constant for urinary excretion (k_e) was larger than in sheep with slower urinary flow rates. This confirmed previous findings in man and in the dog, and

underlines the need to measure urinary pH, urine volume and urinary drug concentrations in pharmacokinetic studies (Cunningham *et al.*, 1981; Goodman & Gilman, 1980).

The present pharmacokinetic analysis indicated that the rate of diffusion of probenecid between the central and peripheral compartment was rapid. This rapid diffusion of probenecid is probably due to its high lipophilicity (Cunningham *et al.*, 1981). The values for V_C and Vd_{ss} and the ratio of k_{12} to k_{21} also suggested that there was extravascular distribution of probenecid in sheep. The Vd_{ss} value found for probenecid in man (0.16 l/kg) (Dayton *et al.*, 1963) compared well with the values found in sheep (0.18 l/kg).

Previous experiments carried out in man and in the dog indicated that probenecid was completely absorbed when given by the oral route (Cunningham *et al.*, 1981). In the present study, probenecid given i.m. or s.c. was incompletely absorbed with only one-half

to one-third of the administered dose reaching the circulation intact. However after 2 h, the s.c. injection provided higher and more prolonged plasma probenecid concentration than did the i.m. administration. Some of the drug may have precipitated at the site of injection and continued to be released from the site of injection after blood collection was terminated. There was no visible discomfort or damage done to the sheep. The subcutaneous administration of 1 g of probenecid in sheep provided similar plasma concentrations to those found in man after oral administration. Subcutaneous administration of drugs avoids muscle damage and is generally less painful; thus, this route is a useful alternative for the administration of probenecid in animals. Precipitated drug could result in tissue residues if the drug is given intramuscularly.

In conclusion, the present results suggest that one of the reasons why probenecid has a short biological half-life in the ruminant is because the urinary pH is normally high. Urine output also appeared to influence the renal elimination of probenecid. The s.c. administration of probenecid in animals is preferred because muscle damage is avoided and it provided useful plasma concentrations.

ACKNOWLEDGMENTS

This project was partially supported by the Australian Meat Research Committee. V. H. Guerrini is the recipient of a Australian Meat Research Committee pre-doctoral scholarship.

REFERENCES

- Bourne, D.W.A. & Wright, A.W. (1981) Modification of the digital computer program NONLIN which allows it to run on the PDP-11 mini-computer. *Australian Journal of Pharmaceutical Sciences*, **10**, 23-24.
- Cunningham, R.F., Israili, Z.H. & Dayton, P.G. (1981) Clinical pharmacokinetics of probenecid. *Clinical Pharmacokinetics*, **6**, 135-151.
- Dayton, P.G., Yu, T.F., Chen, W., Berger, L., West, L.A. & Gutman, A.B. (1963) The physiological disposition of probenecid including renal clearance, in man, studied by an improved method for its estimation in biological material. *Journal of Pharmacology and Experimental Therapeutics*, **140**, 278-285.
- Filippich, L.J. (1982) An evaluation of renal function tests in normal sheep and sheep with reductions in functional renal mass. *Ph.D. Thesis*. University of Queensland, Australia.
- Gibaldi, M. & Schwartz, A. (1968) Apparent effect of probenecid on the distribution of penicillins in man. *Clinical Pharmacology and Therapeutics*, **9**, 345-348.
- Goodman, A.G. & Gilman, (1980) *The Pharmacological Basis of Therapeutics*, 6th edn. Macmillan, New York.
- Guerrini, V.H., Filippich, L.J., English, P.B. & Bourne, D.W.A. (1983) Pharmacokinetics of cefotaxime in sheep. *American Journal of Veterinary Research*, **44**, 1488-1491.
- Kalager, T., Digranes, A., Bakke, K., Hellum, K.B., Bergan, T. & Solberg, C.O. (1982) Cefotaxime in serious infections — a clinical and pharmacokinetic study. *Journal of Antimicrobial Chemotherapy*, **9**, 157-163.
- Martindale, W. (1982) *The Extra Pharmacopoeia*, Ed. Reynolds, E.F. & Prasad, A.B., 28th edn, p. 1118. The Pharmaceutical Press, London.
- Melethil, S. & Conway, W.D. (1976) Urinary excretion of probenecid and its metabolites in humans as a function of dose. *Journal of Pharmaceutical Sciences*, **65**, 861-865.
- Metzler, C.M., Elfring, G.L. & McEwen, A.J. (1974) A package of computer programs for pharmacokinetic modeling. *Biometrics*, **30**, 562-563.
- Perel, J.M., Dayton, P.G., Fu, T.F. & Gutman, A.B. (1971) Studies of the renal excretion of probenecid acylglucuronide in man. *European Journal of Clinical Pharmacology*, **3**, 106-112.
- Welling, P.G., Dean, S., Selen, A., Kendall, M.J. & Wise R. (1979) Probenecid: an unexplained effect on cephalosporin pharmacology. *British Journal of Clinical Pharmacology*, **8**, 491-495.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 January 2003 (23.01.2003)

PCT

(10) International Publication Number
WO 03/006064 A1

- (51) International Patent Classification⁷: A61K 47/14, 47/44, 47/10, 31/565
- (21) International Application Number: PCT/GB02/03092
- (22) International Filing Date: 3 July 2002 (03.07.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0116619.8 7 July 2001 (07.07.2001) GB
60/315,711 30 August 2001 (30.08.2001) US
- (71) Applicant (for all designated States except MG, US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Sodertalje (SE).
- (71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CAHILL, Julie, Kay** [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). **GELLERT, Paul, Richard** [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). **IRVING, Alan, Marshall** [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
- (74) Agents: **GILES, Allen, Frank** et al.; AstraZeneca, Global Intellectual Property, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/006064 A1

(54) Title: PHARMACEUTICAL FORMULATION FOR THE INTRAMUSCULAR ADMINISTRATION OF FULVESTRANT

(57) Abstract: The invention relates to a sustained release pharmaceutical formulation adapted for administration by injection containing the compound fulvestrant, 7 a-[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 B-diol, at concentration of at least 100mg/ml in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

PHARMACEUTICAL FORMULATION FOR THE INTRAMUSCULAR ADMINISTRATION OF FULVESTRANT

The invention relates to a sustained release pharmaceutical formulation adapted for administration by injection containing the compound fulvestrant, 7α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, at concentration of at least 100mg/ml in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

An alternative approach to oestrogen withdrawal is to antagonise oestrogens with antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Furr and Jordan, *Pharmacology & Therapeutics*, 25:127-206, 1984; May and Westley, *J Biol Chem* 262:15894-15899, 1987).

The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

Steroidal analogues of oestradiol, with an alkylsulphinyl side chain in the 7α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989). One of these, 7α -[9-(4,4,5,5,5-pentafluoropentyl sulphanyl)nonyl]oestra-1,3,5-(10)triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with

7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphinyl)nonyl]oestra-1,3-5(10)-triene-3,17 β -diol, 5 or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is 10 more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may 15 offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If 20 also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier.

25 European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also 30 disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a

pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml⁻¹ (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present.

In US 5,183,814 Example 3 an oil based injection formulation of fulvestrant is described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

The Table below shows the solubility of fulvestrant in a number of different solvents.

SOLUBILITY OF FULVESTRANT

20

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79

- 4 -

Ethanol	>200
Benzyl Alcohol	>200

As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major
5 constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

Our earlier application PCT/GB01/00049, WO 01/51056, describes certain fulvestrant formulations at a most preferred concentration of 50mg/ml. This application disclosed one formulation with a solubility up to 102 mg/ml – see the last formulation in Table 3 thereof
10 with 15 % weight of ethanol per volume of formulation, 15 % weight of benzyl alcohol per volume of formulation, 15 % weight of benzyl benzoate per volume of formulation in a ricinoleate vehicle. However there is a need for further formulations of fulvestrant that contain high concentrations of fulvestrant to facilitate administration thereof at higher doses or less frequent intervals.

15 According to another aspect of the invention there is provided a pharmaceutical formulation adapted for intramuscular injection comprising 100 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of
20 formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle and provided that the following formulation is excluded: fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per
25 volume of formulation vehicle.

A preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 105 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a
30 pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation comprises at least 5 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 110 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 115 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 120 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 130 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 140 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 12.5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the

formulation vehicle comprises at least 10 % weight of ethanol per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 150 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 17.5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 10 % weight of ethanol per volume of formulation vehicle.

Another aspect of the invention provides any of the formulations described herein stated as having any minimum ethanol content removed. For example, the formulation described in the paragraph immediately above becomes: a pharmaceutical formulation adapted for intramuscular injection is one comprising 150 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 17.5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle.

According to another aspect of the invention there is provided a pharmaceutical formulation having a solubility for fulvestrant of at least Y mg/ml adapted for intramuscular injection comprising;

100 mg/ml or more of fulvestrant;

5% (w/v) or more castor oil per volume of formulation vehicle;

and at least the following amounts (% weight/volume of formulation vehicle) of ethanol (ETOH), benzyl alcohol (BA), benzyl benzoate (BB) determined by the algorithm:

$$Y = -29.77 + 5.44 \times \text{ETOH} + 2.38 \times \text{BA} + 1.57 \times \text{BB}$$

wherein x is at least 100, ETOH is at least 5, BA is at least 5 and BB is at least 5.

A preferred pharmaceutical formulation is one wherein Y is selected from the group consisting of 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 170, 180, 190, and 200.

A more preferred pharmaceutical formulation is one wherein Y is selected from the group consisting of 120, 125, 130, 135, 140, 145, 150, 155, 160, 170, 180, 190, and 200.

A more preferred pharmaceutical formulation is one wherein Y is selected from the group consisting of 150, 155, 160, 170, 180, 190 and 200.

A more preferred pharmaceutical formulation is one wherein Y is selected from 150, 155, 160, 170, 180, 190 and 200 and the formulation comprises at least 150mg/ml of fulvestrant.

A more preferred pharmaceutical formulation is one wherein Y is 200 and the
5 formulation comprises at least 200mg/ml of fulvestrant.

According to another aspect of the present invention there is provided a pharmaceutical formulation having a solubility for fulvestrant of at least 100 mg/ml adapted for intramuscular injection comprising;

100 mg/ml or more of fulvestrant;

10 5% (w/v) or more castor oil per volume of formulation vehicle;

and at least the following amounts (% weight/volume of formulation vehicle) of ethanol (EtOH), benzyl alcohol (BA), benzyl benzoate (BB) determined by the algorithm:

$100 = -29.77 + 5.44 \times \text{EtOH} + 2.38 \times \text{BA} + 1.57 \times \text{BB}$; and

provided that the following formulation is excluded: fulvestrant up to 102 mg/ml, 15 %
15 weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of castor oil per volume of formulation vehicle.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant at a concentration of at least 100 mg/ml in which the
20 formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months and provided that the following formulation is excluded:
fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 %
weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate
25 per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per volume of formulation vehicle.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically
30 significant blood plasma fulvestrant concentration in a human for at least 2 months.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant at a concentration of at least 100 mg/ml in which the formulation is adapted for intra-muscular injection into a human and which is capable after

injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant at a concentration of at least 150 mg/ml in which the
5 formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant at a concentration of at least 200 mg/ml in which the
10 formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months.

According to another aspect of the invention there is provided a pharmaceutical formulation comprising fulvestrant at a concentration of at least 300 mg/ml in which the
15 formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months.

According to one aspect of the invention there is provided any one of the following pharmaceutical formulations comprising about:

20 i)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

15% weight per volume of benzyl benzoate

500-555mg of fulvestrant for each 5ml of finished formulation

25 and the remaining amount as castor oil;

ii)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

30 500-700mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

iii)

10% weight per volume of ethanol

- 20% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
500-750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- 5 iv)
20% weight per volume of ethanol
20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
500-1175mg of fulvestrant for each 5ml of finished formulation
10 and the remaining amount as castor oil;
- v)
15% weight per volume of ethanol
10% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
15 500-810 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- vi)
15% weight per volume of ethanol
20% weight per volume of benzyl alcohol
20 50% weight per volume of benzyl benzoate
500 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- vii)
15% weight per volume of ethanol
25 20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
500-630mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- viii)
30 10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation

- and the remaining amount as castor oil;
- ix)
- 20% weight per volume of ethanol
- 20% weight per volume of benzyl alcohol
- 5 30% weight per volume of benzyl benzoate
- 750mg of fulvestrant for each 5ml of finished formulation
- and the remaining amount as castor oil;
- x)
- 15% weight per volume of ethanol
- 10 10% weight per volume of benzyl alcohol
- 50% weight per volume of benzyl benzoate
- 750mg of fulvestrant for each 5ml of finished formulation
- and the remaining amount as castor oil;
- xi)
- 15 9% weight per volume of ethanol
- 19% weight per volume of benzyl alcohol
- 47% weight per volume of benzyl benzoate
- 700mg of fulvestrant for each 5ml of finished formulation
- and the remaining amount as castor oil;
- 20 xii)
- 14% weight per volume of ethanol
- 19% weight per volume of benzyl alcohol
- 48% weight per volume of benzyl benzoate
- 700mg of fulvestrant for each 5ml of finished formulation
- 25 and the remaining amount as castor oil;
- xiii)
- 15% weight per volume of ethanol
- 20% weight per volume of benzyl alcohol
- 45% weight per volume of benzyl benzoate
- 30 750mg of fulvestrant for each 5ml of finished formulation
- and the remaining amount as castor oil;
- xiv)
- 9% weight per volume of ethanol

- 19% weight per volume of benzyl alcohol
47% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- 5 xv)
19% weight per volume of ethanol
19% weight per volume of benzyl alcohol
28% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
10 and the remaining amount as castor oil;
- xvi)
14% weight per volume of ethanol
9% weight per volume of benzyl alcohol
47% weight per volume of benzyl benzoate
15 750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- xvii)
14% weight per volume of ethanol
19% weight per volume of benzyl alcohol
20 47% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- xviii)
10% weight per volume of ethanol
25 20% weight per volume of benzyl alcohol
45% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;
- xix)
30 15% weight per volume of ethanol
10% weight per volume of benzyl alcohol
45% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xx)

20% weight per volume of ethanol

20% weight per volume of benzyl alcohol

5 25% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxi)

10% weight per volume of ethanol

10 30% weight per volume of benzyl alcohol

25% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxii)

15 10% weight per volume of ethanol

25% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

20 xxiii)

10% weight per volume of ethanol

30% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

25 and the remaining amount as castor oil;

xxiv)

15% weight per volume of ethanol

25% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

30 750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxv)

15% weight per volume of ethanol

- 25% weight per volume of benzyl alcohol
- 25% weight per volume of benzyl benzoate
- 750mg of fulvestrant for each 5ml of finished formulation
- and the remaining amount as castor oil; and
- 5 xxvi)
- 15% weight per volume of ethanol
- 20% weight per volume of benzyl alcohol
- 30% weight per volume of benzyl benzoate
- 750mg of fulvestrant for each 5ml of finished formulation
- 10 and the remaining amount as castor oil.

The term “comprising about” in this context means that the numerical value assigned to each component of the formulation may be varied independently to accommodate manufacturing specifications encountered by a skilled person when making up the formulations. Typically this means plus or minus 5%, more preferably plus or minus 4%, more preferably plus or minus 3%, more preferably plus or minus 2%, more preferably plus or minus 1%. In a preferred embodiment, more variation in drug level is allowed compared with other components. For example:

Drug (+/- %)	Other components (+/- %)
5	4, 3, 2 or 1
4	3, 2 or 1
3	2 or 1
2	1

20 The individual formulations described herein may comprise further excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant.

According to another aspect of the invention there is provided any one of the following pharmaceutical formulations:

- 25 i)
- 10% weight per volume of ethanol
- 20% weight per volume of benzyl alcohol

- 15% weight per volume of benzyl benzoate
500-555mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- ii)
- 5 10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
500-700mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- 10 iii)
- 10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
500-750mg of fulvestrant for each 5ml of finished formulation
- 15 and the remaining amount as castor oil
- iv)
- 20% weight per volume of ethanol
20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
- 20 500-1175mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- v)
- 15% weight per volume of ethanol
10% weight per volume of benzyl alcohol
- 25 50% weight per volume of benzyl benzoate
500-810 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- vi)
- 15% weight per volume of ethanol
- 30 20% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
500 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil

- vii)
15% weight per volume of ethanol
20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
5 500-630mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- viii)
10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
10 50% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- ix)
20% weight per volume of ethanol
15 20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- x)
20 15% weight per volume of ethanol
10% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil.
- 25 A preferred pharmaceutical formulation described herein is one wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
- A preferred pharmaceutical formulation described herein is one wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
- 30 A preferred pharmaceutical formulation described herein is one wherein the pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
- A preferred pharmaceutical formulation described herein is one wherein the ricinoleate excipient is castor oil.

According to one aspect of the present invention there is provided a pharmaceutical formulation adapted for intramuscular injection comprising 100 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of pharmaceutical formulation, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of pharmaceutical formulation and 5 % or more weight of ricinoleate excipient per volume of pharmaceutical formulation provided:

- a) the pharmaceutical formulation comprises at least 5 % weight of ethanol per volume of pharmaceutical formulation;
- b) if the pharmaceutically acceptable alcohol is less than or equal to 13%, then the pharmaceutical formulation must comprise at least 50 % non-aqueous ester solvent; and
- c) if the pharmaceutically acceptable alcohol is greater than 20 % but less than or equal to 25 %, then the pharmaceutical formulation must comprise at least 30 % non-aqueous ester solvent;

and also provided that the following pharmaceutical formulation is excluded: fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per volume of formulation vehicle.

A preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 100 mg/ml or more of fulvestrant, 20 % or more weight of a pharmaceutically acceptable alcohol per volume of pharmaceutical formulation, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of pharmaceutical formulation and 5 % or more weight of ricinoleate excipient per volume of pharmaceutical formulation provided:

- a) the pharmaceutical formulation comprises at least 10 % weight of ethanol per volume of pharmaceutical formulation;
- b) if the pharmaceutically acceptable alcohol is 20%, then the pharmaceutical formulation must comprise at least 22.5 % non-aqueous ester solvent; and
- c) if the pharmaceutically acceptable alcohol is greater than 20 % but less than or equal to 25 %, then the pharmaceutical formulation must comprise at least 15 % non-aqueous ester solvent;

and also provided that the following pharmaceutical formulation is excluded: fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl

alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per volume of formulation vehicle.

A more preferred pharmaceutical formulation adapted for intramuscular injection is one comprising 150 mg/ml or more of fulvestrant, 25 % or more weight of a pharmaceutically acceptable alcohol per volume of pharmaceutical formulation, 30 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of pharmaceutical formulation and 5 % or more weight of ricinoleate excipient per volume of pharmaceutical formulation provided:

- a) the pharmaceutical formulation comprises at least 10 % weight of ethanol per volume of pharmaceutical formulation;
- b) if the pharmaceutically acceptable alcohol is less than 30 %, then the pharmaceutical formulation must comprise at least 35 % non-aqueous ester solvent.

A particularly preferred pharmaceutical formulation is one which comprises 15% w/v or less of ethanol and in which the solubility of fulvestrant is at least 155mg/ml.

According to another aspect of the invention there is provided a unit dose of a pharmaceutical formulation as described herein wherein the total volume of the formulation is 6ml or less.

According to another aspect of the invention there is provided a pharmaceutical formulation adapted for intramuscular injection, as defined in any preceding claim for use in medical therapy.

According to another aspect of the invention there is provided use of fulvestrant in the preparation of a pharmaceutical formulation, as defined herein for the treatment of a benign or malignant disease of the breast or reproductive tract.

According to another aspect of the invention there is provided use of fulvestrant in the preparation of a pharmaceutical formulation, as defined in any preceding claim for the treatment of a benign or malignant disease of the breast or reproductive tract in a human with dosage intervals of at least 8 weeks.

According to another aspect of the invention there is provided a sterile syringe or vial comprising a pharmaceutical formulation as defined in any preceding claim.

The term "pharmaceutical formulation" as used herein means the combination of drug plus formulation vehicle. The terms "finished formulation" and "finished pharmaceutical formulation" mean the same as "pharmaceutical formulation".

The term "formulation vehicle" as used herein means the combination of all excipients used in the pharmaceutical formulation (and therefore excludes drug per se).

The distinction between pharmaceutical formulation and formulation vehicle is important for the following reason. For example, if the concentration (y % w/v) of an excipient "A" is measured by its concentration in formulation vehicle and then drug is added, the addition of drug will result in a concentration of excipient A that is lower than concentration y in the finished pharmaceutical formulation. To convert a concentration expressed in terms of "formulation vehicle" into a concentration of "finished pharmaceutical formulation" it is necessary to use a displacement value.

The "displacement value" is defined as the number of parts by weight of compound that displaces one part by weight of the formulation vehicle. The displacement value allows determination of the amount of formulation vehicle displaced by the compound. The displacement value is used to calculate the actual composition of the finished formulation in terms of proportions of excipients. The density of the compound affects the amount of formulation vehicle required to make the pharmaceutical formulation to the correct concentration. One part by weight of the compound with a density equal to the formulation vehicle will displace an equivalent volume of the formulation vehicle. A compound with twice the density of the formulation vehicle will displace half the volume. It is therefore necessary to make allowance for the compound in terms of the particular formulation vehicle, using the displacement value.

For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

Where whole numbers are used for % weight per volume of formulation, these refer to rounded numbers where appropriate. For example, 4.6% would be rounded to 5%.

It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, 5 when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of 10 two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol.

The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably 15 just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are 20 described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

25 It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

30 By the use of the term ricinoleate excipient we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as

triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

We have surprisingly found that the above formulations of the invention provide, after intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

5 We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still achieved by the formulation of the invention.

10 By use of the term "extended release" we mean at least 4 weeks, at least 5 weeks, and, preferably at least 8 weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for at least 8 weeks or 2 months, more preferably for at least 12 weeks or 3 months.

It will be understood that the attendant physician may wish to administer the
15 intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile.

Preferably 5ml of the intramuscular injection is administered.

20 Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant, more preferably an antioxidant.

As described above fulvestrant is useful in the treatment of oestrogen-dependent indications such as breast cancer and gynaecological conditions, such as endometriosis.

25 In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11 β -fluoro- 7 α -(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17 β -diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical structure. It is believed that the compound will also share with fulvestrant similar physical
30 properties and therefore the current invention will also have application with this compound.

Further features of the invention are those as described above but in which SH-646 is substituted for fulvestrant.

References

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. *Steroids*, 54: 71-100, 1989.
2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. *American New York Academy Science* 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. *Regulatory mechanisms in breast cancer*. Boston: Kluwer Academic, 1990b: 239-57.
4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry* 1990c; 37: 771-5.
- 10 5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. *Journal Endocrinology* 1987; 112: R7-10.
6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry* 1988; 3: 141-7.

15 The invention will now be illustrated by the following non-limiting Examples in which:

 Figure 1 shows plasma profiles obtained following IM injection (data normalised for rabbit weight, based on 3.2 kg rabbit) in which the y-axis is conc (ng/ml) and the x-axis is time (days);

20 Figure 2 shows comparison of plasma profiles in which:

 Figure 2A shows plasma profiles from group A in which the y-axis is conc (ng/ml) and the x-axis is time (days); and

 Figure 2B shows plasma profiles from group B in which the y-axis is conc (ng/ml) and the x-axis is time (days);

25 Figure 3 shows plasma profiles from formulations 1, 5 and Control (normalised for rabbit weight, based on 3.2 kg rabbit in which the y-axis is conc (ng/ml) and the x-axis is time (days);

 Figure 4 shows muscle residue data from 3 month PK study in which the y-axis is % fulvestrant remaining per injection site and the x-axis is formulation number. Each bar

30 represents one injection site (2 sites per animal).

 Figure 5 shows predicted versus actual solubility

 Figure 6 shows a confidence interval for predicted solubility

Figure 7 shows plasma profiles obtained following IM injection (data normalised for rabbit weight, based on 3.2 kg rabbit) in which the y-axis is conc (ng/ml) and the x-axis is time (days);

Abbreviations

5	IM	intramuscular
	PK	pharmacokinetic
	AUC	area under curve
	SD	standard deviation

10 Reference Example 1

Measurement of Solubility of Fulvestrant in Formulations

1. Materials and Apparatus

Balance

2 mL glass vials with screw caps

15 Magnetic stirrer bars

Temperature control reaction block with magnetic stirring facility

Positive displacement pipette (PDP) 20 - 25 μ L with appropriate microsyringe tips

Polycarbonate ultracentrifuge tubes

Ultracentrifuge

20 Pipette 0.5 - 200 μ L PDP with appropriate microsyringe tips

Pipette 200 μ L - 1 mL with appropriate plastic tips

2 mL amber Snap Top glass HPLC vials

1 mm Snap Caps for HPLC vials

HPLC kit with diode array detector

25 Methanol (MeOH) HPLC Grade

Acetonitrile (ACN) Far UV HPLC Grade

Ultrapure de-ionised water

25 cm H5 ODS 5 μ 4.6 mm i.d. HPLC column

Vortex mixer

30 Ultrasonic bath

20 - 200 μ L pipette with appropriate plastic tips

Aluminium weigh pans 5 mL glass volumetric flasks

2. Experimental procedure

- 2.1 1ml formulation vehicles were made up in triplicate by adding the appropriate volumes of alcohols and benzyl benzoate, and then adding castor oil by weight
- 5 2.2 Fulvestrant was then added to excess, until no more drug was seen to visibly dissolve. The weight of fulvestrant added was noted.
- 2.3 A magnetic stirrer bar was placed in each vial.
- 2.4 All the samples were overlaid with nitrogen and the vials were capped placed in the reaction block and stirred at a speed of 1000 at a temperature of 4°C
- 10 2.5 Using the PDP 20 - 200 µL pipettor, 200 µL aliquots were removed from each vial after 6 days and transferred to ultracentrifuge tubes. These tubes were centrifuged at a speed of 80,000 r.p.m. for 30 minutes at 25°C.
- 2.6 HPLC eluent was prepared by adding 1400 mL methanol, 450 mL water and 150 mL acetonitrile to a 2 litre plastic-coated solvent bottle.
- 15 2.7 990 µL HPLC eluent was added to 60 amber glass HPLC vials using a 1mL Pipette.
- 2.8 3 x 10 µL of supernatant were removed from each ultracentrifuge tube using the Pipette PDP 0.5-25 µL pipette and added to the vials containing eluent.
- 2.9 The samples were diluted again 1 in 10. 100µl sample was added to 900µl HPLC eluent
- 2.10 The amber vials were capped, vortex mixed for 10 seconds, sonicated for 10 minutes and
- 20 then placed in the HPLC autosampler tray.

3. Calibration preparation

- 3.1 Approximately 10 mg fulvestrant was accurately weighed into an aluminium weigh pan on the microbalance and placed in a 5 mL glass volumetric flask The actual weight was recorded.
- 25 3.2 Approximately 4.5 mL HPLC eluent was added to the flask using a plastic pasteur pipette. The flask was then sonicated for 5 minutes prior to making accurately to volume (to give a spiking solution of approximately 2 mg.mL⁻¹).
- 3.3 0 - 250 µL spiking solution was added to 2 mL amber HPLC vials using the appropriate Pipette and the volume made to 1 mL with HPLC eluent using a 1 mL Pipette
- 30 as shown in the table below:

Volume Spiking Solution (μL)	Volume HPLC eluent (μL)	Theoretical fulvestrant Concentration ($\mu\text{g}\cdot\text{mL}^{-1}$)
0	1000	0
5	995	10
25	975	50
50	950	100
100	900	200
150	850	300
200	800	400
250	750	500

3.4 The HPLC vials were capped, vortexed for 10 seconds and placed on the HPLC autosampler tray.

5 3.5 A calibration was prepared (as per 3.1 - 3.4) for both batches of ICI 182780.

4. HPLC Conditions

Eluent : 70% MeOH / 22.5% Water / 7.5% ACN
 Column : 25 cm 5μ Hypersil ODS 4.6 mm i.d. with guard column
 10 Detection wavelength : 280 nm
 Flow rate : 1.2 mL.min⁻¹
 Temperature : Ambient
 Injection volume : 50 μL
 Retention time : 12 minutes approximately

15

Example 1

Pharmaceutical Formulations

Fulvestrant is mixed with ethanol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil
 20 and stirred, (for convenience weight is used rather than volume by using the weight to volume

- 25 -

ratio). The bulk solution is overlaid with nitrogen. The solution is sterilised by filtration using one or two filters of 0.2µm porosity. The sterile filtrate is kept under a nitrogen overlay as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary pack to
5 facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing. The process flow diagram below depicts the manufacturing process.

Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of
10 each component to prepare the following formulations:

- a)
10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
15% weight per volume of benzyl benzoate
15 500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- b)
10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
20 30% weight per volume of benzyl benzoate
500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- c)
10% weight per volume of ethanol
25 20% weight per volume of benzyl alcohol
50% weight per volume of benzyl benzoate
500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil
- d)
30 20% weight per volume of ethanol
20% weight per volume of benzyl alcohol
30% weight per volume of benzyl benzoate

- 26 -

500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil

e)

15% weight per volume of ethanol

5 10% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil

f)

10 15% weight per volume of ethanol


20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

500mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil

15

FLOW DIAGRAM OF MANUFACTURING



Example 2**Stability studies to 4 months for selection of stable 100mg/ml Pharmaceutical formulations**

Formulation No.	Fulvestrant (mg/ml)	96% EtOH % w/v	Benzyl Alcohol % w/v	Benzyl Benzoate % w/v	Castor oil % w/v	Observations (visual) 4 months
Sample 1	100	5	10	15	to 100%	Precipitated
Sample 2	100	5	10	30	to 100%	Precipitated
Sample 3	100	7.5	10	15	to 100%	Precipitated
Sample 4	100	7.5	10	30	to 100%	Precipitated
Sample 5	100	10	10	15	to 100%	Precipitated
Sample 6	100	10	10	17.5	to 100%	Precipitated
Sample 7	100	10	10	20	to 100%	Precipitated
Sample 8	100	10	10	22.5	to 100%	Solution
Sample 9	100	10	10	25	to 100%	Solution
Sample 10	100	10	10	27.5	to 100%	Solution
Sample 11	100	10	10	30	to 100%	Solution
Sample 12	100	10	10	40	to 100%	Solution
Sample 13	100	10	10	50	to 100%	Solution
Sample 14	100	10	15	15	to 100%	Solution
Sample 15	100	10	15	30	to 100%	Solution
Sample 16	100	10	15	40	to 100%	Solution
Sample 17	100	10	15	50	to 100%	Solution
Sample 18	100	10	20	15	to 100%	Solution
Sample 19	100	10	20	30	to 100%	Solution
Sample 20	100	10	20	40	to 100%	Solution
Sample 21	100	10	20	50	to 100%	Solution
Sample 22	100	15	10	15	to 100%	Solution
Sample 23	100	15	10	30	to 100%	Solution
Sample 24	100	15	10	40	to 100%	Solution
Sample 25	100	15	10	50	to 100%	Solution
Sample 26	100	20	5	15	to 100%	Solution
Sample 27	100	20	5	30	to 100%	Solution
Sample 28	100	20	10	15	to 100%	Solution
Sample 29	100	20	10	30	to 100%	Solution
Sample 30	100	20	10	40	to 100%	Solution
Sample 31	100	20	10	50	to 100%	Solution
Sample 32	100	15	15	15	to 100%	Solution
Sample 33	100	15	15	30	to 100%	Solution
Sample 34	100	15	15	40	to 100%	Solution
Sample 35	100	15	15	50	to 100%	Solution
Sample 36	100	15	20	15	to 100%	Solution
Sample 37	100	15	20	30	to 100%	Solution
Sample 38	100	15	20	50	to 100%	Solution
Sample 39	100	20	20	15	to 100%	Solution

- 29 -

Sample 40	100	20	20	30	to 100%	Solution
Control Sample	52.16	10	10	15	60	Solution

The Control Sample refers to the following Pharmaceutical formulation: fulvestrant 50mg/ml, ethanol 10% w/v, benzylalcohol 10% w/v, benzyl benzoate 15% w/v and made to volume with castor oil.

5

Example 3

Pharmaceutical formulations selected for *in vivo* deposition and *in vitro* precipitation studies

The pharmaceutical formulations below were selected from Example 2 for further study.

Formulation	Fulvestrant	95% ETOH	Benzyl Alcohol	Benzyl Benzoate	Castor oil
	% w/v	%w/v	%w/v	%w/v	% w/v
Sample 1	10	10	10	30	to 100%
Sample 2	10	10	10	50	to 100%
Sample 3	10	10	20	15	to 100%
Sample 4	10	10	20	30	to 100%
Sample 5	10	10	20	50	to 100%
Sample 6	10	20	5	15	to 100%
Sample 7	10	20	5	30	to 100%
Sample 8	10	20	20	15	to 100%
Sample 9	10	20	20	30	to 100%
Sample 10	10	15	10	15	to 100%
Sample 11	10	15	10	30	to 100%
Sample 12	10	15	10	50	to 100%
Sample 13	10	15	20	15	to 100%
Sample 14	10	15	20	50	to 100%
Sample 15	10	15	20	30	to 100%
Sample 16	5	10	10	15	to 100%

10

A matrix of 7 pharmaceutical formulations (samples 3, 4, 5, 9, 12, 14 and 16 – see Example 3 below) was identified for further evaluation from *in vitro* precipitation and deposition studies.

- 30 -

Sample 16 was a control. The precipitation experiment involved visual inspection of each sample under conditions where evaporation of alcohols led to precipitation of drug.

Example 4

5 In vivo Studies

Pharmaceutical formulations identified for further *in vivo* evaluation

No.	Fulvestrant (%w/v)	Excipients (% w/v)			
		Ethanol 96%	Benzyl alcohol	Benzyl benzoate	Castor oil
10	F1	10	20	15	45
	F2	10	20	30	30
	F3	10	20	50	10
	F4	20	20	30	20
	F5	15	10	50	15
15	F6	15	20	50	5
	F7	15	20	30	25
	Control	10	10	15	60

(a) **An *in vivo* pharmacokinetic (PK) study on these 7 pharmaceutical formulations was performed over 3 months duration; results are shown in Figures 1, 2 and 3.**

Analysis of PK results

Plasma levels were more variable than Control over the first 30 days; variability was similar to control thereafter. After 2 months, drug levels were equivalent to Control at 1 month indicating a prolonged period of action over Control. This release profile was surprising because, compared with Control which does not precipitate drug locally, local precipitation at the site of injection of the tested formulations was expected to impair their release profile.

Some differences in profiles were noted over the first 30 days such that they were divided into 2 groups (with Formulation F7 showing intermediate behaviour).

Group A, rapid release early time points (50% Benzyl benzoate and low castor oil $\leq 15\%$) – see figure 2A

Group B, lower release, flatter profile ($\leq 30\%$ Benzyl benzoate and higher castor oil $\geq 20\%$) – see figure 2B

35

- 31 -

(b) Histopathology

Local tolerance was assessed to IM injection of the 7 pharmaceutical formulations and Control. Lesions were observed over a 51 day period. All pharmaceutical formulations 5 caused tissue reactions greater than Control.

(c) Plasma levels for 100mg dose at critical time points (ng/ml)

	Form.	Dose (mg)	Time point (days)		
			28	56	84
10	1	100	8.7	4.6	3.5
	2	100	8.0	3.6	2.6
	3	100	9.4	3.5	2.0
	4	100	7.7	5.0	3.4
15	5	100	9.1	4.5	2.6
	6	100	9.9	3.3	1.9
	7	100	9.9	5.5	3.2
	Control	50	3.3	1.7	

20 Summary

Duration for 100mg dose = min 2 months.

Duration for 150mg dose = min 3 months.

(d) Measurement of Fulvestrant solubility in 7 pharmaceutical formulations after 6 days

	Formulation	Solubility (mg/ml)
25	F1	111
	F2	140
	F3	175
	F4	235
30	F5	162
	F6	212
	F7	126

35

Example 5**Model for Fulvestrant Solubility**

Formulation No.	Formulation vehicles for Fulvestrant - solubility				Measured	
	96% ETOH % w/v	Benzyl Alcohol % w/v	Benzyl Benzoate % w/v	Castor oil % w/v	Fulvestrant Solubility (mg/ml)	Predicted solubility (mg/ml)
Sample 1	5	5	0	to 100%	27	9.4
Sample 2	5	5	15	to 100%	36	32.8
Sample 3	10	5	0	to 100%	46	48.5
Sample 4	10	5	15	to 100%	54	36.6
Sample 5	10	10	0	to 100%	45	60.1
Sample 6	10	10	15	to 100%	65	72
Sample 7	15	15	0	to 100%	76	87.6
Sample 8	15	15	15	to 100%	102	111.1
Sample 9	11	22	17	to 100%	111	109.8
Sample 10	11	22	33	to 100%	140	166.1
Sample 11	11	22	56	to 100%	175	135.8
Sample 12	22	22	33	to 100%	235	174.4
Sample 13	17	11	56	to 100%	162	170.6
Sample 14	17	22	56	to 100%	212	200.9
Sample 15	17	22	33	to 100%	126	196.3

5

A linear regression model was fitted to solubility data from 15 samples using as independent variables the % ethanol, benzyl alcohol and benzyl benzoate levels in the formulations. The following model was obtained which had an R-Squared value of 93.2%:

$$\text{SOLUBILITY} = -29.77 + 5.44 \times \text{ETOH} + 2.38 \times \text{BA} + 1.57 \times \text{BB}$$

10 Benzyl alcohol = BA, benzyl benzoate = BB, Ethanol = ETOH.

Solubility measured as mg/ml

See Figures 5 and 6 based on the following data

measured	lower C.L.	predicted	upper C.L.
27	0	9.4	31.1
36	10	32.8	55.7
45	31.6	48.5	65.4
46	17.1	36.6	56.1
54	40.9	60.1	79.2
65	59	72	85
76	64.7	87.6	110.6
102	95.6	111.1	126.7
111	85.4	109.8	134.1
126	149	166.1	183.1

- 33 -

140	114.2	135.8	157.5
162	142.5	174.4	206.3
175	143.4	170.6	197.9
212	179.6	200.9	222.2
235	168.7	196.3	224

The table below shows predicted solubilities for a matrix of pharmaceutical formulations tested for stability. A good correlation was obtained with visual observations.

Formulation of Fulvestrant - stability

Formulation No.	Fulvestrant (mg/ml)	95% ETOH % w/v	Benzyl Alcohol % w/v	Benzyl Benzoate % w/v	Castor oil % w/v	Observations 4 months	Predicted solubility (mg/ml)
Sample 1	100	5	10	15	60	Precipitated	53.1
Sample 2	100	5	10	30	45	Precipitated	79.2
Sample 3	100	7.5	10	15	57.5	Precipitated	68.2
Sample 4	100	7.5	10	30	42.5	Precipitated	94.3
Sample 5	100	10	10	15	55	Precipitated	83.3
Sample 6	100	10	10	17.5	52.5	Precipitated	87.6
Sample 7	100	10	10	20	50	Precipitated	92.0
Sample 8	100	10	10	22.5	47.5	Solution	96.4
Sample 9	100	10	10	25	45	Solution	100.7
Sample 10	100	10	10	27.5	42.5	Solution	105.1
Sample 11	100	10	10	30	40	Solution	109.5
Sample 12	100	10	10	40	30	Solution	126.9
Sample 13	100	10	10	50	20	Solution	144.3
Sample 14	100	10	15	15	50	Solution	96.5
Sample 15	100	10	15	30	35	Solution	122.7
Sample 16	100	10	15	40	25	Solution	140.1
Sample 17	100	10	15	50	15	Solution	157.6
Sample 18	100	10	20	15	45	Solution	109.7
Sample 19	100	10	20	30	30	Solution	135.9
Sample 20	100	10	20	40	20	Solution	153.3
Sample 21	100	10	20	50	10	Solution	170.8
Sample 22	100	15	10	15	50	Solution	113.5
Sample 23	100	15	10	30	35	Solution	139.7
Sample 24	100	15	10	40	25	Solution	157.1
Sample 25	100	15	10	50	15	Solution	174.6
Sample 26	100	20	5	15	50	Solution	130.5
Sample 27	100	20	5	30	35	Solution	156.7
Sample 28	100	20	10	15	45	Solution	143.7
Sample 29	100	20	10	30	30	Solution	169.9
Sample 30	100	20	10	40	20	Solution	187.3
Sample 31	100	20	10	50	10	Solution	204.8
Sample 32	100	15	15	15	45	Solution	126.7
Sample 33	100	15	15	30	30	Solution	152.9
Sample 34	100	15	15	40	20	Solution	170.3
Sample 35	100	15	15	50	10	Solution	187.8
Sample 36	100	15	20	15	40	Solution	140.0
Sample 37	100	15	20	30	25	Solution	166.1
Sample 38	100	15	20	50	5	Solution	201.0

- 34 -

Sample 39	100	20	20	15	35	Solution	170.2
Sample 40	100	20	20	30	20	Solution	196.3
Control Sample [Faslodex]	52.16	10	10	15	60	Solution	72.0

The Tables below show predicted formulations for various solubilities of fulvestrant; where an "X" means in solution. Note that the Tables include some impractical formulations where the sum of components becomes greater than 100%. The principal purpose is to illustrate the wide combinations of ethanol/ benzyl alcohol / benzyl benzoate taught by the invention to achieve different solubilities of fulvestrant.

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG			
1				BB																																
2	SOLUBILITY	ETHANOL	BA	3	5	8	10	13	15	18	20	23	25	28	30	33	35	38	40	43	45	48	50	53	55	58	60	63	65	68	70	73	75			
3	100																																			
4		5	5																									X	X	X	X	X	X	X		
5		5	10																							X	X	X	X	X	X	X	X	X		
6		5	15																				X	X	X	X	X	X	X	X	X	X	X	X		
7		5	20																	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
8		10	5																			X	X	X	X	X	X	X	X	X	X	X	X	X		
9		10	10																	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
10		10	15												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
11		10	20								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
12		15	5												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
13		15	10								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
14		15	15				X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
15		15	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
16		20	5			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
17		20	10	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
18		20	15	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
19		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
20	105																																			
21		5	5																														X	X	X	X
22		5	10																							X	X	X	X	X	X	X	X	X	X	
23		5	15																				X	X	X	X	X	X	X	X	X	X	X	X	X	
24		5	20																	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
25		10	5																			X	X	X	X	X	X	X	X	X	X	X	X	X	X	
26		10	10																	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
27		10	15													X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
28		10	20										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
29		15	5												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
30		15	10								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
31		15	15					X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
32		15	20		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
33		20	5			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
34		20	10	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
35		20	15	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
36		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
37	110																																			
38		5	5																														X	X	X	X
39		5	10																								X	X	X	X	X	X	X	X	X	
40		5	15																					X	X	X	X	X	X	X	X	X	X	X	X	X
41		5	20																			X	X	X	X	X	X	X	X	X	X	X	X	X	X	
42		10	5																			X	X	X	X	X	X	X	X	X	X	X	X	X	X	
43		10	10																		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
44		10	15														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
45		10	20								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
46		15	5												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG			
47				BB																																
48	SOLUBILITY	ETHANOL	BA	3	5	8	10	13	15	18	20	23	25	28	30	33	35	38	40	43	45	48	50	53	55	58	60	63	65	68	70	73	75			
49		15	10										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
50		15	15							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
51		15	20			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
52		20	5					X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
53		20	10		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
54		20	15	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
55		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
56	115																																			
57		5	5																														X	X	X	
58		5	10																								X	X	X	X	X	X	X	X	X	
59		5	15																					X	X	X	X	X	X	X	X	X	X	X	X	X
60		5	20																			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
61		10	5																					X	X	X	X	X	X	X	X	X	X	X	X	X
62		10	10																		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
63		10	15														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
64		10	20											X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
65		15	5														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
66		15	10										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
67		15	15							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
68		15	20			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
69		20	5					X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
70		20	10			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
71		20	15	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
72		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
73	120																																			
74		5	5																																X	
75		5	10																														X	X	X	
76		5	15																								X	X	X	X	X	X	X	X	X	X
77		5	20																				X	X	X	X	X	X	X	X	X	X	X	X	X	X
78		10	5																							X	X	X	X	X	X	X	X	X	X	X
79		10	10																			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
80		10	15																		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
81		10	20														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
82		15	5																								X	X	X	X	X	X	X	X	X	X
83		15	10														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
84		15	15							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
85		15	20						X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
86		20	5							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
87		20	10					X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
88		20	15	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
89		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
90	130																																			
91		5	5																																	
92		5	10																															X	X	

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG	
93				BB																														
94	SOLUBILITY	ETHANOL	BA	3	5	8	10	13	15	18	20	23	25	28	30	33	35	38	40	43	45	48	50	53	55	58	60	63	65	68	70	73	75	
95		5	15																							X	X	X	X	X	X	X	X	
96		5	20																							X	X	X	X	X	X	X	X	
97		10	5																							X	X	X	X	X	X	X	X	
98		10	10																						X	X	X	X	X	X	X	X	X	
99		10	15																			X	X	X	X	X	X	X	X	X	X	X	X	
100		10	20																X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
101		15	5																		X	X	X	X	X	X	X	X	X	X	X	X	X	
102		15	10															X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
103		15	15												X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
104		15	20								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
105		20	5										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
106		20	10							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
107		20	15			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
108		20	20	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
109	140																																	
110		5	5																															
111		5	10																															
112		5	15																													X	X	
113		5	20																							X	X	X	X	X	X	X	X	
114		10	5																													X	X	
115		10	10																								X	X	X	X	X	X	X	
116		10	15																						X	X	X	X	X	X	X	X	X	
117		10	20																		X	X	X	X	X	X	X	X	X	X	X	X	X	
118		15	5																				X	X	X	X	X	X	X	X	X	X	X	
119		15	10																		X	X	X	X	X	X	X	X	X	X	X	X	X	
120		15	15															X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
121		15	20										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
122		20	5																							X	X	X	X	X	X	X	X	
123		20	10										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
124		20	15							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
125		20	20			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
126	150																																	
127		5	5																															
128		5	10																															
129		5	15																															
130		5	20																												X	X	X	
131		10	5																														X	
132		10	10																											X	X	X	X	
133		10	15																							X	X	X	X	X	X	X	X	
134		10	20																					X	X	X	X	X	X	X	X	X	X	
135		15	5																							X	X	X	X	X	X	X	X	
136		15	10																		X	X	X	X	X	X	X	X	X	X	X	X	X	
137		15	15																		X	X	X	X	X	X	X	X	X	X	X	X	X	
138		15	20																		X	X	X	X	X	X	X	X	X	X	X	X	X	

WO 03/006064

PCT/GB02/03092

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG				
139				BB																																	
140	SOLUBILITY	ETHANOL	BA	3	5	8	10	13	15	18	20	23	25	28	30	33	35	38	40	43	45	48	50	53	55	58	60	63	65	68	70	73	75				
141		20	5																	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		
142		20	10														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
143		20	15										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
144		20	20							X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
145	200																																				
146		5	5																																		
147		5	10																																		
148		5	15																																		
149		5	20																																		
150		10	5																																		
151		10	10																																		
152		10	15																																		
153		10	20																																		
154		15	5																																		
155		15	10																																		
156		15	15																																		X
157		15	20																																	X	X
158		20	5																																	X	X
159		20	10																																	X	X
160		20	15																							X	X	X	X	X	X	X	X	X	X	X	X
161		20	20																					X	X	X	X	X	X	X	X	X	X	X	X	X	X

Example 6**In-vivo Pharmacokinetic study using compositions at 140mg/ml of fulvestrant**

Formulations F3 and F6 as described in Example 4 were modified to contain an increased level of fulvestrant to 140mg/ml. The modified formulations were named F8 and

5 F9 as described below.

Formulation Composition

The compositions of the formulations dosed in the PK study are shown in the table below.

Formulation No.	Fulvestrant (%w/v)	Ethanol 96% (%w/v)	Benzyl alcohol (%w/v)	Benzyl Benzoate (%w/v)	Castor Oil (%w/v)
F8	14	9	19	47	To 100%
F9	14	14	19	48	To 100%

10 **PK Profile**

The results are set out in Figure 7. The composition of the Control is the same as described in Example 4. Compositions F8 and F9 gave similar profiles with improved performance in terms of extended release of higher levels fulvestrant compared with Control.

15 **Example 7****Compositions at 150mg/ml fulvestrant**

Compositions analogous or similar to F3, F4, F5 and F6 (see Example 4) but comprising 150mg/ml of fulvestrant are prepared as follows.

Formulation No.	Fulvestrant (%w/v)	Ethanol 96% (%w/v)	Benzyl alcohol (%w/v)	Benzyl Benzoate (%w/v)	Castor Oil (%w/v)
F10	15	10	20	50	To 100%
F11	15	20	20	30	To 100%
F12	15	15	10	50	To 100%
F13	15	15	20	45	To 100%
F14	15	9	19	47	To 100%
F15	15	19	19	28	To 100%

- 40 -

F16	15	14	9	47	To 100%
F17	15	14	19	47	To 100%
F18	15	10	20	45	To 100%
F19	15	15	10	45	To 100%
F20	15	20	20	25	To 100%
F21	15	10	30	25	To 100%
F22	15	10	25	30	To 100%
F23	15	10	30	30	To 100%
F24	15	15	25	30	To 100%
F25	15	15	25	25	To 100%
F26	15	15	20	30	To 100%

Claims

1. A pharmaceutical formulation adapted for intramuscular injection comprising 100 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle and provided that the following formulation is excluded: fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per volume of formulation vehicle.
2. A pharmaceutical formulation adapted for intramuscular injection according to claim 2 comprising 105 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation comprises at least 5 % weight of ethanol per volume of formulation vehicle.
3. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 110 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.
4. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 115 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the

- 42 -

formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

5. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 120 mg/ml or more of fulvestrant, 10 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

6. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 130 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 5 % weight of ethanol per volume of formulation vehicle.

7. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 140 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 12.5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the formulation vehicle comprises at least 10 % weight of ethanol per volume of formulation vehicle.

8. A pharmaceutical formulation adapted for intramuscular injection according to claim 1 comprising 150 mg/ml or more of fulvestrant, 15 % or more weight of a pharmaceutically acceptable alcohol per volume of formulation vehicle, 17.5 % or more weight of a pharmaceutically acceptable non-aqueous ester solvent per volume of formulation vehicle and 5 % or more weight of ricinoleate excipient per volume of formulation vehicle provided the

- 43 -

formulation vehicle comprises at least 10 % weight of ethanol per volume of formulation vehicle.

9. A pharmaceutical formulation having a solubility for fulvestrant of at least Y mg/ml adapted for intramuscular injection comprising;

100 mg/ml or more of fulvestrant;

5% (w/v) or more castor oil per volume of formulation vehicle;

and at least the following amounts (% weight/volume of formulation vehicle) of ethanol (ETOH), benzyl alcohol (BA), benzyl benzoate (BB) determined by the algorithm:

$$Y = -29.77 + 5.44\text{ETOH} + 2.38\text{BA} + 1.57\text{BB}$$

wherein Y is at least 100, ETOH is at least 5, BA is at least 5 and BB is at least 5.

10. A pharmaceutical formulation according to claim 9 wherein Y is selected from the group consisting of 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 170, 180, 190, and 200.

11. A pharmaceutical formulation according to claim 9 wherein Y is selected from the group consisting of 120, 125, 130, 135, 140, 145, 150, 155, 160, 170, 180, 190, and 200.

12. A pharmaceutical formulation according to claim 9 wherein Y is selected from the group consisting of 150, 155, 160, 170, 180, 190 and 200.

13. A pharmaceutical formulation according to claim 9 wherein Y is selected from 150, 155, 160, 170, 180, 190 and 200 and the formulation comprises at least 150mg/ml of fulvestrant.

14. A pharmaceutical formulation according to claim 9 wherein Y is 200 and the formulation comprises at least 200mg/ml of fulvestrant.

15. A pharmaceutical formulation comprising fulvestrant at a concentration of at least 100 mg/ml in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months and provided that the following

- 44 -

formulation is excluded: fulvestrant up to 102 mg/ml, 15 % weight of ethanol per volume of formulation vehicle, 15 % weight of benzyl alcohol per volume of formulation vehicle, 15 % weight of benzyl benzoate per volume of formulation vehicle and 30 % or more weight of ricinoleate excipient per volume of formulation vehicle.

16. A pharmaceutical formulation comprising fulvestrant at a concentration of at least 150 mg/ml in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration in a human for at least 2 months.

17. Any one of the following pharmaceutical formulations comprising about:

i)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

15% weight per volume of benzyl benzoate

500-555mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

ii)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

500-700mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

iii)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

500-750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

iv)

20% weight per volume of ethanol

20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

- 45 -

500-1175mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

v)

15% weight per volume of ethanol

10% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

500-810 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

vi)

15% weight per volume of ethanol

20% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

500 mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

vii)

15% weight per volume of ethanol

20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

500-630mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

viii)

10% weight per volume of ethanol

20% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

ix)

20% weight per volume of ethanol

20% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

- 46 -

x)

15% weight per volume of ethanol

10% weight per volume of benzyl alcohol

50% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xi)

9% weight per volume of ethanol

19% weight per volume of benzyl alcohol

47% weight per volume of benzyl benzoate

700mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xii)

14% weight per volume of ethanol

19% weight per volume of benzyl alcohol

48% weight per volume of benzyl benzoate

700mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xiii)

15% weight per volume of ethanol

20% weight per volume of benzyl alcohol

45% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xiv)

9% weight per volume of ethanol

19% weight per volume of benzyl alcohol

47% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xv)

19% weight per volume of ethanol

19% weight per volume of benzyl alcohol

- 47 -

28% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

xvi)

14% weight per volume of ethanol
9% weight per volume of benzyl alcohol
47% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

xvii)

14% weight per volume of ethanol
19% weight per volume of benzyl alcohol
47% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

xviii)

10% weight per volume of ethanol
20% weight per volume of benzyl alcohol
45% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

xix)

15% weight per volume of ethanol
10% weight per volume of benzyl alcohol
45% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

xx)

20% weight per volume of ethanol
20% weight per volume of benzyl alcohol
25% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil;

- 48 -

xxi)

10% weight per volume of ethanol

30% weight per volume of benzyl alcohol

25% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxii)

10% weight per volume of ethanol

25% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxiii)

10% weight per volume of ethanol

30% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxiv)

15% weight per volume of ethanol

25% weight per volume of benzyl alcohol

30% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil;

xxv)

15% weight per volume of ethanol

25% weight per volume of benzyl alcohol

25% weight per volume of benzyl benzoate

750mg of fulvestrant for each 5ml of finished formulation

and the remaining amount as castor oil; and

xxvi)

15% weight per volume of ethanol

20% weight per volume of benzyl alcohol

- 49 -

30% weight per volume of benzyl benzoate
750mg of fulvestrant for each 5ml of finished formulation
and the remaining amount as castor oil.

18. A pharmaceutical formulation as claimed in any preceding claim wherein the pharmaceutically-acceptable alcohol is a mixture of ethanol and benzyl alcohol.
19. A pharmaceutical formulation as claimed in any preceding claim wherein the pharmaceutically-acceptable non-aqueous ester solvent is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.
20. A pharmaceutical formulation as claimed in any preceding claim wherein the pharmaceutically-acceptable non-aqueous ester solvent is benzyl benzoate.
21. A pharmaceutical formulation as claimed in any preceding claim wherein the ricinoleate excipient is castor oil.
22. A unit dose of a pharmaceutical formulation as claimed in any preceding claim wherein the total volume of the formulation is 6ml or less.
23. A pharmaceutical formulation adapted for intramuscular injection, as defined in any preceding claim for use in medical therapy.
24. Use of fulvestrant in the preparation of a pharmaceutical formulation, as defined in any preceding claim for the treatment of a benign or malignant disease of the breast or reproductive tract.
25. Use of fulvestrant in the preparation of a pharmaceutical formulation, as defined in any preceding claim for the treatment of a benign or malignant disease of the breast or reproductive tract in a human with dosage intervals of at least 8 weeks.
26. A sterile syringe or vial comprising a pharmaceutical formulation as defined in any preceding claim.





Fig.2A.






Fig.3.





Fig.7.







Fig.4.






Fig.6.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/03092

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K47/14 A61K47/44 A61K47/10 A61K31/565

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
 EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 21440 A (ZENECA) 19 June 1997 (1997-06-19) claims 1,5 examples 1,3,4 page 4, line 15 -page 6, line 30 ---	1-26
Y	DR H. FIEDLER EDITOR: "Lexicon der Hilfstoffe für Pharmazie, Kosmetik und angrenzende Gebiete" 1981, EDITIO CANTOR AULENDORF, D-7960 AULENDORF XP002221673 page 788 -page 789 ---	1-26
Y	EP 0 346 014 A (IMPERIAL CHEMICAL INDUSTRIES) 13 December 1989 (1989-12-13) page 9; example 3 --- -/--	1-26

Further documents are listed in the continuation of box C. Patent family members are listed in annex.

° Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document but published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
O document referring to an oral disclosure, use, exhibition or other means	* & * document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 20 November 2002	Date of mailing of the international search report 05/12/2002
---	--

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016	Authorized officer Giacobbe, S
--	---------------------------------------

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/03092

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOHN C. WATERTON; ET AL.: "A Case of Adenomyosis in a Pigtailed Monkey Diagnosed by Magnetic Resonance Imaging and treated with the Novel Pure Antiestrogen, ICI 182,780" LABORATORY ANIMAL SCIENCE, vol. 43, no. 3, 1993, pages 247-251, XP000998289 page 247, column 2, line 32 - line 35 ---	1-26
P,X	WO 01 51056 A (ASTRAZENECA UK LTD ;EVANS JOHN RAYMOND (GB); GRUNDY ROSALIND URSUL) 19 July 2001 (2001-07-19) page 17, line 5 - line 13 tables 3,4 claims 1-23 ---	1-26
P,X	WO 01 74366 A (THURLIMANN BEAT ;ASTRAZENECA UK LTD (GB); ASTRAZENECA AB (SE)) 11 October 2001 (2001-10-11) page 3, line 20 -page 4, line 8 page 16, paragraph 2.3 - paragraph 2.4 -----	1-26

INTERNATIONAL SEARCH REPORT

International Application No
 PCT/GB 02/03092

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9721440	A	19-06-1997	AU 1039097 A	03-07-1997
			WO 9721440 A1	19-06-1997
			ZA 9610426 A	23-06-1997

EP 346014	A	13-12-1989	AT 109351 T	15-08-1994
			AU 622184 B2	02-04-1992
			AU 3592389 A	07-12-1989
			CA 1337591 A1	21-11-1995
			DE 68917219 D1	08-09-1994
			DE 68917219 T2	15-12-1994
			DK 273489 A	07-12-1989
			EP 0346014 A1	13-12-1989
			ES 2057124 T3	16-10-1994
			FI 892762 A	07-12-1989
			IE 64368 B1	26-07-1995
			IL 90410 A	31-07-1994
			JP 2042024 A	13-02-1990
			JP 2918042 B2	12-07-1999
			NO 892293 A	07-12-1989
			NZ 229392 A	29-01-1992
			PT 90764 A ,B	29-12-1989
US 5183814 A	02-02-1993			
ZA 8903892 A	28-02-1990			

WO 0151056	A	19-07-2001	AU 2386301 A	24-07-2001
			BE 1013477 A3	05-02-2002
			BR 0107445 A	17-09-2002
			DE 10100779 A1	26-07-2001
			EP 1250138 A1	23-10-2002
			FR 2803516 A1	13-07-2001
			WO 0151056 A1	19-07-2001
			GB 2359254 A ,B	22-08-2001
			NL 1017075 C2	25-09-2001
			NL 1017075 A1	11-07-2001
			NO 20023227 A	03-07-2002
			PT 102548 A	31-07-2001
			US 2001020016 A1	06-09-2001

WO 0174366	A	11-10-2001	AU 4437201 A	15-10-2001
			WO 0174366 A1	11-10-2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION of:)	Confirmation No. 2093
)	
EVANS et al.)	
)	
Application No.: 10/872,784)	Group Art Unit: 1617
)	
Filed: June 22, 2004)	Examiner: Hui, San-Ming R
)	
FOR: FORMULATION)	
)	

DECLARATION UNDER 35 U.S.C. § 1.132
OF PAUL RICHARD GELLERT

PAUL RICHARD GELLERT of AstraZeneca, Alderley Park, Macclesfield, Cheshire, UK
declares:

1. I graduated from the University of Oxford in Chemistry in 1984. I undertook postgraduate research with Professor Brian Howard in the Physical Chemistry Laboratory at the University of Oxford leading to the award of a D.Phil in 1988. From February 1988 until the present I have been employed by AstraZeneca, (formerly Zeneca and ICI) initially as a Senior Research Scientist and subsequently as a Team Leader/Manager, Principal Scientist and, since 2004, a Senior Principal Scientist.
2. I have worked in the formulation and drug delivery area throughout my career with AstraZeneca, where my research and development work has covered a range of formulation types including sustained released injections, including fulvestrant.
3. During the course of my study of the subject application (hereinafter "the Evans Application") and the underlying data, I have become aware of several transcription or other errors between certain disclosures of the subject application and the underlying laboratory notebook data. One purpose of this Declaration is to point out the existence

and nature of these errors and to report further testing that has been carried out under my guidance to obtain additional data (paragraphs 4-10 below and Attachments A-D). A further purpose of this Declaration is to set out and document the manner in which an experienced formulator would likely have approached the task of developing a sustained release injectable formulation suitable for human use for a steroidal compound such as fulvestrant in about early 2000, which I understand is when the priority applications supporting the Evans Application were filed (paragraphs 11 - 25 below and Attachment E). Citations to literature and patent references in this Declaration will be in the format Lead Author (Date), and the full citations are given in the Table of References at the end of this Declaration. A copy of each cited reference (or cited portions of the longer references) is included in Attachment F under the Tab number noted in the Table of References.

4. In Table 2 of the Evans Application, the solubility of fulvestrant in castor oil appears to have been transcribed incorrectly from the original source, the laboratory notebook. The value in the latter is 24.5 mg/ml and not 20 mg/ml. In other experiments to determine the solubility of fulvestrant in castor oil and also in benzyl benzoate, some variability was observed.
5. In Table 3 of the Evans Application, the given solubility values were generated at 4°C and not at 25°C as is stated in the title of Table 3. For fulvestrant formulations, it is preferable that the fulvestrant remains completely in solution at both 4°C and 25°C. The 4°C temperature corresponds to the storage temperature (2°C to 8°C in the FDA approved label for Faslodex), and the 25°C temperature corresponds to the administration temperature (ambient temperature). In addition, the specified solubility values on this Table 3 are mean values calculated from analysis of replicate samples from one or more trials. The individual values are shown in handwriting in the amended version of Table 3 in Attachment A. In addition, it appears that the mean values for the last three compositions have been incorrectly calculated. The corrected mean values, together with the correction of the temperature from "25°C" to read "4°C", are also shown in handwriting in the amended version of Table 3 in Attachment A.

6. I have evaluated the transcription and other errors against the original application disclosures and conclude that these do not change the ultimate conclusions made from the data as originally reported. The addition of 15% w/v benzyl benzoate to compositions having total alcohol concentrations in castor oil of 10%, 15%, 20% and 30% w/v unexpectedly provides a positive effect on fulvestrant solubility, significantly increasing the solubility of fulvestrant in the compositions despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.
7. An additional set of experiments has been conducted at 25°C under my guidance to obtain consistent data with reduced variability from a single set of rigorously controlled solubility experiments and to demonstrate that the unexpected increase of solubility of fulvestrant by adding benzyl benzoate into compositions containing ethanol, benzyl alcohol and castor oil, is present across the broader range of composition encompassed by the claims being presented with this Declaration. The solubility of fulvestrant in benzyl benzoate and in castor oil was also measured in the same set of experiments using the same batch of benzyl benzoate and the same batch of castor oil as were used to make up the compositions. The Experimental Test Procedure is described in Attachment B.
8. The results from these solubility experiments are shown in the table in Attachment C. These results show that the solubility of fulvestrant in castor oil alone (21.4 mg/ml) is significantly greater than the solubility of fulvestrant in benzyl benzoate alone (3.8 mg/ml) and demonstrate the unexpected increase in fulvestrant solubility on the addition of 10, 15 and 25% w/v benzyl benzoate, in place of an equivalent amount of castor oil, to compositions having total alcohol concentrations in castor oil of 10%, 15%, 20%, 25% and 30% w/v.
9. Thus, the results that were obtained from experiments conducted under rigorously controlled conditions and with an expanded range of compositions, as shown in Attachment C, confirm the ultimate conclusions drawn from the results shown in Table 3 of the original application disclosure, namely that the addition of 10% to 25% w/v benzyl

benzoate to compositions having total alcohol concentrations in castor oil of between 10% to 30% w/v unexpectedly provides a positive effect on fulvestrant solubility, significantly increasing the solubility of fulvestrant in the compositions despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

10. During the course of my study of the Evans Application and the underlying source materials it was drawn to my attention that some of the composition data given for Delestrogen and Delalutin somehow had been shifted one column to the right. Thus, for Delestrogen, the 78% and 58% figures shown under the BzBz column should have been under the OIL column; the 20% and 40% figures shown under the BzOH column should have been under the BzBz column; and the 2% figures shown under EtOH should have been under the BzOH column. Similarly for Delalutin, the “up to 2%” shown under the EtOH column should have been under the BzOH column. This table reports that the source of this data was J.Pharm.Sci (1964) 53(8) 891, which is Riffkin (1964) elsewhere referred to in this Declaration, and I have also verified the corrected data from the entries for Delalutin and Delestrogen in PDR (1973). A copy of Table 1 from the Evans Application is reproduced as Attachment D, on which these corrections have been made in handwriting, and I have additionally more correctly noted that Delalutin is 17-hydroxy progesterone *caproate*, and that the “COMP” designation for Delalutin should be “BMS” (Bristol-Myers Squibb). Attachment D also includes a one page explanation of the corrections to this Table 1.

11. In about early 2000, a person responsible for developing a sustained release injectable formulation suitable for administration to humans for a new steroidal compound such as fulvestrant, would have had specialized training and experience in developing pharmaceutical formulations and methods for their administration. In developing such a formulation for fulvestrant, the objective would have been to formulate an intramuscular (IM) injection that would provide for the satisfactory sustained release of fulvestrant over a period of at least two weeks and preferably over a period of at least four weeks to reduce the frequency of administration, and would have a target fulvestrant content of at

least 45 mg/mL so as to provide a fulvestrant dose of at least 250 mg in a single 5-6 mL injection. From my personal experience and knowledge of the literature at about that time, I believe that such an experienced formulator would likely have approached the task of developing a formulation for fulvestrant in about the following manner.

12. Given the foregoing objective, the experienced formulator would have appreciated that the traditional administration options to explore were intramuscular (IM) injection of a sustained release aqueous or oil suspension or an oil-based solution (depot) containing at least 250 mg of fulvestrant in a volume of vehicle that is tolerable for injection, *i.e.*, no more than 5 or 6 mL.
13. Because of the extremely low solubility of fulvestrant in water, a reasonable starting point would have been to investigate intramuscular injection of an aqueous or oil suspension of fulvestrant. However, the formulator would have found that injection of an aqueous suspension of fulvestrant resulted in extensive local tissue irritation at the injection site as well as a poor release profile, such as reported in paragraph [0042] of the Evans Application. Since suspensions thus were not an acceptable option for fulvestrant, the experienced formulator would have moved on to further explore whether 250 mg of fulvestrant could be solubilised in no more than 5-6 mL of an oil-based vehicle, *i.e.*, to achieve the target fulvestrant concentration of at least 45 mg/mL.
14. In the preformulation phase, the experienced formulator would have conducted a literature review or otherwise would have become familiar with commercially marketed injectable formulations, particularly injectable sustained release formulations of steroids or other relatively insoluble compounds such as those listed in Table I of the Evans Application, with the objective of identifying potential oil vehicles, co-solvents and other excipients that already had been found to be tolerated and/or to have passed through regulatory review, and which might be candidates for further consideration and testing for the fulvestrant formulation. This review also would have provided guidance with respect to concentration levels of such co-solvents and other excipients that generally had been found acceptable in sustained release oil-based intramuscular injections administered to

humans. This objective is confirmed, for example, in Nema (1997) at page 166:

Generally, a knowledge of which excipients have been deemed safe by the FDA or are already present in a marketed product provides increased assurance to the formulator that these excipients will probably be safe for their new drug product. ... Regulatory bodies may view an excipient previously approved in an injectable dosage form favorably, and will frequently require less safety data.

The purpose of this Nema paper was thus “to present the various excipients that have been included in the formulation of injectable products marketed in the USA.”¹ Similar objectives were intended to be served by the compilations of commercial formulations in Strickley I (1999), Strickley II (2000) and Strickley III (2000):

This compilation will also be useful for those interested in knowing what additives are currently used in injectable products and at what concentrations they are administered in practice. This compilation only focuses on marketed formulations and does not delve into the subject of preclinical or drug discovery formulations associated with early-stages pharmacokinetics or proof-of-concept pharmacodynamics, where the formulation scientist is not bound by regulatory constraints.

(Strickley I (1999) at 324).

Powell (1998) similarly states at page 238 with respect to its compilation of commercially used excipients:

Thus, the formulation scientist is often faced with a dilemma -- which excipients are truly available for use (based on what has been used previously), and which are not? ... And at what concentrations, and by what route? ...

Herein are listed the excipients found in most of the approved and marketed parenteral formulations, given systematically by excipient name. In this format it is easy to determine what concentrations were used, the route of administration, the main rationale for addition of that excipient, the drug that was formulated, the manufacturer, brand name, etc.

15. From the literature review, the formulator would have noted reference to a number of intramuscular injectable sustained release oil-based steroidal formulations that had been

¹ Nema (1997) does caution, however, that there is no guarantee that the new drug product will be safe as excipients are combined with other additives and/or with a new drug, creating unforeseen potentiation or synergistic toxic effects.

commercially marketed:

- Strickley I (1999), Table VII:
 - Haloperidol Decanoate/Haldol decanoate (50-100 mg/mL in sesame oil, benzyl alcohol 1.2%);
 - Testosterone Enanthate/Delatestryl (200 mg/mL in sesame oil, chlorobutanol 5 mg/mL);
- PDR (1973) at pages 1277-1278
 - Proluton/progesterone (50 mg/mL in sesame oil, 150 mg/ml benzyl benzoate, 5 mg/ml benzyl alcohol, 1 mg/ml propylparaben);
- PDR (1973) at pages 1349-1354
 - Deladumone/Testosterone Enanthate & Estradiol Valerate (90 & 4 mg/mL in sesame oil, 0.5% chlorobutanol);
 - Deladumone OB/Testosterone Enanthate & Estradiol Valerate (180 & 8 mg/mL in sesame oil, 2% benzyl alcohol);
 - Delalutin/hydroxyprogesterone caproate (250 mg/mL in 52% castor oil, 46% benzyl benzoate, 2% benzyl alcohol);
 - Delestrogen/estradiol valerate (20 mg/mL in 78% castor oil, 20% benzyl benzoate, 2% benzyl alcohol and 40 mg/mL in 58% castor oil, 40% benzyl benzoate, 2% benzyl alcohol);
 - Delatestryl/Testosterone Enanthate (200 mg/mL in sesame oil, 0.5% chlorobutanol);
 - Delalutecval 2X/hydroxyprogesterone caproate & estradiol valerate (250 mg/mL & 5 mg/mL in castor oil, 45% benzyl benzoate, 1.6% benzyl alcohol);
- PDR (1973) at pages 1391-1392
 - Prolixin Enanthate/FluphenazineEnanthate (25 mg/mL in sesame oil, 1.5% benzyl alcohol);
- Wang (1980):
 - Depo-Testosterone/testosterone cypionate (100 mg/mL in 87.4% cottonseed oil, 0.1 mL benzyl benzoate, 9.45 mg benzyl alcohol as a preservative);
- Mackey (1995):
 - Testoviron Depot/testosterone enanthate (250 mg/mL in castor oil and benzyl

benzoate);

as well as a number of other commercialized oil based long-acting IM injectable formulations reported on Table 1 of the Evans Application.

16. As a further part of the preformulation phase, the experienced formulator would have conducted a preformulation solubility screen, separately measuring the solubility of fulvestrant in a range of pure solvents, including the potential oil and co-solvent candidates that had been identified in the above literature review as being suitable for inclusion in intramuscular injection formulations. See, for example, Gupta (1999), Chapter 17 at page 402, under the heading “Formulation Development”:

The activities necessary to develop a parenteral product can be placed into the following three broad areas: preformulation, formulation, and scale-up. While there are alternative development perspectives, all development ultimately needs to accomplish the same activities. Preformulation includes the characteristics of the bulk drug plus initial screening for excipient compatibility with the drug.

“Preformulation studies” are said to “provide fundamental data and experience necessary to develop formulations for a specific compound” including, as item 8.1 in the outline of areas of specific interest, a determination of “solubility” in “selected solvents” (at 403). “Significant formulation activities begin with initial preformulation data and knowledge of the specific route of administration” (at 405), which “formulation activities include the identification and selection of a suitable vehicle (aqueous, nonaqueous or co-solvent system) ...” (at 404). It is further noted that “injection volume is one of the most important considerations in the formulation development of a commercial product” (at 405). When carrying out such a preformulation solubility screen with fulvestrant, the formulator would have found that fulvestrant had extremely low solubility in water, low solubility in most oils (but highest in castor oil), low solubility in benzyl benzoate, and the highest solubility in ethanol and benzyl alcohol, such as reported in Table 2 of the Evans Application.

17. With the information on prior commercialized formulations and the fulvestrant solubility data from the preformulation screen (such as reported in Table 2 of the Evans

Application), the experienced formulator would have selected castor oil as the oil vehicle because of the higher solubility of fulvestrant in castor oil relative to the other oils tested. Nevertheless, he would have appreciated that the target fulvestrant concentration of at least 45 mg/mL could not be achieved with castor oil alone, and that a co-solvent would be required.

18. A number of the commercialized formulations that would have been identified in the literature review (including the castor oil-based formulations) have a substantial benzyl benzoate component, which may be present as a co-solvent. See, for example, Delalutin noted in paragraph 15 above, which is reported in PDR (1973) and noted in Table I of the Evans Application, and is one of the formulations discussed in Riffkin (1964), "Castor Oil as a Vehicle for Parenteral Administration of Steroid Hormones" (see Riffkin n. 6). Delalutin is 250 mg/mL 17-hydroxyprogesterone caproate dissolved in 52% castor oil, 46% benzyl benzoate and 2% benzyl alcohol. However, Riffkin Table II reports that the solubility of 17-hydroxyprogesterone caproate in castor oil alone is only 55.6 mg/mL, but the solubility of 17-hydroxyprogesterone caproate in benzyl benzoate is substantially higher, being at least 250 mg/mL (see example 4 of Huber (US '520) and Attachment E discussed below). Even if not needed as a cosolvent, Riffkin (1964) notes that "the addition of benzyl alcohol or benzyl benzoate to castor oil resulted in a lower and more favorable viscosity, making it easier to inject" (paragraph bridging pages 893-894).

19. However, the skilled formulator would have appreciated from the fulvestrant solubility data generated in the preformulation screen that fulvestrant had very different solubility characteristics relative to the steroids of previous commercial formulations. Attachment E is a compilation showing the chemical structures and relative solubilities in castor oil and sesame oil of the compounds named in Riffkin (1964) Table II compared to the structure and the solubility of fulvestrant in these oils. It can be seen that the solubility of fulvestrant in castor oil and in sesame oil (20 mg/mL and 0.58 mg/mL, respectively, from Table 2 of the Evans Application) is appreciably lower than the solubility of the other steroids in these oils (taken from Table II of Riffkin (1964)). The second page of Attachment E tabulates the concentration in benzyl benzoate of five named steroids, taken

from Examples 1-5 of Huber (US '520), ranging from 200 to 400 mg/mL.² By comparison, the solubility of fulvestrant in benzyl benzoate is reported in Table 2 of the Evans Application as being only 6.15 mg/mL, and only 3.8 mg/mL as determined in the recently conducted tests reported in Attachment C.

20. The experienced formulator thus would have expected that benzyl benzoate would *not* act as a co-solvent for fulvestrant in castor oil because the solubility of fulvestrant in benzyl benzoate was significantly lower than its solubility in castor oil. The addition of benzyl benzoate to castor oil, for whatever reason, would have been expected to *decrease, rather than increase*, the solubility of fulvestrant in the resulting castor oil/benzyl benzoate mixture. This is confirmed in Table 4 of the Evans Application, which reports a fulvestrant solubility of only 12.6 mg/mL in the castor oil vehicle containing only 15% benzyl benzoate, compared to the 20 mg/mL solubility of fulvestrant in castor oil alone as reported in Table 2.³
21. Based on the solubility data determined in the preformulation screen (such as reported in Table 2 of the Evans Application), ethanol and/or benzyl alcohol would have been seen as the best co-solvent candidates for raising the fulvestrant solubility to the 45 mg/mL target in the castor oil vehicle, and would also function to lower the viscosity of the resulting formulation and make it easier to inject. Consistent with this solubility data, Dukes (US '814) added 40% w/v benzyl alcohol in order to dissolve 50 mg/mL fulvestrant in the castor oil-based formulation used in the experimental rat studies of his Example 3. It thus would have been apparent that 40% w/v benzyl alcohol could function as a co-solvent in castor oil to achieve the target fulvestrant concentration. Nevertheless, the skilled formulator would have been concerned with using such a high alcohol content in intramuscular injectable formulations for administration to a human.

² Data taken from the Examples of Huber (US '520); these are concentrations used in the examples and not necessarily the actual maximum solubility of each steroid in benzyl benzoate, which may be higher. Huber was a co-author on Riffkin (1964).

³ It should be noted that in the further tests that were recently conducted under my guidance (paragraphs 7-9 above and Attachments B and C hereto), the solubility of fulvestrant in castor oil alone was again tested and found to be 21.4 mg/mL, and the solubility of fulvestrant in benzyl benzoate alone was again tested and found to be only 3.8 mg/mL, which further confirms that benzyl benzoate would not be expected to act as a cosolvent for fulvestrant in castor oil.

22. First of all, the experienced formulator would want to minimize the amount of co-solvents and excipients in any injectable formulation. For example, as stated in Gupta (1999), Chapter 17, "Formulation and Administration Techniques to Minimize Injection Pain and Tissue Damage Associated with Parental Products" at page 414:

Cosolvents are commonly used to enhance drug solubility and stability. Cosolvents may include ethanol, propylene glycol, polyethylene glycols, and glycerine. These components have intrinsic effects on biologic tissue and can alter the properties of other excipients, thus influencing the tissue damage or pain caused by a product. There is a dearth of literature on the pain caused by cosolvents, but there is also a growing body of knowledge on the tissue damage that they can cause. It is not certain that tissue damage is always directly correlated with the injection pain, but minimization of both pain on injection and potential for tissue damage should be included in the product development plan.

See also Gupta (1999), Chapter 11, titled Cosolvent Use in Injectable Formulations, page 217:

Ideally, it is best to select and use solvents that would maximize the solubility of the compound. Maximizing the solubility of a compound in a particular cosolvent system would result in lower total levels of the non-aqueous solvent(s) being administered to the patient, thereby lowering the chance for potential side effects.

This objective would have applied to aqueous and oil-based systems alike, in that the precedent of commercialized formulations identified in the literature review would have confirmed that fixed oils, such as castor oil, have long been commercially used and accepted as the major component of oil-based sustained release intramuscular injectable steroidal formulations. On the other hand, co-solvents such as ethanol or benzyl alcohol have generally been used only in far lesser concentrations, as discussed in the following paragraph.

23. Thus, use of such a high content of either benzyl alcohol or ethanol would have been contrary to precedent as shown from the review of commercialized oil-based intramuscular injectable sustained release formulations. The literature review as of early 2000 would have shown that any benzyl alcohol in such formulations was almost always

present as a preservative in a concentration of about 2% or less, occasionally at a concentration of up to 5%, but only rarely at higher concentrations. With respect to benzyl alcohol see, for example:

- Gupta (1999), Chapter 11 at page 229 stating that benzyl alcohol “is typically used in concentrations of up to 2 percent as a preservative and up to 5 percent as a solvent,” and then discussing reported toxicities.
- Nema (1997), Table V at page 168, reporting that benzyl alcohol was present as an antimicrobial preservative in 74 injectable formulations (not limited to oil-based IM formulations) at concentrations of from 0.75-5% (note that benzyl alcohol is not included at all in Nema Table I, “Solvents and Co-solvents”);
- Powell (1998), the benzyl alcohol listing at pages 244-246, particularly those indicated as being used in IM formulations;
- Strickley I (1999) at page 329 notes the inclusion of 2% benzyl alcohol in an IM lorazepam formulation in a propylene glycol vehicle, but does not include benzyl alcohol at all in Table VI listing “Cosolvents Used in Parenteral Formulations;”
- Lopatin (1972) noting in Table 3 at page 727 opposite Benzyl alcohol, “Toxic. Used in concentration of not over 3%. Has irritant action in concentration of 5%;”
- Cornelius (US ‘863), col. 1, lines 30-35 stating, “It is known that the solubility of steroids in vegetable or animal oils can be increased by the addition of excipients such as benzyl alcohol and benzyl benzoate. An objection to the use of such excipients, and specifically benzyl alcohol in somewhat higher concentrations, is that these agents may irritate the tissues.”

The literature review as of early 2000 also would have shown that, with few exceptions, ethanol was not included in such formulations in excess of about 10%. See, for example:

- Gupta (1999), Chapter 11 at page 225 noting that ethanol has been used at levels up to 50 percent, but these levels typically are associated with pain on injection;
- Strickley I (1999), Table VI, “List of Cosolvents Used in Parenteral Formulations” more specifically lists the ethanol content in IM formulations for specifically identified drugs, which concentrations range only from 2.5 to 10%; an IM/IV lorazepam formulation in a propylene glycol vehicle is noted at page 329 as having 18% alcohol, but is not included with the IM formulations in Table VI;

- Nema (1997), Table I, “Solvents and Co-solvents” at page 167, lists ethanol as being in 24 formulations with a concentration range of 0.6-80% (for Prograf); note that this is misleading, however, since Prograf is a *concentrate* for intravenous infusion only, and is to be diluted 250 to 1000 times before administration;
- Powell (1998), lists “alcohol” at page 242 and “ethyl alcohol” at page 255, wherein the ethanol concentration for IM formulations ranges from 0.61-10%.

24. Thus, even though Dukes (US '814) had demonstrated that the target 45 mg/mL fulvestrant concentration could be achieved by adding 40% benzyl alcohol to the castor oil vehicle, the precedent of commercialized IM oil-based systems would have motivated the experienced formulator to substantially reduce the benzyl alcohol content of the formulation intended for human use, and this commercial precedent would have made him very reluctant to replace benzyl alcohol with the substantial amount of ethanol that would be needed to maintain the target fulvestrant concentration. Benzyl benzoate clearly would not be considered to solve this dilemma, but rather would be expected to have a negative effect on fulvestrant solubility since fulvestrant was even less soluble in benzyl benzoate than in castor oil, that is, one would have expected that adding benzyl benzoate would require still *more* alcohol to maintain the target fulvestrant concentration.⁴

25. Under these circumstances, the discovery by Evans *et al.*, that the addition of benzyl benzoate to the castor oil/alcohol mixture actually increases the solubility of fulvestrant such that more fulvestrant could be dissolved in a given volume of formulation, was unexpected and truly surprising. This positive benzyl benzoate effect on fulvestrant solubility in the resulting formulation is shown in Table 3 of the specification (and is not changed by the above-noted corrections), and is confirmed and demonstrated over a broader range of formulation composition by the additional set of experiments conducted under my guidance and discussed in paragraphs 7-9 above, the results of which are reported in Attachments C.

⁴ It should be noted that even apart from this solubility issue, there would have been no motivation to add benzyl benzoate for viscosity reduction since the significant quantity of alcohol would serve the dual function of acting as a co-solvent as well as reducing the injection viscosity and making it easier to inject, whereas the benzyl benzoate would be expected to have a negative effect on the fulvestrant solubility.

The undersigned declares further that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punished by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardise the validity of the application or any patent issuing thereon.

Signature: P. R. Markel.

Date: 8th August 2008.

TABLE OF REFERENCES

Tab	Author/Inventor	Reference Citation/Patent
1	Cornelius (US '863)	US Patent 4,212,863
2	Dukes (EP '014)	EP 0 346 014 A1 (corresponds to US Patent 5,183,814)
3	Dukes (US '814)	US Patent 5,183,814 (corresponds to EP 0 346 013 A1)
4	Gupta (1999)	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
5	Huber (US '520)	US Patent 3,164,520
6	Lopatin (1972)	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
7	Mackey (1995)	M.A. Mackey, A.J. Conway and D.J. Handelsman. Tolerability of intramuscular injections of testosterone ester in oil vehicle. <i>Hum. Reprod.</i> 10 : 862-865 (1995)
8	Nema (1997)	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
9	PDR (1973)	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
10	Powell (1998)	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 (1998)
11	Riffkin (1964)	C. Riffkin, R. Huber and C.H. Keysser. Castor oil as a vehicle for parenteral administration of steroid hormones. <i>J. Pharm. Sci.</i> 53 : 891-5 (1964)
12	Strickley I (1999)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
13	Strickley II (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
14	Strickley III (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
15	Wang (1980)	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

ATTACHMENT A

TABLE 3

		EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C							
		4°C							
		% w/v							
	Ethanol (96%)	5	5	10	10	10	10	15	15
	Benzyl Alcohol	5	5	5	5	10	10	15	15
	Benzyl Benzoate		15		15		15		15
Mean	Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
	Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	68 64	76 77	102 103
Individual values		27.8	35.5	54.4	64.1	48.4	68.9	65.8	80.6
		25.8	36.1	38.6	47.3	61.3	60.2	76.9	101.9
					53.0		63.2	90.0	121.6
					80.3		72.9	73.4	107.4

ATTACHMENT B:

Experimental Test Procedure for measuring the solubility of fulvestrant in different solvent vehicles at 25°C

1. Solvent vehicles for the solubility experiments were prepared by weighing the required amount of benzyl benzoate, benzyl alcohol and ethanol into a 20 ml volumetric flask and then diluting to volume with castor oil.
2. For each solvent vehicle in which the solubility of fulvestrant was to be determined, 1.0-1.5g of fulvestrant was weighed into each of 3 separate vials (2 dram size) and 5mls of the solvent vehicle was added to each vial, except for the pure castor oil vehicle, where 80mg of fulvestrant were weighed into each of the 3 separate vials and 2mls of the castor oil added to each vial. The reduced amount of fulvestrant and lower volume of solvent vehicle was needed to maintain stirring and achieve adequate mixing with the pure castor oil vehicle due to the combination of its higher viscosity and lower fulvestrant solubility/higher undissolved fulvestrant levels compared to the other solvent vehicles.
3. A magnetic stirrer bar was placed into each vial and the vials were capped and then placed on a magnetic stirrer block maintained at $25 \pm 0.5^{\circ}\text{C}$.
4. After 5 days of stirring at $25 \pm 0.5^{\circ}\text{C}$, an aliquot of each fulvestrant/solvent vehicle mixture was removed from each vial and placed into an Eppendorf tube which was then centrifuged at 12000 rpm for 5 minutes at ambient temperature.
5. For all but the fulvestrant/castor oil mixture, 1 ml of the supernatant was then removed from the Eppendorf tube and pipetted into a 10ml or 20ml volumetric flask and then diluted to volume with methanol and mixed to give a sample for analysis. The choice of whether to use a 10ml or 20ml volumetric flask for a particular sample was dependent on the likely concentration of fulvestrant in the sample and the quantifiable concentration range of the HPLC assay method used. For the fulvestrant/castor oil mixture, 100 μl of the supernatant was removed from the Eppendorf tube and pipetted into a 1ml volumetric flask and then diluted to volume with methanol and mixed to give a sample for analysis.
6. Step 5 was repeated to give a duplicate sample for analysis. Thus, this gave 2 samples for each of the 3 vials, giving a total of 6 samples for analysis for each solvent vehicle tested.
7. The resultant samples were analysed for fulvestrant content by reverse phase High

Performance Liquid Chromatography (HPLC). The HPLC method that was used is described below at point 9. The fulvestrant content obtained for each sample was used to calculate a value for the concentration of fulvestrant dissolved in the corresponding solvent vehicle after stirring for 5 days at 25°C.

8. The mean solubility of fulvestrant for each different solvent vehicle tested was calculated as the arithmetic mean of the 6 individual values for the concentration of fulvestrant dissolved in the corresponding solvent vehicle.

9. HPLC Method details:

Gradient HPLC Method

Eluent A : 27% Methanol / 32% Acetonitrile / 41% Water

Eluent B : 41% Methanol / 49% Acetonitrile / 10% Water

Column : 15cm 3.5um Symmetry C8 4.6mm i.d.

Detection wavelength : 225 nm

Flow rate : 2 mL min⁻¹

Temperature : 40°C

Injection volume : 10 µL

Gradient programme :

Time (min)	Eluent A (%)	Eluent B (%)
0	100	0
25	100	0
55	0	100
65	0	100
66	100	0
70	100	0

Retention time of fulvestrant: 21 minutes approximately

ATTACHMENT C:

EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C

		% w/v																				
Ethanol (96%)	0	0	5	5	5	5	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Benzyl Alcohol	0	0	5	5	5	5	5	5	5	5	10	10	10	10	12.5	12.5	12.5	12.5	12.5	12.5	12.5	12.5
Benzyl Benzoate	0	100		10	15	25		10	15	25		10	15	25		10	15	25		10	15	25
Castor oil	100	0	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Mean Fulvestrant solubility [mgml ⁻¹]	21.4	3.8	27.6	29.2	43.3	47.5	64.6	71.6	84.2	94.0	68.1	87.2	93.4	118.9	96.6	107.7	116.1	139.6	121.3	144.6	143.8	166.2
Individual values [mgml ⁻¹]	23.2	3.9	29.5	31.2	43.9	48.3	64.2	76.2	83.8	95.2	68.6	90.0	92.5	122.1	104.1	106.1	115.5	138.9	110.0	129.8	148.2	163.3
	17.8	4.0	28.3	26.3	45.1	50.7	66.8	72.1	81.9	97.8	68.9	84.9	92.1	120.3	74.0	86.6	117.9	141.0	120.0	133.5	147.1	164.8
	21.5	3.9	24.5	31.5	44.3	45.4	61.2	66.2	93.2	95.6	71.6	87.6	93.9	120.4	102.0	112.6	118.8	139.4	124.4	150.1	144.4	168.5
	21.8	3.8	26.6	29.3	45.4	45.2	66.0	65.7	84.6	96.1	67.6	88.1	93.0	118.3	98.6	117.9	116.1	142.1	125.6	151.7	144.4	169.7
	22.2	4.0	27.0	29.1	36.9	47.6	65.8	75.4	82.4	88.2	67.0	90.7	93.8	116.8	102.1	107.9	117.0	138.7	123.3	151.2	139.5	165.5
	22.0	3.2	29.6	27.8	44.3	47.6	63.6	73.9	79.1	91.0	64.8	82.1	95.3	115.7	98.4	115.1	111.5	137.9	124.6	151.1	139.1	165.5

ATTACHMENT D

TABLE 1

OIL-BASED LONG-ACTING ESTROGENIC INJECTIONS										
PRODUCT NAME	STEROID	DOSE	TYPE	CLASS ¹	SOURCE	OH ²	Safe	Safe	Exact	Other Details
METHAMON 100	Dexamethasone propionate	20 mg	Androgen	Organic	ABPI Data	Astra	-	-	-	1 ml 3 weeks
	Testosterone phenylpropionate	60 mg								
	Dexamethasone isocaproate	60 mg								
	Testosterone decanoate	100 mg								
PROGESTIN DEPOT	Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	Schering BC	ABPI Data	Cantor	up to 40%	-	-	1 or 2 ml 2 week
	Testosterone	100 mg								2 ml
TODIGESTAN	Hydroxyprogesterone succinate	300 mg	Progestin	Diamant	Dist. Vidal 1999	Vidal	-	-	-	2 ml 4-6 weeks
	Progesterone	50 mg or 150 mg								
TROPICOLIMS	Enoxolone	1.5 mg	Mixed	Diamant	Dist. Vidal 1997	Off-c	40%	-	-	1 ml 10 to 30 days
	Methocort	50 mg								
	Hydroxyprogesterone laurate	80 mg								
NOBENTIN	Mestranolone succinate	200 mg	Contraceptive	Schering BC	ABPI Data	Cantor	YES	-	1 ml 8 weeks	
BENZO-ETHYNYL	Retarded benzylpropionate	5 mg	Retarded	Bioss	Dist. Vidal 1998	Astra	-	-	1 ml 3 week	
PROGESTERONE-RETARD	Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	Fisher	Dist. Vidal 1999	Cantor	YES	-	1 or 2 ml 1 week	
ORAVIBIAN	Retarded 17- β -valerate	5 mg/ml ³	Mixed	Schering BC	Dist. Vidal 1998	Cantor	YES	-	-	1 or 2 ml
	Hydroxyprogesterone caproate	250 mg/ml ³								2 ml 1 week
PARASOLAN	Decalone	76 mg	Androgen	Negon	Dist. Vidal 1997	Astra	77 mg	40 mg	1.5 ml 2 weeks	
OIL ESTROGEN	Retarded valerate	20 mg/ml ³ 40 mg/ml ³	Retarded	BMS	I Pharm. (1964) S(19) 491	Cantor	74 88	10% 10%	77 88	1 or 2 ml
	Hydroxyprogesterone caproate	250 mg/ml ³								
OBLALLEN	17-Hydroxyprogesterone caproate	250 mg/ml ³	Progestin	BMS	I Pharm. (1964) S(19) 491	Cantor	YES	YES	-	up to 2 ml

OH₂ = benzylbenzene
 BC = benzylbenzene
 OH₁ = ethyl ester
 Vidal = Dist. Vidal % are w/w and
³mg/ml = mg/ml directly from a single sample

Corrections to Table 1

In Table 1, the given values for the benzyl benzoate, benzyl alcohol and ethanol levels for the Delestrogen and Delalutin products have been incorrectly entered into the wrong columns. The entries are shown in their correct form in the attached corrected version of Table 1. The error is apparent from a review of the reference J.Pharm Sci (1964) 53 (8) 891 (Riffkin) which is stated in Table 1 as being the Source of the information for the Delestrogen and Delalutin products:

- In the Summary on page 895 of Riffkin, Delestrogen and Delalutin are identified as castor oil based commercially available products containing estradiol valerate at 20 & 40 mg/ml and 17-hydroxy-progesterone caproate at 250 mg/ml respectively.
- Furthermore, details of particular vehicle compositions for estradiol valerate and 17-hydroxy-progesterone caproate are given in Tables V and VI
 - In Table VI, the only 20 mg/ml formulation of estradiol valerate, also referred to as commercially available, has the composition castor oil 78%, benzyl benzoate 20% and benzyl alcohol 2%.
 - In Table VI, the only 40 mg/ml castor oil based formulation of estradiol valerate, has the composition castor oil 58%, benzyl benzoate 40% and benzyl alcohol 2%.
 - In Table V, there are three 250/mg/ml castor oil based formulations of 17-hydroxy-progesterone caproate that all contain benzyl benzoate. Two of these formulations also contain 2% benzyl alcohol and the other formulation does not contain benzyl alcohol ie they all contain up to 2% benzyl alcohol.
- None of the vehicle compositions disclosed in Tables V and VI in Riffkin contain ethanol. Therefore the entries in the Ethanol column of Table 1 for the Delestrogen and Delalutin products must have been incorrectly entered in the wrong column and should have been entered into the Benzyl Alcohol column.
- It is also apparent from Table VI that the 78% and 58% entries in the Benzyl Benzoate column of Table 1 for the Delestrogen products should have been entered into the Oil column and the 20% and 40% entries in the Benzyl Alcohol column should have been entered into the Benzyl Benzoate column
- The exact compositions for the Delestrogen and Delalutin products are confirmed in the Physicians Desk Reference (Edition 27, 1973) on page 1352.

In addition, the name of the steroid given in Table 1 for the Delalutin product should have been 17-hydroxy-progesterone caproate and not just 17-hydroxy-progesterone. Also the entry under the Company column for the same product should read BMS rather than DMS.

ATTACHMENT E

Structure of compounds disclosed in Riffkin et al.


17-Hydroxypregesterone caproate:




Testosterone:




Estradiol valerate:



Pregesterone:



On the other hand, fulvestrant has the following structure:



From Riffkin et al. Table II:

Steroid	Solubility [mg/ml] at 25°C	
	Castor oil	Sesame oil
Fulvestrant	28	0.38
17-Hydroxypregesterone caproate	55.5	23.4
Testosterone	38.6	5.4
Estradiol valerate	60.6	16.1
Pregesterone	52.0	22.9

Tabulation of data from Examples of Huber, 3,164,520:

Example	Steroid	Steroid concentration in benzyl benzoate (mg/ml)
1	16,17-dihydroxyprogesterone	200
2	testosterone palmitate	200
3	progesterone	250
4	Progesterone + 17-hydroxyprogesterone caproate	250 + 250
5	Testosterone enanthate	400

ATTACHMENT F

TABLE OF REFERENCES

Tab	Author/Inventor	Reference Citation/Patent
1	Cornelius (US '863)	US Patent 4,212,863
2	Dukes (EP '014)	EP 0 346 014 A1 (corresponds to US Patent 5,183,814)
3	Dukes (US '814)	US Patent 5,183,814 (corresponds to EP 0 346 013 A1)
4	Gupta (1999)	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
5	Huber (US '520)	US Patent 3,164,520
6	Lopatin (1972)	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
7	Mackey (1995)	M.A. Mackey, A.J. Conway and D.J. Handelsman. Tolerability of intramuscular injections of testosterone ester in oil vehicle. <i>Hum. Reprod.</i> 10 : 862-865 (1995)
8	Nema (1997)	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
9	PDR (1973)	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
10	Powell (1998)	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 [pages 238-255 provided] (1998)
11	Riffkin (1964)	C. Riffkin, R. Huber and C.H. Keysser. Castor oil as a vehicle for parenteral administration of steroid hormones. <i>J.Pharm.Sci.</i> 53 : 891-5 (1964)
12	Strickley I (1999)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
13	Strickley II (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
14	Strickley III (2000)	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
15	Wang (1980)	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

ATTACHMENT F

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875	Application or Docket Number 12/285,887	Filing Date 10/15/2008	<input type="checkbox"/> To be Mailed
---	---	----------------------------------	---------------------------------------

APPLICATION AS FILED – PART I			OTHER THAN SMALL ENTITY				
(Column 1)		(Column 2)	SMALL ENTITY <input type="checkbox"/>		OR	SMALL ENTITY	
FOR	NUMBER FILED	NUMBER EXTRA	RATE (\$)	FEE (\$)		RATE (\$)	FEE (\$)
<input type="checkbox"/> BASIC FEE <small>(37 CFR 1.16(a), (b), or (c))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> SEARCH FEE <small>(37 CFR 1.16(k), (l), or (m))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> EXAMINATION FEE <small>(37 CFR 1.16(o), (p), or (q))</small>	N/A	N/A	N/A			N/A	
TOTAL CLAIMS <small>(37 CFR 1.16(j))</small>	minus 20 =	*	X \$ =		OR	X \$ =	
INDEPENDENT CLAIMS <small>(37 CFR 1.16(h))</small>	minus 3 =	*	X \$ =			X \$ =	
<input type="checkbox"/> APPLICATION SIZE FEE <small>(37 CFR 1.16(s))</small>	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).						
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENT <small>(37 CFR 1.16(j))</small>							
* If the difference in column 1 is less than zero, enter "0" in column 2.			TOTAL			TOTAL	

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY			
(Column 1)		(Column 2)	(Column 3)	SMALL ENTITY		OR	SMALL ENTITY	
AMENDMENT	01/17/2012	CLAIMS REMAINING AFTER AMENDMENT	MINUS	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA		RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	* 20	Minus	** 30	= 0		X \$ =	
	Independent <small>(37 CFR 1.16(h))</small>	* 2	Minus	*** 8	= 0		X \$ =	
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>							
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>							
							TOTAL ADD'L FEE	0

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY			
(Column 1)		(Column 2)	(Column 3)	SMALL ENTITY		OR	SMALL ENTITY	
AMENDMENT		CLAIMS REMAINING AFTER AMENDMENT	MINUS	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA		RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	*	Minus	**	=		X \$ =	
	Independent <small>(37 CFR 1.16(h))</small>	*	Minus	***	=		X \$ =	
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>							
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>							
							TOTAL ADD'L FEE	

* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.
 ** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".
 *** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".
 The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.

Legal Instrument Examiner:
/DESHONNE T. MARTINO/

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**
 If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

AMENDMENTS TO THE CLAIMS

Please amend claims 24, 32, 34, 36, 44, and 46. Please add new claims 54-57. Please cancel claims 25, 28, 31, 33, 37, 40, 43, 45, and 48-53 without prejudice or disclaimer. This listing of claims will replace all prior versions and listings of claims in the application.

Claims 1-23 (Cancelled)

24. (Currently amended) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation comprising:

~~at least 45 mgml⁻¹ of fulvestrant;~~

~~a mixture of from 17 – 23% w/v of ethanol and benzyl alcohol;~~

~~12 – 18% w/v of benzyl benzoate; and~~

about 50 mgml⁻¹ of fulvestrant;

about 10% w/v of ethanol;

about 10% w/v of benzyl alcohol;

about 15% w/v of benzyl benzoate; and

a sufficient amount of castor oil vehicle;

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least ~~two~~ four weeks.

25. (Cancelled)

26. (Previously presented) The method of claim 24, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml^{-1} .
27. (Previously presented) The method of claim 24, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
28. (Cancelled)
29. (Previously presented) The method of claim 24, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
30. (Previously presented) The method of claim 24, wherein the method further comprises once monthly administration of the formulation.
31. (Cancelled)
32. (Currently amended) The method of ~~claim 31~~ claim 26, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
33. (Cancelled)
34. (Currently amended) The method of ~~claim 33~~ claim 32, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.

35. (Previously presented) The method of claim 34, wherein the method further comprises once monthly administration of the formulation.
36. (Currently amended) A method for treating a hormonal dependent benign or malignant disease of the breast or reproductive tract comprising administering intramuscularly to a human in need of such treatment a formulation consisting essentially of:

~~at least 45 mgml⁻¹ of fulvestrant;~~
~~a mixture of from 17—23% w/v of ethanol and benzyl alcohol;~~
~~12—18% w/v of benzyl benzoate; and~~
~~a sufficient amount of castor oil vehicle;~~
about 50 mgml⁻¹ of fulvestrant;
about 10% w/v of ethanol;
about 10% w/v of benzyl alcohol;
about 15% w/v of benzyl benzoate; and

wherein the method achieves a therapeutically significant blood plasma fulvestrant concentration of at least 2.5 ngml⁻¹ for at least ~~two~~four weeks.

37. (Cancelled)
38. (Previously presented) The method of claim 36, wherein the therapeutically significant blood plasma fulvestrant concentration is at least 8.5 ngml⁻¹.

39. (Previously presented) The method of claim 36, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
40. (Cancelled)
41. (Previously presented) The method of claim 36, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
42. (Previously presented) The method of claim 36, wherein the method further comprises once monthly administration of the formulation.
43. (Cancelled)
44. (Currently amended) The method of ~~claim 43~~ claim 38, wherein the hormonal dependent benign or malignant disease of the breast or reproductive tract is breast cancer.
45. (Cancelled)
46. (Currently amended) The method of ~~claim 45~~ claim 44, wherein the method comprises administering intramuscularly to a human in need of such treatment 5 mL of the formulation.
47. (Previously presented) The method of claim 46, wherein the method further comprises once monthly administration of the formulation.

Claims 48-53 (Cancelled)

54. (New) The method according to claim 24, wherein the formulation is administered in a divided dose.
55. (New) The method according to claim 35, wherein the formulation is administered in a divided dose.
56. (New) The method according to claim 36, wherein the formulation is administered in a divided dose.
57. (New) The method according to claim 47, wherein the formulation is administered in a divided dose.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 5 columns: APPLICATION NO., FILING DATE, FIRST NAMED INVENTOR, ATTORNEY DOCKET NO., CONFIRMATION NO.
12/285,887 10/15/2008 John R. Evans 11285.0056-00000 1199

22852 7590 03/20/2012
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

EXAMINER

HUI, SAN MING R

Table with 2 columns: ART UNIT, PAPER NUMBER

1628

Table with 2 columns: MAIL DATE, DELIVERY MODE

03/20/2012

PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Office Action Summary	Application No. 12/285,887	Applicant(s) EVANS ET AL.	
	Examiner SAN-MING HUI	Art Unit 1628	

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) OR THIRTY (30) DAYS, WHICHEVER IS LONGER, FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) Responsive to communication(s) filed on 17 January 2012.
- 2a) This action is **FINAL**. 2b) This action is non-final.
- 3) An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.
- 4) Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

Disposition of Claims

- 5) Claim(s) 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47 and 54-57 is/are pending in the application.
- 5a) Of the above claim(s) _____ is/are withdrawn from consideration.
- 6) Claim(s) _____ is/are allowed.
- 7) Claim(s) 24, 26-27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46-47, and 54-57 is/are rejected.
- 8) Claim(s) _____ is/are objected to.
- 9) Claim(s) _____ are subject to restriction and/or election requirement.

Application Papers

- 10) The specification is objected to by the Examiner.
- 11) The drawing(s) filed on _____ is/are: a) accepted or b) objected to by the Examiner.
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
- 12) The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

Priority under 35 U.S.C. § 119

- 13) Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
a) All b) Some * c) None of:
1. Certified copies of the priority documents have been received.
2. Certified copies of the priority documents have been received in Application No. _____.
3. Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

* See the attached detailed Office action for a list of the certified copies not received.

Attachment(s)

- 1) Notice of References Cited (PTO-892)
- 2) Notice of Draftsperson's Patent Drawing Review (PTO-948)
- 3) Information Disclosure Statement(s) (PTO/SB/08)
Paper No(s)/Mail Date 6/2011, 1/17/12
- 4) Interview Summary (PTO-413)
Paper No(s)/Mail Date. _____
- 5) Notice of Informal Patent Application
- 6) Other: _____

DETAILED ACTION

Continued Examination Under 37 CFR 1.114

A request for continued examination under 37 CFR 1.114, including the fee set forth in 37 CFR 1.17(e), was filed in this application after final rejection. Since this application is eligible for continued examination under 37 CFR 1.114, and the fee set forth in 37 CFR 1.17(e) has been timely paid, the finality of the previous Office action has been withdrawn pursuant to 37 CFR 1.114. Applicant's submission filed on 1/17/2012 has been entered.

Claims 24, 26-27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46-47, and 54-57 are pending.

The outstanding rejection under 35 USC 103(a) is withdrawn in view of the arguments along with the declaration of Dr. Sawchuk filed 1/17/2012.

However, in view of the decision dated 2/15/2012 with regard to the terminal disclaimer, the obviousness double patenting rejections are maintained with the new added claims are also rejected.

Double Patenting

The nonstatutory double patenting rejection is based on a judicially created doctrine grounded in public policy (a policy reflected in the statute) so as to prevent the unjustified or improper timewise extension of the "right to exclude" granted by a patent and to prevent possible harassment by multiple assignees. A nonstatutory obviousness-type double patenting rejection is appropriate where the conflicting claims

Art Unit: 1628

are not identical, but at least one examined application claim is not patentably distinct from the reference claim(s) because the examined application claim is either anticipated by, or would have been obvious over, the reference claim(s). See, e.g., *In re Berg*, 140 F.3d 1428, 46 USPQ2d 1226 (Fed. Cir. 1998); *In re Goodman*, 11 F.3d 1046, 29 USPQ2d 2010 (Fed. Cir. 1993); *In re Longi*, 759 F.2d 887, 225 USPQ 645 (Fed. Cir. 1985); *In re Van Ornum*, 686 F.2d 937, 214 USPQ 761 (CCPA 1982); *In re Vogel*, 422 F.2d 438, 164 USPQ 619 (CCPA 1970); and *In re Thorington*, 418 F.2d 528, 163 USPQ 644 (CCPA 1969).

A timely filed terminal disclaimer in compliance with 37 CFR 1.321(c) or 1.321(d) may be used to overcome an actual or provisional rejection based on a nonstatutory double patenting ground provided the conflicting application or patent either is shown to be commonly owned with this application, or claims an invention made as a result of activities undertaken within the scope of a joint research agreement.

Effective January 1, 1994, a registered attorney or agent of record may sign a terminal disclaimer. A terminal disclaimer signed by the assignee must fully comply with 37 CFR 3.73(b).

Claims 24, 26-27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46-47, and 54-57 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-9 of U.S. Patent No. 6,774,122 ('122). Although the conflicting claims are not identical, they are not patentably distinct from each other because '122 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of

Art Unit: 1628

the solvents and the excipients are within the range taught in '122. The optimization of result effect parameters (e.g., dosing regimen [single dosing or multiple divided dosing], weight ratio of the actives and the excipients) is obvious as being within the skill of the artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

Claims 24, 26-27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46-47, and 54-57 are rejected on the ground of nonstatutory obviousness-type double patenting as being unpatentable over claims 1-12 of U.S. Patent No. 7,456,160 ('160). Although the conflicting claims are not identical, they are not patentably distinct from each other because '160 teaches the method of treating hormonal dependent benign or malignant disease of reproductive tract by employing the herein claimed composition. The ratio of the solvents and the excipients are within the range taught in '160. The optimization of result effect parameters (e.g., dosing regimen [single dosing or multiple divided dosing], weight ratio of the actives and the excipients) is obvious as being within the skill of the

Art Unit: 1628

artisan. The optimization of known effective amounts of known active agents to be administered, is considered well in the competence level of an ordinary skilled artisan in pharmaceutical science, involving merely routine skill in the art. It has been held that it is within the skill in the art to select optimal parameters, such as amounts of ingredients, in a composition in order to achieve a beneficial effect. See *In re Boesch*, 205 USPQ 215 (CCPA 1980). It is also noted that “[W]here the general conditions of a claim are disclosed in the prior art, it is not inventive to discover the optimum or workable ranges by routine experimentation.” *In re Aller*, 220 F.2d 454, 456, 105 USPQ 233, 235 (CCPA 1955).

No claims are allowed.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SAN-MING HUI whose telephone number is (571)272-0626. The examiner can normally be reached on Mon - Fri from 9:00 to 5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner’s supervisor, Brandon Fetterolf can be reached on (571) 272-2919. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Art Unit: 1628

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

San-ming Hui
Primary Examiner
Art Unit 1628

/San-ming Hui/
Primary Examiner, Art Unit 1628

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known		
				Application Number	12/285,887	
				Filing Date	October 15, 2008	
				First Named Inventor	John R. EVANS	
				Art Unit	1628	
				Examiner Name	HUI, San Ming R.	
Sheet	1	of	1	Attorney Docket Number	11285.0056-00000	

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS					
Examiner Initials	Cite No. ¹	Document Number	Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)			

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS						
Examiner Initials	Cite No. ¹	Foreign Patent Document	Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁶
		Country Code ³ Number ⁴ Kind Code ⁵ (if known)				
	1	WO 03/006064	23-JAN-2003	Astrazeneca AB		

NONPATENT LITERATURE DOCUMENTS			
Examiner Initials	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation ⁶
	2	The Merck Index, 12th Ed., Merck & Co., Inc., pgs. xiv, 189-190, 641-642 and 1715 (1996).	
	3	Guerrini, et al., "Pharmacokinetics of probenecid in sheep", J Vet Pharmacol Ther., 128-135 (1985).	
	4	Lavy, et al., "Pharmacokinetics of clindamycin HCl administered intravenously, intramuscularly and subcutaneously to dogs", J Vet Pharmacol Ther., 22(4):261-265 (1999).	
	5	Ismail, "Disposition kinetics of difloxacin after intravenous, intramuscular and subcutaneous administration in calves", Vet Res Commun., 31(4):467-476 (2007).	
	6	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) from August 27, 2009 to December 15, 2011.	
	7	Documents from the prosecution of European Application No. 10180667.7 (EP 2 266 573) from November 23, 2010 to December 19, 2011.	
	8	Documents from the prosecution of European Application No. 10180661.0 (EP 2 286 818) from January 19, 2011 to December 19, 2011.	
	9	Declaration Under 35 U.S.C §1.132 of Dr. Paul Gellert filed in August 2008 in U.S. Application No. 10/872,784.	
Examiner Signature	/San Ming Hui/		Date Considered 03/09/2012

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /S.H./

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known	
				Application Number	12/285,887
				Filing Date	October 15, 2008
				First Named Inventor	John R. EVANS
				Art Unit	1628
				Examiner Name	San Ming R. Hui
Sheet	1	of	1	Attorney Docket Number	11285.0056-00000

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS						
Examiner Initials ⁷	Cite No. ¹	Document Number		Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)				
		US-				
		US-				
		US-				
		US-				
		US-				

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS							
Examiner Initials ⁷	Cite No. ¹	Foreign Patent Document		Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁸
		Country Code ³	Number ⁴ Kind Code ⁵ (if known)				

NONPATENT LITERATURE DOCUMENTS			
Examiner Initials ⁷	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation ⁸
	1	McLeskey et al., "Tamoxifen-resistant fibroblast growth factor-transfected MCF-7 cells are cross-resistant <i>in vivo</i> to the antiestrogen ICI 182,780 and two aromatase inhibitors," Clin. Cancer Res., 4:697-711 (1998).	
	2	JRF Robertson, et al., "Fulvestrant: pharmacokinetics and pharmacology," British Journal of Cancer, 90(1):S7-S10 (2004).	
	3	John F. R. Robertson, "Fulvestrant (Faslodex®)--how to make a good drug better," The Oncologist, 12:774-784 (2007).	
	4	Search Report for European Patent Application No. 10180667.7 dated November 23, 2010.	
	5	Search Report for European Patent Application No. 10180661.0 dated January 19, 2011.	
	6	Documents from the Opposition against European Patent Application No. 01900186.6 from April 21, 2009 to September 7, 2009.	

Examiner Signature	/San Ming Hui/	Date Considered	03/09/2012
--------------------	----------------	-----------------	------------

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /S.H./

EAST Search History

EAST Search History (Prior Art)

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	86164	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L2	451	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L3	2807	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L4	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L5	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L6	16129	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L7	1884456	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L8	8391	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L9	4	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L10	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L11	14	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L12	1971	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33

L13	2	"6774122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L14	982	514/177.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L15	1418	514/178.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L16	2168462	castor oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L17	86164	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L18	451	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L19	2807	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L20	16129	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L21	1884456	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L22	8391	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L23	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L24	14	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L25	1971	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L26	86164	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L27	5153	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT;	OR	ON	2012/03/09 16:33

			IBM_TDB			
L28	2905	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L29	1495	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L30	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L31	3641	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L32	3641	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L33	86164	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L34	451	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L35	2807	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L36	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L37	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L38	16129	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L39	1884456	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L40	8391	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L41	4	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L42	7	((benzyl adj benzoate) or	US-PGPUB;	OR	ON	2012/03/09

		(phenylmethyl adj benzoate) same solvent) same (testosterone)	USPAT; EPO; JPO; DERWENT; IBM_TDB			16:33
L43	14	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L44	1971	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L45	86164	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L46	5153	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L47	2905	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L48	1495	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L49	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L50	3641	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L51	96080	breast adj cancer	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L52	2487	breast adj cancer and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L53	366	breast adj cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L54	1549	cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L55	2	"7456160".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33
L56	2	"6,774,122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/03/09 16:33

EAST Search History (Interference)

< This search history is empty >

3/ 9/ 2012 4:41:31 PM

C:\ Users\ shui\ Documents\ EAST\ Workspaces\ 12-285887.wsp

Index of Claims 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed

-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011	03/09/2012					
	1	✓							
	2	✓							
	3	✓							
	4	✓							
	5	✓							
	6	✓							
	7	✓							
	8	✓							
	9	✓							
	10	✓							
	11	✓							
	12	✓							
	13	✓							
	14	✓							
	15	✓							
	16	✓							
	17	✓							
	18	✓							
	19	✓							
	20	✓							
	21	✓							
	22	✓							
	23	✓							
	24		✓	✓					
	25		✓	-					
	26		✓	✓					
	27		✓	✓					
	28		✓	-					
	29		✓	✓					
	30		✓	✓					
	31		✓	-					
	32		✓	✓					
	33		✓	-					
	34		✓	✓					
	35		✓	✓					
	36		✓	✓					

Index of Claims 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed


-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011	03/09/2012					
	37		✓	-					
	38		✓	✓					
	39		✓	✓					
	40		✓	-					
	41		✓	✓					
	42		✓	✓					
	43		✓	-					
	44		✓	✓					
	45		✓	-					
	46		✓	✓					
	47		✓	✓					
	48		✓	-					
	49		✓	-					
	50		✓	-					
	51		✓	-					
	52		✓	-					
	53		✓	-					
	54			✓					
	55			✓					
	56			✓					
	57			✓					


Search Notes 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

SEARCHED			
Class	Subclass	Date	Examiner
514	177, 178	12/19/10	SH
514	177, 178	9/6/11	SH
514	177, 178	3/9/12	SH

SEARCH NOTES		
Search Notes	Date	Examiner
EAST and inventor search in PALM	12/19/10	SH
update search in EAST and inventor search in PALM	9/6/11	SH
EAST in EAST and inventor search in PALM	3/9/12	SH

INTERFERENCE SEARCH			
Class	Subclass	Date	Examiner

--	--

Application Number 	Application/Control No. 12/285,887	Applicant(s)/Patent under Reexamination EVANS ET AL.	

Document Code - DISQ	Internal Document – DO NOT MAIL
-----------------------------	--

TERMINAL DISCLAIMER	<input checked="" type="checkbox"/> APPROVED	<input type="checkbox"/> DISAPPROVED
Date Filed : 01/17/12	This patent is subject to a Terminal Disclaimer	

Approved/Disapproved by:

Lawana Hixon

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
)
John R. Evans et al.) Group Art Unit: 1628
)
Application No.: 12/285,887) Examiner: HUI, San Ming R.
)
Filed: October 15, 2008) Confirmation No.: 1199
)
For: FORMULATION) **VIA EFS-WEB**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

RESPONSE UNDER 37 C.F.R. § 1.111

In reply to the non-final Office Action mailed March 20, 2012 ("Office Action"),
and pursuant to 37 C.F.R. § 1.111, Applicants hereby respectfully request
reconsideration of this application in view of the following remarks.

REMARKS

I. Status of the claims and amendments

Claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 are pending in this application. No claims are being amended in this response.

II. Statement of Substance of Interview under 37 C.F.R. § 1.133(b)

Applicants would like to thank Examiner San Ming Hui for granting a personal interview to Applicants on March 1, 2012. Applicants present this Statement of Substance of Interview in connection with that interview conducted between Examiner San Ming Hui, Dr. Ronald J. Sawchuk, Dr. Paul R. Gellert (AstraZeneca Pharmaceuticals), Mr. Allen F. Giles (AstraZeneca Pharmaceuticals), and the undersigned.

During the interview, the participants discussed the outstanding rejections, the arguments Applicants presented in the Response filed on January 17, 2012, as well as the contents of the declaration under 37 C.F.R. §1.132 by Dr. Sawchuk filed on January 17, 2012.

No agreement was reached at the interview and the Examiner indicated he would consider the information submitted with the January 17th Response and discussed at the interview in the preparation of the next Office Action.

III. Double Patenting Rejection

The Office rejected claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 under the nonstatutory obviousness-type double patenting doctrine as

being unpatentable over: (a) claims 1-9 of U.S. Patent No. 6,774,122 (“the ’122 patent”) and (b) claims 1-12 of U.S. Patent No. 7,456,160 (“the ’160 patent”).

Applicants filed a Terminal Disclaimer on January 17, 2012, to obviate a double patenting rejection. In the Office Action, the Examiner refers to a decision dated February 15, 2012, which disapproved Applicants’ Terminal Disclaimer for failure to comply with 37 C.F.R. 3.73(b).

On March 23, 2012, the undersigned contacted Ms. Lawana Hixon, who was the paralegal who issued the decision dated February 15, 2012. During the call with Ms. Hixon, the undersigned pointed out that Applicants complied with 37 C.F.R. 3.73(b) because the Terminal Disclaimer: (1) recites the chain of title from the inventors to the current assignee, AstraZeneca AB, including citation to the relevant assignment records, and (2) states that the person signing the Terminal Disclaimer is authorized to act on behalf of AstraZeneca AB. Ms. Hixon agreed that the requirements of Section 3.73 had been met and a review of the image file wrapper of this application available through PAIR confirms Ms. Nixon approved the Terminal Disclaimer.

Thus, the instant double patenting rejections are moot and Applicants respectfully request that they be withdrawn.

IV. Conclusion

In view of the foregoing remarks, Applicants respectfully request reconsideration of this application and the timely allowance of the pending claims.


The Office is encouraged to contact the undersigned at the phone number below should the Office consider a telephonic conversation will facilitate prosecution of the application.

Please grant any extensions of time required to enter this response and charge any required fees not included with this Response to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated: April 9, 2012

By: 

Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
)
John R. Evans et al.) Group Art Unit: 1628
)
Application No.: 12/285,887) Examiner: HUI, San Ming R.
)
Filed: October 15, 2008) Confirmation No.: 1199
)
For: FORMULATION) **VIA EFS-WEB**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

INFORMATION DISCLOSURE STATEMENT UNDER 37 C.F.R. § 1.97(c)

Pursuant to 37 C.F.R. §§ 1.56 and 1.97(c), Applicant brings to the attention of the Examiner the documents on the attached listing. This Information Disclosure Statement is being filed after the events recited in Section 1.97(b) but, to the undersigned's knowledge, before the mailing date of either a Final action, Quayle action, or a Notice of Allowance. Under the provisions of 37 C.F.R. § 1.97(c), this Information Disclosure Statement is accompanied by a fee of \$180.00 as specified by Section 1.17(p)].

Copies of the listed foreign and non-patent literature documents are attached. It is the undersigned's understanding that the cited U.S. copending application is available to the Examiner through the PTO's Image File Wrapper system. Accordingly, a copy of that application is not enclosed. See M.P.E.P. § 609.04.

Applicants respectfully request that the Examiner consider the listed documents and indicate they were considered by making appropriate notations on the attached form.

This submission does not represent that a search has been made or that no better art exists and does not constitute an admission that each or all of the listed documents are material or constitute "prior art." If the Examiner applies any of the documents as prior art against any claims in the application and Applicants determine that the cited documents do not constitute "prior art" under United States law, applicants reserve the right to present to the office the relevant facts and law regarding the appropriate status of such documents.


Applicants further reserve the right to take appropriate action to establish the patentability of the disclosed invention over the listed documents, should one or more of the documents be applied against the claims of the present application.

If there is any fee due in connection with the filing of this Statement, please charge the fee to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated: April 9, 2012

By: 

Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known			
				Application Number		12/285,887	
				Filing Date		October 15,2008	
				First Named Inventor		John R. EVANS	
				Art Unit		1628	
				Examiner Name		HUI, San Ming R.	
Sheet	1	of	1	Attorney Docket Number		11285.0056-00000	

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS						
Examiner Initials ⁷	Cite No. ¹	Document Number		Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)				
	1	U.S. Application No. 13/387,584		Filing date: 27-Jan-2012	DIMERY et al.	

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS								
Examiner Initials ⁷	Cite No. ¹	Foreign Patent Document			Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁶
		Country Code ³	Number ⁴	Kind Code ⁵ (if known)				
	2	WO	2011/012885		03-Feb-2011	AstraZeneca UK Ltd.		

NONPATENT LITERATURE DOCUMENTS			
Examiner Initials ⁷	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation ⁶
	3	Buzdar, A. U., "Fulvestrant - A novel estrogen receptor antagonist for the treatment of advanced breast cancer," <i>Drugs of Today</i> , 44(9):679-692 (2008).	
	4	"Comparison of fulvestrant (faslodex) 250 mg and 500 mg in postmenopausal women with estrogen receptor-positive advanced breast cancer progressing or relapsing after previous endocrine therapy," <i>Clinicaltrials.gov</i> (20-May-2009) retrieved 24-Jan- 2012.	
	5	Di Leo A., et al., "Confirm: a phase III, randomized, parallel-group trial comparing fulvestrant 250 mg vs fulvestrant 500 mg in postmenopausal women with estrogen receptor-positive advanced breast cancer," <i>Cancer Res.</i> , 69(24) Supp. 3, (2009).	
	6	International Search Report for PCT Application No. PCT/GB10/51228 (WO 2011/012885) mailed December 20, 2012.	
	7	International Preliminary Report on Patentability for PCT Application No. PCT/GB10/51228 (WO 2011/012885) mailed December 20, 2012.	
	8	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) dated December 15, 2011.	
	9	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) dated February 27, 2012.	
Examiner Signature		Date Considered	

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 February 2011 (03.02.2011)

PCT

(10) International Publication Number
WO 2011/012885 A9

(51) International Patent Classification:
A61K 31/565 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:
PCT/GB2010/051228

(22) International Filing Date:
26 July 2010 (26.07.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0912999.0 27 July 2009 (27.07.2009) GB

(71) Applicant (for all designated States except MG, US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DIMERY, Isaiah, William** [US/US]; AstraZeneca Intellectual Property, AstraZeneca R & D Wilmington, 1800 Concord Pike, P.O. Box 15437, Wilmington, DE 19850-5437 (US). **WEBSTER, Alan** [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(74) Agent: **ASTRAZENECA INTELLECTUAL PROPERTY**; AstraZeneca AB, S -151 85 Södertälje (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IIR, IIU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(48) Date of publication of this corrected version:
24 March 2011

(15) Information about Correction:
see Notice of 24 March 2011



WO 2011/012885 A9

(54) Title: FULVESTRANT IN A DOSAGE OF 500MG FOR THE TREATMENT OF ADVANCED BREAST CANCER

(57) Abstract: The present invention relates to fulvestrant at a dosage of 500mg for use in the treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy.

FULVESTRANT IN A DOSAGE OF 500MG FOR THE TREATMENT OF ADVANCED BREAST CANCER

The present invention relates to fulvestrant at a dosage of 500mg for use in the treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy.

5 Breast cancer is one of the most common malignancies in women, comprising 18% of female cancers worldwide (Mcpherson et al 2000), and the most common cause of cancer deaths. The incidence varies among populations with about half of all cases occurring in North America and Western Europe. It has long been acknowledged that many breast cancers are hormone dependent and that hormonal manipulation can affect the
10 progress of the disease (Beatson 1896). The most important factor determining response to hormonal manipulation is the presence of the oestrogen receptor (ER) in the target tissue (Fisher et al 2001).

The antioestrogen (AO) tamoxifen has been the most widely used endocrine therapy for breast cancer in both premenopausal and postmenopausal women. However,
15 despite its demonstrated efficacy, *de novo* or acquired resistance may occur during treatment. In some patients, the disease progresses during therapy because tumour growth may be stimulated by tamoxifen, due to its partial agonist activity on the ER (Wiebe et al 1993).

The search for a pure AO, devoid of the agonist activity of tamoxifen, resulted in
20 the discovery and clinical development of ICI 182,780 (also known as fulvestrant or FASLODEX™). Fulvestrant is an ER antagonist without known agonistic properties that down-regulates cellular levels of the ER in a dose-dependent manner (Howell et al 2000, Robertson et al 2001, Wakeling et al 1991). Fulvestrant is well tolerated and has demonstrated efficacy in women whose breast cancer had progressed following endocrine
25 therapy (Howell et al 2002, Osborne et al 2002, Chia et al 2008).

Women diagnosed with early breast cancer are generally treated with tamoxifen or an aromatase inhibitor if endocrine therapy is appropriate. However if the cancer recurs or progresses there is a need for alternative therapeutics. Fulvestrant (FASLODEX™) is presently approved at a dose of 250mg as an alternative endocrine therapy. The present

invention is based on the discovery that increasing the dose of fulvestrant to 500mg is more advantageous for patients than the 250mg dose.

One feature of the invention provides fulvestrant at a dosage of 500mg for use in the treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy. Preferably the fulvestrant is administered monthly. Preferably an additional dose of 500mg is administered during the first month of treatment. Preferably the additional dose is administered at about day 14. Preferably the woman is oestrogen receptor positive or progesterone receptor positive; more preferably oestrogen receptor positive. Preferably the progression or recurrence on endocrine therapy comprised therapy with tamoxifen or an aromatase inhibitor. Preferably the aromatase inhibitor is selected from anastrozole, letrozole or exemestane; more preferably anastrozole or letrozole. Preferably the use use of fulvestrant at 500mg dosage provides an increase the time to progression compared with fulvestrant at a dosage of 250mg; in particular the doses are preferably administered monthly with an additional dose at 500mg in the first month. Tamoxifen, anastrozole, letrozole and excmestane are all commercially available drugs with regulatory approval for administration to women with breast cancer.

Another feature of the invention provides the use fulvestrant at a dosage of 500mg for preparation of a medicament for treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy. This feature may be combined with any of the preferred features described herein.

Another feature of the invention provides the treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy with fulvestrant at a dosage of 500mg. This feature may be combined with any of the preferred features described herein.

The invention is exemplified by the following non-limiting Example, in which Figure 1 shows a Kaplan-Meier plot of time to progression comparing fulvestrant at 250mg with 500mg. The x-axis shows the time in months and y-axis shows proportion of patients progression free. Tick marks indicate censored observations.

LIST OF ABBREVIATIONS AND DEFINITIONS OF TERMS

Abbreviation or special term	Explanation
AE	Adverse event
AI	Aromatase inhibitor
ALT	Alanine aminotransferase
AO	Antioestrogen
AST	Aspartate aminotransferase
BOR	Best objective/overall response
CBR	Clinical benefit rate
CI	Confidence interval
CR	Complete response
CRA	Clinical research associate
CRF	Case report form
CSP	Clinical Study Protocol
CSR	Clinical Study Report
CT	Computed tomography
CTCAE	Common terminology criteria for adverse events
DAE	Premature discontinuation of treatment with investigational product due to an adverse event (adverse events).
DCO	Data cut-off
DoCB	Duration of clinical benefit
DoR	Duration of response
ECG	Electrocardiogram
EDoCB	Expected duration of clinical benefit
EDoR	Expected duration of response
Endpoint	A status of the patient that constitutes the 'endpoint' of a patient's participation in a clinical study and that is used as the final outcome.
ER	Oestrogen receptor
EU	European Union
FACT-B	Functional Assessment of Cancer Therapy - breast cancer
FSH	Follicle stimulating hormone
GCP	Good clinical practice
HER	Human epidermal growth factor receptor

Abbreviation or special term	Explanation
HRQoL	Health-related quality of life
ICH	International Conference on Harmonisation
IDMC	Independent Data Monitoring Committee
IEC	Independent Ethics Committee
im	Intramuscular
INR	International normalised ratio
IRB	Institutional Review Board
International Co-ordinating investigator	An Investigator assigned the responsibility for the co-ordination of investigators across all Study Sites participating in a multinational, multicentre study.
LD	Longest diameter
LHRH	Luteinising hormone releasing hormone
MedDRA	Medical dictionary for regulatory activities
MRI	Magnetic resonance imaging
NCCN	National Comprehensive Cancer Network
OAE	Other significant adverse event (ie, significant AEs, other than SAEs and DAEs, which are of particular clinical importance in this development program).
OR	Objective response
ORR	Objective response rate
OS	Overall survival
Outcome variable	A variable (usually a derived variable) specifically defined to be used in the analysis of a study objective.
Patient identifier	Only one variable is used to identify each patient within the reporting database. This identifier is a concatenation of the Study Number, and the enrolment Code (eg, D1234C00001/E0010001). Within an individual study report, the enrolment code alone (eg, E0010001) may be used to reference individual patients in-text within the CSR, including tables and listings. With respect to individual Patient Narratives, and the higher level documents, the full unique patient identifier should be used.
PD	Progressive disease
PgR	Progesterone receptor
PPS	Per Protocol Set
PR	Partial response
Principal investigator	A person responsible for the conduct of a clinical study at an investigational study site. Every investigational study site has a principal investigator.
PRO	Patient reported outcomes
PT	Preferred term

Abbreviation or special term	Explanation
RECIST	Response evaluation criteria in solid tumours
SAE	Serious adverse event
SAP	Statistical Analysis Plan
SD	Stable disease
sd	Standard deviation
SE	Standard error
SOC	System organ class
TOI	Trial outcome index
TTP	Time to progression. The definition of TTP used in this clinical study is also commonly termed progression free survival (PFS).
TTR	Time to response
ULRR	Upper limit reference range
US	United States of America
Variable	A characteristic or a property of a patient that may vary eg from time to time or between patients.
WHO	World Health Organisation

Example 1

A Randomised, Double-Blind, Parallel-group, Multicentre, Phase III Study

5 **Comparing the Efficacy and Tolerability of Fulvestrant (FASLODEX™) 500 mg with Fulvestrant (FASLODEX™) 250 mg in Postmenopausal Women with Oestrogen Receptor Positive Advanced Breast Cancer Progressing or Relapsing after Previous Endocrine Therapy**

10 This study assessed the relationship between fulvestrant dose and efficacy. It compared the current approved dose and dosing schedule of fulvestrant (250 mg every 28 days) with a higher dose regimen (500 mg every 28 days plus an additional 500 mg on Day 14 of the first month only). The study is also referred to as CONFIRM.

Study centres

15 One-hundred and twenty-eight centres in 17 countries (Belgium, Brazil, Chile, Colombia, Czech Republic, Hungary, India, Italy, Malta, Mexico, Poland, Russia, Slovakia, Spain, USA, Ukraine and Venezuela). The US, Mexico, Italy, Brazil, Spain, Chile, Colombia and

Venezuela also participated in health-related quality of life (HRQoL) assessments during the study.

Objectives

The primary objective of the study was to compare the efficacy of fulvestrant 500 mg treatment with fulvestrant 250 mg treatment in terms of time to progression (TTP).

The secondary objectives of the study were:

- To compare the objective response rate (ORR) of patients treated with fulvestrant 500 mg with the objective response rate of patients treated with fulvestrant 250 mg.
- To compare clinical benefit rate (CBR) of patients treated with fulvestrant 500 mg with the clinical benefit rate of patients treated with fulvestrant 250 mg.
- To compare duration of response (DoR) of patients treated with fulvestrant 500 mg with the duration of response of patients treated with fulvestrant 250 mg.
- To compare the duration of clinical benefit (DoCB) of patients treated with fulvestrant 500 mg with the duration of clinical benefit of patients treated with fulvestrant 250 mg.
- To compare the overall survival (OS) of patients treated with fulvestrant 500 mg with the overall survival of patients treated with fulvestrant 250 mg.
- To assess the tolerability of fulvestrant 500 mg treatment compared with fulvestrant 250 mg treatment.
- To assess the health-related quality of life (HRQoL) of patients treated with fulvestrant 500mg as compared to fulvestrant 250 mg in a subgroup of patients.

Study design

This was a randomised, double-blind, parallel-group, multicentre, phase III study to compare 2 dose levels of fulvestrant in postmenopausal women with oestrogen receptor positive (ER+ve) advanced breast cancer who had either relapsed whilst on adjuvant endocrine therapy, or progressed whilst on first endocrine therapy for advanced disease.

Target patient population and sample size

A total of 720 postmenopausal women with histological/cytological confirmation of ER+ve breast cancer who had relapsed or progressed on previous endocrine therapy were planned to be recruited; a total of 736 were actually randomised.

The sample size calculation was based on the primary variable, TTP, and assumed exponential progression times. The sample size was driven by the number of required events. In order to detect a hazard ratio of ≤ 0.8 (or ≥ 1.25) for fulvestrant 500 mg compared to fulvestrant 250 mg, at a 2-sided significance level of 5%, with 80% power, approximately 632 events were required to have occurred in the study (ie, approximately 632 patients to have progressed or died).

Investigational product and comparator: dosage, mode of administration and batch numbers

Fulvestrant 500 mg was given as two 5 ml intramuscular (im) injections, one in each buttock, on days 0, 14, 28 and every 28 (± 3) days thereafter.

Fulvestrant 250 mg was given as two 5 ml im injections (1 fulvestrant injection plus 1 placebo injection), one in each buttock, on days 0, 14 (2 placebo injections only), 28 and every 28 (± 3) days thereafter.

Duration of treatment

Treatment was to continue until disease progression occurred, unless any of the criteria for treatment discontinuation were met first.

Criteria for evaluation - efficacy and pharmacokinetics (main variables)

Efficacy

The primary outcome variable TTP; secondary variables were ORR, CBR, DoR, DoCB and OS.

Patient reported outcomes

The primary patient reported outcome for HRQoL was the Trial Outcome Index (TOI) derived from the Functional Assessment of Cancer Therapy - Breast cancer (FACT-B) questionnaire.

Criteria for evaluation - safety (main variables)

Outcome variables for safety were frequency and severity of adverse events (AEs), including pre-specified AEs of interest.

Statistical methods

For the primary endpoint TTP, the primary analysis was an unadjusted log-rank test and the secondary analysis was a Cox proportional hazard model, adjusted for treatment and other predefined covariates.

For OS, the unadjusted log-rank test was performed. For ORR and CBR, a logistic regression model with treatment factor only was fitted. DoR and DoCB were analysed in those patients who had an OR and CB, respectively. For HRQoL endpoints, a longitudinal model with treatment and other covariates was used.

5 The hypotheses for TTP, ORR, CBR, DoR, DoCB, OS, FACT-B score and TOI score were:

H₀: fulvestrant 500 mg is not different from fulvestrant 250 mg, vs.

H₁: fulvestrant 500 mg is different from fulvestrant 250 mg

For efficacy and HRQoL endpoints, summaries and analyses were carried out according to
10 the randomised treatment ie, using the Full Analysis Set. For safety endpoints, summaries and analyses were carried out according to the treatment actually received, ie, using the safety analysis set. The primary endpoint was also analysed in the per protocol set (PPS).

Patient population

A total of 720 patients were planned to be recruited; 736 were actually randomised.

15 Diagram S1 shows the number of patients randomised to each of the 2 treatment groups and the number in each of the populations analysed. In addition, HRQoL was analysed in 145 of the patients in the Full Analysis Set (72 patients in the fulvestrant 500 mg group and 73 patients in the fulvestrant 250 mg group). The patient population was consistent with the one intended to be recruited. In the fulvestrant 500 mg group, 41 patients were
20 ongoing study treatment at data cut off (DCO) compared with 31 patients in the fulvestrant 250 mg group.

1.1 Selection of study population

Before entering the study, patients were assessed to ensure that they met the eligibility criteria. Investigators had to keep a record of patients who were considered for enrolment
25 but were never randomised (patient screening log). This information is necessary to establish that the patient population was selected without bias. The patient screening log had to be filed in the Investigator study file at each centre.

1.1.1 Inclusion criteria

For inclusion in the study patients had to fulfil all of the following criteria:

- 30 1. Provision of written informed consent
2. Histological/cytological confirmation of breast cancer

3. Documented ER+ve status of primary or metastatic tumour tissue, according to the local laboratory parameters
4. Requiring endocrine therapy:
 - Relapsing during, or within 12 months of completion of, adjuvant endocrine therapy (tamoxifen, toremifene or AIs such as anastrozole, letrozole and exemestane), or
 - Progressing on an endocrine therapy (tamoxifen, toremifene or AIs such as anastrozole, letrozole and exemestane) provided that this endocrine treatment was started at least 12 months after the completion of adjuvant endocrine treatment, or
 - Progressing on an endocrine therapy (tamoxifen, toremifene or AIs such as anastrozole, letrozole and exemestane) given as first treatment for patients with *de novo* advanced¹ breast cancer
5. Fulfilling one of the following criteria:
 - Patients with measurable disease as per RECIST criteria. This is defined as at least one lesion that can be accurately measured in at least one dimension (longest diameter to be recorded) as ≥ 20 mm with conventional techniques or as ≥ 10 mm with spiral CT scan.
 - Patients with bone lesions, lytic or mixed (lytic and sclerotic), in the absence of measurable disease as defined by RECIST.
6. Postmenopausal woman, defined as a woman fulfilling any 1 of the following criteria:
 - Age ≥ 60 years.
 - Age ≥ 45 years with amenorrhoea ≥ 12 months with an intact uterus.
 - Having undergone a bilateral oophorectomy
 - Follicle stimulating hormone (FSH) and oestradiol levels in postmenopausal range (utilising ranges from the local laboratory facility).
 - In patients who had previously been treated with a luteinising hormone releasing hormone (LHRH) analogue, the last depot must have been

¹ Advanced breast cancer: Metastatic disease or locally advanced disease which is not amenable to treatment with curative intent.

administered more than 4 months prior to randomisation, menses must not have restarted, and FSH and oestradiol levels must also have been in the postmenopausal range (utilising ranges from the local laboratory facility).

7. WHO performance status 0, 1 or 2.

5 **Rationale for inclusion criteria**

1. This criterion was set as part of the ethical conduct of the study, which complies with GCP.

2. This criterion was set to objectively confirm breast cancer.

3. This criterion was set to select a patient population expected to respond to fulvestrant based on its mechanism of action.

10 4. This criterion was set to clarify the history of hormonal therapy for breast cancer in this study.

5. This criterion was set to enable the conduct of efficacy assessments according to modified RECIST.

15 6. This criterion was set because the effect of fulvestrant on pre-menopausal breast cancer patients had not been fully assessed.

7. This criterion was set to conduct efficacy assessments properly and to ensure the safety of patients.

1.1.2 Exclusion criteria

20 Any of the following was regarded as a criterion for exclusion from the study:

1. Presence of life-threatening metastatic visceral disease, defined as extensive hepatic involvement, or any degree of brain or leptomeningeal involvement (past or present), or symptomatic pulmonary lymphangitic spread. Patients with discrete pulmonary parenchymal metastases were eligible, provided their respiratory function was not compromised as a result of disease.

2. More than one regimen of chemotherapy for advanced disease.²

3. More than one regimen of endocrine therapy for advanced disease.³

² Patients previously treated with one regimen of chemotherapy for advanced disease were allowed as long as their last treatment was an AO or an AI.

³ Oophorectomy, ovarian ablation, or LHRH analogue therapy did not count as endocrine treatments in this context and also did not render the patient ineligible for this study.

4. Extensive radiation therapy within the last 4 weeks (greater than or equal to 30% marrow or whole pelvis or spine) or cytotoxic treatment within the past 4 weeks prior to screening laboratory assessment, or strontium-90 (or other radiopharmaceuticals) within the past 3 months.
5. Treatment with a non-approved or experimental drug within 4 weeks before randomisation.
6. Current or prior malignancy within previous 3 years (other than breast cancer or adequately treated basal cell or squamous cell carcinoma of the skin or in-situ carcinoma of the cervix).
- 10 7. Any of the following laboratory values:
 - Platelets $<100 \times 10^9/L$
 - Total bilirubin $>1.5 \times$ upper limit reference range (ULRR)
 - ALT or AST $>2.5 \times$ ULRR if no demonstrable liver metastases or $>5 \times$ ULRR in presence of liver metastases.
- 15 8. History of:
 - Bleeding diathesis (ie, disseminated intravascular coagulation, clotting factor deficiency), or
 - Long-term anticoagulant therapy (other than antiplatelet therapy and low dose warfarin (see Section 3.7 of the CSP [Appendix 12.1.1 of this report])).
- 20 9. History of hypersensitivity to active or inactive excipients of fulvestrant and/or castor oil.
10. Any severe concomitant condition which made it undesirable for the patient to participate in the trial or which would jeopardize compliance with the CSP, eg,
25 uncontrolled cardiac disease or uncontrolled diabetes mellitus.

Rationale for exclusion criteria

The exclusion criteria for concurrent diseases, concomitant drugs and patients' conditions were set because they were considered to affect the safety of patients or the efficacy assessment of fulvestrant in hormone receptor positive, postmenopausal advanced or
30 recurrent breast cancer.

1.1.3 Restrictions

The following restrictions were applied to patients in this trial:

1. Patients who were blood donors were not to donate blood during the study and
5 for 12 weeks following their last dose of randomised treatment.
2. Patients who had confirmed disease progression must have been discontinued from
their randomised treatment.
3. Concomitant treatments listed in Section 3.7 of the CSP.


Rationale for restrictions

- 10 1. This restriction was included to ensure that anaemia was not induced by blood
donation following the additional blood sampling requirement of the study.
2. This restriction was included to protect patients who were not receiving or who
ceased to receive clinical benefit from their study treatment and is in line with
current clinical practice.
- 15 3. This restriction was included because the concomitant treatments listed in
Section 3.7 of the CSP were considered to effect the safety of patients or the
efficacy assessment of the study drugs.

1.1.4 Discontinuation of patients from treatment or assessment

Patients could be discontinued from study treatment and assessments at any time at the
20 discretion of the investigators. Patients were also free to discontinue their participation in
the study at any time, without prejudice to further treatment. Specific reasons for
discontinuing a patient from this study, and the procedures to be followed when a patient
discontinued or was incorrectly enrolled, are listed in Section 3.3.5 of the CSP. For
patients who discontinued, it was noted whether they were assessed after study medication
25 was stopped, and whether they were asked about the reason(s) for their discontinuation and
about the presence of any adverse events (AEs). If possible, they were seen and assessed
by an investigator. AEs were followed up for 56 days after the last injection.

Diagram S1 Analysis sets



^a The patient who was excluded from the safety analysis set was also classified as a deviator, therefore these n values are not mutually exclusive.

5 **Summary of demographics and baseline characteristics**

A total of 96.1% of patients randomised into the study were Caucasian. The mean age of patients was 60.9 years and the mean weight of patients was approximately 70 kg.

Tumour characteristics were well balanced across the 2 treatment groups. Most patients (507 [68.9%]) were ER+ve and PgR+ve at primary diagnosis and almost all

10 patients (721 [98%]) had metastatic disease at baseline. In this study, 42.5% of patients had relapsed or progressed on AI therapy and 57.5% had relapsed or progressed on AOs. Most patients had relapsed or progressed either during previous adjuvant endocrine cancer therapy (344 patients [46.7%]) or during endocrine therapy given as a first treatment for *de*

novo advanced disease (255 patients [34.6%]). Approximately two thirds of patients had shown a response⁴ to their last endocrine therapy.

Summary of efficacy results

A summary of efficacy data is presented in Table S1.

Table S1 Summary of efficacy results for the main outcome variables

Variable	Result
Primary outcome variable	
TTP ^a	Hazard ratio=0.80 (95% CI 0.68–0.94); p=0.006 Median TTP: fulvestrant 500 mg =6.5 months; fulvestrant 250 mg =5.5 months % patients progression free at 12 months: fulvestrant 500 mg=34%; fulvestrant 250 mg = 25%
Secondary outcome variables	
ORR	Odds ratio=0.94 (95% CI 0.57–1.55); p=0.795 ORR: fulvestrant 500 mg=13.8%; fulvestrant 250 mg=14.6%
CBR	Odds ratio=1.28 (95% CI 0.95–1.71); p=0.100 CBR: fulvestrant 500 mg=45.6%; fulvestrant 250 mg=39.6%
DoR ^b	Ratio of EDoR=0.894 (95% CI 0.479–1.667); p=0.724 Median DoR ^c : fulvestrant 500 mg=19.4 months; fulvestrant 250 mg=16.4 months
DoCB	Ratio of EDoCB=1.357 (95% CI 1.067–1.726); p=0.013 Median DoCB: fulvestrant 500 mg=16.6 months; fulvestrant 250 mg=13.9 months
OS	Hazard ratio=0.84 (95% CI 0.69–1.03); p=0.091 Median OS: fulvestrant 500 mg=25.1 months; fulvestrant 250 mg=22.8 months % patients alive at 24 months: fulvestrant 500 mg=53%; fulvestrant 250 mg=49%

^a TTP ≡ progression-free survival. At data cut-off, 84% of patients had progressed or died in the absence of progression.

^b measured from randomisation to progression

^c from randomisation.

⁴ Defined as patients who experienced recurrence after ≥2 years on adjuvant endocrine therapy and/or patients who received clinical benefit (CR, PR or SD ≥24 weeks) from first-line therapy for advanced disease.

TTP:time to progression; ORR:objective response rate; CBR:clinical benefit rate; DoR:duration of response; DoCB:duration of clinical benefit; OS:overall survival; EDoR:expected duration of response; EDoCB:expected duration of clinical benefit. Fulvestrant 500 mg was associated with a significantly longer TTP compared with
5 fulvestrant 250 mg (hazard ratio=0.80 [95% CI 0.68–0.94]; p=0.006) corresponding to a reduction in risk of progression of 20%. Subgroup analyses showed a consistent treatment effect across all 6 predefined baseline covariates, including patients treated previously with either an aromatase inhibitor (AI) or antioestrogen (AO).

The ORR for fulvestrant 500 mg and fulvestrant 250 mg were similar (13.8% and 14.6%
10 respectively, odds ratio=0.94 [95% CI 0.57 to 1.55]; p=0.795) but there was a trend for an increased CBR in patients receiving fulvestrant 500 mg compared to those receiving fulvestrant 250 mg (45.6% vs. 39.6%, odds ratio=1.28 [95% CI 0.95 to 1.71]; p=0.100).

There was no statistically significant difference between the 2 treatment groups in expected DoR (EDoR); however, there was a statistically significant improvement in expected
15 DoCB (EDoCB) in patients randomised to receive fulvestrant 500 mg compared with patients randomised to receive fulvestrant 250 mg (9.83 months vs. 7.24 months, ratio of EDoCB=1.357 [95% CI 1.067 to 1.726]; p=0.013).

There was a trend for improved survival for patients treated with fulvestrant 500 mg compared with fulvestrant 250 mg (hazard ratio=0.84 [95% CI 0.69 to 1.03]; p=0.091); this
20 corresponds to a 16% reduction in risk of death.

In the subgroup of patients where it was measured, on-treatment HRQoL for both fulvestrant 500 mg and fulvestrant 250 mg was good (mean TOI score of approximately 60 out of 92). Patients treated with fulvestrant 500 mg had a similar on-treatment HRQoL to patients treated with fulvestrant 250 mg and there were no statistically significant
25 differences between the 2 treatment groups in terms of change in on treatment HRQoL as measured by both the TOI and FACT-B score, although there was a numerical advantage in TOI in favour of fulvestrant 500 mg.

Efficacy results

Primary variable: Time to progression

30 The primary objective of this study was to compare TTP between patients treated with fulvestrant 500 mg and those treated with fulvestrant 250 mg. The primary analysis set was the Full Analysis Set. An analysis of TTP in the PPS was also performed as a

secondary analysis. Table S2 shows the TTP data for patients in the fulvestrant 500 mg and fulvestrant 250 mg groups in the Full Analysis Set; Figure 1 shows a Kaplan-Meier plot of these data.

At DCO 618/736 (84.0%) patients had progressed or died in the absence of progression (297 [82.0%] in the fulvestrant 500 mg group and 321 [85.8%] in the fulvestrant 250 mg group). The unadjusted log rank test indicates that the TTP for patients in the fulvestrant 500 mg group was significantly longer than for those in the fulvestrant 250 mg group (hazard ratio=0.80 [95% CI 0.68 to 0.94]; p=0.006). Median TTP was 6.5 months in the fulvestrant 500 mg group and 5.5 months in the fulvestrant 250 mg group. The Kaplan-Meier plot for TTP in the Full Analysis Set shows a separation between the 2 treatment groups from approximately 3 months, favouring the fulvestrant 500 mg group.

Month	0	4	8	12	16	20	24	28	32	36	40	44	48
Fulvestrant 500mg at risk	362	216	163	113	90	54	37	19	12	7	3	2	0
Fulvestrant 250mg at risk	374	199	144	85	60	35	25	12	4	3	1	1	0

Table S2 Summary of time to progression: Full Analysis Set

	Fulvestrant 500 mg N=362	Fulvestrant 250 mg N=374
Number progressed (%)	297 (82.0)	321 (85.8)
Median (months)	6.5	5.5
Time to progression (months): 25% quartile	2.8	2.7
Time to progression (months): 75% quartile	16.6	11.9
Percentage of patients progression free at:		
6 months	51%	45%
12 months	34%	25%
18 months	23%	14%
24 months	16%	11%
Hazard ratio (95% CI)	0.80 (0.68–0.94)	
p-value	0.006	

Time to progression is the time between randomisation and the earliest of progression or death from any cause.

A hazard ratio <1 indicates fulvestrant 500 mg is associated with a longer time to disease progression than fulvestrant 250 mg

5 A hazard ratio >1 indicates fulvestrant 500 mg is associated with a shorter time to disease progression than fulvestrant 250 mg

Data source: Tables 11.2.1.1, 11.2.1.2 and 11.2.1.5.

The primary analysis of TTP is supported by the Cox proportional hazards regression analysis, adjusted for treatment and 6 specified covariates (hazard ratio=0.78 [95% CI 0.67
10 to 0.92]; $p=0.003$).

Summary of safety results

Fulvestrant 500 mg was well tolerated and its safety profile was consistent with the known safety profile of fulvestrant 250 mg. The most commonly reported pre-specified AEs of interest were gastrointestinal disturbances and joint disorders (approximately 20% and
15 19% of patients, respectively, in each of the treatment groups). There were no differences between treatment groups in the incidence or type of AEs, serious AEs and AEs leading to discontinuation. There was no evidence for dose dependence for any AE. There were no clinically important changes in haematology, clinical chemistry, vital signs or physical findings.

20 Conclusions

This study demonstrates that fulvestrant 500 mg provides a clinically meaningful benefit over fulvestrant 250 mg, in terms of TTP, in the treatment of postmenopausal women with ER+ve advanced breast cancer who have progressed or recurred on endocrine therapy. Further analyses demonstrated that the TTP data obtained in the study are robust. The
25 results show that fulvestrant 500 mg reduces the risk of disease progression by 20% compared with fulvestrant 250 mg. The risk in progression appears to be reduced in the fulvestrant 500 mg group compared to the 250 mg group by 3 observed factors:

- a reduction in the proportion of patients with a best objective response of progressive disease (38.7% in the fulvestrant 500 mg group vs 44.7% in the
30 fulvestrant 250 mg group)
- an increase in the proportion of patients who achieved clinical benefit (45.6% vs 39.6%, respectively)

- an increase in the duration of clinical benefit in patients receiving clinical benefit (median of 16.6 months vs 13.9 months, respectively).

There was also a trend towards improved survival in the fulvestrant 500 mg group (median of 25.1 months compared with 22.8 months in the 250 mg group), indicating that the
5 observed treatment comparison for overall survival supports the advantage observed for TTP and suggesting that the benefit provided by treatment, in terms of progression, is maintained past progression.

In the subgroup of patients where it was measured, on-treatment HRQoL remained stable while patients were receiving study treatment; there was no detrimental effect of the
10 fulvestrant 500 mg dose compared with 250 mg.

In the registration trials for fulvestrant, Studies 20/21, fulvestrant 250 mg was shown to be non-inferior to anastrozole (Robertson et al 2003). Demographic characteristics of patients in the CONFIRM study were broadly similar to those of patients in the combined analysis of Studies 20/21 and the efficacy results for fulvestrant 250 mg were consistent across the
15 studies (median TTP of 5.5 months in CONFIRM and the combined analysis of Studies 20/21). Data from these studies give further reassurance of the significant benefit that fulvestrant 500 mg offers over an already effective 250 mg dose.

The treatment effect for TTP, favouring fulvestrant 500 mg, was consistent across all subgroups analysed. The consistency of the TTP treatment effect in the aromatase
20 inhibitor (AI) and antiestrogen (AO) subgroups is of particular interest, given that in many markets the current regulatory approval for fulvestrant 250 mg is limited to patients who have progressed on AO therapy. Since the first regulatory approval for the use of non-steroidal AIs in breast cancer, changes in clinical practice have meant that there has been a considerable increase in the proportion of patients being treated upfront with these
25 drugs in both the adjuvant and the advanced setting (see National Comprehensive Cancer Network [NCCN], Inc. 2009 and references therein for more details). There are few endocrine treatment options available to patients who progress on AI therapy and it is therefore important to identify agents that effectively prolong the time to progression after failing on such therapy. Although guidelines like NCCN support the use of a same class
30 agent with a steroidal structure (steroidal AIs) in patients who have progressed on a non-steroidal AI, there are currently no agents of this type with regulatory approval for this treatment sequence. Fulvestrant 500 mg has a different mechanism of action to AIs and is

the first agent to show consistent benefit in a phase III setting in patients who have progressed during either AO or AI therapy.

The safety profile of fulvestrant 500 mg is consistent with the known safety profile of fulvestrant 250 mg with no evidence for dose dependence for any AE. The 2 SAEs that
5 were considered by the investigator to be possibly causally related to study treatment were confounded by other factors in the patients' medical histories and concomitant medications. The incidence of pre-specified AEs was well balanced between the 2 treatment groups. Although the incidence of injection site reactions was similar between treatment groups, a full assessment of the injection procedure was not possible to evaluate due to the double
10 blind design. However, it is reassuring to observe that there is no increase in the AE incidence with doubling the dose of fulvestrant.

Overall, fulvestrant 500 mg provides improved efficacy without any detrimental effect on safety, tolerability or HRQoL compared with fulvestrant 250 mg.

Overall conclusions

15 The CONFIRM study demonstrated a clear improvement in the efficacy of fulvestrant 500 mg when compared with the currently approved dose of fulvestrant 250 mg. There was a statistically significant prolongation of the TTP with a 20% reduction in the risk of progressing for patients receiving fulvestrant 500mg. Given the superior efficacy, similar safety, tolerability and HRQoL that fulvestrant 500mg offers over fulvestrant 250mg we
20 conclude that there is a superior benefit-risk profile for fulvestrant 500mg in patients recurring or progressing on endocrine therapy.

References

Addo et al 2002

25 Addo S, Yates RA, Laight A. A phase I trial to assess the pharmacology of the new oestrogen receptor antagonist fulvestrant on the endometrium in healthy postmenopausal volunteers. Br J Cancer 2002;87:1354-9.

Beatson 1896

30 Beatson GT. On the treatment of inoperable cases of carcinoma of the mamma; suggestions for a new method of treatment with illustrative cases. Lancet 1896; 2:104-7.

Chia et al 2008

Chia S, Gradishar W, Mauriac L, Bines J, Amant F, Federico M, et al. A double blind, randomized placebo controlled trial of fulvestrant versus exemestane following prior non-steroidal aromatase inhibitor therapy in post-menopausal women with hormone receptor
5 positive advanced breast cancer: results from EFACT. *J Clin Oncol* 2008; 26: 1664–70.

Collet 1997

Collet D. *Modelling Survival Data in Medical Research*. Chapman and Hall, 1997.

Dancey et al 2009

Dancey JE, Dodd LE, Ford R, Kaplan R, Mooney M, Rubinstein L et al.

10 Recommendations for the assessment of progression in randomised cancer treatment trials. *European Journal of Cancer* 2009;45:281-289.

DeFriend et al 1994

DeFriend DJ, Howell A, Nicholson RI, Anderson E, Dowsett M, Mansel RE, et al.

15 Investigation of a new pure antiestrogen (ICI 182780) in women with primary breast cancer. *Cancer Res.* 1994;54:408-414.

Dodd et al 2008

Dodd LE, Korn EL, Freidlin B, Jaffe CC, Rubinstein LV, Dancey J, et al. Blinded independent central review of progression-free survival in phase III clinical trials: important design element or unnecessary expense? *J Clin Oncol.* 2008;26:3791-6.

Ellis et al 2008

20 Ellis S, Carroll KJ, Pemberton K. Analysis of duration of response in oncology trials, *Contemp Clin Trials* 2008; 29:456-65.

Fisher et al 2001

Fisher B, Anderson S, Tan-Chiu E, Wolmark N, Wicherham DL, Fisher ER, et al.

25 Tamoxifen and chemotherapy for axillary node-negative, estrogen receptor-negative breast cancer findings from the National Surgical Adjuvant Breast and Bowel Project B-23. *J Clin Oncol.* 2001;19(4):931-42.

Howell et al 2000

Howell A, Osborne CK, Morris C, Wakeling AE. ICI 182,780 (FASLODEX):

30 development of a novel, “pure” anti-estrogen. *Cancer* 2000;89:817-25.

Howell et al 2002

Howell A, Robertson JFR, Quaresma Albano J, et al. Fulvestrant, formerly ICI 182,780, is as effective as anastrozole in postmenopausal women with advanced breast cancer progressing after prior endocrine treatment J Clin Oncol, 2002; 20: 3396-3403.

5 **Howell et al 2005**

Howell A, Pippen J, Elledge RM, Mauriac L, Vergote I, Jones SE, et al. Fulvestrant versus anastrozole for the treatment of advanced breast carcinoma: a prospectively planned combined survival analysis of two multicenter trials. Cancer 2005; 104: 236–239.

Mcpherson et al 2000

10 McPherson K, Steel CM, Dixon JM. ABC of breast diseases. Breast cancer – epidemiology, risk factors and genetics. BMJ 2000;321:624–8.

National Comprehensive Cancer Network [NCCN], Inc. 2009

The NCCN Clinical Practice Guidelines in Oncology™ Breast Cancer (Version 1.2009)© Available at: NCCN.org. Accessed [June 22, 2009]. To view the most recent and complete
15 version of the NCCN Guidelines, go online to NCCN.org.

Osborne et al 2002

Osborne CK, Pippen J, Jones SE, et al. Double-blind, randomized trial comparing the efficacy and tolerability of fulvestrant versus anastrozole in postmenopausal women with advanced breast cancer progressing on prior endocrine therapy: Results of a North
20 American trial. J Clin Oncol, 2002; 20: 3386-3395.

Robertson et al 2001

Robertson JF, Nicholson RI, Bundred NJ, Anderson E, Rayter Z, Dowsett M, et al., Comparison of the store-term biological effects of 7alpha-[9-(4,4,5,5,5-pentafluoropentylsulfinyl)-nonyl] estra-1,3,5, (10)-triene-3, 17beta-diol (FASLODEX)
25 versus tamoxifen in postmenopausal women with primary breast cancer. Cancer Res 2001;6:6739-46.

Robertson et al 2003

Robertson JFR, Osborne CK, Howell A, Jones SE, Mauriac L, Ellis M, Kleeberg UR, Come SE, Vergote I, Gertler S, Buzdar A, Webster A, Morris C. Fulvestrant versus
30 anastrozole for the treatment of advanced breast carcinoma in postmenopausal women. A prospective combined analysis of two multicenter trials. Cancer 2003;98(2):229-38.

Wakeling et al 1991

Wakeling AE, Dukes M, Bowler J. A potent specific pure anti-estrogen with clinical potential. *Cancer Research* 1991;51:3867-73.

Whitehead 1989

- 5 Whitehead J. The analysis of relapse clinical trials, with application to a comparison of two ulcer treatments. *Stat Med* 1989; 8(12):1439-54.

Wiebe et al 1993

Weibe VJ, Osborne CK, Fuqua SAW, DE, Gregoria MW. Tamoxifen resistance in breast cancer. *Crit Rev Oncol Hematol* 1993;14:173-88.

Claims:

1. Fulvestrant at a dosage of 500mg for use in the treatment of a postmenopausal woman with advanced breast cancer who has progressed or recurred on endocrine therapy.

5

2. A use according to claim 1 wherein the fulvestrant is administered monthly.

3. A use according to claim 2 wherein an additional dose of 500mg is administered during the first month of treatment.

10

4. A use according to claim 3 wherein the additional dose is administered at about day 14.

5. A use according to any preceding claim wherein the woman is oestrogen receptor positive or progesterone receptor positive.

15

6. A use according claim 5 wherein the woman is oestrogen receptor positive.

7. A use according to any preceding claim wherein the progression or recurrence on endocrine therapy comprised therapy with tamoxifen or an aromatase inhibitor.

20

8. A use according to claim 7 wherein the aromatase inhibitor is selected from anastrozole, letrozole or exemestane.

9. A use according to any preceding claim whereby to increase the time to progression compared with fulvestrant at a dosage of 250mg.

25

30




Figure 1

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2010/051228

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K31/565 A61P35/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BIOSIS, EMBASE, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>BUZDAR AMAN U: "FULVESTRANT - A NOVEL ESTROGEN RECEPTOR ANTAGONIST FOR THE TREATMENT OF ADVANCED BREAST CANCER" DRUGS OF TODAY, vol. 44, no. 9, September 2008 (2008-09), pages 679-692, XP002612295 ISSN: 1699-3993 page 684 page 686, column 2, paragraph 1 page 688, column 2, paragraph 2-3 figure 8</p> <p align="center">----- -/--</p>	1-9
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 2 December 2010		Date of mailing of the international search report 20/12/2010
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Rodríguez-Palmero, M

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2010/051228

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ROBERTSON J F R: "Fulvestrant (Faslodex(R)) - How to make a good drug better" ONCOLOGIST 200707 US LNKD- DOI:10.1634/THEONCOLOGIST.12-7-774, vol. 12, no. 7, July 2007 (2007-07), pages 774-784, XP002612296 ISSN: 1083-7159 figure 1 page 778, column 1, paragraph 2 page 780, column 2, last paragraph page 782, column 1, last paragraph - column 2, paragraph 1</p>	1-9
X	<p>"Comparison of Fulvestrant (Faslodex) 250 mg and 500 mg in postmenopausal women with oestrogen receptor-positive advanced breast cancer progressing or relapsing after previous endocrine therapy." [Online] 20 May 2009 (2009-05-20), XP002612297 Clinicaltrials.gov Retrieved from the Internet: URL: http://clinicaltrials.gov/archive/NCT0099437/2009_05_20 [retrieved on 2010-11-26] the whole document</p>	1-9
X,P	<p>Di Leo A et al.: "CONFIRM: A Phase III, Randomized, Parallel-Group Trial Comparing Fulvestrant 250 mg vs Fulvestrant 500 mg in Postmenopausal Women with Estrogen Receptor-Positive Advanced Breast Cancer." Cancer Res, [Online] vol. 69, no. 24 Suppl 3, 25, 15 December 2009 (2009-12-15), pages 1-2, XP002612298 DOI: 10.1158/0008-5472.SABCS-09-25 Retrieved from the Internet: URL: http://cancerres.aacrjournals.org/cgi/content/abstract/69/24_MeetingAbstracts/25 > [retrieved on 2010-12-01] abstract</p>	1-9

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY
(Chapter I of the Patent Cooperation Treaty)

(PCT Rule 44bis)

Applicant's or agent's file reference 103803-1P WO	FOR FURTHER ACTION		See item 4 below
International application No. PCT/GB2010/051228	International filing date (<i>day/month/year</i>) 26 July 2010 (26.07.2010)	Priority date (<i>day/month/year</i>) 27 July 2009 (27.07.2009)	
International Patent Classification (8th edition unless older edition indicated) See relevant information in Form PCT/ISA/237			
Applicant ASTRAZENECA AB			

1. This international preliminary report on patentability (Chapter I) is issued by the International Bureau on behalf of the International Searching Authority under Rule 44 bis.1(a).

2. 'This REPORT' consists of a total of 7 sheets, including this cover sheet.

In the attached sheets, any reference to the written opinion of the International Searching Authority should be read as a reference to the international preliminary report on patentability (Chapter I) instead.

3. This report contains indications relating to the following items:

<input checked="" type="checkbox"/>	Box No. I	Basis of the report
<input checked="" type="checkbox"/>	Box No. II	Priority
<input type="checkbox"/>	Box No. III	Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
<input type="checkbox"/>	Box No. IV	Lack of unity of invention
<input checked="" type="checkbox"/>	Box No. V	Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
<input checked="" type="checkbox"/>	Box No. VI	Certain documents cited
<input type="checkbox"/>	Box No. VII	Certain defects in the international application
<input checked="" type="checkbox"/>	Box No. VIII	Certain observations on the international application

4. The International Bureau will communicate this report to designated Offices in accordance with Rules 44bis.3(c) and 93bis.1 but not, except where the applicant makes an express request under Article 23(2), before the expiration of 30 months from the priority date (Rule 44bis .2).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No. +41 22 338 82 70	Date of issuance of this report 31 January 2012 (31.01.2012)
	Authorized officer Athina Nickitas-Etienne e-mail: pt04.pct@wipo.int

PATENT COOPERATION TREATY

From the
INTERNATIONAL SEARCHING AUTHORITY

PCT

**WRITTEN OPINION OF THE
INTERNATIONAL SEARCHING AUTHORITY
(PCT Rule 43bis.1)**

To:

see form PCT/ISA/220

Date of mailing
(day/month/year) see form PCT/ISA/210 (second sheet)

Applicant's or agent's file reference
see form PCT/ISA/220

FOR FURTHER ACTION
See paragraph 2 below

International application No.
PCT/GB2010/051228

International filing date (day/month/year)
26.07.2010

Priority date (day/month/year)
27.07.2009

International Patent Classification (IPC) or both national classification and IPC
INV. A61K31/565 A61P35/00

Applicant
ASTRAZENECA AB

1. This opinion contains indications relating to the following items:

- Box No. I Basis of the opinion
- Box No. II Priority
- Box No. III Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- Box No. IV Lack of unity of invention
- Box No. V Reasoned statement under Rule 43bis.1(a)(i) with regard to novelty, inventive step and industrial applicability; citations and explanations supporting such statement
- Box No. VI Certain documents cited
- Box No. VII Certain defects in the international application
- Box No. VIII Certain observations on the international application

2. **FURTHER ACTION**

If a demand for international preliminary examination is made, this opinion will usually be considered to be a written opinion of the International Preliminary Examining Authority ("IPEA") except that this does not apply where the applicant chooses an Authority other than this one to be the IPEA and the chosen IPEA has notified the International Bureau under Rule 66.1bis(b) that written opinions of this International Searching Authority will not be so considered.

If this opinion is, as provided above, considered to be a written opinion of the IPEA, the applicant is invited to submit to the IPEA a written reply together, where appropriate, with amendments, before the expiration of 3 months from the date of mailing of Form PCT/ISA/220 or before the expiration of 22 months from the priority date, whichever expires later.

For further options, see Form PCT/ISA/220.

3. For further details, see notes to Form PCT/ISA/220.

Name and mailing address of the ISA:



European Patent Office
D-80298 Munich
Tel. +49 89 2399 - 0
Fax: +49 89 2399 - 4465


Date of completion of this opinion

see form PCT/ISA/210

Authorized Officer

Rodríguez-Palmero, M

Telephone No. +49 89 2399-7871



**WRITTEN OPINION OF THE
INTERNATIONAL SEARCHING AUTHORITY**

International application No.
PCT/GB2010/051228

Box No. I Basis of the opinion

1. With regard to the **language**, this opinion has been established on the basis of:
 - the international application in the language in which it was filed
 - a translation of the international application into , which is the language of a translation furnished for the purposes of international search (Rules 12.3(a) and 23.1 (b)).
2. This opinion has been established taking into account the **rectification of an obvious mistake** authorized by or notified to this Authority under Rule 91 (Rule 43bis.1(a))
3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, this opinion has been established on the basis of a sequence listing filed or furnished:
 - a. (means)
 - on paper
 - in electronic form
 - b. (time)
 - in the international application as filed
 - together with the international application in electronic form
 - subsequently to this Authority for the purposes of search
4. In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
5. Additional comments:

Box No. II Priority

1. The validity of the priority claim has not been considered because the International Searching Authority does not have in its possession a copy of the earlier application whose priority has been claimed or, where required, a translation of that earlier application. This opinion has nevertheless been established on the assumption that the relevant date (Rules 43*bis*.1 and 64.1) is the claimed priority date.
2. This opinion has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid (Rules 43*bis*.1 and 64.1). Thus for the purposes of this opinion, the international filing date indicated above is considered to be the relevant date.
3. Additional observations, if necessary:

**WRITTEN OPINION OF THE
INTERNATIONAL SEARCHING AUTHORITY**

International application No.
PCT/GB2010/051228

Box No. V Reasoned statement under Rule 43bis.1(a)(i) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes: Claims	<u>1, 5-9</u>
	No: Claims	<u>2-4</u>
Inventive step (IS)	Yes: Claims	
	No: Claims	<u>1-9</u>
Industrial applicability (IA)	Yes: Claims	<u>1-9</u>
	No: Claims	

2. Citations and explanations

see separate sheet

Box No. VI Certain documents cited

1. Certain published documents (Rules 43bis.1 and 70.10)

and / or

2. Non-written disclosures (Rules 43bis.1 and 70.9)

see form 210

Box No. VIII Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

see separate sheet

Re Item V

Reasoned statement with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1 Documents

Reference is made to the following documents; the numbering will be adhered to in the rest of the procedure:

- D1 BUZDAR AMAN U: "FULVESTRANT - A NOVEL ESTROGEN RECEPTOR ANTAGONIST FOR THE TREATMENT OF ADVANCED BREAST CANCER",
DRUGS OF TODAY,
vol. 44, no. 9, September 2008 (2008-09), pages 679-692,
ISSN: 1699-3993
- D2 ROBERTSON J F R: "Fulvestrant (Faslodex(R)) - How to make a good drug better",
ONCOLOGIST 200707 US LNKD- DOI:10.1634/THEONCOLOGIST.
12-7-774,
vol. 12, no. 7, July 2007 (2007-07), pages 774-784,
ISSN: 1083-7159
- D3 "Comparison of Fulvestrant (Faslodex) 250 mg and 500 mg in postmenopausal women with oestrogen receptor-positive advanced breast cancer progressing or relapsing after previous endocrine therapy.",
Clinicaltrials.gov
, 20 May 2009 (2009-05-20),
Retrieved from the Internet:
URL:http://clinicaltrials.gov/archive/NCT00099437/2009_05_20
[retrieved on 2010-11-26]
- D4 Di Leo A et al.: "CONFIRM: A Phase III, Randomized, Parallel-Group Trial Comparing Fulvestrant 250 mg vs Fulvestrant 500 mg in Postmenopausal Women with Estrogen Receptor-Positive Advanced Breast Cancer.",
Cancer Res,
vol. 69, no. 24 Suppl 3, 25, 15 December 2009 (2009-12-15), pages 1-1,
DOI: 10.1158/0008-5472.SABCS-09-25
Retrieved from the Internet:

URL:http://cancerres.aacrjournals.org/cgi/content/abstract/69/24_MeetingAbstracts/25
[retrieved on 2010-12-01]

1.1 Unless indicated, reference is made to the passages indicated in the international search report.

2 Novelty (Art. 33(2) PCT)

2.1 Both D1 and D2 report on the results of the EFECT study. This study evaluated fulvestrant administered in a loading-dose regimen (500 mg at day 0, 250 mg at days 14 and 28, and then 250 mg every 28 days thereafter) for the treatment of postmenopausal women with oestrogen receptor positive advanced breast cancer progressing or relapsing after previous endocrine therapy. The results indicate that the treatment is effective and safe. Since a dose of 500 mg fulvestrant is used in this study, D1 and D2 take away the novelty of present claim 1. The subject-matter of dependent claims 5-8 is also disclosed in D1 or D2 and is thus also not novel (see in particular page 684 in D1 and page 778, column 1, paragraph 2 in D2).

Claim 9 concerns the result of a comparison which has not been done in the EFECT study. However, this result does not characterize the claimed use (fulvestrant at 500 mg for the treatment of postmenopausal women with oestrogen receptor positive breast cancer progressing or relapsing after previous endocrine therapy). Therefore, this claim does not comprise any feature that further characterizes the use defined in claim 1 and anticipated in D1 and D2. Consequently, claim 9 is also not novel in the light of D1 or D2.

3 Inventive Step (Art. 33(3) PCT)

D1-D3 all mention the CONFIRM study, which evaluates the use of fulvestrant at 500 mg/month) for the treatment of postmenopausal women with oestrogen receptor positive advanced breast cancer progressing or relapsing after previous endocrine therapy with anti-oestrogen hormonal treatment such as tamoxifen or an aromatase inhibitor. Therefore, these documents are the closest prior art for the claimed subject-matter. The difference with present claims 1-9 is that the results of this study are not mentioned in D1-D3. The **problem** can therefore be defined as to provide a regimen for the treatment of

breast cancer progressing or relapsing after previous endocrine therapy in postmenopausal women. The **solution** given in present claims 1-9 is to use fulvestrant at 500mg. Since this solution is already mentioned in D1-D3, the person skilled in the art only needed to perform the instructions given in D1-D3 to obtain the results included in the present application. This does not involve an inventive step and Art. 33(3) PCT is thus not fulfilled.

4 Industrial applicability (Art. 33(4) PCT)

Present claims 1-9 are susceptible of industrial application and thus do not contravene Art. 33(4) PCT.

Re Item VI

Certain documents cited

The current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the priority document. If it later turns out that this is not the case, D4 could become relevant for the questions of novelty and/or inventive step of the present patent application.

Re Item VIII

Certain observations on the international application

- 5 The term "advanced" in "advanced breast cancer" in claim 1 is not considered to be clear. This term is vague and there is no general accepted definition in the field, so that the scope of the claim is uncertain. The description does not provide any further clarification of the term. For the purpose of the present opinion, this term has been ignored.
- 6 It appears that the word "whereby" in present claim 9 had to be deleted.

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Carlos M. Tellez
Attorney Docket Number:	11285.0056-00000

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Pages:				
Claims:				
Miscellaneous-Filing:				
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Extension-of-Time:				

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Miscellaneous:				
Submission- Information Disclosure Stmt	1806	1	180	180
Total in USD (\$)				180

Electronic Acknowledgement Receipt

EFS ID:	12500599
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	Formulation
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Carlos M. Tellez
Filer Authorized By:	
Attorney Docket Number:	11285.0056-00000
Receipt Date:	09-APR-2012
Filing Date:	15-OCT-2008
Time Stamp:	17:18:20
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Credit Card
Payment was successfully received in RAM	\$180
RAM confirmation Number	3893
Deposit Account	
Authorized User	

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
-----------------	----------------------	-----------	----------------------------------	------------------	------------------

1		11285-0056-00000--09- APR-2012--Response_20- Mar-2012_OA.pdf	110796 ef73257122a429e023f5524934f990a07139 36f1	yes	4
Multipart Description/PDF files in .zip description					
		Document Description	Start	End	
		Amendment/Req. Reconsideration-After Non-Final Reject	1	1	
		Applicant Arguments/Remarks Made in an Amendment	2	4	
Warnings:					
Information:					
2		11285-0056-00000--09- APR-2012--IDS-SB-08.pdf	130853 d001d0d3f802db7b45efba454f210fd8592 9b3c	yes	3
Multipart Description/PDF files in .zip description					
		Document Description	Start	End	
		Transmittal Letter	1	2	
		Information Disclosure Statement (IDS) Form (SB08)	3	3	
Warnings:					
Information:					
3		11285-0056-00000-09- APR-2012--WO-2011-012885-- spec_for-0059_with_ISR.pdf	1122686 dbb1b7f1640ac4a25c243083dd978bb8794 303bc	yes	27
Multipart Description/PDF files in .zip description					
		Document Description	Start	End	
		Foreign Reference	1	25	
		Non Patent Literature	26	27	
Warnings:					
Information:					
4	Non Patent Literature	11285-0056-00000-09- APR-2012--Fulvestrant-Buzdar. pdf	877405 6e842fa192b8e62c613a5f0b7382c9ed29c2 787e	no	14
Warnings:					
Information:					
5	Non Patent Literature	11285-0056-00000--09- APR-2012- ComparisonofFulvestrant.pdf	103280 1dc5c1fd7b3c3cee08cb7060c69ef6da5c81 1a69	no	3
Warnings:					

Information:					
6	Non Patent Literature	11285-0056-00000--09- APR-2012--DiLeo.pdf	75882 <small>ae06797d8cbf7b1e30df9dccc314a3ffb978f73</small>	no	2
Warnings:					
Information:					
7	Non Patent Literature	11285-0056-00000-09- APR-2012-- IPRP_related_app-0059.pdf	281691 <small>488c6a22ae56ca12023dcf82377994633d06d146</small>	no	7
Warnings:					
Information:					
8	Non Patent Literature	11285-0056-00000--09- APR-2012-- EP12501383rdpartyobservation sreMcLeskeypaper.pdf	1813766 <small>de1d96bb02a6281ed8113271edd52b9c3a8c1c75</small>	no	30
Warnings:					
Information:					
9	Non Patent Literature	11285-0056-00000--09- APR-2012-- EP205138Appealstatement27F eb2012.pdf	2988672 <small>3a22cf51f7e2e38617325437d48f9f47643c379e</small>	no	82
Warnings:					
Information:					
10	Fee Worksheet (SB06)	fee-info.pdf	29656 <small>e0e62a09a961af8a596438098546f778384f3b6</small>	no	2
Warnings:					
Information:					
Total Files Size (in bytes):			7534687		
<p>This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.</p> <p><u>New Applications Under 35 U.S.C. 111</u> If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.</p> <p><u>National Stage of an International Application under 35 U.S.C. 371</u> If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.</p> <p><u>New International Application Filed with the USPTO as a Receiving Office</u> If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.</p>					

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875	Application or Docket Number 12/285,887	Filing Date 10/15/2008	<input type="checkbox"/> To be Mailed
---	---	----------------------------------	---------------------------------------

APPLICATION AS FILED – PART I			OTHER THAN SMALL ENTITY				
	(Column 1)	(Column 2)	SMALL ENTITY <input type="checkbox"/>	OR			
FOR	NUMBER FILED	NUMBER EXTRA	RATE (\$)	FEE (\$)	OR	RATE (\$)	FEE (\$)
<input type="checkbox"/> BASIC FEE <small>(37 CFR 1.16(a), (b), or (c))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> SEARCH FEE <small>(37 CFR 1.16(k), (l), or (m))</small>	N/A	N/A	N/A			N/A	
<input type="checkbox"/> EXAMINATION FEE <small>(37 CFR 1.16(o), (p), or (q))</small>	N/A	N/A	N/A			N/A	
TOTAL CLAIMS <small>(37 CFR 1.16(j))</small>	minus 20 =	*	X \$ =		OR	X \$ =	
INDEPENDENT CLAIMS <small>(37 CFR 1.16(h))</small>	minus 3 =	*	X \$ =			X \$ =	
<input type="checkbox"/> APPLICATION SIZE FEE <small>(37 CFR 1.16(s))</small>	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).						
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENT <small>(37 CFR 1.16(j))</small>							
* If the difference in column 1 is less than zero, enter "0" in column 2.			TOTAL			TOTAL	

APPLICATION AS AMENDED – PART II					OTHER THAN SMALL ENTITY				
	(Column 1)	(Column 2)	(Column 3)		SMALL ENTITY	OR			
AMENDMENT	04/09/2012	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	OR	RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	* 20	Minus	** 30	=	0	OR	X \$60=	0
	Independent <small>(37 CFR 1.16(h))</small>	* 2	Minus	***8	=	0	OR	X \$250=	0
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>						OR		
					TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	0

	(Column 1)	(Column 2)	(Column 3)		SMALL ENTITY	OR			
AMENDMENT		CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)	ADDITIONAL FEE (\$)	OR	RATE (\$)	ADDITIONAL FEE (\$)
	Total <small>(37 CFR 1.16(i))</small>	*	Minus	**	=		OR	X \$ =	
	Independent <small>(37 CFR 1.16(h))</small>	*	Minus	***	=		OR	X \$ =	
	<input type="checkbox"/> Application Size Fee <small>(37 CFR 1.16(s))</small>								
	<input type="checkbox"/> FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM <small>(37 CFR 1.16(j))</small>						OR		
					TOTAL ADD'L FEE		OR	TOTAL ADD'L FEE	

* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.
 ** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".
 *** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".
 The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.

Legal Instrument Examiner:
 /MARGARET BYARS/

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**
 If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.



NOTICE OF ALLOWANCE AND FEE(S) DUE

22852 7590 06/04/2012
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

Table with 2 columns: EXAMINER (HUI, SAN MING R), ART UNIT (1628), PAPER NUMBER.

DATE MAILED: 06/04/2012

Table with 5 columns: APPLICATION NO. (12/285,887), FILING DATE (10/15/2008), FIRST NAMED INVENTOR (John R. Evans), ATTORNEY DOCKET NO. (11285.0056-00000), CONFIRMATION NO. (1199)

TITLE OF INVENTION: FORMULATION

Table with 7 columns: APPLN. TYPE (nonprovisional), SMALL ENTITY (NO), ISSUE FEE DUE (\$1740), PUBLICATION FEE DUE (\$300), PREV. PAID ISSUE FEE (\$0), TOTAL FEE(S) DUE (\$2040), DATE DUE (09/04/2012)

THE APPLICATION IDENTIFIED ABOVE HAS BEEN EXAMINED AND IS ALLOWED FOR ISSUANCE AS A PATENT. PROSECUTION ON THE MERITS IS CLOSED. THIS NOTICE OF ALLOWANCE IS NOT A GRANT OF PATENT RIGHTS. THIS APPLICATION IS SUBJECT TO WITHDRAWAL FROM ISSUE AT THE INITIATIVE OF THE OFFICE OR UPON PETITION BY THE APPLICANT. SEE 37 CFR 1.313 AND MPEP 1308.

THE ISSUE FEE AND PUBLICATION FEE (IF REQUIRED) MUST BE PAID WITHIN THREE MONTHS FROM THE MAILING DATE OF THIS NOTICE OR THIS APPLICATION SHALL BE REGARDED AS ABANDONED. THIS STATUTORY PERIOD CANNOT BE EXTENDED. SEE 35 U.S.C. 151. THE ISSUE FEE DUE INDICATED ABOVE DOES NOT REFLECT A CREDIT FOR ANY PREVIOUSLY PAID ISSUE FEE IN THIS APPLICATION. IF AN ISSUE FEE HAS PREVIOUSLY BEEN PAID IN THIS APPLICATION (AS SHOWN ABOVE), THE RETURN OF PART B OF THIS FORM WILL BE CONSIDERED A REQUEST TO REAPPLY THE PREVIOUSLY PAID ISSUE FEE TOWARD THE ISSUE FEE NOW DUE.

HOW TO REPLY TO THIS NOTICE:

I. Review the SMALL ENTITY status shown above.

If the SMALL ENTITY is shown as YES, verify your current SMALL ENTITY status:

- A. If the status is the same, pay the TOTAL FEE(S) DUE shown above.
B. If the status above is to be removed, check box 5b on Part B - Fee(s) Transmittal and pay the PUBLICATION FEE (if required) and twice the amount of the ISSUE FEE shown above, or

If the SMALL ENTITY is shown as NO:

- A. Pay TOTAL FEE(S) DUE shown above, or
B. If applicant claimed SMALL ENTITY status before, or is now claiming SMALL ENTITY status, check box 5a on Part B - Fee(s) Transmittal and pay the PUBLICATION FEE (if required) and 1/2 the ISSUE FEE shown above.

II. PART B - FEE(S) TRANSMITTAL, or its equivalent, must be completed and returned to the United States Patent and Trademark Office (USPTO) with your ISSUE FEE and PUBLICATION FEE (if required). If you are charging the fee(s) to your deposit account, section "4b" of Part B - Fee(s) Transmittal should be completed and an extra copy of the form should be submitted. If an equivalent of Part B is filed, a request to reapply a previously paid issue fee must be clearly made, and delays in processing may occur due to the difficulty in recognizing the paper as an equivalent of Part B.

III. All communications regarding this application must give the application number. Please direct all communications prior to issuance to Mail Stop ISSUE FEE unless advised to the contrary.

IMPORTANT REMINDER: Utility patents issuing on applications filed on or after Dec. 12, 1980 may require payment of maintenance fees. It is patentee's responsibility to ensure timely payment of maintenance fees when due.

PART B - FEE(S) TRANSMITTAL

**Complete and send this form, together with applicable fee(s), to: Mail Mail Stop ISSUE FEE
 Commissioner for Patents
 P.O. Box 1450
 Alexandria, Virginia 22313-1450
 or Fax (571)-273-2885**

INSTRUCTIONS: This form should be used for transmitting the ISSUE FEE and PUBLICATION FEE (if required). Blocks 1 through 5 should be completed where appropriate. All further correspondence including the Patent, advance orders and notification of maintenance fees will be mailed to the current correspondence address as indicated unless corrected below or directed otherwise in Block 1, by (a) specifying a new correspondence address; and/or (b) indicating a separate "FEE ADDRESS" for maintenance fee notifications.

CURRENT CORRESPONDENCE ADDRESS (Note: Use Block 1 for any change of address)

Note: A certificate of mailing can only be used for domestic mailings of the Fee(s) Transmittal. This certificate cannot be used for any other accompanying papers. Each additional paper, such as an assignment or formal drawing, must have its own certificate of mailing or transmission.

22852 7590 06/04/2012

**FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
 LLP
 901 NEW YORK AVENUE, NW
 WASHINGTON, DC 20001-4413**

Certificate of Mailing or Transmission

I hereby certify that this Fee(s) Transmittal is being deposited with the United States Postal Service with sufficient postage for first class mail in an envelope addressed to the Mail Stop ISSUE FEE address above, or being facsimile transmitted to the USPTO (571) 273-2885, on the date indicated below.

(Depositor's name)
(Signature)
(Date)

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
-----------------	-------------	----------------------	---------------------	------------------

12/285,887 10/15/2008 John R. Evans 11285.0056-00000 1199

TITLE OF INVENTION: FORMULATION

APPLN. TYPE	SMALL ENTITY	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
-------------	--------------	---------------	---------------------	----------------------	------------------	----------

nonprovisional NO \$1740 \$300 \$0 \$2040 09/04/2012

EXAMINER	ART UNIT	CLASS-SUBCLASS
----------	----------	----------------

HUI, SAN MING R 1628 514-177000

<p>1. Change of correspondence address or indication of "Fee Address" (37 CFR 1.363).</p> <p><input type="checkbox"/> Change of correspondence address (or Change of Correspondence Address form PTO/SB/122) attached.</p> <p><input type="checkbox"/> "Fee Address" indication (or "Fee Address" Indication form PTO/SB/47; Rev 03-02 or more recent) attached. Use of a Customer Number is required.</p>	<p>2. For printing on the patent front page, list</p> <p>(1) the names of up to 3 registered patent attorneys or agents OR, alternatively, 1 _____</p> <p>(2) the name of a single firm (having as a member a registered attorney or agent) and the names of up to 2 registered patent attorneys or agents. If no name is listed, no name will be printed. 2 _____</p> <p>3 _____</p>
---	---

3. ASSIGNEE NAME AND RESIDENCE DATA TO BE PRINTED ON THE PATENT (print or type)

PLEASE NOTE: Unless an assignee is identified below, no assignee data will appear on the patent. If an assignee is identified below, the document has been filed for recordation as set forth in 37 CFR 3.11. Completion of this form is NOT a substitute for filing an assignment.

(A) NAME OF ASSIGNEE (B) RESIDENCE: (CITY and STATE OR COUNTRY)

Please check the appropriate assignee category or categories (will not be printed on the patent) : Individual Corporation or other private group entity Government

<p>4a. The following fee(s) are submitted:</p> <p><input type="checkbox"/> Issue Fee</p> <p><input type="checkbox"/> Publication Fee (No small entity discount permitted)</p> <p><input type="checkbox"/> Advance Order - # of Copies _____</p>	<p>4b. Payment of Fee(s): (Please first reapply any previously paid issue fee shown above)</p> <p><input type="checkbox"/> A check is enclosed.</p> <p><input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.</p> <p><input type="checkbox"/> The Director is hereby authorized to charge the required fee(s), any deficiency, or credit any overpayment, to Deposit Account Number _____ (enclose an extra copy of this form).</p>
---	--

5. Change in Entity Status (from status indicated above)

a. Applicant claims SMALL ENTITY status. See 37 CFR 1.27. b. Applicant is no longer claiming SMALL ENTITY status. See 37 CFR 1.27(g)(2).

NOTE: The Issue Fee and Publication Fee (if required) will not be accepted from anyone other than the applicant; a registered attorney or agent; or the assignee or other party in interest as shown by the records of the United States Patent and Trademark Office.

Authorized Signature _____ Date _____
 Typed or printed name _____ Registration No. _____

This collection of information is required by 37 CFR 1.311. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, Virginia 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 5 columns: APPLICATION NO., FILING DATE, FIRST NAMED INVENTOR, ATTORNEY DOCKET NO., CONFIRMATION NO.
12/285,887 10/15/2008 John R. Evans 11285.0056-00000 1199

22852 7590 06/04/2012
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

EXAMINER

HUI, SAN MING R

ART UNIT PAPER NUMBER

1628

DATE MAILED: 06/04/2012

Determination of Patent Term Adjustment under 35 U.S.C. 154 (b)
(application filed on or after May 29, 2000)

The Patent Term Adjustment to date is 248 day(s). If the issue fee is paid on the date that is three months after the mailing date of this notice and the patent issues on the Tuesday before the date that is 28 weeks (six and a half months) after the mailing date of this notice, the Patent Term Adjustment will be 248 day(s).

If a Continued Prosecution Application (CPA) was filed in the above-identified application, the filing date that determines Patent Term Adjustment is the filing date of the most recent CPA.

Applicant will be able to obtain more detailed information by accessing the Patent Application Information Retrieval (PAIR) WEB site (http://pair.uspto.gov).

Any questions regarding the Patent Term Extension or Adjustment determination should be directed to the Office of Patent Legal Administration at (571)-272-7702. Questions relating to issue and publication fee payments should be directed to the Customer Service Center of the Office of Patent Publication at 1-(888)-786-0101 or (571)-272-4200.

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. Accordingly, pursuant to the requirements of the Act, please be advised that: (1) the general authority for the collection of this information is 35 U.S.C. 2(b)(2); (2) furnishing of the information solicited is voluntary; and (3) the principal purpose for which the information is used by the U.S. Patent and Trademark Office is to process and/or examine your submission related to a patent application or patent. If you do not furnish the requested information, the U.S. Patent and Trademark Office may not be able to process and/or examine your submission, which may result in termination of proceedings or abandonment of the application or expiration of the patent.

The information provided by you in this form will be subject to the following routine uses:

1. The information on this form will be treated confidentially to the extent allowed under the Freedom of Information Act (5 U.S.C. 552) and the Privacy Act (5 U.S.C. 552a). Records from this system of records may be disclosed to the Department of Justice to determine whether disclosure of these records is required by the Freedom of Information Act.
2. A record from this system of records may be disclosed, as a routine use, in the course of presenting evidence to a court, magistrate, or administrative tribunal, including disclosures to opposing counsel in the course of settlement negotiations.
3. A record in this system of records may be disclosed, as a routine use, to a Member of Congress submitting a request involving an individual, to whom the record pertains, when the individual has requested assistance from the Member with respect to the subject matter of the record.
4. A record in this system of records may be disclosed, as a routine use, to a contractor of the Agency having need for the information in order to perform a contract. Recipients of information shall be required to comply with the requirements of the Privacy Act of 1974, as amended, pursuant to 5 U.S.C. 552a(m).
5. A record related to an International Application filed under the Patent Cooperation Treaty in this system of records may be disclosed, as a routine use, to the International Bureau of the World Intellectual Property Organization, pursuant to the Patent Cooperation Treaty.
6. A record in this system of records may be disclosed, as a routine use, to another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)).
7. A record from this system of records may be disclosed, as a routine use, to the Administrator, General Services, or his/her designee, during an inspection of records conducted by GSA as part of that agency's responsibility to recommend improvements in records management practices and programs, under authority of 44 U.S.C. 2904 and 2906. Such disclosure shall be made in accordance with the GSA regulations governing inspection of records for this purpose, and any other relevant (i.e., GSA or Commerce) directive. Such disclosure shall not be used to make determinations about individuals.
8. A record from this system of records may be disclosed, as a routine use, to the public after either publication of the application pursuant to 35 U.S.C. 122(b) or issuance of a patent pursuant to 35 U.S.C. 151. Further, a record may be disclosed, subject to the limitations of 37 CFR 1.14, as a routine use, to the public if the record was filed in an application which became abandoned or in which the proceedings were terminated and which application is referenced by either a published application, an application open to public inspection or an issued patent.
9. A record from this system of records may be disclosed, as a routine use, to a Federal, State, or local law enforcement agency, if the USPTO becomes aware of a violation or potential violation of law or regulation.

Notice of Allowability

Application No.

12/285,887

Examiner

SAN-MING HUI

Applicant(s)

EVANS ET AL.

Art Unit

1628

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address--

All claims being allowable, PROSECUTION ON THE MERITS IS (OR REMAINS) CLOSED in this application. If not included herewith (or previously mailed), a Notice of Allowance (PTOL-85) or other appropriate communication will be mailed in due course. **THIS NOTICE OF ALLOWABILITY IS NOT A GRANT OF PATENT RIGHTS.** This application is subject to withdrawal from issue at the initiative of the Office or upon petition by the applicant. See 37 CFR 1.313 and MPEP 1308.

- 1. This communication is responsive to 4/9/2012.
- 2. An election was made by the applicant in response to a restriction requirement set forth during the interview on ____; the restriction requirement and election have been incorporated into this action.
- 3. The allowed claim(s) is/are 24,26,27,29,30,32,34-36,38,39,41,42,44,46,47 and 54-57.
- 4. Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
 - a) All b) Some* c) None of the:
 - 1. Certified copies of the priority documents have been received.
 - 2. Certified copies of the priority documents have been received in Application No. 10/872784 .
 - 3. Copies of the certified copies of the priority documents have been received in this national stage application from the International Bureau (PCT Rule 17.2(a)).

* Certified copies not received: ____.

Applicant has THREE MONTHS FROM THE "MAILING DATE" of this communication to file a reply complying with the requirements noted below. Failure to timely comply will result in ABANDONMENT of this application.

THIS THREE-MONTH PERIOD IS NOT EXTENDABLE.

- 5. A SUBSTITUTE OATH OR DECLARATION must be submitted. Note the attached EXAMINER'S AMENDMENT or NOTICE OF INFORMAL PATENT APPLICATION (PTO-152) which gives reason(s) why the oath or declaration is deficient.
 - 6. CORRECTED DRAWINGS (as "replacement sheets") must be submitted.
 - (a) including changes required by the Notice of Draftsperson's Patent Drawing Review (PTO-948) attached
 - 1) hereto or 2) to Paper No./Mail Date ____.
 - (b) including changes required by the attached Examiner's Amendment / Comment or in the Office action of Paper No./Mail Date ____.
- Identifying indicia such as the application number (see 37 CFR 1.84(c)) should be written on the drawings in the front (not the back) of each sheet. Replacement sheet(s) should be labeled as such in the header according to 37 CFR 1.121(d).**
- 7. DEPOSIT OF and/or INFORMATION about the deposit of BIOLOGICAL MATERIAL must be submitted. Note the attached Examiner's comment regarding REQUIREMENT FOR THE DEPOSIT OF BIOLOGICAL MATERIAL.

Attachment(s)

- 1. Notice of References Cited (PTO-892)
- 2. Notice of Draftsperson's Patent Drawing Review (PTO-948)
- 3. Information Disclosure Statements (PTO/SB/08), Paper No./Mail Date 4/9/12, 6/4/12
- 4. Examiner's Comment Regarding Requirement for Deposit of Biological Material
- 5. Notice of Informal Patent Application
- 6. Interview Summary (PTO-413), Paper No./Mail Date ____ .
- 7. Examiner's Amendment/Comment
- 8. Examiner's Statement of Reasons for Allowance
- 9. Other ____.

/San-ming Hui/
Primary Examiner, Art Unit 1628

DETAILED ACTION

Applicant's response filed 4/9/2012 has been entered. Claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 are pending.

The outstanding double patenting rejection is withdrawn in view of the terminal disclaimer filed 1/17/2012.

REASONS FOR ALLOWANCE

The following is an examiner's statement of reasons for allowance: The outstanding rejection is withdrawn and therefore, the pending claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 are in condition of allowance.

Any comments considered necessary by applicant must be submitted no later than the payment of the issue fee and, to avoid processing delays, should preferably accompany the issue fee. Such submissions should be clearly labeled "Comments on Statement of Reasons for Allowance."

Allowable Subject Matter

Claims 24, 26, 27, 29, 30, 32, 34-36, 38, 39, 41, 42, 44, 46, 47, and 54-57 are allowed.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SAN-MING HUI whose telephone number is (571)272-0626. The examiner can normally be reached on Mon - Fri from 9:00 to 5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Brandon Fetterolf can be reached on (571) 272-2919. The fax phone

Art Unit: 1628

number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

San-ming Hui
Primary Examiner
Art Unit 1628

/San-ming Hui/
Primary Examiner, Art Unit 1628

Index of Claims 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed

-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011	03/09/2012	04/17/2012				
	1	✓							
	2	✓							
	3	✓							
	4	✓							
	5	✓							
	6	✓							
	7	✓							
	8	✓							
	9	✓							
	10	✓							
	11	✓							
	12	✓							
	13	✓							
	14	✓							
	15	✓							
	16	✓							
	17	✓							
	18	✓							
	19	✓							
	20	✓							
	21	✓							
	22	✓							
	23	✓							
	24		✓	✓	=				
	25		✓	-					
	26		✓	✓	=				
	27		✓	✓	=				
	28		✓	-					
	29		✓	✓	=				
	30		✓	✓	=				
	31		✓	-					
	32		✓	✓	=				
	33		✓	-					
	34		✓	✓	=				
	35		✓	✓	=				
	36		✓	✓	=				

Index of Claims 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

✓	Rejected
=	Allowed


-	Cancelled
÷	Restricted

N	Non-Elected
I	Interference

A	Appeal
O	Objected

Claims renumbered in the same order as presented by applicant
 CPA
 T.D.
 R.1.47

CLAIM		DATE							
Final	Original	12/19/2010	09/06/2011	03/09/2012	04/17/2012				
	37		✓	-					
	38		✓	✓	=				
	39		✓	✓	=				
	40		✓	-					
	41		✓	✓	=				
	42		✓	✓	=				
	43		✓	-					
	44		✓	✓	=				
	45		✓	-					
	46		✓	✓	=				
	47		✓	✓	=				
	48		✓	-					
	49		✓	-					
	50		✓	-					
	51		✓	-					
	52		✓	-					
	53		✓	-					
	54			✓	=				
	55			✓	=				
	56			✓	=				
	57			✓	=				

Search Notes 	Application/Control No. 12285887	Applicant(s)/Patent Under Reexamination EVANS ET AL.
	Examiner San-ming Hui	Art Unit 1628

SEARCHED			
Class	Subclass	Date	Examiner
514	177, 178	12/19/10	SH
514	177, 178	9/6/11	SH
514	177, 178	3/9/12	SH
514	177, 178	4/17/12	SH

SEARCH NOTES		
Search Notes	Date	Examiner
EAST and inventor search in PALM	12/19/10	SH
update search in EAST and inventor search in PALM	9/6/11	SH
EAST in EAST and inventor search in PALM	3/9/12	SH
update search in EAST and inventor search in PALM	4/17/12	SH

INTERFERENCE SEARCH			
Class	Subclass	Date	Examiner
514	177, 178	4/17/12	SH

--	--

INFORMATION DISCLOSURE CITATION (Use several sheets if necessary) PTO Form 1449 June 4, 2009	Attorney Docket No. 056291-5004-02	Application No. 12/285,887
	Applicants: John R. EVANS et al.	
	Filing Date: October 15, 2008	Group Art Unit: 1617

U.S. PATENT DOCUMENTS

Initial	Document No.	Date	Name	Class	Sub-Class	Filing Date
	1. US 3,164,520	January 5, 1965	Huber			
	2. US 4,212,863	July 15, 1980	Cornelius			
	3. US 4,388,307	June 14, 1983	Cavanak			

FOREIGN PATENT DOCUMENTS

	Document No.	Date	Country	Class	Sub-Class	Translation
	4. EP 0310542A1	April 5, 1989	EPO			Yes

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)

	5.	P.K. Gupta and G.A. Brazeau (eds). <i>Injectable Drug Development: Techniques to Reduce Pain and Irritation</i> . Chapters 11 & 17 Interpharm Press, Denver, Colorado (1999)
	6.	P.V. Lopatin, V. P. Safonov, T. P. Litvinova and L. M. Yakimenko. Use of nonaqueous solvents to prepare injection solutions. <i>Pharm. Chem. J.</i> 6 :724-733 (1972)
	7.	S. Nema, R.J. Washkuhn, and R.J. Brendel. Excipients and their use in injectable products. <i>PDA J. Pharm. Sci. Technol.</i> 51 :166-71 (1997)
	8.	<i>Physicians' Desk Reference (27th edition)</i> . 1277-1278, 1350-1354, 1391-1392 Medical Economics Company, Oradell, NJ (1973)
	9.	M. F. Powell, T. Nguyen, and L. Baloian. Compendium of excipients for parenteral formulations. <i>PDA J. Pharm. Sci. Technol.</i> 52 :238-311 [pages 238-255 provided] (1998)
	10.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) -Part I. <i>PDA J. Pharm. Sci. Technol.</i> 53 :324-349 (1999)
	11.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part II <i>PDA J. Pharm. Sci. Technol.</i> 54 :69-96 (2000)
	12.	R. G. Strickley. Parenteral formulations of small molecule therapeutics marketed in the United States (1999) - Part III. <i>PDA J. Pharm. Sci. Technol.</i> 54 :152-169 (2000)
	13.	Y.C. J. Wang and R. R. Kowal. Review of excipients and pH's for parenteral products used in the United States. <i>J. Parenteral Drug Assoc.</i> 34 :452-462 (1980).

Examiner	/San Ming Hui/	Date Considered	/S.H./
----------	----------------	-----------------	--------

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

INFORMATION DISCLOSURE STATEMENT BY APPLICANT <i>(Use as many sheets as necessary)</i>				Complete if Known			
				Application Number		12/285,887	
				Filing Date		October 15,2008	
				First Named Inventor		John R. EVANS	
				Art Unit		1628	
Examiner Name		HUI, San Ming R.					
Sheet	1	of	1	Attorney Docket Number		11285.0056-00000	

U.S. PATENTS AND PUBLISHED U.S. PATENT APPLICATIONS						
Examiner Initials	Cite No. ¹	Document Number		Issue or Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear
		Number-Kind Code ² (if known)				
	1	U.S. Application No. 13/387,584		Filing date: 27-Jan-2012	DIMERY et al.	

Note: Submission of copies of U.S. Patents and published U.S. Patent Applications is not required.

FOREIGN PATENT DOCUMENTS								
Examiner Initials	Cite No. ¹	Foreign Patent Document			Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages or Relevant Figures Appear	Translation ⁶
		Country Code ³	Number ⁴	Kind Code ⁵ (if known)				
	2	WO	2011/012885		03-Feb-2011	AstraZeneca UK Ltd.		

NONPATENT LITERATURE DOCUMENTS					
Examiner Initials	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.			Translation ⁶
	3	Buzdar, A. U., "Fulvestrant - A novel estrogen receptor antagonist for the treatment of advanced breast cancer," <i>Drugs of Today</i> , 44(9):679-692 (2008).			
	4	"Comparison of fulvestrant (faslodex) 250 mg and 500 mg in postmenopausal women with estrogen receptor-positive advanced breast cancer progressing or relapsing after previous endocrine therapy," <i>Clinicaltrials.gov</i> (20-May-2009) retrieved 24-Jan- 2012.			
	5	Di Leo A., et al., "Confirm: a phase III, randomized, parallel-group trial comparing fulvestrant 250 mg vs fulvestrant 500 mg in postmenopausal women with estrogen receptor-positive advanced breast cancer," <i>Cancer Res.</i> , 69(24) Supp. 3, (2009).			
	6	International Search Report for PCT Application No. PCT/GB10/51228 (WO 2011/012885) mailed December 20, 2012.			
	7	International Preliminary Report on Patentability for PCT Application No. PCT/GB10/51228 (WO 2011/012885) mailed December 20, 2012.			
	8	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) dated December 15, 2011.			
	9	Documents from the prosecution of European Application No. 01900186.6 (EP 1 250 138) dated February 27, 2012.			
Examiner Signature	/San Ming Hui/			Date Considered	04/17/2012

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /S.H./

EAST Search History

EAST Search History (Prior Art)

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
S264	86836	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S265	459	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S266	2864	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S267	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S268	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S269	16301	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S270	1895366	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S271	8503	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S272	4	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S273	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S274	14	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S275	2008	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47

S276	2	"6774122".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S277	982	514/177.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S278	1426	514/178.ccls.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S279	2183348	castor oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S280	86836	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S281	459	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S282	2864	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S283	16301	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S284	1895366	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S285	8503	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S286	7	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (testosterone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S287	14	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S288	2008	((((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S289	86836	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S290	5214	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT;	OR	ON	2012/04/16 14:47

			IBM_TDB			
S291	2934	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S292	1513	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S293	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S294	3712	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S295	3712	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S296	86836	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S297	459	fulvestrant and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S298	2864	oil and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S299	3	"4659516".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S300	7	"346014".pn.	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S301	16301	(benzyl adj benzoate) or (phenylmethyl adj benzoate)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S302	1895366	solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S303	8503	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S304	4	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (estrogen or estradiol or estrone)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16: 14:47
S305	7	((benzyl adj benzoate) or	US-PGPUB;	OR	ON	2012/04/16:

		(phenylmethyl adj benzoate) same solvent) same (testosterone)	USPAT; EPO; JPO; DERWENT; IBM_TDB			14:47
S306	14	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) same (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S307	2008	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same solvent) and (steroid)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S308	86836	castor adj oil	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S309	5214	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) and (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S310	2934	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S311	1513	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S312	3	((benzyl adj benzoate) or (phenylmethyl adj benzoate)) same (castor adj oil)) same solvent) same steroid	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S313	3712	fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S314	97131	breast adj cancer	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S315	2541	breast adj cancer and fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S316	368	breast adj cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47
S317	1579	cancer same fulvestrant	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2012/04/16 14:47

EAST Search History (Interference)

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	973	514/177.ccls.	US-PGPUB; USPAT	OR	ON	2012/04/17 09:28
L2	1395	514/178.ccls.	US-PGPUB; USPAT	OR	ON	2012/04/17 09:28

4/ 17/ 2012 9:28:47 AM

C:\Users\shui\Documents\EAST\Workspaces\12-285887.wsp



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

Table with 5 columns: APPLICATION NO., FILING DATE, FIRST NAMED INVENTOR, ATTORNEY DOCKET NO., CONFIRMATION NO.
12/285,887 10/15/2008 John R. Evans 11285.0056-00000 1199

22852 7590 07/13/2012
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

EXAMINER

HUI, SAN MING R

ART UNIT PAPER NUMBER

1628

MAIL DATE DELIVERY MODE

07/13/2012

PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.



UNITED STATES PATENT AND TRADEMARK OFFICE

Commissioner for Patents
United States Patent and Trademark Office
P. O. Box 1450
Alexandria, VA 22313-1450
www.uspto.gov

Application No. : 12285887
Applicant : Evans
Filing Date : 10/15/2008
Date Mailed : 07/13/2012

NOTICE TO FILE CORRECTED APPLICATION PAPERS

Notice of Allowance Mailed

This application has been accorded an Allowance Date and is being prepared for issuance. The application, however, is incomplete for the reasons below.

Applicant is given 1 month(s) from the mail date of this Notice, or the time remaining from the Notice of Allowance and Fee(s) Due, whichever is longer, within which to respond.

The informalities requiring correction are indicated in the attachment(s). If the informality pertains to the abstract, specification (including claims) or drawings, the informality must be corrected with an amendment in compliance with 37 CFR 1.121 (or, if the application is a reissue application, 37 CFR 1.173). Such an amendment may be filed after payment of the issue fee if limited to correction of informalities noted herein. See Waiver of 37 CFR 1.312 for Documents Required by the Office of Patent Publication, 1280 Off. Gaz. Patent Office 918 (March 23, 2004). In addition, if the informality is not corrected until after payment of the issue fee, for purposes of 35 U.S.C. 154(b)(1)(iv), "all outstanding requirements" will be considered to have been satisfied when the informality has been corrected. A failure to respond within the above-identified time period will result in the application being ABANDONED. **This period for reply is NOT extendable under 37 CFR 1.136(a).**

See attachment(s).

*A copy of this notice **MUST** be returned with the reply. Please address response to "Mail Stop Issue Fee, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450".*

/Patricia Pacenski/
Publication Branch
Office of Data Management
(571) 272-4200

IDENTIFICATION OF APPLICATION DEFICIENCIES

- Applicant must provide legible text for the following item(s).
 - Specification filed , page(s) .
 - Claims filed , claim(s) .
 - Oath/declaration filed .
 - Other: .
- Applicant must provide missing information on the following page(s) of the specification by amending the specification to add the missing text. No new matter may be added. line 1 of page 6 appears to be incomplete
- The specification refers to one or more applications by attorney docket number and does not show the U.S. application number(s). Applicant must supply the U.S. application number in place of each attorney docket number.
- Applicant must provide an Abstract of the Disclosure.
- Applicant has submitted a DECLARATION (37 CFR 1.63) FOR A UTILITY OR DESIGN APPLICATION USING AN APPLICATION DATA SHEET (37 CFR 1.76) (e.g., form PTO/SB/01A). The Application Data Sheet, however, is not present with the filed application. Applicant must submit an Application Data Sheet or file a new oath or declaration (e.g., PTO/SB/01) executed by the inventors and containing the information required in 37 CFR 1.63.
- Applicant must provide an executed declaration.
- Applicant must provide the missing page(s) of the oath/declaration or Application Data Sheet filed
- Applicant must provide a declaration signed by inventor(s) .
- The oath/declaration filed shows non-initialed and/or non-dated alterations. Applicant must file a new oath/declaration in compliance with 37 CFR 1.67(a).
- Applicant(s) in the latest-filed oath/declaration or Application Data Sheet (ADS) did not show the inventor's residence at all, or did not show both a city and state in the U.S. inventor's residence, or did not show both a city and country in the non-U.S. inventor's residence. Applicant must supply an oath/declaration or Application Data Sheet (ADS) that shows each U.S. inventor's city and state of residence and each non-U.S. inventor's city and country of residence.

PART B - FEE(S) TRANSMITTAL

**Complete and send this form, together with applicable fee(s), to: Mail Mail Stop ISSUE FEE
 Commissioner for Patents
 P.O. Box 1450
 Alexandria, Virginia 22313-1450
 or Fax (571)-273-2885**

INSTRUCTIONS: This form should be used for transmitting the ISSUE FEE and PUBLICATION FEE (if required). Blocks 1 through 5 should be completed where appropriate. All further correspondence including the Patent, advance orders and notification of maintenance fees will be mailed to the current correspondence address as indicated unless corrected below or directed otherwise in Block 1, by (a) specifying a new correspondence address; and/or (b) indicating a separate "FEE ADDRESS" for maintenance fee notifications.

CURRENT CORRESPONDENCE ADDRESS (Note: Use Block 1 for any change of address)

Note: A certificate of mailing can only be used for domestic mailings of the Fee(s) Transmittal. This certificate cannot be used for any other accompanying papers. Each additional paper, such as an assignment or formal drawing, must have its own certificate of mailing or transmission.

22852 7590 06/04/2012

**FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
 LLP
 901 NEW YORK AVENUE, NW
 WASHINGTON, DC 20001-4413**

Certificate of Mailing or Transmission

I hereby certify that this Fee(s) Transmittal is being deposited with the United States Postal Service with sufficient postage for first class mail in an envelope addressed to the Mail Stop ISSUE FEE address above, or being facsimile transmitted to the USPTO (571) 273-2885, on the date indicated below.

(Depositor's name)
(Signature)
(Date)

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
12/285,887	10/15/2008	John R. Evans	11285.0056-00000	1199

TITLE OF INVENTION: FORMULATION

APPLN. TYPE	SMALL ENTITY	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
nonprovisional	NO	\$1740	\$300	\$0	\$2040	09/04/2012

EXAMINER	ART UNIT	CLASS-SUBCLASS
HUI, SAN MING R	1628	514-177000

1. Change of correspondence address or indication of "Fee Address" (37 CFR 1.363).
 Change of correspondence address (or Change of Correspondence Address form PTO/SB/122) attached.
 "Fee Address" indication (or "Fee Address" Indication form PTO/SB/47; Rev 03-02 or more recent) attached. Use of a **Customer Number is required.**

2. For printing on the patent front page, list
 (1) the names of up to 3 registered patent attorneys or agents OR, alternatively,
 (2) the name of a single firm (having as a member a registered attorney or agent) and the names of up to 2 registered patent attorneys or agents. If no name is listed, no name will be printed.
 1 Finnegan, Henderson,
 2 Farabow, Garrett &
 3 Dunner LLP

3. ASSIGNEE NAME AND RESIDENCE DATA TO BE PRINTED ON THE PATENT (print or type)

PLEASE NOTE: Unless an assignee is identified below, no assignee data will appear on the patent. If an assignee is identified below, the document has been filed for recordation as set forth in 37 CFR 3.11. Completion of this form is NOT a substitute for filing an assignment.

(A) NAME OF ASSIGNEE Astrazeneca AB (B) RESIDENCE: (CITY and STATE OR COUNTRY) Södertälje, Sweden

Please check the appropriate assignee category or categories (will not be printed on the patent): Individual Corporation or other private group entity Government

4a. The following fee(s) are submitted:

Issue Fee
 Publication Fee (No small entity discount permitted)
 Advance Order - # of Copies _____


4b. Payment of Fee(s): (Please first reapply any previously paid issue fee shown above)

A check is enclosed.
 Payment by credit card. ~~Payment by credit card~~ EFS Web
 The Director is hereby authorized to charge the ~~FEES~~ FEES any deficiency, or credit any overpayment, to Deposit Account Number 06-0916 (enclose an extra copy of this form).

5. Change in Entity Status (from status indicated above)

a. Applicant claims SMALL ENTITY status. See 37 CFR 1.27. b. Applicant is no longer claiming SMALL ENTITY status. See 37 CFR 1.27(g)(2).

NOTE: The Issue Fee and Publication Fee (if required) will not be accepted from anyone other than the applicant; a registered attorney or agent; or the assignee or other party in interest as shown by the records of the United States Patent and Trademark Office.

Authorized Signature 
 Typed or printed name Carlos M. Tellez

Date September 4, 2012
 Registration No. 48,638

This collection of information is required by 37 CFR 1.311. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, Virginia 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450.


Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Please grant any extensions of time required to enter this response and charge any required fees not included with this Response to Deposit Account No. 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated: September 4, 2012

By: 

Carlos M. Téllez
Reg. No. 48,638
(202) 408-4123



UNITED STATES PATENT AND TRADEMARK OFFICE

Commissioner for Patents
United States Patent and Trademark Office
P.O. Box 1450
Alexandria, VA 22313-1450
www.uspto.gov

Application No. : 12285887
Applicant : Evans
Filing Date : 10/15/2008
Date Mailed : 07/13/2012

NOTICE TO FILE CORRECTED APPLICATION PAPERS

Notice of Allowance Mailed

This application has been accorded an Allowance Date and is being prepared for issuance. The application, however, is incomplete for the reasons below.

Applicant is given 1 month(s) from the mail date of this Notice, or the time remaining from the Notice of Allowance and Fee(s) Due, whichever is longer, within which to respond.

The informalities requiring correction are indicated in the attachment(s). If the informality pertains to the abstract, specification (including claims) or drawings, the informality must be corrected with an amendment in compliance with 37 CFR 1.121 (or, if the application is a reissue application, 37 CFR 1.173). Such an amendment may be filed after payment of the issue fee if limited to correction of informalities noted herein. See Waiver of 37 CFR 1.312 for Documents Required by the Office of Patent Publication, 1280 Off. Gaz. Patent Office 918 (March 23, 2004). In addition, if the informality is not corrected until after payment of the issue fee, for purposes of 35 U.S.C. 154(b)(1)(iv), "all outstanding requirements" will be considered to have been satisfied when the informality has been corrected. A failure to respond within the above-identified time period will result in the application being ABANDONED. **This period for reply is NOT extendable under 37 CFR 1.136(a).**

See attachment(s).

*A copy of this notice **MUST** be returned with the reply. Please address response to "Mail Stop Issue Fee, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450".*

/Patricia Pacenski/
Publication Branch
Office of Data Management
(571) 272-4200

IDENTIFICATION OF APPLICATION DEFICIENCIES

- Applicant must provide legible text for the following item(s).
 - Specification filed , page(s) .
 - Claims filed , claim(s) .
 - Oath/declaration filed .
 - Other: .
- Applicant must provide missing information on the following page(s) of the specification by amending the specification to add the missing text. No new matter may be added. line 1 of page 6 appears to be incomplete
- The specification refers to one or more applications by attorney docket number and does not show the U.S. application number(s). Applicant must supply the U.S. application number in place of each attorney docket number.
- Applicant must provide an Abstract of the Disclosure.
- Applicant has submitted a DECLARATION (37 CFR 1.63) FOR A UTILITY OR DESIGN APPLICATION USING AN APPLICATION DATA SHEET (37 CFR 1.76) (e.g., form PTO/SB/01A). The Application Data Sheet, however, is not present with the filed application. Applicant must submit an Application Data Sheet or file a new oath or declaration (e.g., PTO/SB/01) executed by the inventors and containing the information required in 37 CFR 1.63.
- Applicant must provide an executed declaration.
- Applicant must provide the missing page(s) of the oath/declaration or Application Data Sheet filed
- Applicant must provide a declaration signed by inventor(s) .
- The oath/declaration filed shows non-initialed and/or non-dated alterations. Applicant must file a new oath/declaration in compliance with 37 CFR 1.67(a).
- Applicant(s) in the latest-filed oath/declaration or Application Data Sheet (ADS) did not show the inventor's residence at all, or did not show both a city and state in the U.S. inventor's residence, or did not show both a city and country in the non-U.S. inventor's residence. Applicant must supply an oath/declaration or Application Data Sheet (ADS) that shows each U.S. inventor's city and state of residence and each non-U.S. inventor's city and country of residence.

Substituted page 6

described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

SUMMARY OF THE INVENTION

The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by injection containing the compound

10 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, more particularly to a formulation adapted for administration by injection containing the compound
7 α -[9-(4,4,5,5,5-pentafluoropentylsulphinyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle.

BRIEF DESCRIPTION OF THE DRAWING

15 Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 below, and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits.

DETAILED DESCRIPTION OF THE INVENTION

20 Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

25

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06

Electronic Patent Application Fee Transmittal

Application Number:	12285887
Filing Date:	15-Oct-2008
Title of Invention:	FORMULATION
First Named Inventor/Applicant Name:	John R. Evans
Filer:	Carlos M. Tellez
Attorney Docket Number:	11285.0056-00000

Filed as Large Entity

Utility under 35 USC 111(a) Filing Fees

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Basic Filing:				
Pages:				
Claims:				
Miscellaneous-Filing:				
Petition:				
Patent-Appeals-and-Interference:				
Post-Allowance-and-Post-Issuance:				
Utility Appl issue fee	1501	1	1740	1740
Publ. Fee- early, voluntary, or normal	1504	1	300	300

Description	Fee Code	Quantity	Amount	Sub-Total in USD(\$)
Extension-of-Time:				
Miscellaneous:				
Total in USD (\$)				2040

Electronic Acknowledgement Receipt

EFS ID:	13649794
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	FORMULATION
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Carlos M. Tellez
Filer Authorized By:	
Attorney Docket Number:	11285.0056-00000
Receipt Date:	04-SEP-2012
Filing Date:	15-OCT-2008
Time Stamp:	15:03:13
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	yes
Payment Type	Credit Card
Payment was successfully received in RAM	\$2040
RAM confirmation Number	1646
Deposit Account	
Authorized User	

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
-----------------	----------------------	-----------	----------------------------------	------------------	------------------

1	Issue Fee Payment (PTO-85B)	11285-0056-00000ISSUEFEEPAYMENT.pdf	36618 4abbd63ba121833238721bbc8c421b8fc4733ea8	no	1
Warnings:					
Information:					
2	Post Allowance Communication - Incoming	11285-0056-00000RESPONSENOTICEANDCOPYOFNOTICEOFFILECORRECTEDAPPLICATION.pdf	78402 d1f784346701d68691d6581e727f6e46e8027f7b	no	6
Warnings:					
Information:					
3	Fee Worksheet (SB06)	fee-info.pdf	31526 eee9ce3badbbbf35eb96456cca5ffd70bfdab3d	no	2
Warnings:					
Information:					
Total Files Size (in bytes):				146546	
<p>This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.</p> <p><u>New Applications Under 35 U.S.C. 111</u> If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.</p> <p><u>National Stage of an International Application under 35 U.S.C. 371</u> If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.</p> <p><u>New International Application Filed with the USPTO as a Receiving Office</u> If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.</p>					



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
12/285,887	10/15/2008	John R. Evans	11285.0056-00000	1199
22852	7590	09/06/2012	EXAMINER	
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER LLP 901 NEW YORK AVENUE, NW WASHINGTON, DC 20001-4413			HUI, SANMING R	
			ART UNIT	PAPER NUMBER
			1628	
			MAIL DATE	DELIVERY MODE
			09/06/2012	PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Response to Rule 312 Communication	Application No.	Applicant(s)
	12/285,887	JOHN R.
	Examiner	Art Unit

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

1. The amendment filed on 04 September 2012 under 37 CFR 1.312 has been considered, and has been:

- a) entered.
- b) entered as directed to matters of form not affecting the scope of the invention.
- c) disapproved because the amendment was filed after the payment of the issue fee.
Any amendment filed after the date the issue fee is paid must be accompanied by a petition under 37 CFR 1.313(c)(1) and the required fee to withdraw the application from issue.
- d) disapproved. See explanation below.
- e) entered in part. See explanation below.

B.Crittenden

Publishing Division



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	ISSUE DATE	PATENT NO.	ATTORNEY DOCKET NO.	CONFIRMATION NO.
12/285,887	12/11/2012	8329680	11285.0056-00000	1199

22852 7590 11/21/2012
FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER
LLP
901 NEW YORK AVENUE, NW
WASHINGTON, DC 20001-4413

ISSUE NOTIFICATION

The projected patent number and issue date are specified above.

Determination of Patent Term Adjustment under 35 U.S.C. 154 (b) (application filed on or after May 29, 2000)

The Patent Term Adjustment is 338 day(s). Any patent to issue from the above-identified application will include an indication of the adjustment on the front page.

If a Continued Prosecution Application (CPA) was filed in the above-identified application, the filing date that determines Patent Term Adjustment is the filing date of the most recent CPA.

Applicant will be able to obtain more detailed information by accessing the Patent Application Information Retrieval (PAIR) WEB site (<http://pair.uspto.gov>).

Any questions regarding the Patent Term Extension or Adjustment determination should be directed to the Office of Patent Legal Administration at (571)-272-7702. Questions relating to issue and publication fee payments should be directed to the Application Assistance Unit (AAU) of the Office of Data Management (ODM) at (571)-272-4200.

APPLICANT(s) (Please see PAIR WEB site <http://pair.uspto.gov> for additional applicants):

John R. Evans, Macclesfield, UNITED KINGDOM;
Rosalind U. Grundy, Macclesfield, UNITED KINGDOM;

The United States represents the largest, most dynamic marketplace in the world and is an unparalleled location for business investment, innovation, and commercialization of new technologies. The USA offers tremendous resources and advantages for those who invest and manufacture goods here. Through SelectUSA, our nation works to encourage and facilitate business investment. To learn more about why the USA is the best country in the world to develop technology, manufacture products, and grow your business, visit SelectUSA.gov.

PATENT
Customer No. 22,852
Attorney Docket No. 11285.0056-00000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
U.S. Patent No. 8,329,680) Application No.: 12/285,887
Issue Date: December 11, 2012) Filed: October 15, 2008
Inventors: John R. EVANS et al.) Confirmation No.: 1199
For: FORMULATION) **EFS-Web Filing**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Dear Commissioner:

**FEE ADDRESS FOR MAINTENANCE FEE PURPOSES
IN ACCORDANCE WITH 37 C.F.R. § 1.363**


In accordance with the provisions of 37 C.F.R. § 1.363, the fee address set forth below is being supplied for purposes of receiving notices, receipts, and other correspondence relating to the payment of maintenance fees:

Thomson Scientific IP Management Services
300 Franklin Center, 29100 Northwestern Highway
Southfield, Michigan 48034
Payor Number: 00124

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, LLP

Dated: January 17, 2013

By: 
Mark D. Sweet
Reg. No. 41,469
(202) 408-4162

Electronic Acknowledgement Receipt

EFS ID:	14725009
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	FORMULATION
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Abhay Ashok Watwe/andrea pinkney
Filer Authorized By:	Abhay Ashok Watwe
Attorney Docket Number:	11285.0056-00000
Receipt Date:	17-JAN-2013
Filing Date:	15-OCT-2008
Time Stamp:	13:39:18
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	no
------------------------	----

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1	Change of Address	feeaddress.pdf	49101 <small>936c65a9d42a66083d27210c0fbc9b453577275a</small>	no	1

Warnings:

Information:

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)
U.S. Patent No. 8,329,680) Application No.: 12/285,887
Issue Date: December 11, 2012) Filed: October 15, 2008
Inventors: John R. EVANS et al.) Confirmation No.: 1199
For: FORMULATION) **EFS-Web Filing**

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Dear Commissioner:

**FEE ADDRESS FOR MAINTENANCE FEE PURPOSES
IN ACCORDANCE WITH 37 C.F.R. § 1.363**

In accordance with the provisions of 37 C.F.R. § 1.363, the fee address set forth below is being supplied for purposes of receiving notices, receipts, and other correspondence relating to the payment of maintenance fees:

CPA Global Limited
Liberation House
Castle Street
Jersey JE1 1BL
Channel Islands
Customer Number: 000197

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, LLP

Dated: February 15, 2013

By: 

Mark D. Sweet
Reg. No. 41,469
(202) 408-4162

Electronic Acknowledgement Receipt

EFS ID:	14976524
Application Number:	12285887
International Application Number:	
Confirmation Number:	1199
Title of Invention:	FORMULATION
First Named Inventor/Applicant Name:	John R. Evans
Customer Number:	22852
Filer:	Abhay Ashok Watwe/Margie Harris
Filer Authorized By:	Abhay Ashok Watwe
Attorney Docket Number:	11285.0056-00000
Receipt Date:	15-FEB-2013
Filing Date:	15-OCT-2008
Time Stamp:	17:49:31
Application Type:	Utility under 35 USC 111(a)

Payment information:

Submitted with Payment	no
------------------------	----

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)/ Message Digest	Multi Part /.zip	Pages (if appl.)
1	Change of Address	FeeAddress.pdf	49405 <small>6cb1e2c68f5730b06c81a99b6de1beb246375150</small>	no	1

Warnings:

Information:

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

AO 120 (Rev. 08/10)

TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
-----	---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the U.S. District Court for the District of New Jersey on the following:
 ___ Trademarks or Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 3:14-cv-03547-FLW-LHG	DATE FILED 6/3/2014	U.S. DISTRICT COURT TRENTON, NJ
PLAINTIFF ASTRAZENECA PHARMACEUTICALS LP		DEFENDANT SANDOZ INC.

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 US 6,774,122 B2	August 10, 2004	AstraZeneca AB
2 US 7,456,160 B2	November 25, 2008	AstraZeneca AB
3 US 8,329,680 B2	December 11, 2012	AstraZeneca AB
4 US 8,466,139 B2	June 18, 2013	AstraZeneca AB
5		

In the above—entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY ___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above—entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ Marlene Kalbach	DATE 6/3/2014
---------------------------	---	------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

AO 120 (Rev. 08/10)

TO: Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the U.S. District Court Northern District of Illinois on the following
 Trademarks or Patents. (the patent action involves 35 U.S.C. § 292.):

DOCKET NO. 14-cv-7358	DATE FILED 9/22/2014	U.S. DISTRICT COURT Northern District of Illinois
PLAINTIFF AstraZeneca Pharmaceuticals LP, AstraZeneca UK Limited, AstraZeneca AB		DEFENDANT Sagent Pharmaceuticals, Inc.
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 6,774,122	8/10/2004	AstraZeneca AB
2 7,456,160	11/25/2008	AstraZeneca AB
3 8,329,680	12/11/2012	AstraZeneca AB
4 8,466,139	6/18/2013	AstraZeneca AB
5		

In the above—entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY <input type="checkbox"/> Amendment <input type="checkbox"/> Answer <input type="checkbox"/> Cross Bill <input type="checkbox"/> Other Pleading		
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK	
1			
2			
3			
4			
5			

In the above—entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK Thomas G. Bruton	(BY) DEPUTY CLERK Melissa Rivera	DATE 9/23/14
---------------------------	-------------------------------------	-----------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

<i>AO 120 (Rev. 08/10)</i>		
TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the **U.S. District Court for the District of New Jersey** on the following:
 ___ Trademarks or **X** Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 1:15-cv-00615-RMB-KMW	DATE FILED 1/29/2015	U.S. DISTRICT COURT CAMDEN, NJ
PLAINTIFF ASTRAZENECA PHARMACEUTICALS LP		DEFENDANT GLENMARK PHARMACEUTICALS LTD.

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 6,774,122	8/10/2004	AstraZeneca AB
2 7,456,160	11/25/2008	AstraZeneca AB
3 8,329,680	12/11/2012	AstraZeneca AB
4 8,466,139	6/18/2013	AstraZeneca AB
5		

<u>In the above—entitled case, the following patent(s)/ trademark(s) have been included:</u>		
DATE INCLUDED	INCLUDED BY ___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

<u>In the above—entitled case, the following decision has been rendered or judgement issued:</u>	
DECISION/JUDGEMENT	

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ Nicholas Zotti	DATE 1/29/2015
---------------------------	--	-------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

AO 120 (Rev. 08/10)		
TO:	<p style="text-align: center;">Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450</p>	<p style="text-align: center;">REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK</p>

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the U.S. District Court for the District of New Jersey on the following:
 ___ Trademarks or Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 1:15-cv-07009-RMB-KMW	DATE FILED 9/21/2015	U.S. DISTRICT COURT CAMDEN, NJ
PLAINTIFF ASTRAZENECA PHARMACEUTICALS LP		DEFENDANT MYLAN PHARMACEUTICALS INC.

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 US 6,774,122 B2	Aug. 10, 2004	AstraZeneca AB,
2 US 7,456,160 B2	Nov. 25, 2008	AstraZeneca AB,
3 US 8,329,680 B2	Dec. 11, 2012	AstraZeneca AB
4 US 8,466,139 B2	Jun. 18, 2013	AstraZeneca AB
5		

In the above--entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY	
	___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above--entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ JAIME KASSELMAN	DATE 9/21/2015
---------------------------	---	-------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

AO 120 (Rev. 08/10)

TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
-----	---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the U.S. District Court for the District of New Jersey on the following:
 ___ Trademarks or Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 1:15-cv-06990-NLH-AMD	DATE FILED 9/22/2015	U.S. DISTRICT COURT CAMDEN, NJ
PLAINTIFF HORIZON PHARMA IRELAND LIMITED		DEFENDANT AMNEAL PHARMACEUTICALS LLC

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 US 9,132,110 B2	9/15/2015	HZNP LIMITED
2		
3		
4		
5		

In the above--entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY	
	___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above--entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ Brian D. Kemner	DATE 9/22/2015
---------------------------	---	-------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

AO 120 (Rev. 08/10)

TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
-----	---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the **U.S. District Court for the District of New Jersey** on the following:
 ___ Trademarks or **X** Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 1:15-cv-06039-RMB-KMW	DATE FILED 8/7/2015	U.S. DISTRICT COURT CAMDEN, NJ
PLAINTIFF ASTRAZENECA PHARMACEUTICALS LP		DEFENDANT AGILA SPECIALTIES, INC

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 6,774,122 B2	08/10/2004	ASTRAZENECA AB
2 7,456,160 B2	11/25/2008	ASTRAZENECA AB
3 8,329,680 B2	12/11/2012	ASTRAZENECA AB
4 8,466,139 B2	06/18/2013	AstraZeneca AB
5		

In the above--entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY	
	___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above--entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ JAIME KASSELMAN	DATE 8/7/2015
---------------------------	---	------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
-----	---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the **U.S. District Court for the District of New Jersey** on the following:
 ___ Trademarks or Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 3:15-cv-06075-PGS-DEA	DATE FILED 8/6/2015	U.S. DISTRICT COURT TRENTON, NJ
PLAINTIFF MERCK SHARP & DOHME CORP.		DEFENDANT ACTAVIS LABORATORIES FL, INC.

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 5,661,151	8/26/1997	Schering Corporation
2		
3		
4		
5		

In the above--entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY	
	___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above--entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ Karen McGonigle	DATE 8/6/2015
---------------------------	---	------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

TO: <p style="text-align: center;">Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450</p>	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
--	--

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the U.S. District Court Northern District of Texas, Dallas Division on the following

Trademarks or Patents. (the patent action involves 35 U.S.C. § 292.):

DOCKET NO. 3:15-cv-2607-M	DATE FILED 8/7/2015	U.S. DISTRICT COURT Northern District of Texas, Dallas Division
PLAINTIFF Pathway Senior Living LLC		DEFENDANT Pathways Senior Living LLC
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 3,432,946	5/20/2008	Pathway Senior Living LLC
2		
3		
4		
5		

In the above—entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY <input type="checkbox"/> Amendment <input type="checkbox"/> Answer <input type="checkbox"/> Cross Bill <input type="checkbox"/> Other Pleading		
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK	
1			
2			
3			
4			
5			

In the above—entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK Karen Mitchell	(BY) DEPUTY CLERK s/A. Lowe	DATE 8/10/2015
-------------------------	--------------------------------	-------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy

AO 120 (Rev. 08/10)

TO:	Mail Stop 8 Director of the U.S. Patent and Trademark Office P.O. Box 1450 Alexandria, VA 22313-1450	REPORT ON THE FILING OR DETERMINATION OF AN ACTION REGARDING A PATENT OR TRADEMARK
-----	---	---

In Compliance with 35 U.S.C. § 290 and/or 15 U.S.C. § 1116 you are hereby advised that a court action has been filed in the **U.S. District Court for the District of New Jersey** on the following:
 ___ Trademarks or **X** Patents. (___ the patent action involves 35 U.S.C. § 292.)

DOCKET NO. 1:16-cv-01962-RMB-KMW	DATE FILED 4/7/2016	U.S. DISTRICT COURT CAMDEN, NJ
PLAINTIFF ASTRAZENECA PHARMACEUTICALS LP		DEFENDANT INNOPHARMA LICENSING LLC

PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1 US 6,774,122 B2	Aug. 10, 2004	AstraZeneca AB,
2 US 7,456,160 B2	Nov. 25, 2008	AstraZeneca AB,
3 US 8,329,680 B2	Dec. 11, 2012	AstraZeneca AB,
4 US 8,466,139 B2	Jun. 18, 2013	AstraZeneca AB
5		

In the above—entitled case, the following patent(s)/ trademark(s) have been included:

DATE INCLUDED	INCLUDED BY ___ Amendment ___ Answer ___ Cross Bill ___ Other Pleading	
PATENT OR TRADEMARK NO.	DATE OF PATENT OR TRADEMARK	HOLDER OF PATENT OR TRADEMARK
1		
2		
3		
4		
5		

In the above—entitled case, the following decision has been rendered or judgement issued:

DECISION/JUDGEMENT

CLERK William T. Walsh	(BY) DEPUTY CLERK s/ Ryan Merrigan	DATE 4/7/2016
---------------------------	---------------------------------------	------------------

Copy 1—Upon initiation of action, mail this copy to Director Copy 3—Upon termination of action, mail this copy to Director
 Copy 2—Upon filing document adding patent(s), mail this copy to Director Copy 4—Case file copy