Home → Treatment & Side Effects → Hormonal Therapy → Selective Estrogen Receptor Modulators (SERMs)

Treatment & Side Effects

Tamoxifen in Pill Form (Brand Name: Nolvadex)

Tamoxifen in Liquid Form (Brand Name: Soltamox)

Fvista

Fareston





Scroll for Important Safety Information and Indication

IBRANCE may cause serious side effects. including: Low white blood cell counts Low white plood cen code (neutropenia). Low white blood cell counts are very common when taking IBRANCE and may cause serious infections that can lead to death. Your healthcare provider should

Full Prescribing and Patient Product Information



Did you know there's another way to get your Neulasta®?



LEARN MORE >

Selective Estrogen Receptor Modulators (SERMs)

Selective estrogen receptor modulators, called SERMs for short, block the

effects of estrogen in the breast tissue. SERMs work by sitting in the estrogen

for estrogen and it can't attach to the cell. If estrogen isn't attached to a breast

Cells in other tissues in the body, such as bones and the uterus, also have

estrogen receptors. But each estrogen receptor has a slightly different structure, depending on the kind of cell it is in. So breast cell estrogen receptors are

different from bone cell estrogen receptors and both of those estrogen receptors

are different from uterine estrogen receptors. As their name says, SERMs are "selective" - this means that a SERM that blocks estrogen's action in breast cells can activate estrogen's action in other cells, such as bone, liver, and uterine

cell, the cell doesn't receive estrogen's signals to grow and multiply.

receptors in breast cells. If a SERM is in the estrogen receptor, there is no room



Sign in to receive recommendations (<u>Learn more</u>)











LEARN MORE >

Email Updates

Stay informed about current research, online events, and more

Email Address

Sign Up

Read our privacy policy

There are three SERMs:

cells

- . tamoxifen in pill form (also called tamoxifen citrate; brand name: Nolvadex); tamoxifen in liquid form (brand name: Soltamox)
- · Evista (chemical name: raloxifene)
- <u>Fareston</u> (chemical name: toremifene)

Each is a pill, usually taken once a day, except for tamoxifen in liquid form. Tamoxifen is the oldest, most well-known, and most-prescribed SERM.

SERMs can be used to treat women both before and after menopause.

Benefits of SERMs

Because tamoxifen is the most commonly used SERM, most of the studies comparing SERMs to aromatase inhibitors have looked at tamoxifen versus aromatase inhibitors. Several studies have compared tamoxifen with aromatase inhibitors to see which type of medicine was more effective in treating earlystage, hormone-receptor-positive breast cancer in postmenopausal women. Based on the results, most doctors go by the following recommendations:

- . An aromatase inhibitor is the best type of hormonal therapy to start with for postmenopausal women. When treating early-stage, hormone-receptorpositive breast cancer, aromatase inhibitors have more benefits and fewer serious side effects than tamoxifen.
- · Switching to an aromatase inhibitor after taking tamoxifen for 2 to 3 years (for a total of 5 years of hormonal therapy) offers more benefits than 5 years of tamoxifen. Taking an aromatase inhibitor for 5 years after taking tamoxifen for 5 years continues to reduce the risk of the cancer coming back, compared to no treatment after tamoxifen.
- . Taking an aromatase inhibitor for 5 years after taking tamoxifen for 5 years







endometrial cancer. If you and your doctor are considering tamoxifen or another SERM as part of your treatment plan, tell your doctor if you smoke or have a history of blood clots or heart attack. If you're taking a SERM, call your doctor immediately if you have any of these symptoms:

- · abnormal vaginal bleeding or discharge
- · pain or pressure in the pelvis
- · leg swelling or tenderness
- · chest pain
- · shortness of breath
- · weakness, tingling, or numbness in your face, arm, or leg
- sudden difficulty seeing
- dizziness
- sudden severe headache

The most common side effects of SERMs are:

- fatigue
- hot flashes
- night sweats
- vaginal discharge
- · mood swings

You should not take a SERM if you are breastfeeding, pregnant, trying to get pregnant, or if there is any chance that you could be pregnant. These medicines may cause damage to developing embryos. You should use an effective non-hormonal type of birth control -- such as condoms, a diaphragm along with spermicide, or a non-hormonal I.U.D. —while you are taking a SERM. Ask your doctor which type of non-hormonal birth control would be best for you, as well as how long you should use this type of birth control after you stop taking a SERM.

As a benefit, SERMs also can improve bone density, which reduces the risk of osteoporosis.

Last modified on November 5, 2015 at 11:50 AM Leer esta página en español

More From Our Site



Estrogen Receptor Downregulators (ERDs)



Hormonal Therapy Side Effects Comparison Chart

Recommended by Outbrain



Symptoms & Diagnosis

Back to Top

Breastcancer.org is a non-profit organization dedicated to providing information and community to those touched by this disease. Learn more about our commitment to providing complete, accurate, and private breast cancer information.

Breastcancer.org 120 East Lancaster Avenue, Suite 201 Ardmore, PA 19003 © 2016 Breastcancer.org - All rights reserved.

Stay informed about current research, online events, and more.

Email address

Sign up

earn more about our commitment to your privacy



http://www.breastcancer.org/treatment/hormonal/serms Fri Jun 10 2016 10:53:38 GMT-0700 (Pacific Daylight Time)

