

A *v-erbB*-related protooncogene, *c-erbB-2*, is distinct from the *c-erbB-1*/epidermal growth factor-receptor gene and is amplified in a human salivary gland adenocarcinoma

(*src* gene family/kinase family/gene amplification/cancer)

KENTARO SEMBA*, NOBUYUKI KAMATA*†, KUMAO TOYOSHIMA*, AND TADASHI YAMAMOTO*

*The Institute of Medical Science, The University of Tokyo 4-6-1, Shirokanedai, Minato-ku, Tokyo 108, Japan; and †Tokyo Medical and Dental University, Faculty of Dentistry 1-5-45, Yushima, Bunkyo-ku, Tokyo 113, Japan

Communicated by Takashi Sugimura, June 10, 1985

ABSTRACT From a human genomic library, we obtained six *v-erbB*-related DNA clones. A DNA probe prepared from one of the clones, λ 107, hybridized to *Eco*RI fragments of 6.4 and 13 kilobase pairs of human DNA. Neither of these fragments was amplified in A431 vulva carcinoma cells, in which the gene encoding the epidermal growth factor receptor is amplified. In addition, the probe from λ 107 hybridized with a single, 4.8-kilobase poly(A)⁺ RNA species and did not react with EGF receptor mRNA. Thus, we conclude that clone λ 107 represents a *v-erbB*-related gene (*c-erbB-2*) that is distinct from the EGF receptor gene. In contrast, the other five clones were shown to represent the EGF receptor gene (*c-erbB-1*). Partial nucleotide sequence analysis of the λ 107 insert showed that this clone contained at least seven putative exons and that six of them could encode the kinase domain characteristic of protein products of the *src* oncogene family. Southern blot analysis showed close similarity of the restriction patterns of the rat *c-erbB-2* gene and the rat *neu* oncogene, suggesting possible involvement of *c-erbB-2* in human cancer. In fact, \approx 30-fold amplification of *c-erbB-2* was observed in a human adenocarcinoma of the salivary gland.

At least 19 genes have been identified as retroviral oncogenes that are responsible for inducing tumors *in vivo* and transforming cells *in vitro* (1). Ten of them apparently encode transforming proteins that share a kinase domain homologous to that of pp60^{src}, a tyrosine-specific protein kinase. The cellular cognate, encoded by the *c-src* gene, also exhibits tyrosine-specific kinase activity. Of particular interest is the fact that tyrosine-specific kinases are also encoded by the genes for several receptors for polypeptide growth factors, including the receptors for epidermal growth factor (EGF) (2), platelet-derived growth factor (PDGF) (3), insulin (4), and insulin-like growth factor I (5). This implies a possible link between the action of the growth factor-receptor complex and the oncogene product with tyrosine-specific kinase activity. In fact, recent analysis of the *v-erbB* gene and the EGF receptor gene indicated that the *v-erbB* gene is a part of the EGF receptor gene and codes for the internal domain and transmembrane portion of the receptor (6-8). These findings, together with the extensive identity of the amino acid sequences of the *v-sis* protein and platelet-derived growth factor (9, 10), suggest that some viral oncogene products mimic the action of the polypeptide growth factor-receptor complex in activating a cellular pathway involved in cell proliferation.

We examined details of the relation between the *v-erbB* gene and the EGF receptor gene and the possible involvement of this gene in human cancer. During this study we identified two *v-erbB*-related genes, *c-erbB-1* and *c-erbB-2*, in the human genome.

MATERIALS AND METHODS

Cells and Tissues. A431 vulva carcinoma cells and human embryo fibroblasts were maintained in Dulbecco's modified Eagle's medium with 10% fetal calf serum. K562 chronic myelogenous leukemia cells and MT2 adult T-cell leukemia cells were maintained in RPMI 1640 medium with 10% fetal calf serum. Mouse FM3A cells were from M. C. Yoshida (Hokkaido University) and primary tumors were provided by K. Rikimaru (Tokyo Medical and Dental University).

Isolation of Clones. A human genomic library was constructed from placental DNA as described (11). The library was screened for the *v-erbB*-related sequence by plaque-hybridization as described (12) in 30% (vol/vol) formamide/4 \times NaCl/Cit (1 \times is 0.15 M NaCl/15 mM sodium citrate)/50 mM Hepes, pH 7.0/denatured salmon sperm DNA (20 μ g/ml)/10 \times Denhardt's solution (1 \times is 0.02% polyvinylpyrrolidone/0.02% Ficoll/0.02% bovine serum albumin) at 42°C for 16 hr. After hybridization, the filters were washed with 2 \times NaCl/Cit/0.1% NaDodSO₄ at room temperature and then with 0.6 \times NaCl/Cit/0.1% NaDodSO₄ at 50°C. Plaques were purified by successive plaque-hybridization. The probe used was a 1.7-kilobase-pair (kbp) *Sst* I-*Stu* I DNA fragment that represents the *v-erbB* gene of avian erythroblastosis virus strain H (6); it was labeled with [α -³²P]dCTP by nick-translation (13) to a specific activity of 2 \times 10⁸ cpm/ μ g of DNA. Phage DNAs were prepared as described (14).

Nucleotide Sequence Analysis. The nucleotide sequence was determined by the procedure of Maxam and Gilbert (15) and the dideoxy chain-termination method (16, 17) in conjunction with bacteriophage M13 mp19 (18).

Blot-Hybridization Analysis of DNA and RNA. High molecular weight DNAs were prepared from chicken blood, human embryo fibroblasts, mouse FM3A cells, rat spleen, and primary tumors. The DNAs (10 μ g per lane) were digested with restriction endonucleases under the conditions recommended by suppliers (Takara Shuzo) and fractionated by electrophoresis in 1% agarose gels. The fragments were subjected to Southern blot hybridization (19) at 42°C for 16 hr in 4 \times NaCl/Cit/50 mM Hepes, pH 7.0/10 \times Denhardt's solution/denatured salmon sperm DNA (20 μ g/ml)/30% (relaxed conditions) or 50% (stringent conditions) formamide. After hybridization, the filters were washed with 2 \times




FIG. 1. Identification and cloning of the *v-erbB*-related sequence that is distinct from the EGF receptor gene. (a) Hybridization with the *v-erbB* probe. DNAs prepared from A431 cells (lanes 1, 3, and 5) and placenta (lanes 2, 4, and 6) were digested with *EcoRI* (lanes 1 and 2), *Sac I* (lanes 3 and 4), or *Pvu II* (lanes 5 and 6). Because the placental DNA was contaminated with RNA, samples in lanes 2, 4, and 6 contained less DNA than those in lanes 1, 3, and 5. The digests were subjected to Southern blot hybridization using a *v-erbB* probe under the relaxed conditions described in *Materials and Methods*. Autoradiographs obtained by longer exposures of lanes 2, 4, and 6 are shown to the right of the respective lanes. Arrowheads indicate positions of fragments of *HindIII*-digested λ DNA used as standards. Stars indicate bands that were not amplified in the A431 samples. (b) Restriction map of the λ 107 insert. E, K, S, and X represent restriction sites of *EcoRI*, *Kpn I*, *Sac I*, and *Xba I*, respectively. The thick line indicates the *Kpn I*-*Xba I* fragment (KX fragment) used as a specific probe of the *c-erbB-2* gene. (c) Hybridization with the KX fragment. The same filter represented in a was washed and then hybridized with the KX probe under the stringent conditions described in *Materials and Methods*.

NaCl/Cit/0.1% NaDodSO₄ and then with either 0.2 \times NaCl/Cit/0.1% NaDodSO₄ (stringent) or 0.6 \times NaCl/Cit/0.1% NaDodSO₄ (relaxed). RNAs were prepared by the guanidinium isothiocyanate/cesium chloride method (20). Poly(A)⁺ RNA selected by oligo(dT)-cellulose (P-L Biochemicals type 7) column chromatography was denatured with 50% formamide/2.2 M formaldehyde and 5 μ g of each RNA sample was subjected to electrophoresis in 1% agarose gel containing 2.2 M formaldehyde (21). RNAs on the gel were transferred directly to a nitrocellulose filter and subjected to blot hybridization under stringent conditions, as described (22). The DNA probes used for blot hybridization were *v-erbB* DNA (described above), an EGF receptor cDNA clone (pE7) (22), and *c-erbB-2*-specific DNA (described below).

RESULTS

Isolation of a Human *v-erbB*-Related Sequence Distinct from the EGF Receptor Gene. High molecular weight DNAs prepared from A431 cells and human placenta were digested with either *EcoRI*, *Sac I*, or *Pvu II*, and the digests were analyzed by Southern blot hybridization under relaxed conditions with the *v-erbB* probe, which covers almost 90% of the *v-erbB* gene. Most of the *v-erbB*-related sequences were amplified in A431 cells at least 20-fold compared with those in placenta (Fig. 1a). These sequences were observed as 8.1-, 5.9-, 5.4-, and 3.5-kbp *EcoRI* fragments; 20-, 4.7-, 3.3-, 1.7-, 0.9-, and 0.5-kbp *Sac I* fragments; and 6.6-, 3.3-, 2.9-, 2.3-, 1.9-, and 0.9-kbp *Pvu II* fragments. The amplified sequences were of the EGF receptor gene as reported (8, 23). However, we observed several restriction fragments that hybridized to the *v-erbB* probe but that were not amplified in A431 cells (13- and 6.4-kbp *EcoRI* fragment, a 6.9-kbp *Sac I* fragment, and

another *v-erbB*-related gene, from which the unamplified restriction fragments are generated.

We searched a human genomic library for *v-erbB*-related sequences, using the *v-erbB* probe, under the relaxed hybridization and washing conditions, and isolated six independent clones. Restriction map analysis showed that all the cloned inserts except the λ 107 insert represented one gene, *c-erbB-1*. In these clones, we identified four *EcoRI* fragments as exon-containing sequences by the criterion that they hybridized to the *v-erbB* probe. All these fragments were




FIG. 2. Expression of the EGF receptor gene and the *c-erbB-2* gene in human cells. Duplicate samples of poly(A)⁺ RNA were subjected to blot hybridization with the ³²P-labeled KX fragment (a) or with the ³²P-labeled EGF receptor cDNA clone pE7 (22) (b). Sizes of RNA species that hybridize with the probes are given. Chicken ribosomal RNAs (28S, about 4.8 kb, and 18S, about 2.0 kb) and Rous sarcoma virus RNA (39S, about 10.0 kb) served as size standards. RNAs were isolated from human placenta (lanes 1), A431 cells (lanes




FIG. 3. Nucleotide sequence of the exons for the kinase domain of *c-erbB-2*. (a) Putative exons in the $\lambda 107$ clone are indicated by thick vertical bars. The two *EcoRI* (E) sites are shown by vertical lines. (b) Nucleotide sequence of the putative exons. Restriction fragments that hybridized to *v-erbB* DNA were sequenced. The exons were defined by the splicing consensus sequence and by the high homology of the deduced amino acid sequences to those of the EGF receptor gene *EGFR*. The nucleotide sequence of putative exons of *c-erbB-2* (in capital letters) is compared with that of the EGF receptor gene (8). Only nonidentical nucleotides are shown for the EGF receptor gene. Intron sequences at the splicing sites that flank the exons are shown in lowercase letters. The splicing donor site of the last exon was not identified. The predicted amino acid sequence is shown above the nucleotide sequence.

amplified in A431 cells (data not shown), suggesting that the *c-erbB-1* gene is the same as the EGF receptor gene.

A restriction map of the $\lambda 107$ DNA insert is shown in Fig. 1b. The 440-bp *Kpn I-Xba I* fragment (KX fragment) was used as a probe in the following hybridization experiments. This fragment did not contain human repetitive sequences and hybridizes with the *v-erbB* probe under the relaxed conditions. The *v-erbB* probe on the filter represented in Fig. 1a was washed off and the filter was rehybridized with the ^{32}P -labeled KX fragment. Fig. 1c shows that the KX probe reacted with the restriction fragments (13- and 6.4-kbp *EcoRI* fragments, 6.9-kbp *Sac I* fragment, and 5.1-kbp *Pvu II* fragment) that were not amplified in A431 cells, suggesting that the $\lambda 107$ insert does not represent the EGF receptor gene. Therefore, we tentatively concluded that the $\lambda 107$ insert represented a *v-erbB*-related gene (*c-erbB-2*) that differs from the EGF receptor gene.

Then we examined the expression of the *v-erbB*-related genes. RNAs were prepared from human cell lines (K562, MT2, and A431 cells), human embryo fibroblasts, and human placenta. Hybridization was carried out under stringent

conditions, most transcripts detected with the EGF receptor cDNA probe were of 10 kilobases (kb) and 5.6 kb and were apparently overproduced in A431 cells (Fig. 2b and refs. 8 and 23). In addition, variant 2.9-kb mRNA was detected in A431 cells, as reported previously (8, 23). On the other hand, the mRNA that hybridized with the KX probe was a single species of about 4.8 kb (Fig. 2a). The level of expression of this mRNA was virtually the same in A431 cells as in placenta and was slightly higher in human embryo fibroblasts than in other cells. Interestingly, transcription of the EGF receptor gene and *c-erbB-2* genes was not detected in the leukemic cell lines K562 and MT2. The above results strongly indicate that the human genome contains two *v-erbB*-related genes, the *c-erbB-1*/EGF receptor gene and the *c-erbB-2* gene.

***c-erbB-2* Encodes a Protein with a Kinase Domain.** We determined the nucleotide sequence of the $\lambda 107$ insert that hybridized with the *v-erbB* DNA and identified seven putative exons flanked by a consensus sequence of splicing junctions (data not shown). Fig. 3 shows that the nucleotide sequences of all seven exons (total, 885 bp) were highly homologous with those of the EGF receptor gene.




FIG. 5. Conservation of the *c-erbB-2* gene in vertebrates. (a) Conserved *c-erbB-2* sequences in chicken and mouse. Nitrocellulose filters containing *EcoRI* digests of DNA from human placenta (lane 1), mouse FM3A cells (lane 2), and chicken blood (lane 3) were probed with the KX DNA. (b) Close similarity of the *c-erbB-2* gene to the *neu* oncogene. Rat DNA was cleaved with either *EcoRI* (lane 1) or *EcoRI* plus *BamHI* (lane 2). The digests were analyzed by Southern blot hybridization with the KX probe. Arrowheads indicate positions of fragments from *HindIII* digestion of bacteriophage λ DNA (see Fig. 1a).

These findings lead us to speculate that there is a close relation between *c-erbB-2* and *neu*.

Amplification of protooncogenes such as *myc* and *ras* has been reported to be associated with various human cancers (26–30). This amplification suggests that the protein products of the amplified protooncogenes are involved in some stage of tumorigenesis. Of particular interest is the role(s) of growth factor receptors in tumorigenesis, which is inferred from the observations that EGF receptors are expressed in large numbers on the surface of cells of epidermoid carcinomas (31–33). Since the *c-erbB-2* gene could code for a protein similar to the EGF receptor or at least its intracellular domain and was amplified in an adenocarcinoma of the salivary gland, the gene products expressed abnormally in this tumor




FIG. 6. Amplification of the *c-erbB-2* gene in human primary tumors. (a) Nitrocellulose filters containing *EcoRI* digests of tumor DNAs were probed with *c-erbB-2*-specific KX DNA. (b) After removal of the first probe, the filters were then hybridized with the EGF receptor cDNA probe. Hybridization was carried out under stringent conditions. DNAs were isolated from a neuroblastoma (lane 1), an adenocarcinoma of the salivary gland (lane 2), two squamous cell carcinomas (lanes 3 and 4), placenta (lane 5), and the A431 cell line (lane 6). Markers at left in a and b indicate positions

might function in establishing and maintaining the cancerous state.

Note Added in Proof. King et al. (34) have independently isolated an *erbB-2* gene amplified in a human mammary carcinoma.

We thank Y. Miura and H. Kawano for excellent technical assistance, M. Nishizawa and N. Miyajima for helpful discussions, and S. Sasaki for typing this manuscript. We also wish to thank N. Nomura for DNA from a human primary neuroblastoma and I. Pastan for recombinant plasmid pE7. This work was supported in part by a Grant-in-Aid for Special Project Research Cancer-Bioscience from the Ministry of Education, Science, and Culture of Japan.

- Land, H., Parada, L. F. & Weinberg, R. A. (1983) *Science* **222**, 771–778.
- Cohen, S., Carpenter, G. & King, L. (1980) *J. Biol. Chem.* **255**, 4834–4842.
- Nishimura, J., Hung, H. S. & Deuel, T. F. (1982) *Proc. Natl. Acad. Sci. USA* **79**, 4303–4307.
- Kasuga, M., Zick, Y., Blithe, D. L., Crettaz, M. & Kahn, C. R. (1982) *Nature (London)* **298**, 667–669.
- Rubin, J. B., Shia, M. A. & Pilch, P. F. (1983) *Nature (London)* **305**, 438–440.
- Yamamoto, T., Nishida, T., Miyajima, N., Kawai, S., Ooi, T. & Toyoshima, K. (1983) *Cell* **35**, 71–78.
- Downward, J., Yarden, Y., Mayes, E., Scrace, G., Totty, N., Stockwell, P., Ulrich, A., Schlessinger, J. & Waterfield, M. D. (1984) *Nature (London)* **307**, 521–527.
- Ulrich, A., Coussens, L., Hayflick, J. S., Dull, T. J., Gray, A., Tam, A. W., Lee, J., Yarden, Y., Libermann, T. A., Schlessinger, J., Downward, J., Mayes, E. L. V., Whittle, N., Waterfield, M. D. & Seeburg, P. H. (1984) *Nature (London)* **309**, 418–425.
- Waterfield, M. D., Scrace, G. T., Whittle, N., Stroobant, P., Johnson, A., Wasteson, A., Westermark, B., Heldin, C.-H., Huang, J. S. & Deuel, T. F. (1983) *Nature (London)* **304**, 35–39.
- Doolittle, R. F., Hunkapiller, M. W., Hood, L. E., Devare, S. G., Robbins, K. C., Aaronson, S. A. & Antoniades, H. N. (1983) *Science* **221**, 275–277.
- Maniatis, T., Hardison, R. C., Lacy, E., Lauer, J., O'Connell, C., Quon, D., Sim, G. K. & Efstratiadis, A. (1978) *Cell* **15**, 687–701.
- Benton, W. D. & Davis, R. W. (1977) *Science* **196**, 180–182.
- Maniatis, T., Jeffrey, A. & Kleid, D. G. (1975) *Proc. Natl. Acad. Sci. USA* **72**, 1184–1188.
- Davis, R. W., Botstein, D. & Roth, J. R. (1982) *Advanced Bacterial Genetics* (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), pp. 109–111.
- Maxam, A. M. & Gilbert, W. (1980) *Methods Enzymol.* **65**, 499–560.
- Sanger, F., Nicklen, S. & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463–5467.
- Messing, J., Crea, R. & Seeburg, P. H. (1981) *Nucleic Acids Res.* **9**, 309–321.
- Messing, J. & Vieira, J. (1982) *Gene* **19**, 269–276.
- Southern, E. M. (1975) *J. Mol. Biol.* **98**, 503–517.
- Chirgwin, J. M., Pizybyla, A. E., MacDonald, R. J. & Ruffer, W. J. (1979) *Biochemistry* **18**, 5294–5299.
- Lehrach, H., Diamond, K., Wozney, J. M. & Boedtker, H. (1977) *Biochemistry* **16**, 4743–4751.
- Xu, Y.-H., Ishii, S., Clark, A. J. L., Sullivan, M., Wilson, R. K., Ma, D. P., Roe, B. A., Melino, G. T. & Pastan, I. (1984) *Nature (London)* **309**, 806–810.
- Merino, G. T., Xu, Y.-H., Ishii, S., Clark, A. J. L., Semba, K., Toyoshima, K., Yamamoto, T. & Pastan, I. (1984) *Science* **224**, 417–419.
- Schechter, A. L., Stern, D. F., Vaidyanathan, L., Decker, S. J., Drebin, J. A., Greene, M. I. & Weinberg, R. A. (1984) *Nature (London)* **312**, 513–516.
- Derbin, J. A., Stern, D. F., Link, V. C., Weinberg, R. A. & Green, M. I. (1984) *Nature (London)* **312**, 545–548.
- Collins, S. J. & Groudine, M. (1982) *Nature (London)* **298**, 679–681.
- Dalla-Favera, R., Wong-Staal, F. & Gallo, R. C. (1982) *Nature (London)* **299**, 61–63.
- Alitalo, K., Schwab, M., Lin, C. C., Varmus, H. E. & Bishop, J. M. (1983) *Proc. Natl. Acad. Sci. USA* **80**, 1707–1711.
- McCoy, M. S., Toole, J. J., Cunningham, J. M., Chang, E. H., Lowy, D. R. & Weinberg, R. A. (1983) *Nature (London)* **302**, 79–81.
- Schwab, M., Alitalo, K., Varmus, H. E., Bishop, J. M. & George, D. (1983) *Nature (London)* **303**, 497–501.
- Carpenter, G., King, L. E. & Cohen, S. (1979) *J. Biol. Chem.* **254**, 4884–4891.
- Fabricant, R., Delarco, J. E. & Todaro, G. J. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 565–569.
- Cowly, J., Gutason, B., Smith, J., Hendler, F. & Ozanne, B. (1984) in *Cancer Cells*, eds. Levine, A. J., Vande Woude, G. F., Topp, W. C. & Watson, J. D. (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), Vol. 1, pp. 5–10.