



17 May 2002



Science

Vol. 296 · No. 5571
Pages 1189–1352 \$9

DO NOT REMOVE FROM
CURRENT PERIODICALS
ROOM



RNA Silencing and Noncoding RNA



#BXNCCNG *****AUTO**5-DIGIT 53706
#041014213# AS 12/06/02 N 5571 92
UNIV OF WISCONSIN
MEMORIAL LIB SOCS CTS 7275
728 STATE ST 2
MADISON WI 53706-1418

**DOCKET
ALARM**

Find authenticated court documents without watermarks at docketalarm.com.

MTY1056

RESEARCH

BREVIA

- 1275 **Rebirth of Novae as Distance Indicators Due to Efficient, Large Telescopes** M. Della Valle and R. Gilmozzi

RESEARCH ARTICLES

- 1276 **Premature Aging in Mice Deficient in DNA Repair and Transcription** J. de Boer, J. O. Andressoo, J. de Wit, J. Huijman, R. B. Beems, H. van Steeg, G. Weeda, G. T. J. van der Horst, W. van Leeuwen, A. P. N. Themmen, M. Meradji, J. H. J. Hoeijmakers

- 1280 **Structural Basis of Transcription Initiation: RNA Polymerase Holoenzyme at 4 Å Resolution** K. S. Murakami, S. Masuda, S. A. Darst

- 1285 **Structural Basis of Transcription Initiation: An RNA Polymerase Holoenzyme-DNA Complex** K. S. Murakami, S. Masuda, E. A. Campbell, O. Muzzini, S. A. Darst

REPORTS

- 1290 **Formation of a Matter-Wave Bright Soliton** L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, C. Salomon

- 1293 **Electrochemistry and Electrogenated Chemiluminescence from Silicon Nanocrystal Quantum Dots** Z. Ding, B. M. Quinn, S. K. Haram, L. E. Pell, B. A. Korgel, A. J. Bard

- 1297 **Global Azimuthal Anisotropy in the Transition Zone** J. Trampert and H. Jan van Heijst

- 1300 **Seismic Evidence for Olivine Phase Changes at the 410- and 660-Kilometer Discontinuities** S. Lebedev, S. Chevrot, R. D. van der Hilst

- 1302 **Identity and Search in Social Networks** D. J. Watts, P. S. Dodds, M. E. J. Newman

- 1305 **Ascent of Dinosaurs Linked to an Iridium Anomaly at the Triassic-Jurassic Boundary** P. E. Olsen, D. V. Kent, H.-D. Sues, C. Koeberl, H. Huber, A. Montanari, E. C. Rainforth, S. J. Fowell, M. J. Szajna, B. W. Hartline

- 1308 **C-Cadherin Ectodomain Structure and Implications for Cell Adhesion Mechanisms** T. J. Boggon, J. Murray, S. Chappuis-Flament, E. Wong, B. M. Gumbiner, L. Shapiro

RNA SILENCING AND NONCODING RNA

- 1259 **The Other RNA World**

VIEWPOINTS


- 1260 **An Expanding Universe of Noncoding RNAs** G. Storz
- 1263 **RNA Silencing: The Genome's Immune System** R. H. A. Plasterk
- 1265 **Ancient Pathways Programmed by Small RNAs** P. D. Zamore

REVIEW

- 1270 **RNA-Dependent RNA Polymerases, Viruses, and RNA Silencing** P. Ahlquist

See also *Science's STKE* on p. 1195 and *Report* on p. 1319

COVER
1259



RNA silencing of green fluorescent protein (GFP) (center) in leaves from *Nicotiana benthamiana* is suppressed by an animal (left; B2 protein of flock house virus) or a plant (right) viral suppressor, leading to enhanced GFP expression (lighter green/yellow areas). The role of RNA silencing in defending both plant and animal genomes from invading foreign nucleic acids, the mechanisms underlying RNA silencing, and noncoding RNAs are considered in this special section. [Image: Shou-Wei Ding]

- 1313 **Vitamin D Receptor As an Intestinal Bile Acid Sensor** M. Makishima, T. T. Lu, W. Xie, G. K. Whitfield, H. Domoto, R. M. Evans, M. R. Haussler, D. J. Mangelsdorf

- 1316 **Heterotopic Shift of Epithelial-Mesenchymal Interactions in Vertebrate Jaw Evolution** Y. Shigetani, F. Sugahara, Y. Kawakami, Y. Murakami, S. Hirano, S. Kuratani


- 1319 **Induction and Suppression of RNA Silencing by an Animal Virus** H. Li, W. X. Li, S. W. Ding

- 1321 **Is Face Processing Species-Specific During the First Year of Life?** O. Pascalis, M. de Haan, C. A. Nelson

- 1323 **Direct Recognition of Cytomegalovirus by Activating and Inhibitory NK Cell Receptors** H. Arase, E. S. Mocarski, A. E. Campbell, A. B. Hill, L. L. Lanier

1305

Tracking the rise of dinosaurs



New on Science Express

Catching lymphocytes in the act



AMERICAN
ASSOCIATION FOR THE
ADVANCEMENT OF
SCIENCE

SCIENCE (ISSN 0036-8075) is published weekly on Friday, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue, NW, Washington, DC 20005. Periodicals Mail postage (publication No. 484460) paid at Washington, DC, and additional mailing offices. Copyright © 2002 by the American Association for the Advancement of Science. The title SCIENCE is a registered trademark of the AAAS. Domestic individual membership and subscription (51 issues): \$120 (\$66 allocated to subscription). Domestic institutional subscription (51 issues): \$390; Foreign postage extra: Mexico, Caribbean (surface mail) \$55; other countries (air assist delivery) \$85. First class, airmail, student, and emeritus rates on request. Canadian rates with GST available upon request, GST #1254 88122. Publications Mail Agreement Number 1069624. Printed in the U.S.A.

Change of address: allow 4 weeks, giving old and new addresses and 8-digit account number. Postmaster: Send change of address to *Science*, P.O. Box 1811, Danbury, CT 06813-1811. Single copy sales: \$9.00 per issue prepaid includes surface postage; bulk rates on request. Authorization to photocopy material for internal or personal use under circumstances not falling within the fair use provisions of the Copyright Act is granted by AAAS to libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that \$8.00 per article is paid directly to CCC, 222 Rosewood Drive, Danvers, MA 01923. The identification code for *Science* is 0036-8075/83 \$8.00. *Science* is indexed in the *Reader's Guide to Periodical Literature* and in several specialized indexes.

DOCKET
ALARM

Find authenticated court documents without watermarks at docketalarm.com.

An Expanding Universe of Noncoding RNAs

Gisela Storz

Noncoding RNAs (ncRNAs) have been found to have roles in a great variety of processes, including transcriptional regulation, chromosome replication, RNA processing and modification, messenger RNA stability and translation, and even protein degradation and translocation. Recent studies indicate that ncRNAs are far more abundant and important than initially imagined. These findings raise several fundamental questions: How many ncRNAs are encoded by a genome? Given the absence of a diagnostic open reading frame, how can these genes be identified? How can all the functions of ncRNAs be elucidated?

Over the years, a number of RNAs that do not function as messenger RNAs (mRNAs), transfer RNAs (tRNAs), or ribosomal RNAs (rRNAs) have been discovered, mostly fortuitously. The non-mRNAs have been given a variety of names (1, 2); the term small RNAs (sRNAs) has been predominant in bacteria, whereas the term noncoding RNAs (ncRNAs) has been predominant in eukaryotes and will be used here. ncRNAs range in size from 21 to 25 nt for the large family of microRNAs (miRNAs) that modulate development in *Caenorhabditis elegans*, *Drosophila*, and mammals (3–8), up to ~100 to 200 nt for sRNAs commonly found as translational regulators in bacterial cells (9, 10) and to >10,000 nt for RNAs involved in gene

silencing in higher eukaryotes (11–13). The functions described for ncRNAs thus far are extremely varied (Table 1).

Some ncRNAs affect transcription and chromosome structure. The *Escherichia coli* 6S RNA binds to the bacterial σ^{70} holoenzyme and modulates promoter use (14), and the human 7SK RNA binds and inhibits the transcription elongation factor P-TEFb (15, 16). Another human ncRNA, SRA RNA, was identified as interacting with progesterone steroid hormone receptor and may serve as a coactivator of transcription (17). Several extremely long ncRNAs detected in insect and mammalian cells have been implicated in silencing genes and changing chromatin structure across large chromosomal regions (11–13). Examples include the human *Xist* RNA required for X chromosome inactivation and mouse *Air* RNA required for autosomal gene imprinting. The *Xist* RNA is produced by the inactive X chromosome and spreads in cis along the chromosome (13). The chromosome-associated RNA has been

proposed to recruit proteins that affect chromatin structure; however, much remains to be learned about the mechanism by which *Xist* and other long ncRNAs establish and/or maintain gene silencing. Another eukaryote-specific RNA that is required for proper chromosome replication and structure is the telomerase RNA. This ncRNA is an integral part of the telomerase enzyme and serves as the template for the synthesis of the chromosome ends (18).

ncRNAs play roles in RNA processing and modification. The catalytic ribonuclease P (RNase P) RNA, found in organisms from all kingdoms, is responsible for processing the 5' end of precursor tRNAs and some rRNAs (19). In eukaryotes, small nuclear RNAs (snRNAs) are central to splicing of pre-mRNAs (20), and small nucleolar RNAs (snoRNAs) direct the 2'-O-ribose methylation (C/D-box type) and pseudouridylation (H/ACA-box type) of rRNA, tRNA, and ncRNAs by forming base pairs with sequences near the sites to be modified (21). Homologs of the two classes of snoRNAs have been found in archaea (22); however, counterparts have not yet been identified in bacteria, even though the rRNAs are modified. The less ubiquitous guide RNAs (gRNAs) present in kinetoplasts direct the insertion or deletion of uridine residues into mRNA (RNA editing) by mechanisms that involve base-pairing as well (23, 24).

ncRNAs also regulate mRNA stability

Cell Biology and Metabolism Branch, National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD 20892, USA. E-mail: storz@helix.nih.gov

Table 1. Processes affected by ncRNAs.

Process	Example	Function	Reference
Transcription	184-nt <i>E. coli</i> 6S	Modulates promoter use	(9, 14)
	331-nt human 7SK	Inhibits transcription elongation factor P-TEFb	(15, 16, 46)
	875-nt human SRA	Steroid receptor coactivator	(12, 17)
Gene silencing	16,500-nt human <i>Xist</i>	Required for X-chromosome inactivation	(12, 13)
	~100,000-nt human <i>Air</i>	Required for autosomal gene imprinting	(11)
Replication	451-nt human telomerase RNA	Core of telomerase and telomere template	(18, 46)
RNA processing	377-nt <i>E. coli</i> RNase P	Catalytic core of RNase P	(9, 19)
	186-nt human U2 snRNA	Core of spliceosome	(20, 46)
RNA modification	102-nt <i>S. cerevisiae</i> U18 C/D snoRNA	Directs 2'-O-ribose methylation of target rRNA	(21, 47)
	189-nt <i>S. cerevisiae</i> snR8 H/ACA snoRNA	Directs pseudouridylation of target rRNA	(21, 47)
	68-nt <i>T. brucei</i> gCYb gRNA	Directs the insertion and excision of uridines	(23, 24, 48)
RNA stability	80-nt <i>E. coli</i> RyhB sRNA	Targets mRNAs for degradation?	(27)
	Eukaryotic miRNA?	Targets mRNAs for degradation?	(7, 8)
mRNA translation	109-nt <i>E. coli</i> OxyS	Represses translation by occluding ribosome binding	(9, 10)
	87-nt <i>E. coli</i> DsrA sRNA	Activates translation by preventing formation of an inhibitory mRNA structure	(9, 10)
	22-nt <i>C. elegans</i> <i>lin-4</i> miRNA	Represses translation by pairing with 3' end of target mRNA	(7, 8)
Protein stability	363-nt <i>E. coli</i> tmRNA	Directs addition of tag to peptides on stalled ribosomes	(9, 28)
Protein translocation	114-nt <i>E. coli</i> 4.5S RNA	Integral component of signal recognition particle central to protein translocation across membranes	(9, 29)

and translation. The first discovered miRNAs, *C. elegans lin-4* and *let-7*, repress translation by forming base pairs with the 3' end of target mRNAs (7, 8). Many of the recently identified miRNAs are likely to act in a similar fashion. However, it is conceivable that some members of this large family target mRNAs for degradation, as is the case for the similarly sized small interfering RNAs (siRNAs) that are processed and amplified from exogenously added, double-stranded RNA and lead to gene suppression in a process termed RNA interference (25, 26). As yet there is no evidence for miRNAs in bacteria, archaea, or fungi, but it might be fruitful to search for RNAs of <25 nt in these organisms. Several ncRNAs have been found to regulate translation and possibly mRNA stability in *E. coli* (9, 10, 27). These sRNAs form base pairs at various positions with their target mRNAs, and they have been shown to repress translation by occluding the ribosome binding site and to activate translation by

preventing the formation of inhibitory mRNA structures.

Finally, ncRNAs affect protein stability and transport. One unique bacterial sRNA is recognized as both a tRNA and an mRNA by stalled ribosomes (tmRNA) (28). Alanylated tmRNA is delivered to the A site of a stalled ribosome; the nascent polypeptide is transferred to the alanine-charged tRNA portion of tmRNA. The problematic transcript then is replaced by the mRNA portion of tmRNA, which encodes a tag for degradation of the stalled peptide. It is not yet clear whether there is a counterpart to this coding RNA in archaeal and eukaryotic cells. In contrast, a small cytoplasmic RNA that forms the core of the signal recognition particle (SRP) required for protein translocation across membranes is found in organisms from all kingdoms (29).

The mechanisms of action for the characterized ncRNAs can be grouped into several general categories (Fig. 1). There are

ncRNAs where base-pairing (often <10 base pairs and discontinuous) with another RNA or DNA molecule is central to function. The snoRNAs that direct RNA modification, the bacterial RNAs that modulate translation by forming base pairs with specific target mRNAs, and probably most of the miRNAs are examples of this category. Some ncRNAs mimic the structures of other nucleic acids; the 6S RNA structure is reminiscent of an open bacterial promoter, and the tmRNA has features of both tRNAs and mRNAs. Other ncRNAs, such as the RNase P RNA, have catalytic functions. Although synthetic RNAs have been selected to have a variety of biochemical functions, the number of natural ncRNAs shown to have catalytic function is limited. Most, if not all, ncRNAs are associated with proteins that augment their functions; however, some ncRNAs, such as the snRNAs and the SRP RNA, serve key structural roles in RNA-protein complexes. Several ncRNAs fit into more than one mechanistic category; the telomerase RNA provides the base-pairing template for telomere synthesis and is an integral part of the telomerase ribonucleoprotein complex. The mechanisms of action for a number of ncRNAs (such as the 7SK RNA) are not known, and it is probable that some ncRNAs act in ways that have not yet been established. Some investigators have suggested that many ncRNAs are vestiges of a world in which RNA carried out all of the functions in a primitive cell. However, given the versatility of RNA and the fact that the properties of RNA provide advantages over peptides for some mechanisms, it is likely that a number of ncRNAs have evolved more recently (30, 31).

How Many ncRNAs Exist?

The first ncRNAs were identified in the 1960s on the basis of their high expression; these RNAs were detected by direct labeling and separation on polyacrylamide gels. Others were later found by subfractionation of nuclear extracts or by association with specific proteins. A few were identified by mutations or phenotypes resulting from overexpression. The serendipitous discoveries of many of these ncRNAs were the first glimpses of their existence, but this work did not presage the vast numbers that appear to be encoded by a genome.

Several systematic searches for ncRNA genes have been carried out in the past 4 years. Among the computation-based searches, there have been screens of the yeast *Saccharomyces cerevisiae* and archaeal *Pyrococcus* genomes for the short conserved motifs present in snoRNAs (32, 33). In other searches, the intergenic regions of *S. cerevisiae*, *E. coli*, *Methanococcus jannaschii*, and *Pyrococcus furiosus* chromosomes have been scanned for properties indicative of an




Fig. 1. Different mechanisms of ncRNA (red) action. (A) Direct base-pairing with target RNA or DNA molecules is central to the function of some ncRNAs: Eukaryotic snoRNAs direct nucleotide modifications (green star) by forming base pairs with flanking sequences, and the *E. coli* OxyS RNA represses translation by forming base pairs with the Shine-Dalgarno sequence (green box) and occluding ribosome binding. (B) Some ncRNAs mimic the structure of other nucleic acids: Bacterial RNA polymerase may recognize the 6S RNA as an open promoter, and bacterial ribosomes recognize tmRNA as both a tRNA and an mRNA. (C) ncRNAs also can function as an integral part of a larger RNA-protein complex, such as the signal recognition particle, whose structure has been partially determined (49).

RNA SILENCING AND NONCODING RNA

ncRNA gene. Criteria for identifying candidate intergenic regions have included large gaps between protein-coding genes (34), extended stretches of conservation between species with the same gene order (35, 36), orphan promoter or terminator sequences (34, 36, 37), presence of GC-rich regions in an organisms with a high AT content (38), and conserved RNA secondary structures (39, 40). Other searches for ncRNAs have involved large-scale cloning efforts that have taken into account specific ncRNA properties. In studies of mouse (41, 42) and the archaeon *Archaeoglobus fulgidus* (22), total RNA between 50 to 500 nt was isolated, and arrays of cDNA clones obtained from the RNA were screened with oligonucleotides corresponding to the most abundant known RNAs. Clones showing the lowest hybridization signal then were randomly sequenced. In recent screens for *C. elegans*, *Drosophila*, and human miRNAs, RNA molecules of less than 30 nt were isolated, and cDNA clones were generated upon the ligation of primers to the 5' and 3' ends of the RNA (3, 4) or upon RNA tailing (5). Other miRNAs were isolated and cloned on the basis of their association with a complex composed of the human Gemin3, Gemin4, and IF2C proteins (6). In most studies, Northern blots have been carried out to confirm that the cloned genes are expressed as small transcripts. These blots also have provided information about spatial and temporal expression patterns as well as potential precursor and degradation products.

Despite the success of the recent systematic efforts, it is certain that not all ncRNAs have been detected. Estimates for the number of sRNAs in *E. coli* range from 50 to 200 (1, 35), and estimates for the number of miRNAs in *C. elegans* range from hundreds to thousands (7). There also are many non-protein-coding regions of the bacterial and eukaryotic chromosomes for which transcription is detected (43, 44), but it is not known how many of these regions encode defined, functional ncRNAs. Extensions of the various systematic searches should lead to the identification of more ncRNAs. However, limitations of the current approaches should be noted. Most of the computation methods have focused on the intergenic regions. It has already been shown that some of the ncRNAs are processed from longer protein- or rRNA-encoding transcripts (42). It also is quite possible that ncRNAs are expressed from the opposite strand of protein-coding genes. On the other hand, expression-based methods may miss ncRNAs that are synthesized under very defined conditions, such as in response to a specific environmental signal, during a specific stage in development, or in a specific cell type. Much attention has been focused on characterizing the "proteome" of a sequenced organism. The

recent discovery of hundreds of new ncRNAs illustrates that the "RNome" also will need to be characterized before a complete tally of the number of genes encoded by a genome can be achieved.

What Are All the Functions of ncRNAs?

An astonishing variety of ncRNA functions have already been found, but there are many ncRNAs for which the cellular roles are still unknown. For instance, Y RNAs, small cytoplasmic RNAs associated with the Ro autoantigen in several different organisms, are still enigmatic even after many years of study (45). With the more systematic identification of increasing numbers of ncRNAs, the question of how to elucidate the functions of all ncRNAs is becoming more and more prominent.

Approaches that have succeeded previously are an obvious place to start in answering the question of function, but it is likely that new approaches also will need to be developed. For genetically tractable organisms, ncRNA knock-out or overexpression strains can be screened for differences in phenotypes (such as viability) or whole-genome expression patterns. The functions of several ncRNAs were identified by the biochemical identification of associated proteins, and the development of more systematic methods for characterizing ncRNA-associated proteins should be fruitful. As the knowledge base of what sequences are critical for the formation of specific structures or for base-pairing expands, and as computer programs for predicting structures improve, computational approaches should become an increasingly important avenue for elucidating the functions of ncRNAs. The three-dimensional structures of only a limited number of RNAs and RNA-protein complexes have been solved. An increase in the structural database may bring to light recognizable RNA or RNA-protein domains associated with specific functions.

Information about when ncRNAs are expressed and where ncRNAs are localized is useful for all experiments aimed at probing function. Many of the *C. elegans* miRNAs are synthesized only at very specific times in development, and thus they have also been called small temporal RNAs (stRNAs). Among the snoRNAs, some are expressed exclusively in the brain (41), and one of the bacterial sRNAs is only detected upon oxidative stress (9, 10). It is likely that other ncRNAs will be found to have very defined expression and localization patterns and that these will be critical to function.

There are many more ncRNAs than was ever suspected. A big challenge for the future will be to identify the whole complement of ncRNAs and to elucidate their functions. This is an exciting time for investigators whose work has focused on ncRNAs. However, scientists

studying all aspects of biology should keep ncRNAs in mind. The phenotypes associated with specific mutations may be due to defects in a ncRNA instead of being due to defects in a protein, as is usually expected. Investigators developing purification schemes for specific proteins or activities should be aware of the possible presence of an ncRNA component; many purification procedures are designed to remove nucleic acids. There may be ncRNAs lurking behind many an unexplained phenomenon.

References and Notes

1. S. R. Eddy, *Nature Rev. Genet.* **2**, 919 (2001).
2. Non-mRNAs have been denoted ncRNA = noncoding RNA, snmRNA = small non-mRNA, sRNA = small RNA, fRNA = functional RNA, and oRNA = other RNA, and it is likely that the nomenclature of these RNAs will need to be revisited.
3. M. Lagos-Quintana, R. Rauhut, W. Lendeckel, T. Tuschl, *Science* **294**, 853 (2001).
4. N. C. Lau, L. P. Lim, E. G. Weinstein, D. P. Bartel, *Science* **294**, 858 (2001).
5. R. C. Lee, V. Ambros, *Science* **294**, 862 (2001).
6. Z. Mourelatos *et al.*, *Genes Dev.* **16**, 720 (2002).
7. G. Ruvkun, *Science* **294**, 797 (2001).
8. H. Grosshans, F. J. Slack, *J. Cell Biol.* **156**, 17 (2002).
9. K. M. Wassarman, A. Zhang, G. Storz, *Trends Microbiol.* **7**, 37 (1999).
10. S. Altuvia, E. G. H. Wagner, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 9824 (2000).
11. F. Sletstels, R. Zwart, D. P. Barlow, *Nature* **415**, 810 (2002).
12. V. A. Erdmann, M. Szymanski, A. Hochberg, N. de Groot, J. Barciszewski, *Nucleic Acids Res.* **28**, 197 (2000).
13. P. Avner, E. Heard, *Nature Rev. Genet.* **2**, 59 (2001).
14. K. M. Wassarman, G. Storz, *Cell* **101**, 613 (2000).
15. Z. Yang, Q. Zhu, K. Luo, Q. Zhou, *Nature* **414**, 317 (2001).
16. V. T. Nguyen, T. Kiss, A. A. Michels, O. Bensaude, *Nature* **414**, 322 (2001).
17. R. B. Lánz *et al.*, *Cell* **97**, 17 (1999).
18. J.-L. Chen, M. A. Blasco, C. W. Greider, *Cell* **100**, 503 (2000).
19. D. N. Frank, N. R. Pace, *Annu. Rev. Biochem.* **67**, 153 (1998).
20. C. L. Will, R. Lührmann, *Curr. Opin. Cell Biol.* **13**, 290 (2001).
21. T. Kiss, *EMBO J.* **20**, 3617 (2001).
22. T.-H. Tang *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, in press.
23. M. L. Kable, S. Heidmann, K. D. Stuart, *Trends Biochem. Sci.* **22**, 162 (1997).
24. L. Simpson, O. H. Thiemann, N. J. Savill, J. D. Alfonzo, D. A. Maslov, *Proc. Natl. Acad. Sci. U.S.A.* **97**, 6986 (2000).
25. K. Nishikura, *Cell* **107**, 415 (2001).
26. P. D. Zamore, *Science* **296**, 1265 (2002).
27. E. Massé, S. Gottesman, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 4620 (2002).
28. R. Gillet, B. Felden, *Mol. Microbiol.* **42**, 879 (2001).
29. R. J. Keenan, D. M. Freymann, R. M. Stroud, P. Walter, *Annu. Rev. Biochem.* **70**, 755 (2001).
30. V. Y. Kuryshv, B. V. Skryabin, J. Kremerskothen, J. Jurka, J. Brosius, *J. Mol. Biol.* **309**, 1049 (2001).
31. W. Wang, F. G. Brunet, E. Nevo, M. Long, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 4448 (2002).
32. T. M. Lowe, S. R. Eddy, *Science* **283**, 1168 (1999).
33. C. Gaspin, J. Cavaille, G. Erauso, J.-P. Bachellerie, *J. Mol. Biol.* **297**, 895 (2000).
34. W. M. Olivas, D. Muhlrad, R. Parker, *Nucleic Acids Res.* **25**, 4619 (1997).
35. K. M. Wassarman, F. Repoila, C. Rosenow, G. Storz, S. Gottesman, *Genes Dev.* **15**, 1637 (2001).
36. L. Argaman *et al.*, *Curr. Biol.* **11**, 941 (2001).

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.