



[For Healthcare Professionals \(/tablet/hcp\)](#) [Important Safety Information](#) [Prescribing Information \(PDF\) \(http://shared.salix.com/sha](#)



A different kind of treatment for UC

UCERIS is a different kind of treatment for ulcerative colitis. It is specifically designed to target the colon and helps eliminate the symptoms of UC, with a safety profile similar to placebo (sugar pill). UCERIS is a prescription corticosteroid medicine used to help get mild to moderate UC under control.

[LEARN MORE ABOUT UCERIS \(/TABLET/ABOUT-UCERIS/\)](#)

Long-time use of UCERIS extended release tablets can cause you to have too much glucocorticosteroid medicine in your system. Talk to your healthcare provider if you have any of the following signs and symptoms of hypercorticism:

- acne
- bruise easily
- rounding of your face (moon face)
- ankle swelling
- thicker or more hair on your body and face
- a fatty pad or hump between your shoulders (buffalo hump)
- pink or purple stretch marks on the skin of your abdomen, thighs, breasts and arms

Schedule a doctor visit

Ask if UCERIS can help put you back in control. Be prepared to discuss details about your symptoms and how it's affecting your life.

[DOWNLOAD DOCTOR DISCUSSION GUIDE \(/PORTALS/106/ASSETS/PDF/UC-DISCUSSION-GUIDE.PDF\)](#)

Exhibit 1017

MOST ELIGIBLE PATIENTS WITH COMMERCIAL INSURANCE

PAY ONLY \$25 FOR EACH UCERIS PRESCRIPTION

(1)

Indication for UCERIS

UCERIS® extended release tablets are a prescription corticosteroid medicine used to help get mild to moderate ulcerative colitis (UC) under control (induce remission).

Important Safety Information about UCERIS

UCERIS® (budesonide) extended release tablets are not for everyone. Do not take UCERIS extended release tablets if you are allergic to budesonide or any of the ingredients in UCERIS.

UCERIS is a steroid. When UCERIS extended release tablets are taken for a long period of time, the adrenal glands do not make enough steroid hormones. Tell your healthcare provider if you are under stress or have any of the following symptoms: tiredness, vomiting, weakness, low blood pressure, and nausea.

UCERIS may weaken your immune system. Taking medicines that weaken your immune system makes you more likely to get infections. Avoid contact with people who have contagious diseases such as chicken pox or measles while taking UCERIS extended release tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter vitamins, and herbal supplements. UCERIS extended release tablets and other medicines may affect each other and cause side effects. If you take certain other steroid medicines to treat allergies, switching to UCERIS extended release tablets may cause your allergies to come back. These allergies may include eczema (a skin disease) or rhinitis (inflammation inside your nose). Tell your healthcare provider if any of your allergies become worse while taking UCERIS extended release tablets.


Before you take UCERIS extended release tablets tell your healthcare provider if you have liver problems, are planning to have surgery, have an infection, have or had a family history of diabetes, cataracts or glaucoma, have or had tuberculosis, have high blood pressure (hypertension), have decreased bone mineral density (osteoporosis), stomach ulcers, or any other medical condition. Do not eat grapefruit or drink grapefruit juice while taking UCERIS. Eating grapefruit or drinking grapefruit juice can increase the level of this medicine in your blood.


In clinical studies, the most common side effects (occurring in $\geq 2\%$ of people) were headache, nausea, decreased blood cortisol (a hormone made by the adrenal glands), stomach-area pain, tiredness, stomach or intestinal gas, bloating, acne, urinary tract infection, joint pain, and constipation.


You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch/ (<http://www.fda.gov/medwatch/>) or call 1-800-FDA-1088.


For product information, adverse event reports, and product complaint reports, please contact:

Salix Product Information Call Center
Phone: 1-800-508-0024
Fax: 1-510-595-8183
Email: salix@medcomsol.com (<mailto:salix@medcomsol.com>)

Please see complete Prescribing Information for UCERIS extended release tablets
(<http://shared.salix.com/shared/pi/uceris-pi.pdf>).  (<http://shared.salix.com/shared/pi/uceris-pi.pdf>)

- [About UCERIS \(/tablet/about-uceris/\)](/tablet/about-uceris/)
- [How UCERIS Works \(/tablet/how-uceris-provides-relief/\)](/tablet/how-uceris-provides-relief/)
- [Why UCERIS \(/tablet/uceris-benefits/\)](/tablet/uceris-benefits/)
- [Taking UCERIS \(/tablet/taking-uceris/\)](/tablet/taking-uceris/)
- [Back To Top](#) 

- [Understanding UC \(/tablet/what-is-ulcerative-colitis/\)](/tablet/what-is-ulcerative-colitis/)
- [Resources \(/tablet/ulcerative-colitis-resources/\)](/tablet/ulcerative-colitis-resources/)
- [Back To Top](#) 

- [Back To Top](#) 

The health information contained herein is provided for educational purposes only and is not intended to replace discussions with a healthcare provider. All decisions regarding patient care must be made with a healthcare provider, considering the unique characteristics of the patient.

The product information provided in this site is intended only for residents of the United States. The products discussed herein may have different product labeling in different countries.

UCERIS is a registered trademark of Santarus, Inc, a wholly owned subsidiary of Salix Pharmaceuticals, Inc.



Follow Salix on:



(<http://www.facebook.com/SalixPharma>)



(<http://www.twitter.com/SalixPharma>)

Use of this site signifies your agreement to the Terms of Use (</tablet/terms-of-use/>). Privacy Policy (</tablet/privacy-policy/>).

©2015 Salix Pharmaceuticals, Inc. All rights reserved.

8510 Colonnade Center Drive, Raleigh, NC 27615 USA

T: 919-862-1000 (tel:9198621000), F: 919-862-1095 (tel:9198621095)

UCE3-0514