



US008784888B2

(12) **United States Patent**
Villa et al.

(10) **Patent No.:** **US 8,784,888 B2**
(45) **Date of Patent:** ***Jul. 22, 2014**

(54) **CONTROLLED RELEASE AND TASTE MASKING ORAL PHARMACEUTICAL COMPOSITION**

(75) Inventors: **Roberto Villa**, Lecco (IT); **Massimo Pedrani**, Gignese (IT); **Mauro Ajani**, Milan (IT); **Lorenzo Fossati**, Milan (IT)

(73) Assignee: **Cosmo Technologies Limited**, Dublin (IE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

5,955,502 A	9/1999	Hansen et al.	
5,965,167 A	10/1999	Sanghvi	
6,140,308 A	10/2000	Brattsand	
6,190,692 B1	2/2001	Busetti	
6,239,120 B1 *	5/2001	Hallgren et al.	514/174
6,258,377 B1	7/2001	New	
6,363,635 B1	4/2002	England	
6,368,629 B1	4/2002	Watanabe et al.	
6,562,363 B1	5/2003	Mantelle	
7,410,651 B2	8/2008	Villa et al.	
7,410,652 B2	8/2008	Villa et al.	
7,431,943 B1	10/2008	Villa et al.	
8,029,823 B2	10/2011	Villa et al.	
2006/0134208 A1 *	6/2006	Villa et al.	424/468
2012/0021052 A1	1/2012	Villa et al.	
2012/0021053 A1	1/2012	Villa et al.	

FOREIGN PATENT DOCUMENTS

CA	2119253	11/1998
EP	0 375 063 B1	6/1990
EP	0 453 001 A1	10/1991
EP	0482576	4/1992
EP	0 514 008	11/1992
EP	0 514 008 A1	11/1992
EP	0514008	11/1992
GB	935639	9/1963
WO	93/00889 A1	1/1993
WO	96/13273	5/1996
WO	9800169	1/1998
WO	WO 99/11245	3/1999
WO	WO 99/17752	4/1999

(21) Appl. No.: **13/617,138**

(22) Filed: **Sep. 14, 2012**

(65) **Prior Publication Data**

US 2013/0022679 A1 Jan. 24, 2013

Related U.S. Application Data

(63) Continuation of application No. 13/462,409, filed on May 2, 2012, now Pat. No. 8,293,273, which is a continuation of application No. 13/249,839, filed on Sep. 30, 2011, which is a continuation of application No. 12/210,969, filed on Sep. 15, 2008, now Pat. No. 8,029,823, which is a continuation-in-part of application No. 10/009,532, filed as application No. PCT/EP00/05356 on Jun. 9, 2000, now Pat. No. 7,431,943.

(30) **Foreign Application Priority Data**

Jun. 14, 1999 (IT) MI99A1317
Mar. 3, 2000 (IT) MI2000A0422

(51) **Int. Cl.**
A61K 9/28 (2006.01)
A61K 9/20 (2006.01)

(52) **U.S. Cl.**
USPC **424/474**; 424/464; 424/465; 424/469

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,800,051 A	3/1974	Barnhart et al.	
4,608,248 A	8/1986	Knecht et al.	
5,320,848 A	6/1994	Geyer	
5,342,625 A	8/1994	Hauer	
5,534,501 A	7/1996	Samain	
5,597,844 A	1/1997	Chauhan	
5,643,602 A	7/1997	Ulmius	
5,681,584 A *	10/1997	Savastano et al.	424/473
5,811,388 A *	9/1998	Friend et al.	424/474
5,840,332 A *	11/1998	Lerner et al.	424/464
5,874,063 A	2/1999	Briggner et al.	
5,908,833 A	6/1999	Brattsand et al.	

OTHER PUBLICATIONS

Travis, S. et al., Poster, "Induction of Clinical and Endoscopic Remission with Budesonide MMX in Mild-to-Moderately Active Ulcerative Colitis, Magnitude of Response in Two Phase III Studies," Oct. 20-24, 2012, Amsterdam, UEG Week, 1 page.
Lichtenstein, G. et al., Poster, "Effect of Budesonide MMX 6 mg on the Hypothalamic-Pituitary-Adrenal (HPA) Axis in Patients with Ulcerative Colitis: Results from Phase III, 12 Month Safety and Extended Use Study," May 2012, San Diego, CA, 1 page.

(Continued)

Primary Examiner — Susan Tran

(74) Attorney, Agent, or Firm — Rothwell, Figg, Ernst & Manbeck P.C.

(57) **ABSTRACT**

Controlled release and taste masking compositions containing one or more active principles inglobated in a three-component matrix structure, i.e. a structure formed by successive amphiphilic, lipophilic or inert matrices and finally inglobated or dispersed in hydrophilic matrices. The use of a plurality of systems for the control of the dissolution of the active ingredient modulates the dissolution rate of the active ingredient in aqueous and/or biological fluids, thereby controlling the release kinetics in the gastrointestinal tract.

9 Claims, No Drawings

(56)

References Cited

OTHER PUBLICATIONS

- Sandborn, W.J. et al., "Once-Daily Budesonide MMX® Extended Release Tablets Induce Remission in Patients With Mild to Moderate Ulcerative Colitis: Results From the CORE I Study," *Gastroenterology* 2012, vol. 143, pp. 1218-1226.
- Jantzen, G.M. et al., "Sustained- and Controlled-Release Drug Delivery Systems," *Modern Pharmaceutics*, 3rd Edition, Revised and Expanded, pp. 575-609, © 1996 by Marcel Dekker, Inc., 37 pages.
- Angelucci et al., "Budesonide for Inflammatory Bowel Disease Treatment," *Current Medicinal Chemistry*, 2008, vol. 15, No. 14, pp. 2-9.
- D'Haens, G.R. et al., "Budesonide MMX™ Is Active and Safe in Patients With Active Left-Sided Ulcerative Colitis," *Br J Clin Pharmacol.*, 2005, vol. 61, 3 pages.
- Maejima, T., "Application of Tumbling Melt Granulation Method to Prepare Controlled-Release Beads by Coating with Mixture of Functional Non-Meltable and Meltable Materials," *Chem. Pharm. Bull.*, 1998, Vol. 46, No. 3, pp. 531-533, © 1998 Pharmaceutical Society of Japan.
- Sandborn, W.J. et al., "Budesonide MXX® 9 mg for the Induction of Remission of Mild-to-Moderate Ulcerative Colitis (UC): Data From a Multicenter, Randomized, Double-Blind Placebo-Controlled Study in North America and India," Presentation at DDW 2011, Poster, 1 page.
- D'Haens, G.R., et al., "Clinical Trial: Preliminary Efficacy and Safety Study of a New Budesonide-MMX® 9 mg Extended-Release Tablets in Patients With Active Left-Sided Ulcerative Colitis," *Journal of Crohn's and Colitis*, 2010, vol. 4, pp. 153-160, © copyright 2009 European Crohn's and Colitis Organisation.
- Flanders, P. et al., *The Control of Drug Release From Conventional Melt Granulation Matrices*, Drug Development and Industrial Pharmacy, 1987, vol. 13, No. 6, pp. 1001-1022, © 1987 Marcel Dekker, Inc.
- Ferraboschi, P. et al., "Estimation and Characterisation of Budesonide Tablets Impurities," *Journal of Pharmaceutical and Biomedical Analysis*, 2008, vol. 47, pp. 636-640, © 2008 Elsevier B.V.
- Fiorino, G. et al., "New Drug Delivery Systems in Inflammatory Bowel Disease: MMX™ and Tailored Delivery to the Gut," *Current Medicinal Chemistry*, 2010, vol. 17, pp. 1851-1857, © 2010 Bentham Science Publishers Ltd.
- Koutroubakis, I., "Recent Advances in the Management of Distal Ulcerative Colitis," *World Journal of Gastrointestinal Pharmacology and Therapeutics*, 2010, vol. 1, No. 2, pp. 43-50, © 2010 Baishideng.
- Steward, P., "Review of Pharmaceutical Controlled Release Methods and Devices", 1995, pp. 1-9.
- Physical Pharmacy*, Chapter 19: Drug Product Design, Oct. 1993, pp. 515-519.
- Moro, et al., "Drug Delivery Systems: Diffusion Controlled Systems", *Il Prodotto Chimico & Aerosol Selezione (The Chemical & Aerosol Selection)*, Apr. 1985, pp. 16-24.
- Brunner, M. et al., "Gastrointestinal Transit, Release and Plasma Pharmacokinetics of a New Oral Budesonide Formulation," *British Journal of Clinical Pharmacology*, DOI:10.1111/j.1365-2125.2005.02517.x, pp. 1-8, copyright 2005 Blackwell Publishing Ltd., 8 pages.
- Brunner, M. et al., "Gastrointestinal Transit and 5-ASA Release From a New Mesalazine Extended-Release Formulation," *Alimentary Pharmacology and Therapeutics*, vol. 17, pp. 395-402, copyright 2003 Blackwell Publishing Ltd., 8 pages.

* cited by examiner

1

**CONTROLLED RELEASE AND TASTE
MASKING ORAL PHARMACEUTICAL
COMPOSITION**

CROSS REFERENCE TO RELATED
APPLICATIONS

This application is a continuation of application Ser. No. 13/462,409 filed on May 2, 2012, now U.S. Pat. No. 8,293,273; which is a continuation of Ser. No. 13/249,839 filed on Sep. 30, 2011; which is a continuation of application Ser. No. 12/210,969 filed on Sep. 15, 2008, which reissued as U.S. Pat. No. RE43,799 from U.S. Pat. No. 8,029,823; which is a continuation-in-part of application Ser. No. 10/009,532 filed on Dec. 12, 2001, now U.S. Pat. No. 7,431,943; which is the 35 U.S.C. 371 national stage of International application PCT/EP00/05356 filed on Jun. 9, 2000; which claimed priority to Italian applications MI2000A000422 and MI99A001317 filed Mar. 3, 2000 and Jun. 14, 1999, respectively. The entire contents of each of the above-identified applications are hereby incorporated by reference.

The present invention relates to controlled release and taste-masking compositions containing one or more active principles incorporated in a three-component matrix structure, i.e. a structure formed by successive amphiphilic, lipophilic or inert matrices and finally incorporated or dispersed in hydrophilic matrices. The use of a plurality of systems for the control of the dissolution of the active ingredient modulates the dissolution rate of the active ingredient in aqueous and/or biological fluids, thereby controlling the release kinetics in the gastrointestinal tract, and it also allows the oral administration of active principles having unfavourable taste characteristics or irritating action on the mucosae of the administration site, particularly in the buccal area.

The compositions of the invention can contain active principles belonging to the therapeutical classes of analgesics, antiinflammatories, cardioactives, tranquillizers, antihypertensives, disinfectants and topical antimicrobials, antiparkinson drugs, antihistamines and are suitable to the oral administration or for acting topically at some areas of the gastrointestinal tract.

TECHNOLOGICAL BACKGROUND

The preparation of a sustained, controlled, delayed or anyhow modified release form can be carried out according to different known techniques:

1. The use of inert matrices, in which the main component of the matrix structure opposes some resistance to the penetration of the solvent due to the poor affinity towards aqueous fluids; such property being known as lipophilia.
2. The use of hydrophilic matrices, in which the main component of the matrix structure opposes high resistance to the progress of the solvent, in that the presence of strongly hydrophilic groups in its chains, mainly branched, remarkably increases viscosity inside the hydrated layer.
3. The use of bioerodible matrices, which are capable of being degraded by the enzymes of some biological compartment.

All the procedures listed above suffer, however, from drawbacks and imperfections.

Inert matrices, for example, generally entail non-linear, but exponential, release of the active ingredient.

2

Hydrophilic matrices have a linear behaviour until a certain fraction of active ingredient has been released; then they significantly deviate from linear release.

Bioerodible matrices are ideal to carry out the so-called "site-release", but they involve the problem of finding the suitable enzyme or reactive to degradation. Furthermore, they frequently release in situ metabolites that are not wholly toxicologically inert.

A number of formulations based on inert lipophilic matrices have been described: Drug Dev. Ind. Pharm. 13 (6), 1001-1022, (1987) discloses a process making use of varying amounts of colloidal silica as a porization element for a lipophilic inert matrix in which the active ingredient is incorporated.

The same notion of canalization of an inert matrix is described in U.S. Pat. No. 4,608,248 in which a small amount of a hydrophilic polymer is mixed with the substances forming an inert matrix, in a non sequential compenetration of different matrix materials.

EP 375,063 discloses a technique for the preparation of multiparticulate granules for the controlled-release of the active ingredient which comprises co-dissolution of polymers or suitable substances to form an inert matrix with the active ingredient and the subsequent deposition of said solution on an inert carrier which acts as the core of the device. Alternatively, the inert carrier is kneaded with the solution containing the inert polymer and the active ingredient, then the organic solvent used for their dissolution is evaporated off to obtain a solid residue. The resulting structure is a "reservoir", i.e. is not macroscopically homogeneous along all the symmetry axis of the final form.

The same "reservoir" structure is also described in Chem. Pharm. Bull. 46 (3), 531-533, (1998) which improves the application through an annealing technique of the inert polymer layer which is deposited on the surface of the pellets.

To the "reservoir" structure also belong the products obtained according to the technique described in WO 93/00889 which discloses a process for the preparation of pellets in hydrophilic matrix which comprises:

- dissolution of the active ingredient with gastro-resistant hydrophilic polymers in organic solvents;
- drying of said suspension;
- subsequent kneading and formulation of the pellets in a hydrophilic or lipophilic matrix without distinction of effectiveness between the two types of application.

EP 0 453 001 discloses a multiparticulate with "reservoir" structure inserted in a hydrophilic matrix. The basic multiparticulate utilizes two coating membranes to decrease the release rate of the active ingredient, a pH-dependent membrane with the purpose of gastric protection and a pH-independent methacrylic membrane with the purpose of slowing down the penetration of the aqueous fluid.

WO 95/16451 discloses a composition only formed by a hydrophilic matrix coated with a gastro-resistant film for controlling the dissolution rate of the active ingredient.

When preparing sustained-, controlled-release dosage forms of a medicament topically active in the gastrointestinal tract, it is important to ensure a controlled release from the first phases following administration, i.e. when the inert matrices have the maximum release rate inside the logarithmic phase, namely the higher deviation from linear release.

Said object has been attained according to the present invention, through the combination of an amphiphilic matrix inside an inert matrix, the latter formulated with a lipophilic polymer in a superficial hydrophilic matrix. The compositions of the invention are characterized by the absence of a first phase in which the medicament superficially present on

3

the matrix is quickly solubilized, and by the fact the amphiphilic layer compensates the lack of affinity of the aqueous solvent with the lipophilic compounds forming the inner inert matrix.

DISCLOSURE OF THE INVENTION

The invention provides controlled release and taste masking oral pharmaceutical compositions containing an active ingredient, comprising:

- a) a matrix consisting of lipophilic compounds with melting point lower than 90° C. and optionally by amphiphilic compounds in which the active ingredient is at least partially incorporated;
- b) optionally an amphiphilic matrix;
- c) an outer hydrophilic matrix in which the lipophilic matrix and the optional amphiphilic matrix are dispersed;
- d) optionally other excipients.

A particular aspect of the invention consists of controlled release oral compositions containing one or more active ingredients comprising:

- a) a matrix consisting of amphiphilic compounds and lipophilic compounds with melting point below 90° C. in which the active ingredient is at least partially incorporated;
- b) an outer hydrophilic matrix in which the lipophilic/amphiphilic matrix is dispersed;
- c) optional other excipients.

A further aspect of the invention provides taste masking oral pharmaceutical compositions containing one or more active ingredients comprising:

- an inert or lipophilic matrix consisting of C6-C20-alcohols or C8-C20 fatty acids or esters of fatty acids with glycerol or sorbitol or other polyalcohols with carbon atom chain not higher than six;
- an amphiphilic matrix consisting of polar lipids of type I or II or glycols partially etherified with C1-C4 alkyl chains;
- an outer hydrophilic matrix containing the above matrices, mainly formed by saccharide, dextrin, polyalcohol or cellulose compounds or by hydrogels;
- optional excipients to give stability to the pharmaceutical formulation.

DETAILED DISCLOSURE OF THE INVENTION

The compositions of the invention can be prepared by a method comprising the following steps:

a) the active ingredient is first inglobated by simple kneading or mixing in a matrix or coating consisting of compounds having amphiphilic properties, which will be further specified below. The active principle(s) can be mixed with the amphiphilic compounds without the aid of solvents or with small amounts of water-alcoholic solvents.

b) The matrix obtained in a) is incorporated in a low melting lipophilic excipient or mixture of excipients, while heating to soften and/or melt the excipient itself, which thereby incorporates the active ingredient by simple dispersion. After cooling at room temperature an inert matrix forms, which can be reduced in size to obtain inert matrix granules containing the active ingredient particles.

c) The inert matrix granules are subsequently mixed together with one or more hydrophilic water-swellaable excipients. The mixture is then subjected to compression or tableting. This way, when the tablet is contacted with biological fluids, a high viscosity swollen layer is formed, which coordinates the solvent molecules and acts as a barrier to penetration of the aqueous fluid itself inside the new structure. Said barrier antagonizes the starting "burst effect"

4

caused by the dissolution of the medicament inglobated inside the inert matrix, which is in its turn inside the hydrophilic matrix.

The amphiphilic compounds which can be used according to the invention comprise polar lipids of type I or II (lecithin, phosphatidylcholine, phosphatidylethanolamine), ceramides, glycol alkyl ethers such as diethylene glycol monomethyl ether (Transcutol^(R)).

The lipophilic matrix consists of substances selected from unsaturated or hydrogenated alcohols or fatty acids, salts, esters or amides thereof, fatty acids mono-, di- or triglycerides, the polyethoxylated derivatives thereof, waxes, ceramides, cholesterol derivatives or mixtures thereof having a melting point within the range of 40 to 90° C., preferably from 60 to 70° C.

If desired, a fatty acid calcium salt may be incorporated in the lipophilic matrix which is subsequently dispersed in a hydrophilic matrix prepared with alginic acid, thus remarkably increasing the hydrophilic matrix viscosity following penetration of the solvent front until contact with the lipophilic matrix granules dispersed inside.

According to an embodiment of the invention, an amphiphilic matrix with high content in active ingredient, typically from 5 to 95% w/w, is first prepared by dispersing the active ingredient or the mixture of active ingredients in a mixture of amphiphilic compounds, such as lecithin, other type II polar lipids, surfactants, or in diethylene glycol monoethyl ether; the resulting amphiphilic matrix is then mixed or kneaded, usually while hot, with lipophilic compounds suitable to form an inert matrix, such as saturated or unsaturated fatty acids, such as palmitic, stearic, myristic, lauric, laurylic, or oleic acids or mixtures thereof with other fatty acids with shorter chain, or salts or alcohols or derivatives of the cited fatty acids, such as mono-, di-, or triglycerides or esters with polyethylene glycols, alone or in combination with waxes, ceramides, cholesterol derivatives or other apolar lipids in various ratios so that the melting or softening points of the lipophilic compounds mixtures is within the range of 40 to 90° C., preferably from 60 to 70° C.

Alternatively, the order of formation of the inert and amphiphilic matrices can be reversed, incorporating the inert matrix inside the amphiphilic compounds.

The resulting inert lipophilic matrix is reduced into granules by an extrusion and/or granulation process, or any other known processes which retain the homogeneous dispersion and matrix structure of the starting mixture.

The hydrophilic matrix consists of excipients known as hydrogels, i.e. substances which when passing from the dry state to the hydrated one, undergo the so-called "molecular relaxation", namely a remarkable increase in mass and weight following the coordination of a large number of water molecules by the polar groups present in the polymeric chains of the excipients themselves.

Examples of hydrogels which can be used according to the invention are compounds selected from acrylic or methacrylic acid polymers or copolymers, alkylvinyl polymers, hydroxyalkyl celluloses, carboxyalkyl celluloses, polysaccharides, dextrans, pectins, starches and derivatives, natural or synthetic gums, alginic acid.

In case of taste-masking formulations, the use of polyalcohols such as xylitol, maltitol and mannitol as hydrophilic compounds can also be advantageous.

The lipophilic matrix granules containing the active ingredient are mixed with the hydrophilic compounds cited above in a weight ratio typically ranging from 100:0.5 to 100:50 (lipophilic matrix: hydrophilic matrix). Part of the active ingredient can optionally be mixed with hydrophilic sub-

stances to provide compositions in which the active ingredient is dispersed both in the lipophilic and the hydrophilic matrix, said compositions being preferably in the form of tablets, capsules and/or minitabets.

The compression of the mixture of lipophilic and/or amphiphilic matrix, hydrogel-forming compound and, optionally, active ingredient not inglobated in the lipophilic matrix, yields a macroscopically homogeneous structure in all its volume, namely a matrix containing a dispersion of the lipophilic granules in a hydrophilic matrix. A similar result can also be obtained by coating the lipophilic matrix granules with a hydrophilic polymer coating.

The tablets obtainable according to the invention can optionally be subjected to known coating processes with a gastro-resistant film, consisting of, for example, methacrylic acids polymers (Eudragit^(R)) or cellulose derivatives, such as cellulose acetophthalate.

Active ingredients which can conveniently be formulated according to the invention comprise:

analgesics, such as acetaminophen, phenacetin, sodium salicylate; antitussives, such as dextromethorphan, codeine phosphate;

bronchodilators, such as albuterol, procaterol;

antipsychotics, such as haloperidol, chlorpromazine;

antihypertensives and coronary-dilators, such as isosorbide mono- and dinitrate, captopril;

selective β 2 antagonists such as salbutamol, terbutaline, ephedrine, orciprenaline sulfate;

calcium antagonists, such as nifedipine, nicardipine, diltiazem, verapamil;

antiparkinson drugs, such as pergolide, carbidopa, levodopa;

non steroid anti-inflammatory drugs, such as ketoprofen, ibuprofen, diclofenac, diflunisal, piroxicam, naproxen, ketorolac, nimesulide, thiaprophenic acid, mesalazine (5-aminosalicylic acid); antihistamines, such as terfenadine, loratadine;

antidiarrheals and intestinal antiinflammatories, such as loperamide, 5-aminosalicylic, olsalazine, sulfasalazine, budesonide;

spasmolytics such as octylonium bromide;

anxiolytics, such as chlordiazepoxide, oxazepam, medazepam, alprazolam, donazepam, lorazepam;

oral antidiabetics, such as glipizide, metformin, phenformin, gliclazide, glibenclamide;

cathartics, such as bisacodil, sodium picosulfate;

antiepileptics, such as valproate, carbamazepine, phenytoin, gabapentin;

antitumorals, such as flutamide, etoposide;

oral cavity disinfectants or antimicrobials, such as benzalkonium chloride, cetylpyridinium chloride or tibeonium iodide, and some amino derivatives such as benzydamine and chlorhexidine as well as the salts and derivatives thereof;

sodium fluoride.

The compositions of the invention can further contain conventional excipients, for example bioadhesive excipients such as chitosans, polyacrylamides, natural or synthetic gums, acrylic acid polymers.

The compositions of the invention can contain more than one active ingredient, each of them being optionally contained in the hydrophilic matrix or in the inert amphiphilic matrix, and are preferably in the form of tablets, capsules or minitabets.

In terms of dissolution characteristics, contact with water or aqueous fluids causes the immediate penetration of water inside the more superficial layer of the matrix which, thanks to the presence of the aqueous solvent, swells due to the

distension of the polymeric chains of the hydrogels, giving rise to a high viscosity hydrated front which prevents the further penetration of the solvent itself linearly slowing down the dissolution process to a well determined point which can be located at about half the thickness, until the further penetration of water would cause the disintegration of the hydrophilic layer and therefore the release of the content which, consisting of inert matrix granules, however induces the diffusion mechanism typical of these structures and therefore further slows down the dissolution profile of the active ingredient.

The presence of the amphiphilic matrix inside the lipophilic matrix inert allows to prevent any unevenness of the release profile of the active ingredient. The surfactants present in the amphiphilic portion promote wettability of the porous canaliculuses which cross the inert matrix preventing or reducing resistance to penetration of the solvent inside the inert matrix.

To obtain taste masking tablets, the components of the hydrophilic matrix are carefully selected to minimize the active substance release time through penetration accelerated by the canalization induced by the hydrophilic compound.

The following Examples illustrate the invention in greater detail.

EXAMPLE 1

500 g of 5-aminosalicylic-acid and 20 g of octylonium bromide are mixed with 10 g of soy lecithin dissolved in 50 g of a water:ethyl alcohol 1:3 mixture at about 50° C. After homogenization and drying, the granules of the resulting matrix are treated in a kneader with 20 g of carnauba wax and 50 g of stearic acid, heating until homogeneous dispersion, then cold-extruded into small granules. The inert matrix granules are loaded into a mixer in which 30 g of carbopol 971 P and 65 g of hydroxypropyl methylcellulose are sequentially added. After a first mixing step for homogeneously dispersing the powders, 60 g of microcrystalline cellulose and 5 g of magnesium stearate are added. After mixing, the final mixture is tableted to unitary weight of 760 mg/tablet. The resulting tablets are film-coated with cellulose acetophthalate or polymethacrylates and a plasticizer to provide gastric resistance and prevent the early release of product in the stomach.

The resulting tablets, when subjected to dissolution test in simulated enteric juice, have shown a release of the active principles having the following profile: after 60 minutes no more than 30%, after 180 minutes no more than 60%, after 5 hours no more than 80%.

EXAMPLE 2

50 g of diethylene glycol monoethyl ether are homogeneously distributed on 500 g of microcrystalline cellulose; then 100 g of Budesonide are added, mixing to complete homogenization. This mix is further added with 400 g of Budesonide, then dispersed in a blender containing 100 g of carnauba wax and 100 g of stearic acid preheated at a temperature of 60° C. After kneading for 5 minutes, the mixture is cooled to room temperature and extruded in granules of size below 1 mm.

A suitable mixer is loaded with the matrix granules prepared as above and the following amounts of hydrophilic excipients: 1500 g of hydroxypropyl methylcellulose and 500 g of polycarbofil.

The components are mixed until homogeneous dispersion of the matrices, then added with 2450 g of microcrystalline cellulose, 400 g of lactose, 100 g of colloidal silica and 50 g

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.