



Journal of Pharmacy and Pharmacology



The Pharmaceutical Society
of Great Britain

Volume 35 Number 1 January 1983

CODEN JPPMAB 35 (1) 1-64

ISSN 0022-3573

Page 1

Anacor Exhibit 2031
Flatwing Pharmaceuticals, Inc. v. Anacor Pharmaceuticals, Inc
IPR2018-00169

Journal of Pharmacy and Pharmacology

Published by The Pharmaceutical Society of Great Britain
1 Lambeth High Street, London SE1 5JN Telephone 01-735 9141 Telegrams Pharmakon London SE1

EDITOR

J. R. FOWLER

EDITORIAL BOARD

Advisory Editors

Professor A. M. BARRETT, The University of Leeds

Dr. B. A. CALLINGHAM, University of Cambridge

Professor P. H. ELWORTHY, The University of Manchester

Professor D. W. MATHIESON, University of Bradford

Professor M. J. H. SMITH, King's College Hospital Medical School, University of London

Other Board Members

Dr. L. C. BLABER, Roche Products Ltd., Welwyn Garden City

Professor M. R. W. BROWN, The University of Aston in Birmingham

Dr. G. P. CARR, British Pharmacopoeia Commission

Dr. J. H. COLLETT, The University of Manchester

D. H. DORKEN, Smith Kline & French Ltd., Welwyn Garden City

Professor A. T. FLORENCE, University of Strathclyde

Dr. J. D. PHILLIPSON, School of Pharmacy, University of London

Dr. N. PILPEL, Chelsea College, University of London

Professor R. L. SMITH, St. Mary's Hospital Medical School, University of London

Professor S. E. SMITH, St. Thomas' Hospital Medical School, University of London

Secretary

D. F. LEWIS

Copyright © 1983 *Journal of Pharmacy and Pharmacology*. The appearance on the code on the first page of an article in this Journal indicates the copyright owner's consent that copies of the article may be made for personal or internal use, or for the personal or internal use of specific clients. This consent is given on the condition, however, that the copier pay the per-copy fee of \$2.50 through the Copyright Clearance Center Inc., Operations Center, 21 Congress Street, Salem, Mass. 01970, USA, for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

Annual subscription for 1983: UK £51.00 (inc. postage); overseas £59.00 (air speeded).
Single copies £5.00 (overseas £6.00).

Claims for missing copies cannot be considered unless received within three months of publication.

Original Papers

- 1-6 S. MALAMATARIS, N. PILPEL
Tensile strength and compression of coated pharmaceutical powders: tablets
- 7-11 I. G. JOLLIFFE, J. M. NEWTON
The effect of dosator nozzle wall texture on capsule filling with the mG2 simulator
- 12-14 R. H. GUY, J. HADGRAFT, M. J. TAYLOR, I. W. KELLAWAY
Release of non-electrolytes from liposomes
- 15-18 L. LENNARD, J. L. MADDOCKS
Assay of 6-thioguanine nucleotide, a major metabolite of azothiaprime, 6-mercaptopurine and 6-thioguanine, in human red blood cells
- 19-22 K. YAMAOKA, T. NAKAGAWA, T. UNO
Moment analysis for disposition kinetics of several cephalosporin antibiotics in rats
- 23-27 P. LABRUDE, L. VIGNERON
Stability and functional properties of haemoglobin freeze-dried in the presence of four protective substances after prolonged storage: dose-effect relationships
- 28-33 K. A. WALTERS, G. L. FLYNN, J. R. MARVEL
Physicochemical characterization of the human nail: permeation pattern for water and the homologous alcohols and differences with respect to the stratum corneum
- 34-37 R. MATHISON
Actions of neurotransmitters and peptides on longitudinal and circular muscle of the rat portal vein
- 38-42 T. R. MACGREGOR, M. A. DRUM, S. E. HARRIGAN, J. N. WILEY, R. H. REUNING
Naltrexone metabolism and sustained release following administration of an insoluble complex to rhesus monkeys and guinea-pigs
- 43-44 R. C. ROWE
The orientation and alignment of particles in tablet film coatings
- 44-45 J. HARVEY, H. PARISH, P. P. K. HO, J. R. BOOT, W. DAWSON
The preferential inhibition of 5-lipoxygenase product formation by benoxaprofen
- 46-48 F. FRANCONI, S. MANZINI, I. STENDARDI, F. BENNARDINI, G. ANTONINI, P. FAILLI, R. MATUCCI, A. GIOTTI
Differential inhibitory effect of taurine on contractile responses to potassium and noradrenaline in rabbit ear artery
- 48-49 I. F. STAMFORD, M. A. CARROLL, A. CIVIER, C. N. HENSBY, A. BENNETT
Identification of arachidonate metabolites in normal, benign and malignant human mammary tissues
- 50-51 S. HARA, T. SATOH, H. KITAGAWA
Dose-dependence of the effect of hydralazine on the central nervous system in rats
- 52-53 J. R. FRY, C. G. WILSON
The effect of adrenalectomy on hepatic mixed function oxidase activity in female rats
- 54-56 R. C. SMALL, V. W. YONG
The failure of morphine to depress selectively non-adrenergic neural inhibition of the guinea-pig taenia caeci
- 57-58 R. W. FULLER, K. W. PERRY
Effect of pergolide on MOPEG sulphate levels in rat brain regions
- 59-61 A. IBRAHIM, P. COUVREUR, M. ROLAND, P. SPEISER
New magnetic drug carrier
- 62-64 R. IENTILE, A. DE SARRO, D. ROTIROTI, G. B. DE SARRO, G. NISTICO
Powerful stimulation of rat caudate nucleus adenylate cyclase activity by BW 245C, a prostaglandin analogue with prostacyclin-like activity

Communications

Physicochemical characterization of the human nail: permeation pattern for water and the homologous alcohols and differences with respect to the stratum corneum*

KENNETH A. WALTERS†, GORDON L. FLYNN*, AND JOHN R. MARVEL‡

College of Pharmacy, University of Michigan, Ann Arbor, Michigan 48109, U.S.A., and ‡ Dermatological Division, Ortho Pharmaceutical Corporation, Raritan, New Jersey, U.S.A.

In order to develop a basic concept of the permeability of the human nail plate and thus create a better understanding of the toxic potentials and therapeutic possibilities of substances applied to the nail, avulsed cadaver nails have been placed in specially constructed diffusion chambers and their permeation by water and the n-alkanols through dodecanol, all in high aqueous dilution, has been investigated. The permeability coefficient of water is 16.5×10^{-3} cm h⁻¹ and that for methanol is 5.6×10^{-3} cm h⁻¹. Ethanol's permeability coefficient measured 5.8×10^{-3} cm h⁻¹. Permeability coefficients decreased systematically thereafter to a low value of 0.27×10^{-3} cm h⁻¹ at n-octanol. The middle chain length alkanols, n-pentanol through n-octanol, have similar permeability coefficients but n-decanol and n-dodecanol show higher rates of permeation. The data suggest that, as a membrane, the hydrated human nail plate behaves like a hydrogel of high ionic strength to the polar and semipolar alcohols. Declining permeability rates appear linked to decreased partitioning into the complex matrix of the plate as the compounds become hydrophobic. The results for n-decanol and n-dodecanol introduce the possibility that a parallel lipid pathway exists which favours the permeation of these exceedingly hydrophobic species.

Apparently, no evidence exists concerning fundamental permeation mechanisms and possible influences of chemical structure on transport across the nail plate. To an extent its permeability properties have been inferred without foundation from the behaviour of other horny tissues. In order to make *a priori* judgements concerning toxic risk and therapeutic benefit of substances brought in contact with the nail, some baseline information on this tissue is needed.

We have shown it possible to determine nail plate permeability coefficients using standard diffusion cell techniques (Walters et al 1981). Results obtained for water agreed well with literature data on water transpiration through the nail plate (Burch & Winsor 1946; Spruit 1971; Baden et al 1973). In pursuant studies the techniques have been extended to the permeation of some n-alkanols. These are useful prototype compounds with systematically varying oil/water (o/w) distribution coefficients and diffusion

coefficients. Such structural influences on physicochemical properties, when considered together with relative permeabilities, have helped decipher the barrier mechanisms of several membranes (Blank 1964; Scheuplein 1965; Hwang et al 1976; Ho et al 1977; Behl et al 1980; Durrheim et al 1980; Flynn et al 1981). Previous studies of the alkanol's permeation of skin are especially notable as these provide evidence that the stratum corneum acts to some extent as a hydrophobic continuum (barrier) (Blank 1964; Scheuplein 1965; Behl et al 1980; Durrheim et al 1980; Flynn et al 1981). Similar studies on the human nail plate presented here are comparably revealing as, unlike the stratum corneum, the nail becomes less permeable to the n-alkanols as their hydrophobicity is increased. At extreme hydrophobicity there is increased permeability. The mechanistic significance of these general observations is considered.

MATERIALS AND METHODS

Materials

Tritiated water and radiolabelled alcohols were obtained from New England Nuclear ([³H]water, [³H]methanol, [¹⁴C]ethanol, [¹⁴C]butanol), California Bionuclear ([¹⁴C]propanol, [¹⁴C]pentanol,

* Correspondence. This work supported through the generosity of Ortho Pharmaceutical Corporation, Raritan, N.J., U.S.A.

† Present address, Fisons Limited, Pharmaceutical Division, Research & Development Laboratories, Bakewell Road, Loughborough, Leicestershire LE11 0QY, U.K.

[¹⁴C]heptanol, [¹⁴C]dodecanol) and ICN ([¹⁴C]hexanol, [¹⁴C]octanol, [¹⁴C]decanol). All radiolabelled compounds were diluted with saline (0.9% NaCl Irrigation Solution, Abbott Labs) before use. The alkanols were diluted to trace concentrations, 10⁻⁴ molar or less.

Permeation procedures

Details of the diffusion cell and permeation procedures have been given previously (Walters et al 1981). Briefly, trimmed human nail plate sections* were placed between two halves of a diffusion cell. A known amount of a radiolabelled permeant was placed in the donor chamber and samples were taken at predetermined intervals from the receptor chamber. Isotope activity was monitored using a Beckman LS 9000 liquid scintillation counter.

The permeation behaviours of [³H]water and [³H]methanol and [¹⁴C]alkanols in dilute solution were followed as a function of time at 37 °C. In all cases two permeants were applied with different radiolabels. Generally methanol was run as a tritiated compound along with a ¹⁴C-labelled co-permeant. Methanol thus served as a reference and it is important to note that the increased values for the permeability coefficient of decanol and dodecanol were obtained concurrently with normal methanol data.

Permeability coefficients (P) were calculated from:

$$P = \frac{V(dC/dt)}{A \cdot \Delta C} \quad (1)$$

where V is the volume of the receiver half cell, dC/dt is the rate of change in concentration in the pseudo-steady state portion of the receiver concentration versus time plot, A is the diffusional area and ΔC is the concentration differential of permeant across the membrane. V(dC/dt) gives the diffusional flux in mass per unit time. The diffusion cells with nail plate membranes in place were scrupulously checked for intercompartmental leakage using soluble but impenetrable polyethyleneglycol markers and no leaks were evident.

Diffusivities of the permeants in the nail plate tissue were calculated from the non-stationary state periods using:

$$D_{\text{eff}} = \frac{h^2}{6t_L} \quad (2)$$

Where D_{eff} is the effective diffusivity for a given

* Fresh cadaver nails generously supplied by Dr T. M. Oelrich, University of Michigan, School of Medicine.

compound, h is the nail plate thickness and t_L is the diffusional lag time obtained by linear regression of the steady state slope of uptake versus time plots. The nail plates used in these studies were measured with a micrometer and averaged 0.54 mm in thickness.

RESULTS AND DISCUSSION

Permeability coefficients of water and the saline diluted n-alkanols are given along with diffusion lag times in Table 1. Fig. 1 shows the relationship between the logarithms of the permeability coefficients and the alkyl chain lengths of the alcohols. An unusual pattern is observed with minimum permeability coefficient values at intermediate alkyl chain length.

Table 1. Nail plate permeability data for water and n-alkanols.

Permeant	Permeability ^a (cm h ⁻¹ × 10 ³)	Lag. time (t _L) (s)	Effective diffusion ^b constant (D _{eff}) cm ² s ⁻¹ × 10 ⁷
Water	16.5 ± 5.9 (6)	900 ± 100	5.4
Methanol	5.6 ± 1.2 (26)	1790 ± 200	2.7
Ethanol	5.8 ± 3.1 (8)	2730 ± 200	1.3
n-Propanol	0.83 ± 0.15 (4)	4020 ± 350	1.2
n-Butanol	0.61 ± 0.27 (4)	3470 ± 350	1.4
n-Pentanol	0.35 ± 0.07 (6)	2700 ± 250	1.8
n-Hexanol	0.36 ± 0.23 (5)	3540 ± 300	1.4
n-Heptanol	0.42 ± 0.12 (4)	2520 ± 300	1.9
n-Octanol	0.27 ± 0.03 (4)	2120 ± 150	2.2
n-Decanol	2.5 ± 1.7 (10)	2090 ± 150	2.1
n-Dodecanol	4.1 ± 2.7 (8)	2300 ± 150	2.1

a. Data include standard deviation and () number of experiments.

b. From t_L = $\frac{h^2}{6D}$ (Mean value for h = 0.54 mm)

Fig. 2 shows the effective diffusivities of the permeants in the nail plate tissue as a function of alkyl chain length.

Theoretical considerations

The nail plate's barrier properties are governed by its anatomical construction and its physicochemical properties and a proposed model must be supportable in terms of both. The model developed here, although speculative, fulfills these requirements. The plate consists of a laminate of sheets of keratinized cells (Caputo & Dadati 1968; Forslind 1970; Forslind & Thyresson 1975). Like the stratum

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.