



US005532241A

United States Patent [19][11] **Patent Number:** **5,532,241****Böttcher et al.**[45] **Date of Patent:** **Jul. 2, 1996**[54] **PIPERIDINES AND PIPERAZINES**

94/13659 6/1994 WIPO .


[75] Inventors: **Henning Böttcher**, Darmstadt;
Christoph Seyfried,
 Seeheim-Jugenheim; **Gerd Bartoszyk**;
Hartmut Greiner, both of Darmstadt,
 all of Germany

Primary Examiner—Emily Bernhardt
Attorney, Agent, or Firm—Millen, White, Zelano & Branigan

[73] Assignee: **Merck Patent Gesellschaft mit
 beschränkter Haftung**, Darmstadt,
 Germany

[57] **ABSTRACT**

Piperidine and piperazine derivatives of the formula I

[21] Appl. No.: **314,734**[22] Filed: **Sep. 29, 1994**[30] **Foreign Application Priority Data**

Sep. 30, 1993 [DE] Germany 43 33 254.4

[51] **Int. Cl.**⁶ **A61K 31/495**; A61K 31/445;
C07D 405/10[52] **U.S. Cl.** **514/254**; 544/373; 546/201;
514/323[58] **Field of Search** 544/373; 514/254[56] **References Cited****U.S. PATENT DOCUMENTS**

5,002,948 3/1991 Perregaard et al. 544/373
 5,242,925 9/1993 Böttcher et al. 514/254
 5,418,237 5/1995 Böttcher et al. 514/253

FOREIGN PATENT DOCUMENTS

0490772 6/1992 European Pat. Off. .
 4127849 2/1993 Germany .

wherein

Ind is an indol-3-yl radical which is unsubstituted or
 mono- or polysubstituted by OH, OA, CN, Hal, COR²
 or CH₂R²,

R¹ is benzofuran-5-yl or 2,3-dihydrobenzofuran-5-yl,
 chroman-6-yl, chroman-4-on-6-yl, 3-chromen-6-yl or
 chromen-4-on-6-yl, which is unsubstituted or mono-
 substituted by CN, CH₂OH, CH₂OA or COR²,

Q is C_mH_{2m},N or CR³,

A is alkyl having 1–6 C atoms,

Hal is F, Cl, Br or I,

R² is OH, OA, NH₂, NHA or NA₂,R³ is H, OH or OA and

m is 2, 3 or 4,

and their physiologically acceptable salts, are active on the
 central nervous system.


17 Claims, No Drawings

1

PIPERIDINES AND PIPERAZINES

SUMMARY OF THE INVENTION

The invention relates to novel piperidine and piperazine derivatives of the formula I



wherein

Ind is an indol-3-yl radical which is unsubstituted or mono- or polysubstituted by OH, OA, CN, Hal, COR² or CH₂R²,

R¹ is benzofuran-5-yl or 2,3-dihydrobenzofuran-5-yl, chroman-6-yl, chroman-4-on-6-yl, 3-chromen-6-yl or chromen-4-on-6-yl, which is unsubstituted or mono-substituted by CN, CH₂OH, CH₂OA or COR²,

Q is C_mH_{2m},

Z is N or CR³,

A is alkyl having 1-6 C atoms,

Hal is F, Cl, Br or I,

R² is OH, OA, NH₂, NHA or NA₂,

R³ is H, OH or OA and

m 2, 3 or 4,

and to their physiologically acceptable salts.

An object of the invention is to provide novel compounds capable of being used for the preparation of drugs.

Upon further study of the specification and appended claims, further objects and advantages of this invention will become apparent to those skilled in the art.

It has been found that the compounds of the formula I and their physiologically acceptable acid addition salts possess valuable pharmacological properties. Thus, in particular, they are active on the central nervous system, especially in terms of 5-HT_{1A}-agonist and 5-HT-reuptake inhibition. The compounds are furthermore active as serotonin agonists and antagonists. They inhibit the binding of tritiated serotonin ligands to hippocampal receptors (Cossery et al., European J. Pharmacol., 140:143-155 (1987)). They also modify the accumulation of DOPA in the corpus striatum and the accumulation of 5-HTP in the nuclei raphes (Seyfried et al., European J. Pharmacol., 160:31-41 (1989)). They also have analgesic and hypotensive effects; thus, in catheterized, conscious, spontaneously hypertensive rats (strain: SHR/Okamoto/NIH-MO-CHB-Kisslegg; method: q.v. Weeks and Jones, Proc. Soc. Exptl. Biol. Med., 104:646-648 (1960)), the directly measured blood pressure is lowered after oral administration of the compounds. They are also useful for prophylaxis and control of the sequelae of cerebral infarction (apoplexia cerebri) such as stroke and cerebral ischaemia.

Compounds of the formula I and their physiologically acceptable acid addition salts can, therefore, be used as active ingredients for anxiolytics, antidepressants, antipsychotics, neuroleptics, and/or antihypertensives, and also as intermediates for the preparation of other pharmaceutical active ingredients.

The invention relates to the piperidine and piperazine derivatives of the formula I and to their physiologically acceptable acid addition salts.

The radical A is alkyl having 1, 2, 3, 4, 5 or 6 C atoms, especially 1 or 2 C atoms, preferably methyl and also ethyl,

2

preferably methoxy and also ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or tert-butoxy. NHA is preferably methylamino and also ethylamino, isopropylamino, n-butylamino, isobutylamino, sec-butylamino or tert-butylamino. NA₂ is preferably dimethylamino and also N-ethyl-N-methylamino, diethylamino, di-n-propylamino, diisopropylamino or di-n-butylamino.

Analogously, CO-NHA is preferably N-methylcarbamoyl or N-ethylcarbamoyl; CO-NA₂ is preferably N,N-dimethylcarbamoyl or N,N-diethylcarbamoyl.

The radical Ind is an indol-3-yl radical which is unsubstituted or mono- or, for example, disubstituted by the radicals indicated. Preferably, it is substituted in the 5-position. Substitution in the 4-, 6- or 7-position is also suitable. Furthermore, substitution in the 1- or 2-position is possible. Preferred substituents on the indol-3-yl radical are OH, OA, CN, CONH₂, CH₂OH, but also CO₂H, F, Cl, Br, I, CH₂NH₂, CONHA or CONA₂, where A preferably corresponds to methyl or ethyl.

The radical R¹ is preferably benzofuran-5-yl, 2,3-dihydrobenzofuran-5-yl, chroman-6-yl or chromen-4-on-6-yl, which is unsubstituted or monosubstituted by -CH₂OH, -CONH₂, -CO₂A or -CO₂NHA.

Q is preferably -(CH₂)₄-, but also -(CH₂)₂- or -(CH₂)₃-, while Z is preferably -N-, -C(OH)- or -CH-.

Accordingly, the invention relates particularly to those compounds of the formula I in which at least one of said radicals has one of the meanings indicated above, especially one of the preferred meanings indicated above. Some preferred groups of compounds can be expressed by the following partial formulae Ia to Ig, which correspond to formula I and in which the radicals and parameters not described in greater detail are as defined for formula I, but in which:

in Ia, Ind is an indol-3-yl radical substituted in the 5-position by OH or OA;

in Ib, Ind is an indol-3-yl radical substituted in the 5-position by CONH₂ or by CN;

in Ic, Z is N and R¹ is substituted or unsubstituted benzofuran-5-yl;

in Id, Z is -C(OH)- and R¹ is substituted or unsubstituted benzofuran-5-yl;

in Ie, Z is N and R¹ is 2,3-dihydrobenzofuran-5-yl;

in If, Z is N and R¹ is chroman-6-yl;

in Ig, Z is N and R¹ is chromen-4-on-6-yl.

Especially preferred compounds are those of partial formulae Ih and Iah to Igh, which correspond to partial formulae I and Ia to Ig, but in which additionally: Q is -(CH₂)₄-.

The invention further relates to a process for the preparation of indole derivatives of the formula I and their salts, characterized in that a compound of the formula II




wherein

X¹ is X or NH₂,

X is Cl, Br, I, OH or an OH group functionally modified to form a reactive group, and

Ind and Q are as defined, is reacted with a compound of the formula III



3

X^2 and X^3 can be identical or different and are each X if $X^1 = \text{NH}_2$ or are together NH in other cases, and Z and R^1 are as defined, or in that to prepare a compound of the formula I in which Z is N, a compound of the formula IV



wherein

X, Q and Ind are as defined, is reacted with a compound of the formula V



wherein

R^1 is as defined, or in that a compound which has formula I except that one or more hydrogen atoms have been replaced by one or more reducible groups and/or one or more additional C—C and/or C—N bonds are treated with a reducing agent,

or in that a compound which has formula I except that one or more hydrogen atoms have been replaced by one or more solvolyzable groups is treated with a solvolyzing agent, and/or in that an OA group is optionally cleaved to form an OH group, and/or an Ind group and/or an Ar group is converted into another Ind and/or Ar group, and/or in that a resulting base or acid of the formula I is converted into one of its salts by treatment with an acid or base.

The compounds of the formula I are otherwise prepared by methods known per se, such as those described in the literature (e.g. in the standard works such as Houben-Weyl, Methoden der Organischen Chemie (Methods of Organic Chemistry), Georg-Thieme-Verlag, Stuttgart; Organic Reactions, John Wiley & Sons, Inc., New York; German Offenlegungsschrift 41 01 686), namely under reaction conditions such as those which are known and suitable for said reactions. It is also possible to make use of variants known per se, which are not mentioned in greater detail here.

If desired, the starting materials for the claimed process can also be formed in situ in such a way that they are not isolated from the reaction mixture but are immediately reacted further to give the compounds of the formula I.

In the compounds of the formula II, X^1 is preferably X; accordingly, in the compounds of the formula III, X^2 and X^3 are together preferably NH. The radical X is preferably Cl or Br, but it can also be I, OH or an OH group functionally modified to form a reactive group, especially alkylsulfonyloxy having 1–6 C atoms (e.g., methanesulfonyloxy) or arylsulfonyloxy having 6–10 C atoms (e.g., benzenesulfonyloxy, p-toluenesulfonyloxy, naphthalene-1- or -2-sulfonyloxy).

Accordingly, the indole derivatives of the formula I can be obtained especially by reacting compounds of the formula $\text{Ind}-\text{Q}-\text{Cl}$ or $\text{Ind}-\text{Q}-\text{Br}$ with piperidine/piperazine derivatives of the formula III in which X^2 and X^3 together are an NH group (designated as IIIa hereafter).

Some of the compounds of the formulae II and, in particular, III are known; the unknown compounds of the formulae II and III can easily be prepared analogously to the known compounds.

Primary alcohols of the formula $\text{Ind}-\text{Q}-\text{OH}$ can be obtained, e.g., by reducing the appropriate carboxylic acids or their esters. Treatment with thionyl chloride, hydrogen bromide, phosphorus tribromide or similar halogen compounds yields the corresponding halides of the formula $\text{Ind}-\text{Q}-\text{Hal}$. The corresponding sulfonyloxy compounds can be obtained from the alcohols $\text{Ind}-\text{Q}-\text{OH}$ by reaction

4

The iodine compounds of the formula $\text{Ind}-\text{Q}-\text{I}$ can be obtained, e.g., by reacting potassium iodide with the appropriate p-toluenesulfonic acid esters. The amines of the formula $\text{Ind}-\text{Q}-\text{N}_2$ can be prepared, e.g., from the halides with potassium phthalimide or by reducing the appropriate nitriles.

Most of the piperazine derivatives IIIa are known and can be obtained, e.g., by reacting bis(2-chloroethyl)amine or bis(2-chloroethyl)ammonium chloride with 5-aminobenzofuran, 2,3-dihydro-5-aminobenzofuran, 6-aminochroman or 6-aminochromen-4-one or an appropriately substituted derivative of the compounds mentioned. Compounds of the formula III (X^2 and $X^3 = \text{X}$ in each case) can be prepared, e.g., by reducing diesters of the formula $\text{alky} 100\text{C}-\text{CH}_2-\text{ZR}^1-\text{CH}_2-\text{COO}-\text{alkyl}$ to give compounds of the formula $\text{HO}-\text{CH}_2-\text{CH}_2-\text{ZR}^1-\text{CH}_2-\text{CH}_2\text{OH}$ (III, $X^2 = X^3 = \text{OH}$), this being followed, if desired, by reaction with SOCl_2 or PBr_3 .

The reaction of the compounds of formulae II and III proceeds according to methods such as those known from the literature for the alkylation of amines. The components can be melted together in the absence of a solvent, in a sealed tube or an autoclave if necessary. It is also possible, however, to react the compounds in the presence of an inert solvent. Examples of suitable solvents are hydrocarbons such as benzene, toluene or xylene; ketones such as acetone or butanone; alcohols such as methanol, ethanol, isopropanol or n-butanol; ethers such as tetrahydrofuran (THF) or dioxane; amides such as dimethylformamide (DMF) or N-methylpyrrolidone; or nitriles such as acetonitrile, or else, if desired, mixtures of these solvents with one another or mixtures with water. It can be favorable to add an acid-binding agent, for example an alkali metal or alkaline earth metal hydroxide, carbonate or bicarbonate or another alkali metal or alkaline earth metal salt of a weak acid, preferably a potassium, sodium or calcium salt, or to add an organic base such as triethylamine, dimethylaniline, pyridine or quinoline, or an excess of the amine component $\text{Ind}-\text{Q}-\text{NH}_2$ or of the piperidine or piperazine derivative of the formula IIIa. The reaction time is between about a few minutes and 14 days, depending on the conditions used, and the reaction temperature is preferably about 0° – 150° , normally 20° – 130° .

It is also possible to obtain a compound of the formula I by reacting a compound of the formula $\text{Ind}-\text{Q}-\text{N}(\text{CH}_2-\text{CH}_2-\text{X})_2$ (IV) with a compound of the formula R^1-NH_2 (V).

Most of the compounds of the formula V are known; the unknown compounds can easily be prepared analogously to the known compounds. For example, starting from the appropriately substituted nitro compounds, they can be converted into the amines of the formula V by reduction. The compounds of the formula IV can be prepared by reaction of $\text{Ind}-\text{Q}-\text{Cl}$, $\text{Ind}-\text{Q}-\text{Br}$ or $\text{Ind}-\text{Q}-\text{I}$ with secondary amines of the formula $\text{HN}(\text{CH}_2-\text{CH}_2-\text{X})_2$.

The reaction of compounds IV and V proceeds according to methods which are known from the literature and were given above for the alkylation of amines.


A compound of the formula I can also be obtained by treating a precursor, in which hydrogen atoms have been replaced by one or more reducible groups and/or one or more additional C—C and/or C—N bonds, with a reducing agent, preferably at temperatures of about -80 to 250° , in the presence of at least one inert solvent.

Reducible groups (groups replaceable by hydrogen) are, in particular, oxygen in a carbonyl group, hydroxyl, arylsulfonyloxy (e.g. p-toluenesulfonyloxy), N-benzenesulfo-

5

In principle, compounds containing only one of the above-mentioned groups or additional bonds, or compounds containing two or more of the above-mentioned groups or additional bonds adjacent to one another, can be converted into a compound of the formula I by reduction, it being possible simultaneously to reduce substituents in the Ind group which are present in the starting compound. This is preferably carried out using nascent hydrogen or complex metal hydrides or by means of a Wolff-Kishner reduction or the reductions with hydrogen gas under transition metal catalysis.

Preferred starting materials for the reduction have formula VI



wherein

Ind' is an Ind radical which can additionally be substituted in the 1-position by an arylsulfonyl group or an alkoxycarbonyl group,

L is Q or a chain which corresponds to the radical Q except that one or more $-\text{CH}_2-$ groups have been replaced by $-\text{CO}-$ and/or one or more hydrogen atoms have been replaced by one or more OH groups or a double bond, and

R^1 has the meaning given, but wherein the following meanings cannot apply simultaneously: $\text{Ind}=\text{Ind}$ and $\text{L}=\text{Q}$.

In the compounds of the formula VI, L is preferably $-\text{CO}-(\text{CH}_2)_{n-2}-\text{CO}-$, wherein n is 2, 3 or 4 [specifically $-\text{COCO}-$, $-\text{COCH}_2\text{CO}-$, $-\text{CO}-(\text{CH}_2)_2-\text{CO}-$, $-\text{CO}-(\text{CH}_2)_3-\text{CO}-$], $-(\text{CH}_2)_{n-1}-\text{CO}-$, wherein n is 2, 3 or 4 [specifically $-\text{CH}_2-\text{CO}-$, $-\text{CH}_2\text{CH}_2-\text{CO}-$, $-(\text{CH}_2)_3-\text{CO}-$ or $-(\text{CH}_2)_4-\text{CO}-$], further examples being $-\text{CO}-\text{CH}_2\text{CH}_2-$, $-\text{CO}-(\text{CH}_2)_3-$, $-\text{CH}_2-\text{CO}-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-\text{CO}-\text{CH}_2-$.

Compounds of the formula VI can be prepared, e.g., by reacting 4- R^1 -piperazine or 4- R^1 -piperidine with a compound of the formula VII



wherein

R^1 Ind', L and X^1 are as defined above, under the conditions indicated above for the reaction of II with III.

If nascent hydrogen is used as the reducing agent, this can be produced, e.g., by treating metals with weak acids or with bases. Thus, it is possible, e.g., to use a mixture of zinc with an alkali metal hydroxide solution or a mixture of iron with acetic acid. It is also appropriate to use sodium or another alkali metal dissolved in an alcohol such as ethanol, isopropanol, butanol, amyl or isoamyl alcohol or phenol. It is also possible to use an aluminum-nickel alloy in aqueous-alkaline solution, ethanol being added if necessary. Sodium amalgam or aluminum amalgam in aqueous-alcoholic or aqueous solution is also suitable for producing the nascent hydrogen. The reaction can also be carried out in the heterogeneous phase, in which case it is convenient to use an aqueous phase and a benzene or toluene phase.

Other reducing agents which can be used to particular advantage are complex metal hydrides such as LiAlH_4 , NaBH_4 , diisobutylaluminum hydride or $\text{NaAl}(\text{OCH}_2\text{CH}_2\text{OCH}_2)_2\text{H}_2$, and diborane. catalysts such as

6

are suitable for this purpose are, in particular, ethers such as diethyl ether, di-n-butyl ether, THF, dioxane, diglyme or 1,2-dimethoxyethane, and hydrocarbons such as benzene. Solvents which are suitable for a reduction with NaBH_4 are primarily alcohols such as methanol or ethanol, as well as water and aqueous alcohols. Reduction by these methods is preferably carried out at temperatures of about -80 to $+150^\circ$, especially about 0° - 100° .

The reduction of $-\text{CO}-$ groups in acid amides (e.g., those of the formula VI in which L is a $-(\text{CH}_2)_{n-1}-\text{CO}-$ group) to CH_2 groups can be carried out to particular advantage with LiAlH_4 in THF at temperatures of preferably about 0° - 66° . Arylsulfonyl protecting groups located in the 1-position of the indole ring can be simultaneously eliminated by reduction. N-Benzyl groups can be eliminated by reduction with sodium in liquid ammonia.

It is also possible to reduce one or more carbonyl groups to CH_2 groups according to the Wolff-Kishner method, e.g., by treatment with anhydrous hydrazine in absolute ethanol, under pressure, at temperatures of preferably about 150° - 250° . A sodium alcoholate is advantageously used as the catalyst. The reduction can also be varied according to the Huang-Minlon method by carrying out the reaction with hydrazine hydrate in a high-boiling water-miscible solvent such as diethylene glycol or triethylene glycol, in the presence of an alkali such as sodium hydroxide. The reaction mixture is normally boiled for about 3-4 hours. The water is then distilled off and the hydrazone formed is decomposed at temperatures of up to about 200° . The Wolff-Kishner reduction can also be carried out with hydrazine in dimethyl sulfoxide at room temperature.

Moreover, it is possible to carry out certain reductions by using H_2 gas under the catalytic action of transition metals, such as, e.g., Raney Ni or Pd. In this way, e.g., Cl, Br, I, SH or, in certain cases, even OH groups can be replaced by hydrogen. Nitro groups can also be converted into NH_2 groups by catalytic hydrogenation with Pd/H_2 in methanol.

Compounds which have formula I except that one or more H atoms have been replaced by one or more solvolyzable groups can be solvolyzed, especially hydrolyzed, to give the compounds of the formula I.

The starting materials for the solvolysis can be obtained for example by reacting IIIa with compounds which have formula II ($\text{X}^1=\text{X}$) except that one or more H atoms have been replaced by one or more solvolyzable groups. Thus, in particular, 1-acylindole derivatives (which have formula I except that, in the 1-position of the Ind radical, they contain an acyl group, preferably an alkoxycarbonyl, alkanoyl, alkylsulfonyl or arylsulfonyl group having up to 10 C atoms in each case, such as methanesulfonyl, benzenesulfonyl or p-toluenesulfonyl) can be hydrolyzed to give the corresponding indole derivatives unsubstituted in the 1-position of the indole ring, e.g. in an acidic or, preferably, neutral or alkaline medium at temperatures of preferably about 0° - 200° . Sodium, potassium or calcium hydroxide, sodium or potassium carbonate, or ammonia, is conveniently used as the base. The chosen solvents are preferably water; lower alcohols such as methanol or ethanol; ethers such as THF or dioxane; sulfones such as tetramethylene sulfone; or mixtures thereof, especially mixtures containing water. Hydrolysis can also be carried out simply by treatment with water alone, especially at the boiling point.

A compound of the formula I can furthermore be converted to another compound of the formula I by methods known per se.

Compounds of the formula I in which Ind is an indol- 3-yl

appropriate carboxyindol-3-yl compounds. It is possible, e.g., to esterify the acids with appropriate alcohols or alcoholates, using methods known per se. It is also possible to amidate acids or esters with primary or secondary amines. It is preferred to react the free carboxylic acid with the amine under the conditions of a peptide synthesis. This reaction is preferably carried out in the presence of a dehydrating agent, e.g., a carbodiimide such as dicyclohexylcarbodiimide or else N-(3-dimethylaminopropyl)-N-ethylcarbodiimide, or propanephosphonic anhydride (q.v. *Angew. Chem.* 92, 129 (1980)), diphenylphosphoryl azide or 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline, in an inert solvent, e.g., a halogenated hydrocarbon such as methylene chloride, an ether such as THF or dioxane, an amide such as DMF or dimethylacetamide, or a nitrile such as acetonitrile, at temperatures of preferably about -10 to 40, preferably about 0°-30°. Instead of the acid or amide, it is also possible to use reactive derivatives of these substances in the reaction, e.g., those in which reactive groups are blocked by protecting groups in an intermediate step. The acids can also be used in the form of their activated esters, which are conveniently formed in situ, e.g., by the addition of 1-hydroxybenzotriazole or N-hydroxysuccinimide.

Furthermore, cyano-substituted indol-3-yl radicals can be hydrolyzed to give carboxy-indol-3-yl or carbamido-indol-3-yl radicals.

Conversely, however, it is particularly convenient to prepare the nitriles by elimination of water, starting from the amides, e.g., by means of trichloroacetyl chloride/ Et_3N [*Synthesis* (2), 184, (1985)] or with POCl_3 (*J. Org. Chem.* 26, 1003 (1961)).

A base of the formula I can be converted with an acid into the corresponding acid addition salt. Acids which produce physiologically acceptable salts are suitable for this reaction. Thus, it is possible to use inorganic acids, e.g., sulfuric acid, hydrohalic acids such as hydrochloric acid or hydrobromic acid, phosphoric acids such as orthophosphoric acid, nitric acid and sulfamic acid, as well as organic acids, i.e., specifically aliphatic, alicyclic, araliphatic, aromatic or heterocyclic monobasic or polybasic carboxylic, sulfonic or sulfuric acids, such as formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, benzoic acid, salicylic acid, 2-phenylpropionic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methanesulfonic or ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenemonosulfonic acid and naphthalenedisulfonic acids and laurylsulfuric acid.

If desired, the free bases of the formula I can be liberated from their salts by treatment with strong bases such as sodium or potassium hydroxide or sodium or potassium carbonate provided there are no other acid groups in the molecule. In those cases where the compounds of the formula I have free acid groups, salt formation can also be achieved by treatment with bases. Suitable bases are alkali metal hydroxides, alkaline earth metal hydroxides or organic bases in the form of primary, secondary or tertiary amines.

The invention further relates to the use of the compounds of the formula I and their physiologically acceptable salts for the manufacture of pharmaceutical preparations, especially by a non-chemical route. For this purpose, they can be converted into a suitable dosage form together with at least one excipient or adjunct and, if appropriate, in combination with one or more additional active ingredients.

The invention further relates to compositions, especially

pound of the formula I and/or one of their physiologically acceptable salts. These preparations can be used as drugs in human or veterinary medicine. Possible excipients are organic or inorganic substances which are suitable for enteral (e.g., oral), parenteral or topical administration and which do not react with the novel compounds, examples of such excipients being water, vegetable oils, benzyl alcohols, polyethylene glycols, gelatin, carbohydrates such as lactose or starch, magnesium stearate, talc and petroleum jelly. Tablets, coated tablets, capsules, syrups, juices, drops or suppositories are used in particular for enteral administration, solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions or implants are used for parenteral administration, and ointments, creams or powders are used for topical administration. The novel compounds can also be lyophilized and the resulting lyophilizates used, e.g., to manufacture injectable preparations.

The preparations indicated can be sterilized and/or can contain adjuncts such as lubricants, preservatives, stabilizers and/or wetting agents, emulsifiers, salts for influencing the osmotic pressure, buffer substances, colorants, taste correctors and/or flavorings. If desired, they can also contain one or more additional active ingredients, e.g. one or more vitamins.

The compounds of the formula I and their physiologically acceptable salts can be used for the therapeutic treatment of the human or animal body and for controlling diseases. They can be used for treating disorders of the central nervous system, such as tension, depressions and/or psychoses, and side-effects in the treatment of hypertension (e.g., with α -methyl dopa). The compounds can also be used in endocrinology and gynecology, e.g., for the therapeutic treatment of acromegaly, hypogonadism, secondary amenorrhea, premenstrual syndrome and undesired puerperal lactation, and also for the prophylaxis and therapy of cerebral disorders (e.g., migraine), especially in geriatrics in a manner similar to certain ergot alkaloids and for controlling the sequelae of cerebral infarction (apoplexia cerebri), such as stroke and cerebral ischemia.

In these treatments, the substances of the invention are normally administered analogously to known, commercially available preparations (e.g., bromocriptine, dihydroergocorine), preferably in dosages of about 0.2-500 mg, especially 0.2-50 mg per dosage unit. The daily dosage is preferably about 0.001-10 mg/kg of body weight. The low dosages (about 0.2-1 mg per dosage unit; about 0.001-0.005 mg/kg of body weight) are particularly suitable for use as anti-migraine preparations; dosages of about 10-50 mg per dosage unit are preferred for the other indications. However, the particular dose for each individual patient depends on a very wide variety of factors, for example, the activity of the particular compound used, age, body weight, general state of health, sex, diet, time and method of administration, rate of excretion, drug combination and severity of the particular disease to which the therapy is applied. Oral administration is preferred.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius and unless otherwise indicated, all parts and percentages are by weight.

The entire disclosure of all applications, patents and

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.