Diagnosis and Treatment of Acne

STEPHEN TITUS, MD, and JOSHUA HODGE, MD, Fort Belvoir Community Hospital Family Medicine Residency, Fort Belvoir, Virginia

Acne is a chronic inflammatory skin disease that is the most common skin disorder in the United States. Therapy targets the four factors responsible for lesion formation: increased sebum production, hyperkeratinization, colonization by Propionibacterium acnes, and the resultant inflammatory reaction. Treatment goals include scar prevention, reduction of psychological morbidity, and resolution of lesions. Grading acne based on lesion type and severity

can help guide treatment. Topical retinoids are effective in treating inflammatory and noninflammatory lesions by preventing comedones, reducing existing comedones, and targeting inflammation. Benzoyl peroxide is an over-the-counter bactericidal agent that does not lead to bacterial resistance. Topical and oral antibiotics are effective as monotherapy, but are more effective when combined with topical retinoids. The addition of benzoyl peroxide to antibiotic therapy reduces the risk of bacterial resistance. Oral isotretinoin is approved for the treatment of severe recalcitrant acne and can be safely administered using the iPLEDGE program. After treatment goals are reached, maintenance therapy should be initiated. There is insufficient evidence to recommend the use of laser and light therapies. Referral to a dermatologist should be considered if treatment goals are not met. (Am Fam Physician. 2012;86(8):734-740. Copyright © 2012 American Academy of Family Physicians.)



cne is the most common skin disorder in the United States, affecting 40 to 50 million persons of all ages and races.1 Potential outcomes include physical scars, persistent hyperpigmentation, and psychological sequelae.

Pathogenesis

Acne is a chronic inflammatory disease involving the pilosebaceous unit. It is typified by the eruption of a comedo within the follicle, which is preceded by a microcomedo.1 Four main factors lead to the formation of acne lesions: (1) increased sebum production by sebaceous glands, in which androgens have an important role; (2) hyperkeratinization of the follicle, leading to a microcomedo that eventually enlarges into a comedo; (3) colonization of the follicle by the anaerobe Propionibacterium acnes; and (4) an inflammatory reaction.² The inflammatory events may begin before hyperkeratinization of the follicle.3 Current therapies target these four factors for acute control of flare-ups and long-term maintenance.



Evaluation

Acne is diagnosed by the identification of lesions. The spectrum of acne lesions ranges from noninflammatory open or closed comedones (blackheads and whiteheads; Figure 1) to inflammatory lesions, which may be papules, pustules, or nodules (Figures 2 through 4). Lesions are most likely to occur on the face, neck, chest, and back, where there is a higher concentration of sebaceous glands. Other conditions can mimic acne, and even include the term acne in their nomenclature, but they lack the presence of comedones. Table 1 outlines the differential diagnosis for acne.4 Grading acne based on the type of lesions and their severity can help in deciding which therapies are warranted (Figure 5); however, there is no consensus on the best grading system.5

Treatment

TOPICAL THERAPIES: PRESCRIPTION

Topical retinoids are versatile agents in the treatment of acne (Table 2).6,7 They prevent the formation and reduce the number

Downloaded from the American Family Physician Web site at www.aafp.org/afp. Copyright © 2012 American Academy of Family Physicians. For the private, noncommercial use of one individual user of the Web site. All other rights reserved. Contact copyrights@aafp.org for copyright questions and/or permission requests.






Figure 1. Noninflammatory acne lesions consisting of open and closed comedones.



Figure 2. Mild inflammatory acne lesions with comedones and few papules and pustules.



Topical antibiotics are used predominantly for the treatment of mild to moderate inflammatory or mixed acne. Clindamycin and erythromycin are the most studied (*Table 3*).^{2,5,7} They are sometimes used



Figure 3. Moderate inflammatory acne lesions with comedones, several papules and pustules, and few nodules.



Figure 4. Severe inflammatory acne lesions with comedones, several papules and pustules, multiple nodules, and scarring.

Table 1. Differential Diagnosis of Acne

Diagnosis	Distinguishing features
Bacterial folliculitis	Abrupt eruption; spreads with scratching or shaving; variable distribution
Drug-induced acne	Use of androgens, adrenocorticotropic hormone, bromides, corticosteroids, oral contraceptives, iodides, isoniazid, lithium, phenytoin (Dilantin)
Hidradenitis suppurativa	Double comedo; starts as a painful boil; sinus tracts
Miliaria	"Heat rash" in response to exertion or heat exposure; nonfollicular papules, pustules, and vesicles
Perioral dermatitis	Papules and pustules confined to the chin and nasolabia folds; clear zone around the vermilion border
Pseudofolliculitis barbae	Affects curly-haired persons who regularly shave closely
Rosacea	Erythema and telangiectasias; no comedones
Seborrheic dermatitis	Greasy scales and yellow-red coalescing macules or papules

October 15, 2012 * Volume 86, Number 8

www.aafp.org/afp

American Family Physician 735






Figure 5. Severity-based approach to treating acne.

as monotherapy, but are more effective in combination with topical retinoids.⁵ Because of the possibility that topical antibiotics may induce resistance, it is recommended that benzoyl peroxide be added to these regimens.²

Table 4 summarizes the additional topical therapies

that are available.^{5,8-11}Azelaic acid should be considered for use in pregnant women. The cream formulation (Azelex) is approved by the U.S Food and Drug Administration (FDA) for the treatment of acne vulgaris, but the gel (Finacea) has significantly better bioavailability.⁸ It has mixed antimicrobial and anticomedonal effects,

(Differin) dryness, pruritus, stinging Gel (0.1%, 0.3%) Adapalene/benzoyl peroxide (Epiduo) gel (0.1%/2.5%) Tazarotene X Local erythema, peeling, dryness, pruritus, stinging Cream, gel (0.05%, 0.1%) NA (\$240) Tretinoin C Local erythema, peeling, Cream (0.025%, 0.05%, 0.1%) \$27 (\$130)	Agent	FDA pregnancy category	Adverse effects	Available formulations	Estimated cost generic (brand)*
(Epiduo) gel (0.1%/2.5%) Tazarotene X Local erythema, peeling, dryness, pruritus, stinging Tretinoin C Local erythema, peeling, Cream (0.025%, 0.05%, 0.1%) \$27 (\$130)	Adapalene (Differin)	С			\$125 (\$363)
(Tazorac) dryness, pruritus, stinging Tretinoin C Local erythema, peeling, Cream (0.025%, 0.05%, 0.1%) \$27 (\$130)					NA (\$269)
		X		Cream, gel (0.05%, 0.1%)	NA (\$240)
(Retin-A) dryness, pruritus, stinging Gel (0.01%, 0.025%, 0.05%) \$24 (\$19 to \$105)	Tretinoin (Retin-A)	С	, , , , , , , , , , , , , , , , , , , ,	Cream (0.025%, 0.05%, 0.1%)	\$27 (\$130)
				Gel (0.01%, 0.025%, 0.05%)	\$24 (\$19 to \$105)
Microsphere gel (0.04%, 0.1%) NA (\$170)				Microsphere gel (0.04%, 0.1%)	NA (\$170)

736 American Family Physician

www.aafp.org/afp

Volume 86, Number 8 * October 15, 2012



Agent	FDA pregnancy category	Adverse effects	Available formulations	Estimated cost generic (brand)*
Clindamycin	В	Local erythema, peeling, dryness, pruritus, burning, oiliness	Foam, gel, lotion, solution (1.0%)	\$12 to \$96, depending on formulation (\$46 to \$213)
			Clindamycin/benzoyl peroxide (Benzaclin) gel (1%/5%, 1.2%/2.5%)	\$107 (\$210)
			Clindamycin/tretinoin gel (Veltin, Ziana; 1.2%/0.025%)	NA (\$180 Veltin, \$250 Ziana)
Erythromycin	В	Local erythema, peeling,	Gel, solution, ointment (2%)	\$25 (NA)
		dryness, pruritus, burning, oiliness	Erythromycin/benzoyl peroxide (Benzamycin) gel (3%/5%)	\$62 (\$313)

NOTE: Topical antibiotics are more effective when combined with a topical retinoid.

FDA = U.S. Food and Drug Administration; NA = not available.

and may be effective for the treatment of mild to moderate inflammatory or mixed acne.⁵

Dapsone is the first agent in a new class of topical acne medications to achieve FDA approval in the past 10 years. Although it is an antibiotic, it likely improves acne by inhibiting inflammation. In studies, dapsone was minimally more effective than placebo in reducing inflammatory and noninflammatory lesions, but it has never been compared with other topical agents. Unlike oral dapsone, there is no evidence that the

topical formulation causes hemolytic anemia or severe skin reactions.9

TOPICAL THERAPIES: OVER THE COUNTER

Benzoyl peroxide is an over-the-counter bactericidal agent that comes in a wide array of concentrations and formulations. No particular form has been proven better than another.⁵ Benzoyl peroxide is unique as an antimicrobial because it is not known to increase bacterial resistance.¹¹ It is most effective for the treatment of mild

Table 4. Selected Nonantibiotic	Topical Therapies for the	Treatment of Acne Vulgaris
---------------------------------	---------------------------	----------------------------

Agent	FDA pregnancy category	Adverse effects	Available formulations	Estimated cost generic (brand)*
Azelaic acid	В	Hypopigmentation, burning, stinging, tingling, pruritus	Cream (Azelex, 20%; approved for acne vulgaris) Gel (Finacea, 15%; approved for rosacea)	NA (\$210)
Benzoyl peroxide	С	Dry skin, local erythema	Bar, cream, gel, lotion, pad, wash (2.5% to 10%)	\$5 over the counter \$8 to \$36 prescription (NA)
Dapsone	С	Local oiliness, peeling, dryness, erythema	Gel (Aczone, 5%)	NA (\$193)
Salicylic acid	С	Dryness, mild skin irritation	Cream, dressing, foam, gel, liquid, lotion, ointment, pad, paste, shampoo, soap, solution, stick (0.5% to 3%)	\$5 over the counter

FDA = U.S. Food and Drug Administration; NA = not available.

Information from references 5, and 8 through 11.

October 15, 2012 * Volume 86, Number 8

www.aafp.org/afp

American Family Physician 737



^{*—}Estimated retail price of one month's treatment based on information obtained at http://www.lowestmed.com (accessed September 18, 2012). Information from references 2, 5, and 7.

^{*—}Estimated retail price of one month's treatment based on information obtained at http://www.lowestmed.com and http://www.drugstore.com (accessed September 18, 2012).

Table 5. Selected Oral Antibiotics for the Treatment of Acne Vulgaris

Agent	FDA pregnancy category	Adverse effects	Dosage	Estimated cost generic (brand)*
Doxycycline	D	Photosensitivity, pseudotumor cerebri, esophageal irritation	50 to 100 mg once or twice per day	\$15 (\$71 to \$363)
Erythromycin	В	Gastrointestinal upset	250 to 500 mg two to four times per day	\$73 to \$340 (NA)
Minocycline (Minocin)	D	Vestibular dysfunction, photophobia, hepatotoxicity, lupus-like reaction, pseudotumor cerebri	50 to 100 mg once or twice per day	\$21 to \$59 (\$173 to \$675)
Tetracycline	С	Gastrointestinal upset, photosensitivity, pseudotumor cerebri	250 to 500 mg once or twice per day	\$8 (NA)
Trimethoprim/ sulfamethoxazole (Bactrim, Septra)	С	Allergic reactions	160/800 mg twice per day	\$33 (\$194)

FDA = U.S. Food and Drug Administration; NA = not available.

to moderate mixed acne when used in combination with topical retinoids.² Benzoyl peroxide may also be added to regimens that include topical and oral antibiotics to decrease the risk of bacterial resistance.²

Salicylic acid is present in a variety of over-the-counter cleansing products. These products have anticomedonal properties and are less potent than topical retinoids, but there have been only limited high-quality studies examining their effectiveness.⁵

ORAL THERAPIES

Oral antibiotics are effective for the treatment of moderate to severe acne⁵ (*Table 5* ^{2,5,10,12}). The best-studied antibiotics include tetracycline and erythromycin. Based on expert consensus on relative effectiveness, the American Academy of Dermatology recommends using doxycycline and minocycline (Minocin) rather than tetracycline.⁵ Trimethoprim/sulfamethoxazole (Bactrim, Septra) and trimethoprim alone may be used if tetracycline or erythromycin cannot be tolerated. Because of the potential for

Topical dapsone is the first drug in a new class of acne therapy to receive approval in the past 10 years. bacterial resistance with the use of an oral antibiotic, it is recommended that benzoyl peroxide be added to any regimen of oral

antibiotics.² Tetracycline is preferred over erythromycin because of the higher rates of resistance associated with erythromycin.⁵

After individual treatment goals have been met, oral antibiotics can be discontinued and replaced with topical

retinoids for maintenance therapy.² Topical retinoids are sufficient to prevent relapses in most patients with acne vulgaris, especially if the disease was originally classified as mild or moderate. If the patient's acne was initially classified as severe inflammatory, benzoyl peroxide with or without an antibiotic can be added for maintenance therapy.²

Oral isotretinoin is FDA-approved for the treatment of severe recalcitrant acne. Evidence suggests that it is also useful for less severe acne that is treatment resistant.5 The usual dosage for severe treatment-resistant acne is 0.5 to 1.0 mg per kg per day for about 20 weeks, or a cumulative dose of 120 mg per kg.13 Initial flareups can be minimized with a beginning daily dosage of 0.5 mg or less per kg.5 Total cumulative doses of less than 120 mg increase relapse rates, and doses of more than 150 mg increase the incidence of adverse effects without producing greater benefits.¹³ Approximately 40 percent of patients achieve long-term remission with a 120-mg cumulative dose, 40 percent require retreatment with topical therapy or oral antibiotics, and 20 percent require retreatment with isotretinoin.14,15 Patients with moderate acne may respond to lower dosages (0.3 mg per kg per day) and experience fewer adverse effects.¹⁶

Physicians, distributors, pharmacies, and patients must register in the iPLEDGE program (http://www.ipledge program.com) before using isotretinoin. This program was established to prevent pregnancy in patients taking the medication. Isotretinoin is a potent teratogen and is associated with abnormalities of the face, eyes, ears, skull, central nervous system, cardiovascular system, thymus, and parathyroid glands. Negative pregnancy

738 American Family Physician

www.aafp.org/afp

Volume 86, Number 8 * October 15, 2012



^{*—}Estimated retail price of one month's treatment based on information obtained at http://www.lowestmed.com (accessed September 18, 2012). Information from references 2, 5, 10, and 12.

DOCKET

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

