



US 20050031546A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0031546 A1**

**Bartholomaus et al.**

(43) **Pub. Date: Feb. 10, 2005**

(54) **ABUSE-PROFFED DOSAGE FORM**

(30) **Foreign Application Priority Data**

Aug. 6, 2003 (DE)..... 103 36 400.5

(76) Inventors: **Johannes Bartholomaus**, Aachen (DE);  
**Heinrich Kugelmann**, Aachen (DE)

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 49/00**; A61K 9/14

(52) **U.S. Cl.** ..... **424/10.1**; 424/486

Correspondence Address:

**PERMAN & GREEN**  
**425 POST ROAD**  
**FAIRFIELD, CT 06824 (US)**

(57) **ABSTRACT**

The present invention relates to an abuse-proofed, thermoformed dosage form containing, in addition to one or more active ingredients with abuse potential optionally together with physiologically acceptable auxiliary substances, at least one synthetic or natural polymer with a breaking strength of at least 500 N and to a process for the production thereof.

(21) Appl. No.: **10/718,112**

(22) Filed: **Nov. 20, 2003**

**ABUSE-PROFFED DOSAGE FORM**

[0001] The present invention relates to an abuse-proofed, thermoformed dosage form containing, in addition to one or more active ingredients with abuse potential (A) optionally together with physiologically acceptable auxiliary substances (B), at least one synthetic or natural polymer (C) and optionally at least one wax (D), wherein component (C) exhibits a breaking strength of at least 500 N, and to a process for the production of the dosage form according to the invention.

[0002] Many pharmaceutical active ingredients, in addition to having excellent activity in their appropriate application, also have abuse potential, i.e. they can be used by an abuser to bring about effects other than those intended. Opiates, for example, which are highly active in combating severe to very severe pain, are frequently used by abusers to induce a state of narcosis or euphoria.

[0003] In order to make abuse possible, the corresponding dosage forms, such as tablets or capsules are comminuted, for example ground in a mortar, by the abuser, the active ingredient is extracted from the resultant powder using a preferably aqueous liquid and the resultant solution, optionally after being filtered through cotton wool or cellulose wadding, is administered parenterally, in particular intravenously. An additional phenomenon of this kind of administration, in comparison with abusive oral administration, is a further accelerated increase in active ingredient levels giving the abuser the desired effect, namely the "kick" or "rush". This kick is also obtained if the powdered dosage form is administered nasally, i.e. is sniffed. Since controlled-release dosage forms containing active ingredients with abuse potential do not give rise to the kick desired by the abuser when taken orally even in abusively high quantities, such dosage forms are also comminuted and extracted in order to be abused.

[0004] U.S. Pat. No. 4,070,494 proposed adding a swellable agent to the dosage form in order to prevent abuse. When water is added to extract the active ingredient, this agent swells and ensures that the filtrate separated from the gel contains only a small quantity of active ingredient.

[0005] The multilayer tablet disclosed in WO 95/20947 is based on a similar approach to preventing parenteral abuse, said tablet containing the active ingredient with abuse potential and at least one gel former, each in different layers.

[0006] WO 03/015531 A2 discloses another approach to preventing parenteral abuse. A dosage form containing an analgesic opioid and a dye as an aversive agent is described therein. The colour released by tampering with the dosage form is intended to discourage the abuser from using the dosage form which has been tampered with.

[0007] Another known option for complicating abuse involves adding antagonists to the active ingredients to the dosage form, for example naloxone or naltrexone in the case of opiates, or compounds which cause a physiological defence response, such as for example Radix ipecacuanha=ipecac root.

[0008] However, since in most cases of abuse it is still necessary to pulverise the dosage form comprising an active ingredient suitable for abuse, it was the object of the present invention to complicate or prevent the pulverisation preced-

ing abuse of the dosage form comprising the agents conventionally available for potential abuse and accordingly to provide a dosage form for active ingredients with abuse potential which ensures the desired therapeutic effect when correctly administered, but from which the active ingredients cannot be converted into a form suitable for abuse simply by pulverisation.

[0009] Said object has been achieved by the provision of the abuse-proofed, thermoformed dosage form according to the invention which contains, in addition to one or more active ingredients with abuse potential (A), at least one synthetic or natural polymer (C) and optionally at least one wax (D), wherein component (C) exhibits a breaking strength of at least 500 N.

[0010] The use of polymers having the stated minimum breaking strength, preferably in quantities such that the dosage form also exhibits such a minimum breaking strength, means that pulverisation of the dosage form is considerably more difficult using conventional means, so considerably complicating or preventing the subsequent abuse.

[0011] If comminution is inadequate, parenteral, in particular intravenous, administration cannot be performed safely or extraction of the active ingredient therefrom takes too long for the abuser or there is no "kick" when taken orally, as release is not spontaneous.

[0012] According to the invention, comminution is taken to mean pulverisation of the dosage form with conventional means which are available to an abuser, such as for example a mortar and pestle, a hammer, a mallet or other usual means for pulverisation by application of force.

[0013] The dosage form according to the invention is thus suitable for preventing parenteral, nasal and/or oral abuse of pharmaceutical active ingredients with abuse potential.

[0014] Pharmaceutical active ingredients with abuse potential are known to the person skilled in the art, as are the quantities thereof to be used and processes for the production thereof, and may be present in the dosage form according to the invention as such, in the form of the corresponding derivatives thereof, in particular esters or ethers, or in each case in the form of corresponding physiologically acceptable compounds, in particular in the form of the salts or solvates thereof, as racemates or stereoisomers. The dosage form according to the invention is also suitable for the administration of several active ingredients. It is preferably used to administer a specific active ingredient.

[0015] The dosage form according to the invention is in particular suitable for preventing abuse of a pharmaceutical active ingredient selected from the group consisting of opiates, opioids, tranquillisers, preferably benzodiazepines, barbiturates, stimulants and other narcotics.

[0016] The dosage form according to the invention is very particularly suitable for preventing abuse of an opiate, opioid, tranquilliser or another narcotic selected from the group consisting of N-{1-[2-(4-ethyl-5-oxo-2-tetrazolin-1-yl)ethyl]-4-methoxymethyl-4-piperidyl}propionanilide (alfentanil), 5,5-diallylbarbituric acid (allobarbital), allylprodine, alphaprodine, 8-chloro-1-methyl-6-phenyl-4H-[1,2,4]triazolo[4,3-a][1,4]-benzodiazepine (alprazolam), 2-diethylaminopropiophenone (amfepramone), ( $\pm$ )- $\alpha$ -methyl-

phenethylamine (amphetamine), 2- $\alpha$ -methylphenethylamino)-2-phenylacetone nitrile (amphetaminil), 5-ethyl-5-isopentylbarbituric acid (amobarbital), anileridine, apocodeine, 5,5-diethylbarbituric acid (barbital), benzylmorphine, bezitramide, 7-bromo-5-(2-pyridyl)-1H-1,4-benzodiazepine-2(3H)-one (bromazepam), 2-bromo-4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo-[4,3-a][1,4]diazepine (brotizolam), 17-cyclopropylmethyl-4,5 $\alpha$ -epoxy-7 $\alpha$ [(S)-1-hydroxy-1,2,2-trimethyl-propyl]-6-methoxy-6,14-endo-ethanomorphinan-3-ol (buprenorphine), 5-butyl-5-ethylbarbituric acid (butobarbital), butorphanol, (7-chloro-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepine-3-yl)-dimethylcarbamate (camazepam), (1S,2S)-2-amino-1-phenyl-1-propanol (cathine/D-norpseudoephedrine), 7-chloro-N-methyl-5-phenyl-3H-1,4-benzodiazepine-2-ylamine-4-oxide (chlorodiazepoxide), 7-chloro-1-methyl-5-phenyl-1H-1,5-benzodiazepine-2,4(3H,5H)-dione (clobazam), 5-(2-chlorophenyl)-7-nitro-1H-1,4-benzodiazepine-2(3H)-one (clonazepam), clonitazene, 7-chloro-2,3-dihydro-2-oxo-5-phenyl-1H-1,4-benzodiazepine-3-carboxylic acid (clorazepate), 5-(2-chlorophenyl)-7-ethyl-1-methyl-1H-thieno[2,3-e][1,4]diazepine-2(3H)-one (clotiazepam), 10-chloro-11b-(2-chlorophenyl)-2,3,7,11b-tetrahydrooxazol[3,2-d][1,4]benzodiazepine-6(5H)-one (cloxazolam), (-)-methyl-3 $\beta$ -benzoyloxy-2 $\beta$ (1 $\alpha$ (H,5 $\alpha$ H)-tropancarboxylate] (cocaine), 4,5 $\alpha$ -epoxy-3-methoxy-17-methyl-7-morphinene-6 $\alpha$ -ol (codeine), 5-(1-cyclohexenyl)-5-ethylbarbituric acid (cyclobarbital), cyclorphan, cyprenorphine, 7-chloro-5-(2-chlorophenyl)-1H-1,4-benzodiazepine-2(3H)-one (delorazepam), desomorphine, dextromoramide, (+)-(1-benzyl-3-dimethylamino-2-methyl-1-phenylpropyl)propionate (dextropropoxyphen), dezocine, diampromide, diamorphine, 7-chloro-1-methyl-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (diazepam), 4,5 $\alpha$ -epoxy-3-methoxy-17-methyl-6 $\alpha$ -morphinanol (dihydrocodeine), 4,5 $\alpha$ -epoxy-17-methyl-3,6 $\alpha$ -morphinandiol (dihydromorphine), dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, (6aR,10aR)-6,6,9-trimethyl-3-pentyl-6a,7,8,10a-tetrahydro-6H-benzo[c]chromene-1-ol (dronabinol), eptazocine, 8-chloro-6-phenyl-4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine (estazolam), ethoheptazine, ethylmethylthiambutene, ethyl [7-chloro-5-(2-fluorophenyl)-2,3-dihydro-2-oxo-1H-1,4-benzodiazepine-3-carboxylate](ethyl loflazepate), 4,5 $\alpha$ -epoxy-3-ethoxy-17-methyl-7-morphinene-6 $\alpha$ -ol (ethylmorphine), etonitazene, 4,5 $\alpha$ -epoxy-7 $\alpha$ -(1-hydroxy-1-methylbutyl)-6-methoxy-17-methyl-6,14-endo-etheno-morphinan-3-ol (etorphine), N-ethyl-3-phenyl-8,9,10-trinorboman-2-ylamine (fencamfamine), 7-[2-(1-methyl-phenethylamino)ethyl]-theophylline (fenethylline), 3-( $\alpha$ -methylphenethylamino)propionitrile (fenproporex), N-(1-phenethyl-4-piperidyl)propionanilide (fentanyl), 7-chloro-5-(2-fluorophenyl)-1-methyl-1H-1,4-benzodiazepine-2(3H)-one (fludiazepam), 5-(2-fluorophenyl)-1-methyl-7-nitro-1H-1,4-benzodiazepine-2(3H)-one (flunitrazepam), 7-chloro-1-(2-diethylaminoethyl)-5-(2-fluorophenyl)-1H-1,4-benzodiazepine-2(3H)-one (flurazepam), 7-chloro-5-phenyl-1-(2,2,2-trifluoroethyl)-1H-1,4-benzodiazepine-2(3H)-one (halazepam), 10-bromo-11b-(2-fluorophenyl)-2,3,7,11b-tetrahydro[1,3]oxazolyl[3,2-d][1,4]benzodiazepine-6(5H)-one (haloxazolam), heroin, 4,5 $\alpha$ -epoxy-3-methoxy-17-methyl-6-morphinanone (hydrocodone), 4,5 $\alpha$ -epoxy-3-hydroxy-17-methyl-6-morphinanone (hydromorphone), hydroxypethi-

dine, isomethadone, hydroxymethyl morphinanone, 11-chloro-8,12b-dihydro-2,8-dimethyl-12b-phenyl-4H-[1,3]oxazino[3,2-d][1,4]benzodiazepine-4,7(6H)-dione (ketazolam), 1-[4-(3-hydroxyphenyl)-1-methyl-4-piperidyl]-1-propanone (ketobemidone), (3S,6S)-6-dimethylamino-4,4-diphenylheptan-3-yl acetate (levacetylmethadol (LAAM)), (-)-6-dimethyl-amino-4,4-diphenol-3-heptanone (levomethadone), (-)-17-methyl-3-morphinanol (levorphanol), levophenacylmorphane, lofentanil, 6-(2-chlorophenyl)-2-(4-methyl-1-piperazinylmethylene)-8-nitro-2H-imidazo[1,2-a][1,4]benzodiazepine-1(4H)-one (loprazolam), 7-chloro-5-(2-chlorophenyl)-3-hydroxy-1H-1,4-benzodiazepine-2(3H)-one (lorazepam), 7-chloro-5-(2-chlorophenyl)-3-hydroxy-1-methyl-1H-1,4-benzodiazepine-2(3H)-one (lormetazepam), 5-(4-chlorophenyl)-2,5-dihydro-3H-imidazo[2,1-a]isoindol-5-ol (mazindol), 7-chloro-2,3-dihydro-1-methyl-5-phenyl-1H-1,4-benzodiazepine (medazepam), N-(3-chloropropyl)- $\alpha$ -methylphenethylamine (mefenorex), meperidine, 2-methyl-2-propyltrimethylene dicarbamate (meprobamate), meptazinol, metazocine, methylmorphine, N, $\alpha$ -dimethylphenethylamine (methamphetamine), (O)-6-dimethylamino-4,4-diphenyl-3-heptanone (methadone), 2-methyl-3-o-tolyl-4(3H)-quinazolinone (methaqualone), methyl [2-phenyl-2-(2-piperidyl)acetate] (methylphenidate), 5-ethyl-1-methyl-5-phenylbarbituric acid (methylphenobarbital), 3,3-diethyl-5-methyl-2,4-piperidinedione (methpyrrolon), metopon, 8-chloro-6-(2-fluorophenyl)-1-methyl-4H-imidazo[1,5-a][1,4]benzodiazepine (midazolam), 2-(benzhydrylsulfinyl)-acetamide (modafinil), 4,5 $\alpha$ -epoxy-17-methyl-7-morphinene-3,6 $\alpha$ -diol (morphine), myrophine, ( $\pm$ )-trans-3-(1,1-dimethylheptyl)-7,8,10,10 $\alpha$ -tetrahydro-1-hydroxy-6,6-dimethyl-6H-dibenzo[b,d]pyrane-9 (6 $\alpha$ H)-one (nabilone), nalbuphine, nalorphine, narceine, nicomorphine, 1-methyl-7-nitro-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (nimetazepam), 7-nitro-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (nitrazepam), 7-chloro-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (nordazepam), norlevorphanol, 6-dimethylamino-4,4-diphenyl-3-hexanone (normethadone), normorphine, norpipanone, the exudation of plants belonging to the species *Papaver somniferum* (opium), 7-chloro-3-hydroxy-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (oxazepam), (cis-trans)-10-chloro-2,3,7,11b-tetrahydro-2-methyl-11b-phenyloxazol[3,2-d][1,4]benzodiazepine-6(5H)-one (oxazolam), 4,5 $\alpha$ -epoxy-14-hydroxy-3-methoxy-17-methyl-6-morphinanone (oxycodone), oxymorphone, plants and parts of plants belonging to the species *Papaver somniferum* (including the subspecies *setigerum*), papaveretum, 2-imino-5-phenyl-4-oxazolidinone (pernoline), 1,2,3,4,5,6-hexahydro-6,11-dimethyl-3-(3-methyl-2-butenyl)-2,6-methano-3-benzazocin-8-ol (pentazocine), 5-ethyl-5-(1-methylbutyl)-barbituric acid (pentobarbital), ethyl-(1-methyl-4-phenyl-4-piperidine carboxylate) (pethidine), phenadoxone, phenomorphan, phenazocine, phenoperidine, pimindine, pholcodine, 3-methyl-2-phenylmorpholine (phenmetrazine), 5-ethyl-5-phenylbarbituric acid (phenobarbital),  $\alpha,\alpha$ -dimethylphenethylamine (phentermine), 7-chloro-5-phenyl-1-(2-propynyl)-1H-1,4-benzodiazepine-2(3H)-one (pinazepam),  $\alpha$ -(2-piperidyl)benzhydryl alcohol (pipradrol), 1'-(3-cyano-3,3-diphenylpropyl)[1,4'-bipiperidine]-4'-carboxamide (piritramide), 7-chloro-1-(cyclopropylmethyl)-5-phenyl-1H-1,4-benzodiazepine-2(3H)-one (prazepam), profadol, proheptazine, promedol, propripidine, propoxyphene, N-(1-methyl-2-piperidinoethyl)-N-(2-pyridyl)propionamide, methyl {3-[4-methoxycar-

bonyl-4-(N-phenylpropanamido)piperidino]propanoate} (remifentanyl), 5-sec-butyl-5-ethylbarbituric acid (secbutabarbital), 5-allyl-5-(1-methylbutyl)-barbituric acid (secobarbital), N-{4-methoxymethyl-1-[2-(2-thienyl)ethyl]-4-piperidyl}-propionanilide (sufentanyl), 7-chloro-2-hydroxymethyl-5-phenyl-1H-1,4-benzodiazepin-2(3H)-one (temazepam), 7-chloro-5-(1-cyclohexenyl)-1-methyl-1H-1,4-benzodiazepine-2(3H)-one (tetrazepam), ethyl(2-dimethylamino-1-phenyl-3-cyclohexene-1-carboxylate) (tilidine (cis and trans)), tramadol, 8-chloro-6-(2-chlorophenyl)-1-methyl-4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine (triazolam), 5-(1-methylbutyl)-5-vinylbarbituric acid (vinylbital), (1R\*,2R\*)-3-(3-dimethylamino-1-ethyl-2-methylpropyl)-phenol, (1R,2R,4S)-2-(dimethylamino)methyl-4-(p-fluorobenzyloxy)-1-(m-methoxyphenyl)cyclohexanol, (1R,2R)-3-(2-dimethylaminomethyl-cyclohexyl)phenol, (1S,2S)-3-(3-dimethylamino-1-ethyl-2-methylpropyl)phenol, (2R,3R)-1-dimethylamino-3-(3-methoxyphenyl)-2-methylpentan-3-ol, (1RS,3RS,6RS)-6-dimethylaminomethyl-1-(3-methoxyphenyl)-cyclohexane-1,3-diol, 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)phenyl 2-(4-isobutoxyphenyl)-propionate, 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)phenyl 2-(6-methoxy-naphthalen-2-yl)-propionate, 3-(2-dimethylamino-methyl-cyclohex-1-enyl)-phenyl 2-(4-isobutyl-phenyl)-propionate, 3-(2-dimethylaminomethyl-cyclohex-1-enyl)-phenyl 2-(6-methoxy-naphthalen-2-yl)-propionate, (RR—SS)-2-acetoxy-4-trifluoromethyl-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-2-hydroxy-4-trifluoromethyl-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-4-chloro-2-hydroxy-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-2-hydroxy-4-methyl-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-2-hydroxy-4-methoxy-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-2-hydroxy-5-nitro-benzoic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester, (RR—SS)-2,1,4'-difluoro-3-hydroxy-biphenyl-4-carboxylic acid 3-(2-dimethylaminomethyl-1-hydroxy-cyclohexyl)-phenyl ester and for corresponding stereoisomeric compounds, the corresponding derivatives thereof in each case, in particular esters or ethers, and the physiologically acceptable compounds thereof in each case, in particular the salts and solvates thereof.

[0017] The compounds (1R\*,2R\*)-3-(3-dimethylamino-1-ethyl-2-methylpropyl)-phenol, (1R,2R,4S)-2-(dimethylamino)methyl-4-(p-fluorobenzyloxy)-1-(m-methoxyphenyl)cyclohexanol or the stereoisomeric compounds thereof or the physiologically acceptable compounds thereof, in particular the hydrochlorides thereof, the derivatives thereof, such as esters or ethers, and processes for the production thereof are known, for example, from EP-A-693475 or EP-A-780369. The corresponding descriptions are hereby introduced as a reference and are deemed to be part of the disclosure.

[0018] In order to achieve the necessary breaking strength of the dosage form according to the invention, at least one synthetic or natural polymer (C) is used which has a breaking strength, measured using the method disclosed in the present application, of at least 500 N. At least one polymer selected from the group consisting of polymethylene oxide, polyethylene oxide, polypropylene oxide, polyethylene,

polypropylene, polyvinyl chloride, polycarbonate, polystyrene, polyacrylate, copolymers thereof, and mixtures of at least two of the stated polymers is preferably used for this purpose. The polymers are distinguished by a molecular weight of at least 0.5 million, determined by rheological measurements. Thermoplastic polyalkylene oxides, such as polyethylene oxides, with a molecular weight of at least 0.5 million, preferably of up to 15 million, determined by Theological measurements, are very particularly preferred. These polymers have a viscosity at 25° C. of 4500 to 17600 cP, measured on a 5 wt. % aqueous solution using a model RVF Brookfield viscosimeter (spindle no. 2/rotational speed 2 rpm), of 400 to 4000 cP, measured on a 2 wt. % aqueous solution using the stated viscosimeter (spindle no. 1 or 3/rotational speed 10 rpm) or of 1650 to 10000 cP, measured on a 1 wt. % aqueous solution using the stated viscosimeter (spindle no. 2/rotational speed 2 rpm).

[0019] The polymers are used in powder form.

[0020] In order to achieve the necessary breaking strength of the dosage form according to the invention, it is furthermore possible additionally to use at least one natural or synthetic wax (D) with a breaking strength, measured using the method disclosed in the present application, of at least 500 N. Waxes with a softening point of at least 60° C. are preferred. Carnauba wax and beeswax are particularly preferred. Carnauba wax is very particularly preferred. Carnauba wax is a natural wax which is obtained from the leaves of the carnauba palm and has a softening point of >80° C. When the wax component is additionally used, it is used together with at least one polymer (C) in quantities such that the dosage form has a breaking strength of at least 500 N.

[0021] The dosage forms according to the invention are distinguished in that, due their hardness, they cannot be pulverised, for example by grinding in a mortar. This virtually rules out oral or parenteral, in particular intravenous or nasal abuse. However, in order to prevent any possible abuse in the event of comminution and/or pulverisation of the dosage form according to the invention which has nevertheless been achieved by application of extreme force, the dosage forms according to the invention may, in a preferred embodiment, contain further agents which complicate or prevent abuse as auxiliary substances (B).

[0022] The abuse-proofed dosage form according to the invention, which comprises, apart from one or more active ingredients with abuse potential, at least one hardening polymer (C) and optionally at least one wax (D), may accordingly also comprise at least one of the following components (a)-(f) as auxiliary substances (B):

[0023] (a) at least one substance which irritates the nasal passages and/or pharynx,

[0024] (b) at least one viscosity-increasing agent, which, with the assistance of a necessary minimum quantity of an aqueous liquid, forms a gel with the extract obtained from the dosage form, which gel preferably remains visually distinguishable when introduced into a further quantity of an aqueous liquid,

[0025] (c) at least one antagonist for each of the active ingredients with abuse potential,

[0026] (d) at least one emetic,

[0027] (e) at least one dye as an aversive agent,

[0028] (f) at least one bitter substance.

[0029] Components (a) to (f) are additionally each individually suitable for abuse-proofing the dosage form according to the invention. Accordingly, component (a) is preferably suitable for proofing the dosage form against nasal, oral and/or parenteral, preferably intravenous, abuse, component (b) is preferably suitable for proofing against parenteral, particularly preferably intravenous and/or nasal abuse, component (c) is preferably suitable for proofing against nasal and/or parenteral, particularly preferably intravenous, abuse, component (d) is preferably suitable for proofing against parenteral, particularly preferably intravenous, and/or oral and/or nasal abuse, component (e) is suitable as a visual deterrent against oral or parenteral abuse and component (f) is suitable for proofing against oral or nasal abuse. Combined use according to the invention of at least one of the above-stated components makes it possible still more effectively to prevent abuse of dosage forms according to the invention.

[0030] In one embodiment, the dosage form according to the invention may also comprise two or more of components (a)-(f) in a combination, preferably (a), (b) and optionally (c) and/or (f) and/or (e) or (a), (b) and optionally (d) and/or (f) and/or (e).

[0031] In another embodiment, the dosage form according to the invention may comprise all of components (a)-(f).

[0032] If the dosage form according to the invention comprises component (a) to counter abuse, substances which irritate the nasal passages and/or pharynx which may be considered according to the invention are any substances which, when administered via the nasal passages and/or pharynx, bring about a physical reaction which is either so unpleasant for the abuser that he/she does not wish to or cannot continue administration, for example burning, or physiologically counteracts taking of the corresponding active ingredient, for example due to increased nasal secretion or sneezing. These substances which conventionally irritate the nasal passages and/or pharynx may also bring about a very unpleasant sensation or even unbearable pain when administered parenterally, in particular intravenously, such that the abuser does not wish to or cannot continue taking the substance.

[0033] Particularly suitable substances which irritate the nasal passages and/or pharynx are those which cause burning, itching, an urge to sneeze, increased formation of secretions or a combination of at least two of these stimuli. Appropriate substances and the quantities thereof which are conventionally to be used are known per se to the skilled person or may be identified by simple preliminary testing.

[0034] The substance which irritates the nasal passages and/or pharynx of component (a) is preferably based on one or more constituents or one or more plant parts of at least one hot substance drug.

[0035] Corresponding hot substance drugs are known per se to the person skilled in the art and are described, for example, in "Pharmazeutische Biologie—Drogen und ihre Inhaltsstoffe" by Prof. Dr. Hildebert Wagner, 2nd., revised edition, Gustav Fischer Verlag, Stuttgart-New York, 1982,

pages 82 et seq. The corresponding description is hereby introduced as a reference and is deemed to be part of the disclosure.

[0036] The dosage form according to the invention may preferably contain the plant parts of the corresponding hot substance drugs in a quantity of 0.01 to 30 wt. %, particularly preferably of 0.1 to 0.5 wt. %, in each case relative to the total weight dosage unit.

[0037] If one or more constituents of corresponding hot substance drugs are used, the quantity thereof in a dosage unit according to the invention preferably amounts to 0.001 to 0.005 wt. %, relative to the total weight of the dosage unit.

[0038] A dosage unit is taken to mean a separate or separable administration unit, such as for example a tablet or a capsule.

[0039] One or more constituents of at least one hot substance drug selected from the group consisting of *Allii sativi* bulbus (garlic), *Asari rhizoma cum herba* (*Asarum* root and leaves), *Calami rhizoma* (*calamus* root), *Capsici fructus* (*capsicum*), *Capsici fructus acer* (cayenne pepper), *Curcuma longae rhizoma* (turmeric root), *Curcuma xanthorrhizae rhizoma* (Javanese turmeric root), *Galangae rhizoma* (galangal root), *Myristicae semen* (nutmeg), *Piperis nigri fructus* (pepper), *Sinapis albae semen* (white mustard seed), *Sinapis nigri semen* (black mustard seed), *Zedoariae rhizoma* (zedoary root) and *Zingiberis rhizoma* (ginger root), particularly preferably from the group consisting of *Capsici fructus* (*capsicum*), *Capsici fructus acer* (cayenne pepper) and *Piperis nigri fructus* (pepper) may preferably be added as component (a) to the dosage form according to the invention.

[0040] The constituents of the hot substance drugs preferably comprise o-methoxy(methyl)phenol compounds, acid amide compounds, mustard oils or sulfide compounds or compounds derived therefrom.

[0041] Particularly preferably, at least one constituent of the hot substance drugs is selected from the group consisting of myristicin, elemicin, isoeugenol,  $\beta$ -asarone, safrole, gingerols, xanthorrhizol, capsaicinoids, preferably capsaicin, capsaicin derivatives, such as N-vanillyl-9E-octadecanamide, dihydrocapsaicin, nordihydrocapsaicin, homocapsaicin, norcapsaicin and nomorcapsaicin, piperine, preferably trans-piperine, glucosinolates, preferably based on non-volatile mustard oils, particularly preferably based on p-hydroxybenzyl mustard oil, methylmercapto mustard oil or methylsulfonyl mustard oil, and compounds derived from these constituents.

[0042] Another option for preventing abuse of the dosage form according to the invention consists in adding at least one viscosity-increasing agent as a further abuse-preventing component (b) to the dosage form, which, with the assistance of a necessary minimum quantity of an aqueous liquid, forms a gel with the extract obtained from the dosage form, which gel is virtually impossible to administer safely and preferably remains visually distinguishable when introduced into a further quantity of an aqueous liquid.

[0043] For the purposes of the present invention visually distinguishable means that the active ingredient-containing gel formed with the assistance of a necessary minimum quantity of aqueous liquid, when introduced, preferably with

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.