

Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

**0 235 796
A2**

EUROPEAN PATENT APPLICATION

Application number: 87102983.1

Int. Cl.: C07D 313/12 , C07D 405/06 ,
C07D 405/12 , A61K 31/335

Date of filing: 03.03.87

Priority: 03.03.86 JP 45676/86

Date of publication of application:
09.09.87 Bulletin 87/37

Designated Contracting States:
BE CH DE ES FR GB IT LI NL

Applicant: KYOWA HAKKO KOGYO CO., LTD.
6-1, Ohte-Machi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Oshima, Etsuo c/o KYOWA HAKKO
KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Kumazawa, Toshiaki c/o KYOWA
HAKKO KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Otaki, Shizuo c/o KYOWA HAKKO
KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Obase, Hiroyuki c/o KYOWA HAKKO
KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Ohmori, Kenji c/o KYOWA HAKKO
KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Ishii, Hidee c/o KYOWA HAKKO
KOGGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Manabe, Haruhiko c/o KYOWA
HAKKO KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Tamura, Tadafumi c/o KYOWA
HAKKO KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Inventor: Shuto, Katsuichi c/o KYOWA HAKKO
KOGYO CO., LTD

Patent Dept. 6-1 Ohtemachi Itchome
Chiyoda-ku Tokyo(JP)

Representative: Casalonga, Axel et al
BUREAU D.A. CASALONGA - JOSSE
Morassistrasse 8
D-8000 Munich 5(DE)

EP 0 235 796 A2

Xerox Copy Centre

⑤ Dibenzo [b,e] oxepin derivative and antiallergic and antiinflammatory agent.

⑥ Novel dibenz[b,e]oxepin derivatives are employed in the treatment and control of allergic conditions such as allergic asthma and also employed in the treatment of inflammation.

DIBENZ[b,e]OXEPIN DERIVATIVE AND ANTIALLERGIC AND ANTIINFLAMMATORY AGENT

Background of the Invention

Heretofore, it has been known that 11-unsubstituted, 11-hydroxy or 11-oxodibenz[b,e]oxepin derivative is used for antiinflammatory agents [J. Med. Chem., 21, 633-639 (1978)].

Further, it is known that dibenz[b,e]oxepin derivative wherein substituents Ra and Rb at 11-position have the following definitions, is employed in the treatment and control of allergic conditions (USP 4,282,365).


Ra : H, OH, lower alkoxy, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, arylthio, NH₂, NHCHO or imidazolyl;

Rb : H or lower alkyl;

or Ra and Rb taken together are = O, = CH-Rc wherein Rc is H or aryl.


Furthermore, it is known that 11-(4-methylpiperazino) dibenz[b,e]oxepin derivative has an antiasthmatic activity (USP 4,396,550, USP 4,465,835, EP-A-3856).

It is also known that dibenz[b,e]oxepin derivative having the following formula:




wherein R_d and R_e are lower alkyl and R_f is lower alkyl or halogen, has an antiasthmatic activity (EP-A-85870).

Dibenz[b,e]oxepin derivative having an antiallergic activity and having the following structural formula:




wherein R_g and R_h are alkyl, r is 2 or 3 and R_i is alkyl or halogen is known (JP-A-227879/84).

Dibenz[b,e]oxepin derivative having an antiallergic activity and having the following structural formula:




wherein R_j is 4-alkylpiperazino, 3-quinuclidylamino or -X_a-(CH₂)_s-NR_lR_m wherein X_a is -NH-, -S- or -O-, s is 2 or 3 and R_l and R_m are alkyl, and R_k is CN, 5-tetrazolyl, CONH₂ or CO₂R_n wherein R_n is H, alkyl or l-(ethoxycarbonyloxy)ethyl is known (EP-A-130555).

Doxepin having an antidepressant activity and having the following structural formula is known [Drugs, 13, 161 (1977)].



Dothiepin having an antidepressant activity and having the following structural formula is known [Arz.-Forsch., 13 1039 (1963); *ibid.*, 14 100 (1964)].




20 As the compound having both an antiallergic activity and an antiinflammatory activity, steroids are known.

It is always desired that a novel compound having an antiallergic activity or an antiinflammatory activity be developed.

25 Summary of the Invention

The present invention relates to a dibenz[b,e]oxepin derivative represented by the formula (I):



35

wherein A represents a hydroxymethyl, a lower alkoxyethyl, a triphenylmethyloxymethyl, a lower alkanoyloxymethyl, a lower alkanoyl, a carboxy, a lower alkoxy carbonyl, a triphenylmethyloxycarbonyl, -CONR₁R₂ (wherein R₁ and R₂ are the same or different and represent hydrogen atom or lower alkyl) 4,4-dimethyl-2-oxazoline-2-yl group or -CONHOH; Y represents -(CH₂)_m, -CHR₃(CH₂)_m-or -CR₄=CR₅(CH₂)_m which is substituent at 2-or 3-position of the mother nucleus (wherein R₃ represents a lower alkyl, R₄ and R₅ are the same or different and represent a hydrogen atom or a lower alkyl, m is 0, 1, 2, 3 or 4, and the left side of the group of Y mentioned above is bound to benzen nucleus); X represents = N-, = CH-or -CH₂-; n is 0, 1, 2, 3 or 4; Z represents 4-methylpiperazino, 4-methylhomopiperazino, piperidino, pyrrolidino, thiomorpholino, morpholino, or -NR₆R₇ (wherein R₆ and R₇ are the same or different and represent a hydrogen atom or a lower alkyl); and --- means a single bond or double bond [hereinafter referred to as Compound (I) and Compounds with other formula numbers are hereinafter likewise referred to], and a pharmaceutically acceptable salt thereof. The present invention further pertains to a pharmaceutical composition containing an effective amount of Compound (I) or a pharmaceutically acceptable salt thereof as an active ingredient, and a carrier or an excipient.

50 The present Compound (I) is useful for treatment of allergic conditions and inflammation.

55

Detailed Description of the Invention

In the definition of each group of formula (I), the lower alkyl group includes straight or branched chain alkyl groups having 1 to 6 carbon atoms, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, etc. In the definition of the group A, lower alkyl moiety of lower alkoxyethyl group and lower alkoxyethyl group has the same meaning as previously defined.

The lower alkoxyethyl group includes methoxyethyl, ethoxyethyl, n-propoxyethyl, isopropoxy, etc. and the lower alkoxyethyl group includes methoxyethyl, ethoxyethyl, etc.


In the definition of the group A, the lower alkyl moiety of lower alkanoyl group and lower alkanoyloxymethyl group has the same meaning as previously defined.

The lower alkanoyl group includes formyl, acetyl, etc. and the lower alkanoyloxymethyl group includes formyloxymethyl, acetyloxymethyl, etc.


The pharmaceutically acceptable salt of Compound (I) includes pharmaceutically acceptable acid addition salt, metal salt, ammonium salt, organic amine addition salt, amino acid addition salt, etc.

The pharmaceutically acceptable acid addition salt of Compound (I) includes inorganic acid salts such as hydrochloride, sulfate, phosphate, etc., and organic acid salts such as acetate, maleate, fumarate, tartrate, citrate, etc. The pharmaceutically acceptable metal salt includes alkalimetal salts such as sodium salt, potassium salt, etc., alkaline earth metal salts such as magnesium salt, calcium salt, etc., and aluminium salt, zinc salt, etc. The pharmaceutically acceptable organic amine addition salt includes addition salt of morpholine and piperidine and the pharmaceutically acceptable amino acid addition salt includes addition salt of lysine, glycine, phenylalanine, etc.

Compound (I) is prepared by using a compound represented by the formula (II):



30 wherein Y and A have the same meanings as previously defined or a compound represented by the formula (III):



40 wherein Y and A have the same meanings as previously defined as the starting compound. Compound (II) is disclosed in J. Med. Chem., 19, 941 (1976), ibid., 20, 1499 (1977) and JP-A-21679/83.

45 Compound (III) wherein -Y-A is -COOH is disclosed in JP-A-21679/83 and the other Compounds (III) can be prepared according to the method described in the publication though they do not occur in the publication.

The process for preparing Compound (I) is explained, depending on the kind of the group X.

Process A

50 [Synthesis of Compound (I) wherein X is =CH-(Part I)]

The carboxy group of Compound (IIa) is protected according to the following reaction scheme.

55

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.