



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07F 5/02, A01N 55/00, C09D 5/14, C08K 5/55</p>	A1	<p>(11) International Publication Number: <b>WO 95/33754</b></p> <p>(43) International Publication Date: 14 December 1995 (14.12.95)</p>
<p>(21) International Application Number: PCT/GB95/01206</p> <p>(22) International Filing Date: 26 May 1995 (26.05.95)</p> <p>(30) Priority Data: 9411587.0                      9 June 1994 (09.06.94)                      GB</p> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): AUSTIN, Peter, William [GB/GB]; 45 Randale Drive, Bury, Lancashire BL9 8NF (GB). KNEALE, Christopher, Juan [GB/GB]; 8 Dalefields, Delph, Oldham, Lancashire OL3 5HZ (GB). CROWLEY, Patrick, Jelf [GB/GB]; 56 Ellis Road, Crowthorne, Berkshire RG11 6PT (GB). CLOUGH, John, Martin [GB/GB]; 7 Gypsy Lane, Marlow, Buckinghamshire SL7 3JT (GB).</p> <p>(74) Agents: FAWKES, David, Melville; Intellectual Property Group, Zeneca Specialties, P.O. Box 42, Hexagon House, Blackley, Manchester M9 8ZS (GB) et al.</p>	<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p><b>Published</b> <i>With international search report.</i></p>	
<p>(54) Title: OXABORoles AND SALTS THEREOF, AND THEIR USE AS BIOCIDES</p> <p>(57) Abstract</p> <p>The use of oxaboroles and salts thereof as industrial biocides especially fungicides for the protection of plastics materials such as plasticised PVC. Preferred compounds are 5- and 6-fluoro or bromo- 1,3-dihydro-1-hydroxy-2,1-benzoxaborole including O-esters thereof.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## OXABORoles AND SALTS THEREOF, AND THEIR USE AS BIOCIDES

The present invention relates to the use of oxaboroles and salts thereof as industrial biocides, especially fungicides, biocidal compositions containing the oxaboroles including their salts and certain oxaboroles.


5 No single industrial biocide is ideal for all applications and new biocides are constantly being sought with better activity against individual spoilage micro-organisms, wider spectrum of activity, improved compatibility with the medium in which they are used and improved persistence in use. Safety in use is another important  
10 consideration.

A small number of compounds containing an oxaborole ring (hereinafter "oxaborole") have already been described in the literature. These are N-(1,3-dihydro-1-hydroxy-2,1-benzoxaborol-6-yl)-succinamic acid (CA 55 23423c); 4-(1,3-dihydro-1-hydroxy-2,1-benzoxaborol-6-ylazo)-  
15 2-naphthoic acid (CA 55 23423c); 1,3-dihydro-1-hydroxy-6-nitro-2,1-benzoxaborole (CA 55 23423b); 6-amino-1,3-dihydro-1-hydroxy-2,1-benzoxaborole and its hydrochloride (CA 55 23423c); 1,3-dihydro-1-hydroxy-7-methyl-2,1-benzoxaborole (CA 55 6473f); 1-(benzyloxy)-1,3-dihydro-2,1-benzoxaborole (CA 61 16084f); 1,3-dihydro-1-hydroxy-N,N-dimethyl-2,1-benzoxaborol-6-amine (CA 103(3) 22633f); 4-bromo-1,3-dihydro-1-hydroxy-2,1-benzoxaborole (CA 103(3) 22633f); 1,1'-oxybis[4-bromo-1,3-dihydro-2,1-benzoxaborole (CA 103(3) 22633f); 1-(cyclohexyloxy)-1,3-dihydro-2,1-benzoxaborole (CA 61 16084f); 1-ethoxy-1,3-dihydro-2,1-benzoxaborole (CA 61 16084f); 3,7-dihydro-1,5-dihydroxy-  
25 1H,3H-benzo[1,2-c: 4,5-c']bis[1,2]oxaborole (CA 61 14698a); 1,3-dihydro-1-hydroxy-6-methyl-2,1-benzoxaborole (CA 61 14698b); 5-bromo-1,3-dihydro-1-hydroxy-2,1-benzoxaborole-6-methanol (CA 51 14698b); 1,1'-oxybis[1,3-dihydro-2,1-benzoxaborole] (CA 103(3) 22633f); and boronophthalide (CA 116(13) 129587q). French certificate of utility No  
30 73 29370 discloses boronophthalide (1-hydroxy-3H-1,2-benzoxaborole) and this is the only citation known which discloses that an oxaborole is biologically active. It is disclosed as being useful in inhibiting the growth of micro organisms in aviation fuels. However, at least 100ppm of the boronophthalide is required to protect the fuel.

35 It has now been found that compounds containing an oxaborole ring are particularly effective against micro-organisms such as bacteria, algae, yeasts and particularly fungi, especially fungi which cause degradation of plastics materials. The level of microbiological activity now found is surprising in the light of the disclosure in the  
40 above utility certificate.

According to the present invention there is provided a method for the protection of a medium susceptible to microbial attack by the treatment of the medium with an effective amount of an oxaborole of general formula (1)

5




or a salt thereof  
wherein

10 A and D are each independently, hydrogen, optionally substituted  $C_{1-18}$ -alkyl, aralkyl, aryl, or heterocyclyl or where A and D together with the carbon atoms to which they are attached form a 5,6 or 7-membered fused ring which itself may be substituted;

15 X is a group  $-CR^1R^2$  wherein  $R^1$  and  $R^2$  are each, independently, hydrogen, optionally substituted  $C_{1-6}$ -alkyl, nitrile, nitro, aryl or aralkyl or  $R^1$  and  $R^2$  together with the carbon atom to which they are attached form an alicyclic ring;

20 R is hydrogen, optionally substituted  $C_{1-18}$ -alkyl, aralkyl, aryl, heteroaryl, cycloalkyl or a radical of formula (2)

20



25 wherein A, D and X are as hereinbefore defined except where the medium is aviation fuel and the only oxaborole is boronophthalide.

When A and/or D is alkyl, it may be linear or branched and is preferably  $C_{1-12}$ -, more preferably  $C_{1-8}$ - and especially  $C_{1-4}$ -alkyl.

30 When A and/or D is substituted alkyl, the substituent may be  $C_{1-6}$ -alkoxy, hydroxy, halogen, nitrile, amino, substituted amino, carboxy, acyl, aryloxy or carbonylamino optionally substituted by  $C_{1-6}$ -alkyl.

When A and/or D is alkyl the alkyl group or groups are preferably unsubstituted.

35 When A and/or D is aryl, it is preferably phenyl which may itself be substituted.

When A and/or D is aralkyl, it is preferably benzyl or 2-ethylphenyl, where the phenyl ring may be substituted.

40 When the phenyl ring is substituted, the substituents include  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy, aryloxy, hydroxy, halogen, nitro, carbonamido, sulphonamido, trifluoromethyl or amino optionally substituted by one or more  $C_{1-6}$ -alkyl groups.

Aryloxy is preferably phenoxy.

When A and D together with the two carbon atoms to which they are attached form a fused ring the ring may be alicyclic as in cyclopentene, cyclohexene or cycloheptene or it may be aromatic such as phenyl, pyridyl, thienyl or furanyl. The fused ring may also carry substituents as described hereinbefore for substituted phenyl and substituted alkyl. The fused ring may also contain more than one ring system, for example, a naphthyl or quinolinyl ring system or the fused ring may also link two oxaborole rings as for example in 1H,3H-benzo[1,2-c: 4,5-c']bis[1,2]oxaborole.

When R<sup>1</sup> and/or R<sup>2</sup> is aryl it is preferably phenyl.

When R<sup>1</sup> and/or R<sup>2</sup> is aralkyl it is preferably benzyl.

Preferably, at least one of R<sup>1</sup> and R<sup>2</sup> is hydrogen and it is especially preferred that both are hydrogen.

When R is alkyl it may be linear or branched and is preferably C<sub>1-12</sub>- and especially C<sub>1-6</sub>-alkyl.

When R is substituted alkyl, the substituent may be C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, amino, substituted amino, carboxy, aryl, aryloxy, carbonamido optionally substituted by C<sub>1-6</sub>-alkyl, aryl such as phenyl and aralkyl such as benzyl.

When R is aralkyl it is preferably benzyl or 2-ethylphenyl.

When R is aryl it is preferably phenyl.

When R is heteroaryl it is preferably quinolinyl and particularly quinolin-8-yl.

When R is cycloalkyl it is preferably cyclohexyl.

When the substituent is halogen, it is preferably bromine, chlorine and especially fluorine.

One preferred class of oxaborole is a benzoxaborole of formula 1 wherein A and D together with the carbon atoms to which they are attached form a fused phenyl, naphthyl or thienyl ring.

When the fused ring is phenyl, the oxaborole is a benzoxaborole and the substituent or substituents may be in any of positions 4,5,6 or 7 of the benzoxaborole. Preferably the substituent or substituents is/are in the 5 and/or 6 position. Preferred substituents are amino, alkyl, alkoxy, phenyl, phenoxy, sulphonamide, carbonamide, each of which may be substituted, and also trifluoromethyl, chlorine, bromine and especially fluorine.

When the fused ring is naphthyl, the other fused phenyl ring is attached to the benzoxaborole ring system in either the 4,5- or 5,6-position.

In one preferred class of oxaborole, R is hydrogen.

Another preferred class of oxaboroles for use in the present invention is where R is substituted alkyl, especially where the substituent is a primary, secondary or tertiary amino group and

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.