

Lucilla Parnetti  
Maurizio Paciaroni  
Virgilio Gallai

## Headache and ischemic stroke

Received: 27 September 2001  
Accepted in revised form: 27 December 2001

L. Parnetti (✉) • M. Paciaroni • V. Gallai  
Department of Neuroscience,  
University of Perugia,  
Via Enrico dal Pozzo, I-06126 Perugia, Italy  
e-mail: parnetti@unipg.it  
Tel.: +39-075-5783564  
Fax: +39-075-5783583

**Abstract** Migraine is a common neurological condition affecting yearly 1%–10% of all men and 3%–20% of all women. Focal neurological symptoms (auras), most commonly visual and sensory, occur in 4% of migraine attacks. Migraine with and without aura seems to be associated with an increased stroke risk. Migraine with aura may mimic transient ischemic attacks and may induce stroke (migrainous stroke). Headache is also a common symptom during ischemic stroke. In this review, we present the evidence about each of these circumstances to better understand the relationship between headache, especially migraine, and ischemic stroke.

**Key words** Headache • Stroke • Migraine • Cerebrovascular diseases

### Migraine and subsequent risk of ischemic stroke

Large-scale prospective epidemiological studies have identified major risk factors for stroke, including advanced age, race, family history of stroke, hypertension, diabetes mellitus, heart disease, elevated cholesterol levels, current smoking, obesity, heavy alcohol intake, sedentary lifestyle, use of exogenous hormones, and hyperhomocysteinemia [1, 2]. Although several clinical reports claim the existence of an association between migraine and stroke, only a few epidemiological studies have addressed this specific issue. The U.S. Physicians' Health Study showed that physicians aged

40–84 years with a history of migraine had increased risk of stroke; after adjustment for age and other cardiovascular risk factors, the relative risk was 1.84 (95% CI, 1.06 to 3.20) for total strokes and 2.0 (95% CI, 1.10 to 3.64) for ischemic strokes [3]. Merikangas et al. [4] also found that, after controlling for established risk factors for stroke (e.g. hypertension, diabetes, heart disease, gender), both migraine and severe nonspecific headache were associated with a significantly increased risk for stroke (RR, 1.5) that was higher in young patients: at the age of 40 years, the relative risk was 2.81 (95% CI, 1.45–5.43), at 50 years it was 2.07 (95% CI, 1.30–3.30), at 60 years it was 1.69 (95% CI, 1.10–2.60), and at 90 years it was 1.16 (95% CI, 0.63–2.11). Therefore,

migraine plays a more critical role in stroke of the young. In other studies the risk of ischemic stroke was not excessive among migraineurs aged  $\geq 60$  years [5, 6].

In recent years, several studies have been published on the association between migraine and stroke in young women under 45 years of age. Tzourio et al. [7] found that stroke was strongly associated with migraine, both without aura (odds ratio, 3.0; 95% CI, 1.5–5.8) and with aura (odds ratio, 6.2; 95% CI, 2.1–18.0). In another study [8], a history of migraine was more frequent in stroke patients than in controls (odds ratio, 1.9; 95% CI, 1.1–1.3). In the prospectively designed subgroup analyses, a history of migraine reached the highest odds ratio (3.7; 95% CI, 1.5–9.0) and was the only significant risk factor in women below age 35 years. Chang et al. [9] reported that in young women (20–44 years of age) a personal history of migraine was strongly associated with stroke (for all stroke: adjusted OR, 1.78; 95% CI, 1.14–2.77; for ischemic stroke: OR, 3.54; 95% CI, 1.30–9.61). In agreement with previous studies [5, 10], ischemic stroke was associated with both migraine with aura (OR, 3.81; 95% CI, 1.26–1.15) and migraine without aura (OR, 2.97; 95% CI, 0.66–1.35).

The risk of stroke in young migrainous women seems to apply not only to ischemic stroke, but also to hemorrhagic stroke, although hemorrhagic stroke has been not extensively studied and the results are less consistent [11].

In this category of patients, the risk of ischemic stroke was substantially increased by the use of oral contraceptives (OR, 13.9), for which a dose-effect relationship between risk of stroke and dose of estrogen was found (for pills containing 50  $\mu\text{g}$  estrogen: OR, 4.8; for pills with 30–40  $\mu\text{g}$ : OR, 2.7; pills with 20  $\mu\text{g}$ : OR, 1.7; pills with progesterone; OR, 1) [7]; the risk of ischemic stroke was also elevated in heavy smokers ( $\geq 20$  cigarettes per day: OR, 10.2). In migrainous women, coexistent use of oral contraceptives, history of high blood pressure and smoking habit had greater than multiplicative effects on the odds ratio for ischemic stroke (OR, 34.4; 95% CI, 3.27–3.61). However, this dramatic increase was based on only nine cases and two controls [9]. Other conditions predisposing to stroke, namely minor cardiac abnormalities like patent foramen ovale [12] and mitral valve prolapse [13] or the presence of anti-cardiolipin antibodies [14], may cause a stroke when combined with migraine, but this has not been definitely established [11].

Several hypotheses have been raised to explain the association between migraine and stroke: vasospasm, endothelial dysfunction, congenital thrombophilia, platelet hyperaggregability, and association with cardiac abnormalities predisposing to ischemic stroke. However, no fully convincing evidence has been produced [11].

The strength of the association should not, however, lead to the conclusion that all young women with migraine are at high risk of stroke. The incidence of ischemic stroke in

young women is low (approximately 10 cases in 100 000 women/years) [15, 16], and the risk of ischemic stroke is only 17–19 per 100 000 women with migraine per year [7]. Furthermore, it is not known whether this increased risk relates to all young migrainous women or only to a subgroup of them (i.e. patients with MELAS, cardiac abnormalities or antiphospholipid antibodies syndrome) [17–19]. However, young women, especially those <35 years of age, should be firmly advised to avoid smoking, and if they use oral contraceptives, to choose pills with a low estrogen content or only progesterone [11].

### Migrainous stroke

Stroke is a rare but potentially devastating complication of migraine. In 1988, the International Headache Society [20] classified migrainous infarction under the complications of migraine with aura (IHS 1.6.2), where the diagnosis of migraine-induced stroke in patients with migraine without aura is not allowed. To meet this definition, patients must have: (i) neurological symptoms and signs that are identical or similar to those of other migraine attacks and not completely reversible within 7 days and/or neuroimaging evidence of ischemic infarction in a relevant area; (ii) the stroke must have occurred during a typical migraine attack; (iii) other causes of stroke must be excluded, although stroke risk factors may be present. Several authors [21, 22] proposed that migrainous infarction should be redefined in the IHS classification as a possible complication of both migraine with and without aura because they described patients who had an ischemic stroke during a typical migraine attack (even if migraine with aura is substantially present (80%) in patients with migrainous stroke) [23]. In the WHO study [9], migrainous stroke was defined as any stroke occurring in presence of headaches in the 3 days before the stroke; following this rule, up to 40% of the strokes in migrainous women can be considered migrainous stroke.

The incidence of migrainous infarction has been estimated to be 3.36 per 100 000 persons per year, but in the absence of other stroke risk factors, this was reduced to 1.44 per 100 000 persons per year [24]. Migraine-induced stroke accounts for 0.8% of all strokes and accounts for as few as 4% [25] to as many as 20% of ischemic strokes in the young [26, 27].

Bogousslavsky et al. [28] found a marked female preponderance in the migrainous stroke group with respect to controls, and Iglesias et al. [29] claimed that the combination of smoking and migraine was highly associated with stroke. Oral contraceptives are also recognized to increase stroke risk in migraine sufferers [7]. Usually, headache fre-

quency and severity decrease after stroke, probably due to reduced nociceptive transmission as a result of loss in vasoreactivity of the affected cerebral blood vessel [30].

The prognosis of cerebral infarction associated with migraine is generally good. Hoekstra-van Dalen et al. [31] found that no patient had recurrent stroke during an average follow-up period of 5.8 years. Strokes related to migraine are commonly found in the territory of the posterior cerebral artery [27, 32]. The mechanism by which migraine causes a cerebral infarction is unknown; it has been proposed that during migraine attack there is a reduction of cerebral blood flow secondary to arteriolar vasoconstriction [33–35]. A slowly spreading wave of cortical neuronal depression with spreading oligoemia is also possible. Prolonged vasoconstriction and oligoemia may lead to hemostasis and predispose to intravascular thrombosis and migraine-induced cerebral infarction. Increased platelet aggregability, presence of antiphospholipid antibodies and use of oral contraceptives may contribute to the risk of enhanced coagulation [36–38]. The increase in procoagulant effects, the decrease in anticoagulant effects, and the hyperactivation of the antifibrinolytic system indicate that oral contraceptives do have a net prothrombotic effect [39]. In his model of neurogenic inflammation due to pathological activation of the trigeminovascular system during migraine attack, Moskowitz [40] showed that platelet aggregation occurs in the lumen of blood vessels. Another hypothesis suggests that repeated episodes of migraine-induced vasoconstriction may weaken the internal elastic lamina of cerebral vessels and predispose to arterial dissection [41]. Paradoxical embolism is another postulated cause [42]. Both migraine and stroke may also result from underlying genetic disorders such as MELAS or CADASIL [43, 44].

### Headache during ischemic stroke

Headache occurs frequently in acute ischemic stroke (IHS 6.1.2) or during a transient ischemic attack (IHS 6.1.1), but its frequency varies widely among different studies, ranging from 18% to 41% [45–55]. Overall, an underestimation of the frequency of headache in ischemic stroke patients can be claimed, as patients with language dysfunction, altered mental status or other factors impeding reliable determination of a headache complaint are excluded from most studies. In our series, headache was present in more than one-third of the patients with ischemic stroke and was much more common among patients with infarct in the posterior circulation (73.3%) than in those in whom the anterior circulation was involved (25.8%) [55]; this finding has also been described in other reports and may be due to the rich innervation by nociceptive afferents of vessels in the posterior circulation

[40, 51]. Stroke-related headache was frequently associated with large artery disease (40.7%) and was higher in patients with carotid artery occlusion. Similar results have been obtained by other authors, who reported headache frequencies ranging from 26% to 35% in patients with symptomatic carotid artery disease [46, 48]. Gorelick et al. [48] reported that there is no significant difference in the frequency of onset headache between patients with extracranial carotid artery disease, disease of the carotid siphon, middle cerebral artery, or carotid siphon and middle cerebral artery in tandem. As already reported, in the great majority of papers, headache is less common in lacunar infarction. In our series, the frequency of headache in lacunar infarction was 12.9%; similar frequencies have been observed by others [47, 49, 51, 56]. A history of headache, was present only in stroke patients with headache, and headache anticipated (on average by 2 days) the stroke in 24% of subjects. This observation is in agreement with previous studies and demonstrates that stroke in many cases is the result of a long-lasting pathological vascular process, in which headache merely serves as a warning sign of ischemic stroke [48, 51]. In several series, a higher frequency of onset headache was present in women and young people [57, 58], who generally have a higher frequency of headaches. The mechanisms underlying headache are not known. Several studies have suggested that headache is associated with dilation of some arteries at the base of the brain [59]. Headache is probably related to activation of nociceptive trigeminovascular afferents; pain ensues when a sufficient amount of nociceptors have been recruited [60]. The release of amino-acid neurotransmitters [61] and platelet activation [62] may also play a role in the pathogenesis of headache occurring at the onset of ischemic stroke.

Headache is also common in patients with cervical artery dissection [63]. Unilateral facial or orbital pain is present in half of patients with internal carotid artery dissection. The characteristic unilateral headache develops in two-thirds of patients, most commonly in the frontotemporal area, but it occasionally involves the hemicranium or the occipital area [64]. The onset of headache is usually gradual, but it may be an instantaneous, excruciating, “thunderclap” headache that mimics a subarachnoid hemorrhage [64, 65]. The headache is most commonly described as a constant steady aching, but it may also be throbbing or steady and sharp [63, 64]. About one-fourth of patients with a history of migraine claim that the headache is similar to a migraine attack. In case of vertebral artery dissection, headache occurs in two-thirds of patients, almost always in the occipital area, but in rare cases it involves the hemicranium or the frontal area or is bilateral [64]. Pain is usually the initial manifestation of cervical artery dissection and the mean time to the appearance of other symptoms is four days for carotid dissection and 15 hours for vertebral dissection [63]. In our experience [66],

headache was present in 10% of patients with cervical artery dissection before the focal neurological deficits, while 66.7% of the patients had headache during the dissection-related stroke.

---

### Migraine syndromes that mimic stroke

Migrainous syndromes that may mimic conventional cerebrovascular syndromes include hemiplegic migraine and basilar artery migraine. The IHS classifies hemiplegic migraine under migraine with typical aura (IHS 1.2.1) or prolonged aura (IHS 1.2.2). Familial hemiplegic migraine is classified as a subgroup of migraine with aura (IHS 1.2.3) and the definition includes the criteria for migraine with aura with hemiplegic features and at least one first-degree relative with identical attacks. Hemiplegic migraine attacks are characterized by hemiparesis or hemiplegia. There is an autosomal dominant inheritance pattern of the disorder. Hemiplegic migraine attacks may also be part of other familial disorders, namely mitochondrial encephalomyopathy with lactic acidosis and stroke-like episodes (MELAS), and cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL). Basilar migraine (IHS 1.2.4) is another disorder that may mimic a transient cerebrovascular accident. The diagnostic criteria include those for migraine with aura plus two or more aura symptoms of the following types: visual symptoms in both the temporal and nasal fields of both eyes, dysarthria, vertigo, tinnitus, decreasing hearing, double vision, ataxia, bilateral paresthesias, bilateral paresis, and decreased level of consciousness. The diagnosis of basilar migraine is supported by the pattern of evolution of the neurological deficit and the accompanying headache, the history of previous similar attacks, a positive family history and the often negative diagnostic workup [67]. It is not unusual for patients who have experienced migraine with aura to suffer identical auras at other times, but without headache, particularly as they get older [68, 69]. Diagnostic difficulty arises when a patient over 40 years of age with no previous history of classic migraine presents a first-ever episode of transient symptoms of focal neurological dysfunction that are typical of a

migrainous aura, but without any associated headache [70]. The differential diagnosis of this circumstance is important especially for the prognosis: after 10 years of follow-up, the relative risk of a stroke occurring in a patient with a history of transient ischemic attack (TIA) compared with that in a patient with “migrainous aura without headache” was 7.6 (95% CI, 0.8–74), and the relative risk of any serious vascular event (e.g. stroke, myocardial infarction, or vascular death) was 3.8 (95% CI, 0.8–19) [71].

---

### Conclusions

Headache is an underemphasized feature of ischemic cerebrovascular disease; the thorough evaluation of its clinical features can be of help in diagnosing correctly different cerebrovascular pathological conditions.

The relationship between headache, namely migraine, and stroke is difficult to disentangle. Epidemiological studies indicate that the comorbidity of migraine and stroke in the young is an important issue and has implications for patient management.

Migraine is associated with an increased incidence of stroke in young migraineur women who smoke or use oral contraceptives; these subjects should be strongly advised to discontinue smoking and they should be discouraged to use oral contraceptives; in any case, only pills with a low (<50 µg) estrogen content should be used.

No randomized study of primary stroke prevention in migraineurs or of secondary prevention in patients with a migraine-associated stroke has been performed. In patients who have had a migraine-induced stroke, daily use of an antiplatelet agent is reasonable although unproven, while the use of vasoconstrictive agents (e.g. ergotamine, triptans) for treating migraine attack should be avoided.

The mechanism by which migraine causes a cerebral infarction – i.e. why focal oligoemia sometime progresses to cerebral infarction – is substantially unknown. Pathophysiological studies aimed at identifying the role played by each different component (endothelium, platelets, leukocytes, coagulation factors, genetic susceptibility) in the ischemic cascade are highly recommended.

---

### References

1. Warlow CP, Dennis MS, van Gijn J, Hankey GJ, Sandercock PAG, Bamford JM, Wardlaw JM (2001) What caused this transient or persisting ischaemic event? In: *Stroke. A practical guide to management*, 2nd edn. Blackwell Science, Oxford, pp 223–300
2. Gallai V, Caso V, Paciaroni M, Cardaioli G, Arning E, Bottiglieri T, Parnetti L (2001) Mild hyperhomocysteinemia. A possible risk factor for cervical artery dissection. *Stroke* 32:714–718
3. Buring JE, Hebert P, Romero J, Kittross A, Cook N, Manson J, Peto R, Hennekens C (1995) Migraine and subsequent risk of stroke in the Physicians' Health Study. *Arch Neurol* 52:129–134

4. Merikangas KR, Fenton BT, Cheng SH, Stolar MJ, Risch N (1997) Association between migraine and stroke in a large-scale epidemiological study of the United States. *Arch Neurol* 54:362–368
5. Tzourio C, Iglesias S, Hubert JB, Visy JM, Alperovitch A, Tehindrazanarivelo A, Bioussé V, Woimant F, Bousser MG (1993) Migraine and risk of ischemic stroke: a case-control study. *BMJ* 307:289–292
6. Mosek A, Marom R, Korczyn A, Bornstein N (2001) A history of migraine is not a risk factor to develop an ischemic stroke in the elderly. *Headache* 41:399–401
7. Tzourio C, Tehindrazanarivelo A, Iglesias S, Alperovitch A, Chedru F, d'Anglejan-Chatillon J, Bousser MG (1995) Case-control study of migraine and risk of ischemic stroke in young women. *BMJ* 310:830–833
8. Carolei A, Marini C, De Matteis G (1996) History of migraine and risk of cerebral ischaemia in young adults. The Italian National Research Council Study Group on Stroke in the Young. *Lancet* 347:1503–1506
9. Chang CL, Donaghy M, Poulter N (1999) Migraine and stroke in young women: case-control study. *BMJ* 318:13–18
10. Lidegaard O (1995) Oral contraceptives, pregnancy and the risk of cerebral thromboembolism: the influence of diabetes, hypertension, migraine and previous thrombotic disease. *Br J Obstet Gynaecol* 102:153–159
11. Tzourio C, Kittner SJ, Bousser MG, Alperovitch (2000) Migraine and stroke in young women. *Cephalalgia* 20:190–199
12. Anzola GP, Magoni M, Guindani M, Rozzini L, Dalla Volta G (1999) Potential source of cerebral embolism in migraine with aura – a transcranial Doppler study. *Neurology* 52:1622–1625
13. Spence JD, Wong DG, Melendez LJ, Nichol PM, Brown JD (1984) Increased prevalence of mitral valve prolapse in patients with migraine. *Can Med Assoc J* 131:1457–1460
14. Welch KMA (1994) Relationship of stroke and migraine. *Neurology* 44[Suppl 7]:S33–S36
15. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception (1995) A multinational study of cardiovascular description and validation of methods. *J Clin Epidemiol* 48:1513–1547
16. Petitti DB, Quesenberry CP, Bernstein A (1997) Incidence of stroke and myocardial infarction in women of reproductive age. *Stroke* 28:280–283
17. Montagna P, Galassi R, Medori R, Govoni E, Zeviani M, Di Mauro S, Lugaresi E, Andermann F (1988) MELAS syndrome: characteristics migrainous and epileptic features and maternal transmission. *Neurology* 38:751–754
18. Levine SR, Welch KMA (1989) Antiphospholipid antibodies. *Ann Neurol* 26:386–389
19. Pearson AC, Nagelhout D, Castello R, Gomez C, Labovitz A (1991) Atrial septal aneurysm and stroke: a transesophageal echocardiographic study. *J Am Coll Cardiol* 18:1223–1229
20. – (1988) Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. Headache Classification Committee of the International Headache Society. *Cephalalgia* 8[Suppl 7]:1–96
21. Welch KMA, Levine SR (1990) Migraine-related stroke in the context of the International Headache Society classification of head pain. *Arch Neurol* 47:458–462
22. Narbone MC, Leggiadro N, La Spina P, Rao R, Grugno R, Musolino R (1996) Migraine stroke: a possible complication of both migraine with and without aura. *Headache* 36:481–483
23. Rothrock J, North J, Madden K, Lyden P, Fleck P, Dittrich H (1993) Migraine and migrainous stroke: risk factors and prognosis. *Neurology* 43:2473–2476
24. Henrich JB, Sandercock PAG, Warlow CP, Jones LN (1986) Stroke and migraine in the Oxfordshire Community Stroke Project. *J Neurol* 233:257–262
25. Adams HP Jr, Kappelle J, Biller J, Gordon DL, Love BB, Gomez F, Heffner M (1995) Ischemic stroke in young adults: experience in 329 patients enrolled in the Iowa Registry of Stroke in Young Adults. *Arch Neurol* 52:491–495
26. Bogousslavsky J, Pierre P (1992) Ischemic stroke in patients under age 45. *Neurol Clin* 10:113–124
27. Broderick JP, Swanson JM (1987) Migraine-related strokes: clinical profile and prognosis in 20 patients. *Arch Neurol* 44:868–871
28. Bogousslavsky J, Regli F, van Melle G, Payot M, Uske A (1988) Migraine stroke. *Neurology* 38:223–227
29. Iglesias S, Visy JM, Hubert JB, Tehindrazanarivelo A, Tzourio C, Bousser MG (1993) Migraine as a risk factor for ischemic stroke: a case-control study. *Stroke* 24:171
30. Linetsky E, Leker RR, Ben-Hur T (2001) Headache characteristics in patients after migrainous stroke. *Neurology* 57:130–132
31. Hoekstra-van Dalen RA, Cillessen JP, Kappelle LJ, van Gijn J (1996) Cerebral infarcts associated with migraine: clinical features, risk factors and follow-up. *J Neurol* 243:511–515
32. Sacquegna T, Andreoli A, Baldrati A, Lamieri C, Guttman S, de Carolis P, Di Pasquale G, Pinelli G, Testa C, Lugaresi E (1989) Ischemic stroke in young adults: the relevance of migrainous infarction. *Cephalalgia* 9:255–258
33. Levine SR, Welch KMA, Ewing JR, Robertson WM (1987) Asymmetric cerebral blood flow patterns in migraine. *Cephalalgia* 7:245–248
34. Lauritzen M, Olesen J (1984) Regional cerebral blood flow during migraine attacks by xenon 133 inhalation and emission tomography. *Brain* 107:447–461
35. Olesen J (1991) Cerebral and extracranial circulatory disturbances in migraine: pathophysiological implications. *Cerebrovasc Brain Metab Rev* 3:1–28
36. Joseph R, Welch KMA (1987) Migraine and the platelet: nonspecific association. *Headache* 27:375–380
37. Brey RL, Hart RG, Sherman DG, Tegeler CH (1990) Antiphospholipid antibodies and cerebral ischemia in young people. *Neurology* 40:1190–1196
38. Conard J, Samama MM (2000) Oral contraceptives, hormone replacement therapy and hemostasis. *Cephalalgia* 20:175–182
39. Vandenbroucke JP, Rosing J, Bloemenkamp KWM, Middeldorp S, Helmerhorst FM, Bouma BN, Rosendaal FR (2001) Oral contraceptives and the risk of venous thrombosis. *N Engl J Med* 344:1527–1535

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.