

Hand grip strength in patients with type 2 diabetes mellitus

Ercan Cetinus^{a,*}, Mehmet Akif Buyukbese^b, Murat Uzel^a,
Hasan Ekerbicer^c, Ahmet Karaoguz^a

^a Kahramanmaraş Sutcu Imam University, Faculty of Medicine, Department of Orthopedics, 46050 Kahramanmaraş, Turkey

^b Kahramanmaraş Sutcu Imam University, Faculty of Medicine, Department of Internal Medicine, 46050 Kahramanmaraş, Turkey

^c Kahramanmaraş Sutcu Imam University, Faculty of Medicine, Department of Public Health, 46050 Kahramanmaraş, Turkey

Received 6 December 2004; received in revised form 21 February 2005; accepted 23 March 2005

Abstract

Aim: The aim of the present study was to compare hand grip strength and pinch power, which are important parameters of hand function, in 76 patients with type 2 diabetes mellitus (T2DM) (mean age: 50.11 ± 7.6) with 47 non-diabetic control subjects (mean age: 46.93 ± 10.2).

Methods: Grip strength was assessed with a Jamar dynamometer and pinch power was measured with a pinch gauge. Body composition was measured using a Tanita body composition analyzer. Mann–Whitney, Fisher’s exact and chi-square tests were used to determine the differences within groups and a p -value <0.05 was taken as statistically significant.

Results: Hand grip strength test values were significantly lower in the diabetic group compared with the control group. Key pinch power value for the right hand was significantly lower in the diabetic group than in the control group whereas the left hand value was similar.

Conclusion: Hand grip strength and key pinch power values were found to be lower in patients with T2DM than in age-matched control subjects. Hands, as well as feet, are also affected by diabetes and physicians should be aware of this.

© 2005 Published by Elsevier Ireland Ltd.

Keywords: Type 2 diabetes; Hand grip strength; Key pinch power; Jamar dynamometer; Pinch gauge

1. Introduction

Type 2 diabetes mellitus (T2DM), is the most common endocrine disorder worldwide, and it is

characterized by metabolic abnormalities and by chronic complications involving the eyes, kidneys, nerves, and blood vessels [1]. These complications can cause morbidity and premature mortality, and lead to serious social and cause economic problems due to loss of employment.

Foot ulcers and joint problems in T2DM are the most significant causes of morbidity and admittance to orthopedic outpatient clinics. The major predisposing

* Corresponding author. Tel.: +90 344 2212337/364-226;

fax: +90 344 2212371.

E-mail addresses: ercancetinus@hotmail.com, ecetinus2000@yahoo.com (E. Cetinus).

41 cause is diabetic polyneuropathy because the sensory
42 denervation impairs the perception of trauma after
43 wearing ill-fitting shoes. Alterations in proprioception
44 may give rise to an abnormal pattern of weight bearing
45 and sometimes to the development of Charcot's joints.
46 In addition to sensory neuropathy, motor neuropathy is
47 often emphasized, considering that diabetic foot
48 pathology, which is characterized by intrinsic muscle
49 atrophy, can result in a motor imbalance and diffuse
50 claw-toe. This pathology affects both the foot
51 function and postural stability [2].

52 In diabetic patients, the strength of flexor and
53 extensor muscles at the elbow, wrist, knee, and ankle
54 have been evaluated clinically using manual muscle
55 testing (MMT) and isokinetic dynamometry [3,4].
56 The volume of ankle dorsal and plantar flexors, and
57 intrinsic muscle atrophy of the foot have been
58 investigated radiologically using magnetic reso-
59 nance imaging (MRI) [2,5]. In contrast to the
60 measurement of the strength of the lower extremity
61 muscles; hand grip strength has seldom been studied
62 in patients with diabetes mellitus (DM) [6]. In the
63 present study, our aims were to establish, using a
64 Jamar dynamometer and a pinch gauge, whether the
65 grip and pinch power of the hand in patients with
66 DM were different than those of healthy non-diabetic
67 control subjects.

68 2. Patients and methods

69 Seventy-six patients with T2DM (mean age:
70 50.11 ± 7.6 years) were recruited from outpatient
71 clinics of the Department of Internal Medicine, at
72 Kahramanmaraş Sutcu Imam University. Forty-seven
73 healthy volunteers (mean age: 46.93 ± 10.2) without
74 diabetes, established by an oral glucose tolerance test
75 (OGTT), served as the control group.

76 DM was diagnosed according to American
77 Diabetes Association (ADA) diagnostic criteria as
78 follows: a fasting plasma glucose ≥ 7.0 mmol/L or 2-h
79 plasma glucose ≥ 11.1 mmol/L after a 75 g oral
80 glucose load [7]. Criteria for inclusion in the study
81 were that the patients had T2DM (known or newly
82 diagnosed after glucose challenge test or those
83 receiving oral hypoglycemic pills) and that the control
84 subjects had no glucose abnormality, no history of
85 pain in the shoulder, arm or hand, no documented

86 history of trauma or cervical radiculopathy in the
87 previous 12 months.

88 A calibrated, Jamar dynamometer (Smith and
89 Nephew, Irvington, NY 10533, USA) was used to
90 assess grip strength at the first three settings. A pinch
91 gauge (PG-30, B&L Engineering Santa Fe, CA, USA)
92 was used to assess the key pinch. Both the
93 dynamometer and pinch gauge were reset to zero
94 prior to each reading and were read to the nearest
95 increment of the two scale divisions. The American
96 Society of Hand Therapists' recommendations for
97 testing both grip and pinch strengths were followed
98 [8]. Subjects were seated comfortably on a chair
99 without armrests. The shoulder was adducted and
100 neutrally rotated, with the elbow at 90° flexion, and the
101 forearm and wrist in a neutral position. Standard
102 verbal encouragement in the same tone of voice
103 ("squeeze the handle/button as hard as possible") was
104 used during the measurements. Three measurements
105 of each grip and pinch were obtained at 15 s intervals
106 and mean values were analyzed. Measurements started
107 with the dominant hand. The right hand was dominant
108 in 67 (88.2%) of T2DM patients, whereas in 2 (2.6%)
109 the left hand was dominant, and the remaining 7
110 (9.2%) were ambidextrous. In the control group the
111 dominant hand was the right in 38 (80.9%) subjects,
112 the left in 6 (12.8%), and 3 subjects (6.4%) were
113 ambidextrous.

114 Percentages of body fat (BF), the basal metabolism
115 rate (BMR), and fat mass of the subjects were obtained
116 using a Tanita body composition analyzer TBF-300
117 (Tanita Corp., Tokyo, Japan). Tanita TBF-300 is a
118 commercially available foot-to-foot bioelectrical
119 impedance analysis (BIA) system. The manufac-
120 turer-supplied equations incorporate gender, mass,
121 height, activity category and a measured impedance
122 value to determine the percentages of BF, BMR, and
123 fat mass. In order to assess these measurements, girth-
124 hip ratio (G/H), height, body weight, and body mass
125 index (BMI) were all measured.

126 All T2DM patients were examined for hyperten-
127 sion, smoking and diabetes duration, and were
128 investigated for diabetic complications using clinical
129 examination and laboratory findings (Urine protein,
130 HbA1c).

131 The study was reviewed and approved by the local
132 research and ethics committee and all subjects gave
133 written consent.

DIAB 3346 1–9

Sanofi Exhibit 2176.002

134 Parametric or nonparametric tests were chosen to
 135 test for statistical significance depending on the data
 136 distribution. Mann–Whitney, chi-square, Fisher’s
 137 exact, *T*-test, Kruskal–Wallis analysis of variance,
 138 Wilcoxon’s signed rank test and Pearson’s correlation
 139 coefficient were used to determine the differences and
 140 relations between groups. A *p*-value of <0.05 was
 141 taken as statistically significant. Statistical analysis
 142 was performed using SPSS 9.0 for Windows (SPSS
 143 Inc., Chicago, IL, USA).

144 3. Results

145 The characteristics and body composition values of
 146 the subjects were given in Tables 1 and 2. There were
 147 no significant difference between the groups with
 148 respect to age, sex, hypertension, proteinuria, and
 149 smoking (*p* > 0.05). However HbA1c values in
 150 diabetic patients were significantly higher than those
 151 of the control group ($7.14 \pm 1.64\%$ versus
 152 $5.16 \pm 0.62\%$, *p* < 0.001) (Table 1). BMI, G/H, BF,
 153 BMR, and fat mass were similar in both groups
 154 (Table 2).

155 Working status of subjects is given in Fig. 1. All
 156 subjects in both groups were classified as non-manual
 157 workers (housewives, civil servants, tradesmen,
 158 retired etc.) (Fig. 1).

159 The results of the hand grip strength test with the
 160 Jamar dynamometer were significantly lower in the
 161 diabetic group compared with the control group
 162 (*p* < 0.05). The key pinch strength value for the
 163 right hand was significantly lower in the diabetic
 164 group than in the control group (*p* < 0.05), whereas
 165 the left hand value was lower than in the control
 166 group but this was not statistically significant

(*p* > 0.05) (Table 3). However, when the subjects
 of both the diabetic and control group were classified
 according to age intervals, hand grip and pinch
 strength values were found to be lower in diabetic
 patients in both the 30–49 and the >50 age groups
 (*p* < 0.05) (Table 4).

54.5% of the diabetic patients reported that daily
 activities and hand grasping power were not affected
 and 10.6% said that decreased hand power did not
 affect their daily activities. However, patients who
 reported that their hand power affected daily activities
 comprised 34.9% of the diabetic study group.

The relationship between HbA1c level and values
 of hand grip and key pinch strength of diabetic and
 control groups were analyzed using Pearson’s
 correlation coefficient. There was no relationship
 found among HbA1c levels, proteinuria, hypertension
 and values of hand grip and key pinch strength in
 neither the diabetic patients nor the controls
 (*p* > 0.05).

The relationship between age, BMI and values of
 hand grip and key pinch strength of both the diabetic
 and control group were analyzed using Pearson’s
 correlation coefficient but no significant relationship
 was (*p* > 0.05).

Both the hand grip and key pinch strength values
 were found to be higher in males than in females in
 both the diabetic and the control group (*p* < 0.05)
 (Table 5). Furthermore, there was a relationship
 between the grip strength and key pinch power in both
 the diabetic and control groups (*p* < 0.001). In the
 diabetic group, the relationship between grip strength
 and key pinch power was relatively higher than those
 of the control group (Table 6).

There were significant differences in the hand grip
 and key pinch strength values of the subjects between

Table 1
 Characteristics of subjects

	Diabetic patients (<i>n</i> = 76)	Control (<i>n</i> = 47)	<i>p</i> -Value
Age (years) (mean ± S.D.)	50.11 ± 7.6	46.93 ± 10.2	>0.05
Sex F/M (<i>n</i> and %)	51 (67.1%)/25 (32.9%)	28 (59.6%)/19 (40.4%)	>0.05
Hypertension	(<i>n</i> = 49) 65.3%	(<i>n</i> = 20) 42.5%	>0.05
Proteinuria	(<i>n</i> = 13) 17.1%	(<i>n</i> = 6) 12.8%	>0.05
Smoking	(<i>n</i> = 1) 1.3%	(<i>n</i> = 3) 6.4%	>0.05
HbA1c (%) (mean ± S.D.)	7.14 ± 1.64 (4.6 – 10.9)	5.16 ± 0.62 (4.1 – 7.9)	<0.001
Diabetes duration (years)	5.94 ± 6.18	–	–

No significant difference between groups (*p* > 0.05) in age, sex, hypertension, proteinuria, smoking, but significant difference between groups (*p* < 0.001) in HbA1c.

Table 2

Comparison of the body composition values (values are mean \pm S.D.)

	Diabetic patients (n = 76)	Control (n = 47)	p-Value
BMI	30.59 \pm 6.03	31.22 \pm 5.0	>0.05
Girth–hip ratio (G/H)	0.88 \pm 0.07	0.86 \pm 0.06	>0.05
Percentages of body fat (BF)	34.86 \pm 9.19	34.89 \pm 7.85	>0.05
Basal metabolism rate (BMR)	6341.77 \pm 795.49	6565.51 \pm 1283.85	>0.05
Fat mass	28.39 \pm 12.19	29.08 \pm 10.82	>0.05

No significant difference between groups ($p > 0.05$) in BMI, G/H, BF, BMR, and fat mass.

203 the dominant and non-dominant hand ($p < 0.05$)
204 (Table 7).

205 Implications on life and activities, as a conse-
206 quence of the patients' lower hand grip and pinch
207 strength values were investigated. All the diabetic
208 patients were studied. 54.5% of the diabetic patients
209 reported that daily activities and hand grasping power
210 were not affected and 10.6% stated that decreased
211 hand power did not affect their daily activities.
212 However, subjects with affected hand power and daily
213 activities comprised 34.9% of the diabetic population.

214 4. Discussion

215 It is well known that mild distal muscle weakness
216 can accompany predominant distal symmetrical

217 sensory neuropathy in DM patients [9]. While there
218 are numerous quantitative studies on sensory neuro-
219 pathy and autonomic disturbances, there is little data
220 about motor function in diabetic patients [10,11].
221 Dyck et al. [3] indicated that clinically apparent
222 muscle weakness was a severe disturbance in type 1
223 diabetes (T1DM) patients with more advanced
224 neuropathy. However, neither the severity nor the
225 distribution of the muscle weakness due to manual
226 muscle testing (MMT) was reported in their clinical
227 studies. Andersen and Jakobsen stated that the
228 sensitivity of MMT was low and dynamometry should
229 be considered in clinical trials of motor function in
230 neuropathic patients [12]. Some investigators reported
231 that there was a significant reduction in the muscle
232 strength of the ankle dorsal and plantar flexors, and the
233 knee extensors and flexors in 56 T1DM patients using




Fig. 1. Working status of subjects.

DIAB 3346 1–9

Sanofi Exhibit 2176.004

Table 3
Hand grip strength and key pinch values (kg) of both groups

Strength	Diabetic patients	Control
R Jamar first setting	27.61 ± 9.76	31.89 ± 8.88 ^a
R Jamar second setting	31.53 ± 11.82	36.34 ± 11.01 ^a
R Jamar third setting	28.92 ± 10.86	33.22 ± 10.53 ^a
L Jamar first setting	25.91 ± 9.53	31.10 ± 9.08 ^a
L Jamar second setting	29.77 ± 11.15	35.48 ± 10.35 ^a
L Jamar third setting	27.54 ± 10.51	32.05 ± 9.30 ^a
R key pinch	8.47 ± 2.56	9.37 ± 1.89 ^b
L key pinch	8.15 ± 2.50	8.92 ± 1.83 ^c

R: right; L: left.

^a $p < 0.05$. Hand grip strength was significantly lower in the diabetic group than those of control group.

^b $p < 0.05$. Key pinch strength value for right hand was significantly lower in the diabetic group than value of control group.

^c $p > 0.05$. Key pinch strength value for left hand was lower in diabetic patients than value of control group. But there was no statistically significant difference.

isokinetic dynamometer, but a reduction in muscle strength of the wrist flexors and extensors was not significant [4]. Lord et al. [13] found impaired muscle strength of the knee extension in a group of aged women with T2DM. Andersen et al. [14] pointed out that T2DM patients may have weakness of the extensors and flexors at the ankle and of the knee flexors and extensors, with a preservation of muscle strength at the wrist and elbow. It was thought that the distribution of muscular weakness indicated a distal neuropathic process underlying the impaired motor performance, and this assumption was supported by the observation that muscular strength at the ankle and knee was related to the degree of severity of neuropathy.

In addition to clinically determined lower extremity muscular weakness in DM patients, it was reported that there was a 32% reduction in the volume of dorsal and plantar flexors [5], and also using MRI, remarkable atrophy in the intrinsic muscles of the foot in neuropathic patients was reported [2]. Both biochemical and structural changes in the plantar foot muscles of DM patients with neuropathic ulcers and a reduction in high-energy metabolites with an increase in fat content were also demonstrated via magnetic resonance spectroscopy [15]. Significant relationships between motor nerve conduction velocity, and these physiological variables were suggested as atrophy in the intrinsic muscles of the foot was seen as secondary to motor nerve dysfunction. Remarkable atrophy of the foot and ankle muscles was thought to be secondary to diabetic neuropathy [2,15].

A number of investigations related to the evaluation of the muscle strength in DM patients were carried out on the lower extremity muscles, and mild distal muscle weakness in the lower extremity, due to diabetic neuropathy, was identified. However, hand grip strength and pinch power values in diabetic patients are unclear in the literature.

In the present study, we evaluated the grip and pinch power of the hand in T2DM. Grip strength and pinch power are important parameters of hand function. The grip strength test was commonly used to evaluate the integrated performances of hand muscles by determining maximal grip force that could be produced in one muscular contraction [16]. Hand strength can be used to determine a treatment [17], to assess nutrition [18], to assess risk of mortality in

Table 4
Comparisons of hand grip and pinch strength (kg) values (mean ± S.D.) of subjects according to age intervals

Strength (kg)	Diabetic patients		Control	
	30–49	>50	30–49	>50
R Jamar first setting	28.32 ± 10.78	27.06 ± 9.01	32.35 ± 9.31 ^a	30.62 ± 7.81 ^b
R Jamar second setting	32.49 ± 12.65	30.79 ± 11.24	35.87 ± 10.78 ^a	37.65 ± 12.03 ^b
R Jamar third setting	29.44 ± 11.26	28.52 ± 10.66	32.55 ± 9.90 ^a	35.04 ± 12.39 ^b
L Jamar first setting	26.91 ± 10.03	25.13 ± 9.18	31.69 ± 9.49 ^a	29.48 ± 8.00 ^b
L Jamar second setting	30.46 ± 11.96	29.25 ± 10.61	35.18 ± 10.48 ^a	36.32 ± 10.38 ^b
L Jamar third setting	27.54 ± 10.88	27.55 ± 10.36	31.78 ± 9.38 ^a	32.82 ± 9.46 ^b
R key pinch	8.78 ± 2.66	8.21 ± 2.49	9.42 ± 1.95 ^a	9.24 ± 1.80 ^b
L key pinch	8.11 ± 2.36	8.18 ± 2.65	8.94 ± 1.72 ^a	8.85 ± 2.22 ^b

R: right; L: left.

^a $p < 0.05$. Hand grip and pinch strength values of diabetic patients were lower than those of controls in 30–49 age interval.

^b $p < 0.05$. Hand grip and pinch strength values of diabetic patients were lower than those of controls in >50 age.

DIAB 3346 1–9

Sanofi Exhibit 2176.005

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.