



US007476652B2

(12) **United States Patent**
Brunner-Schwarz et al.

(10) **Patent No.:** **US 7,476,652 B2**
(45) **Date of Patent:** **Jan. 13, 2009**

(54) **ACIDIC INSULIN PREPARATIONS HAVING IMPROVED STABILITY**

(75) Inventors: **Anette Brunner-Schwarz**, Frankfurt (DE); **Norbert Lill**, Kronberg (DE)

(73) Assignee: **Sanofi-Aventis Deutschland GmbH**, Frankfurt (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 40 days.

(21) Appl. No.: **11/089,777**

(22) Filed: **Mar. 25, 2005**

(65) **Prior Publication Data**

US 2005/0171009 A1 Aug. 4, 2005

Related U.S. Application Data

(63) Continuation of application No. 10/461,740, filed on Jun. 13, 2003, now abandoned.

(60) Provisional application No. 60/409,338, filed on Sep. 9, 2002.

(30) **Foreign Application Priority Data**

Jun. 18, 2002 (DE) 102 27 232

(51) **Int. Cl.**
A61K 38/28 (2006.01)

(52) **U.S. Cl.** 514/3; 514/4

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,758,683 A 9/1973 Jackson
3,868,358 A 2/1975 Jackson
4,153,689 A * 5/1979 Hirai et al. 514/3
4,608,364 A 8/1986 Grau
4,614,730 A 9/1986 Hansen et al.
4,644,057 A 2/1987 Bicker et al.
4,701,440 A 10/1987 Grau
4,731,405 A 3/1988 Kirsch et al.
4,783,441 A 11/1988 Thurow
4,839,341 A * 6/1989 Massey et al. 514/4
4,885,164 A 12/1989 Thurow
4,959,351 A 9/1990 Grau
4,994,439 A 2/1991 Longenecker et al.
5,008,241 A 4/1991 Markussen et al.
5,034,415 A 7/1991 Rubin
5,101,013 A 3/1992 Dorschug et al.
5,177,058 A 1/1993 Dorschug
5,358,857 A 10/1994 Stengelin et al.
5,397,771 A 3/1995 Bechgaard et al.
5,428,006 A 6/1995 Bechgaard et al.
5,473,049 A 12/1995 Obermeier et al.
5,474,978 A 12/1995 Bakaysa et al.
5,496,924 A 3/1996 Habermann et al.
5,506,203 A 4/1996 Backstrom et al.

5,547,929 A 8/1996 Anderson, Jr. et al.
5,559,094 A 9/1996 Brems et al.
5,597,796 A 1/1997 Brange
5,614,219 A 3/1997 Wunderlich et al.
5,656,722 A 8/1997 Dorschug
5,663,291 A 9/1997 Obermeier et al.
5,693,608 A 12/1997 Bechgaard et al.
5,700,662 A 12/1997 Chance et al.
5,707,641 A 1/1998 Gertner et al.
5,783,556 A * 7/1998 Clark et al. 514/4
5,824,638 A * 10/1998 Burnside et al. 514/3
5,948,751 A 9/1999 Kimer et al.
5,985,309 A 11/1999 Edwards et al.
6,034,054 A 3/2000 DeFelippis et al.
6,043,214 A * 3/2000 Jensen et al. 514/3
6,051,551 A 4/2000 Hughes et al.
6,100,376 A 8/2000 Doerschug
6,174,856 B1 1/2001 Langballe et al.
6,211,144 B1 4/2001 Havelund
6,248,363 B1 6/2001 Patel et al.
6,267,981 B1 * 7/2001 Okamoto et al. 424/426
6,309,663 B1 10/2001 Patel et al.
6,310,038 B1 10/2001 Havelund
6,335,316 B1 1/2002 Hughes et al.
6,468,959 B1 10/2002 Wunderlich et al.
6,489,292 B1 12/2002 Havelund et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 62066/86 3/1987
AU 75916/87 1/1988
CA 1173388 8/1984
CA 1258427 8/1989
CA 1336329 7/1995
CA 1341203 3/2001

(Continued)

OTHER PUBLICATIONS

Brange, et al., Toward Understanding Insulin Fibrillation, Journal of Pharmaceutical Sciences, 86:517-525 (1997).
Schubert-Zsilavecz, et al., Insulin Glargin Ein Langwirksames Insulinanalogon, Pharmazie 2:125-130 (2001).
Sluzky, et al., Kinetics Of Insulin Aggregation In Aqueous Solutions Upon Agitation In The Presence Of Hydrophobic Surfaces, Proc. Natl. Acad. Sci. USA. vol. 88, pp. 9377-9381, Nov. 1991.

(Continued)

Primary Examiner—Jeffrey E Russel
(74) *Attorney, Agent, or Firm*—Balaram Gupta

(57) **ABSTRACT**

The invention relates to a pharmaceutical formulation comprising a polypeptide selected from the group consisting of insulin, an insulin metabolite, an insulin analog, an insulin derivative and combinations thereof; a surfactant or combinations of two or more surfactants; optionally a preservative or combinations of two or more preservatives; and optionally an isotonicizing agent, buffers or further excipients or combinations thereof, the pharmaceutical formulation having a pH in the acidic range.

U.S. PATENT DOCUMENTS

6,734,162	B2	5/2004	Van Antwerp et al.
6,818,738	B2	11/2004	Havelund
6,908,897	B2	6/2005	Brandenburg et al.
6,960,561	B2*	11/2005	Boderke 514/3
7,205,276	B2	4/2007	Boderke
7,205,277	B2	4/2007	Boderke
2001/0033868	A1	10/2001	Rossling et al.
2001/0039260	A1	11/2001	Havelund
2002/0107265	A1	8/2002	Chen et al.
2002/0198140	A1	12/2002	Havelund
2003/0004096	A1	1/2003	Boderke
2007/0155653	A1	7/2007	Boderke

FOREIGN PATENT DOCUMENTS

DE	2 219 635	11/1972
DE	3 240 177	5/1983
EP	0 018 609	11/1980
EP	0 046 979	3/1982
EP	0 166 529	1/1986
EP	0 180 920	5/1986
EP	0 194 864	9/1986
EP	0 200 383	12/1986
EP	0 211 299	2/1987
EP	0 224 885 A1	6/1987
EP	0 227 938	7/1987
EP	0 229 956	7/1987
EP	0 254 516	1/1988
EP	0 305 760	3/1989
EP	0 368 187	5/1990
EP	0 383 472	8/1990
EP	0419504	1/1994
EP	0 600 372	6/1994
EP	0214826	10/1994
EP	0 668 292	8/1995
EP	0678522	10/1995
EP	0 837 072	4/1998
EP	0375437	9/1998
EP	1 172 114	1/2002
FR	2456522	12/1980
GB	1527605 *	10/1978
WO	WO 83/00288	7/1982
WO	WO 88/06599	9/1988
WO	WO 90/07522	7/1990
WO	WO 91/16929	11/1991
WO	WO 92/00321	1/1992
WO	WO 94/14461	7/1994
WO	WO 96/04307	2/1996
WO	WO 96/07399	3/1996
WO	WO 96/11705	4/1996
WO	WO 96/41606	12/1996
WO	WO 97/01331	1/1997
WO	WO 98/42749	10/1998
WO	WO 98/56406	12/1998
WO	WO 99/24071	5/1999
WO	WO 00/23098	4/2000
WO	WO 00/23099	4/2000
WO	WO 00/29013	5/2000
WO	WO 00/74736	12/2000
WO	WO 01/00223	1/2001
WO	WO 01/12155	2/2001
WO	WO 01/21154	3/2001
WO	WO 01/28555	4/2001
WO	WO 01/37808	5/2001
WO	WO 01/43762	6/2001
WO	WO 01/52937	7/2001
WO	WO 01/93837	12/2001
WO	WO 02/064115	8/2002

WO WO 03/035051 5/2003

OTHER PUBLICATIONS

The Diabetes Control and Complications Trial Research Group, The Effect Of Intensive Treatment Of Diabetes On The Development And Progression Of Long-Term Complications In Insulin-Dependent Diabetes Mellitus, *The New England Journal of Medicine*, 329, 977-986 (1993).

Aoki, K, et al., Hydrolysis of Nonionic Surfactants, *Ann. Rept. Takeda Res. Lab.* 27, 172-176 (1968).

Garriques, L. N., et al., The Effect of Mutations on the Structure of Insulin Fibrils Studied by Fourier Transform Infrared (FTIR) Spectroscopy and Electron Microscopy, *Journal of Pharmaceutical Sciences*, vol. 91, No. 12 (2002), pp. 2473-2480.

Whittingham, J. L., et al., Insulin at PH2: Structural Analysis of the Conditions Promoting Insulin Fibre Formation, *J. Mol. Biol.*, (2002), vol. 318, pp. 479-490.

Bakaysa et al., "Physicochemical basis for the rapid time-action of LysB28ProB29-insulin: Dissociation of a protein-ligand complex?" *Protein Science*, 1996, 5:2521-31.

Beintema and Campagne, "Molecular Evolution of Rodent Insulins," *Mol. Biol. Evol.* 4(1): 10-18, 1987.

Berger, "Towards more physiological insulin therapy in the 1990s—A comment," *Diabetes Research and Clinical Practice*, 6(1989), pp. S25-S31.

Bolli, "The pharmacokinetic basis of insulin therapy in diabetes mellitus," *Diabetes Research and Clinical Practice*, 9(1989), pp. S3-S16.

Brange et al., "Monomeric insulins and their experimental and clinical implications," *Diabetes Care*, 13(9): 923-45 (1990).

Brange, "Galenics of Insulin" 1987, p. 35-36.

Burgermeister et al. "D: The Isolation of Insulin from the Pancreas," *Insulin*, Part 2, 1975, p. 715-727.

Burke et al., "Nature of the B10 amino acid residue," *Int. J. Peptide Protein Res.*, 23, 1984, p. 394-401.

Dixon et al., "Regeneration of Insulin Activity From the Separated and Inactive A and B Chains," *Nature*, vol. 188, No. 4752 (1960), pp. 721-724.

Drury et al., "Diabetic nephropathy," *British Medical Bulletin*, vol. 45, No. 1, 1989, pp. 127-147.

EP Search Report for Application 98110889.7-2105, Oct. 14, 1998.

Geiger, *Chem. Zeitung*, 100(3), p. 54-56. (Jan. 1976).

German Search Report for Application 19726167.1, Nov. 24, 1997.

Home et al., "Insulin treatment: a decade of change," *British Medical Bulletin*, 1989, vol. 45, No. 1, pp. 92-110.

Kang et al., "Subcutaneous Insulin Absorption Explained by Insulin's Physicochemical Properties-Evidence from Absorption Studies of Soluble Human Insulin and Insulin Analogues in Humans," *Diabetes Care*, vol. 14, No. 11, 1991, pp. 942-948.

Kemmler et al., "Studies on the Conversion of Proinsulin to Insulin," *The Journal of Biological Chemistry*, vol. 246, No. 22, 1971, pp. 6786-6791.

Kohner, "Diabetic retinopathy," *British Medical Bulletin*, vol. 45, No. 1, 1989, pp. 148-173.

Markussen et al., "Soluble, prolonged-acting insulin derivatives. I. Degree of protraction and crystallizability of insulins substituted in the termini of the B-chain," *Prot. Eng.* 1(3), 1987, pp. 205-213.

Markussen et al., "Soluble, prolonged-acting insulin derivatives. II. Degree of protraction and crystallizability of insulins substituted in positions A17, B8, B13, B27 and B30," *Prot. Eng.* 1(3), 1987, pp. 215-223.

Markussen et al., "Soluble, prolonged-acting insulin derivatives. III. Degree of protraction, crystallizability and chemical stability of insulins substituted in positions A21, B13, B23, B27 and B30," *Prot. Eng.* 2(2), 1988, pp. 157-166.

Müller et al., "Insulin Signaling in the Yeast *Saccharomyces cerevisiae*—I. Stimulation of Glucose Metabolism and snf 1 Kinase

Pillion et al., "Dodecylmaltoside-mediated Nasal and Ocular Absorption of Lyspro-Insulin: Independence of Surfactant from Multimer Dissociation," *Pharmaceutical Research*, vol. 15, No. 10, 1998, pp. 1637-1639.

Schartz et al., "A superactive insulin: [B10-Aspartic acid]insulin(human)," *Proc. Natl. Acad. Sci. USA*, v84, 1987, pp. 6408-6411.

Sundby, "Separation and Characterization of Acid-Induced Insulin Transformation Products by Paper Electrophoresis in 7 M Urea," *J. Biol. Chem.* 237(11), 1962, p. 3406-3411.

Volund et al., "In Vitro and In Vivo Potency of Insulin Analogues Designed for Clinical Use," *Diab. Med.* 8, 1991, p. 839-847.

Ward, "Diabetic neuropathy," *British Medical Bulletin*, vol. 45, No. 1, 1989, pp. 111-126.

Zinman, "The Physiologic Replacement of Insulin," *the New England J. Med.* 321(6), 1989, p. 363-370.

Thurrow and Geisen, "Stabilisation of dissolved proteins against denaturation at hydrophobic interfaces," *Diabetologia*, 27: 212-18 (1984).

Hinds et al., "Synthesis and Characterization of Poly(ethylene-glycol)-Insulin Conjugates," 2000, vol. 11, pp. 195-201.

Brange and Langkjaer, "Chemical stability of insulin—3. Influence of excipients, formulation and pH," *Acta Pharm. Nord.*, 1992, 4(3): 149-158.

Brange et al. "Neutral insulin solutions physically stabilized by addition of Zn²⁺," *Diabetic Medicine*, 1986, pp. 532-536.

Lougheed et al. "Physical Stability of Insulin Formulations," *Diabetes*, 1983, 32:424-32.

Kadima "Role of Metal Ions in the T- to R-Allosteric Transition in the Insulin Hexamer," *Biochem.* 38(41), 1999, p. 13443.

Nettleton et al. "Characterization of the Oligomeric States of Insulin in Self-Assembly and Amyloid Fibril Formation by Mass Spectrometry," *Biophysical J.*, v79, 2000, p. 1053-1065.

* cited by examiner

1

ACIDIC INSULIN PREPARATIONS HAVING IMPROVED STABILITY

This application is entitled to the benefit of U.S. Provisional Application 60/409,338, filed Sep. 9, 2002, and Federal Republic of Germany Application 10227232.8-41, filed Jun. 18, 2002.

SUMMARY OF THE INVENTION

The invention relates to a pharmaceutical formulation comprising a polypeptide selected from the group consisting of insulin, an insulin metabolite, an insulin analog, an insulin derivative or combinations thereof; a surfactant or combinations of two or more surfactants; optionally a preservative or combinations of two or more preservatives; and optionally an isotonicizing agent, buffers or further excipients or combinations thereof, the pharmaceutical formulation having a pH in the acidic range. These formulations can be employed for the treatment of diabetes, and are particularly suitable for preparations in which a high stability to thermal and/or physico-mechanical stress is necessary. The invention likewise relates to parenteral preparations which contain such formulations and can be used in diabetes and to methods for producing the preparations and for improving the stability of insulin preparations.

BACKGROUND OF THE INVENTION

Worldwide, approximately 120 million people suffer from diabetes mellitus. Among these, approximately 12 million are type I diabetics, for whom the substitution of the lacking endocrine insulin secretion is the only currently possible therapy. The affected persons are dependent lifelong on insulin injections, as a rule a number of times daily. In contrast to type I diabetes, there is not basically a deficiency of insulin in type II diabetes, but in a large number of cases, especially in the advanced stage, treatment with insulin, optionally in combination with an oral antidiabetic, is regarded as the most favorable form of therapy.

In the healthy person, the release of insulin by the pancreas is strictly coupled to the concentration of blood glucose. Elevated blood glucose levels, such as occur after meals, are rapidly compensated by a corresponding increase in insulin secretion. In the fasting state, the plasma insulin level falls to a basal value which is adequate to guarantee a continuous supply of insulin-sensitive organs and tissue with glucose and to keep hepatic glucose production low at night. The replacement of endogenous insulin secretion by exogenous, mostly subcutaneous administration of insulin, as a rule does not approximate the quality of the physiological regulation of the blood glucose described above. Often, deviations of blood glucose upward or downward occur, which in their severest forms can be life-threatening. In addition, however, blood glucose levels which are increased for years without initial symptoms are a considerable health risk. The large-scale DCCT study in the USA (The Diabetes Control and Complications Trial Research Group (1993) *N. Engl. J. Med.* 329, 977-986) demonstrated clearly that chronically elevated blood glucose levels are essentially responsible for the development of diabetic late damage. Diabetic late damage is microvascular and macrovascular damage which is manifested, under certain circumstances, as retinopathy, nephropathy or neuropathy and leads to loss of sight, kidney failure and the loss of extremities and is moreover accompanied by

2

the blood glucose as closely as possible in the physiological range. According to the concept of intensified insulin therapy, this should be achieved by repeated daily injections of rapid- and slow-acting insulin preparations. Rapid-acting formulations are given at meals in order to level out the postprandial increase in the blood glucose. Slow-acting basal insulins should ensure the basic supply with insulin, in particular during the night, without leading to hypoglycemia.

Insulin is a polypeptide of 51 amino acids, which are divided into 2 amino acid chains: the A chain having 21 amino acids and the B chain having 30 amino acids. The chains are connected to one another by means of 2 disulfide bridges. Insulin preparations have been employed for diabetes therapy for many years. Not only are naturally occurring insulins used, but recently also insulin derivatives and analogs.

Insulin analogs are analogs of naturally occurring insulins, namely human insulin or animal insulins, which differ by substitution of at least one naturally occurring amino acid residue with other amino acids and/or addition/removal of at least one amino acid residue from the corresponding, otherwise identical, naturally occurring insulin. The amino acids can in this case also be those which do not occur naturally.

Insulin derivatives are derivatives of naturally occurring insulin or an insulin analog which are obtained by chemical modification. This chemical modification can consist, for example, of the addition of one or more specific chemical groups to one or more amino acids. As a rule, insulin derivatives and insulin analogs have a somewhat modified action compared with human insulin.

Insulin analogs having an accelerated onset of action are described in EP 0 214 826, EP 0 375 437 and EP 0 678 522. EP 0 124 826 relates, inter alia, to substitutions of B27 and B28. EP 0 678 522 describes insulin analogs which in position B29 have various amino acids, preferably proline, but not glutamic acid. EP 0 375 437 includes insulin analogs with lysine or arginine in B28, which can optionally be additionally modified in B3 and/or A21.

In EP 0 419 504, insulin analogs are disclosed which are protected against chemical modifications, in which asparagine in B3 and at least one further amino acid in the positions A5, A15, A18 or A21 are modified.

In WO 92/00321, insulin analogs are described in which at least one amino acid of the positions B1-B6 is replaced by lysine or arginine. According to WO 92/00321, insulins of this type have a prolonged action. The insulin analogs described in EP-A 0 368 187 also have a delayed action.

The insulin preparations of naturally occurring insulins on the market for insulin substitution differ in the origin of the insulin (e.g. bovine, porcine, human insulin), and also the composition, whereby the profile of action (onset of action and duration of action) can be influenced. By combination of various insulin preparations, very different profiles of action can be obtained and blood sugar values which are as physiological as possible can be established. Recombinant DNA technology today makes possible the preparation of such modified insulins. These include insulin glargine (Gly(A21)-Arg(B31)-Arg(B32)-human insulin) with a prolonged duration of action. Insulin glargine is injected as an acidic, clear solution and precipitates on account of its solution properties in the physiological pH range of the subcutaneous tissue as a stable hexamer associate. Insulin glargine is injected once daily and is distinguished compared with other long-acting insulins by its flat serum profile and the reduction of the danger of nightly hypoglycemia associated therewith (Schubert-Zsilavec et al., 2: 125-130(2001)).

to previously described preparations, by a clear solution having an acidic pH. Especially at acidic pH, insulins, however, show a decreased stability and an increased proneness to aggregation on thermal and physicochemical stress, which can make itself felt in the form of turbidity and precipitation (particle formation) (Brange et al., J. Ph. Sci 86:517-525 (1997)).

The proneness to aggregation can additionally be promoted by hydrophobic surfaces which are in contact with the solution (Sluzky et al., Proc. Natl. Acad. Sci. 88:9377-9381 (1991)). Surfaces which can be considered as hydrophobic are the glass vessels of the preparations, the stopper material of the sealing caps or the boundary surface of the solution with the air supernatant. In addition, very fine silicone oil droplets can function as additional hydrophobic aggregation nuclei in the taking of the daily insulin dose by means of customary, siliconized insulin syringes and accelerate the process.

WO 01/43762 describes aqueous, parenteral pharmaceutical preparations comprising a polypeptide and glycerol, in which the stabilization of the preparation is to be achieved by purifying off destabilizing constituents of the glycerol.

WO 00/23098 describes insulin preparations stabilized using polysorbate 20 or poloxamer 188 for pulmonary administration, but does not describe the stabilization in an acidic solution against aggregation nuclei.

Published International patent application WO 02/076495 describes zinc-free and low-zinc insulin preparations having improved stability at room and body temperature and to mechanical stress by the addition of surfactants, but does not describe the stabilization of acidic insulin preparations against hydrophobic aggregation nuclei.

The present invention was thus based on the object of finding preparations for acid-soluble insulins containing surfactants, which are distinguished by a high long-term stability to stress due to temperature or physicochemical stressing and tolerate a high stress with hydrophobic aggregation nuclei.

DETAILED DESCRIPTION OF THE INVENTION

It has now surprisingly been found that the addition of surfactants can greatly increase the stability of acidic insulin preparations and thus preparations can be produced which guarantee superior stability to hydrophobic aggregation nuclei for several months under temperature stress.

The pharmaceutical preparations of the present invention contain 60-6000 nmol/ml, preferably 240-3000 nmol/ml, of an insulin, an insulin metabolite, an insulin analog or an insulin derivative.

The surfactants which can be used are, inter alia, nonionic surfactants. In particular, pharmaceutically customary surfactants are preferred, such as, for example: partial and fatty acid esters and ethers of polyhydric alcohols such as of glycerol, sorbitol and the like (Span®, Tween®, in particular Tween® 20 and Tween® 80, Myrj®, Brij®), Cremophor® or poloxamers. The surfactants are present in the pharmaceutical composition in a concentration of 5-200 µg/ml, preferably of 5-120 µg/ml and particularly preferably of 20-75 µg/ml.

The preparation can additionally optionally contain preservatives (e.g. phenol, cresol, parabens), isotonicizing agents (e.g. mannitol, sorbitol, lactose, dextrose, trehalose, sodium chloride, glycerol), buffer substances, salts, acids and alkalis and also further excipients. These substances can in each case be present individually or alternatively as mixtures.

Glycerol, dextrose, lactose, sorbitol and mannitol are cus-

150 mM. Buffer substances, such as, for example, phosphate, acetate, citrate, arginine, glycylglycine or TRIS (i.e. 2-amino-2-hydroxymethyl-1,3-propanediol) buffer and corresponding salts, are present in a concentration of 5-250 mM, preferably 10-100 mM. Further excipients can be, inter alia, salts or arginine.

The invention therefore relates to a pharmaceutical formulation comprising a polypeptide selected from the group consisting of insulin, an insulin analog, an insulin derivative, an active insulin metabolite and combinations thereof; a surfactant or combinations of two or more surfactants; optionally a preservative or combinations of two or more preservatives; and optionally an isotonicizing agent, buffer substances and/or further excipients or combinations thereof, the pharmaceutical formulation being a clear solution which has a pH in the acidic range (pH 1-6.8), preferably pH 3.5-6.8, very particularly preferably 3.5-4.5.

Preferred pharmaceutical formulations of the present invention are those wherein the surfactant is selected from the group consisting of partial and fatty acid esters and ethers of polyhydric alcohols such as of glycerol and sorbitol, and polyols; the partial and fatty acid esters and ethers of glycerol and sorbitol being selected from the group consisting of Span®, Tween®, Myrj®, Brij®, Cremophor®; the polyols being selected from the group consisting of polypropylene glycols, polyethylene glycols, poloxamers, Pluronic®, and Tetronics®; the preservative being selected from the group consisting of phenol, cresol, and parabens; the isotonicizing agent being selected from the group consisting of mannitol, sorbitol, sodium chloride, and glycerol; the excipients being selected from the group consisting of buffer substances, acids, and alkalis; the insulin analog being selected from the group consisting of Gly(A21)-Arg(B31)-Arg(B32)-human insulin; Lys(B3)-Glu(B29)-human insulin; Lys^{B28}Pro^{B29} human insulin, B28 Asp-human insulin, human insulin in which proline in position B28 has been substituted by Asp, Lys, Leu, Val or Ala and where in position B29 Lys can be substituted by Pro; AlaB26-human insulin; des(B28-B30)-human insulin; des(B27)-human insulin and des(B30)-human insulin; the insulin derivative being selected from the group consisting of B29-N-myristoyl-des(B30) human insulin, B29-N-palmitoyl-des(B30) human insulin, B29-N-myristoyl human insulin, B29-N-palmitoyl human insulin, B28-N-myristoyl Lys^{B28}Pro^{B29} human insulin, B28-N-palmitoyl-Lys^{B28}Pro^{B29} human insulin, B30-N-myristoyl-Thr^{B29}Lys^{B30} human insulin, B30-N-palmitoyl-Thr^{B29}Lys^{B30} human insulin, B29-N-(N-palmitoyl-γ-glutamyl)-des(B30) human insulin, B29-N-(N-lithocholy-γ-glutamyl)-des(B30) human insulin, B29-N-(ω-carboxyheptadecanoyl)-des(B30) human insulin and B29-N-(ω-carboxyheptadecanoyl) human insulin.

A further subject of the invention is a pharmaceutical formulation such as described above, in which the insulin, the insulin analog, the active insulin metabolite and/or the insulin derivative is present in a concentration of 60-6000 nmol/ml, preferably in a concentration of 240-3000 nmol/ml (this corresponds approximately to a concentration of 1.4-35 mg/ml or 40-500 units/ml);

in which the surfactant is present in a concentration of 5-200 µg/ml, preferably of 5-120 µg/ml and particularly preferably of 20-75 µg/ml.

A further subject of the invention is a pharmaceutical formulation such as mentioned above, in which glycerol and/or mannitol is present in a concentration of 100-250 mM, and/or

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.