

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2006/033672

International filing date: 28 August 2006 (28.08.2006)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/711,950
Filing date: 26 August 2005 (26.08.2005)


Date of receipt at the International Bureau: 20 November 2006 (20.11.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1540603



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

November 07, 2006

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/711,950
FILING DATE: August 26, 2005
RELATED PCT APPLICATION NUMBER: PCT/US06/33672

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US60/711,950



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

082605

18351 U.S. PTO

PTO/SB/16 (07-05)

Approved for use through 7/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EV 664382224 US

119277 U.S. PTO
60/711950

082605

INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
David C. William H. Daniel I.	YEOMANS FREY II JACOBS	Sunnyvale, California White Bear Lake, Minnesota Palo Alto, California
Additional inventors are being named on the _____ separately numbered sheets attached hereto		
TITLE OF THE INVENTION (500 characters max)		
THERAPY PROCEDURE FOR DRUG DELIVERY FOR TRIGEMINAL PAIN		
Direct all correspondence to: CORRESPONDENCE ADDRESS		
<input checked="" type="checkbox"/> The address corresponding to Customer Number:	<input type="text" value="25226"/>	
OR		
<input type="checkbox"/> Firm or Individual Name		
Address		
City	State	Zip
Country	Telephone	Email
ENCLOSED APPLICATION PARTS (check all that apply)		
<input checked="" type="checkbox"/> Application Data Sheet. See 37 CFR 1.76 (3 pages)	<input type="checkbox"/> CD(s), Number of CDs _____	
<input checked="" type="checkbox"/> Specification Number of Pages _____ 46	<input checked="" type="checkbox"/> Other (specify)	<input type="text" value="Return Receipt Postcard"/>
<input checked="" type="checkbox"/> Drawing(s) Number of Sheets _____ 1		
Application Size Fee: Filing Fee of \$200 (\$100 for small entity). If the specification and drawings exceed 100 sheets of paper, an application size fee due, which is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).		
METHOD OF PAYMENT OF FILING FEES AND APPLICATION SIZE FEE FOR THIS PROVISIONAL APPLICATION FOR PATENT		
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.	TOTAL FEE AMOUNT (\$)	
<input type="checkbox"/> A check or money order is enclosed to cover the filing fee and application size fee (if applicable).	<input type="text" value="100.00"/>	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.		
<input checked="" type="checkbox"/> The Director is hereby authorized to charge the filing fee and application size fee (if applicable) or credit any overpayment to Deposit Account Number: _____ 03-1952	A duplicative copy of this form is enclosed for fee processing.	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.		
<input checked="" type="checkbox"/> No.	<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____	

SIGNATURE diane pardi Date August 26, 2005

TYPED or PRINTED NAME Diane Pardi REGISTRATION NO. 56,824
(if appropriate)

TELEPHONE (650) 813-5636 Docket Number: 286003023500

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

[Number 1 of 1]

I hereby certify that this correspondence is being deposited with the U.S. Postal Service as Express Mail, Airbill No. EV 664382224 US, in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on the date shown below.

Dated: August 26, 2005 Signature: Georgina Matos (Georgina Matos)

pa-1005151

THERAPY PROCEDURE FOR DRUG DELIVERY FOR TRIGEMINAL PAIN

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] Not applicable.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] Not applicable.

TECHNICAL FIELD

[0003] The present invention relates generally to methods and compositions for the treatment of pain. More specifically, the present invention relates to methods for the treatment or prevention of trigeminal nerve-associated procedural, acute and chronic pain by administration and targeted delivery of analgesic agents to the trigeminal nerve system resulting in localized pain relief with minimal untoward central nervous system effects or systemic side effects.

BACKGROUND OF THE INVENTION

[0004] Pain is experienced when the free nerve endings which constitute the pain receptors in the skin as well as in certain internal tissues are subjected to mechanical, thermal, chemical or other noxious stimuli. The pain receptors (nociceptors) can transmit signals along afferent neurons into the central nervous system and then to the brain. The causes of pain can include inflammation, injury, disease, muscle spasm and the onset of a neuropathic event or syndrome. Ineffectively treated pain can be devastating to the person experiencing it by limiting function, reducing mobility, complicating sleep, and dramatically interfering with the quality of life.

[0005] The trigeminal sensory nerves (afferents) innervate and transmit to the brain most of the sensory signals from the face and anterior head. Pain involving the trigeminal nerve and ganglion arises in many different medical situations and presents unique problems to pain therapists and doctors. Chronic pain due to syndromes such as trigeminal neuralgia, atypical facial pain,

anesthesia dolorosa, post-herpetic neuralgia, cancer of the head and neck, migraine headaches, and temporomandibular joint pain are examples of very different pain syndromes that all involve the trigeminal system and which present clinical challenges that are peculiar to this nerve distribution. In addition to chronic pain states, there are clinical situations where facial and head pain is associated with acute trauma such as an abscessed tooth, a headache or a direct injury to the face and/or head such as a laceration or a burn. Further, medical procedures such as common dental work and facial plastic and/or cosmetic surgery may elicit considerable pain, as well as discomfort and anxiety.

[0006] Among syndromes associated with facial pain is Trigeminal neuralgia, also called “tic duloreaux” which is among the most common facial pain syndromes. Trigeminal neuralgia usually begins after the age of 40, is slightly more common in women and has an incidence of approximately 4-5 per 100,000 persons (Khorami and Totah, 2001). The primary symptom of trigeminal neuralgia is the sudden onset of severe, sharp facial pain, usually without warning. The quick bursts of pain are described as “lightening bolt-like”, “machine gun-like” or “electric shock-like”. The pain is generally on one side of the face and is spasmodic, coming in short bursts lasting a few seconds which may repeat many times over the course of a day. Trigeminal neuralgia can involve one or more branches of the trigeminal nerve and the causes are varied. Pharmacologic treatments include anti-seizure medications such as carbamazepine (Tegretol, Carbatrol), phenytoin (Dilantin), clonazepam (Klonopin), gabapentin (Neurontin), and lamtrigine (Lamictal), tricyclic antidepressants such as amitriptyline (Elavil) and muscle relaxants such as baclofen. The treatments generally have limited efficacy and many patients eventually undergo an invasive procedure. The procedural interventions often involve the direct manipulation of the trigeminal ganglion and include microvascular decompression, alcohol injection aimed at destroying pain fibers, glycerol injection aimed at selectively destroy pain-transmitting fibers, percutaneous radiofrequency rhizotomy, pulse radio frequency and gamma-knife. The pain relief from these procedures can be successful in a percentage of these patients, but the relief can be short-lived and often facial pain returns. Significant procedural pain and long term morbidity may also be associated with such treatments.

[0007] Atypical Facial Pain (ATFP) is a syndrome encompassing a wide group of facial pain problems. ATFP can have many different causes but the symptoms are all similar. Facial pain, often described as burning, aching or cramping, occurs on one side of the face, often in the region of

the trigeminal nerve and can extend into the upper neck or back of the scalp. Although rarely as severe as trigeminal neuralgia, facial pain is continuous for ATRP patients, with few, if any periods of remission. Some studies propose that ATRP is an early form of trigeminal neuralgia, but there is no agreement at this time. Drug treatments for ATRP are similar to what is prescribed for trigeminal neuralgia including anti-seizure medications and tricyclic antidepressants with limited effectiveness.

[0008] Anesthesia dolorosa is one of the most dreaded complications of neurosurgery and is considered to be non-reversible. The two main symptoms of anesthesia dolorosa are facial numbness (much like the numbness from a dental anesthetic injection) and constant pain. The pain is usually burning, pulling or stabbing but can also include a sharp, stinging, shooting or electrical component. Pressure and "heaviness" can also be part of the pain symptoms and often there is eye pain. Cold can increase the feeling of numbness sometimes making the face feel frozen. Anesthesia dolorosa occurs when the trigeminal nerve is damaged by surgery, physical trauma or as a complication of surgery to correct a condition such as trigeminal neuralgia. Topical treatments with capsaicin are used to help manage the pain and discomfort, while topical clonidine has been tested in a few cases but no single treatment has been found that resolves all of the pain of this condition.

[0009] Post-herpetic neuralgia is pain that remains after the rash from shingles (herpes zoster) has healed. Shingles is an infection of the nerves caused by the varicella-zoster virus, which is the same virus that causes chickenpox. About one-third of the people who get shingles will get post herpetic neuralgia. The pain of post herpetic neuralgia may be constant, stabbing, aching, or burning and can last for months to years after the shingles outbreak.

[0010] It is predicted that approximately 65,000 Americans will be diagnosed with head and neck cancers this year, this represents about 3% of all cancers diagnosed in the United States (American Cancer Society). Close to 60% of head and neck cancer patients report long-term pain with up to 25% claiming moderate or severe pain (List and Stracks (2000) *Curr Opin Oncol.*, 12:215-20). The trigeminal nerves and ganglion are likely to mediate most of the head and facial pain in these patients and sometimes are directly affected by the cancerous growth. The recommended treatment for most cancer patients with mild to severe pain is opioid therapy such as hydrocodone, codeine, oxycodone, morphine, fentanyl and hydromorphone. Opioid therapy has a multitude of problems including systemic effects away from the site of pain stimulation. Furthermore, opioids are highly addictive and patients build up tolerance to the drugs quickly resulting in higher and higher doses being administered.

[0011] Migraine headaches affect more than 23 million people in the United States. The typical migraine headache is throbbing or pulsatile, it builds up over a period of 1-2 hours and lasts from several hours to a whole day. Pain intensity is moderate to severe and can be debilitating and often causes nausea and vomiting. Of particular interest to clinicians who study migraine headaches is the superior trigeminal division (the ophthalmic division). This division innervates the forehead, eyebrow, eyelid, anterior scalp, nose and contents of the orbit thus giving an explanation for the pain localization along with the visual aura that is common with migraine headaches. Common treatments for migraine headaches include beta-blockers such as propranolol (Inderal) and Atenolol, tricyclic antidepressants, ergotamines, anti-seizure drugs and calcium channel blockers. Many of these drugs have systemic side effects and limited effectiveness.

[0012] Acute facial pain can arise in patients undergoing common dental procedures such as tooth extraction, root canal surgery and surgery for dental implants and dental prostheses. Acute dental pain can also arise from dental/gingival disease, other conditions such as an abscessed tooth or a bacterial infection or injury, that arise separately from planned dental procedures. Most dentists use topical anesthetics such as Benzocaine, Eugenol and forms of Xylocaine to numb various areas for minor procedures or before injection of a local anesthetic. For most procedures a dentist will inject a local anesthetic such as Lidocaine, Xylocaine and Marcaine to create a nerve block at or around the site where dental work needs to be done. Local anesthetics numb the area where they are injected and eliminate the acute pain of most procedures. In addition to the pain of administration, another main disadvantage of local anesthetics, especially for routine dental procedures, is that numbness and loss of sensation in the facial region will usually last for several hours after the dental procedure is finished.

[0013] Facial plastic surgery is becoming a very common procedure with several million procedures done in the United States each year. The procedures range from necessary repair of damage such as lacerations or broken bones to elective cosmetic surgeries such as face lifts, rhinoplasties, skin rejuvenation, etc. For many of these procedures local anesthetics are used (the patients are not under general anesthetic) and as with dental procedures, the local anesthetics can be painful to administer and include the problem of lingering numbness lasting for hours after the procedure is finished. In addition, depending on the surgery performed, patients experience varying levels of post-operative pain after the anesthetic wears off.

[0014] Pain treatment of almost any type usually includes some form of analgesic agent or drug. Analgesic drugs are usually classified into three groups: non-opioid drugs, opioid drugs, and co-analgesic drugs, also known as adjuvants. Non-opioid analgesic drugs include acetaminophen and non-steroidal anti-inflammatory drugs or NSAIDs. Opioid drugs, sometimes referred to as "narcotics", include natural substances such as morphine, and semi-synthetic and synthetic substances. Co-analgesic medications are drugs that have a primary use other than pain relief, but also help produce analgesia for some painful conditions.

[0015] Opioid drugs are commonly used to relieve pain however their usefulness is limited by the tolerance and dependence that normally develops on chronic treatment. Opioid drugs such as morphine can be addictive and can have central nervous system-mediated side effects such as respiratory and cardiac depressions and drowsiness. Additionally, opioid drugs suffer from frequent side effects such as nausea, vomiting and constipation.

[0016] Therapeutic drugs are delivered by a number of routes including, for example, oral, intravenous injection, intramuscular injection and subcutaneous injection. For patients suffering procedural, acute or chronic pain associated with the trigeminal nerve, one of the main problems with convention drug delivery with analgesic agents is the lack of localized pain relief due to systemic distribution of the agent. Often larger dosages need to be administered to achieve an effective concentration of the drug at a desired site. With higher doses of an analgesic agent, there is the additional problem of limited efficacy relative to the increase in undesired side effects due to the systemic distribution of the agent. Treatments consisting of localized but invasive interventions directly to the trigeminal nerve have a significant disadvantage due to the lack of selectivity and/or reversibility of the intervention and the fact that these procedures can, by themselves, cause additional facial nerve problems including anesthesia dolorosa, persistent numbness and nerve deafferentation. An additional problem with conventional treatments for trigeminal nerve-associated pain, especially with invasive procedures, is the high level of skill, training and equipment required by the medical team which can make treatment expensive and impractical for widespread use.

[0017] Intranasal administration has been used for systemic delivery of several therapeutic agents, for example, insulin, thyrotropin-releasing hormone, and vasopressin. In addition, using an intranasal or other mucosal route for systemic delivery of a therapeutic agent allows for ease of administration and the ability to bypass intestinal degradation and first pass hepatic metabolism of the therapeutic agent. However, there are times when it is desirable to not have systemic distribution

of a therapeutic agent or to have a therapeutic agent targeted to a localized or regional area. For examples, intranasal drug delivery has been used to bypass the blood-brain barrier and deliver substances to the central nervous system (CNS) and the brain. It has been demonstrated that large molecules such as polypeptides, peptides, oligonucleotides or DNA plasmids can be delivered directly to the CNS via specific uptake routes within the nose such as the axonal and perineural vascular/lymphatic pathways of the olfactory and trigeminal nerves (Frey II (2002) Drug Delivery Technology, 2:46-49; Thorne et al. (2004) Neuroscience, 127:481-496). Thus while there is evidence that various therapeutic agents can be delivered to the brain by an intranasal route and that the agents may travel along perineural pathways, there is no known method utilizing these pathways to specifically target the trigeminal nerve system for localized or regional analgesia in individuals suffering from trigeminal nerve-associated pain.

[0018] Despite a wide range of medical treatments, trigeminal nerve-associated pain, in many different forms and situations, continues to affect millions of people. Thus new methods for treating an individual for trigeminal nerve-associated pain are needed to directly target the trigeminal nerve system with analgesic agents and deliver analgesia to facial or head regions with minimal central nervous system effects or systemic side effects.

BRIEF SUMMARY OF THE INVENTION

[0019] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. Some aspects of the invention include methods wherein the trigeminal nerve-associated pain is selected from the group consisting of chronic, acute and procedural-related pain and combinations thereof. In some examples, the chronic pain is selected from the group consisting of trigeminal neuralgia, atypical facial pain, anesthesia dolorosa, post-herpetic neuralgia, cancer of the head and neck, migraine headaches, and temporomandibular joint pain. In some examples, the procedural-related pain is pain arising from dental, medical, surgical or cosmetic procedures. In yet other examples, the acute pain is pain arising from a laceration, a burn, a broken bone, an injury, a headache, an abscessed tooth, dental disease, a bacterial infection or a sinus infection.

[0020] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region and wherein the analgesic agent is administered via mucosal or dermal administration. In some examples the analgesic agent is administered intranasally. In other examples the analgesic agent is administered via buccal or sublingual administration. In other examples the analgesic agent is administered to conjunctiva or other mucosal tissues around the eye. In yet other examples the analgesic agent is administered to the skin or dermal surface.

[0021] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. Some aspects of the invention include methods wherein the analgesic agent includes, but is not limited to, a peptide, an amino acid, or a polypeptide. In some examples the analgesic agent is an opioid peptide selected from a group comprising enkephalins, endorphins, dynorphins, endomorphins, casomorphins, dermorphin, oxytocin and analogues and derivatives thereof. In some examples the analgesic agent is a peptide which inhibits peptidergic enzymes. In other examples the peptide is a peptidergic receptor agonist. In yet other examples the peptide is a peptidergic receptor antagonist. In further examples the analgesic agent is an antibody directed against proalgesic antigens such as endothelin, nerve growth factor, vasoactive intestinal polypeptide (VIP) or pituitary adenylate cyclase-activating polypeptide (PACAP). In some examples the analgesic agent is an antibody directed against calcitonin gene-related peptide (CGRP), cholecystokinin (CCK), Substance P or galanin.

[0022] Some aspects of the invention include methods wherein the analgesic agent is administered as a pharmaceutical composition. Accordingly, provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of a pharmaceutical composition comprising an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. Some aspects of the invention include methods wherein the pharmaceutical composition is administered in a dosage form selected from a group comprising a powder, a liquid, a gel, an ointment, a suspension, a cream or a bioadhesive. Some aspects of the invention include methods wherein the pharmaceutical composition further comprises a protease

inhibitor, an absorption enhancer, a vasoconstrictor or combinations thereof. In some examples, the protease inhibitor is selected from a group comprising antipain, arphamenine A and B, benzamidine HCl, AEBSF, CA-074, calpain inhibitor I and II, calpeptin, pepstatin A, actinonin, amastatin, bestatin, chloroacetyl-HOLeu-Ala-Gly-NH₂, DAPT, diprotin A and B, ebelactone A and B, foroxymithine, leupeptin, pepstatin A, phosphoramidon, aprotinin, BBI, soybean trypsin inhibitor, phenylmethylsulfonyl fluoride, E-64, chymostatin, 1,10-phenanthroline, EDTA and EGTA. In other examples the absorption enhancer is selected from a group comprising surfactants, bile salts, bioadhesive agents, phospholipid additives, mixed micelles, liposomes, or carriers, alcohols, enamines, cationic polymers, NO donor compounds, long-chain amphipathic molecules, small hydrophobic penetration enhancers; sodium or a salicylic acid derivatives, glycerol esters of acetoacetic acid, cyclodextrin or beta-cyclodextrin derivatives, medium-chain fatty acids, chelating agents, amino acids or salts thereof, N-acetylamino acids or salts thereof, mucolytic agents, enzymes specifically targeted to a selected membrane component, inhibitors of fatty acid synthesis and inhibitors of cholesterol synthesis.

[0023] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual i) an effective amount of a pharmaceutical composition comprising an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region and ii) a vasoconstrictor wherein administration of the vasoconstrictor reduces systemic distribution of the analgesic agent. In some examples the vasoconstrictor is selected from the group comprising phenylephrine hydrochloride, tetrahydrozoline hydrochloride, naphazoline nitrate, oxymetazoline hydrochloride, tramazoline hydrochloride, endothelin-1, endothelin-2, epinephrine, norepinephrine and angiotensin. In some examples the vasoconstrictor is administered prior to the administration of the pharmaceutical composition. In other examples the vasoconstrictor is co-administered with the pharmaceutical composition. In some examples administration of the vasoconstrictor results in a decreased effective dosage of the analgesic agent.

[0024] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of a pharmaceutical composition comprising an analgesic peptide wherein the peptide is administered by buccal or sublingual administration to the oral cavity and wherein the peptide preferentially binds to opioid receptors within the trigeminal nerve system and results predominantly in analgesia to the facial or

head region. Some aspects provide methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of a pharmaceutical composition comprising an analgesic peptide wherein the peptide is administered by transdermal administration to the skin and wherein the peptide preferentially binds to opioid receptors within the trigeminal nerve system and results predominantly in analgesia to the facial or head region

[0025] Provided herein are methods for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of a pharmaceutical composition comprising an analgesic peptide wherein the peptide is administered by intranasal administration to the nasal cavity and wherein the peptide preferentially binds to opioid receptors within the trigeminal nerve system and results predominantly in analgesia to the facial or head region. In some examples the administration is directed to the inferior two-thirds of the nasal cavity. In other examples the administration is directed to the inferior two-thirds of the nasal cavity and is directed away from the olfactory region.

BRIEF DESCRIPTION OF THE DRAWINGS

[0026] Figure 1 depicts data demonstrating withdrawal latencies after noxious thermal stimulation to the ears or hindpaws in a rat model after intranasal administration of met-enkephalin. Panel A shows baseline and treated withdrawal latencies after thermal stimulation to the ear. Panel B shows baseline and treated withdrawal latencies after thermal stimulation to the hindpaw. After taking baseline withdrawal latencies, rats were intranasally administered 10 nmoles/kg met-enkephalin and withdrawal latencies were retested. Each bar represents the average, across 4 animals, of latencies in response to stimulation at a particular time after the beginning of the set. Thus, the first white bar in each graph represents responses at the beginning of the baseline testing set; the first black bar represents responses at approximately five minutes after administering met-enkephalin. Each successive bar represents responses at approximately 15 minutes after the previous bar.

DETAILED DESCRIPTION OF THE INVENTION

[0027] As used herein, unless otherwise specified, the term “treatment” or “treating pain” refers to administration to an individual an agent of interest wherein the agent alleviates or prevents a pathology for which the subject is being treated. Treatment for trigeminal nerve-associated pain refers to the alleviation or prevention of trigeminal nerve-associated pain.

[0028] As used herein, “central nervous system” or “CNS” refers to that part of the nervous system that consists of the brain and spinal cord. The CNS is one of the two major divisions of the nervous system. The other is the peripheral nervous system which is outside of the brain and spinal cord and includes the cranial nerves - of which the trigeminal nerve is a member.

[0029] Although analgesia in the strictest sense is an absence of pain, as used herein, “analgesia” refers to reduction in the intensity of the pain perceived by an individual without causing general numbness.

[0030] As used herein, “analgesia agent”, “analgesic agent” or “analgesic” refers to any biomolecule that alleviates or prevents pain.

[0031] As used here, “analgesic peptide” refers to any peptide molecule that alleviates or prevents pain.

[0032] As used herein, “opioid peptide” refers to a peptide having a opioid receptor binding moiety and the capacity to bind to an opioid receptor. An opioid peptide can be a naturally occurring endogenous peptide, fragments, analogues or derivatives thereof. An opioid peptide can also be a non-endogenous peptide, fragments, analogues or derivatives thereof.

[0033] As used herein, “analogues and derivatives” refers to any peptide analogous to naturally occurring opioid peptides wherein one or more amino acids within the peptide have been substituted, deleted, or inserted. The term also refers to any peptide wherein one or more amino acids have been modified, for example by chemical modification. In general, the term covers all peptides which bind to an opioid receptor and exhibit an opioid activity but which may, if desired, have a different potency or pharmacological profile.

[0034] As used herein, “acute pain” refers to sudden, severe pain from a specific cause (injury, infection, inflammation, etc) that lasts a limited time (as opposed to chronic pain). As used herein “chronic pain” refers to a persistent state of pain whereby the cause of the pain cannot be easily removed. Chronic pain is often associated with long-term incurable or intractable medical

conditions or diseases. As used herein “procedural pain” refers to pain arising from a medical, dental or surgical procedure wherein the procedure is usually planned or associated with acute trauma.

[0035] As used herein “systemic side effects” include, but are not limited to, cardiovascular including peripheral vasodilation, reduced peripheral resistance, and inhibition of baroreceptors; dermatologic including pruritus (itching), flushing and red eyes; gastrointestinal including nausea and vomiting, decreased gastric motility in stomach, decreased biliary, pancreatic and intestinal secretions and delays in food digestion in small intestine, diminished peristaltic waves in large intestine contributing to constipation, epigastric distress or biliary colic in biliary tract; respiratory including depressed respiratory rate; and urinary including urinary urgency and difficulty with urination, and peripheral limb heaviness.

[0036] As used herein, “central nervous system effects” or “CNS effects” include, but are not limited to, narcosis, euphoria, drowsiness, apathy, psychotic ideation, mental confusion, alteration in mood, reduction in body temperature, feelings of relaxation, dysphoria (an emotional state characterized by anxiety, depression, or unease), nausea and vomiting (caused by direct stimulation of chemoreceptors in the medulla).

[0037] As used herein, “mucosal administration” or “administered transmucosally” refers to delivery to the mucosal surfaces of the nose, nasal passageways, nasal cavity; the mucosal surfaces of the oral cavity including the gingiva (gums), the floor of the oral cavity, the cheeks, the lips, the tongue, the teeth; and the mucosal surfaces of or around the eye including the conjunctiva, the lacrimal gland, the nasolacrimal ducts, the mucosa of the upper or lower eyelid and the eye.

[0038] As used herein, “intranasal administration” or “administered intranasally” refers to delivery to the nose, nasal passageways or nasal cavity by spray, drops, powder, gel, inhalant or other means.

[0039] The nasal cavity contains turbinate bones which protrude in to the nasal cavity and generally separate it into three regions. As used herein, the “inferior two-thirds of the nasal cavity” refers to the portion of the nasal cavity where the middle and inferior turbinate bones protrude and is the region of the nasal cavity that is innervated by the trigeminal nerve system. The superior third of the nasal cavity is defined by the superior turbinate bone wherein the olfactory region is located.

[0040] As used herein, “transdermal administration” or “dermal administration” refers to delivery to the skin of the face, neck, scalp or combinations thereof.

[0041] As used herein, “pharmaceutically acceptable carrier” or “suitable carrier” refers to a carrier that is conventionally used in the art to facilitate the storage, administration, and/or the healing effect of the agent.

[0042] As used herein, “therapeutically effective dose”, “therapeutically effective amount” or “an effective amount” refers to an amount of an analgesic agent that is useful for treating pain.

[0043] As used herein, “visual analogue scale” refers to a commonly used scale in pain assessment. It is a 10 cm horizontal or vertical line with word anchors at each end, such as “no pain” and “pain as bad as it could be”. A subject or patient is asked to make a mark on the line to represent pain intensity. This mark is converted to distance in either centimeters or millimeters from the “no pain” anchor to give a pain score that can range from 0-10 cm or 0-100 mm.

[0044] Described here are methods for treating an individual for trigeminal nerve-associated pain. In general, the methods are based on the finding that molecules can travel along perineural pathways to the trigeminal nerve and to the brain. Without wishing to be bound by theory, it is believed that analgesic agents can be targeted to the trigeminal nerve and that administration of the agents can result in analgesia and pain relief to an individual suffering acute, chronic or procedural facial or head pain. Furthermore, it is believed that targeted drug delivery to the trigeminal nerve system can limit systemic distribution of an analgesic agent which may decrease or eliminate undesirable central nervous system (CNS) effects or systemic side effects. In particular, it is believed that higher concentrations of an analgesic agent at a targeted site will allow for administration of lower dosages of the analgesic agent to the individual.

Analgesic Agents

[0045] The methods described herein can administer a variety of different analgesic agents to an individual for treatment of trigeminal nerve-associated pain. In general, the methods can administer to the trigeminal nerve an analgesic agent for prevention or treatment of facial or head pain. Administration of analgesic agents targeted for a predominantly regional analgesic effect can result in prevention or alleviation of pain without numbness as compared to local anesthetics. Since the trigeminal nerve transmits most of the sensory signals of the face and head, administration of analgesic agents targeted to the trigeminal nerve can localize the analgesic effect to the face and head region. Targeted delivery can also decrease the amount of agent administered to an individual to achieve an analgesic effect, and can decrease the undesirable CNS effects or systemic side effects

of many analgesic agents. More effective or efficient delivery of an analgesic agent to the trigeminal nerve can decrease the total dose of an agent administered to a subject suffering from trigeminal nerve-associated pain. Effective targeted delivery of an analgesic agent to the trigeminal nerve can decrease the systemic distribution of the agent wherein CNS effects or systemic side effects are minimized or eliminated.

[0046] Opioids are one of the classes of analgesic drugs commonly used for treatment of moderate to severe pain. These compounds include both plant-derived and synthetic alkaloids and also include endogenous peptides found in mammals as well as in lower animals. Examples of opioid analgesics include, but are not limited to, codeine, oxycodone, meperidine (Demerol), propoxyphene (Darvon), fentanyl, hydrocodone, tramadol (Ultram), morphine, hydromorphone (Dilaudid), methadone, levorphanol (Levo-Dromoran), oxymorphone (Numorphan), heroin, remifentanyl, butorphanol (Torbugesic), buprenorphine (Suboxone).

[0047] Naturally occurring endogenous opioid peptides can generally be referred to as “endorphins” (endogenous morphines) and include, but are not limited to, beta-endorphin, endomorphins, enkephalins, dynorphins, deltorphins, casomorphins, dermorphin and oxytocin.

[0048] Analgesic activity is mediated by opiate receptors found within the central nervous system and on peripheral neurons throughout the body. Opioid peptides bind to the same opiate receptors as narcotic opioid drugs. Both endogenous peptide opioids and narcotic morphine-like analgesics can alter the central release of neurotransmitters from afferent nerves sensitive to noxious, i.e. painful, stimuli. After binding with a receptor, opioid drugs or peptides may act to initiate or block various biochemical and physiological sequences.

[0049] Several major categories of opioid receptors are known: mu (μ), kappa (κ), delta (δ), and epsilon (ϵ). In general, mu-receptors mediate analgesia, euphoria, respiratory and physical depression, miosis, and reduced GI motility, delta-receptors mediate analgesia, dysphoria, psychotomimetic and respiratory effects and kappa-receptors mediate analgesia, sedation, miosis, respiratory depression and dysphoria. There is a differential distribution of these opiate receptors on nerves throughout the CNS and the peripheral neural system. In particular, mu and delta opioid receptors are found on the nociceptors in the trigeminal nerve but not on the olfactory nerve fibers within the nasal cavity. The differential distribution of opioid receptors can allow for targeted administration of opioid peptides to receptors within the trigeminal nerve while delivery to the olfactory nerve and the brain is minimized.

[0050] In some aspects of the invention are included methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. The analgesic agent can be a peptide, in particular an opioid peptide. The opioid peptide can be selected from a group comprising enkephalins, endorphins, α -neoendorphins, dynorphins, endomorphins, casomorphins, deltorphins, dermorphin, oxytocin and analogues and derivatives thereof. Further, the peptide can be targeted to opioid receptors on the trigeminal nerves and more than one opioid peptide can be administered.

[0051] The peptides for use in the herein described methods can be natural or synthetic, therapeutically or prophylactically active, peptide fragments, peptide analogues, and chemically modified derivatives or salts of active peptides. A variety of peptide analogues and derivatives are available and others can be contemplated for use within the invention and can be produced and tested for biological activity according to known methods. Peptides for use within the invention can be peptides that are obtainable by partial substitution, addition, or deletion of amino acids within a naturally occurring or native peptide sequence. Peptides can be chemically modified, for example, by amidation of the carboxyl terminus (-NH₂), the use of D amino acids in the peptide, incorporation of small non-peptidyl moieties, as well as the modification of the amino acids themselves (e.g. alkylation or esterification of side chain R-groups). Such analogues, derivatives and fragments should substantially retain the desired biological activity of the native peptide.

[0052] All peptides described and/or contemplated herein can be prepared by chemical synthesis using either automated or manual solid phase synthetic technologies, generally known in the art. The peptides can also be prepared recombinantly, using techniques known in the art.

[0053] A list of peptides is included in Table I, however one skilled in the art would know this list is not complete and one could contemplate and produce additional peptides, analogues and derivatives. Enkephalin was isolated from mammalian brains and found to be a mixture of two pentapeptides which differ only in the amino acid present at the 5-position. The two pentapeptides are methionine enkephalin (also known as met-enk or met-enkephalin) and leucine enkephalin (also known as leu-enk or leu-enkephalin). Met-enkephalin has an amino acid sequence of Tyr-Gly-Gly-Phe-Met and leu-enkephalin has an amino acid sequence of Tyr-Gly-Gly-Phe-Leu wherein the Tyr, Met and Leu residues are all L-amino acids. The tyrosine moiety is important for activity and probably corresponds to the 3-hydroxyl group on the morphine molecule. Proenkephalin A is the

precursor for met-enkephalin, leu-enkephalin and several other larger peptides. The structure of proenkephalin A contains four copies of met-enkephalin and one copy of leu-enkephalin, along with a heptapeptide (met-enk-Arg-Phe) and an octapeptide (met-enk-Arg-Gly-Leu). A series of peptides containing met-enkephalin at the N-terminus also possess opioid activity, and include peptide F and peptide E. Peptide F contains two met-enkephalin sequences one at each end, while peptide E contains met-enkephalin at the N-terminus and leu-enkephalin at the C-terminus.

[0054] In addition to naturally occurring enkephalins, enkephalin analogues and derivatives are known in the art, including, for example, derivatives and analogues which are specific for different types of opiate receptors. (Hruby and Gehrig (1989) Medicinal Research Reviews 9:343-401). Furthermore, enkephalin peptides can be modified by replacement and/or modification of specific amino acids, as it is known in the art that these modifications decrease the rate of hydrolysis and degradation of enkephalins by proteases.

[0055] β -endorphin is a 31-amino acid peptide formed from a larger precursor, pro-opiomelanocortin. β -endorphin contains a tetrapeptide sequence (Tyr-Gly-Gly-Phe) which is common to the enkephalin peptides and this tetrapeptide sequence appears to be essential to the function of these peptides. α -endorphin is a 16-amino acid peptide that is also formed from precursor pro-opiomelanocortin.

[0056] Dynorphins are another class of endogenous opioids that exist in multiple forms in the central nervous system. Dynorphins derive from precursor prodynorphin (proenkephalin B). Dynorphin, also known as Dynorphin A1-17, is a well-known opioid peptide that has the sequence Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile-Arg-Pro-Lys-Leu-Lys-Trp-Asp-Asn-Gln. Smaller peptides such as dynorphin A1-8 and three leu-enkephalins are also contained within proenkephalin B and are abundant in the neural lobe of the pituitary gland. Dynorphin A1-13 has been found in the striatonigral pathway and may provide a feedback mechanism for regulating dopaminergic activity in the striatum.

[0057] Endomorphins are amidated tetrapeptides and are structurally unrelated to the other endogenous opioid peptides. Two peptides, endomorphin-1 and endomorphin-2 have been isolated from mammalian brain. Both peptides have similar characteristics including analgesia against heat stimuli, mechanical stimuli and inflammatory and neuropathic pain. (Zadina et al. (1997) Nature 386:499-502; Hackler et al. (1997) Peptides 18:1635-1639)

[0058] Casomorphin peptides are novel opioid peptides derived from casein. Beta-casomorphins are the more extensively studied opioid peptides arising from the proteolytic breakdown of food proteins. The β -casomorphin peptide, Tyr-Pro-Phe-Pro-Gly-Pro-Ile was originally isolated from bovine beta-casein, and subsequently peptides with the same sequence were identified originating from ovine and buffalo beta-caseins. A human beta-casomorphin has been identified and has two amino acid differences from the bovine sequence, Tyr-Pro-Phe-Val-Glu-Pro-Ile. Several other casomorphin peptides have been isolated including β -casomorphin 1-3, β -casomorphin 1-4, β -casomorphin 1-5 and β -casomorphin 1-8.

[0059] Dermorphin is a seven amino acid peptide, originally isolated from *Phylomedusa sauvagei* frog skin. It is a ligand which binds with high affinity to the mu opioid receptor and has many biological roles including analgesia, endocrine modulation, immunomodulation, increased K⁺ conductance and inhibition of action potentials.

[0060] Oxytocin is a nine amino acid cyclic peptide hormone that is released from the posterior lobe of the pituitary gland and stimulates the contraction of smooth muscle of the uterus during labor and facilitates release of milk from the breast during nursing. Studies have shown that oxytocin can also play an important role in nociceptive modulation and that mu receptors appear to be the predominant receptor bound by oxytocin. (Wang et al. (2003) Regul Pept., 115:153-159; Zubrzycka et al. (2005) Brain Res. 1035:67-72).

[0061] Accordingly, in some aspects of the invention, the analgesic agent can be an opioid peptide selected from the group comprising leu-enkephalin, met-enkephalin, met-enk-Arg-Phe, met-enk-Arg-Gly-Leu, peptide E, peptide F, β -endorphin, α -endorphin, dynorphin A1-17, dynorphin B, beta-neoendorphin, α -neoendorphin, dynorphin A1-8, dynorphin A1-13, endomorphin-1, endomorphin-2, β -casomorphin, β -casomorphin 1-3, β -casomorphin 1-4, β -casomorphin 1-5, β -casomorphin 1-8, dermorphin, deltorphin I, deltorphin II, dermenkephalin, morphiceptin, oxytocin and analogues and derivatives thereof. In some examples more than one peptide is administered.

Table I

Name	Amino Acid Sequence	
Leu-enkephalin	Tyr-Gly-Gly-Phe-Leu	SEQ ID NO:1
Met-enkephalin	Tyr-Gly-Gly-Phe-Met	SEQ ID NO:2
Peptide F	Tyr-Gly-Gly-Phe-Met-Lys-Lys-Met-Asp-Glu-Leu-Tyr-Pro-Leu-Glu-Val-Glu-Glu-Glu-Ala-Asn-Gly-Gly-Phe-Val-Leu-Gly-Lys-Arg-Try-Gly-Gly-Phe-Met	SEQ ID NO:3
β -endorphin (human) GenBank 764134	Tyr-Gly-Gly-Phe-Met-Thr-Ser-Glu-Lys-Ser-Gln-Thr-Pro-Leu-Val-Thr-Leu-Phe-Lys-Asn-Ala-Ile-Ile-Lys-Asn-Ala-Tyr-Lys-Lys-Gly-Glu	SEQ ID NO:4
α -endorphin	Tyr-Gly-Gly-Phe-Met-Thr-Ser-Glu-Ser-Gln-Thr-Pro-Leu-Val-Thr-NH ₂	SEQ ID NO:5
Dynorphin A	Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile-Arg-Pro-Lys-Leu-Lys-Trp-Asp-Asn-Gln	SEQ ID NO:6
Dynorphin B	Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Gln-Phe-Lys-Val-Val-Thr	SEQ ID NO:7
α -neoendorphin	Tyr-Gly-Gly-Phe-Leu-Arg-Lys-Tyr-Pro-Lys	SEQ ID NO:8
β -neoendorphin	Tyr-Gly-Gly-Phe-Leu-Arg-Lys-Tyr-Pro	SEQ ID NO:9
Dynorphin A1-8	Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile	SEQ ID NO:10
Dynorphin A1-13	Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile-Arg-Pro-Lys-Leu-Lys	SEQ ID NO:11

Endomorphin-1	Tyr-Pro-Trp-Phe-NH ₂	SEQ ID NO:12
Endomorphin-2	Tyr-Pro-Phe-Phe-NH ₂	SEQ ID NO:13
Dermorphin	Tyr-(D)Ala-Phe-Gly-Tyr-Pro-Ser-NH ₂	SEQ ID NO:14
β -casomorphin (bovine)	Tyr-Pro-Phe-Pro-Gly-Pro-Ile	SEQ ID NO:15
β -casomorphin (human)	Tyr-Pro-Phe-Val-Glu-Pro-Ile	SEQ ID NO:16
β -Casomorphin 1-3	Tyr-Pro-Phe	SEQ ID NO:17
β -Casomorphin 1-4	Tyr-Pro-Phe-Pro	SEQ ID NO:18
β -Casomorphin 1-4, amide	Tyr-Pro-Phe-Pro-NH ₂	SEQ ID NO:19
β -Casomorphin 1-5	Tyr-Pro-Phe-Pro-Gly	SEQ ID NO:20
β -Casomorphin 1-8	Tyr-Pro-Phe-Pro-Gly-Pro-Ile-Pro	SEQ ID NO:21
Deltorphin I	Tyr-(D)Ala-Phe-Asp-Val-Val-Gly-NH ₂	SEQ ID NO:22
Deltorphin II	Tyr-(D)Ala-Phe-Glu-Val-Val-Gly-NH ₂	SEQ ID NO:23
Dermenkephalin	Tyr-(D)Met-Phe-His-Leu-Met-Asp-NH ₂	SEQ ID NO:24
Dermorphin	Tyr-(D)Ala-Phe-Gly-Tyr-Pro-Ser	SEQ ID NO:25
Morphiceptin	Tyr-Pro-Phe-Pro-NH ₂	SEQ ID NO:26
Oxytocin	Cys-Tyr-Ile-Gln-Asn-Cys-Pro-Leu-Gly	SEQ ID NO:27

[0062] Other analgesic or potentially analgesic agents can include amino acids, non-opioid peptides and polypeptides. These agents can include, but are not limited to, peptidergic channel modulators, peptidergic enzyme inhibitors, analgesic enzymes, trophic factors, peptidergic receptor agonists, peptidergic receptor antagonists, amino acid receptor agonists, nicotinic agonists, antibodies directed toward proalgesic antigens and antibodies directed to other neuropeptides. Channel modulators may include snail toxins and their derivatives and tetrodotoxin. Enzyme inhibitors may include cyclosporin A, bestatin, bestatin analogue Z4212 (N-[(2S, 3R)-3-Amino-2-hydroxy-4-(4-methylsulphonyl-phenyl)-1-oxobutyl]-1-aminocyclopentanecarboxylic) and bestatin analogue Z 1796 ((2S)-N-[(2S,3R)-3-Amino-2-hydroxy-4-(4-methylsulphonyl-phenyl)-1-oxobutyl]-L-leucine). Analgesic enzymes may include endothelin-1 peptidase. Trophic factors may include glial-derived neurotrophic factor (GDNF) and brain-derived neurotrophic factor (BDNF). Peptidergic receptor agonists may include somatostatin and its synthetic analogue octreotide, nocistatin, galanin and neuropeptide Y. Peptidergic receptor antagonists may include calcitonin gene-related peptide receptor antagonist CGRP(8-37), cholecystokinin (CCK) receptor antagonists such as Tyr-(D)Phe-Gly-(D)Trp-NMeNle-Asp-Phe-NH₂ or PD134308, neurokinin-1 receptor (substance P receptor) antagonists such as spantide II ((D)-NicLys1, 3-Pal3, D-Cl₂Phe5, Asn6, D-Trp7,9, Nle11-substance P), vasoactive intestinal peptide (VIP) receptor antagonists such as (Ac-Try1, D-Phe2)-GRF-(1-29) (where GRF is growth hormone releasing factor) and galanin receptor antagonists such as RWJ-57408). Amino acid receptor agonists may include gamma-amino butyric acid (GABA) and glycine. Nicotinic agonists may include nicotine and epibatidine. Antibodies directed toward proalgesic antigens may include antibodies to endothelin, nerve growth factor, vasoactive intestinal peptide (VIP) and pituitary adenylate cyclase-activating polypeptide (PACAP). Antibodies directed toward other neuropeptides may include antibodies to CGRP, CCK, substance P, and galanin. One skilled in the art would know this list is not complete and it is believed that one could contemplate and produce additional peptides, polypeptides, analogues and derivatives which have analgesic properties.

Administration

[0063] Chronic, acute or procedural pain associated with the trigeminal nerve system is experienced in many syndromes and diseases including, but not limited to, trigeminal neuralgia, atypical facial pain, anesthesia dolorosa, post-herpetic neuralgia, cancer of the head and neck,

migraine headaches, other types of headaches, temporomandibular joint pain, injuries to the face and/or head, injuries or infections of the teeth, common dental procedures and facial surgeries such as cosmetic plastic surgery. It is believed that analgesic agents can be targeted to the trigeminal nerve system and that this directed administration can result in analgesia and pain relief for a individual suffering acute, chronic or procedural facial or head pain.

[0064] The trigeminal nerve (fifth cranial nerve or CN V) is the largest of the 12 cranial nerves and it is the principal general sensory nerve to the head, particularly the face and is the motor nerve to the muscles of mastication. The trigeminal nerve innervates tissues of a mammal's (e.g. human) head including skin of the face and scalp, oral tissues and tissues of and surrounding the eye. The trigeminal nerve has three major branches or divisions: the ophthalmic, the maxillary, and the mandibular divisions. Thus, some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent wherein the administration of the analgesic agent is targeted to one or more of the three major branches of the trigeminal nerve including the ophthalmic, maxillary, and mandibular divisions.

[0065] The ophthalmic division is the superior division of the trigeminal nerve, it is the smallest of the three branches and is wholly sensory. The ophthalmic nerve has three branches known as the nasociliary nerve, the frontal nerve, and the lacrimal nerve which participate in the sensory supply to the skin of the forehead, upper eyelid and nose. The nasociliary nerve further divides into the anterior ethmoidal nerve and the infratrochlear nerve, while the frontal nerve divides into the supratrochlear and supraorbital nerves. The supratrochlear nerve supplies the middle part of the forehead, and the supraorbital nerve supplies the lateral part and the front of the scalp. The lacrimal nerve, supplies the lacrimal gland and the lateral part of the upper eyelid. Thus, the methods can administer an analgesic agent to the one or more of the nerves branching from the ophthalmic nerve including the nasociliary nerve, frontal nerve, lacrimal nerve, anterior ethmoidal nerve, infratrochlear nerve, supratrochlear nerve and supraorbital nerve.

[0066] The maxillary division is the intermediate division of the trigeminal nerve. It has three cutaneous branches: the infraorbital nerve which participates in the sensory supply to the skin on the lateral aspect of the nose, upper lip, and lower eyelid; the zygomaticofacial nerve, which supplies the skin of the face over the zygomatic bone; and the zygomaticotemporal nerve which supplies the skin over the temporal region. Thus, the methods can administer an analgesic agent to

the one or more of the nerves branching from the maxillary nerve including infraorbital, zygomaticofacial and zygomaticotemporal.

[0067] The mandibular division is the inferior division of the trigeminal nerve. It has three sensory branches: the buccal nerve supplies the skin of the cheek over the buccinator muscle. It also supplies the mucous membrane lining of the cheek and the posterior part of the buccal surface of the gingiva (gum). The auriculotemporal nerve supplies parts of the auricle, the external acoustic meatus, the tympanic membrane (eardrum) and the skin in the temporal region. The inferior alveolar nerve further divides into the incisive nerve and the mental nerve. The incisive nerve supplies the incisor teeth, the adjacent gingiva, and the mucosa of the lower lip and the mental nerve supplies the skin of the chin, the skin and mucosa membrane of the lower lip and gingiva. The lingual nerve supplies general sensory fibers to the anterior two-thirds of the tongue, the floor of the mouth, and the gingiva of the mandibular teeth. Thus the methods can administer an analgesic agent to one or more of the nerves branching from the mandibular nerve including buccal, auriculotemporal, inferior alveolar, incisive, mental and lingual.

[0068] Accordingly, some aspects of the invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent to mucosa tissue or epithelium within the oral cavity, within or around the eye or to the skin. The methods can include administering an agent to oral tissues wherein the analgesic agent is targeted to mucosal tissue innervated by a trigeminal division, for example the mandibular division. The oral mucosal tissues include, but are not limited to, the gingiva (gums), the floor of the oral cavity, the cheeks, the lips, the tongue, the teeth or a combination thereof. The methods can include administering an agent to conjunctiva or other mucosal tissues around the eye wherein the analgesic agent is targeted to mucosal tissue or epithelium innervated by a trigeminal division, for example the ophthalmic or maxillary division. The tissues or epithelium include, but are not limited to, the conjunctiva, the lacrimal gland, the nasolacrimal ducts, the mucosa of the upper or lower eyelid, the eye, or a combination thereof. An agent that is administered to the conjunctiva but not absorbed completely through the conjunctival mucosa can drain through the nasolacrimal ducts into the nose wherein it can be absorbed by mucosal tissue innervated by the trigeminal nerve within the nasal cavity. The methods can include administering an agent to skin of the face or head wherein the analgesic agent is targeted to tissue innervated by one of the trigeminal divisions. The agent can be administered to the skin of the face, scalp or temporal region. Suitable skin of the face includes skin

of the chin, the upper lip, the lower lip, the forehead, the nose, the cheek, the skin around the eyes, the upper eyelid, the lower eyelid or combinations thereof. Suitable skin of the scalp includes the front of the scalp, the scalp over the temporal region, the lateral part of the scalp, or combinations thereof. Suitable skin of the temporal region includes the temple and the scalp over the temporal region and combinations thereof.

[0069] Within the nasal cavity, the trigeminal nerve innervates mainly the inferior two-thirds of the nasal mucosa, while the olfactory nerve innervates the superior upper third of the nasal mucosa. There are primary afferents somatosensory neuronal fibers in the trigeminal nerve which allows for craniofacial somatosensory information including touch, temperature, proprioception (position sense) and pain. Those that are involved in nociception are termed “nociceptors”. In contrast, there are no nociceptors or other somatosensory primary afferents in the olfactory nerve which is essentially devoted to the sense of smell and pheromone detection. The anterior ethmoidal nerve, a branch of the nasociliary nerve, innervates, among other tissues, the ethmoidal sinus and regions of the inferior two-thirds of the nasal mucosa, including the anterior portion of the nasal septum and the lateral wall of the nasal cavity. The maxillary division has several branches that innervate the nasal cavity and sinuses, including the nasopalatine nerve, the greater palatine nerve, the posterior superior alveolar nerves, the middle superior alveolar nerve and the anterior superior alveolar nerve. The maxillary sinus is innervated by the posterior, middle and anterior superior alveolar nerves. The mucous membrane of the nasal septum is supplied chiefly by the nasopalatine nerve and the lateral wall of the nasal cavity is supplied by the greater palatine nerve.

[0070] Accordingly, some aspects of the invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent to mucosa tissue within the nasal cavity. The methods include administration of an analgesic agent to the inferior two-thirds of the nasal cavity wherein the analgesic agent is targeted to mucosal tissue innervated by the trigeminal nerve and away from the olfactory nerve. The methods include administration of an analgesic agent to the inferior two-thirds of the nasal cavity wherein the analgesic agent preferentially binds to opioid receptors within the trigeminal nerves. The methods can administer an analgesic agent to one or more of the nerves branching from the maxillary division that innervate the nasal cavity including nasopalatine, greater palatine, posterior superior alveolar, middle superior alveolar and anterior superior alveolar.

[0071] Within the oral cavity, the buccal or sublingual delivery routes are convenient choices for drug delivery as they are user-friendly and non-invasive. Some of the advantages include i) less proteolytic activity in the oral cavity as compared to some other routes thereby avoiding the problems of enzymatic degradation of peptide and protein drugs and ii) bypassing the liver first pass effect. In particular, buccal or sublingual delivery can allow for targeted delivery of an analgesic agent to the oral mucosa and to the trigeminal divisions that innervate the oral mucosa to treat or prevent trigeminal nerve-associated pain. Targeted delivery to a trigeminal division can reduce the effective dosage necessary to achieve analgesia in the facial or head regions wherein lower effective dosages will further reduce CNS effects or systemic side effects.

[0072] Drug delivery to the mucosal tissue around the eye or to the conjunctiva is another convenient choice for drug delivery that is non-invasive. In particular, administration to the mucosa or epithelium of the eyelids, the conjunctiva or the lacrimal system can allow for targeted delivery of an analgesic agent to the mucosa and tissues innervated by trigeminal divisions to treat or prevent trigeminal nerve-associated pain. Targeted delivery to a trigeminal division can reduce the effective dosage necessary to achieve analgesia in the facial or head regions wherein lower effective dosages will further reduce CNS effects or systemic side effects.

[0073] Transdermal drug delivery or administration of a therapeutic agent to the skin has become a proven technology over the last 20 years. Transdermal drug delivery offers controlled release of a drug to the patient and transdermal patches are user-friendly, convenient, painless, and offer multi-day dosing which usually results in improved patient compliance. Administration to the skin by transdermal delivery can allow for targeted delivery of an analgesic agent to the skin innervated by any one of the trigeminal divisions or a combination thereof to treat or prevent trigeminal nerve-associated pain in the facial or head regions.

[0074] Intranasal drug delivery has been a topic of research and development for many years, although it has been only within the past decade that carrier systems have been devised which make delivery of substances effective. (Sayani. and Chien (1996) Critical Reviews in Therapeutic Drug Carrier Systems, 13:85-184.)

[0075] Like the delivery systems discussed above, intranasal delivery of analgesic agents has a number of advantageous features including comparatively high bioavailability, rapid kinetics of absorption and avoidance of liver first-pass effect. In regard to patient compliance and ease of use, intranasal administration provides a simple, rapid and non-invasive mode of application. In

particular, intranasal delivery can allow for targeted delivery of an analgesic agent to the nasal cavity and to the trigeminal nerve to treat or prevent trigeminal nerve-associated pain. Furthermore, targeted delivery to the trigeminal nerve and preferably not the olfactory region can reduce the amount of drug entering the CNS or systemic circulation thereby reducing or eliminating CNS effects or systemic side effects. Targeted delivery to the trigeminal nerve can reduce the effective dosage necessary to achieve analgesia in the facial or head regions wherein lower effective dosages will further reduce CNS or systemic side effects.

[0076] In some aspects of the invention a vasoconstrictor is used to decrease systemic distribution of the analgesic agent. The vasoconstrictor can be included in a pharmaceutical composition to decrease systemic distribution of the analgesic agent. Alternatively, the vasoconstrictor may be delivered to the mucosal or dermal surface separately from the pharmaceutical composition. Vasoconstrictors are compounds that constrict blood vessels and capillaries and decrease blood flow. They can be used to increase concentration of an agent at a desired site by inhibiting movement of the analgesic agent into the bloodstream and thereby reducing systemic distribution of the agent. Vasoconstrictors can be used to decrease the effective dosage of agent needed to achieve analgesia by limiting systemic distribution and concentrating the agent in the trigeminal nerve. A vasoconstrictor can be administered before administration of the analgesic agent or can be co-administered with the analgesic agent. Vasoconstrictors may include, but are not limited to, phenylephrine hydrochloride, tetrahydrozoline hydrochloride, naphazoline nitrate, oxymetazoline hydrochloride, tramazoline hydrochloride, endothelin-1, endothelin-2, epinephrine, norepinephrine and angiotensin.

[0077] Accordingly, some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent by buccal or sublingual administration wherein the administration is targeted to a trigeminal division and results predominantly in analgesia to the facial or head region. The methods can administer an analgesic agent to the oral cavity of an individual to promote delivery to the trigeminal nerve with minimal systemic distribution. The methods can administer a vasoconstrictor to the oral cavity of an individual prior to administration of an analgesic agent to the oral cavity, wherein administration of the vasoconstrictor decreases systemic distribution of the analgesic agent thereby minimizing undesirable CNS effects or systemic side effects. The methods can co-administer a vasoconstrictor and an analgesic agent to the oral cavity of an individual. The methods

can administer a vasoconstrictor to the oral cavity of an individual prior to or co-administer with an analgesic agent, wherein administration of the vasoconstrictor decreases systemic distribution of the analgesic agent thereby decreasing the effective dosage amount of analgesic agent necessary to achieve analgesia to the facial or head region.

[0078] Some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an effective amount of an analgesic agent to the conjunctiva or other mucosal tissues around the eye wherein the administration is targeted to a trigeminal division and results predominantly in analgesia to the facial or head region.

[0079] Some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an effective amount of an analgesic agent to the skin of the face, head or scalp wherein the administration is targeted to the trigeminal nerve and results predominantly in analgesia to the facial or head region. In some examples, the analgesic agent is administered to particular sites on the face or scalp to promote delivery to particular trigeminal divisions.

[0080] Some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an analgesic agent by intranasal administration wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. The methods can administer an analgesic agent to the nasal cavity of an individual, in particular to the inferior two-thirds of the nasal cavity, to promote delivery to the trigeminal nerve with minimal delivery to the olfactory nerve. The methods can administer a vasoconstrictor to the nasal cavity of an individual prior to administration of an analgesic agent to the nasal cavity, wherein administration of the vasoconstrictor decreases systemic distribution of the analgesic agent thereby minimizing undesirable CNS effects or systemic side effects. The methods can co-administer a vasoconstrictor and an analgesic agent to the nasal cavity of an individual, wherein administration of the vasoconstrictor decreases systemic distribution of the analgesic agent thereby minimizing undesirable CNS or systemic side effects. The methods can administer a vasoconstrictor to the nasal cavity of an individual prior to or co-administer with an analgesic agent, wherein administration of the vasoconstrictor decreases systemic distribution of the analgesic agent thereby decreasing the

effective dosage amount of analgesic agent necessary to achieve analgesia to the facial or head region.

Pharmaceutical Composition

[0081] While it is possible to administer an analgesic agent alone, there are situations wherein it is advantageous to present it as part of a pharmaceutical composition. Thus, in some aspects of the present invention, the analgesic agent is administered as a pharmaceutical composition. The pharmaceutical composition can comprise an analgesic agent at a therapeutically effective dose together with one or more pharmaceutically acceptable carriers and optionally other ingredients. A suitable carrier is one which does not cause an intolerable side effect, but which allows the analgesic agent to retain its pharmacological activity in the body. A carrier may also reduce any undesirable side effects of the agent. A suitable carrier should be stable, i.e., incapable of reacting with other ingredients in the formulation. A suitable carrier should have minimal odor or fragrance or a positive (pleasant) odor. A suitable carrier should not irritate the mucosa, epithelium, underlying nerves or provide a health risk. It may be an accepted transcutaneous or percutaneous carrier or vehicle, because any carrier that can effectively penetrate the stratum corneum of the skin should be highly efficacious in not only penetrating mucosa, but also allowing rapid absorption of substances into the submucosal tissues, nerve sheaths and nerves.

[0082] Suitable nontoxic pharmaceutically acceptable carriers will be apparent to those skilled in the art of pharmaceutical formulations. Also see *Remington: The Science and Practice of Pharmacy*, 20th Edition, Lippincott, Williams & Wilkins (2000). Typical pharmaceutically acceptable carriers include, but are not limited to, mannitol, urea, dextrans, lactose, potato and maize starches, magnesium stearate, talc, vegetable oils, polyalkylene glycols, ethyl cellulose, poly(vinylpyrrolidone), calcium carbonate, chitosan, ethyl oleate, isopropyl myristate, benzyl benzoate, sodium carbonate, gelatin, potassium carbonate, silicic acid, and other conventionally employed acceptable carriers. Other carriers include, but are not limited to, phosphatidylcholine, phosphatidylserine, and sphingomyelins.

[0083] The choice of a suitable carrier will depend on the exact nature of the particular dosage form desired, e.g., whether the drug is to be formulated into a liquid solution (e.g., for use as drops, as a spray or impregnated in a nasal tampon, or other agent-impregnated solid), a suspension, a ointment or a gel. If desired, sustained-release compositions, e.g. sustained-release gels,

transdermal patches, etc. can be readily prepared. The particular dosage form will also depend on the route of administration. The agent can be administered to the nasal cavity as a powder, a granule, a solution, a cream, a spray, a gel, an ointment, an infusion, a drop or a sustained-release composition. For buccal administration, the composition can take the form of tablets or lozenges formulated in a conventional manner. For sublingual administration, the composition can take the form of a bioadhesive, a spray, paint or a swab applied to or under the tongue. For administration to the conjunctiva or other mucosal tissues around the eye, the composition can be applied as an ointment, a solution or a drop. For administration to the skin, the composition can be applied as a topical ointment, a topical gel, a lotion, a solution, a spray, a paint, a cosmetic, a patch or a bioadhesive.

[0084] Liquid carriers include, but are not limited to, water, saline, aqueous dextrose, and glycols particularly (when isotonic) for solutions. The carrier can be also be selected from various oils, including those of petroleum, animal, vegetable or synthetic origin, (e.g. peanut oil, soybean oil, mineral oil, sesame oil, and the like). Suitable pharmaceutical excipients include, but are not limited to, starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol, and the like. The compositions can be subjected to conventional pharmaceutical expedients, such as sterilization, and can contain conventional pharmaceutical additives, such as preservatives, stabilizing agents, reducing agents, anti-oxidants, chelating agents, wetting agents, emulsifying agents, dispersing agents, jelling agents, salts for adjusting osmotic pressure, buffers, and the like. Where the carrier is a liquid, it is preferred that the carrier be hypotonic or isotonic with body fluids and have a pH within the range of 4.5-8.5. Where the carrier is in powdered form, it is preferred that the carrier be within an acceptable non-toxic pH range. The use of additives in the preparation of peptide and/or protein-based compositions, particularly pharmaceutical compositions, is well-known in the art.

[0085] These lists of carriers and additives are by no means complete and a worker skilled in the art can choose excipients from the GRAS (generally regarded as safe) list of chemicals allowed in the pharmaceutical preparations and those that are currently allowed in topical and parenteral formulations. See also Wang et al., 1980, J. Parent. Drug Assn., 34:452-462, Wang et al., 1988, J. Parent. Sci. and Tech., 42:S4-S26.

[0086] Other forms of compositions for administration include a suspension of a particulate, such as an emulsion, a liposome, or in a sustained-release form to prolong the presence of the

pharmaceutically active agent in an individual. The powder or granular forms of the pharmaceutical composition may be combined with a solution and with a diluting, dispersing or surface-active agent. Additional compositions for administration include a bioadhesive to retain the agent at the site of administration, for example a spray, paint, or swab applied to the mucosa or epithelium. A bioadhesive can refer to hydrophilic polymers, natural or synthetic, which, by the hydrophilic designation, can be either water soluble or swellable and which are compatible with the pharmaceutical composition. Such adhesives function for adhering the dosage forms to the mucosal tissues of the oral or nasal cavity. Such adhesives can include, but are not limited to, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hydroxy ethylcellulose, ethylcellulose, carboxymethyl cellulose, dextran, gaur gum, polyvinyl pyrrolidone, pectins, starches, gelatin, casein, acrylic acid polymers, polymers of acrylic acid esters, acrylic acid copolymers, vinyl polymers, vinyl copolymers, polymers of vinyl alcohols, alkoxy polymers, polyethylene oxide polymers, polyethers, and combinations thereof. The composition can also be in the form of lyophilized powder, which can be converted into solution, suspension, or emulsion before administration. The pharmaceutical composition is preferably sterilized by membrane filtration and is stored in unit-dose or multi-dose containers such as sealed vials or ampoules.

[0087] The pharmaceutical composition can be formulated in a sustained-release form to prolong the presence of the active agent in the treated individual. Many methods of preparation of a sustained-release formulation are known in the art and are disclosed in Remington's Pharmaceutical Sciences (see above). Generally, the agent can be entrapped in semi-permeable matrices of solid hydrophobic polymers. The matrices can be shaped into films or microcapsules. Matrices can include, but are not limited to, polyesters, co-polymers of L-glutamic acid and gamma ethyl-L-glutamate, polylactides, polylactate polyglycolate, hydrogels, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers, hyaluronic acid gels, and alginic acid suspensions. Suitable microcapsules can also include hydroxymethylcellulose or gelatin and poly-methyl methacrylate. Microemulsions or colloidal drug delivery systems such as liposomes and albumin microsphere can also be used. Some sustained-release compositions can use a bioadhesive to retain the agent at the site of administration.

[0088] To further enhance the mucosal delivery of a pharmaceutical composition comprising an analgesic agent, an enzyme inhibitor, particularly proteases inhibitors, can be included in the formulation. Protease inhibitors may include, but are limited to, antipain, arphamenine A and B,

benzamidine HCl, AEBSF, CA-074, calpain inhibitor I and II, calpeptin, pepstatin A, actinonin, amastatin, bestatin, boroleucine, captopril, chloroacetyl-HOLeu-Ala-Gly-NH₂, DAPT, diprotin A and B, ebelactone A and B, foroxymithine, leupeptin, pepstatin A, phosphoramidon, aprotinin, puromycin, BBI, soybean trypsin inhibitor, phenylmethylsulfonyl fluoride, E-64, chymostatin, 1,10-phenanthroline, EDTA and EGTA.

[0089] To enhance delivery into or across a mucosal surface and/or absorption of a pharmaceutical composition comprising an analgesic agent, an absorption-enhancing agent can be included in the formulation. These enhancing agents may enhance the release or solubility (e.g., from a formulation delivery vehicle), diffusion rate, penetration capacity and timing, uptake, residence time, stability, effective half-life, peak or sustained concentration levels, clearance and other desired mucosal delivery characteristics (e.g., as measured at the site of delivery) of the composition. Enhancement of mucosal delivery can thus occur by any of a variety of mechanisms, for example by increasing the diffusion, transport, persistence or stability of an analgesic agent, increasing membrane fluidity, modulating the availability or action of calcium and other ions that regulate intracellular or paracellular permeation, solubilizing mucosal membrane components (e.g., lipids), changing non-protein and protein sulfhydryl levels in mucosal tissues, increasing water flux across the mucosal surface, modulating epithelial junctional physiology, reducing the viscosity of mucus overlying the mucosal epithelium, reducing mucociliary clearance rates, and other mechanisms.

[0090] Mucosal absorption enhancing compounds may include, but are not limited to, surfactants, bile salts, dihydrofusidates, bioadhesive agents, phospholipid additives, mixed micelles, liposomes, or carriers, alcohols, enamines, cationic polymers, NO donor compounds, long-chain amphipathic molecules, small hydrophobic penetration enhancers; sodium or a salicylic acid derivatives, glycerol esters of acetoacetic acid, cyclodextrin or beta-cyclodextrin derivatives, medium-chain fatty acids, chelating agents, amino acids or salts thereof, N-acetylamino acids or salts thereof, mucolytic agents, enzymes specifically targeted to a selected membrane component, inhibitors of fatty acid synthesis and inhibitors of cholesterol synthesis.

[0091] These additional agents and compounds can be coordinately administered or combinatorially formulated with the analgesic agents. Accordingly, some aspects of the present invention include methods wherein the analgesic agent is administered as a pharmaceutical composition that comprises protease inhibitors, absorption enhancers, vasoconstrictors or

combinations thereof. The pharmaceutical composition can be administered to the nasal cavity, oral cavity, to conjunctiva or other mucosal tissues around the eye or to the skin. The pharmaceutical composition can be administered by an intranasal route. The pharmaceutical composition can be administered by a buccal or sublingual route. The pharmaceutical composition can be administered by a transdermal route. The pharmaceutical composition can include at least one protease inhibitor, at least one absorption enhancer, at least one vasoconstrictor or combinations thereof. The pharmaceutical composition can be co-administered with a vasoconstrictor or administered after the vasoconstrictor has been delivered.

Delivery Systems

[0092] An analgesia agent or pharmaceutical composition may be dispensed to the buccal or sublingual surfaces in a number of different dosage forms including, but not limited to, fast-melting tablets, liquid-filled capsules, liquid sprays or lozenges. Alternatively, the pharmaceutical composition can be delivered to the mucosa of the oral cavity by direct placement of the composition in the mouth, for example, with a gel, an ointment, a dropper, or a bioadhesive strip or patch.

[0093] In some aspects of the present invention, the methods comprise administering to an individual a pharmaceutical composition wherein administration to the buccal and/or sublingual mucosal surfaces of the oral cavity is by a delivery device. The delivery device can include, but is not limited to, unit dose containers, pump sprays, droppers, squeeze bottles, airless and preservative-free sprays, nebulizers, dose inhalers and pressurized dose inhalers. The delivery device can be metered to administer an accurate effective dosage amount (as described below) to the oral cavity. In some aspects, an accurate effective dosage amount is contained within a capsule, tablet, lozenge, or bioadhesive patch that is placed directly within the oral cavity.

[0094] An analgesia agent or pharmaceutical composition may be dispensed to the conjunctiva or to other mucosal tissues around the eye in a number of different dosage forms such as a liquid drop, a gel, an ointment or a bioadhesive patch or strip. Thus, in some aspects of the present invention the methods comprise administering to an individual a pharmaceutical composition wherein administration is directed to the conjunctiva or other mucosal tissues around the eye. In some aspects, an accurate effective dosage amount is contained within a drop, a gel, an ointment or a bioadhesive patch that is placed directly onto the mucosal tissues around the eye.

[0095] An analgesia agent or pharmaceutical composition may be administered to the skin or scalp in a number of different dosage forms such as a liquid, a spray, a gel, an ointment or a bioadhesive patch or strip. Thus, in some aspects of the present invention the methods comprise administering to an individual a pharmaceutical composition wherein administration is directed to the skin of the face or scalp. In some aspects, an accurate effective dosage amount is contained within a drop, a gel, an ointment or a bioadhesive transdermal patch that is placed directly onto the skin.

[0096] An analgesia agent or pharmaceutical composition may be dispensed intranasally as a powdered or liquid nasal spray, suspension, nose drops, a gel or ointment, through a tube or catheter, by syringe, by packtail, by pledget (a small flat absorbent pad), by nasal tampon or by submucosal infusion. Nasal drug delivery can be carried out using devices including, but not limited to, unit dose containers, pump sprays, droppers, squeeze bottles, airless and preservative-free sprays, nebulizers (devices used to change liquid medication to an aerosol particulate form), metered dose inhalers, and pressurized metered dose inhalers. It is important that the delivery device protect the drug from contamination and chemical degradation. The device should also avoid leaching or absorption as well as provide an appropriate environment for storage. Each drug needs to be evaluated to determine which nasal drug delivery system is most appropriate. Nasal drug delivery systems are known in the art and several are commercially available.

[0097] The composition may be conveniently delivered in the form of an aerosol spray using a pressurized pack or a nebulizer and a suitable propellant including, but not limited to, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, hydrocarbons, compressed air, nitrogen or carbon dioxide. An aerosol system requires the propellant to be inert towards the pharmaceutical composition. In the case of a pressurized aerosol, the dosage unit may be controlled by providing a valve to deliver an accurately metered amount.

[0098] The means to deliver the analgesic agent to the nasal cavity as a powder can be in a form such as microspheres delivered by a nasal insufflator device (a device to blow a gas, powder, or vapor into a cavity of the body) or pressurized aerosol canister. The insufflator produces a finely divided cloud of the dry powder or microspheres. The insufflator may be provided with means to ensure administration of a substantially metered amount of the pharmaceutical composition. The powder or microspheres should be administered in a dry, air-dispensable form. The powder or microspheres may be used directly with an insufflator which is provided with a bottle or container

for the powder or microspheres. Alternatively the powder or microspheres may be filled into a capsule such as a gelatin capsule, or other single dose device adapted for nasal administration. The insufflator can have means such as a needle to break open the capsule or other device to provide holes through which jets of the powdery composition can be delivered to the nasal cavity.

[0099] Nasal delivery devices can be constructed or modified to dispense the pharmaceutical composition wherein the composition is delivered predominantly to the inferior two-thirds of the nasal cavity. For example, the angle of dispersion from a delivery device such as a nebulizer or an insufflator can be set so that the pharmaceutical composition is mechanically directed to the inferior two-thirds of the nasal cavity, and preferably away from the superior region of the nasal cavity. Alternatively, the pharmaceutical composition can be delivered to the inferior two-thirds of the nasal cavity by direct placement of the composition in the nasal cavity, for example, with a gel, an ointment, a nasal tampon, a dropper, or a bioadhesive strip.

[0100] Thus in some aspects of the present invention, the methods comprise administering to an individual a pharmaceutical composition wherein administration to the nasal cavity is by a nasal delivery device. The nasal delivery device can include, but is not limited to, unit dose containers, pump sprays, droppers, squeeze bottles, airless and preservative-free sprays, nebulizers, dose inhalers, pressurized dose inhalers, insufflators, and bi-directional devices. The nasal delivery device can be metered to administer an accurate effective dosage amount (as described below) to the nasal cavity. The nasal delivery device can be for single unit delivery or multiple unit delivery. In some aspects of the present invention, the nasal delivery device can be constructed whereby the angle of dispersion of a pharmaceutical composition is mechanically directed towards the inferior two-thirds of the nasal cavity thereby minimizing delivery to the olfactory region. The nasal delivery device can be constructed whereby the angle of dispersion of a pharmaceutical composition is mechanically directed towards the inferior two-thirds of the nasal cavity thereby maximizing delivery of the agent to opioid receptors in the trigeminal nerve. In some aspects of the present invention, the pharmaceutical composition is a gel, cream, ointment, impregnated in a nasal tampon or bioadhesive strip whereby the composition is placed in the inferior two-thirds of the nasal cavity. In some aspects of the present invention, the methods include intranasal administration of an analgesia agent wherein the administration uses a nasal delivery device with an angle of dispersion that mechanically directs the agent to the inferior two-thirds of the nasal cavity wherein the analgesic agent is administered after a vasoconstrictor. In some aspects of the present invention, the methods

include intranasal administration of an analgesia agent wherein the administration uses a nasal delivery device with an angle of dispersion that mechanically directs the agent to the inferior two-thirds of the nasal cavity wherein the analgesic agent is co-administered with a vasoconstrictor.

Dosages

[0101] An analgesic agent is administered in a dose sufficient to provide a therapeutically effective amount to the trigeminal nerve that results predominantly in analgesia to the facial or head region of an individual suffering from trigeminal nerve-associated pain. In particular, the analgesic agent can be administered in a dose that results in analgesia to the facial or head regions with minimal CNS effects or systemic side effects. A therapeutically effective dose of an analgesic agent can be determined empirically and depends on the analgesic agent, the type and severity of the pain, the route of administration, the state of disease progression, and the size/weight and overall health of the patient. In particular, a therapeutically effective dose of an analgesic agent which results in regional analgesia with minimal CNS effects or systemic side effects can be determined empirically and will depend on the same described parameters.

[0102] The amount of an analgesic agent administered as a unit dose will depend upon the type of pharmaceutical composition being administered, for example, a solution, a suspension, a gel, an emulsion, a powder, or a sustained-release formulation. Generally the effective dosage will be lower than dose amounts needed for oral, intravenous, intramuscular or subcutaneous administration of the analgesic agent involved, since targeted delivery will allow for a more concentrated level of the analgesic agent in the trigeminal nerve. The effective dosage will be lower than dosage amounts generally used for other common analgesic opioid drugs, for example, morphine. The quantity of dosage form needed to deliver the desired dose will depend on the concentration of the analgesic agent in the composition. Such determinations are within the skill of one in the art.

[0103] The therapeutic dosage of an analgesic agent in the pharmaceutical compositions used in the methods of the present invention will depend on a number of factors such as the particular analgesic agent chosen, its bioavailability by the chosen route of administration, its efficacy, and the desired frequency of administration combined with the desired single dosage of the formulation. Particularly, dosage of the analgesic agent will be chosen to maximize analgesia to the facial and head regions and minimize CNS effects or systemic side effects. Such pharmacological data can be

obtained from animal models and clinical trials with normal human volunteers or patients experiencing trigeminal nerve-associated pain by one with skill in the art.

[0104] Experimental models to test for analgesic activity of agents are known in the art. Animal models comprise tests which include, but are not limited to, acetic acid writhing, phenylquinone writhing, tail-flick, paw withdrawal and ear withdrawal wherein the pain receptor activation is induced by such compounds as acetic acid, phenylquinone, formalin or capsaicin, or by thermal activators such as a hot plate or a laser. In particular, models for facial or head pain utilizing tests such as orofacial delivery of capsaicin, orofacial delivery of formalin, or delivery of thermal heat to the ear are available. Models can be used to determine optimal dosage ranges wherein an analgesic agent delivered to the trigeminal nerve results in analgesia in the facial or head region with minimal analgesia at a systemic site, i.e. the paw. Further, models can be used to administer an analgesic agent by a particular delivery route, e.g. intranasally, and test for analgesic effect at the ears and at the hindpaws. Thus, one model can be used to test for analgesic activity of an analgesic agent after administration of a pharmaceutical composition to the trigeminal nerve. Withdrawal latencies at the ear will determine localized analgesia while withdrawal latencies at the hindpaw will determine systemic distribution and analgesia.

[0105] As stated above, an effective amount of an analgesic agent will depend on the analgesic agent being used in the method. Preferably the effective amount of an analgesic agent administered transmucosally or transdermally to the trigeminal nerve is lower than dosages used when the agent is delivered by other routes (e.g. oral, intravenous, intramuscular or subcutaneous). For example, dosages used for administration of an enkephalin peptide can include, but are not limited to, an effective amount within the dosage range of about 0.01 ng per kg body weight to about 50 µg per kg body weight, or within 0.1 ng per kg body weight to about 50 µg per kg body weight, or within 1 ng per kg body weight to about 50 µg per kg body weight, or within about 10 ng per kg body weight to about 50 µg per kg body weight, or within about 0.1 µg per kg body weight to about 50 µg per kg body weight, or within about 1 µg per kg body weight to about 50 µg per kg body weight. Dosages used for administration of an endorphin peptide can include, but are not limited to, an effective amount within the dosage range of about 0.4 µg per kg body weight to about 4 mg per kg body weight, or within 4 µg per kg body weight to about 400 µg per kg body weight, or within 4 µg per kg body weight to about 200 µg per kg body weight, or within 10 µg per kg body weight to about 100 µg per kg body weight. Dosages used for administration of an endomorphin

peptide can include, but are not limited to, an effective amount within the dosage range of about 0.15 nmol per kg body weight to about 1.5 μ mol per kg body weight, or within 1.5 nmol per kg body weight to about 150 nmol per kg body weight, or within 1 nmol per kg body weight to about 100 nmol per kg body weight, or within 1 nmol per kg body weight to about 50 nmol per kg body weight. Dosages used for intranasal administration of a dynorphin peptide can include, but are not limited to, an effective amount within the dosage range of about 10 nmol per kg body weight to about 100 μ mol per kg body weight, or within 100 nmol per kg body weight to about 50 μ mol per kg body weight, or within 250 nmol per kg body weight to about 25 μ mol per kg body weight, or within 0.5 μ mol per kg body weight to about 5 μ mol per kg body weight.

[0106] Dosages can be administered in a single dose or in multiple doses, for example, dosages can be administered two, three, four, up to ten times daily depending on the analgesic agent and the type of pain being treated. Dosages can be administered in a sustained release formulation which allows for the analgesic agent to be administered less frequently such as six times a week, five times a week, four times a week, three times a week, twice a week, or once a week.

[0107] Thus some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain comprising administering to the individual an effective amount of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region with minimal CNS effects or systemic side effects. The analgesic agent can be an enkephalin with a dosage range of about 0.01 ng per kg body weight to about 50 μ g per kg body weight, or within 0.1 ng per kg body weight to about 50 μ g per kg body weight, or within 1 ng per kg body weight to about 50 μ g per kg body weight, or within about 10 ng per kg body weight to about 50 μ g per kg body weight, or within about 0.1 μ g per kg body weight to about 50 μ g per kg body weight, or within about 1 μ g per kg body weight to about 50 μ g per kg body weight. The analgesic agent can be an endorphin with a dosage range of about 0.4 μ g per kg body weight to about 4 mg per kg body weight, or within 4 μ g per kg body weight to about 400 μ g per kg body weight, or within 4 μ g per kg body weight to about 200 μ g per kg body weight, or within 10 μ g per kg body weight to about 100 μ g per kg body weight.

Therapeutic Uses

[0108] Chronic pain in the face and head region can arise from a variety of medical conditions including but not limited to neuropathic pain, headache pain, TMJ, pain from cancer

and/or cancer treatment. These pain syndromes are often not effectively treated with current medications or invasive interventions and new methods for localized pain relief in the face and head regions are needed. Accordingly, some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated chronic pain by administration of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region. The analgesic agent can be administered to a patient with neuropathic pain, including but not limited to, trigeminal neuralgia, atypical facial neuralgia and post herpetic neuralgia. The analgesic agent can be administered to an individual with headache pain, for example, migraine headaches or cluster headaches. The analgesic agent can be administered to an individual with chronic pain arising from head or facial cancer or arising from previous treatment of head or facial cancer.

[0109] Local anesthesia is used for most medical, dental or cosmetic procedures wherein the procedures last a short period of time and wherein the work is isolated to the teeth, face or head. Anesthesia is generally defined as the total or partial loss of sensation, especially tactile sensibility, usually induced by use of an anesthetic compound. In addition to the pain of administration, another main disadvantage of local anesthesia is that loss of sensation results in numbness which often lasts significantly longer than the procedure. Thus there are situations wherein patients undergoing medical, dental or cosmetic procedures would benefit from short-term regional analgesia of the teeth, face or head regions. Some procedures potentially would require no other pain relieving agents or any vasoconstrictors, and the length of time of analgesia, in the absence of numbness, would be much less important than the excessive length of time of facial numbness resulting from an anesthetic. Such medical, dental and cosmetic procedures can include, but are not limited to, microdermabrasion, Botox injection, photodynamic therapy or other skin tumor ablations, hair removal (including electrolysis, laser, waxing, etc.), general facial laser treatments (including pigment removal, vascular lesions), dermal and subdermal injectable fillers (including collagen, hyaluronic acid, methylmethacrylate, hydroxyapatite, etc), facial peels by chemical or laser applications, photofacials, collagen shrinkage procedures (including radiofrequency, HIFU, high intensity light, laser, etc.), routine dental procedures, tattooing, tattoo removal, piercing and treatment of scars, keloids, etc. by steroid injection.

[0110] Thus some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain arising from medical, dental or cosmetic procedures comprising

administration of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia of the facial or head regions. The methods can include an analgesic agent administered to an individual undergoing a procedure selected from the group comprising medical, dental and cosmetic. The methods can include medical, dental or cosmetic procedures selected from the group comprising microdermabrasion, Botox injection, photodynamic therapy or other skin tumor ablations, hair removal (including electrolysis, laser, waxing, etc.), general facial laser treatments (including pigment removal, vascular lesions), dermal and subdermal injectable fillers (including collagen, hyaluronic acid, methylmethacrylate, hydroxyapatite, etc), facial peels by chemical or laser applications, photofacials, collagen shrinkage procedures (including radiofrequency, HIFU, high intensity light, laser, etc.), dental procedures, tattooing, tattoo removal, piercing and treatment of scars and keloids by steroid injection. The analgesic agent can be administered to a patient undergoing a procedure wherein the analgesic effect lasts the length of the procedure. The analgesic agent can be administered to a patient undergoing a procedure wherein the time requirement for the procedure and for analgesia is less than 30 minutes, is less than 45 minutes, is less than 60 minutes or is less than 90 minutes. The analgesic agent can be administered to a patient undergoing a procedure wherein the time requirement for the procedure and for analgesia is more than 90 minutes.

[0111] For medical, dental or cosmetic procedures that are more extensive and will take a longer period of time, a local anesthetic is used usually in combination with a sedative to make the patient drowsy. In some cases, depending on the procedure, the patient is put under a general anesthetic. The use of local or general anesthesia does not effectively mitigate post-operative pain and analgesics are almost always delivered to the patient after the procedure is complete.

[0112] An evolving concept in operative pain management is the use of preemptive analgesia. The pain and inflammation that result from surgery normally causes increased prostaglandin production and sensitization. If analgesic agents are administered before surgery the amount of sensitization may be decreased or prevented and the degree and persistence of post-operative pain may be diminished. Therefore, there are situations wherein patients undergoing medical, dental or cosmetic procedures would benefit from regional trigeminal or facial analgesia prior to the procedure, during the procedure and during the post-operative period. Some procedures typically benefit by the use of vasoconstrictors, for example, before plastic surgery to decrease the amount of bleeding at an incision site. Therefore one level of benefit from facial analgesia would be

the elimination of pain associated with an injection of a vasoconstrictor, wherein the time required for analgesia could be less than 10 minutes. Another level of benefit exists if the facial analgesia starts before the surgery starts, lasts throughout the entire procedure and continues into the post-operative period, wherein the post-operative period could represent hours or days. Examples of dental procedures may include, but are not limited to, major dental procedures such as periodontal, reconstructive, palatal, tooth extraction, root canal surgery, etc. Examples of cosmetic or medical surgical procedures may include, but are not limited to, facelift, blepharoplasty, browlift, rhinoplasty, cheek implant, chin implant, fat injections, lesion removal, excisional biopsies, Mohs surgery (micrographic surgery for skin cancer), flap reconstruction, orthognathic (correction of jaw deformities), ophthalmic and oculoplastic (plastic surgery of the eye), hair replacement surgery, extensive laser resurfacing, trauma such as laceration repair, nasal fracture repair, facial bone fracture repair, burn debridement and wound cleaning.

[0113] Accordingly, some aspects of the present invention include methods for treating an individual for trigeminal nerve-associated pain arising from medical, dental or cosmetic procedures comprising administration of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results in localized analgesia of the face, head or teeth. The methods can include an effective dosage amount wherein the localized analgesia lasts for the length of the procedure and continues into a post-operative period. The methods can include medical, dental or cosmetic procedures selected from the group comprising periodontal surgery, reconstructive tooth surgery, palatal surgery, tooth extraction, root canal surgery, facelifts, blepharoplasties, browlifts, rhinoplasties, cheek implants, chin implants, fat injections, lesion removal, excisional biopsies, Mohs surgery, flap reconstruction, orthognathic surgery, ophthalmic surgery, oculoplastic surgery, hair replacement surgery, extensive laser resurfacing, laceration repair, nasal fracture repair, facial bone fracture repair, burn debridement and wound cleaning. The analgesic agent can be administered to a patient undergoing a medical procedure prior to injection of a vasoconstrictor into the facial or head region. The analgesic agent can be administered to a patient undergoing a medical procedure wherein the analgesia lasts beyond the length of the procedure and into a post-operative time period. The analgesic agent can be administered to a patient undergoing a medical procedure wherein the analgesia lasts for hours to days after the medical procedure is finished.

EXAMPLES

EXAMPLE 1

[0114] One way to test activity of an analgesic agent in a rat model is by treatment-induced changes in latencies (times) of withdrawal in response to noxious heating of the skin, typically using an ear or a hindpaw. Thus, application of coherent or non-coherent (non-laser) radiant heat to the ear or hindpaw will elicit rapid withdrawal movements. Latencies of withdrawal have been demonstrated to be sensitive to analgesic treatments, such that analgesics increase the latency to withdrawal. Transmucosal or transdermal administration of analgesic agents to the trigeminal nerve to reduce trigeminal nerve-associated pain can be tested for regional and/or systemic analgesia. The rostral external part of a rat's ear is innervated by a branch of the mandibular nerve, itself a branch of the trigeminal nerve, thus after treatment an increase in latency to withdrawal time would indicate regional analgesia. A change in the latency to withdrawal time of the hindpaw would indicate whether there was a systemic analgesic effect, i.e. no change in the latency to withdrawal time indicates no systemic effect, while an increase in latency to withdrawal time would indicate a systemic effect.

[0115] Rats are housed in a 12/12-hour light/dark environment and are provided food and water ad libitum. Efforts are made to minimize discomfort and reduce the number of animals used. Rats are lightly anesthetized with urethane and placed with minimal restraint on a heating pad to maintain their body temperature at 37 °C. A laser beam is directed via a fiberoptic cable to the rostral external part of both ears. Characteristic responses to laser irradiation are a retraction or withdrawal of the stimulated ear for 1-3 seconds after a thermal stimulus by the laser. Laser stimulation is terminated rapidly after response of the stimulated ear or after a maximal response (cut-off) latency of 30 seconds to prevent tissue damage.

[0116] For baseline testing of latency withdrawal responses to the ear, 3 pulses are applied to each ear. The stimulation site is changed after each pulse allowing at least 2 minutes in between 2 stimuli on the same ear. For baseline testing of latency withdrawal responses to the hindpaw, 3 pulses are applied to the hindpaw. The stimulation site is changed after each pulse allowing at least 2 minutes in between 2 stimuli on the same hindpaw. Testing sessions are videotaped for off-line

analysis of responses. The off-line analysis is performed by an investigator who determines the latency of withdrawal responses to the laser stimulation and who is blinded to the treatment groups.

[0117] After measuring baseline latencies, analgesic agents are administered intranasally. This involves 5 equal 10 μ l applications to the nose by pipette for a total volume of 50 μ l over 20 minutes. The effect of different doses of an agent (e.g. 10 nmoles/kg met-enkephalin) on latency responses is examined. To assess the local analgesic effect, the latency responses of the ear are tested at various time points after agent administration. To assess the systemic analgesic effect, the latency responses of hindpaws are tested at various timepoints after agent administration.

EXAMPLE 2

[0118] Sprague-Dawley rats (Charles River Laboratories) were lightly anesthetized with urethane and placed with minimal restraint on a heating pad to maintain their body temperature at 37 °C. A laser beam was directed via a fiberoptic cable to the rostral external part of both ears or to the hindpaws as described above. Baseline withdrawal latencies were measured by delivering 4 separate stimuli with a resting period of approximately 15 minutes between each stimulus. 50 μ l of met-enkephalin in phosphate-buffered saline was intranasally administered in 5 equal 10 μ l applications at a dosage of 10 nmoles/kg of body weight. Withdrawal latencies for both ears and hindpaws were tested five minutes after the final application of met-enkephalin. As described above, testing sessions were videotaped and analyzed. Results demonstrated that intranasal administration of met-enkephalin at this dosage achieved a regional analgesic effect in the head region without a systemic analgesic effect at the hindpaw. (Figure 1)

EXAMPLE 3

Normal human volunteers.

[0119] Regional analgesia in the face region after administration of analgesic agents by intranasal delivery can be tested in normal subjects. Study participants are selected based on inclusion/exclusion criteria, history and physical exam, laboratory tests, and other customary procedures. Thermal pain responses are elicited on the face, in particular the cheek, and on the hand of healthy normal volunteers, such that temperature thresholds for evoking pain and/or the temperature of maximal pain tolerance can be assessed and baselines established. Increasing doses of an analgesic agent are administered to the subjects and a dose-response curve is calculated for

each stimulation site. Changes in thermal pain threshold and tolerance at the two sites can be compared so that the efficacy of an analgesic agent at a given dose in affecting facial and whole-body pain can be determined.

[0120] The analgesic agent is delivered intranasally to the subjects by a metered dose nebulizer. For example, a dose of 0.01 $\mu\text{g}/\text{kg}$ of oxytocin in 0.1 ml of saline is administered with each nasal puff application to the subjects. 0.1 ml of saline only with each nasal puff application is administered to control subjects. It is determined what doses of an analgesic agent administered to the trigeminal nerve are effective for establishing regional analgesia in the facial region (i.e. the cheek) with minimal systemic distribution and minimal or no analgesic effect at the peripheral site (i.e. the hand).

EXAMPLE 4

Human patients.

[0121] Patients undergoing cosmetic facial surgery normally experience significant post-operative pain. The patients are treated with an analgesic agent which is administered intranasally by metered dose nebulizer at the cessation of surgery. Patients receive a dose of a test agent (e.g. 0.01 $\mu\text{g}/\text{kg}$ of oxytocin) in 0.1 ml of normal saline or they receive a placebo of saline alone. A patient's facial pain ratings are then determined on the visual analogue scale at 10 minute intervals for 2 hours. A second set of patients undergoing a similar surgical procedure to the hand also receive an intranasal application of either the test agent or saline placebo. Patient's hand pain ratings are then determined on the visual analogue scale at 10 minute intervals for 2 hours.

[0122] Although the invention has been described with respect to particular aspects and examples for purposes of clarity of understanding, it will be apparent to those skilled in the art that various changes and modifications can be made without departing from the invention. Therefore, the descriptions and examples should not be construed as limiting the scope of the invention.

CLAIMS

We claim:

1. A method for treating an individual for trigeminal nerve-associated pain, comprising: administering to the individual an effective amount of an analgesic agent wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia to the facial or head region.
2. The method of claim 1, wherein the analgesic agent is administered via mucosal administration, wherein mucosal administration comprises intranasal administration, buccal administration, sublingual administration and conjunctival administration.
3. The method of claim 1, wherein the analgesic agent is administered via transdermal administration, wherein transdermal administration comprises administering the agent to the skin of the face, forehead, eyelids, nose, cheek, chin, scalp, temple region or a combination thereof.
4. The method of claim 1, wherein the analgesic agent is administered via intranasal administration.
5. The method of claim 1, wherein the analgesic agent is administered via buccal or sublingual administration.
6. The method of claim 1, wherein the analgesic agent is administered via conjunctival administration or via other tissues around the eye.
7. The method of claims 1, 2 or 3, wherein the analgesic agent comprises a peptide.
8. The method of claim 7, wherein the peptide is selected from the group comprising enkephalins, endorphins, dynorphins, endomorphins, casomorphins, dermorphin, oxytocin and analogues and derivatives thereof.
9. The method of claims 1, 2 or 3, wherein the analgesic agent comprises an agent selected from the group comprising peptidergic channel modulators, peptidergic enzyme inhibitors, analgesic enzymes, trophic factors, peptidergic receptor agonists, peptidergic receptor antagonists, amino acid receptor agonists, nicotinic agonists, antibodies directed toward proalgesic antigens and antibodies directed to other neuropeptides.
10. The method of claim 1, wherein the trigeminal nerve-associated pain is selected from the group consisting of chronic, acute and procedural-related pain and combinations thereof.

11. The method of claim 10, wherein the chronic pain is selected from the group consisting of trigeminal neuralgia, atypical facial pain, anesthesia dolorosa, post-herpetic neuralgia, cancer of the head and neck, migraine headaches, and temporomandibular joint pain.

12. The method of claim 10, wherein the procedural-related pain is pain arising from dental, medical, surgical or cosmetic procedures.

13. The method of claim 10, wherein the acute pain is pain arising from a laceration, a burn, a broken bone, a headache, a dental disease, a bacterial infection, an abscessed tooth, or a sinus infection.

14. The method of claim 1, wherein the analgesic agent is administered as a pharmaceutical composition.

15. The method of claim 14, wherein the pharmaceutical composition is administered as a powder, a gel, an ointment, a liquid, a suspension, a cream or a bioadhesive.

16. The method of claim 14, wherein the pharmaceutical composition further comprises at least one protease inhibitor or at least one absorption enhancer.

17. The method of claim 14, wherein the pharmaceutical composition further comprises at least one protease inhibitor and at least one absorption enhancer.

18. The method of claim 1, wherein the method further comprises administration of a vasoconstrictor.

19. The method of 18, wherein the vasoconstrictor is administered prior to the analgesic agent.

20. The method of 18, wherein the vasoconstrictor is co-administered with the analgesic agent.

21. The method of 18, wherein administration of the vasoconstrictor reduces systemic distribution of the analgesic agent.

22. The method of 21, wherein reduced systemic distribution allows for a decreased effective dosage of the analgesic agent to achieve analgesia.

23. The method of claim 1, wherein the method further comprises administration of at least one protease inhibitor and at least one absorption enhancer.

24. The method of claim 1, wherein the method further comprises administration of at least one protease inhibitor, at least one absorption enhancer and a vasoconstrictor.

25. The method of claim 16, 17, 23 and 24, wherein the protease inhibitor is selected from a group comprising antipain, arphamenine A and B, benzamidine HCl, AEBSF, CA-074, calpain inhibitor I and II, calpeptin, pepstatin A, actinonin, amastatin, bestatin, chloroacetyl-HOLeu-Ala-Gly-NH₂, DAPT, diprotin A and B, ebelactone A and B, foroxymithine, leupeptin, pepstatin A, phosphoramidon, aprotinin, BBI, soybean trypsin inhibitor, phenylmethylsulfonyl fluoride, E-64, chymostatin, 1,10-phenanthroline, EDTA and EGTA.

26. The method of claim 16, 17, 23 and 24, wherein the absorption enhancer is selected from a group comprising surfactants, bile salts, bioadhesive agents, phospholipid additives, mixed micelles, liposomes, or carriers, alcohols, enamines, cationic polymers, NO donor compounds, long-chain amphipathic molecules, small hydrophobic penetration enhancers; sodium or a salicylic acid derivatives, glycerol esters of acetoacetic acid, cyclodextrin or beta-cyclodextrin derivatives, medium-chain fatty acids, chelating agents, amino acids or salts thereof, N-acetylamino acids or salts thereof, mucolytic agents, enzymes specifically targeted to a selected membrane component, inhibitors of fatty acid synthesis and inhibitors of cholesterol synthesis.

27. The method of claim 18, wherein the vasoconstrictor is selected from a group comprising phenylephrine hydrochloride, tetrahydrozoline hydrochloride, naphazoline nitrate, oxymetazoline hydrochloride, tramazoline hydrochloride, endothelin-1, endothelin-2, epinephrine, norepinephrine and angiotensin.

28. A method for treating an individual for trigeminal nerve-associated pain, comprising: administering to an individual an effective amount of an analgesic peptide wherein the peptide is administered by transmucosal or transdermal administration and wherein the peptide preferentially binds to opioid receptors within the trigeminal nerve system and results predominantly in analgesia to the facial or head region.

29. A method for treating an individual for trigeminal nerve-associated pain, comprising: administering to an individual an effective amount of an analgesic peptide wherein the administration is targeted to the trigeminal nerve system and results predominantly in localized analgesia to the facial or head region with minimal central nervous system effects or systemic side effects.

30. A method for treating an individual for trigeminal nerve-associated pain, comprising: administering to an individual an effective amount of an analgesic peptide to the nasal cavity of the

individual wherein the administration is targeted to the trigeminal nerve system and results predominantly in analgesia localized to the facial or head region.

31. The method of claim 30, further comprising the administration of a vasoconstrictor.



32. The method of claim 30 or 31, wherein the administration is directed to the inferior two-thirds of the nasal cavity.

33. The method of claim 30 or 31, wherein the administration is directed to the inferior two-thirds of the nasal cavity and away from the olfactory region.

ABSTRACT

The present invention relates to methods for the treatment or prevention of trigeminal nerve-associated pain, in particular chronic, acute and procedural-related pain. The methods comprise administration of analgesic agents to the trigeminal nerve which results in analgesia to the facial or head region.

FIGURE 1



Application Data Sheet

Application Information

Application Type::	Provisional
Subject Matter::	Utility
Suggested Group Art Unit::	Not Yet Assigned
CD-ROM or CD-R?::	None
Sequence submission?::	None
Computer Readable Form (CRF)?::	No
Title::	THERAPY PROCEDURE FOR DRUG DELIVERY FOR TRIGEMINAL PAIN
Attorney Docket Number::	286003023500
Request for Early Publication?::	No
Request for Non-Publication?::	No
Total Drawing Sheets::	1
Small Entity?::	Yes
Petition included?::	No
Secrecy Order in Parent Appl.?::	No

Applicant Information

Applicant Authority Type::	Inventor
Primary Citizenship Country::	US
Status::	Full Capacity
Given Name::	David
Middle Name::	C.
Family Name::	YEOMANS
City of Residence::	Sunnyvale
State or Province of Residence::	CA
Country of Residence::	US
Street of mailing address::	1293 Bedford Court
City of mailing address::	Sunnyvale
State or Province of mailing address::	CA

Postal or Zip Code of mailing address:: 94087

Applicant Authority Type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: William
Middle Name:: H.
Family Name:: FREY
Name Suffix:: II
City of Residence:: White Bear Lake
State or Province of Residence:: MN
Country of Residence:: US
Street of mailing address:: 4800 Centerville Road
Apt. 216
City of mailing address:: White Bear Lake
State or Province of mailing address:: MN
Postal or Zip Code of mailing address:: 55127

Applicant Authority Type:: Inventor
Primary Citizenship Country:: US
Status:: Full Capacity
Given Name:: Daniel
Middle Name:: I.
Family Name:: JACOBS
City of Residence:: Palo Alto
State or Province of Residence:: CA
Country of Residence:: US
Street of mailing address:: 2344 Old Page Mill Road
City of mailing address:: Palo Alto
State or Province of mailing address:: CA
Postal or Zip Code of mailing address:: 94304

Correspondence Information

Correspondence Customer Number:: 25226

Representative Information

Representative Customer Number:: 25226

Domestic Priority Information

Foreign Priority Information

Assignee Information