
Handbook of PHARMACEUTICAL EXCIPIENTS

Third Edition

Edited by

Arthur H. Kibbe, Ph.D.

Professor and Chair
Department of Pharmaceutical Sciences
Wilkes University School of Pharmacy
Wilkes-Barre, Pennsylvania



American Pharmaceutical Association
Washington, D.C.



London, United Kingdom

Published by the American Pharmaceutical Association
2215 Constitution Avenue NW, Washington, DC 20037-2985, USA
www.aphanet.org
and the Pharmaceutical Press
1 Lambeth High Street, London SE1 7JN, UK
www.pharmpress.com

© 1986, 1994, 2000 American Pharmaceutical Association and Pharmaceutical Press

First edition 1986
Second edition 1994
Third edition 2000

Printed in the United States of America

ISBN: 0-85369-381-1 (UK)
ISBN: 0-917330-96-X (USA)

Library of Congress Cataloging-in-Publication Data

Handbook of pharmaceutical excipients / edited by Arthur H. Kibbe.--3rd ed.
p. ; cm.

Includes bibliographical references and index.

ISBN 0-917330-96-X

1. Excipients--Handbooks, manuals, etc. I. Kibbe, Arthur H. II. American
Pharmaceutical Association.

[DNLM: 1. Excipients--Handbooks. QV 735 H236 2000]

RS201.E87 H36 2000

615'.19--dc21

99-044554

A catalogue record for this book is available from the British Library.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, without the prior written permission of the copyright holder. The publisher makes no representation, express or implied, with regard to the accuracy of the information contained in this book and cannot accept any legal responsibility or liability for any errors or omissions that may be made.

Managing Editor: Melanie Segala
Copyeditor: Paul Gottehrer
Indexer: Lillian Rodberg
Compositor: Roy Barnhill
Cover Designer: Tim Kaage

Alcohol

1. Nonproprietary Names

BP: Ethanol (96%)
 JP: Ethanol
 PhEur: Ethanol (96 per centum)
 USP: Alcohol

2. Synonyms

Ethyl alcohol; ethyl hydroxide; grain alcohol; methyl carbinol.


3. Chemical Name and CAS Registry Number

Ethanol [64-17-5]

4. Empirical Formula Molecular Weight

C₂H₆O 46.07

5. Structural Formula



6. Functional Category

Antimicrobial preservative; disinfectant; skin penetrant; solvent.

7. Applications in Pharmaceutical Formulation or Technology

Ethanol and aqueous ethanol solutions of various concentrations (*see* Sections 8 and 18) are widely used in pharmaceutical formulations and cosmetics. Although ethanol is primarily used as a solvent it is also employed in solutions as an antimicrobial preservative.^(1,2) Topical ethanol solutions are also used as penetration enhancers⁽³⁾ and as disinfectants.

Use	Concentration (% v/v)
Antimicrobial preservative	≥ 10
Disinfectant	60-90
Extracting solvent in galenical manufacture	Up to 85
Solvent in film coating	Variable
Solvent in injectable solutions	Variable
Solvent in oral liquids	Variable
Solvent in topical products	60-90

8. Description

In the BP, the term 'ethanol' used without other qualification refers to ethanol containing ≥ 99.5% v/v of C₂H₆O. The term 'alcohol', without other qualification, refers to ethanol 96.0-96.6% v/v. Where other strengths are intended, the term 'alcohol' or 'ethanol' is used, followed by the statement of the strength.

In the USP, the term 'dehydrated alcohol' refers to ethanol ≥ 99.5% v/v. The term 'alcohol', without other qualification refers to ethanol 94.9-96.0% v/v.

In the JP, ethanol (alcohol) contains 95.1-95.6% v/v (by specific gravity) of C₂H₆O at 15°C.

In the *Handbook of Pharmaceutical Excipients*, the term 'alcohol' is used for either ethanol 95% v/v or ethanol 96% v/v.

Alcohol is a clear, colorless, mobile, and volatile liquid with a slight, characteristic odor and burning taste.

See also Section 18.

9. Pharmacopeial Specifications

Test	JP	PhEur	USP
Identification	+	+	+
Specific gravity	0.814-0.816	0.8051-0.8124	0.812-0.816
Acidity	+	+	+
Clarity of solution	+	+	—
Nonvolatile residue	+	≤ 2.5 mg/100 mL	≤ 1 mg/40 mL
Water-insoluble substances	—	—	+
Aldehydes	+	—	+
Amyl alcohol, etc.	—	—	+
Absorbance	—	+	—
Fusel oil constituents	+	—	—
Acetone and propan-2-ol	—	—	+
Methanol	—	—	+
Reducing substances	+	—	—
Organic volatile impurities	+	+	—
Chloride	+	—	—
Heavy metals	≤ 1.2 ppm	—	—

10. Typical Properties

Antimicrobial activity: ethanol is bactericidal in aqueous mixtures at concentrations between 60-95% v/v; the optimum concentration is generally considered to be 70% v/v. Antimicrobial activity is enhanced in the presence of edetic acid or edetate salts.⁽¹⁾ Ethanol is inactivated in the presence of nonionic surfactants and is ineffective against bacterial spores.

Boiling point: 78.15°C

Flammability: readily flammable, burning with a blue, smokeless flame.

Flash point: 14°C (closed cup)

Solubility: miscible with chloroform, ether, glycerin, and water (with rise of temperature and contraction of volume).

Specific gravity: 0.8119-0.8139 at 20°C

Note: the above typical properties are for alcohol (ethanol 95% or 96% v/v). *See* Section 18 for typical properties of dehydrated alcohol.

11. Stability and Storage Conditions

Aqueous ethanol solutions may be sterilized by autoclaving or by filtration and should be stored in airtight containers, in a cool place.

12. Incompatibilities

In acidic conditions, ethanol solutions may react vigorously with oxidizing materials. Mixtures with alkali may darken in

color due to a reaction with residual amounts of aldehyde. Organic salts or acacia may be precipitated from aqueous solutions or dispersions. Ethanol solutions are also incompatible with aluminum containers and may interact with some drugs.

13. Method of Manufacture

Ethanol is manufactured by the controlled enzymatic fermentation of starch, sugar, or other carbohydrates. A fermented liquid is produced containing about 15% ethanol; ethanol 95% v/v is then obtained by fractional distillation. Ethanol may also be prepared by a number of synthetic methods.

14. Safety

Ethanol and aqueous ethanol solutions are widely used in a variety of pharmaceutical formulations and cosmetics. Ethanol is also consumed in alcoholic beverages.

Ethanol is rapidly absorbed from the gastrointestinal tract and vapor may be absorbed through the lungs. Ethanol is metabolized mainly in the liver to acetaldehyde, which is further oxidized to acetate.

Ethanol is a central nervous system depressant and ingestion of low to moderate quantities can lead to symptoms of intoxication including muscle incoordination, visual impairment, slurred speech, etc. Ingestion of higher concentrations may cause depression of medullary action, lethargy, amnesia, hypothermia, hypoglycemia, stupor, coma, respiratory depression, and cardiovascular collapse. The lethal human blood-alcohol concentration is generally estimated to be 400-500 mg/100 mL.

Although symptoms of ethanol intoxication are usually encountered following deliberate consumption of ethanol-containing beverages, many pharmaceutical products contain ethanol as a solvent which, if ingested in sufficiently large quantities, may cause adverse symptoms of intoxication.

Parenteral products containing up to 50% of alcohol (ethanol 95% or 96% v/v) have been formulated. However, such concentrations can produce pain on intramuscular injection and lower concentrations such as 5-10% v/v are preferred. Subcutaneous injection of alcohol (ethanol 95% v/v) similarly causes considerable pain followed by anesthesia. If injections are made close to nerves, neuritis and nerve degeneration may occur. This effect is used therapeutically to cause anesthesia in cases of severe pain although the practice of using alcohol in nerve blocks is controversial. Doses of 1 mL of absolute alcohol have been used for this purpose.⁽⁴⁾

Preparations containing greater than 50% v/v alcohol may cause skin irritation when applied topically.

LD₅₀ (guinea pig, IP): 3.41 g/kg⁽⁵⁾

LD₅₀ (guinea pig, IV): 2.3 g/kg

LD₅₀ (guinea pig, oral): 5.56 g/kg

LD₅₀ (hamster, IP): 5.07 g/kg

LD₅₀ (mouse, IP): 0.93 g/kg

LD₅₀ (mouse, IV): 1.97 g/kg

LD₅₀ (mouse, oral): 7.5 g/kg

LD₅₀ (mouse, SC): 8.29 g/kg

LD₅₀ (rabbit, IP): 0.96 g/kg

LD₅₀ (rabbit, IV): 2.37 g/kg

LD₅₀ (rabbit, oral): 6.3 g/kg

LD₅₀ (rat, IP): 3.75 g/kg

LD₅₀ (rat, IV): 1.44 g/kg

LD₅₀ (rat, oral): 7.06 g/kg

15. Handling Precautions

Observe normal precautions appropriate to the circumstances and quantity of material handled. Ethanol and aqueous ethanol solutions should be handled in a well-ventilated environment. In the UK, the long-term 8-hour TWA exposure limit for ethanol is 1920 mg/m³ (1000 ppm).⁽⁶⁾ Ethanol may be irritant to the eyes and mucous membranes and eye protection and gloves are therefore recommended. Ethanol is flammable and should be heated with care. Fixed storage tanks should be electrically grounded to avoid ignition from electrostatic discharges, when ethanol is transferred.

16. Regulatory Status

Included in the FDA Inactive Ingredients Guide (dental preparations, inhalations, IM and IV injections, nasal and ophthalmic preparations, oral capsules, solutions, suspensions, syrups and tablets, rectal, topical, and transdermal preparations). Included in nonparenteral and parenteral medicines licensed in the UK.

17. Pharmacopeias

China, Eur, Int, Jpn, Pol, and US.

18. Related Substances

Dehydrated alcohol; denatured alcohol; dilute alcohol; isopropyl alcohol.

Dehydrated alcohol

Synonyms: absolute alcohol; anhydrous ethanol; ethanol.

Autoignition temperature: 365°C

Boiling point: 78.5°C

Explosive limits: 3.5-19.0% v/v in air

Flash point: 12°C (closed cup)

Hygroscopicity: absorbs water rapidly from the air.

Melting point: -112°C

Refractive index: n_D²⁰ = 1.361

Specific gravity: 0.7904-0.7935 at 20°C

Surface tension: 22.75 mN/m at 20°C (ethanol/vapor)

Vapor density (relative): 1.59 (air = 1)

Vapor pressure: 5.8 Pa at 20°C

Viscosity (dynamic): 1.22 mPa s (1.22 cP) at 20°C

Comments: dehydrated alcohol is ethanol ≥ 99.5% v/v. See Section 8.

Denatured alcohol

Synonyms: industrial methylated spirit; surgical spirit,

Comments: denatured alcohol is alcohol, for external use only, that has been rendered unfit for human consumption by the addition of a denaturing agent such as methanol or methyl isobutyl ketone.

Dilute alcohol

Synonyms: dilute ethanol.

Specific gravity:

Strength of alcohol (% v/v)	Specific gravity at 20°C
90	0.8289-0.8319
80	0.8599-0.8621
70	0.8860-0.8883
60	0.9103-0.9114
50	0.9314-0.9326
45	0.9407-0.9417
25	0.9694-0.9703
20	0.9748-0.9759

Comments: the term 'dilute alcohol' refers to a mixture of ethanol and water of stated concentration. The BP lists eight strengths of dilute alcohol (dilute ethanol) containing 90, 80, 70, 60, 50, 45, 25, and 20% v/v respectively of ethanol.

19. Comments

Possession and use of nondenatured alcohols are usually subject to close control by excise authorities.

20. Specific References

1. Chiori CO, Ghobashy AA. A potentiating effect of EDTA on the bactericidal activity of lower concentrations of ethanol. *Int J Pharmaceutics* 1983; 17: 121-128.

2. Karabit MS, Juneskans OT, Lundgren P. The determination of antimicrobial characteristics of some pharmaceutical compounds in aqueous solutions. *Int J Pharmaceutics* 1989; 54: 51-56.
3. Liu P, Higuchi WI, Song W, Kurihara-Bergstrom T, Good WR. Quantitative evaluation of ethanol effects on diffusion and metabolism of β -estradiol in hairless mouse skin. *Pharm Res* 1991; 8: 865-872.
4. Lloyd JW. Use of anaesthesia: the anaesthetist and the pain clinic. *Br Med J* 1980; 281: 432-434.
5. Sweet DV, editor. Registry of Toxic Effects of Chemical Substances. Cincinnati, US Department of Health, 1987.
6. Health and Safety Executive. Occupational exposure limits 1998: EH40/98. Sudbury, Health and Safety Executive, 1998.

21. General References

- Lund W, editor. *The Pharmaceutical Codex: Principles and Practice of Pharmaceutics*, 12th edition. London, The Pharmaceutical Press, 1994; 694-695.
- Spiegel AJ, Noseworthy MN. Use of nonaqueous solvents in parenteral products. *J Pharm Sci* 1963; 52: 917-927.
- Wade A, editor. *Pharmaceutical Handbook*, 19th edition. London, The Pharmaceutical Press, 1980; 227-230.

22. Authors

PJ Weller.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.