Chemical Stability of Pharmaceuticals

A Handbook for Pharmacists

Second Edition

Kenneth A. Connors
School of Pharmacy, The University of Wisconsin


Gordon L. Amidon
College of Pharmacy, The University of Michigan

Valentino J. Stella School of Pharmacy, The University of Kansas

A Wiley-Interscience Publication
JOHN WILEY & SONS

New York • Chichester • Brisbane • Toronto • Singapore







Morphine

GENERAL

Names

Morphine; 7,8-didehydro-4,5 α -epoxy-17-methylmorphinan-3,6 α -diol.

Structure

Forms Available

Free base (morphine); morphine hydrobromide, $c_{17} \mu_{19} \mu_{03} \cdot \mu_{Br} \cdot 2\mu_{20}$; morphine hydrochloride, $c_{17} \mu_{19} \mu_{03} \cdot \mu_{C1} \cdot 3\mu_{20}$; morphine methylbromide, $c_{18} \mu_{22} \mu_{20} \mu_{22} \mu_{20} \mu_{20} \mu_{20}$; morphine sulfate, $(c_{17} \mu_{19} \mu_{03})_2 \cdot \mu_{25} \mu_{25} \cdot \mu_{25}$

Physical Properties

Melting point 254 to 256°C (decomposition); it has a metastable phase with melting point 197°C. Solubility: 1 g of morphine dissolves in about 5.0 L of water at 25°C; 1 g in 1075 mL of boiling water; 1 g in 210 mL of ethanol at 25°C; 1 g in 1220 mL of chloroform; 1 g in about 6.5 L of ether; freely soluble in alkaline solutions and acidic solutions. Most morphine salts are soluble to the extent of 1 g in 15 to 20 mL of water at 25°C and are freely soluble in boiling water. A saturated aqueous solution of morphine

604



is basic to litmus (1). $\rm pK_a=9.51$ at 25°C for the phenolic group (2); $\rm pK_a=8.31$ at 25°C for the amino group (2).

Stability Summary

Although morphine contains many functional groups that undergo reaction under certain conditions, the functional group primarily responsible for its instability in aqueous solutions is the phenolic moiety. This group readily undergoes oxidation, the major products being pseudomorphine (I) and morphine N-oxide (II). Aqueous morphine solutions exhibit a strong dependence of stability on pH and also on the presence of oxygen. The pH-rate profile shows that solutions of morphine are relatively stable at low pH values, but degradation proceeds at a much higher rate in neutral or basic media. Degradation is often accompanied by the appearance of a yellow or brown color in the solution.

DRUG KINETICS

Reactions

Oxidation is the most important reaction contributing to the degradation of aqueous morphine solutions. The oxidation of morphine is promoted by oxygen (3-6), sunlight (3), UV radiation (7), organic impurities (3,8), certain fungi (9), and iron (10). The major products of the oxidation are pseudomorphine (I) (oxydimorphine) and morphine N-oxide (II), which are formed in a ratio of 9:1, respectively. A trace of a base, believed to be methylamine (3), is also formed.

The kinetics of the degradation are dependent on both the amount of oxygen and the particular morphine species present (3). The overall reactions can be written

$$M + O_2 \xrightarrow{k_M} products$$
 (1)

$$^{+}$$
HM + $O_2 \xrightarrow{k_{HM}^{+}}$ products (2

where M and $^{+}\mathrm{HM}$ represent the neutral and protonated forms of morphine, respectively. The parameters k_{M}

cy-17-methylmorphinan-

C17H19NO3

CH₃ mol. wt. 285.33

phine hydrobromide, rochloride, $C_{17}H_{19}NO_3$ ·ide, $C_{18}H_{22}NO_3Br$; more $_2SO_4\cdot 5H_2O$; morphine Notinate, $C_{29}H_{25}N_3O_5$.

composition); it has a point 197°C. Solubilves in about 5.0 L of boiling water; 1 g in in 1220 mL of chloroher; freely soluble in solutions. Most morextent of 1 g in 15 to reely soluble in boils solution of morphine



and $k_{\pm HM}$ are second-order rate constants for the oxygen-catalyzed degradation of M and $^{\pm}HM$, respectively. These rate constants become pseudo first order when a large excess of oxygen is maintained.

III

N-CH₃

II

pH-Rate Profile

Figure 1 shows a plot of log k vs. pH for the oxidation of aqueous morphine solutions at $95\,^{\circ}\text{C}$ in the presence of excess oxygen (3). The curve is satisfactorily described by the equation

rate =
$$\left[k_{1}^{'}\left(\frac{K_{a}}{K_{a}+[H^{+}]}\right) + k_{2}^{'}\left(\frac{H^{+}}{K_{a}+[H^{+}]}\right)\right][M]_{t}$$
 (3)

where $k_1^{'}$ and $k_2^{'}$ are pseudo-first-order rate constants for the initial oxidation reactions of M and ^HM to

DOCKET

Explore Litigation Insights



Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

