



US007320999B2

(12) **United States Patent**
Joshi et al.

(10) **Patent No.:** **US 7,320,999 B2**

(45) **Date of Patent:** **Jan. 22, 2008**

- (54) **DIMETHYL FUMARATE FOR THE TREATMENT OF MULTIPLE SCLEROSIS**
- (75) Inventors: **Rajendra Kumar Joshi, Zürich (CH); Hans-Peter Strebhel, Muri (CH)**
- (73) Assignee: **Fumapharm AG, Luzern (CH)**
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 202 days.
- (21) Appl. No.: **10/197,077**
- (22) Filed: **Jul. 17, 2002**
- (65) **Prior Publication Data**
US 2003/0018072 A1 Jan. 23, 2003
- Related U.S. Application Data**
- (62) Division of application No. 09/831,620, filed as application No. PCT/EP99/08215 on Oct. 29, 1999, now Pat. No. 6,509,376.
- (30) **Foreign Application Priority Data**
Nov. 19, 1998 (DE) 198 53 487
- (51) **Int. Cl.**
A61K 31/22 (2006.01)
- (52) **U.S. Cl.** **514/549**
- (58) **Field of Classification Search** None
See application file for complete search history.
- (56) **References Cited**

U.S. PATENT DOCUMENTS

3,832,287	A	8/1974	Gale et al.
4,515,974	A	5/1985	Zecher et al.
4,746,668	A	5/1988	Sato et al.
4,851,439	A	7/1989	Speiser et al.
4,959,389	A	9/1990	Speiser et al.
5,149,695	A	9/1992	Speiser et al.
5,214,196	A	5/1993	Blank
5,242,905	A	9/1993	Blank
5,359,128	A	10/1994	Blank
5,424,332	A	6/1995	Speiser et al.
5,451,667	A	9/1995	Speiser et al.
5,538,968	A	7/1996	Chiesi et al.
5,548,059	A	8/1996	Bayley et al.
5,589,504	A	12/1996	Dannenberg et al.
5,972,363	A	10/1999	Clikeman et al.
6,277,882	B1	8/2001	Joshi et al.
6,355,676	B1	3/2002	Joshi et al.
6,359,003	B1	3/2002	Joshi et al.
6,436,992	B1	8/2002	Joshi et al.
6,509,376	B1	1/2003	Joshi et al.
2003/0013761	A1	1/2003	Joshi et al.
2006/0069161	A1	3/2006	Lee et al.

FOREIGN PATENT DOCUMENTS

CA	2248955	8/2001
CN	1125141	6/1996
DE	25 30 372	A1 1/1977
DE	26 21 214	A1 11/1977
DE	35 31 597	3/1987

DE	38 34794	A1 4/1990
DE	3834794	A1 4/1990
EP	0 188 749	A2 7/1986
EP	0 312 697	4/1989
EP	0 312 697	A2 4/1989
EP	0 518 388	12/1992
EP	0 793 966	9/1997
GB	1 216 699	12/1970
GB	1 422 726	1/1976
GB	2 291 422	1/1996
WO	WO 89/01930	A1 3/1989
WO	WO 94/28883	12/1994
WO	WO 95/25102	9/1995
WO	WO 96/01122	1/1996
WO	WO 96/02244	2/1996
WO	WO 96/27369	9/1996
WO	WO 97/48400	12/1997
WO	WO 98/04290	2/1998
WO	WO 98/27970	7/1998
WO	WO 98/52549	11/1998
WO	WO 99/21565	5/1999
WO	WO 01/59072	8/2001
WO	WO 02/02190	1/2002
WO	WO 03/032969	4/2003
WO	WO 2004/096216	11/2004
WO	WO 2005/027899	3/2005
WO	WO 2005/044272	5/2005
WO	WO 2006/037342	4/2006
WO	WO 2006/050730	5/2006

(Continued)

OTHER PUBLICATIONS

- Anderson, A. et al., Contact Dermatitis, 16:55-78, 1987.
- deJong et al., European Journal of Immunology, 26:2067-2074, 1996.
- Dücker, P. et al., H+Z Zeitschrift für Haut., 65:734-736, 1989.
- Fliegner, L. et al., Hautarzt, 43:554-560, 1992.
- Ghoreschi et al., Arch. Dermatol. Res., 296:P110, 2005.
- Ghoreschi et al., Arch. Dermatol. Res., 294:28, 2002.
- Ghoreschi et al., Current Drug Targets: Inflammation and Allergy 3:193-198, 2004.

(Continued)

Primary Examiner—Raymond J. Henley, III
(74) Attorney, Agent, or Firm—Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P.

(57) **ABSTRACT**

The present invention relates to the use of certain dialkyl fumarates for the preparation of pharmaceutical preparations for use in transplantation medicine or for the therapy of autoimmune diseases and said compositions in the form of micro-tablets or pellets. For this purpose, the dialkyl fumarates may also be used in combination with conventional preparations used in transplantation medicine and immunosuppressive agents, especially cyclosporines.

18 Claims, No Drawings



FOREIGN PATENT DOCUMENTS

WO WO 2006/055871 5/2006

OTHER PUBLICATIONS

- Hunziker et al., "Is Psoriasis an Autoimmune Disease?", excerpt from "Therapeutische Umschau", Derm. Clinic of Univ. Berne, 50:110-113, 1993, translated version 5 pages.
- International Search Report for PCT/EP99/08215 dated Jun. 26, 2000.
- Kiehl, R et al., Acta Derm. Venerol., 72:253-255, 1992.
- Kolbach, D. et al., J. Am. Acad. Derm., 27:769-771, 1992.
- Lahti, A. et al., Contact Dermatitis, 12:139-140, 1985.
- Merck Index, 10th Ed., 1983 Abs. 2748.
- Nieboer, C. et al., J. Am. Acad. Derm., 20:601-608, 1989.
- Peeters et al., Br. J. Rheumatol., 317:502-504, 1992.
- Peeters et al., Ned. Tijdschr. Geneesk., 136:2428-2431, 1992.
- Sadjak, A. et al., Deutsch Med. Wochenschr., 116:478, 1991.
- Schilling et al., Clinical and Experimental Immunology, 145:101-107, 2006.
- Schilling et al., Aktuelle Rheumatologie, 24(6):174-179, 1999.
- Schmirgk et al., European J. of Neurology: The Official Journal of the European Federation of Neurological Societies, 13:604-610, 2006.
- Sebok, B. et al., Skin Pharm., 9:99-103, 1996.
- Thio, H. et al., Br. J. Dermatol., 131:865-861, 1994.
- English Language Derwent Abstract for CN 1125141, date unavailable.
- English Language Derwent Abstract for DE 35 31 597, date unavailable.
- Altmeier, P. et al., "Systemische Therapie der Psoriasis", T & E Dermatologie Jg., 1997, vol. 27, pp. 380-382, 384—not translated.
- M. Bacharach-Buhles et al., "Fumaric Acid Esters (FAEs) Suppress CD 15- and ODP 4-positive Cells in Psoriasis", Acta Derm Venerol (Stockh); 1994; Suppl. 186: 79-82.
- H. M. Ockenfels, et al., "The antipsoriatic agent dimethylfumarate immunomodulates T-cell cytokine secretion and inhibits cytokines of the psoriatic cytokine network", British Journal of Dermatology, 1998, vol. 139, 390-395.
- Gasser, et al., "Host Vs Graft and Graft Vs Host Reactions After Allogeneic Heterotopic Small Bowel Transplantation in the Rat", Transplantation Proceedings, vol. 24, No. 3, Jun. 1992, pp. 1128-1129.
- Hunziker T. et al.; "Is Psoriasis an Autoimmune Disease", Excerpt from "Therapeutische Umschau", Dermatological Clinic of the University of Berne; 1993, vol. 50; 2nd edition; pp. 110-113. (Translated version 5 pages).
- Nathens, et al., "The Glutathione Depleting Agent Diethylmaleate Prolongs Renal Allograft Survival", Journal of Surgical Research, vol. 77, 1998, pp. 75-79.
- Nibbering, P.H. et al., "Intracellular Signalling by Binding Sites for the Antipsoriatic Agent Monomethylfumarate on Human Granulocytes", British J. Dermatol., 1997, vol. 137, pp. 65-75.
- Nibbering, Peter H., "Effects of Monomethylfumarate on Human Granulocytes", Journal of Investigative Dermatology, 1993, vol. 101, pp. 37-42.
- Seböck, Bela et al., "Antiproliferative and Cytotoxic profiles of Antipsoriatic Fumaric Acid Derivatives in Keratinocyte Cultures", European Journal of Pharm., Environ. Toxicol. Pharmacol. Sect., 1994, vol. 270, pp. 79-87.
- Schwinghammer et al., "Pharmacologic prophylaxis of acute graft-versus-host disease after allogeneic marrow transplantation", Therapy Reviews, Clinical Pharmacy, vol. 12, Oct. 1993, pp. 736-761.
- Medline Abstract of Bayard et al., "Peroral long-term treatment of psoriasis using fumaric acid derivatives", Hautarzt, May 1987, 38(5), pp. 279-85.
- "Merck Manual", 1987, Merck XP-002141006, p. 327, paragraph 2-paragraph 6.
- Immunmodulation durch Fumaderm, Das richtungweisende Konzept, Charité-Berlin, Hautklinik, Symposium, 1.-3. Nov. 1996, 28 pages, 4 page english translation of pp. 23-24.
- Amamoto, Toshiro, et al., "Effect of E-64, Thiol Protease Inhibitor on the Secondary Anti-SRBC Response In Vitro", Microbiol. Immunol., vol. 28(1), 1984, pp. 85-97.
- Barrett, Alan J., et al., "L-trans-Epoxy succinyl-leucylamido(4-guanidino)butane (E-64) and its analogues as inhibitors of cysteine proteinases including cathepsins B, H and L", Biochem. J., 1982, vol. 201, pp. 189-198.
- Bellier, Bruno, et al., "Replacement of Glycine with Dicarbonyl and Related Moieties in Analogues of the C-Terminal Pentapeptide of Cholecystokinin: CCK₂ Agonists Displaying a Novel Binding Mode", J. Med. Chem., vol. 43, 2000, pp. 3614-3623.
- Birch, A.J., et al., "Metabolites of *Aspergillus indicus*: The Structure and Some Aspects of the Biosynthesis of Dihydrocanadensolide", Aust. J. Chem., 1968, vol. 21, pp. 2775-2784.
- Choo, Hea-Young, et al., "Design and Synthesis of α,β -unsaturated Carbonyl Compounds as Potential ACE Inhibitors", Short Communication, Eur. J. Med. Chem., vol. 35, 2000, pp. 643-648.
- Dethlefsen, L.A., "Toxic Effects of Acute Glutathione Depletion by Buthionine Sulfoximine and Dimethylfumarate on Murine Mammary Carcinoma Cells", Radiation Research, vol. 114, 1988, pp. 215-224.
- Galpin, I.J., et al., "The Synthesis of an Insulin Active Site Analogue", Tetrahedron, vol. 39, No. 1, 1983, pp. 149-158.
- Gerhard, Ute, et al., "The Free Energy Change of restricting A Bond Rotation in the Binding of Peptide Analogues to Vancomycin Group Antibiotics", Bioorganic & Medicinal Chemistry Letters, vol. 3, No. 5, 1993, pp. 803-808.
- Gordon, G.B., et al., "Induction of NAD(P)H:quinone reductase in human peripheral blood lymphocytes", Carcinogenesis, vol. 12 (12), 1991, pp. 2393-2396.
- Griehl, C., et al., " α -Aspartyl Peptides by Addition of Amines to N-Maleylamino Acid Derivatives", Chemistry of Peptides and Proteins, 1993, 5/6(pt. A), pp. 99-103.
- Hildebrandt, H., "Psyhyrembel Klinisches Woerterbuch Ed. 258", 1998, Walter de Gruyter, New York, XP 002234302, p. 182, col. 1, paragraph 2 and p. 1469, col. 1, paragraph 16-col. 2, paragraph 1. Not translated.
- Hohenegger, M., et al., "Nephrotoxicity of Fumaric Acid Monoethyl ester (FA ME)", Advances in Experimental Medicine and Biology, US 1989, vol. 252, pp. 265-272.
- Holroyd, S.E., et al., "Rational Design and Binding of Modified Cell-Wall Peptides to Vancomycin-Group Antibiotics: Factorising Free Energy Contributions to Binding", Tetrahedron, vol. 49, No. 41, 1993, pp. 9171-9182.
- Kamiyama, T., et al., "Ro 09-1679, A Novel Thrombin Inhibitor", The Journal of Antibiotics, vol. 45, No. 3, Mar. 1992, pp. 424-427.
- Krstenansky, J.L., et al., "Development of MDL 28,050, a Small Stable Antithrombin Agent Based on a Functional Domain of the Leech Protein, Hirudin", Thrombosis and Haemostasis, vol. 63, No. 2, 1990, Stuttgart, De.
- Kuroda, K., et al., "Fumaric Acid Enhances DNA Synthesis of Rat Hepatocytes by Counter Acting the Toxicities of Mitomycin C and Aflatoxin B₁", Jpn. J. Cancer Res. (Gann), Aug. 1986, vol. 77, pp. 750-758.
- Kuroda, K., et al., "Inhibitory Effect of Capsella-bursa-pastoris extract on Growth of Ehrlich Solid Tumor in Mice", Cancer Research, vol. 36, 1976, Abstract only.
- Langlois, M., et al., "Synthesis of symmetrical pseudopeptides as potential inhibitors of the human immunodeficiency virus-1 protease", Eur. J. Med. Chem., vol. 29, 1994, pp. 639-647.
- Lehnert, S., et al., "Radiation Response of Drug-Resistant Variants of a Human Breast Cancer Cell Line: The Effect of Glutathione Depletion", Radiation Research, vol. 124, 1990, pp. 208-215.
- Miller, A.C., et al., "Posttranscriptional Down-Regulation of ras Oncogene Expression by Inhibitors of Cellular Glutathione", Molecular and Cellular Biology, Jul. 1993, vol. 13, No. 7, pp. 4416-4422.
- Mrowietz, U., "Nephrotoxisch Wirkung durch Fumarsaure", Hautarzt, 2000-51:615, Springer-Verlag 2000, pp. 615. Not translated.
- Odom, R.Y., et al., "Cancer Chemoprotective Agents Inhibition of Human HT29 Colon Carcinoma Cell Proliferation is Reversed by

- N-Acetyl Cysteine", Proceedings of the American Assoc. for Cancer Research Annual, No. 41, Mar. 2000, pp. 342, XP008017517.
- Ondrus, V., et al., "A Simple Synthesis of Some analogues of Natural Antibiotics", Preliminary Communication, Chem. Papers, 51(3), 1997, pp. 164-166.
- Orta, T., et al., "Glutathione manipulation and the radiosensitivity of human tumour and fibroblast cell lines", Int. J. Radiat. Biol., 1995, vol. 68, No. 4, pp. 413-419.
- Pearl, J.M., et al., "Fumarate-enriched blood cardioplegia results in complete functional recovery of immature myocardium", Annals of Thoracic Surgery, vol. 57, No. 6, 1994, Abstract only, 1 page.
- Peeters, A.J., et al., "Fumaric Acid Therapy for Psoriatic Arthritis. A Randomized, Double-blind, Placebo-controlled Study", British Journal of Rheumatology, vol XXXI, No. 7, Jul. 1992, pp. 502-504.
- Pereira, M.A., et al., "Use of azoxymethane-induced foci of aberrant crypts in rat colon to identify potential cancer chemopreventive agents", Carcinogenesis, vol. 15, No. 5, 1994, pp. 1049-1054.
- Portoghesi, P.S., et al., "Synthesis and Biological Activity of Analogues of β -Chlornaltrexamine and β -Funnaltrexamine at Opioid Receptors", J. of Medicinal Chem., vol. 29, No. 10, 1986, pp. 1861-1864.
- Prochaska, H.J., et al., "Elevation of Glutathione Levels by Phase II Enzyme Inducers: Lack of Inhibition of Human Immunodeficiency Virus Type 1 Replication in Chronically Infected Monocytoid Cells", Molecular Pharmacology, vol. 45, No. 5, 1994, pp. 916-921.
- Prochaska, H.J., et al., "Oltipraz, an inhibitor of human immunodeficiency virus type 1 replication", Proc. Natl. Acad. Sci., USA, vol. 90, May 1993, pp. 3953-3957.
- Rao, C.V., et al., "Chemoprevention of Azoxymethane-Induced Colon Cancer by Ascorbylpalmitate, Carbenoxolone, Dimethylfumarate and p-Methoxyphenol in Male F344 Rats", Anticancer Research, vol. 15, 1995, pp. 1199-1204.
- Rao, K.S., et al., "Antihepatotoxic activity of monomethyl fumarate isolated from *Fumaria indica*", Journal of Ethnopharmacology, vol. 60, 1998, pp. 207-213.
- Roodnat, J.I., et al., "Akute Niereninsuffizienz bei der Behandlung der Psoriasis mit Fumarsaure-Estern", Schweiz. Med., Wschr., vol. 119, nr 2, 1989, pp. 826-830. Not translated.
- Rossi, Domenico, et al., "Approach to the Use of Benzylpenicillinacylase for Configurational Correlations of Amino Compounds. 2. Hydrolysis of N-(p-Aminophenylacetyl) Derivatives of Some Chiral Primary Amines", J. Org. Chem., vol. 44, No. 13, 1979, pp. 2222-2225.
- Schirmeister, Tanja, "Aziridine-2,3-dicarboxylic Acid Derivatives as Inhibitors of Papain", Arch. Pharm. Pharm. Med. Chem., 329, 1996, pp. 239-244.
- Schmidt, K.N., et al., "Anti-psoriatic drug anthralin activates transcription factor NF-kappa-B in murine keratinocytes", Journal of Immunology, vol. 156, 1996, Abstract Only.
- Spencer, S.R., et al., "Induction of Glutathione Transferases and NAD(P)H:Quinone Reductase by Fumaric Acid Derivatives in Rodent Cells and Tissues", Cancer Research, vol. 50, 1990, pp. 7871-7875.
- Steele, V.E., et al., "Preclinical Efficacy Evaluation of Potential Chemopreventive Agents in Animal Carcinogenesis Models: Methods and Results From the NCI Chemoprevention Drug Development Program", J. of Cellular Biochemistry, Supplement 20, 1994, pp. 32-54.
- Su, Jean Y.C., et al., "Reduction of H₂O₂-evoked, intracellular calcium increases in the rat N18-RE-105 neuronal cell line by pretreatment with an electrophilic antioxidant inducer", Neuroscience Letters, 273, 1999, pp. 109-112.
- Subasinghe, Nalin et al., "Synthesis of Acyclic and Dehydroaspartic Acid Analogues of Ac-Asp-Glu-OH and Their Inhibition of Rat Brain N-Acetylated α -linked Acidic Dipeptidase (NAALA Dipeptidase)", Journal of Medicinal Chemistry, vol. 33, No. 10, 1990, pp. 2734-2744.
- Vandermeeren, M., et al., "Dimethylfumarate is an Inhibitor of Cytokine-Induced Nuclear Translocation of NF-kB1, But not RelA in Normal Human Dermal Fibroblast Cells", The Journal of Investigative Dermatology, vol. 116, No. 1, Jan. 2001, pp. 124-130.
- Vandermeeren, M., et al., "Dimethylfumarate is an Inhibitor of Cytokine-Induced E-Selection, VCAM-1, and ICAM-1 Expression in Human Endothelial Cells", Biochemical and Biophysical Research Communications, vol. 234, 1997, pp. 19-23.
- Wang, X., et al., "Enhanced cytotoxicity of mitomycin C in human tumour cells with inducers of DT-diaphorase", British Journal of Cancer, vol. 80(8), 1999, pp. 1223-1230.
- Weinmann, I., et al., "Influence of Fumaric Acid Derivatives on T Lymphocytes in the Murine Model of HSV-I Keratitis", IOVS, vol. 41, No. 4, Mar. 15, 2000, XP008017516, pp. S146.
- English Language Translation of PCT International Preliminary Examination Report for PCT/EP99/08215, completed Feb. 27, 2001.

DIMETHYL FUMARATE FOR THE TREATMENT OF MULTIPLE SCLEROSIS

REFERENCE TO RELATED APPLICATIONS

This is a Division of commonly-owned application Ser. No. 09/831,620, filed May 10, 2001, now U.S. Pat. No. 6,509,376, which is a 371 continuation of PCT Application PCT/EP99/08215, filed Oct. 29, 1999, the text of which is not in English, which PCT Application claims priority on German Application No. 198 53 487.6, filed Nov. 19, 1998, the text of which is not in English.

DESCRIPTION

The present invention relates to the use of dialkyl fumarates for preparing pharmaceutical preparations for use in transplantation medicine or the therapy of autoimmune diseases and pharmaceutical preparations in the form of micro-tablets or micro-pellets containing dialkyl fumarates.

On the one hand, therefore, it relates especially to the use of dialkyl fumarates for preparing pharmaceutical preparations for the treatment, reduction or suppression of rejection reactions of the transplant by the recipient, i.e. host-versus graft reactions, or rejection of the recipient by the transplant, i.e. graft-versus-host reactions. On the other hand, it relates to the use of dialkyl fumarates for preparing pharmaceutical preparations for treating autoimmune diseases such as polyarthritis, multiple sclerosis, juvenile-onset diabetes, Hashimoto's thyroiditis, Grave's disease, systemic Lupus erythematoses (SLE), Sjogren's syndrome, pernicious anaemia and chronic active (=lupoid) hepatitis.

Both graft rejection and autoimmune diseases are based on medically undesirable reactions or dysregulation of the immune system. Cytokins such as interleukins or tumour necrose factor α (TNF- α) are substantial mediators influencing the immune system. In general, both are treated by the administration of immunosuppressive agents such as cyclosporine.

In the overall result, autoimmune diseases may be defined as the failure of the tolerance of endogenic substances or antigens. As a rule, this tolerance can be maintained only if the antigens keep coming into contact with immunological cells. When this tolerance is lost, autoantibodies are formed, i.e. a humoral immunoreponse against endogenic tissue. The exact nature of the involvement of TNF- α is not known.

Transplantations are tissue or organ transplantations, i.e. the transfer of tissues such as cornea, skin, bones (bone chips), vessels or fasciae, of organs such as kidney, heart, liver, lung, pancreas or intestines, or of individual cells such as islet cells, α -cells and liver cells, the kidney having the greatest significance as a transplanted organ.

According to the degree of relationship between the donor and the recipient we differentiate between autotransplantation (transfer to another part of the body of the same individual), iso-transplantation (transfer to another, genetically identical individual) and allogenic transplantation (transfer to another individual of the same species). Depending on the site of origin and transplantation, we further differentiate between homotopic transplantation (transfer to the same site) and heterotopic transplantation (transfer to a different site). The above-mentioned transplantations play an important role in modern medicine.

A major problem in transplantation medicine is graft rejection after transplantation of the tissue, organ or cell by immunological defense reactions of the recipient. Such a graft rejection is also called host-versus-graft reaction. The

immunological defense reaction of the organism against the heteroprotein often results in rejection or dissolution of the grafts. In host-versus-graft reactions, different stages may be distinguished. Depending on the degree of difference between the recipient and the donor, this reaction takes place at different speeds so that we speak of an acute, sub-acute or chronic reaction. The acute rejection process is accompanied by the irreversible loss of the transplant (necrotisation) as a result of arteriitis or arteriolitis within 48 hours and cannot be influenced by the administration of drugs. The sub-acute rejection reaction becomes manifest as a rejection crisis from day 12 to month 4 with reversible functional disorders as a result of a transplant vasculopathy. Finally, the loss of function of the transplant as a result of vascular changes such as obliterating arteriopathy, which proceeds over weeks or years and can practically not be influenced by drugs, is termed a chronic rejection reaction.

Vice-versa, rejection reactions of the transplant against the recipient, the so-called graft-versus-host reactions, may occur when immunocompetent tissues are transplanted, i.e. primarily in bone marrow transplantation. Again, the severity of the reaction is graded, and substantially similar complications result as in host-versus-graft-reactions, namely arteriopathies and necroses.

To avoid such rejection reactions, i.e. the host-versus-graft reaction and the graft-versus-host reaction, transplantation medicine essentially makes use of immunosuppression, i.e. a weakening of the normal immunoreponse. For this purpose, anti-lymphocyte sera are often used in combination with corticosteroids and so-called anti-metabolites, e.g. purine analogues such as 6-mercaptopurine and thioguanine which affect the nucleic acid and protein synthesis and thus prevent cell division and proliferation. This leads to suppression of the production of antibodies and the cellular immune response. The immunosuppressive agents used for therapy are substances which suppress or weaken the immunoreaction in the body either specifically or non-specifically. Non-specific immunosuppressive agents are cytostatic agents such as, for example, alkylating agents or antimetabolites.

In addition, active ingredients are known which cause at least partial specific immunosuppression, such as corticosteroids, antisera, antibodies FK-506, tacrolimus, mycophenolatemofetil and primarily cyclosporines such as cyclosporine A. As a result of using modern immunosuppressive agents, the most important representatives of which are the cyclosporines, especially cyclosporine A, it was possible to improve the results of transplantation considerably over the last few years. At present, the survival rate after one year is about 60% for liver transplantations, about 80% for heart transplantations and over 90% for kidney transplantations.

Autoimmune diseases where the endogenic immune system attacks endogenic organs, tissues and cells are comparable to graft-versus-host reactions. These are also medically undesirable reactions of the immune system which may be treated with immunosuppressive agents, too.

The danger in using immunosuppressive agents lies in weakening the body's defense against infectious diseases and the increased risk of malignant diseases. Therefore, it is the object of the invention to provide a pharmaceutical preparation to be employed in transplantation medicine which may be used to treat, especially to suppress weaken and/or alleviate host-versus-graft reactions and graft-versus-host reactions, but does not have the above disadvantage.

It is another object of the invention to provide a pharmaceutical preparation which may be employed for treating autoimmune diseases, particularly polyarthritis, multiple

3

sclerosis, juvenile-onset diabetes, Hashimoto's thyroiditis, Grave's disease, systemic Lupus erythematoses (SLE), Sjogren's syndrome, pernicious anaemia and chronic active (=lupoid) hepatitis, without the disadvantages of immunosuppression.

The object of the invention is achieved by using certain dialkyl fumarates for preparing pharmaceutical preparations for use in transplantation medicine and for the therapy of autoimmune diseases and pharmaceutical preparations in the form of micro-tablets and micro-pellets containing these dialkyl fumarates. The individual subject matters of the invention are characterized in detail in the claims. The preparations according to the invention do not contain any free fumaric acids per se.

It is known that pharmaceutical preparations which, upon biological degradation after administration, enter into the citric acid cycle or are part thereof gain increasing therapeutic significance—especially when given in high dosages—since they can alleviate or heal diseases caused cryptogenetically.

Fumaric acid, for example, inhibits the growth of the Ehrlich ascites tumour in mice, reduces the toxic effects of mitomycin C and aflatoxin and displays antipsoriatic and anti-microbial activity. When administered parenterally, transdermally and especially perorally, high dosages of fumaric acid or its derivatives known so far such as dihydroxyl fumaric acid, fumaramide and fumaronitrile have such unacceptably severe side effects and high toxicity that, in most cases, such a therapy had to be abandoned in the past.

Surprisingly, investigations carried out by the applicant have shown that methyl hydrogen fumarate, a metabolite of the dimethyl fumarate, initially increases the endotoxin-stimulated TNF- α secretion in human mononuclear cells of periphery blood (periphery blood mononuclear cells=PBMC cells) and in isolated monocytes. In addition, the applicant was able to show that fumaric acid has an effect on in vitro and in vivo haemagglutination which is comparable to that of cyclosporine.


Surprisingly, it has now been found that dialkyl fumarates are advantageous for preparing pharmaceutical compositions for use in transplantation medicine and for the therapy of autoimmune diseases. This is because compositions containing such dialkyl fumarates surprisingly permit a positive modulation of the immune system in host-versus-graft reactions, graft-versus-host reactions and other autoimmune diseases.

European Patent Application 0188 749 already describes fumaric acid derivatives and pharmaceutical compositions containing the same for the treatment of psoriasis. Pharmaceutical compositions for the treatment of psoriasis containing a mixture of fumaric acid and other fumaric acid derivatives are known from DE-A-25 30 372. The content of free fumaric acid is obligatory for these medicaments.

DE-A-26 21 214 describes medicaments containing the fumaric acid monoethyl ester and its mineral salts as active ingredient for the treatment of psoriasis. The publication "Hautarzt (*Dermatologist*) (1987) 279-285" discusses the use of fumaric acid monoethyl ester salts. Pharmaceutical preparations containing a mixture of fumaric acid monoalkyl ester salts and a fumaric acid diester for the treatment of psoriasis, psoriatic arthritis, neurodermatitis and enteritis regionalis Crohn are known from EP 0 312 697 B1.

Specifically, the object of the invention is achieved by the use of one or more dialkyl fumarates of the formula

4



wherein R_1 and R_2 , which may be the same or different, independently represent a linear, branched or cyclic, saturated or unsaturated C_{1-20} alkyl radical which may be optionally substituted with halogen (Cl, F, I, Br), hydroxy, C_{1-4} alkoxy, nitro or cyano for preparing a pharmaceutical preparation for use in transplantation medicine or for the therapy of autoimmune diseases.

The C_{1-20} alkyl radicals, preferably C_{1-8} alkyl radicals, most preferably C_{1-5} alkyl radicals are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, t-butyl, pentyl, cyclopentyl, 2-ethyl hexyl, hexyl, cyclohexyl, heptyl, cycloheptyl, octyl, vinyl, allyl, 2-hydroxyethyl, 2 or 3-hydroxy propyl, 2-methoxy ethyl, methoxy methyl or 2- or 3-methoxy propyl. Preferably at least one of the radicals R_1 or R_2 is C_{1-5} alkyl, especially methyl or ethyl. More preferably, R_1 and R_2 are the same or different C_{1-5} alkyl radicals such as methyl, ethyl, n-propyl or t-butyl, methyl and ethyl being especially preferred. Most preferably, R_1 and R_2 are identical and are methyl or ethyl. Especially preferred are the dimethyl fumarate, methyl ethyl fumarate and diethyl fumarate.

The dialkyl fumarates to be used according to the invention are prepared by processes known in the art (see, for example, EP 0 312 697).

Preferably, the active ingredients are used for preparing oral preparations in the form of tablets, micro-tablets, pellets or granulates, optionally in capsules or sachets. Preparations in the form of micro-tablets or pellets, optionally filled in capsules or sachets are preferred and are also a subject matter of the invention. The oral preparations may be provided with an enteric coating. Capsules may be soft or hard gelatine capsules.

The dialkyl fumarates used according to the invention may be used alone or as a mixture of several compounds, optionally in combination with the customary carriers and excipients. The amounts to be used are selected in such a manner that the preparations obtained contain the active ingredient in an amount corresponding to 10 to 300 mg of fumaric acid.

Preferred preparations according to the invention contain a total amount of 10 to 300 mg of dimethyl fumarate and/or diethyl fumarate.

According to a preferred embodiment, the size or the mean diameter, respectively, of the pellets or micro-tablets is in the range from 300 to 2,000 μm , especially in the range of 500 or 1,000 μm .

In addition to graft-versus-host reactions (see above), the following autoimmune diseases to be treated may be named: polyarthritis, multiple sclerosis, graft-versus-host reactions, juvenile-onset diabetes, Hashimoto's thyroiditis, Grave's disease, systemic Lupus erythematoses (SLE), Sjogren's syndrome, pernicious anaemia and chronic active (lupoid) hepatitis. Autoimmune diseases in a wider meaning also comprise psoriasis, psoriatic arthritis, neurodermatitis and enteritis regionalis Crohn.

In addition to the preparations for peroral administration in the form of micro-pellets, micro-tablets, capsules (such as soft and hard gelatine capsules), granulates and tablets cited above, suitable pharmaceutical preparations are preparations for cutaneous and transdermal administration in the form of ointments, plasters, lotions or shower preparations and for parenteral administration in the form of aqueous micro-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.