

Structure of Bis(methylguanidinium) Monohydrogen Orthophosphate. A Model for the Arginine-Phosphate Interactions at the Active Site of Staphylococcal Nuclease and Other Phosphohydrolytic Enzymes¹

F. A. Cotton,*^{2a} V. W. Day, E. E. Hazen, Jr.,*^{2a} S. Larsen,^{2b} and S. T. K. Wong

Contribution from the Departments of Chemistry, Texas A & M University, College Station, Texas 77843, and University of Nebraska, Lincoln, Nebraska 68508. Received February 6, 1974

Abstract: A compound has been found which provides an excellent model of certain essential features of the arginyl-phosphate interactions in the complex consisting of *Staphylococcus* nuclease, deoxythymidine 3',5'-diphosphate, and calcium ion. The crystal structure of this substance, bis(methylguanidinium) monohydrogen orthophosphate, $[\text{CN}_3\text{NHC}(\text{NH}_2)_2]_2\text{PO}_3\text{OH}$, has been determined. Crystal data: space group, $Fdd2-C_{2v}^{19}$; unit cell dimensions $a = 23.608(3)$, $b = 24.113(5)$, $c = 7.917(1)$ Å; $Z = 16$ for the formula unit $(\text{C}_2\text{N}_3\text{H}_5)_2\text{HPO}_4$; $d_{\text{calcd}} = 1.436$, $d_{\text{obsd}} = 1.430$ g cm⁻³. Using Zr-filtered Mo K α radiation a total of 5428 reflections having $\lambda^{-1} \sin \theta < 1.030$ were measured with an automated diffractometer. Using 4465 reflections adjudged to be statistically significant, the structure was solved by Patterson and Fourier methods and refined by full-matrix least squares to final unit-weighted and weighted residuals of 0.049 and 0.048, respectively. The crystallographically independent guanidyl groups are planar and each forms two hydrogen bonds to the HPO_4^{2-} ion through separate N-H groups. One phosphate oxygen atom participates in two hydrogen bonds, one from each guanidyl. The overall arrangement is very similar to, though not precisely the same as, that in the enzyme-inhibitor complex and provides an excellent model for the latter. The HPO_4^{2-} ions form hydrogen bonded pairs related by a twofold axis. The O-H-O distances here are relatively short, 2.544 and 2.503 Å, but it would appear that the hydrogens must be considered to be disordered rather than symmetrically located. The P-O distances of 1.514, 1.524, 1.556, and 1.567 Å are values that might be more typically expected for an H_2PO_4^- ion and may be a reflection of the hydrogen-bonding effect of the guanidyl ions.

Only in recent years has it become apparent that the guanidyl groups of arginine residues play an important role in binding and possibly even more active roles at the functional sites of both enzymes and non-catalytic proteins. Limiting reference only to those cases where it has been reasonably demonstrated that the chemical modification of arginine does occur at the binding or functional site of the protein molecule, functionally active arginine residues have been found in *E. coli* alkaline phosphatase,³ yeast inorganic pyrophosphatase,⁴ lactate dehydrogenase,⁵ D-amino acid oxidase,⁶ pepsin,⁷ ribonuclease T₁,⁸ carboxypeptidases A⁹ and B,¹⁰ and antibody combining sites directed against haptens containing such anions as arsonate, phosphate, and carboxylates.¹¹⁻¹⁵ Of particular interest

are those cases where the chemical modifications in solution can be correlated with the results from crystal structure analyses. Thus, the single arginine which is shown by chemical modification to be at the active site in carboxypeptidase A⁹ is very probably Arg-145 which Lipscomb, *et al.*,¹⁶ in their crystallographic studies, have found to bind the terminal carboxylate of peptides. Similarly, the estimation of three essential arginines per subunit of lactate dehydrogenase by Yang and Schwert from their chemical modification studies⁵ correlates very neatly with the recent observations from Rossmann's and Kaplan's laboratories. These observations indicate arginines-101, -109, and -171 are located at the active site in the crystal structure of the abortive lactate dehydrogenase-nicotinamide adenine dinucleotide-pyruvate ternary complex. The first of these arginines bridges the pyrophosphate linkage in the coenzyme, and the other two interact with the substrate.¹⁷

Our direct determination of the high resolution crystal structure of the ternary complex of the Staphylococcal nuclease with its potent competitive inhibitor, thymidine 3',5'-diphosphate and calcium ion has revealed that the 5'-phosphate of the inhibitor forms two hydrogen bonds each to the guanidinium ions of arginines-

(1) This study was supported by Grant GM13300 from the National Institute of General Medical Sciences, National Institutes of Health. The structure has been briefly described in a preliminary note: F. A. Cotton, E. E. Hazen, Jr., V. W. Day, S. Larsen, J. G. Norman, Jr., S. T. K. Wong, and K. H. Johnson, *J. Amer. Chem. Soc.*, **95**, 2367 (1973).

(2) Address correspondence to these authors at the Department of Chemistry, Texas A & M University, College Station, Texas 77843. (b) On leave, 1970-1971, from the Technical University of Copenhagen.

(3) F. J. M. Daemen, *Fed. Proc., Fed. Amer. Soc. Exp. Biol.*, **32**, 473 (1973).

(4) B. S. Cooperman and N. Y. Chiu, *Biochemistry*, **12**, 1676 (1973).

(5) P. C. Yang and G. W. Schwert, *Biochemistry*, **11**, 2218 (1972).

(6) A. Kotaki, M. Harada and K. Yagi, *J. Biochem. (Tokyo)*, **60**, 592 (1966).

(7) W.-Y. Huang and J. Tang, *J. Biol. Chem.*, **247**, 2704 (1972).

(8) A. Takahashi, *J. Biochem. (Tokyo)*, **68**, 659 (1970).

(9) J. F. Riordan, *Biochemistry*, **12**, 3915 (1973).

(10) M. M. Weber and M. Sokolovsky, *Biochem. Biophys. Res. Commun.*, **48**, 384 (1972).

(11) A. L. Grossberg and D. Pressman, *Biochemistry*, **7**, 272 (1968).

(12) M. H. Freedman, A. L. Grossberg, and D. Pressman, *Biochemistry*, **7**, 1941 (1968).


(13) M. H. Freedman, A. L. Grossberg, and D. Pressman, *J. Biol. Chem.*, **243**, 6186 (1968).

(14) G. L. Mayers, A. L. Grossberg, and D. Pressman, *Immunochemistry*, **9**, 196 (1972).

(15) G. L. Mayers, A. L. Grossberg, and D. Pressman, *Immunochemistry*, **10**, 37 (1973).

(16) W. N. Lipscomb, G. N. Reeke, Jr., J. A. Hartsuck, F. A. Quicho, and P. H. Bethge, *Proc. Roy. Soc., Ser. B*, **257**, 177 (1970).

(17) M. J. Adams, M. Buehner, K. Chandrasekhar, G. C. Ford, M. L. Hackert, A. Liljas, M. G. Rossmann, J. E. Smiley, W. S. Allison, J. Everse, N. O. Kaplan, and S. S. Taylor, *Proc. Nat. Acad. Sci. U. S.*, **70**, 1968 (1973).



35 and -87.¹⁸⁻²⁰ Additional hydrogen bonds occur between these guanidinium ions and other parts of the enzyme molecule to effectively lock the 5'-phosphate rigidly in the active site. Thus, our structural observations suggest a rather unique and highly specific functional role for these two arginine residues, a notion that is strongly reinforced by the observation of Chaiken and Anfinsen that the replacement of Arg-35 by either lysine or citrulline in semisynthetic variants of the nuclease completely abolishes enzymatic activity.^{21,22}

Since the chemical modification studies have indicated the presence of arginines at active sites of other phospholytic enzymes, namely, alkaline phosphatase,³ inorganic pyrophosphatase,⁴ and ribonuclease T₁,¹⁸ there may well be a general subclass of enzymes involved in phosphate metabolism having arginines at their active sites.

These structural and chemical observations as to the specificity and functional importance of the guanidinium-phosphate interactions in the Staphylococcal nuclease lead to a search for simpler model systems.^{1b,23} There are two purposes in examining model systems: (1) to confirm the plausibility of the overall interpretation of the enzyme-inhibitor interaction obtained from model fitting to the electron density maps of the nuclease-deoxythymidine 3',5'-diphosphate-Ca²⁺ complex, and (2) to obtain accurate structure parameters which can provide a more refined understanding of the interactions than that obtainable from the enzyme structure itself.

(18) Historically protein chemists have referred to such interactions between two such groups bearing formal charges as "salt linkages" or "salt bridges." Since this terminology was first introduced, there has been ample evidence from the crystal structures of smaller systems that interactions such as these normally occur *via* hydrogen bonds and since extensive charge delocalization is characteristic of most charged groups found in proteins, the use of this rather artificial terminology when the hydrogen bonding groups both have formal charges, could well be discontinued.

(19) A. Arnone, C. J. Bier, F. A. Cotton, V. W. Day, E. E. Hazen, Jr., D. C. Richardson, J. S. Richardson, and A. Yonath, *J. Biol. Chem.*, **246**, 2302 (1971).

(20) F. A. Cotton, C. J. Bier, V. W. Day, E. E. Hazen, Jr., and S. Larsen, *Cold Spring Harbor Symp. Quant. Biol.*, **36**, 243 (1971).

(21) I. M. Chaiken and C. B. Anfinsen, *J. Biol. Chem.*, **246**, 2285 (1971).

(22) In preliminary results from this laboratory, phenyl glyoxal has been found to modify the arginine residues of the nuclease with the concurrent loss of enzymatic activity.

(23) F. A. Cotton, V. W. Day, E. E. Hazen, Jr., and S. Larsen, *J. Amer. Chem. Soc.*, **95**, 4834 (1973).

The search for models was begun with substances obtainable from solutions containing methylguanidinium ion [CH₃NHC(NH₂)₂]⁺ (MGD) which closely resembles the end of the side chain of an arginyl residue and phosphate ions, H₂PO₄ⁿ⁻³. The first substance obtained and examined²³ was (MGD)₂H₂PO₄, which exhibits the type of double hydrogen-bonded interaction that is of interest, **1**, but does not mimic the entire arrangement seen in the enzyme-inhibitor complex, **2**.

When crystals of a second methylguanidinium phosphate, (MGD)₂HPO₄, were obtained, it was considered worthwhile to carry out another structure determination with the objective of finding a more complete facsimile of the enzyme-substrate arrangement, **2**. This objective has been accomplished, since, as shown here, (MGD)₂HPO₄ contains the arrangement, **3**, which very closely resembles **2**, though it does not precisely duplicate it.

Experimental Section

An aqueous solution containing equimolar quantities of methylguanidinium sulfate (Eastman Organic Chemicals) and Ba(OH)₂ was stirred overnight to precipitate BaSO₄ which was then filtered off. Half an equivalent amount of phosphoric acid was added to the filtrate to form an aqueous solution of bis(methylguanidinium) monohydrogen phosphate (pH 7.5). After evaporating this solution to near dryness, ethanol was added until the solution became turbid. Large single crystals of bis(methylguanidinium) monohydrogen phosphate grew over a period of several days.

Anal. Calcd for [(NH₂)₂CNHCH₃]₂(HPO₄): C, 19.67; N, 34.42; H, 7.02. Found: C, 19.42; N, 34.19; H, 7.03.

A spherical specimen 0.70 mm in diameter was ground from a larger crystal and glued to the end of a glass fiber with a tip diameter of 0.10 mm.

Precession photographs, used to determine a preliminary set of lattice constants, indicated orthorhombic, *mmm*, symmetry. The systematically absent reflections were those uniquely required by the noncentrosymmetric space group, *Fdd₂-C_{2v}*.¹⁹ This choice was fully supported by the positive results of sensitive tests for piezoelectricity and by the subsequent structure determination. The crystal was accurately centered on a Syntex PI full-circle goniometer and a total of 15 reflections, chosen to give a good sampling of reciprocal space and diffractometer settings ($2\theta_{\text{MoK}\alpha} > 50^\circ$), were used to align the crystal and calculate angular settings for each reflection. A least-squares refinement of the diffraction geometry for these 15 reflections, recorded at the ambient laboratory temperature of $21 \pm 1^\circ$ with Mo K α radiation ($\lambda(\text{Mo K}\alpha) 0.71069 \text{ \AA}$) gave the lattice constants $a = 23.608 \pm 0.003$, $b = 24.113 \pm 0.005$, and $c = 7.917 \pm 0.001 \text{ \AA}$. A unit cell content of 16 bis(methylguanidinium) monohydrogen phosphate molecules gives a calculated density of 1.436 g/cm³, in good agreement with the observed density of 1.430 g/cm³, measured by flotation in a mixture of dichloromethane and carbon tetrachloride.

Intensity measurements utilized Zr-filtered Mo K α radiation and the θ - 2θ scanning technique with a 3° takeoff angle and a standard-focus X-ray tube on a computer controlled Syntex PI diffractometer. A scanning rate of $3^\circ/\text{min}$ was employed for the scan between 2θ settings 1.0° above and below the calculated K α doublet values ($\lambda(\text{K}\alpha_1) 0.70926$ and ($\text{K}\alpha_2) 0.71354 \text{ \AA}$) of each reflection except for those reflections having $83.3^\circ < 2\theta < 94.1^\circ$ where a $2^\circ/\text{min}$ scanning rate was used. Background counts (each one lasting half the total scan time) were taken at both ends of the scan range. A total of 5428 independent reflections having $(\sin \theta/\lambda) < 1.030$ (four times the number of data in the limiting Cu K α sphere) were measured in concentric shells of increasing 2θ containing approximately 1400 reflections each. The six standard reflections, measured every 300 reflections as a monitor for possible misalignment and/or deterioration of the crystal, gave no indication of either.

The linear absorption coefficient of the crystal for Mo K α radiation is 0.26 cm^{-1} , yielding μR of 0.09 for the spherical crystal used. Since the absorption of X-rays by a spherical crystal having $\mu R = 0.09$ is essentially independent of scattering angle, no absorption correction was made, and the intensities were reduced to relative squared amplitudes, $|F_o|^2$, by means of standard Lorentz and polarization corrections.

Of the 5428 reflections examined, 963 were rejected as objectively unobserved by applying the rejection criterion, $I < \sigma(I)$, where $\sigma(I)$ is the counting statistics standard deviation in the observed intensity computed from

$$\sigma(I) = (C_t + k^2B)^{1/2}$$

C_t being the total count from scanning, k the ratio of scanning time to total background time (in this case $k = 1$), and B the total background count. The remaining 4465 observed intensities were used in the determination and refinement of the structure.

Structure determination was achieved through a combination of the heavy-atom technique, difference Fourier syntheses, and least-squares refinement. The wholly straightforward interpretation of the Patterson synthesis of the 697 $|F_o|^2$ data having $(\sin \theta/\lambda) \leq 0.52$ placed the phosphorus atoms in 16-fold general positions $(0,0,0; 0, 1/2, 1/2; 1/2, 0, 1/2; 1/2, 1/2, p) + (x, y, z; \bar{x}, \bar{y}, \bar{z}; 1/4 - x, 1/4 + y, 1/4 + z; 1/4 + x, 1/4 - y, 1/4 + z)$. These atomic coordinates, an isotropic thermal parameter, and a scale factor were varied in two cycles of isotropic full-matrix least-squares refinement.²⁴ This resulted in a conventional unweighted residual, $R_1 = 0.499$, for these low angle data.

$$R_1 = \frac{\sum ||F_o| - |F_c||}{\sum |F_o|}$$

A difference electron density map at this stage revealed the locations of the four phosphate oxygen atoms. Two cycles of least-squares refinement varying the scale factor, atomic coordinates, and isotropic temperature factors for phosphorus and oxygen atoms, respectively, gave $R_1 = 0.372$. A second electron density difference map clearly revealed the remaining ten non-hydrogen atoms of the asymmetric unit, all of which lie in general positions. Isotropic full-matrix refinement using unit weighting for the 15 non-hydrogen atoms gave $R_1 = 0.072$, for 697 reflections. All of the 4465 reflections were then included in a fully anisotropic least-squares minimization of the function $\sum w(|F_o| - k|F_c|)^2$ to give, with unit weighting (*i.e.*, all $w = 1$) $R_1 = 0.063$. This and all subsequent refinement cycles employed an anomalous dispersion correction²⁵ to the scattering factor of the phosphorus atom and a least-squares refinable extinction coefficient²⁶ of the form $F(x) = 1/(1 + 2x)^{1/2}$, where $x = gI$ and g refined to a final value of 0.36×10^{-7} . A Fourier difference synthesis based on the refined parameters afforded direct evidence for the placement of all hydrogen atoms. Further unit-weighted full-matrix least-squares cycles were used to refine hydrogen atoms isotropically and all other atoms anisotropically to give $R_1 = 0.049$ and a conventional weighted residual, $R_2 = 0.044$.

$$R_2 = \left\{ \frac{\sum w(|F_o| - |F_c|)^2 / \sum w|F_o|^2}{\sum w|F_o|^2} \right\}^{1/2}$$

Empirical weights ($w = 1/\sigma^2$) were then calculated from

$$\sigma = \sum_0^3 a_n |F_o|^n = 1.54 - 0.20 \times 10^{-1}F + 0.23 \times 10^{-3}F^2 - 0.35 \times 10^{-6}F^3$$

the a_n being coefficients derived from the least-squares fitting of the curve

$$||F_o| - |F_c|| = \sum_0^3 a_n |F_o|^n$$

The F_c values were calculated from the fully refined model using unit weighting. The final cycles of least-squares refinement utilized these weights and anomalous dispersion corrections for the phosphorus atom to refine hydrogen atoms isotropically and all other atoms anisotropically together with the scale factor and extinction coefficient to give final values of 0.049 and 0.048 for R_1 and R_2 , respectively. During this last cycle of refinement no parameter for non-hydrogen atoms shifted by more than 0.2σ , with the average shift being 0.03σ .

The following computer programs were employed in this work: MAGTAP and SCTFT2, data reduction programs written by V. Day; FORDAP, Fourier and Patterson synthesis program, a modified version of A. Zalkin's original program; ORFLSE, full-matrix least-squares refinement program, a highly modified version of Busing, Martin, and Levy's original ORFLS; ORFFE, bond lengths and angles

with standard deviations by Busing, Martin, and Levy; ORTEP-II, thermal ellipsoid plotting program by C. K. Johnson; MPLANE, least-squares mean plane calculation program from L. Dahl's group.

Results

The final coordinates and anisotropic thermal parameters for all atoms except hydrogen atoms are listed in Tables I and II, respectively; the refined positions and

Table I. Atomic Coordinates in Crystalline Bis(methylguanidinium) Monohydrogen Phosphate^a

Atom ^b type	Fractional coordinates		
	10 ⁶ x	10 ⁵ y	10 ⁴ z
Cation I			
N ₁	9959 (9)	10244 (7)	-3507 (2)
N ₂	9264 (11)	10208 (8)	-6414 (2)
N ₃	6717 (9)	2542 (6)	-4890 (3)
C ₁	12657 (16)	15602 (11)	-3417 (3)
C ₂	8655 (8)	7711 (6)	-4935 (3)
Cation II			
N ₁	23115 (7)	1538 (7)	1680 (3)
N ₂	28807 (7)	8798 (6)	2535 (3)
N ₃	20656 (7)	10454 (7)	1008 (3)
C ₁	27274 (12)	-2734 (9)	2017 (5)
C ₂	24204 (7)	6942 (7)	1758 (2)
Anion			
P	8675 (1)	-164 (1)	0
O ₁	4917 (5)	-2160 (5)	-1514 (2)
O ₂	5189 (6)	-1066 (7)	1645 (2)
O ₃	14016 (4)	-3632 (4)	73 (2)
O ₄	9965 (5)	5994 (4)	-199 (2)

^a Figures in parentheses are the estimated standard deviations. Coordinate listed without standard deviation is symmetry required. ^b Atoms numbered to agree with Figures 1-5.

isotropic thermal parameters of the hydrogen atoms are listed in Table III.²⁷ The rule used in the atom numbering scheme for bis(methylguanidinium) monohydrogen phosphate is as follows. Atoms of the methylguanidinium ions are grouped according to cation. A numerical subscript is used to differentiate atoms of the same non-hydrogen element. For each hydrogen atom the subscript letter and first subscript number indicate the atom to which it is covalently bonded, while the second numerical subscript distinguishes among hydrogen atoms attached to the same atom.

A projection of one asymmetric unit is presented in Figure 1; each atom is numbered in conformity with Tables I-IX and each non-hydrogen atom is represented by an ellipsoid having shape, orientation, and relative size consistent with the thermal parameters listed in Tables II. Bond lengths and angles in the molecular skeleton are presented in Figure 2 and are listed along with their estimated standard deviations in Tables IV and V; the dimensions of various interionic hydrogen bonds are listed in Table VI. The equations of the mean planes that partially characterize important subgroupings of atoms within the asymmetric unit specified by the coordinates of Tables I and III are given in Table VII,²⁷ and the displacements from these planes of the atoms constituting the asymmetric unit are listed in Tables VIII and IX.²⁷

(24) Scattering factors were taken from D. T. Cromer and J. B. Mann, *Acta Crystallogr., Sect. A*, **24**, 321 (1968).

(25) D. T. Cromer, *Acta Crystallogr.*, **18**, 17 (1965).

(26) W. H. Zachariasen, *Acta Crystallogr.*, **23**, 558 (1967).

(27) See paragraph at end of paper regarding supplementary material.

Table II. Anisotropic Thermal Parameters in Crystalline Bis(methylguanidinium) Monohydrogen Phosphate^a

Atom ^b type	B_{11}	B_{22}	B_{33}	B_{12}	B_{13}	B_{23}	$B_i^c, \text{\AA}^2$
Cation I							
N ₁	4.31 (8)	2.05 (5)	1.64 (5)	-0.60 (5)	0.04 (5)	0.13 (4)	2.40
N ₂	5.83 (11)	2.22 (6)	1.73 (5)	-0.65 (6)	-0.01 (6)	0.26 (4)	2.77
N ₃	5.17 (9)	2.07 (4)	2.09 (5)	-0.86 (5)	-0.10 (7)	0.16 (5)	2.75
C ₁	6.81 (16)	2.97 (8)	2.55 (8)	-1.91 (9)	-0.08 (9)	0.13 (7)	3.48
C ₂	2.93 (5)	1.80 (4)	1.73 (4)	0.06 (4)	0.11 (5)	0.27 (5)	2.07
Cation II							
N ₁	2.31 (5)	1.99 (4)	4.17 (8)	-0.32 (4)	-1.16 (6)	-0.01 (5)	2.52
N ₂	2.74 (5)	2.20 (4)	3.24 (6)	-0.65 (4)	-1.16 (6)	0.19 (5)	2.49
N ₃	2.50 (5)	2.10 (5)	3.83 (8)	0.28 (4)	-0.87 (5)	-0.38 (5)	2.62
C ₁	4.01 (9)	2.19 (6)	5.62 (14)	0.23 (6)	-2.12 (10)	-0.00 (7)	3.39
C ₂	1.98 (5)	2.01 (4)	2.33 (5)	-0.19 (4)	-0.19 (4)	-0.23 (4)	2.08
Anion							
O ₁	1.85 (3)	2.14 (4)	1.49 (3)	-0.12 (3)	-0.19 (3)	-0.19 (3)	1.79
O ₂	1.75 (4)	4.73 (7)	1.33 (3)	0.28 (4)	0.07 (3)	0.25 (4)	2.21
O ₃	1.57 (3)	1.82 (3)	2.35 (4)	0.25 (2)	-0.00 (3)	0.15 (3)	1.87
O ₄	2.76 (4)	1.35 (3)	3.08 (6)	-0.04 (3)	-0.82 (4)	0.06 (3)	2.19
P	1.41 (1)	1.46 (1)	1.38 (1)	0.04 (1)	-0.07 (1)	0.09 (1)	1.41

^a The number in parentheses that follows each B_{ij} value is the estimated standard deviation in the last significant figure. The B_{ij} 's in \AA^2 are related to the dimensionless β_{ij} employed during refinement as $B_{ij} = 4\beta_{ij}a_i a_j^*$. ^b Atoms numbered to agree with Figures 1-5. ^c Isotropic thermal parameter calculated from $B = 4[V^2 \det(\beta_{ij})]^{1/2}$.




Figure 1. A perspective view of one asymmetric unit. The atom numbering scheme is explained in the text. For clarity the actual thermal ellipsoids of the hydrogen atoms are not used.

Discussion

Overall Structure. As emphasized in Figure 3, the structure can be thought of as centered around a pair of monohydrogen phosphate ions which are held together by two strong hydrogen bonds. The O...O distances in these bonds are 2.544 and 2.503 \AA which means that they are close to or perhaps within the range where the hydrogen bonds might be symmetrical. Moreover, there is a crystallographic twofold axis which passes through the midpoints of the two hydrogen-bonded O...O pairs. However, this symmetry requirement could be satisfied either by having the hydrogen atoms on the twofold axis (truly symmetrical hydrogen bonds) or by having them off the axis but dis-

Table III. Refined Parameters for Hydrogen Atoms in Crystalline Bis(methylguanidinium) Monohydrogen Phosphate^a

Atom ^b type	Fractional coordinates			Isotropic thermal parameter, $B, \text{\AA}^2$
	10^3x	10^3y	10^3z	
Cation I				
H _{N1}	99 (1)	85 (1)	-262 (4)	0.7 (5)
H _{N21}	101 (1)	139 (1)	-646 (4)	1.5 (6)
H _{N22}	85 (1)	83 (1)	-731 (5)	1.5 (6)
H _{N31}	54 (2)	12 (1)	-570 (5)	2.1 (7)
H _{N32}	61 (1)	8 (1)	-399 (4)	0.2 (4)
H _{C11}	138 (2)	162 (2)	-243 (7)	4.1 (9)
H _{C12}	157 (1)	159 (1)	-415 (5)	2.1 (7)
H _{C13}	102 (2)	189 (2)	-393 (7)	5.3 (11)
Cation II				
H _{N1}	200 (1)	7 (1)	117 (3)	0.3 (4)
H _{N21}	309 (1)	66 (1)	310 (4)	0.9 (5)
H _{N22}	296 (1)	127 (1)	246 (5)	1.3 (5)
H _{N31}	211 (1)	138 (1)	112 (4)	1.1 (5)
H _{N32}	175 (1)	92 (1)	54 (4)	1.0 (5)
H _{C11}	256 (2)	-62 (3)	196 (8)	7.6 (16)
H _{C12}	309 (2)	-25 (2)	128 (7)	5.8 (13)
H _{C13}	278 (2)	-26 (2)	322 (8)	5.4 (14)
Anion				
H _{O1}	0.0	0.0	-138 (11)	5.8 (19)
H _{O2}	0.0	0.0	143 (12)	8.8 (25)

^a Figures in parentheses are the estimated standard deviations of the last significant digit. Coordinates listed without standard deviations are symmetry required. ^b Atoms numbered to agree with Figures 1-5.

ordered. The experimental data did not allow a choice; both the truly symmetric structure or one in which there are nearly symmetric but disordered bonds are consistent with the data. However, Speakman, in a recent review of short hydrogen bonds,²¹ has concluded that symmetrical hydrogen bonds are likely only when the O-H...O distance is less than 2.44 \AA , and on this basis we believe that the hydrogen bonds in this structure are unsymmetrical and disordered. Should these hydrogen atoms become ordered at lower temperatures, the re-

(28) J. C. Speakman, *Struct. Bonding (Berlin)*, 12, 141 (1972).

Table IV. Bond Lengths in Bis(methylguanidinium) Monohydrogen Phosphate^a

Type ^b	Bond length, Å			Type ^b	Bond length, Å		
	Cation I	Cation II	Av		Cation I	Cation II	Av
C ₁ -N ₁	1.442 (3)	1.448 (3)	1.445	C ₁ -H _{C11}	0.84 (5)	0.93 (6)	0.96
C ₂ -N ₁	1.322 (3)	1.330 (2)	1.327	C ₁ -H _{C12}	0.93 (4)	1.04 (5)	
C ₂ -N ₂	1.324 (3)	1.327 (2)		C ₁ -H _{C13}	1.07 (5)	0.96 (6)	
C ₂ -N ₃	1.328 (2)	1.331 (2)		Anion bond length, Å			
C ₂ -N				P-O ₁	1.567 (1)	1.562 (1.569) ^c	
Av	1.325	1.329		P-O ₂	(1.573) ^c		
				P-O ₃	1.556 (1)		
N ₁ -H _{N1}	0.82 (3)	0.86 (3)	0.86	P-O ₄	(1.565) ^c	1.519 (1.527) ^c	
N ₂ -H _{N21}	0.91 (3)	0.85 (3)					
N ₂ -H _{N22}	0.87 (4)	0.96 (3)					
N ₃ -H _{N31}	0.79 (4)	0.82 (3)					
N ₃ -H _{N32}	0.84 (3)	0.89 (3)					
				O ₁ -H _{O1}	1.276 (7)	1.27	
				O ₃ -H _{O3}	1.263 (13)		

^a The figure in parentheses following each individual distance is the estimated standard deviation. ^b Atoms numbered to agree with Figures 1-5 and Tables I and III. ^c Bond length corrected for libration of HPO₄²⁻ group as a rigid body according to V. Schomaker and K. N. Trueblood, *Acta Crystallogr., Sect. B*, **24**, 63 (1968).

sulting crystals of this compound could prove to be ferroelectric.²⁹

Surrounding each central dimer of phosphate ions is a total of 12 methylguanidinium ions, six from each of the two crystallographically distinct cations. These are linked to the phosphate dimer by a total of 18, 20, or 22 hydrogen bonds, namely, ten from cation II and 8, 10, or 12 from cation I. The hydrogen bonding pattern of a single guanidinium ion is illustrated in Figure 4 for cation I and in Figure 5 for cation II. Considering a single phosphate dimer, cation I and II both form a single hydrogen bond to O₃ of each phosphate ion for a total of four. Cation II forms two pairs of hydrogen bonds bridging across the phosphate dimer, giving altogether four of this pattern and a subtotal of eight. Cation I and cation II each form a pair of hydrogen bonds to two oxygen atoms of one phosphate ion, giving eight H bonds of this type for a subtotal of 16. This type of guanidinium-phosphate interaction, which is best illustrated in Figures 1 and 2, has also been observed in the structure of methylguanidinium dihydrogen orthophosphate^{1b, 23} and in the structure of propylguanidinium diethylphosphate.³⁰ However, the most significant aspect of this type of paired hydrogen bond interaction between a guanidinium ion and a phosphate ion is that it provides an excellent model for the interaction of arginines-35 and -87 of the Staphylococcal nuclease with the 5'-phosphate of its potent inhibitor, thymidine 3',5'-diphosphate, as illustrated diagrammatically by 2.

The final type of guanidinium-phosphate interaction observed in this structure is the N₁₃-H--O₂ of cation I (Figure 4) with an N-O distance of 2.90 Å which is paired with another N-O interaction to a different oxygen atom of the same phosphate, *i.e.*, N₁₂-H--O₄ with an N-O distance of 3.17 Å. This distance is somewhat long to be considered a real hydrogen bond, and the final N-O interaction shown in Figure 4, that of N₁₂-H--O₂ with an N-O distance of 3.27 Å, is even more doubtful. The grand total of N to O hydrogen bonding interactions to a single dimer of phosphate ions is thus 18, 20, or 22, the latter two numbers including one or both of the pair of N-O interaction over 3 Å. This

(29) W. C. Hamilton and J. A. Ibers, "Hydrogen Bonding in Solids," W. A. Benjamin, New York, N. Y., 1968, pp 238-255.

(30) S. Furberg and J. Solbakk, *Acta Chem. Scand.*, **26**, 3699 (1972).




Figure 2. Bond lengths and angles for the asymmetric unit as seen in Figure 1. A complete set of values and standard deviations are listed in Tables IV, V, and VI.

final pattern of guanidinium-phosphate interaction, showing in this structure a third paired cyclic N-O system formed by one strong hydrogen bond and a second rather weak one, may well have some biochemical significance. In the structure of propylguanidinium diethylphosphate³⁰ one paired interaction with two strong hydrogen bonds is observed, but there is also a second pair with the one strong and one weak pattern that we also observe in this structure. The Staphylococcal nuclease is, of course, a phosphodiesterase with a degree of base specificity. Its initial interaction with

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.