

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PAZEO® safely and effectively. See full prescribing information for PAZEO.

**PAZEO (olopatadine hydrochloride ophthalmic solution) 0.7%
For topical ophthalmic administration.
Initial U.S. Approval: 1996**

INDICATIONS AND USAGE
PAZEO is a mast cell stabilizer indicated for the treatment of ocular itching associated with allergic conjunctivitis. (1).

DOSAGE AND ADMINISTRATION
The recommended dose is one drop in each affected eye once a day. (2)

DOSAGE FORMS AND STRENGTHS
Ophthalmic solution: 7.76 mg of olopatadine hydrochloride in one mL of solution (0.7%) in a four mL bottle. (3)

CONTRAINDICATIONS
None.

WARNINGS AND PRECAUTIONS
Contamination of Tip and Solution. To prevent contaminating the dropper tip and solution, do not touch the eyelids or surrounding areas with the dropper tip of the bottle. (5.1)

ADVERSE REACTIONS
The most common adverse reactions (2-5%) were blurred vision, superficial punctate keratitis, dry eye, abnormal sensation in eye, and dysgeusia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Alcon Laboratories, Inc. at 1-800-757-9195 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA approved patient labeling.

Revised: 04/2016

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
- 6 ADVERSE REACTIONS**
- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.3 Nursing Mothers
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use

- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

PAZEO is indicated for the treatment of ocular itching associated with allergic conjunctivitis.

2 DOSAGE AND ADMINISTRATION

The recommended dosage of PAZEO is to instill one drop in each affected eye once a day.

3 DOSAGE FORMS AND STRENGTHS

Ophthalmic solution: 7.76 mg of olopatadine hydrochloride in one mL solution (0.7%) in a 4 mL bottle.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Contamination of Tip and Solution

As with any eye drop, care should be taken not to touch the eyelids or surrounding areas with the dropper tip of the bottle to prevent contaminating the tip and solution. Keep bottle tightly closed when not in use.

5.2 Contact Lens Use

Patients should not wear a contact lens if their eye is red.

The preservative in PAZEO solution, benzalkonium chloride, may be absorbed by soft contact lenses. Patients who wear soft contact lenses and whose eyes are not red, should be instructed to wait at least five minutes after instilling PAZEO before they insert their contact lenses.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In a randomized, double-masked, vehicle-controlled trial, patients at risk for developing allergic conjunctivitis received one drop of either PAZEO (N=330) or vehicle (N=169) in both eyes for 6 weeks. The mean age of the population was 32 years (range 2 to 74 years). Thirty-five percent were male. Fifty-three percent had brown iris color and 23% had blue iris color. The most commonly reported adverse reactions occurred in 2-5% of patients treated with either PAZEO or vehicle. These events were blurred vision, dry eye, superficial punctate keratitis, dysgeusia and abnormal sensation in eye.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate or well-controlled studies with PAZEO in pregnant women. Olopatadine caused maternal toxicity and embryofetal toxicity in rats at levels 1,080 to 14,400 times the maximum recommended human ophthalmic dose (MRHOD). There was no toxicity in rat offspring at exposures estimated to be 45 to 150 times that at MRHOD. Olopatadine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Animal Data

In a rabbit embryofetal study, rabbits treated orally at 400 mg/kg/day during organogenesis showed a decrease in live fetuses. This dose is 14,400 times the MRHOD, on a mg/m² basis.

An oral dose of 600 mg/kg/day olopatadine (10,800 times the MRHOD) was shown to be maternally toxic in rats, producing death and reduced maternal body weight gain. When administered to rats throughout organogenesis, olopatadine produced cleft palate at 60 mg/kg/day (1080 times the MRHOD) and decreased embryofetal viability and reduced fetal weight in rats at 600 mg/kg/day. When administered to rats during late gestation and throughout the lactation period, olopatadine produced decreased neonatal survival at 60 mg/kg/day and reduced body weight gain in offspring at 4 mg/kg/day. A dose of 2 mg/kg/day olopatadine produced no toxicity in rat offspring. An oral dose of 1 mg/kg olopatadine in rats resulted in a range of systemic plasma area under the curve (AUC) levels that were 45 to 150 times higher than the observed human exposure [9.7 ng·hr/mL] following administration of the recommended human ophthalmic dose.

8.3 Nursing Mothers

Olopatadine has been identified in the milk of nursing rats following oral administration. Oral administration of olopatadine doses at or above 4 mg/kg/day throughout the lactation period produced decreased body weight gain in rat offspring; a dose of 2 mg/kg/day olopatadine produced no toxicity. An oral dose of 1 mg/kg olopatadine in rats resulted in a range of systemic plasma area under the curve (AUC) levels that were 45 to 150 times higher than the observed human exposure [9.7 ng·hr/mL] following administration of the recommended human ophthalmic dose. It is not known whether topical ocular administration could result in sufficient systemic absorption to produce detectable quantities in the human breast milk. Nevertheless, caution should be exercised when PAZEO is administered to a nursing mother.

8.4 Pediatric Use

The safety and effectiveness of PAZEO have been established in pediatric patients two years of age and older. Use of PAZEO in these pediatric patients is supported by evidence from adequate and well-controlled studies of PAZEO in adults and an adequate and well controlled study evaluating the safety of PAZEO in pediatric and adult patients.


8.5 Geriatric Use

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

11 DESCRIPTION

PAZEO is a sterile ophthalmic solution containing olopatadine, which is a mast cell stabilizer, for topical administration to the eyes. Olopatadine hydrochloride is a white, crystalline, water-soluble powder with a molecular weight of 373.88 and a molecular formula of $C_{21}H_{23}NO_3 \cdot HCl$.

The chemical structure is presented below:



Chemical Name: 11-[(Z)-3-(dimethylamino) propylidene]-6,11-dihydrodibenz[b,e] oxepin-2-acetic acid, hydrochloride.

Each mL of PAZEO solution contains an active ingredient [7.76 mg of olopatadine hydrochloride (7 mg olopatadine)] and the following inactive ingredients: povidone; hydroxypropyl-gamma-cyclodextrin; polyethylene glycol 400; hypromellose; boric acid; mannitol; benzalkonium chloride 0.015% (preservative); hydrochloric acid/sodium hydroxide (to adjust pH); and purified water.

PAZEO solution has a pH of approximately 7.2 and an osmolality of approximately 300 mOsm/kg.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Olopatadine is a mast cell stabilizer and a histamine H_1 antagonist. Decreased chemotaxis and inhibition of eosinophil activation has also been demonstrated.

12.3 Pharmacokinetics

In healthy subjects, topical ocular dosing of 1 drop of PAZEO once daily for 7 days into both eyes resulted in mean \pm SD (range) steady state plasma olopatadine C_{max} and AUC_{0-12} of 1.6 ± 0.9 ng/mL (0.6 to 4.5 ng/mL) and 9.7 ± 4.4 ng*h/mL (3.7 to 21.2 ng*h/mL), respectively. The olopatadine C_{max} and AUC_{0-12} after the first dose were similar to those measured on day 7 in these subjects, suggesting that there was no systemic accumulation of olopatadine after repeated topical ocular dosing with PAZEO. The median (range) time to achieve peak olopatadine concentrations (T_{max}) was 2.0 hours (0.25 to 4 hours). The mean \pm SD (range) elimination half-life of olopatadine was 3.4 ± 1.2 hours (2 to 8 hours). N-oxide olopatadine (M3) was

detected during the first 4 hours after bilateral topical ocular dosing of PAZEO in approximately half of the subjects and in less than 10% of the total plasma samples collected, at concentrations not exceeding 0.121 ng/mL on day 1 and 0.174 ng/mL on day 7. None of the plasma samples from these subjects had mono-desmethyl olopatadine (M1) concentrations that were above the lower limit of quantitation (0.05 ng/mL) of the PK assay.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity

Olopatadine administered orally was not carcinogenic in mice and rats in doses up to 500 mg/kg/day and 200 mg/kg/day, respectively. Based on a 35 μ L drop size and a 60 kg person, these doses are approximately 4,500 and 3,600 times the MRHOD, on a mg/m² basis.

Mutagenesis

No mutagenic potential was observed when olopatadine was tested in an in vitro bacterial reverse mutation (Ames) test, an in vitro mammalian chromosome aberration assay or an in vivo mouse micronucleus test.

Impairment of fertility

Olopatadine administered at an oral dose of 400 mg/kg/day (approximately 7,200 times the MRHOD) produced toxicity in male and female rats, and resulted in a decrease in the fertility index and reduced implantation rate. No effects on reproductive function were observed at 50 mg/kg/day (approximately 900 times the MRHOD).

14 CLINICAL STUDIES

The efficacy of PAZEO was established in two randomized, double-masked, placebo-controlled, conjunctival allergen challenge (CAC) clinical studies in patients with a history of allergic conjunctivitis (Studies 1 and 2).

In Study 1, patients were randomized to receive one of the following study treatments: PAZEO, PATADAY, or vehicle ophthalmic solutions. In Study 2, patients were randomized to receive one of the following study treatments: PAZEO, PATADAY, PATANOL, or vehicle ophthalmic solutions.

Patients were evaluated with an ocular itching severity score ranging from 0 (no itching) to 4 (incapacitating itch) at several time points after CAC administration. Table 1 displays the mean ocular itching severity scores after ocular administration of a specific antigen using the CAC model in Studies 1 and 2, respectively. A one unit difference compared to vehicle is considered a clinically meaningful change in the ocular itching severity score.

PAZEO demonstrated statistically significantly improved relief of ocular itching compared to vehicle at 30-34 minutes, 16 hours, and 24 hours after study treatment. PAZEO demonstrated statistically significantly improved relief of ocular itching compared to PATADAY at 24 hours after study treatment, but not at 30-34 minutes after study treatment.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.